Fluence rate-dependent kinetics of light-triggered liposomal doxorubicin assessed by quantitative fluorescence-based endoscopic probe

Daniel J. Rohrbach[1], Kevin A. Carter[2], Dandan Luo[2], Shuai Shao[2], Semra Aygun-Sunar[3], Jonathan F. Lovell[2], and Ulas Sunar[3],*

[1]Agilent Technologies, Winooski, Vermont, VT 05404, USA
[2] Department of Biomedical Engineering, University at Buffalo, Buffalo, NY 14260, USA
[3] Department of Biomedical, Engineering, Stony Brook University, Stony Brook, NY, 11794, USA

*Correspondence: ulas.sunar@stonybrook.edu

Abstract: Liposomal doxorubicin (Dox), a treatment option for recurrent ovarian cancer, often suffers from suboptimal biodistribution and efficacy, which might be addressed with precision drug delivery systems. Here, we introduce a catheter-based endoscopic probe designed for multispectral, quantitative monitoring of light-triggered drug release. This tool utilizes red-light photosensitive porphyrin-phospholipid (PoP), which is encapsulated in liposome bilayers to enhance targeted drug delivery. By integrating diffuse reflectance and fluorescence spectroscopy, our approach not only corrects the effects of tissue optical properties but also ensures accurate drug delivery to deep-seated tumors. Preliminary results validate the probe effectiveness in controlled settings, highlighting its potential for future clinical adaptation. This study sets the stage for in vivo applications, enabling the exploration of next-generation treatment paradigms for the management of cancer by optimizing chemotherapy administration with precision and control.

Keywords: doxorubicin drug concentration; diffuse reflectance spectroscopy; diffuse fluorescence spectroscopy; light-triggered release; porphyrins; photodynamic therapy; endoscopic probe.

1. Introduction

Liposomal doxorubicin (Dox) is a commonly utilized therapeutic agent for recurrent ovarian carcinoma, yet its efficacy is often hindered by inefficient drug delivery to tumor cells [1–10]. To address these limitations, nanocarriers such as liposomes have been developed to enhance drug distribution and efficacy. Among these, porphyrin-phospholipid (PoP) liposomes present a unique advantage, allowing controlled drug release triggered by near-infrared (NIR) light with precise spatial and temporal resolution [1,5–8,10–16].

This light-triggered release mechanism not only improves targeted drug delivery but also minimizes systemic side effects by confining the release to specific sites and times. For emerging nanocarriers like PoP-liposomes, obtaining detailed information on local drug concentration distribution is crucial for assessing therapeutic efficacy [5,17–27,27–36]. Both Dox and PoP exhibit fluorescence, which enables the use of fluorescence spectroscopy or imaging for this purpose. However, in turbid media, background optical properties can distort the raw fluorescence signal, making it challenging to accurately quantify drug concentration [24,28,37–46]. Additionally, these optical properties can attenuate the treatment light, delaying drug release.

To overcome the effect of background optical properties, quantitative fluorescence techniques are needed. Several spectroscopic methods have been implemented for quantification of drug concentration *in vivo* [24,28,45,46,37–44]. In this study, we utilized a combination of diffuse reflectance and fluorescence spectroscopy to quantify Dox fluorescence while correcting for variations caused by absorption and scattering at both excitation and emission wavelengths. Using a catheter-based endoscopic probe, we measured the localized concentration of Dox released by NIR light in vitro and in an animal carcass. The probe, which fits into an endoscope's working channel, provides multispectral data for Dox (~590 nm) and PoP (~680–730 nm). This approach enables precise quantification of each component in multi-functional constructs, offering significant potential for intra-tumoral applications in thick and deep-seated tumors.

2. Materials and Methods

2.1. Drug Preparation

The details of our PoP Dox (PoP-D) liposome formulation has been described elsewhere [1,12,47,48] . PoP-liposomes incorporated PoP that was in a manner as recently reported [49]. Briefly, PoP-liposomes were synthesized from pyro-lipid through esterification of pyro with lyso-C16-PC, using 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) and 4-dimethylaminopyridine (DMAP) in chloroform. The liposomes were formed by dispersing porphyrin-lipid, PEGylated-lipid, cholesterol, and distearoylphosphatidylcholine in chloroform, followed by solvent evaporation. A 20 mg/mL lipid solution was extruded through a high-pressure lipid extruder with a 250 mM ammonium sulfate solution using polycarbonate membranes of 0.2, 0.1, and 0.08 μm pore size, sequentially stacked and passed through the extruder 10 times. Free ammonium

sulfate was removed by overnight dialysis in a 10% sucrose solution with 10 mM   83
HEPES at pH 7. Dox was loaded by incubating the liposomes at 60 °C for 1 hour,   84
achieving a loading efficacy of over 95% as confirmed by G-75 column tests. The   85
self-assembly status and elution position of PoP-liposomes were tracked using   86
420 nm excitation and 670 nm emission, while Dox was detected using 480 nm   87
excitation and 590 nm emission in a fluorescence plate reader (TECAN Safire).   88

2.2. Instrumentation   89

The release kinetics of doxorubicin from the PoP-D liposomes was determined   90
using a custom fluorescence spectroscopy setup. A cuvette was placed in a fiber-   91
coupled 4-way holder (Ocean Optics) with one port connected to a 455nm laser   92
(Thorlabs) and a 90-degree port connected to the sensitive channel of a   93
spectrometer (Ocean Optics). The system was controlled with a custom LabView   94
program. For release, a 660nm LED (Mightex) was placed above and focused into   95
the cuvette. Measurements were acquired before PoPD liposomes were added   96
(PBS only) and after each round of treatment light (every 30 seconds for 30   97
minutes). Three measurements were acquired at each time point for averaging.   98

100
101

Fig. 1. Instrument setup. (a). Diagram of interstitial measurement and treatment system (b) Picture   102
of the setup. (c) The picture of the interstitial needle probe. 18-gauge needle is shown for size com-   103
parison. The inset shows the layout of the probe face.   104

To better mimic the *in vivo* case, a custom diffuse optical spectroscopy system   106
combining diffuse reflectance spectroscopy (DRS) and diffuse fluorescence spec-   107
troscopy (DFS) was used as shown in **Figures 1a** and **1b** [31,35,48,50–53]. Briefly,   108
the DRS setup consisted of a tungsten-halogen broadband white light (HL-2000-   109
FHSA, Ocean Optics) as the source and the Master channel of the spectrometer as   110
the detector. The light was directed to the target with one source fiber and the   111
diffusely reflected light was collected with one detector fiber (200 μm dia), with   112
the source detector separation (SD) of 260 μm. For the DFS, a 455 nm laser diode   113
was used as the excitation source and the Slave channel of the spectrometer was   114
used as the detector. A 500 nm long pass filter allowed measuring the fluorescence   115
signal (SD=260 μm) along with a "leakage" signal of the excitation light. The re-   116
flected excitation light of the laser diode and the reflected white light from DRS   117
were used to correct the raw DFS signal. All 4 fibers were contained in a narrow   118
18-gauge needle probe as shown in **Figure 1c.**   119

Fig. 2. Dox and PoP (Pyro) fluorescence quantification. (a) The basis spectra of autofluorescence (Auto), Doxorubicin (Dox), and PoP were used for quantifying fluorescence concentrations. (b) A representative raw fluorescence signal measured pre- and post-Dox release.

2.3 Doxorubicin Quantification

A quantitative fluorescence spectroscopy model was used, similar to Kim *et. al.* [38,41]. Correction is needed to account for the attenuation of the excitation light into the sample as well as the attenuation of emission light out of the sample.

$$FL(\lambda) = \frac{1}{R_{ex}} * \frac{f_{em}(\lambda)}{R_{em}(\lambda)}$$

The raw fluorescence signal ($f_{em}(\lambda)$) was first divided by the white light reflectance spectrum ($R_{em}(\lambda)$), and then scaled by the peak value of the excitation light leakage at 455nm (R_{ex}). The corrected emission (FL(λ)) was then fit to a model of tissue fluorescence, which included background autofluorescence (AF), Dox, and the PoP liposomes (HPPH, also called Pyro),

$$FL(\lambda) = C_{AF}AF(\lambda) + C_{Dox}Dox(\lambda) + C_{Pyro}Pyro(\lambda),$$

whose corresponding basis spectra are shown in **Figure 2a**. The value C_{Dox} is then directly related to the true Dox concentration. Sample measurements taken pre- and post-light treatment show the changes in the raw fluorescence due to Dox release (**Fig. 2b**).

2.3 Phantom and animal preparation

Phantoms were prepared using Intralipid 20% (Fresenius Kabi) for scattering and India Ink (Higgins) for absorption. Four phantoms were prepared with different combinations of optical properties: μ_a = 0.5 and 1.5 cm^{-1}, μ_s' = 20 and 30 cm^{-1} at 455 nm; μ_a = 0.4 and 1.2 cm^{-1}, μ_s' = 15.8 and 23.8 cm^{-1} at 590 nm. Each phantom had a total volume of 120 mL. For the calibration of Dox concentration, a stock solution of 0.5 mg/mL free-Dox was used. After baseline reflectance and fluorescence measurements of each phantom, increasing volumes of free-Dox were added to

each (400, 800, 1200, 1600, and 2000 μL), with reflectance and fluorescence measurements acquired at each addition.

Dox release was performed in phantoms prepared as above with $\mu_s' = 20\ \text{cm}^{-1}$ and $\mu_a = 0.5$ and $1.0\ \text{cm}^{-1}$ at 455 nm and a total volume of 70 mL. A stock solution of 2.59 mg/mL PoP liposomes was prepared, as described in the previous section. The phantom was placed on a stir plate set to 15 rpm. Simulated interstitial measurements were acquired by placing the needle probe into the liquid phantom to acquire measurements. Fluorescence measurements were acquired before and after the addition of PoP liposomes (200 μL). The stir plate was increased to 30 rpm to ensure a uniform release, and the treatment light (1.72 cm diameter spot) was turned on. Fluorescence measurements were acquired every 4 minutes.

Next, a recently sacrificed nude mouse was used for mimicking *in vivo* measurements to check and optimize the signal. The mouse was injected subcutaneously with 50 μL of a lightly scattering medium ($\mu_s' = 5\ \text{cm}^{-1}$ at 455 nm) containing 22 μg/mL PoP-Liposomes to mimic a tumor under the surface and reflectance and fluorescence spectroscopy measurements were performed. Three measurements were acquired for pre-release baseline levels by placing the probe against the tumor-mimicking sample. The treatment light (1.0 cm diameter) was directed onto the injection site, covering it completely. Fluorescence measurements were acquired every 4 minutes to assess the release kinetics during treatment.

3. Results and Discussion

Fig. 3. Phantom calibration. (a) A schematic of the phantom optical properties (b) Raw Dox fluorescence counts show the effect of background optical property variation. Error bars represent the standard deviation between phantoms. (c) Corrected fluorescence shows a linear response to increasing Dox concentration and much less variation between phantoms. Error bars represent the standard deviation between phantoms.

Calibration with free-Dox provided the conversion between corrected fluorescence values and Dox concentrations as shown in **Figure 3**. **Figure 3a** indicates a representative schematics of the four phantoms and their optical properties. **Figure 3b** shows the mean raw fluorescence signal for all four

phantoms at each titration of Dox. The large error bars represents the standard 196
deviation of the four phantoms and highlight the effects of background optical 197
properties on the raw fluorescence. **Figure 3c** shows the corrected fluorescence 198
(C_{Dox} from Equation 2). The corrected fluorescence has much smaller error bars, 199
highlighting the improved quantification. In addition, after the correction, the 200
slope of the line directly relates the relationship between the corrected Dox 201
fluorescence to the true Dox concentration, so that one can extract the local Dox 202
concentration for unknown sample. 203

Next, we investigate the release kinetics with respect to light irradiation fluence 204
rate. Our fluorescence spectroscopy measurements showed a difference in drug 205
release kinetics with fluence-rate dependent manner (**Fig. 4a**). The lower fluence 206
rate only started releasing Dox after 450 seconds and was fully released by 900 207
seconds while the higher fluence rate started releasing Dox after 270 seconds and 208
was fully released by 510 seconds. This is a difference in release rate of 450 209
seconds for the low fluence rate and 240 seconds for the high fluence rate. 210
However, when plotted as a function of light dose (mW/cm^2) the two curve are 211
much more similar (**Fig. 4b**). As **Figures 4c** and **4d** indicate, there was a decrease 212
in the Pyro fluorescence as well, approximately from 4000 counts to 3000 counts, 213
indicates the photobleaching of the porphyrin component. 214

Fig. 4. The release kinetics. The time to complete release depends on the fluence rate. (a) Dow 217
release as a function of time for different fluence rates. (b) Dox release as a function of fluence rate. 218
(c) Pyro-fluorescence signal as a function of time. (d) Pyro-fluorescence signal, as a function of 219
fluence rate, indicates the photobleaching of the porphyrin component. 220

Fig. 5. (a) Background absorption differences showed different final values of raw fluorescence after 223
release. (b) Corrected fluorescence fitting showed the same final dox concentration after release for 224
the two phantoms. (c) Pyro (PoP, HPPH) fluorescence showed a slight decrease throughout 225
treatment. (d) The release in a mouse model. 226

The fluence rate can be precisely controlled for optically clear phantoms in a 227
cuvette. However, for turbid media, the true local-site fluence is unknown since 228
the background optical properties will affect the delivered treatment light. The 229
background optical properties will also affect the determination of fluorophore 230
concentration. To mimic this case, we check the kinetics for the high and low- 231
absorption cases. As **Figure 5a** shows, the raw fluorescence for phantoms with the 232
same PoPD liposome concentration showed different rates of release as well as 233
different values at full release. The phantom with lower background absorption 234
showed a final Dox signal of 2007.1 ± 44.8 counts, while the higher absorbing 235
phantom showed 1656.5 ± 46.1 counts, a difference of 17.5%. However, after 236
quantifying the Dox concentration (**Fig. 5b**), the full release for the low and high- 237
absorbing phantoms was 11.4 ± 1.5 µg/mL vs. 11.5 ± 1.1 µg/mL, respectively, a 238
difference of less than 1%. As **Figure 5c** shows, the signal from Pyro (PoP, HPPH) 239
decreased slightly throughout treatment for both the low and high-absorbing 240
phantom (26.8% and 10.4%, respectively). This decrease is likely due to the 241
photobleaching of the Pyro. 242

3. Discussion 244

For the mouse experiment, the expected value of Dox concentration was 22 µg/mL 245
based on the concentration of the preparation. The corrected fluorescence fitting 246
determined the Dox concentration to be 19.5 µg/mL, a difference of 11.4%. This 247
difference could be due to the partial volume effect due to the skin layer. It is clear 248
that Dox accumulates at the local site with time, which is the main aim for an 249

effective cell kill. It is interesting to note that the PoP amount was photobleached with time. This aligned well with the previous observations of others and ours related to photodynamic therapy (PDT) treatment of tumors with HPPH. Since the release light intensity is on the same order of magnitude as PDT, where photosensitizers (like HPPH) are photobleached due to porphyrin, oxygen, and light interaction. It would be interesting to investigate the combination effect (PDT and Dox-chemo) for improved cell-killing efficacy.

An important part of the fluorescence quantification is the correction for attenuation of excitation light. This was accomplished by allowing some excitation light to "leak" into the detector fiber. This leakage corresponds to the reflected excitation light and depends on the optical properties at the excitation wavelength. Kim et al. used a similar process for fluorescence quantification, but instead of measuring the R_{ex}, they calculated R_{ex} using knowledge of the tissue optical properties. For our method it is not necessary to measure the tissue optical properties, which enables the fluorescence quantification to be performed very quickly. However, in cases of very high absorption or longer source detector separations where the excitation leakage may not be seen, the calculation method will provide more reasonable estimates of R_{ex}. In these studies, the R_{ex} signal was always seen.

In our previous work, we showed that spatial frequency domain imaging (SFDI) can quantify the release of doxorubicin [1,47,48]. While the SFDI technique is well-suited for superficial surfaces, it may not work for tumors located deeper inside the body. In those cases, quantitative fluorescence spectroscopy with a needle probe can be used.

However, this approach comes with several limitations. While the catheter-based probe provides precise point measurements, it lacks the capability for spatially resolved imaging over larger areas, which could be beneficial for identifying heterogeneous drug distribution in tumors. The system's ability to measure fluorescence at depths is limited due to limited source-detector separation (here 260 μm). We used the animal carcass, which may not fully replicate the heterogeneity and complexity of in vivo cases. The study focused on a specific set of light parameters (fluence rate and dose). A broader exploration of different fluence rates and dosimetry schemes is needed to establish universally applicable guidelines for light-triggered drug release. To address these limitations, future research could explore multi-modal imaging approaches combining spatial resolution with the point-based probe, validate the methodology in larger and more clinically relevant animal models, and conduct pilot clinical studies to assess feasibility and efficacy in humans.

5. Conclusions

In this work, we present results from a custom-built DRS and DFS system for monitoring absolute Dox and PoP concentration changes during light-triggered release in phantoms and in a mouse carcass. We corrected for background optical properties to obtain the true dox concentration. Quantitative Dox concentration release kinetics showed a faster release with respect to laser irradiation. These

results show the potential for interstitial measurements to assess Dox release in deeply seated tumors where wide-field imaging techniques cannot reach.

Acknowledgments: The authors acknowledge the funding support from NIH/NCI R01 (5R01CA243164).

Author Contributions: U.S. conceived and designed the experiments; D.J.R. performed the experiments; D.J.R. analyzed the data; D.J.R., U.S., SAS, and J.F. Lovell wrote the paper.

Funding: This research was funded by NIH/NCI R01 (5R01CA243164).

Institutional Review Board Statement: "Not applicable."

Informed Consent Statement: "Not applicable."

Data Availability Statement: The data that support the findings of this study are available from the corresponding author upon reasonable request.

Conflicts of Interest: The authors declare no conflict of interest. The funding sponsors had no role in the design of the study, in the collection, analysis, or interpretation of data, in the writing of the manuscript, and in the decision to publish the results. The research conducted in connection with this article was performed prior to Dr. Rohrbach's employment with Agilent. The research conducted and opinions expressed in this article have not been endorsed or approved by Agilent.

References

1.   J. Kress, D. J. Rohrbach, K. A. Carter, D. Luo, C. Poon, S. Aygun-Sunar, S. Shao, S. Lele, J. F. Lovell, and U. Sunar, "A dual-channel endoscope for quantitative imaging, monitoring, and triggering of doxorubicin release from liposomes in living mice," Sci. Rep. **7**(1), (2017).

2.   H. Ye, A. A. Karim, and X. J. Loh, "Current treatment options and drug delivery systems as potential therapeutic agents for ovarian cancer: a review," Mater Sci Eng C Mater Biol Appl **45**, 609–619 (2014).

3.   M. Zheng, C. Yue, Y. Ma, P. Gong, P. Zhao, C. Zheng, Z. Sheng, P. Zhang, Z. Wang, and L. Cai, "Single-step assembly of DOX/ICG loaded lipid--polymer nanoparticles for highly effective chemo-photothermal combination therapy," ACS Nano **7**(3), 2056–2067 (2013).

4.   G. Kong, G. Anyarambhatla, W. P. Petros, R. D. Braun, O. M. Colvin, D. Needham, and M. W. Dewhirst, "Efficacy of liposomes and hyperthermia in a human tumor xenograft model: Importance of triggered drug release," Cancer Res. **60**(24), 6950–6957 (2000).

5.   D. Luo, K. A. Carter, E. A. G. Molins, N. L. Straubinger, J. Geng, S. Shao, W. J. Jusko, R. M. Straubinger, and J. F. Lovell, "Pharmacokinetics and pharmacodynamics of liposomal chemophototherapy with short drug-light intervals," J Control Release **297**, 39–47 (2019).

6.   D. Luo, J. Geng, N. Li, K. A. Carter, S. Shao, G. E. Atilla-Gokcumen, and J. F. Lovell, "Vessel-Targeted Chemophototherapy with Cationic Porphyrin-Phospholipid Liposomes," Mol Cancer Ther (2017).

7. K. A. Carter, S. Shao, M. I. Hoopes, D. Luo, B. Ahsan, V. M. Grigoryants, W. Song, H. Huang, G. Zhang, R. K. Pandey, J. Geng, B. A. Pfeifer, C. P. Scholes, J. Ortega, M. Karttunen, and J. F. Lovell, "Porphyrin-phospholipid liposomes permeabilized by near-infrared light," Nat. Commun. **5**, (2014).

8. D. Luo, K. A. Carter, A. Razi, J. Geng, S. Shao, C. Lin, J. Ortega, and J. F. Lovell, "Porphyrin-phospholipid liposomes with tunable leakiness," J. Control. Release **220**, 484–494 (2015).

9. G. M. Palmer, R. J. Boruta, B. L. Viglianti, L. Lan, I. Spasojevic, and M. W. Dewhirst, "Non-invasive monitoring of intra-tumor drug concentration and therapeutic response using optical spectroscopy," J. Control. Release **142**(3), 457–464 (2010).

10. K. A. Carter, S. Shao, M. I. Hoopes, D. Luo, B. Ahsan, V. M. Grigoryants, W. Song, H. Huang, G. Zhang, R. K. Pandey, J. Geng, B. A. Pfeifer, C. P. Scholes, J. Ortega, M. Karttunen, and J. F. Lovell, "Porphyrin-phospholipid liposomes permeabilized by near-infrared light," Nat Commun **5**, 3546 (2014).

11. D. Luo, K. A. Carter, A. Razi, J. Geng, S. Shao, D. Giraldo, U. Sunar, J. Ortega, and J. F. Lovell, "Doxorubicin encapsulated in stealth liposomes conferred with light-triggered drug release," Biomaterials **75**, 193–202 (2016).

12. D. Luo, K. A. Carter, A. Razi, J. Geng, S. Shao, D. Giraldo, U. Sunar, J. Ortega, and J. F. Lovell, "Doxorubicin encapsulated in stealth liposomes conferred with light-triggered drug release," Biomaterials **75**, 193–202 (2016).

13. D. Luo, L. Nasi, K. A. Carter, C. Lin, J. Geng, S. Shao, W. C. Huang, Y. Qin, G. E. Atilla-Gokcumen, and J. F. Lovell, "Rapid Light-Triggered Drug Release in Liposomes Containing Small Amounts of Unsaturated and Porphyrin–Phospholipids," Small **12**(22), 3039–3047 (2016).

14. K. A. Carter, D. Luo, A. Razi, J. Geng, S. Shao, J. Ortega, and J. F. Lovell, "Sphingomyelin liposomes containing porphyrin-phospholipid for irinotecan chemophototherapy," Theranostics **6**(13), 2329–2336 (2016).

15. K. A. Carter, S. Wang, J. Geng, D. Luo, S. Shao, and J. F. Lovell, "Metal Chelation Modulates Phototherapeutic Properties of Mitoxantrone-Loaded Porphyrin-Phospholipid Liposomes," Mol. Pharm. **13**(2), 420–427 (2016).

16. S. Shao, J. Geng, H. Ah Yi, S. Gogia, S. Neelamegham, A. Jacobs, and J. F. Lovell, "Functionalization of cobalt porphyrin-phospholipid bilayers with his-tagged ligands and antigens," Nat Chem **7**(5), 438–446 (2015).

17. B. W. Pogue, T. Momma, H. C. Wu, and T. Hasan, "Transient absorption changes in vivo during photodynamic therapy with pulsed-laser light," Br J Cancer **80**(3–4), 344–351 (1999).

18. D. J. Rohrbach, D. Muffoletto, J. Huihui, R. Saager, K. Keymel, A. Paquette, J. Morgan, N. Zeitouni, and U. Sunar, "Preoperative Mapping of Nonmelanoma Skin Cancer Using Spatial Frequency Domain and Ultrasound Imaging," Acad. Radiol. **21**(2), (2014).

19. K. Sano, T. Nakajima, P. L. Choyke, and H. Kobayashi, "Markedly enhanced permeability and retention effects induced by photo-immunotherapy of tumors," ACS Nano **7**(1), 717–724 (2013).

20. A. L. R. de Souza, E. LaRochelle, K. Marra, J. Gunn, S. C. Davis, K. S. Samkoe, M. S. Chapman, E. V Maytin, T. Hasan, and B. W. Pogue, "Assessing daylight & low-light photodynamic therapy efficacy, using biomarkers of photophysical, biochemical and biological damage metrics in situ," Photodiagnosis Photodyn Ther (2017).

21.   U. Sunar, D. J. Rohrbach, J. Morgan, N. Zeitouni, and B. W. Henderson, "Quantification of PpIX concentration in basal cell carcinoma and squamous cell carcinoma models using spatial frequency domain imaging," Biomed. Opt. Express **4**(4), 531 (2013).

22.   N. C. Zeitouni, U. Sunar, D. J. Rohrbach, A. D. Paquette, D. A. Bellnier, Y. Shi, G. Wilding, T. H. Foster, and B. W. Henderson, "A prospective study of pain control by a 2-step irradiance schedule during topical photodynamic therapy of nonmelanoma skin cancer," Dermatologic Surg. **40**(12), (2014).

23.   S. Mallidi, S. Anbil, A. L. Bulin, G. Obaid, M. Ichikawa, and T. Hasan, "Beyond the Barriers of Light Penetration: Strategies, Perspectives and Possibilities for Photodynamic Therapy," Theranostics **6**(13), 2458–2487 (2016).

24.   A. Farhadi, A. Roxin, B. C. Wilson, and G. Zheng, "Nano-enabled SERS reporting photosensitizers," Theranostics **5**(5), 469–476 (2015).

25.   W. Mo, D. Rohrbach, and U. Sunar, " Imaging a photodynamic therapy photosensitizer in vivo with a time-gated fluorescence tomography system ," J. Biomed. Opt. **17**(7), 071306 (2012).

26.   M. S. Ozturk, D. Rohrbach, U. Sunar, and X. Intes, "Mesoscopic Fluorescence Tomography of a Photosensitizer (HPPH) 3D Biodistribution in Skin Cancer," Acad. Radiol. **21**(2), 271–280 (2014).

27.   H. Azais, C. Schmitt, M. Tardivel, O. Kerdraon, A. Stallivieri, C. Frochot, N. Betrouni, P. Collinet, and S. Mordon, "Assessment of the specificity of a new folate-targeted photosensitizer for peritoneal metastasis of epithelial ovarian cancer to enable intraperitoneal photodynamic therapy. A preclinical study," Photodiagnosis Photodyn Ther **13**, 130–138 (2016).

28.   B. W. Pogue, L. Lilge, M. S. Patterson, B. C. Wilson, and T. Hasan, "Absorbed photodynamic dose from pulsed versus continuous wave light examined with tissue-simulating dosimeters," Appl Opt **36**(28), 7257–7269 (1997).

29.   H. Azais, C. Schmitt, M. Tardivel, O. Kerdraon, A. Stallivieri, C. Frochot, N. Betrouni, P. Collinet, and S. Mordon, "Assessment of the specificity of a new folate-targeted photosensitizer for peritoneal metastasis of epithelial ovarian cancer to enable intraperitoneal photodynamic therapy. A preclinical study," Photodiagnosis Photodyn Ther **13**, 130–138 (2016).

30.   B. W. Pogue and T. Hasan, "A theoretical study of light fractionation and dose-rate effects in photodynamic therapy," Radiat Res **147**(5), 551–559 (1997).

31.   D. J. Rohrbach, N. Rigual, E. Tracy, A. Kowalczewski, K. L. Keymel, M. T. Cooper, W. Mo, H. Baumann, B. W. Henderson, and U. Sunar, "Interlesion differences in the local photodynamic therapy response of oral cavity lesions assessed by diffuse optical spectroscopies," Biomed. Opt. Express **3**(9), (2012).

32.   H. Azais, G. Queniat, C. Bonner, O. Kerdraon, M. Tardivel, G. Jetpisbayeva, C. Frochot, N. Betrouni, P. Collinet, and S. Mordon, "Fischer 344 Rat: A Preclinical Model for Epithelial Ovarian Cancer Folate-Targeted Therapy," Int J Gynecol Cancer **25**(7), 1194–1200 (2015).

33.   S. Wang, G. Huttmann, Z. Zhang, A. Vogel, R. Birngruber, S. Tangutoori, T. Hasan, and R. Rahmanzadeh, "Light-Controlled Delivery of Monoclonal Antibodies for Targeted Photoinactivation of Ki-67," Mol Pharm **12**(9), 3272–3281 (2015).

34.   A. Marrero, T. Becker, U. Sunar, J. Morgan, and D. Bellnier, "Aminolevulinic acid-photodynamic therapy combined with topically applied vascular disrupting agent vadimezan leads to enhanced antitumor responses," Photochem. Photobiol. **87**(4),

(2011). 400

35. D. J. Rohrbach, N. Rigual, H. Arshad, E. C. Tracy, M. T. Cooper, G. Shafirstein, G. Wilding, M. Merzianu, H. Baumann, B. W. 401
Henderson, and U. Sunar, "Intraoperative optical assessment of photodynamic therapy response of superficial oral squamous 402
cell carcinoma," J. Biomed. Opt. **21**(1), (2016). 403

36. H. Azaïs, G. Queniat, C. Bonner, O. Kerdraon, M. Tardivel, G. Jetpisbayeva, C. Frochot, N. Betrouni, P. Collinet, and S. 404
Mordon, "Fischer 344 Rat: A preclinical model for epithelial ovarian cancer folate-targeted therapy," Int. J. Gynecol. Cancer 405
25(7), 1194–1200 (2015). 406

37. E. H. Moriyama, A. Kim, A. Bogaards, L. Lilge, and B. Wilson, "A ratiometric fluorescence imaging system for surgical 407
guidance," Adv Opt Tech **532368**, 1–10 (2008). 408

38. A. Kim, M. Roy, F. Dadani, and B. C. Wilson, "A fiberoptic reflectance probe with multiple source-collector separations to 409
increase the dynamic range of derived tissue optical absorption and scattering coefficients," Opt. Express **18**(6), (2010). 410

39. G. A. Wagnieres, W. M. Star, and B. Wilson, "In vivo fluorescence spectroscopy and imaging for oncological applications," 411
Photochem Photobiol **68**(5), 603–632 (1998). 412

40. N. A. Wilson, U. S. RA, K. Lieberman, M. Akeson, and W. B. Dunbar, "Feedback control of a DNA molecule tethered in a 413
nanopore to repeatedly probe DNA-binding enzymes," Conf Proc IEEE Eng Med Biol Soc **2008**, 5745–5748 (2008). 414

41. A. Kim, M. Khurana, Y. Moriyama, and B. C. Wilson, "Quantification of in vivo fluorescence decoupled from the effects of 415
tissue optical properties using fiber-optic spectroscopy measurements," J. Biomed. Opt. **15**(6), 067006 (2010). 416

42. L. Lilge, K. Molpus, T. Hasan, and B. C. Wilson, "Light dosimetry for intraperitoneal photodynamic therapy in a murine 417
xenograft model of human epithelial ovarian carcinoma," Photochem Photobiol **68**(3), 281–288 (1998). 418

43. V. X. Yang, P. J. Muller, P. Herman, and B. C. Wilson, "A multispectral fluorescence imaging system: design and initial clinical 419
tests in intra-operative Photofrin-photodynamic therapy of brain tumors," Lasers Surg Med **32**(3), 224–232 (2003). 420

44. B. C. Wilson and M. S. Patterson, "The physics, biophysics and technology of photodynamic therapy," Phys Med Biol **53**(9), 421
R61-109 (2008). 422

45. J. K. Chan, B. J. Monk, D. Cuccia, H. Pham, S. Kimel, M. Gu, M. J. Hammer-Wilson, L. H. L. Liaw, K. Osann, P. J. DiSaia, M. 423
Berns, B. Tromberg, and Y. Tadir, "Laparoscopic photodynamic diagnosis of ovarian cancer using 5-aminolevulinic acid in a 424
rat model," Gynecol. Oncol. **87**(1), 64–70 (2002). 425

46. A. Bogaards, H. J. Sterenborg, J. Trachtenberg, B. C. Wilson, and L. Lilge, "In vivo quantification of fluorescent molecular 426
markers in real-time by ratio imaging for diagnostic screening and image-guided surgery," Lasers Surg Med **39**(7), 605–613 427
(2007). 428

47. J. Kress, D. J. Rohrbach, K. A. Carter, D. Luo, S. Shao, S. Lele, J. F. Lovell, and U. Sunar, "Quantitative imaging of light- 429
triggered doxorubicin release," Biomed. Opt. Express **6**(9), 3546 (2015). 430

48. U. Sunar, J. Kress, D. J. Rohrbach, K. A. Carter, D. Luo, S. Shao, S. Lele, and J. F. Lovell, "Light-triggered doxorubicin release 431
quantified by spatial frequency domain imaging and diffuse optical spectroscopy," in *Optics InfoBase Conference Papers* (2016). 432

49.   X. He, S. Zhou, W. C. Huang, A. Seffouh, M. T. Mabrouk, M. T. Morgan, J. Ortega, S. I. Abrams, and J. F. Lovell, "A Potent   433
        Cancer Vaccine Adjuvant System for Particleization of Short, Synthetic CD8+ T Cell Epitopes," ACS Nano **15**(3), 4357–4371   434
        (2021).   435

50.   K. Harris, D. J. Rohrbach, K. Attwood, J. Qiu, and U. Sunar, "Optical imaging of tissue obtained by transbronchial biopsies   436
        of peripheral lung lesions," J. Thorac. Dis. **9**(5), (2017).   437

51.   N. Rigual, G. Shafirstein, M. T. Cooper, H. Baumann, D. A. Bellnier, U. Sunar, E. C. Tracy, D. J. Rohrbach, G. Wilding, W.   438
        Tan, M. Sullivan, M. Merzianu, and B. W. Henderson, "Photodynamic therapy with 3-(10-hexyloxyethyl) pyropheophorbide   439
        a for cancer of the oral cavity," Clin. Cancer Res. **19**(23), (2013).   440

52.   U. Sunar, D. Rohrbach, N. Rigual, E. Tracy, K. Keymel, M. T. Cooper, H. Baumann, and B. H. Henderson, "Monitoring   441
        photobleaching and hemodynamic responses to HPPH-mediated photodynamic therapy of head and neck cancer: a case   442
        report," Opt. Express **18**(14), 14969 (2010).   443

53.   N. Rigual, G. Shafirstein, M. T. Cooper, H. Baumann, D. A. Bellnier, U. Sunar, E. C. Tracy, D. J. Rohrbach, G. Wilding, W.   444
        Tan, M. Sullivan, M. Merzianu, and B. W. Henderson, "Photodynamic therapy with 3-(10-hexyloxyethyl) pyropheophorbide   445
        a for cancer of the oral cavity," Clin. Cancer Res. **19**(23), 6605–6613 (2013).   446

                                                                                                                                          449