Mitochondrial Complex I and ROS control synapse function through opposing pre- and

postsynaptic mechanisms

Bhagaban Mallik[1], C. Andrew Frank[1,2]

[1]Department of Anatomy and Cell Biology, University of Iowa Carver College of Medicine, Iowa City,

IA, 52242, USA

[2]to whom correspondence should be addressed

andy-frank@uiowa.edu, @CAndrewFrank

SUMMARY

Neurons require high amounts energy, and mitochondria help to fulfill this requirement. Dysfunctional mitochondria trigger problems in various neuronal tasks. Using the *Drosophila* neuromuscular junction (NMJ) as a model synapse, we previously reported that Mitochondrial Complex I (MCI) subunits were required for maintaining NMJ function and growth. Here we report tissue-specific adaptations at the NMJ when MCI is depleted. In *Drosophila* motor neurons, MCI depletion causes profound cytological defects and increased mitochondrial reactive oxygen species (ROS). But instead of diminishing synapse function, neuronal ROS triggers a homeostatic signaling process that maintains normal NMJ excitation. We identify molecules mediating this compensatory response. MCI depletion in muscles also enhances local ROS. But high levels of muscle ROS cause destructive responses: synapse degeneration, mitochondrial fragmentation, and impaired neurotransmission. In humans, mutations affecting MCI subunits cause severe neurological and neuromuscular diseases. The tissue-level effects that we describe in the *Drosophila* system are potentially relevant to forms of mitochondrial pathogenesis.

Keywords: homeostatic plasticity; mitochondria; Mitochondrial Complex I; *Drosophila*; *ND-20L*; Mito-GFP; rotenone; NACA; *sod2*; ROS

INTRODUCTION

Neurons have vast energy needs. These needs are primarily satisfied by healthy pools of mito-chondria [1,2]. Mitochondria generate energy through the action of the ATP synthase complex in the electron transport chain [3,4]. They also perform complementary functions, including maintaining cal-cium homeostasis [5,6], promoting cell survival [7], triggering reactive oxygen species (ROS) signaling [8], stimulating lipid synthesis [9], and regulating innate immunity [10]. For energy-driven neurons, it is thought that the primary role of mitochondria is to provide ATP. It is less understood how other mi-tochondrial functions contribute to the regulation of normal neurophysiology. It is also not well un-derstood how neural tissues or synaptic sites cope when they are challenged with a loss of mito-chondria. Genetic models can help to address these puzzles.

Mitochondrial Complex I (MCI) (NADH ubiquinone oxidoreductase) is an essential part of the electron transport chain and ATP production. MCI consists of 42 distinct subunits. Much of our un-derstanding about MCI derives from systemic analyses of its assembly. Studies have been per-formed on Complex I components from diverse organisms, including *Neurospora crassa* and *Dro-sophila melanogaster*. Those studies demonstrate that discrete MCI subunits are ancient; indeed, there are few differences between these MCI models from simple organisms and the corresponding human and bovine orthologs [11-14]. For *Drosophila melanogaster*, 13/14 of the core MCI subunits are present, as are the remaining 28 accessory subunits [12].

In humans, MCI dysfunction has been linked to diseases such as Leigh syndrome, mito-chondrial myopathy, and encephalomyopathy, as well as forms of stroke [15-18]. On a cellular level, MCI dysfunction can cause the demise of neurons and muscles; these phenotypes are typically at-tributed to defects in ATP production [19,20]. However, in addition to the ATP production defects, mu-tations affecting MCI subunit components are also associated with excess mitochondrial ROS. Normally, ROS accumulation can be neutralized by the cellular antioxidant system [21]. But if that sys-tem becomes overwhelmed, there can be consequences for cells and organ systems – including progressive neurodegeneration and seizures for the nervous system [22-24]. On the level of synapses, it is possible that MCI loss triggers severe molecular consequences, and it is also possible that ex-cess ROS plays a role.

In a previous study, we depleted MCI function at the *Drosophila* neuromuscular junction (NMJ). Our data suggested fundamental synaptic functions for MCI [13]. Here we expand upon that work, mostly taking advantage of RNAi-mediated depletion of the nuclear DNA-encoded *NADH de-hydrogenase subunit 20* (*ND-20L*) gene, a homolog of human NDUFS7. We also scrutinize loss-of-function mutants of other MCI subunits and pharmacological inhibition of MCI. Our collective data

show that MCI depletion causes *Drosophila* phenotypes reminiscent of mitochondrial diseases, such as progressive degeneration of muscle and presynaptic cytoskeleton, excess ROS production, loss of mitochondria and alteration in mitochondrial morphology.

On single-tissue levels, we were surprised to find that there were opposite effects on synapse activity in the presynaptic motor neurons vs. the postsynaptic muscles. MCI dysfunction in *Drosophila* motor neurons causes profound cytological phenotypes, but there are no significant functional phenotypes. This appears to be because neuronal mitochondrial ROS triggers an adaptive response, demonstrated visually by active zone enhancement. Based on our data, this ROS-driven enhancement of active zones occurs through at least two processes, 1) regulation of calcium flux from intracellular stores (ER) and mitochondria; and 2) use of glycolysis as an alternative energy source. By contrast, postsynaptic depletion of MCI and the associated elevation of muscle ROS triggers a destructive response: disruption of NMJ morphology and the Dlg-Spectrin scaffold that would normally be critical for normal active zone-receptor apposition. To our knowledge, these cellular and molecular mechanisms of MCI deficiency have not previously been elucidated at synapse-specific or tissue-specific levels.

RESULTS

Depletion of MCI affects mitochondrial integrity in multiple *Drosophila* synaptic tissues

In prior work, we reported impairments in neuromuscular junction (NMJ) synapse development and function when Mitochondrial Complex I (MCI) is depleted [13]. To understand those results better, we examined mitochondria by microscopy. We used the GAL4/UAS system to express a *UAS-Mito-GFP* transgene [25] in *Drosophila* tissues. Concurrently, we used tissue-specific GAL4 drivers alone (as controls) or GAL4 drivers + *UAS-ND-20L[RNAi]* to deplete *ND-20L* gene function by RNA interference, as previously published [13]. With these tools, we made qualitative observations of mitochondrial morphology (Fig. 1), and then we quantified those observations in subsequent analyses.

In motor neurons, the Mito-GFP signal localized to the neuropil of the ventral nerve cord (Fig.1: A-A′, B-B′). In control neurons, the neuropil mitochondria had a filamentous appearance. By contrast, *ND-20L*-depleted neurons had punctate and clustered mitochondria. (Fig. 1E-E′, F-F′). We examined mitochondria in the motor axons that innervate proximal and distal NMJs (Fig. 1C-C′, D-D′, G-G′, H-H'). The proximal segment A2 axons had abundant mitochondria in all cases (Fig. 1C-C', G-G'). However, for the distal segment A5 axons, *ND-20L* depletion elicited an obvious decrease in mitochondria number (Fig. 1D-D,′ H-H,′). This A2 vs. A5 discrepancy was consistent with

prior work by others examining defects in mitochondrial trafficking dynamics: distal sites can show
phenotypes more prominently [26,27].

We hypothesized that fewer mitochondria in the A5 axon might correlate with a neurotrans-
mission defect at the NMJ. Yet by NMJ electrophysiology, we found no significant differences in the
evoked amplitude compared to the control NMJs in the distal segment A5 (Fig. 1I-K). These data
matched our prior examination of MCI at the A2 and A3 segments of the NMJ, where neuronal im-
pairment of MCI was not sufficient on its own to reduce evoked NMJ neurotransmission [13].

In muscle, we observed an array of mitochondrial phenotypes. As with neurons, there were
clustered mitochondria when *ND-20L* gene function was depleted (Fig. 1L-M). Additionally, there
was a tissue-level phenotype: *ND-20L*-depleted muscles were developed, but they looked disor-
ganized and fragmented, with oblong-shaped nuclei in the muscle syncytia (Fig. 1L-M). This pheno-
type could explain why we previously observed that muscle impairment of MCI was sufficient to re-
duce evoked NMJ neurotransmission [13].

To examine the mitochondria at presynaptic NMJ release sites, we used the motor neuron
GAL4 driver to label NMJ boutons with Mito-GFP. For image analysis, we marked the presynaptic
membrane boutons with anti-HRP immunostaining. Control NMJs contained abundant and large
clusters of mitochondria in synaptic boutons, but by comparison, *ND-20L[RNAi]* boutons contained
small clusters and few mitochondria (Fig. 1N-P). We measured the mitochondrial volume in a 3-D
stack and compared it to the synaptic volume (Fig. 1Q-S). The Mito-GFP signal occupied a sizeable
proportion of the bouton volume in controls (~30%), but this value was significantly diminished in
ND-20L-depleted animals (~10%) (Fig. 1S). Collectively, our data suggest that the depletion of *ND-*
20L by RNAi leads to abnormal mitochondrial clustering in the neuronal cell body and muscle – as
well as losses of distal axon and synaptic mitochondria.

Loss of MCI phenocopies loss of Mitofusin

The cell-level ND-20L-depletion phenotypes were reminiscent of *Drosophila* mutants impairing mi-
tochondrial dynamics [26,28]. Therefore, we re-examined MCI-depleted mitochondria, this time addi-
tionally impairing genes known to mediate mitochondrial fusion and fission. Mitofusin 1 (Mfn1) and
Mitofusin 2 (Mfn2) are GTPases that regulate outer mitochondrial membrane fusion [29,30]. The *Dro-*
sophila gene encoding the Mitofusin homolog is called *marf*. Dynamin-related protein 1 is a GTPase
that regulates mitochondrial fission. In *Drosophila*, this factor is encoded by the gene *drp1* [31]. Previ-
ous work reported that defective fusion results in fragmented mitochondria, while defective fission
can lead to enlarged mitochondria [32]. We used RNAi-mediated knockdown constructs for each of
these genes.

As before, we observed that wild-type motor neurons had filamentous and oval mitochondria, while *ND-20L* depleted neurons had fewer and smaller clustered mitochondria in the ventral nerve cord (VNC) (Fig. 2A) and axons (Fig. 2B). Knockdown of the fusion gene *marf* phenocopied *ND-20L* loss, revealing small mitochondria in motor neurons, while knockdown of the fission gene *drp1* yielded filamentous mitochondria (Fig. 2A). Simultaneously depleting motor neurons of *marf* and *ND-20L* did not show any additive defect in mitochondrial appearance in the ventral nerve cord (VNC) and axons (Figs. 2A, B). This result could mean that the two genes share a common process to regulate mitochondrial fusion. By contrast, depleting *drp1* and *ND-20L* simultaneously yielded punctate mitochondria. This result likely means that that the punctate *ND-20L* mitochondrial phenotypes (potential fusion phenotypes) are epistatic to *drp1* loss (Fig. 2A, B). We measured mitochondrial branch length from skeletonized images of the mitochondria (Skeletonize3D, ImageJ plugin). Control and *drp1* knockdown showed normal mitochondrial branch length, but knockdown of *ND-20L* or *marf* – or knockdowns using combinations of each – exhibited short branch length (Fig. 2C, Table 1).

To quantify mitochondria in axons, we counted Mito-GFP positive puncta in distal A5 motor axons labeled by anti-GFP. Control axons and *drp1*-depleted axons contained abundant mitochondria (Fig. 2D, Table 1). By contrast, any gene manipulation or combination targeting *ND-20L* or *marf* resulted in diminished numbers of mitochondrial clusters (Fig. 2D, Table 1).

We extended the analysis to NMJ terminals. We counted Mito-GFP clusters in presynaptic boutons apposed by postsynaptic densities, labeled by anti-Discs Large 1 (Dlg1) (Figs. 2E-M). The results matched our earlier observations (Figs. 1N-S). Control NMJs and *drp1*-depleted NMJs contained numerous mitochondrial clusters per bouton (Figs. 2E, M). However, *ND-20L*-depleted boutons contained few mitochondria, and this was phenocopied by *marf*[RNAi] (Fig. 2E-M, Table 2). Collectively, these results suggest that *Drosophila ND-20L* (and hence MCI) contributes to normal mitochondrial fusion, likely in conjunction with the Mitofusin homolog, Marf.

Mitochondrial Reactive Oxygen Species Contribute to Synaptic Phenotypes
Several studies have demonstrated that Complex I loss results in high levels of mitochondrial reactive oxygen species (ROS) [33-37]. This means that excess ROS could be contributing to the cytological and mitochondrial fusion phenotypes that we have described.

We checked if we could observe mitochondrial ROS (superoxide) in living *Drosophila* tissue and if ROS levels corresponded to Complex I function (Fig. S1). We used a commercially available fluorescent mitochondrial superoxide indicator, MitoSOX (MitoSOX[TM], trademarked by ThermoFisher, Materials and Methods) [38-40]. With MitoSOX, we observed mitochondrial superoxide

in many tissues. There was a baseline level of ROS in controls (Fig. S1A, E, I, J, Table 3), and the

level was greatly increased in *ND-20L*-deficient motor neuron cell bodies and muscle (Fig. S1B, F,

I; J, Table 3).

Next, we tested if ROS scavengers could reverse the high Mito-SOX fluorescence levels in

ND-20L-deficient tissues. We fed a pharmacological scavenger, N-Acetyl Cysteine Amide (NACA)

[41-43] to *Drosophila* larvae (Materials and Methods). We also used a transgene, *UAS-sod2* [44], to ex-

press a superoxide dismutase enzyme. Both successfully diminished the high levels of mitochon-

drial ROS that resulted from *ND-20L* depletion at the NMJ, and both worked in muscle and neurons

(Fig. S1C, D, G-J, Table 3).

We wondered if ROS scavengers could reverse mitochondrial phenotypes caused by loss of

MCI. We considered *UAS-ND-20L[RNAi]* knockdown phenotypes. Co-expressing *UAS-sod2* or

rearing larvae with NACA suppressed the mitochondrial morphology defects in the ventral nerve

cord of *ND-20L*-depleted animals; it also restored axonal loss of mitochondria (Fig S2A-D, Table 4).

To test an additional MCI manipulation, we knocked down *Drosophila ND-30* (homologous to hu-

man *NDUFS3*) in motor neurons. As with *ND-20L*, depleting *ND-30* in motor neurons yielded punc-

tate mitochondria in the ventral nerve cord, but the addition of *UAS-sod2* restored a wild-type, fila-

mentous mitochondrial morphology (Fig. S2A, C). Similarly, loss of *ND-30* gene function in neurons

depleted A5 axonal mitochondria and this phenotype was also reversed by *UAS-sod2* transgenic

expression (Fig. S2B, D, Table 4).

Because of the links between MCI and mitochondrial fusion, we considered whether a *marf*

loss of function could also yield high levels of neuronal ROS (Fig. S3). It did – both *marf* and *ND-*

20L loss-of-function conditions showed high levels of mitochondrial superoxide in motor neuron cell

bodies (Fig. S3A-D, A'-D', M, Table 3); in motor axons (Fig. S3E-H, E'-H', N, Table 3); and at NMJ

sites (Fig. S3I-L, I'-L', O, Table 3). These ROS phenotypes were not confined to genetic manipula-

tions. We made similar observations when MCI was impaired pharmacologically by feeding rote-

none to developing larvae (Fig. S3, Table 3). In the case of rotenone, the amount of mitochondrial

ROS in the boutons was high, but it was not increased as much as with the genetic manipulations

(Fig. S3K, O, Table 3).

Finally, because the distal A5 motor axons accumulated high levels of ROS when subjected

to these insults, we examined them for synaptic vesicle trafficking defects. We immunostained for

Cysteine String Protein (CSP), a DNAJ-like co-chaperone and synaptic vesicle-associated protein

(Cysteine string protein: CSP). *ND-20L* depletion caused aberrant accumulation of CSP in the A5

motor axons; and this defect was suppressed upon motor neuron transgene expression of *UAS-*

sod2 or feeding animals with NACA (Fig. S4A-L, Table 5).

ROS scavengers did not reverse all mitochondrial abnormalities. Expressing *UAS-sod2* in

the *UAS-ND-20L[RNAi]* or *UAS-marf[RNAi]*-depletion backgrounds did not restore mitochondrial

clusters to motor neuron terminals (Fig. 2E-M, Table 2). For the remainder of the study, we used

scavengers as complementary tools to test which MCI-loss phenotypes were likely due to mito-

chondrial ROS.

Loss of MCI subunits impairs synaptic cytoskeletal stability

ROS can modulate the cytoskeleton, either through redox modification of cytoskeletal proteins or by

altering pathways that regulate cytoskeletal organization [45]. To test whether the mitochondrial de-

fects and abnormal accumulation of ROS were associated with the altered synaptic cytoskeleton,

we labeled synaptic boutons with an anti-Futsch antibody (Fig. 3). Futsch is a *Drosophila* MAP1B

homolog that associates with microtubules [46].

In motor neuron Gal4-control and *UAS-sod2* overexpression larvae, Futsch organized in pe-

riodic loops, as expected from previous characterizations [46] (Fig.3A-B, Table 5). But in *ND-20L* de-

pleted larvae, the anti-Futsch staining showed a significant reduction in microtubule loops (Figure

3C, Table 5). These *ND-20L* phenotypes were suppressed by motor neuron expression of *UAS-*

sod2 or by raising animals on food containing 0.5 mM NACA (Fig. 3D-E, Table 5). These data indi-

cate that loss of MCI regulates cytoskeletal architecture due to excessive accumulation of ROS in

neurons.

Modest neurotransmission phenotypes after motor neuron-specific loss of MCI or marf

Given the cytological phenotypes after neuronal MCI loss, it was puzzling that there seemed to be

little to no electrophysiological consequence at NMJs ([13] and Fig. 1J). We probed this finding, this

time depleting motor neurons of *marf* and/or *ND-20L* gene function. We recorded spontaneous min-

iature postsynaptic potentials (mEPSP) and evoked excitatory postsynaptic potentials (EPSP).

Phenotypes were normal-to-mild (Fig. S5). For both *ND-20L[RNAi]* and *marf[RNAi]*, there

were small, but statistically significant decreases in mEPSP amplitude (Fig. S5A-C, G, Table 6). But

for *ND-20L[RNAi]*, evoked events (EPSP) and calculated quantal content (QC) were at control lev-

els (Fig. S5A-B, H-I, Table 6). For neuronal *marf[RNAi]*, those measures were near-normal, with a

slight decrease in EPSP amplitude (Fig. S5C, H, Table 6) and a slight increase in QC (Fig. S5I, Ta-

ble 6). Finally, for a double *ND-20L[RNAi]* + *marf[RNAi]* knockdown condition in neurons, we ob-

served a small, but statistically significant decrease in EPSP amplitude.

Synaptic phenotypes caused by mitochondrial dysfunction might be masked until synapses

are challenged with extreme conditions, like high frequency stimulation [47,48]. Therefore, we chal-

lenged neuronally *ND-20L*-depleted *Drosophila* NMJs in several ways (Fig. S6). First, we lowered

recording saline [Ca^{2+}] to 0.15 mM, which is roughly one order of magnitude lower than physiologi-

cal calcium. In low calcium, the motor neuron-driven *ND-20L[RNAi]* NMJs had slightly smaller

evoked potentials compared to driver controls, but the numerical reduction was not statistically sig-

nificant (Fig. S6A-C, Table 7). Next, we lowered extracellular [Ca^{2+}] even further, to 0.1 mM, which

yielded a mix of successful EPSP firing events and failures. Failure analyses revealed an increase

in failure rate at the neuronally depleted *ND-20L[RNAi]* NMJs compared to control animals (Fig

S6D, Table 7), demonstrating a decreased probability of release. Failure rates in 0.1 mM calcium

were restored to baseline levels when *ND-20L*-depleted larvae were raised in a media containing

NACA or genetically expressing *sod2* in motor neurons (Fig S6D, Table 7), indicating that a sensi-

tivity to low calcium could be related to ROS levels.

Finally, we checked if forms of short-term neuroplasticity were affected by *ND-20L* loss in

motor neurons. For two different extracellular [Ca^{2+}] conditions (0.4 to 1.5 mM), we did not observe

any significant changes in paired-pulse ratios (Figs. S6E-J, Table 7). Likewise, we did not see any

notable depreciation of evoked neurotransmission over the course of high frequency stimulus trains

in high calcium (Figs. S6K-P, Table 7). Collectively, these data suggest that there might be a small

effect on NMJ physiology due to defective mitochondrial fusion, and the defect could be sensitive to

low levels of extracellular calcium – but the aggregate data also indicate that neuronal mitochondrial

defects alone do not drastically affect NMJ neurotransmission.

Loss of MCI in neurons controls the level and distribution of the active zone to stabilize syn-

aptic strength

We wondered how synapse might evade severe dysfunction, despite loss of mitochondria in the

motor neurons. One possibility is that *ND-20L* loss/MCI impairment could trigger a form of functional

homeostatic compensation of the NMJ. Another idea is that the mitochondrial ATP generated is su-

perfluous at the NMJ – and that any energy-intensive functions that mitochondria support could be

redundantly covered by glycolysis. These models are not mutually exclusive, and for any scenario,

mitochondrial ROS downstream of defective MCI could be a candidate signal. Recent findings have

demonstrated that ROS intermediates, mitochondrial distribution, and mitochondrial trafficking all

affect development of the *Drosophila* NMJ [45,49,50].

We imaged the presynaptic active zone apparatus in neuronally-depleted *ND-20L[RNAi]*

flies. Third-instar larval active zones showed a decrease in BRP puncta density per unit area in *ND-*

20L-depeleted NMJs compared to control NMJs (Fig.4A-C, G, H-I, Table 8). But they also showed a robust enhancement phenotype: in *ND-20L*-depleted animals, we found a 40% increase in active zone (BRP) immunofluorescence signal per unit area, compared to control (Fig. 4 A-C, F, H-I, Table 8), by laser scanning confocal microscopy. This result was intriguing because NMJ active zone enhancements (or changes in active zone sub-structure) have been proposed by other labs to be molecular correlates of forms of homeostatic plasticity and potentiation of neurotransmitter release [51-55]. This raised the possibility that the NMJs evade severe dysfunction through a form of synaptic homeostasis.

As an independent test, we impaired MCI pharmacologically. To do this, we raised larvae on rotenone-spiked food; and we also incubated wild-type fillet preparations with 500 µM of rotenone for extended time. For both cases, we observed significant increases in BRP protein at the presynaptic active zones (Fig S7A-U, Table 9). For the extended incubation, the fillet preparations required sufficient rotenone incubation time (six hours) and an intact motor nerve to show the active zone enhancement (Fig. S7U, Table 9). This latter result suggested that delivery of active zone material required either trafficking time and/or fully intact neuroanatomy.

Next, we checked if the enhanced active zone signal by was triggered by excess mitochondrial ROS in motor neurons. Indeed, we found that the *ND-20L*-depletion active zone enhancements were fully reversed by ROS scavengers, either by raising larvae in food containing NACA or by neuronally expressing *UAS-sod2* (Fig. 4 D-G, Table 8).

Finally, we assessed synapse function. As with our prior recordings, evoked postsynaptic potentials at the NMJ were not significantly changed by *ND-20L* depletion in motor neurons. But interestingly, scavenging mitochondrial ROS in the *ND-20L[RNAi]* neuronal depletion background with *UAS-sod2* revealed a small decrease in NMJ excitation, compared to controls (Fig. 4J-P, Table 8). This could mean that mitochondrial ROS is helping to maintain synaptic activity. Consistently, neuronal expression of *UAS-sod2* did not restore mitochondrial clusters to the NMJ after *ND-20L* gene function depletion (Fig. 4Q-U, Table 8; like Fig. 2H), meaning that the synaptic sites were still deficient in mitochondria. Our data support a model in which neuronal ROS (nROS) triggers active zone enhancement functional compensation when MCI is limiting (Fig. 4V-X, Table 8).

Neuronal MCI subunits stabilize synaptic strength in conjunction with intracellular calcium signaling proteins

Recent work described a mechanism for local calcium uptake into mitochondria that drives ATP production to maintain synaptic function [56]. The process is governed by the mitochondrial calcium

uniporter (MCU) and its accessory EF-hand MICU proteins [56]. Beyond this role for mitochondrial calcium, there are also known roles for core synaptic functions like vesicle cycling [57,58].

To test if mitochondrial or neuronal calcium could be involved in maintaining synapse function at the NMJ, we acquired genetic reagents to examine depressed MCU function in conjunction with depressed MCI. We also used pharmacological reagents to inhibit release of intracellular sources of calcium, like those from the Ryanodine Receptor (RyR) and the IP_3 receptor (IP_3R) of the endoplasmic reticulum (ER) [59], as well as a genetic reagent that we previously used at the NMJ to deplete IP_3 signaling (*UAS-IP_3-sponge*) (Fig. 5A-D). For this set of experiments, we used the *ND-20L* neuronal knockdown condition as a sensitized genetic background (Fig. 5E-T, Table 10).

We observed no significant differences in EPSP amplitudes when we impaired *mcu* function (Table 10). Similarly, we did not observe deficits in baseline synaptic activity by blocking RyR and IP_3R, alone or in conjunction with *ND-20L*[RNAi] (Fig. 5E-H, Q-T, Table 10). However, when we concurrently impaired a combination of *ND-20L*, *mcu*, and ER calcium store channels, we observed marked decreases in evoked amplitude (Fig. 5K-L, S, Table 10). These results are consistent with a model in which mitochondrial Ca^{2+} uptake, MCU activation, and ER (store) Ca^{2+} efflux combine to stabilize synaptic strength.

If this idea were correct, then it should also be possible to chelate cytoplasmic calcium in a neuronal *ND-20L[RNAi]* background and reveal neurotransmission defects. Direct application of the membrane-permeable chelator BAPTA-AM, followed by a wash to remove chelator residing in the saline, had no significant effect on baseline neurotransmission parameters (DMSO carrier + wash). But in the *ND-20L[RNAi]* background, BAPTA-AM + wash significantly diminished evoked potentials, compared to mock-treated (DMSO + wash) NMJs. (Fig.5 M-T, Table 10).

To check if these effects on neurotransmission correlated with effects on active zone protein accumulation, we also conducted anti-Brp immunostaining experiments (Fig. S8). As before, neuronal knockdown of *ND-20L* gene function triggered a marked, compensatory increase in presynaptic active zone material that was readily apparent by confocal microscopy (Figs. S8A, C, I, Table 11). But this increase was reversed when combined with *mcu* gene function knockdown and pharmacological blockade of store calcium release channels (Figs. S8G-I, Table 11). Together, our data indicate that loss of MCI subunits in neurons sensitizes synapses to decreases in intracellular calcium.

A combination of mitochondria, glycolysis, and the TCA Cycle stabilizes NMJ function

Neuronal calcium handling and MCU play roles in NMJ stability that are uncovered by loss of MCI at the NMJ. Downstream of calcium handling, a logical hypothesis is that synaptic energy would

play a role [56]. If this were the case, would mitochondria be the sole energy source? Alternatively, in
the absence of full mitochondrial function, could glycolysis theoretically substitute, as a homeostatic
(or redundant) means for staving off synapse dysfunction?

We tested these ideas by limiting glycolysis as an energy source in two ways: 1) swapping
out sucrose and trehalose in our recording saline in favor of 2-deoxy-D-glucose (a non-glycolytic
sugar; (Fig. 6A); 2) addition of lonidamine (LDA) to the saline to acutely inhibit hexokinase (Fig. 6A).
Control recordings with these conditions showed little effect on baseline physiology (Fig. 6A, D, F,
H-J, Table 12). However, when Mitochondrial Complex I was impaired neuronally through *ND-*
20L[RNAi] in combination with inhibition of glycolysis, there was a drop in evoked neurotransmis-
sion (Fig. 6C, E, G, H-J, Table 12). This correlated with a failure to increase active zone material
after *ND-20L* gene knockdown (Fig. 6K-Q, Table 12). These results match the idea that a combina-
tion of mitochondrial function or glycolysis can work to maintain normal levels of NMJ output.

We continued this line of investigation genetically. We acquired RNA interference-based
transgenes to target five genes involved in *Drosophila* glycolysis or subsequent ATP generation in
the Citric Acid (TCA) Cycle: *hexokinase A (hex-A)*, *hexokinase C (hex-C)*, *Citrate (Si) Synthase I*,
Isocitrate dehydrogenase (*Idh*), and *Succinyl-coenzyme A synthetase α subunit 1* (*Scsα1*). We
knocked down these genes neuronally, either alone or in combination with *ND-20L[RNAi]* (Figs. S9
and S10). Neuronal impairment of *hex-C* had no effect on baseline neurotransmission, but *hex-A*
impairment reduced it (Figs. S9A-E, Table 13). Impairment of the TCA Cycle enzymes on their own
had little-to-no effect on baseline neurotransmission (Fig. S9A, F-H, Table 13). However, concurrent
impairment of *ND-20L* and most of these genes significantly blunted neurotransmission, with the
exception being *hex-C* (Figs. S9I-N, Table 13). Collectively, the data suggest that MCI works in con-
junction with – or redundantly to – alternative energy-generating pathways to support normal levels
of neurotransmission. In the case of the hexokinases, *Drosophila hex-A* seems more important for
this process than *hex-C*.

We also assayed active zone material accumulation. The results mirrored the neurotrans-
mission tests: as before, neuronal *ND-20L[RNAi]* impairment elicited enhanced active zone material
(Fig. S10, Table 14). But concurrent neuronal impairment of *ND-20L* and any of the glycolysis or
TCA Cycle genes reversed this active zone enhancement (Fig. S10, Table 14).

MCI subunits in muscle are required for proper synapse development

We previously reported that impairments of MCI diminish *Drosophila* NMJ growth [13]. The roles of
MCI in specific tissues for this developmental function were unclear. For the present study, we test-
ed for tissue-specific roles of MCI in NMJ development. To visualize NMJ boutons, we co-stained

larval fillets with anti-Horseradish Peroxidase (HRP) a presynaptic membrane marker, and anti-
Discs Large (Dlg), a postsynaptic density marker [60,61].

On a coarse level, MCI loss in muscle (*BG57-Gal4 > UAS-ND-20L[RNAi]*) caused a severe
reduction in average bouton size, a decrease in bouton number, a notable decrease in Dlg expres-
sion, and a bouton "clustering" phenotype (Fig. 7 A-B), reminiscent of what we previously reported
[13]. To quantify these observations, we measured bouton number, muscle area, and branch number
per muscle in the third-instar larval NMJ synapses (Table 15). We found that *ND-20L* muscle
knockdown resulted in a significant reduction in all these parameters compared to controls (Fig 7J-
L, U-V, Table 15).

In contrast to muscle knockdown, pan-neuronal or motor neuron-specific knockdown of *ND-*
20L showed slight NMJ overgrowth phenotypes (Fig. 7F-L, Table 15). As was the case with the ac-
tive zone enhancement associated with neuronal *ND-20L* knockdown (Fig. 4), this NMJ overgrowth
could represent a developmental mechanism to stave off dysfunction caused by missing neuronal
mitochondria. Taken together, our NMJ immunostaining results indicated to us that blunted NMJ
growth due to MCI loss was likely due to muscle MCI dysfunction.

We tested if the NMJ undergrowth phenotypes could be due to the increased levels of mito-
chondrial reactive oxygen species in muscle (mROS) (Fig. S1F). If that idea were correct, then the
undergrowth phenotypes should be reversed if mitochondrial mROS were scavenged. Consistent
with this idea, bouton number and synaptic undergrowth phenotypes were fully restored to wild-type
levels when the *ND-20L[RNAi]* muscle knockdown animals also had *UAS-sod2* transgenically ex-
pressed in the muscles (Fig. 7C, J-L, R, Table 15). They were also restored to wild-type levels
when muscle *ND-20L* knockdown animals were raised on food containing the antioxidant N-acetyl
cysteine amide (NACA) (Fig. 7J-L, Table 15). By contrast, none of these NMJ growth parameters
were restored to wild-type levels if scavengers *UAS-sod1* or *UAS-catalase* were misexpressed in
the muscle (Fig. 7D-E, J-L, Table 15). SOD1 (cytosol) and Catalase (peroxisomes) localize to dif-
ferent compartments than SOD2 (mitochondrial matrix). These results likely indicates that a scav-
enger needs to access the proper mitochondrial compartment for rescue.

Loss of MCI and mROS in muscle disorganize NMJ postsynaptic densities
The muscle Dlg-Spectrin network functions as an organizing scaffold for synaptic assembly [61,62].
Dysregulation of this network can lead to an aberrant muscle subsynaptic reticulum (SSR) [61].
Therefore, one possible target of excess mROS in the absence of MCI function is Dlg. Dlg is the fly
homologue of PSD-95/SAP97/PSD-93, and it is a member of the membrane-associated guanylate

kinase (MAGUK) family of NMJ scaffolding proteins [61]. It is present both within presynaptic boutons

and in the portion of the SSR closest to the bouton.

Using the same antibodies detailed above (anti-Dlg and anti-HRP), we used a high magnifi-

cation to examine at the postsynaptic densities closely. By confocal immunofluorescence, Dlg area

was significantly reduced when *ND-20L* was depleted postsynaptically by[RNAi] (Fig. 7M-U, Table

15). To quantify the relative Dlg area, we measured Dlg area with respect to HRP (Relative Dlg ar-

ea = Dlg area minus HRP area) in type 1b boutons at muscle 6/7 of the A2 hemisegment (Fig. 7U).

Compared with the control synapses, *ND-20L* knockdown resulted in a significant reduction in the

relative Dlg area (Fig 7M-U, Table 15). Consistently, the relative α-Spectrin area was also reduced

when *ND-20L* was depleted in muscle (Fig. S11A-F, Table 5).

Next, we scavenged mROS, and then we examined the postsynaptic densities. The relative

Dlg and α-Spectrin areas were restored to wild-type levels when animals were grown in a media

containing NACA (*ND-20L[RNAi]/+; BG57-Gal4/+* with NACA) or genetically expressing *UAS-sod2*

in the muscle (*UAS-sod2/ND-20L[RNAi]; BG57-Gal4/BG57-Gal4*) (Fig.7R, U, Table 15; Fig. S11D-

F, Table 5). By contrast, Dlg levels were not restored while expressing other scavengers in the

muscle, encoded by *UAS-sod1* and *UAS-catalase* transgenes (Fig. 7S-U, Table 15). We conclude

that depletion of MCI subunits in the muscle disables postsynaptic density formation via the for-

mation of mitochondrial reactive oxygen species intermediates.

The postsynaptic density (PSD-95/Dlg) has been shown to cluster glutamate receptors at

the SSR [63]. Our results raised the possibility that glutamate receptor clusters could be disrupted

when *ND-20L* was depleted in the muscle (Fig. 8). To test this idea, we simultaneously

immunostained NMJs with antibodies against Brp (neuron, presynaptic active zone) and glutamate

receptor clusters (muscle). In controls, these pre- and postsynaptic structures were directly ap-

posed to one another (Fig. 8A, G). But when *ND-20L* gene function was depleted, we observed

"missing" GluRIIA and GluRIII receptor clusters (Fig. 8 B, F, H, K, Table 5), i.e., Brp puncta without

apposed glutamate receptors. These lack of apposition phenotypes were fully reversed by raising

the larvae with NACA or genetically expressing *UAS-sod2* in the muscle (Fig. 8 D-F, I-K, Table 5).

Together, our data indicate that loss of MCI subunit in the muscle (*ND-20L*) disrupts several as-

pects of the postsynaptic density organization, and these disruptions are likely due to the accumula-

tion of mROS in the muscle.

Loss of MCI subunits in muscle diminishes evoked NMJ neurotransmission through

postsynaptic mROS

Mitochondria and reactive oxygen species influence presynaptic vesicle release and plasticity at
synapses. This has been shown in diverse model systems like flies, mice, and worms [49,50,64-66]. In
our prior work, we identified an NMJ neurotransmission defect when MCI function is impaired [13], but
based on our current study, that defect does not seem like it was dependent upon MCI's neuronal
functions (Figs. 1, 4, 5, 6). Therefore, we turned to analyzing postsynaptic muscle MCI and mROS
to check if these parameters influenced neurotransmission.

We performed sharp electrode electrophysiological recordings of miniature and evoked ex-
citatory postsynaptic potentials (mEPSP and EPSP) at NMJ muscle 6, hemisegment A2. We also
used average mEPSP and EPSP values to estimate quantal content (QC) for each NMJ. For the
most part, mEPSP values remained steady (Figs. 9A-E, Table 16), with some exceptions. The
starkest phenotypes came in terms of evoked amplitudes (Figs. 9A-D, F), sometimes due to chang-
es in QC (Fig. 9G) or combinatorial changes in both mEPSP (Fig. 9E) and QC (Fig. 9G).

Evoked synaptic vesicle release (EPSP) was significantly reduced when *ND-20L* was de-
pleted in muscles by RNAi (Fig. 9A, F, Table 16). That reduction was reversed after scavenging mi-
tochondrial ROS in the muscle. Indeed, transgenic muscle-driven *sod2* gene expression sup-
pressed the *ND-20L[RNAi]* phenotype, but expression of the *catalase* and sod1 did not (Fig. 9A, F,
Table 16). Next, we tested if feeding a ROS scavenger to developing larvae would reverse the
same neurotransmission defect. Carrier feeding alone (10% EtOH) did not affect evoked neuro-
transmission, nor did it influence the neurotransmission loss caused by *ND-20L[RNAi]* (Fig. 9B, F,
Table 16). Feeding larvae 0.5 mM NACA successfully reversed the phenotype, while the nonspecif-
ic additive curcumin had no effect (Fig. 9B, F, Table 16).

We also checked different MCI manipulations. First, we examined hemizygous *ND-
30$^{EY03664/Df}$* genetic mutants. As with muscle-driven *ND-20L[RNAi]*, the *ND-30* mutant NMJs had
blunted evoked neurotransmission, but this defect was successfully reversed by ROS scavengers
(Fig. 9C, Table 16). We also impaired MCI pharmacologically, by feeding larvae 50 µM rotenone (or
0.5% DMSO carrier control), similar to conditions we previously published [13]. As with the prior study,
rotenone blunted neurotransmission (Fig. 9D, Table 16), but this effect was ameliorated by a genet-
ic background overexpressing *UAS-sod2* in the muscle (Fig. 9D, Table 16). Consistent with prior
experiments *UAS-catalase* and *UAS-sod1* were not effective.

Finally, we performed behavioral experiments on *ND-20L* and *ND-30[RNAi]* and mutant an-
imals. Consistent with the electrophysiological recordings, muscle-depleted (by RNAi) and mutant
animals showed severe defects in crawling ability. The abnormal crawling behavior was rescued by
expressing a *sod2* transgene in the muscle or feeding the larvae in a media containing NACA (Fig.
9H, Table 17). By contrast, neuronal depletion of *ND-20L* in larvae did not show any significant

crawling defects (Fig. 9H, Table 17). Together, our data suggest that excess ROS accumulation in
muscle (mROS) diminishes baseline synaptic physiology when MCI activity is lost, and it also trig-
gers aberrant crawling behavior.

DISCUSSION

We uncovered novel aspects of NMJ synapse biology controlled by Mitochondrial Complex I (MCI).
Global impairment of MCI causes profound cytological phenotypes in synaptic tissues (Figs. 1-3) [13].
By examining mitochondria directly, we discovered shared phenotypes between MCI loss and loss
of the *Drosophila* mitofusin, Marf (Fig. 2). Additionally, with MCI loss, we noted an enhancement of
mitochondrial reactive oxygen species (ROS, Fig. S1), consistent with prior work [33-37].
Unexpectedly, these perturbations spur functionally opposite responses in presynaptic neu-
rons vs. postsynaptic muscles. In motor neurons, MCI loss and mitochondrial ROS appear to trigger
a compensatory response, where the underlying cytological problems are offset by an increase in
active zone material, resulting in normal levels of evoked excitation (Fig. 4). This process requires
known intracellular calcium signaling components (Fig. 5). It also appears to require energy stores
because loss of glycolysis – which may function as a supplemental energy source to mitochondria –
abrogates the presynaptic compensation (Fig. 6). By contrast, in the muscle, MCI loss and mito-
chondrial ROS trigger a destructive response, where there is a disassembly of the postsynaptic
density (Fig. 7). This disassembly correlates with mis-apposition of pre- and postsynaptic structures
(Fig. 8) and defective neurotransmission (Fig. 9).

Disruption of mitochondrial dynamics
Energy is needed for normal levels of synaptic transmission [2]. Intuitively, a loss of synaptic mito-
chondria should blunt transmission. Several labs have implicated mitochondrial dynamics in *Dro-*
sophila synapse function, including mitochondrial fission (Dynamin related protein 1, Drp1 [48]), fusion
(Mitofusin/dMarf [28]), trafficking (Miro and Milton [47,67,68]), or quality control (Pink and Parkin [69-72]). Ad-
ditionally, it has been established from various model organisms such as flies, worms, and mice
that any misregulation in mitochondrial distribution could affect synaptic activity [48,56,73,74]. Adding to
that work, we uncovered synaptic transmission and developmental phenotypes after depletion of
Mitochondrial Complex I (MCI) at the NMJ [13].
Separate and apart from those studies, ROS has been studied in the context of mitochon-
drial dysfunction [75]. Excess ROS can trigger mitochondrial calcium uptake and subsequently trigger
apoptosis or degeneration of neurons or neural support cells [76,77]. MCI deficiency elevates ROS

levels, and this process can promote the fragmentation of mitochondria in cells like fibroblasts [78].

Collectively, the prior work suggest that mitochondrial ROS can impinge upon cell survival and mi-

tochondrial dynamics. Our study expands upon these results and supports the idea that MCI has

roles at synaptic sites. Lack of MCI in neurons causes loss of mitochondria at synaptic terminals

(Fig. 1). This defect is linked to defective mitochondrial fusion (Fig. 2), which is essential to maintain

mitochondrial integrity [79,80].

A form of presynaptic homeostatic plasticity triggered by ROS?

ROS has been linked to short-term synaptic plasticity in *Drosophila* [50], as well as long-term potentia-

tion (LTP) in mammals [81-83]. Here, we uncovered a role for ROS in augmentation of active zone ma-

terial when MCI is impaired in neurons. This finding could be considered a form of homeostatic

plasticity: an increase in active zone components likely drives potentiated release to compensate

for defective baseline synaptic transmission, which would be expected after the loss of an energy

source like mitochondria.

Homeostatic augmentations of active zone material have been observed at the *Drosophila*

NMJ. For example, *rab3* mutants have an increase in active zone material to offset decreased syn-

apse growth [84]. Additionally, ROS has been shown to be an obligate signal in *Drosophila* to main-

tain fundamental properties in both pre- and postsynaptic compartments, including at the NMJ [50]. In

our study, we observe an increase in active zone intensity (Fig. 4), which could overlap with mech-

anisms uncovered in those prior studies. Or it could be consistent with a form of homeostatic plas-

ticity at the *Drosophila* NMJ called Presynaptic Homeostatic Potentiation (PHP). PHP is initiated

when the activity of postsynaptic glutamate receptors is impaired. This decreases quantal size. The

synapse detects the impairment, and muscle-to-nerve signaling drives an increase in presynaptic

glutamate release [85,86]. This happens in part through an increase in influx of calcium into the neuron

through Ca_V2 voltage-gated calcium channels [85,87-90]. PHP coincides with an increase in the size of

the readily releasable pool (RRP) of synaptic vesicles [91-94] and increases in active zone protein con-

tent [51,53,95]. The general model is that these modifications drive the neuron to release more gluta-

mate, offsetting the initial synaptic challenge.

Could the excess mitochondrial ROS that is caused by the loss of Complex I be triggering

the same or overlapping homeostatic mechanisms? It is possible. Quenching presynaptic ROS by

expression of *sod2* or by feeding flies NACA led to a reversion of active zone material back down to

the control levels (Fig. 4E). Under ideal conditions, control levels of active zone material would sup-

port normal neurotransmission (Fig. 4A, J). However, when combined with presynaptic *ND-20L* loss

(already depleting synaptic mitochondria; Figs. 1O, 2G), there is a loss in neurotransmission ca-
pacity (Fig. 4M).

The molecular underpinnings of this mitochondrial-loss-induced homeostatic plasticity are
unknown, but prior work might offer clues. One possibility is that *sod2* changes the redox status of
the entire cell, which could potentially affect active zone components [81]. Another possibility is that
ROS could alter the release of calcium from intracellular stores such as ER; and in turn, this could
induce calcium signaling to mitochondria, which would contribute to the formation of ATP and sub-
sequent vesicle fusion [56,57]. Consistent with this idea, we observed a reduced synaptic strength at
the terminals after simultaneous blockade of calcium release and import from ER stores to mito-
chondria through genetic or pharmacological manipulations (Fig. 5). But notably, the link would not
have to be a direct one. Indeed, a recent study indicated that activity-driven mitochondrial Ca^{2+} up-
take does not depend on the ER as a source of Ca^{2+} to maintain normal synaptic strength [56].

Loss of MCI in muscle and subsequent ROS accumulation diminish synaptic excitation
Little is known about what role postsynaptic ROS plays in regulating synaptic plasticity at the NMJ.
It is unlikely that ROS is an abstract signal that triggers wholesale destruction. It likely has specific
targets. There are clues from previous work. For example, ROS signaling plays direct roles in the
activity-dependent structural plasticity of motor neurons and postsynaptic dendrites [50]. Additionally,
postsynaptic ROS plays critical roles in dendritic development. In *Drosophila*, ROS appears to act
as a developmental plasticity signal to regulate the size of the dendritic arbors. [96].

It is slightly surprising that the NMJ fails to compensate for a lack of MCI function in the
muscle. This is because the NMJ employs multiple retrograde signals to stabilize function [97]. One
hypothesis is that MCI deficiency and ROS interfere with important muscle-to-nerve signals normal-
ly required for maintaining NMJ setpoint function. Some of these signals are required to correct
acute challenges (minutes) to NMJ function. Some of the best characterized are: Bone Morphoge-
netic Protein (BMP) signaling [98-100]; the Insomniac (adaptor)-Cul3 (ubiquitin ligase)-Peflin (substrate)
signaling complex [101]; and the endosomal recycling molecules class II PI3 kinase (PI3K) and Rab11
[102]. Additionally, the field has uncovered instructive signaling molecules from the muscle that main-
tain robust neurotransmission, including: muscle-secreted Semaphorin-2b [103]; and Target of
Rapamycin (TOR) [104,105]. Future studies can address whether these or similar targets are substrates
of negative ROS regulation in the muscle.

The roles of ROS in muscle are less understood in terms of synapse regulation and func-
tion. In this study, we showed that excess ROS is sufficient to affect the organization of postsynap-
tic densities (Fig. 7), ionotropic glutamate receptor clusters (Fig. 8), and spectrin cytoskeleton (Fig.

S11) in the muscle. The correlated uncoupling of pre- and postsynaptic structures seems to be re-
sponsible for neurotransmission defects (Fig. 9), and it likely coincides with broader structural insta-
bility [62,106]. These phenotypes are reminiscent of other *Drosophila* mutant phenotypes showing a
degenerative NMJ (e.g., [62,106-108]). Consistent with a model of degeneration, we found behavioral
defects in locomotion (Fig. 9).

Despite these many defects that result from muscle depletion of MCI, we successfully res-
cued many structural and functional defects by culturing animals in an antioxidant-rich media
(0.5mM NACA) or through muscle expression of *UAS-sod2*. This is interesting because it points to
mitochondrial ROS as a signal of interest in the manifestation of synapse dysfunction. In subse-
quent work, we will take advantage of the *Drosophila* neurogenetic toolkit to identify specific targets.
In turn, these targets could be informative about how MCI dysfunction manifests in mitopathies.

RESOURCE AVAILABILITY

Lead Contact

584   The lead contact for this study is Dr. C. Andrew Frank (andy-frank@uiowa.edu).

Materials Availability

587   Dr. Frank's laboratory uses the fruit fly *Drosophila melanogaster* as a model organism. Consistent
588   with ethical conduct of research, the lab maintains *Drosophila* stocks that it generates and publish-
589   es, as well as useful precursors and derivatives of those stocks. These stocks are available to re-
590   searchers who make requests to Dr. Frank – or to the appropriate original source – and they are
591   shipped promptly.

Data and Code Availability

Consistent with the Data Sharing Policy for NIH-funded research, Dr. C. Andrew Frank's laboratory
will share data related to this study. The lab will share final data through publication of summary fig-
ures or tables. In addition, a large amount of final data derives from performing electrophysiological
recordings or synapse imaging experiments and then performing subsequent analyses. Those ex-
periments yield raw data in electronic form. Therefore, in addition to publishing final data, the lab
will share raw data and related materials with researchers who make requests to Dr. Frank (andy-
frank@uiowa.edu). No original coding was required to execute this study.

Acknowledgments

We acknowledge the Developmental Studies Hybridoma Bank (University of Iowa, USA) for anti-
bodies used in this study and the Bloomington Drosophila Stock Center (Indiana University, Bloom-
ington, USA) for fly stocks. We thank members of the Frank lab for their helpful comments and dis-
cussions in this study. BM was supported in part by NIH/NINDS grants to CAF (NS085164,
NS130108, and NS136753). Collectively, the work was supported by those same NIH/NINDS
grants, as well as funds from the University Carver College of Medicine to CAF.

Author contributions

BM and CAF designed the research; BM performed the research; BM and CAF analyzed the data;
BM and CAF wrote the paper.

Declarations of interests

The authors declare no competing financial interests.

SUPPLEMENTAL INFORMATION

Document S1:

• Supplemental Figures S1-S11 and legends.

Document S2:

• Tables 1-17: Summary data for graphs in all Figures and Supplemental Figures.
• Table 18: Detailed information about reagents and materials for items described in the
STAR METHODS section.

FIGURE LEGENDS

Figure 1: MCI-depleted flies harbor fewer mitochondria at neuromuscular junctions

Mitochondrial morphology and trafficking defects in the ventral nerve cord, distal axons, and boutons. RNAi lines and controls were crossed to a motor neuron driver (*D42-GAL4*) and a mitochondrial marker (*UAS-mitoGFP*). (A-A′), (B-B′), (C-C′) and (D-D′) represent control ventral nerve cord (VNC), a magnified section of VNC, proximal (A2) and distal (A5) axons, respectively. These tissues exhibit regular mitochondrial clusters in the soma and axons. (E-E′), (F-F′), (G-G′) and (H-H′) represent *ND-20L* knocked down ventral nerve cord (VNC), a magnified section of VNC, proximal (A2) and distal (A5) axons, respectively. Mitochondria are abnormally clustered in *ND-20L[RNAi]* in the ventral nerve cord and distal segments of A5 axons. *ND-20L[RNAi]* yields fewer mitochondria in the distal segments when compared to the proximal segments. (I-J) Representative electrophysiological traces showing evoked potentials of *mitoGFP, D42-Gal4 × UAS-ND-20*[RNAi] larvae at the A5 hemisegment of muscle 6/7 synapse. Scale bars for EPSPs (mEPSP) are x=50 ms (1000 ms) and y= 10 mV (1 mV). Fewer mitochondria at the presynaptic A5 hemisegment did not affect evoked NMJ excitation. K. Quantification showing EPSP amplitude at NMJ 6/7 in control (*D42-Gal4/+*; EPSP: 19.99 mV ± 2.53, n=6) and RNAi-depleted animals (*ND-20L[RNAi] /+;D42-Gal4/+* ;EPSP: 16.88 mV ± 1.21, n=9). (L-M) Representative images showing mitochondria morphology in control and RNAi-depleted animals in muscle. Mitochondria in *ND-20L[RNAi]*-depleted larvae are clustered compared to the control larvae. (N-Q) *ND-20L[RNAi]* contains almost no mitochondria in boutons when co-stained with pre- (HRP) or post-synaptic markers (Discs Large, Dlg). (A-H′, L-M, N-O) Scale bar: 10 µm. (P) Quantification showing the number of mitochondrial clusters at NMJ 6/7 in control (*D42-Gal4/+*; # clusters: 113.6 ± 11.97, n=7) and RNAi-depleted animals (*ND-20L[RNAi]/+; D42-Gal4/+*; # clusters: 30.33 ± 4.02, n=6). (Q-R) 3D rendered image showing the volume (µm^3) of mitochondria at NMJ 6/7 in RNAi knockdown larvae (*ND-20L[RNAi] /+; D42-Gal4/+* ; 0.11 ± 0.01 µm^3, n=16) compared to the driver control animals (*D42-Gal4/+*; 0.29 ± 0.04 µm^3, n=14). (Q-R) Scale bar: 5 µm. (S) Quantification shows a significantly lower mitochondria volume in boutons at NMJ 6/7 in RNAi-depleted animals. ***$p<0.0001$ and ***$p=0.0003$ for mitochondrial clusters and volume, respectively. Statistical analyses based on Student's t-test. Error bars represent mean ± s.e.m.

Figure 2: Loss of *ND-20L* in motor neurons phenocopies a *marf* depletion

Mitochondrial morphology and trafficking defects in the ventral nerve cord and distal axons. To label neuronal mitochondria, *UAS-[RNAi]* lines and controls were crossed to a motor neuron driver (*D42-Gal4*) and a mitochondrial marker (*UAS-mitoGFP*). (A) Ventral nerve cord (VNC): *UAS-mitoGFP* and *drp1[RNAi]* exhibit normal mitochondrial organization, *ND-20L[RNAi]* and *marf*[RNAi] exhibit clustered mitochondria, *ND-20L[RNAi] ; marf[RNAi]* and *ND-20[RNAi]; drp1[RNAi]* doubles exhibit clustered mito-chondria in the soma. The fluorescent images were skeletonized to measure mitochondrial branch length. (B) Comparison of a proximal axonal segment in A2 and a distal segment in A5. Distal seg-ments of A5 axons in *ND-20L[RNAi]* and *marf[RNAi]* contain many fewer mitochondria than proximal segments. Knocking down *ND-20L[RNAi]* and *marf[RNAi]* together does not show an additive effect. (A-B) Scale bar: 10 μm. (C-D) Histogram showing mitochondrial branch length (μm) and number (μm^2 ar-ea of bouton) in VNC and axons of the third instar larvae in the indicated genotypes. (E-L) Representa-tive images of the A2 hemisegment of muscle 6/7 NMJs in (E) *UAS-mito-GFP, D42-Gal4/+*, (F) *UAS-sod2/+; UAS-mito-GFP, D42-Gal4/+,* (G) *ND-20L[RNAi] /+; UAS-mito-GFP, D42-Gal4/+*, (H) *ND-20L[RNAi] /UAS-sod2;UAS-mito-GFP,D42-Gal4/+*, (I) *UAS-mito-GFP,D42-Gal4/marf[RNAi]* , (J) *UAS-sod2;UAS-mito-GFP,D42-Gal4/marf[RNAi]* , (K) *ND-20L[RNAi] /+;UAS-mito-GFP,D42-Gal4/marf[RNAi]* and (L) *ND-20L[RNAi] /UAS-sod2;UAS-mito-GFP,D42-Gal4/marf[RNAi]* larvae immunostained with an-tibodies against HRP (magenta) and GFP (mito-GFP:green) to label neurons and mitochondria. *ND-20L[RNAi]*- and *marf[RNAi]*-depleted animals harbor fewer mitochondria at the terminals as compared to control animals. Transgenic *UAS-sod2* rescued mitochondrial clustering defects. (E-L) Scale bar: 5 μm. (M) Histograms showing quantification of mitochondrial clusters at the NMJs in the indicated geno-types. ***p <0.0001; ns, not significant. Statistical analysis based on one-way ANOVA followed by post-hoc Tukey's multiple-comparison test. Error bars represent mean ± s.e.m.

Figure 3: *ND-20L* depletion in motor neurons affects synapse stability

Representative confocal images of NMJ synapses at muscle 6/7 of (A-A′) *D42-Gal4* control, (B-B′) *UAS-sod2* overexpression (C-C′) *D42-Gal4*-driven *ND-20L*[RNAi] (*ND-20L[RNAi] /+; D42-Gal4/+*), (D-D′) *ND-20L* knockdown with NACA rescue (*ND-20L[RNAi] /+; D42-Gal4/+* +NACA), (E-E′) *ND-20L* knockdown with *UAS-sod2* (*UAS-ND-20L[RNAi]/UAS-sod2; D42-Gal4/+*). Each condition was double immunolabeled with 22C10 (anti-Futsch, magenta) and anti-HRP (green) antibodies. The motor neuron-depleted *ND-20L[RNAi]* larvae showed a decrease in the number of Futsch-positive loops as compared to the Gal4 control. Futsch-positive loops were significantly restored to the control number when *ND-20L*[RNAi] was raised in media containing NACA or genetically expressing *UAS-sod2* in the *UAS-ND-20L[RNAi]* background. Scale bar: 10 µm. (F) Histograms showing the percentage of Futsch-positive loops in the indicated genotypes. *p=0.01 (Gal4 control vs *sod2* OE neuron), *p=0.0008 (Gal4 control vs *ND-20L*[RNAi] neuron), **p=0.001 (*ND-20L*[RNAi] neuron vs *ND-20L*[RNAi] neuron + NACA) and **p=0.0005 (*ND-20L*[RNAi] neuron vs *sod2* neuron rescue). Statistical analysis based on one-way ANOVA followed by post-hoc Tukey's multiple-comparison test. Error bars represent mean ± s.e.m.

Figure 4: Neuronal ROS (nROS) controls active zone material levels at NMJs

(A) Representative images of the A2 hemisegment of muscle 6/7 NMJs in *UAS-mito-GFP, D42-Gal4/+*, (B) *UAS-sod2/+; UAS-mito-GFP, D42-Gal4/+* (C) *ND-20L[RNAi]/+; UAS-mito-GFP, D42-Gal4/+*, (D) *ND-20L[RNAi] /+; UAS-mito-GFP, D42-Gal4/+* with NACA and (E) *ND-20L[RNAi]/UAS-sod2; UAS-mito-GFP, D42-Gal4/+* larvae immunostained with antibodies against the active zone scaffold Bruchpilot (BRP:fire-LuT) to label the active zones. BRP levels are upregulated at the NMJs in *ND-20L*[RNAi] depleted flies, while overexpression of ROS scavenger *sod2* in the neuron or feeding the larvae with N-Acetyl L-cystine amide (NACA) restores BRP to the control level. (A-E) Scale bar: 2.5 μm. (F-G) Histograms showing quantification of BRP intensity (F) and density (G) in μm^2 area of bouton at muscle 6/7 in the genotypes mentioned above. At least 8 NMJs of each genotype were used for quantification. ***p <0.0001. Error bars denote mean ± s.e.m. Statistical analysis based on one-way ANOVA followed by post-hoc Tukey's multiple-comparison test (H-H'-I-I'). Representative confocal images of muscle 6/7 NMJs in the (H-H') control (*D42-Gal4/+*) and (I-I') motor neuron Gal4 driven *ND-20L[RNAi]* (*ND-20L[RNAi] /+; D42-Gal4/+*) immunostained with antibodies against Bruchpilot (BRP: magenta) and GluRIII (green) to label a glutamate receptor subunit. (H-H'-I-I') Scale bar: 2.5 μm. There are no significant changes of GluRIII-BRP apposed clusters. At least 8 NMJs of each genotype were used for quantification (J-P). Representative traces, quantification of mEPSPs, EPSPs and quantal content in the indicated genotypes. Scale bars for EPSPs (mEPSP) are x=50 ms (1000 ms) and y= 10 mV (1 mV). EPSPs amplitudes were maintained in *ND-20L[RNAi]*-depleted flies due to induction of BRP; however, NMJ with *sod2* rescued *ND-20L[RNAi]* in neurons showed diminished evoked release when compared with *ND-20L[RNAi]*. Minimum 8 NMJs recordings of each genotype were used for quantification. *$p < 0.05$, ***$p <0.0001$; ns, not significant. Statistical analysis based on one-way ANOVA followed by post-hoc Tukey's multiple-comparison test. Error bars denote the standard error of the mean. (Q) Representative images of the A2 hemisegment of muscle 6/7 NMJs in *UAS-mito-GFP, D42-Gal4/+*, (R) *UAS-sod2/+; UAS-mito-GFP, D42-Gal4/+* (S) *ND-20L[RNAi] /+; UAS-mito-GFP, D42-Gal4/+*, and (T) *ND-20L[RNAi]/UAS-sod2; UAS-mito-GFP,D42-Gal4/+* larvae immunostained with antibodies against HRP (magenta) and GFP (mito-GFP:green) to label neurons and mitochondria. *ND-20L*-depleted and *sod2*-rescued *ND-20L*[RNAi] animals harbor fewer mitochondria at the terminals than control animals. (Q-T) Scale bar: 5 μm. (U) Histograms showing quantification of mitochondrial clusters in the above-indicated genotypes. (V-X) Schematic illustration showing ROS (magenta) levels, BRP (grey) and mitochondria (red) number in the indicated genotypes. At least 8 NMJs of each genotype were used for quantification. ***$p<0.0001$. Error bars represent mean ± s.e.m. Statistical analysis based on one-way ANOVA followed by post-hoc Tukey's multiple-comparison test.

Figure 5: Loss of MCI subunit induces ER-mediated calcium release to maintain evoked neuro-transmission at the NMJs

(A-D) Schematics illustrating the role of IP$_3$ receptor (IP$_3$R), Ryanodine receptor (RyR) in the endoplasmic reticulum, mitochondrial calcium uniporter Complex (MCU), Ca$_v$2 calcium channel and synaptic vesicles at the presynaptic nerve terminal. The IP3 and Ryanodine receptors were blocked pharmacologically by using IP$_3$R antagonist Xestospongin C or presynaptically expressing *UAS-IP$_3$-Sponge* and RyR antagonist Dantrolene, while *mcu*[RNAi] was used to block the Mitochondrial calcium uniporter complex. (E-P) Representative traces of mEPSPs and EPSPs in (E) pan-neuronal Gal4 control (*elaV(C155)-Gal4/+*), (F) pan-neuronal Gal4 control (*elaV(C155)-Gal4/+*) with an acute application (10 minutes) of 20 µM Xestospongin C and 10 µM Dantrolene, (G) pan-neuronal Gal4 driven *ND-20L*[RNAi] (*elaV(C155)-Gal4)/+;ND-20L[RNAi] /+*), (H) pan-neuronal Gal4-driven *ND-20L*[RNAi] (*elaV(C155)-Gal4/+;ND-20L[RNAi] /+*) with 20 µM Xestospongin C and 10 µM Dantrolene, (I) pan-neuronal Gal4 driven UAS-*IP3-Sponge.m30* with an acute application of 10 µM Dantrolene (*elaV(C155)-Gal4)/+;ND-20L[RNAi] /+;UAS-IP3-Sponge.m30/+*), (J) pan neuronal Gal4 driven *mcu*[RNAi] with 20 µM Xestospongin C and 10 µM Dantrolene (*elaV(C155)-Gal4)/+;mcu[RNAi] /+),* (K) pan-neuronal *mcu[RNAi] + ND-20L*[RNAi] with 20 µM Xestospongin C and 10 µM Dantrolene and(*elaV(C155)-Gal4)/+;mcu[RNAi] /ND-20L[RNAi]*) and (L) pan neuronal *UAS-IP3-Sponge.m30 + mcu[RNAi] + ND-20L*[RNAi] with 10 µM Dantrolene (*elaV(C155)-Gal4/+;ND-20L[RNAi] /mcu[RNAi] ;UAS-IP$_3$-Sponge.m30/+*), (M) Pan neuronal Gal4 (*elaV(C155)-Gal4/+*) with DMSO, (N) pan neuronal Gal4 (*elaV(C155)-Gal4/+*) with 20 µM BAPTA-AM, (O) pan-neuronal Gal4 driven *ND-20L*[RNAi] (*elaV(C155)-Gal4)/+;ND-20L[RNAi] /+*) with DMSO and (P) pan neuronal Gal4 driven *ND-20L*[RNAi] (*elaV(C155)-Gal4)/+;ND-20L[RNAi] /+*) with 20 µM BAPTA-AM. Scale bars for EPSPs (mEPSP) are x=50 ms (1000 ms) and y= 10 mV (1 mV). Note that EPSPs amplitudes were reduced in pan-neuronal Gal4 driven *mcu[RNAi] + ND-20L*[RNAi] with an acute exposure of 20 µM Xestospongin C, 10 µM Dantrolene or *UAS-IP$_3$-sponge.m30 + mcu[RNAi] + ND-20L*[RNAi] with 10 µM Dantrolene and pan-neuronal Gal4 driven *ND-20L*[RNAi] with 20 µM BAPTA-AM. (Q-T) Histograms showing average mEPSPs, EPSPs amplitude, and quantal content in the indicated genotypes. A minimum of 8 NMJs recordings of each genotype were used for quantification. **p < 0.05 (EPSP and QC: *ND-20L*[RNAi] neuron + DMSO, wash vs *ND-20L*[RNAi] neuron + BAPTA-AM, wash), *p < 0.05, **p=0.001, ***p <0.0001; ns, not significant. Statistical analysis based on one-way ANOVA followed by post-hoc Tukey's multiple-comparison test. Error bars represent mean ± s.e.m.

 Figure 6: Loss of MCI subunit induces glycolysis and regulates levels of active zone materials

to stabilize synaptic strength

(A) Schematic illustrations showing steps of glucose metabolism and ATP production during glycolysis and TCA cycle in the cell. (B-G) Representative traces of mEPSPs and EPSPs in (B) pan-neuronal Gal4 control (*elaV(C155)-Gal4/+* with DMSO), (C) pan-neuronal Gal4 control (*elaV(C155)-Gal4/+*) with an acute application (30 minutes) of 150 µM Lonidamine (LDA) (D) pan-neuronal Gal4 driven *ND-20L*[RNAi] (*elaV(C155)-Gal4)/+; ND-20L[RNAi] /+* with DMSO), (E) pan-neuronal Gal4 driven *ND-20L*[RNAi] (*elaV(C155)-Gal4)/+; ND-20L[RNAi]/+*) with 150 µM Lonidamine, (F) pan-neuronal Gal4 con-trol (*elaV(C155)-Gal4/+*) and HL3 containing 2-Deoxy-D-glucose and (G) pan-neuronal Gal4 driven *ND-20L*[RNAi] (*elaV(C155)-Gal4)/+; ND-20L[RNAi] /+*) with HL3 containing 2-Deoxy-D-glucose. Scale bars for EPSPs (mEPSP) are x=50 ms (1000 ms) and y= 10 mV (1 mV). The EPSPs amplitudes were re-duced in pan-neuronal Gal4-driven *ND-20L*[RNAi] with an acute exposure of 150 µM Lonidamine (LDA) or pan-neuronal Gal4-driven *ND-20L*[RNAi] in HL3 containing 2-Deoxy-D-glucose for 30 minutes. How-ever, we saw a reduction in mEPSP amplitudes when incubated in HL3 containing 2-Deoxy-D-glucose for 30 minutes in pan-neuronal Gal4-driven *ND-20L[RNAi]* -depleted larvae compared to the control an-imals. (H-J) Histograms showing average mEPSPs, EPSPs amplitude, and frequencies in the indicated genotypes. A minimum of 8 NMJ recordings of each genotype were used for quantification. ,**p=0.003 (mEPSP amplitude: *elaV(C155)-Gal4/+*2-Deoxy-D-glucose vs *elaV(C155)-Gal4)/+;ND-20L[RNAi] / +*2-Deoxy-D-glucose), **p=0.002 (EPSP amplitude: *elaV(C155)-Gal4/+*2-Deoxy-D-glucose vs *elaV(C155)-Gal4)/+;ND-20L[RNAi] /+*2-Deoxy-D-glucose),***p=0.0001 (EPSP amplitude: *elaV(C155)-Gal4/+*LDA vs *elaV(C155)-Gal4)/+;ND-20L[RNAi] / +*LDA); ns, not significant. Statistical analysis is based on the Stu-dent's t-test for pairwise sample comparison. Error bars represent mean ± s.e.m. (K-P) Representative images of the A2 hemisegment of muscle 6/7 NMJs in the above-indicated genotypes immunostained with antibodies against the active zone scaffold bruchpilot (BRP:fire-LuT) to label the active zones. The BRP levels downregulated at the NMJs in pan-neuronal Gal4 driven *ND-20L*[RNAi] either incubated with 150 µM LDA or in HL3 containing 2-Deoxy-D-glucose for 30 minutes. (K-P) Scale bar: 5 µm. (Q) Histograms showing quantification of BRP intensity in µm^2 area of bouton at muscle 6/7 in the geno-types mentioned above. At least 8 NMJs of each genotype were used for quantification.***p<0.0001 (BRP levels: *elaV(C155)-Gal4/+*DMSO vs *elaV(C155)-Gal4)/+;ND-20L[RNAi]/+* +DMSO),**p=0.008 (BRP levels: *elaV(C155)-Gal4/+*LDA vs *elaV(C155)-Gal4)/+;ND-20L[RNAi] /+*LDA), ***p=0.0005 (BRP levels: *elaV(C155)-Gal4/+*2-Deoxy-D-glucose vs *elaV(C155)-Gal4)/+;ND-20L[RNAi] /+*2-Deoxy-D-glucose). Error bars denote mean ± s.e.m. Statistical analysis based on one-way ANOVA followed by post-hoc Tukey's multiple-comparison test.

Figure 7: *ND-20L* subunit in muscles is required to promote normal synapse growth

(A-I) Representative confocal images of NMJ synapses at muscle 6/7 of (A) Muscle-Gal4 control (*BG57-Gal4/+*), (B) Muscle Gal4 driven *ND-20L*[RNAi] (*ND-20L[RNAi] /+; BG57-Gal4/+*), (C) *sod2* muscle rescue (*UAS-sod2/ND-20L[RNAi] ; BG57-Gal4/BG57-Gal4*), (D) *sod1* muscle rescue (*UAS-sod1/ND-20L[RNAi] ; BG57-Gal4/BG57-Gal4*), (E) catalase muscle rescue (*UAS-cat/ND-20L[RNAi] ; BG57-Gal4/BG57-Gal4*), (F) Pan neuronal Gal4 control (*elaV(C155)-Gal4)/+*), (G) *elaV(C155)- Gal4)-Gal4* driven *ND-20L*[RNAi] (*elaV(C155)-Gal4/+; ND-20L[RNAi] /+*), (H) Motor neuron Gal4 control *(D42-Gal4/+)* and (I) *D42*-Gal4-driven *ND-20L*[RNAi] (*ND-20L[RNAi] /+; D42-Gal4/+*) double immunolabeled with Dlg (magenta) and HRP (green) antibodies. (A-I) Scale bar: 10 µm. The NMJ morphological defects in *ND-20L*[RNAi] were restored upon co-expression of *UAS-Sod2 in* muscle; however, it did not rescue with *UAS-sod1* or *UAS-catalase* transgene. (J-L) Histograms show the number of boutons, muscle area, and average NMJ length at muscle 6/7 of A2 hemisegment in the indicated genotypes. *p <0.05, *p=0.006 (# of boutons: Muscle *ND-20L*[RNAi] vs *sod2* muscle rescue), **p=0.0002 (# of boutons: Muscle *ND-20L*[RNAi] vs Muscle *ND-20L*[RNAi] +NACA), *p=0.009 (# of branches: Muscle Gal4 vs Muscle *sod2* OE), ***p=0.002 (# of branches: Muscle Gal4 vs Muscle *sod1* OE), *p=0.008 (# of branches: Muscle Gal4 vs Muscle *cat* OE), ***p=0.001 (Muscle Gal4 vs Muscle *ND-20L*[RNAi]), ***p<0.0001; ns, not significant. Statistical analysis based on one-way ANOVA with post-hoc Tukey's test for multiple and Student's t-tests for pairwise comparison. Error bars represent mean ± s.e.m. (M-T) Representative confocal images of boutons at the third instar larval NMJ synapse in (M) Muscle-Gal4 control (*BG57-Gal4/+*), (N) Muscle Gal4 driven *UAS-sod2* (*UAS-sod2/+; BG57-Gal4/+*), (O) *UAS-sod1* (*UAS-sod1/+; BG57-Gal4/+*), (P) *UAS-catalase* (*UAS-catalase/+; BG57-Gal4/+*) (Q) Muscle *ND-20L*[RNAi] (*ND-20L[RNAi] /+; BG57-Gal4* (R) *sod2* muscle rescue (*UAS-sod2/ND-20L[RNAi] ; BG57-Gal4/BG57-Gal4*), (S) *sod1* muscle rescue (*UAS-sod1/ND-20L[RNAi] ; BG57-Gal4/BG57-Gal4*) and (T) catalase muscle rescue (*UAS-cat/ND-20L[RNAi] ; BG57-Gal4/BG57-Gal4*) animals double immunolabeled with anti-HRP (green) and anti-Dlg (magenta) antibodies. (M-T) Scale bar: 5 µm. Note that the gross morphology of SSRs and immunoreactivity of Dlg were reduced in *ND-20L[RNAi]* animals. As mentioned, phenotypes were restored to wild-type level when *ND-20L*[RNAi] -depleted flies were reared in NACA or by genetically expressing *sod2* transgene in muscle. (U-V) Histograms showing normalized synaptic fluorescence of Dlg and bouton area in the indicated genotypes. *p <0.01, **p <0.001, ***p<0.0001; ns, not significant. Error bars represent mean ± s.e.m. Statistical analysis based on one-way ANOVA with post-hoc Tukey's test for multiple and Student's t-tests for pairwise comparison.

Figure 8: *ND-20L* subunit in muscle affects the organization of GluRs cluster in *Drosophila*.
Representative confocal images of boutons at third instar larval NMJ synapse in (A) Muscle-Gal4 con-
trol (*BG57-Gal4/+*), (B) Muscle *ND-20L*[RNAi] (*ND-20L[RNAi] /+; BG57-Gal4/+*), (C) Muscle Gal4 driven
UAS-sod2 (*UAS-sod2/+; BG57-Gal4/+*), (D) *ND-20L[RNAi] /+; BG57-Gal4/+*NACA) and (E) *sod2* mus-
cle rescue (*UAS-Sod2/ND-20L[RNAi] ; BG57-Gal4/BG57-Gal4*) animals immunolabeled with active
zone marker BRP (green) and anti-GluRIII (magenta) antibodies. Scale bar: 5 µm. Note that GluRIII
apposed clusters with BRP are missing in the *ND-20L*[RNAi]-depleted animals (marked in arrow) com-
pared to control. These phenotypes were restored to normal when *ND-20L*[RNAi]-depleted flies were
reared in media containing NACA or genetically expressing *sod2* transgene in muscle. (F) Histograms
showing quantification of the number of missing BRP-GluRIII apposed puncta per bouton in the indicat-
ed genotypes. (G-J) Similar phenotypes were observed when analyzed for BRP-GluRIIA apposed clus-
ters in boutons. (K) Histograms showing quantification of the number of missing BRP-GluRIIA apposed
puncta per bouton in the indicated genotypes. ***p<0.0001. Error bars represent mean ± s.e.m. Statisti-
cal analysis based on one-way ANOVA with post-hoc Tukey's test for multiple comparisons.

Figure 9: Loss of MCI subunits affects synaptic transmission via the formation of excess ROS in the muscle

(A) Representative traces of mEPSPs and EPSPs in muscle-Gal4 control (*BG57-Gal4/+*), muscle Gal4 driven *ND-20L*[RNAi] (*ND-20L[RNAi] /+; BG57-Gal4/+),* muscle *catalase* rescue (2X muscle-Gal4>*UAS-catalase/ND-20L*[RNAi] : *UAS-catalase/ND-20L[RNAi] ; BG57-Gal4/BG57-Gal4*), muscle *sod1* rescue (2X muscle-Gal4>*UAS-sod1/ND-20L*[RNAi] : *UAS-sod1/ND-20L[RNAi] ; BG57-Gal4/BG57-Gal4*), muscle *sod2* rescue (2X muscle-Gal4>*UAS-sod2/ND-20L*[RNAi] : *UAS-sod2/ND-20L[RNAi] ; BG57-Gal4/BG57-Gal4*) animals. Note that EPSPs amplitudes were reduced in *ND-20L[RNAi],* and the phenotype was restored to wild-type levels by expressing *sod2* transgene in the muscle but not with *sod1* and *catalase* transgenes. (B) Representative traces of mEPSPs and EPSPs in muscle Gal4 control (*BG57-Gal4/+*) larvae raised on 10% EtOH, *ND-20L* muscle[RNAi] (*ND-20L[RNAi] /+; BG57-Gal4/+*) raised on 10% EtOH, *ND-20L* muscle[RNAi] (*ND-20L[RNAi] /+; BG57-Gal4/+*) raised on 0.5mM curcumin, *ND-20L* muscle[RNAi] (*ND-20L[RNAi] /+; BG57-Gal4/+*) raised on 0.5mM NACA, (C) Representative traces of mEPSPs and EPSPs in *ND-30^{epgy}/df* mutants, *ND-30^{epgy}/df* mutants raised on 0.5 mM NACA and *UAS-sod2* muscle rescued *ND-30^{epgy}/df* mutant (*UAS-sod2/+; ND-30 (df), BG57-Gal4/ND-30^{epgy}*) animals. The EPSPs amplitudes were restored to wild type when muscle depleted *ND-20L*[RNAi] larvae were raised in food containing NACA or by *UAS-sod2* muscle overexpression in *ND-20L*[RNAi] depleted animals. (D) Representative traces of mEPSPs and EPSPs in muscle-Gal4 control (*BG57-Gal4/+*) larvae raised on DMSO, muscle Gal4 (*BG57-Gal4/+*) larvae raised on 50µM rotenone (complex I inhibitor), *UAS-catalase (UAS-catalase/+; BG57-Gal4/)*, *UAS-sod1(UAS-sod1/+; BG57-Gal4/),* and *UAS-sod2 (UAS-sod2/+; BG57-Gal4/+)* muscle over-expression animals raised on 50µM rotenone. The EPSPs amplitudes were suppressed in rotenone raised larvae overexpressing *UAS-Sod2* in muscle *(UAS-sod2/+; BG57-Gal4/)* due to its free radical scavenging activity. Scale bars for EPSPs (mEPSP) are x=50 ms (1000 ms) and y= 10 mV (1 mV). (E-G) Histograms showing average mEPSPs, EPSPs amplitude, and quantal content in the indicated genotypes. Minimum 8 NMJs recordings of each genotype were used for quantification. (H) Histogram representing crawling behavior (in cm) of the larvae in the indicated genotypes. Knocking down *ND-20L*[RNAi] or *ND-30*[RNAi] in muscle and *ND-30^{epgy}/df* mutants showed a severe defect in crawling behavior. The abnormal crawling behavior was rescued by expressing a *sod2* transgene in the muscle or rearing the larvae in a media containing NACA. Moreover, neuronally depleting *ND-20L* did not show any notable change in crawling defects. Minimum 10 animals were analyzed for crawling behavioral analysis. *$p <$ 0.05 (mEPSP amplitude: Muscle *ND-20L*[RNAi] +10% EtOH vs Muscle *ND-20L*[RNAi] +0.5 mM NACA),**$p=0.006$ (mEPSP amplitude: *ND-30* mutant vs *N-30* mutant+0.5 mM NACA), *$p=0.0004$ (mEPSP amplitude: Muscle-Gal4+0.5% DMSO vs. Muscle *cat* OE+50mM rotenone),*$p=0.039$ (mEPSP

amplitude: Muscle Gal4+50mM rotenone vs Muscle *sod2* OE+50mM rotenone), *p=0.001 (EPSP ampli-

tude: Muscle *ND-20L*[RNAi] vs *sod2* muscle rescue),**p=0.003 (EPSP amplitude: Muscle Gal4+10%

EtOH vs Muscle *ND-20L*[RNAi] +0.5mM curcumin), **p=0.0004 (EPSP amplitude: Muscle *ND-*

20L[RNAi] +10% EtOH vs Muscle *ND-20L*[RNAi] +0.5mM NACA), *p=0.004 (EPSP amplitude: Muscle-

Gal4+0.5% DMSO vs Muscle Gal4+50mM rotenone), *p=0.015 (EPSP amplitude: Muscle-Gal4+0.5%

DMSO vs Muscle *cat* OE+50mM rotenone),**p=0.015 (EPSP amplitude: Muscle-Gal4+0.5% DMSO vs

Muscle *sod1* OE+50mM rotenone), **p=0.001 (QC*: ND-30* mutant vs *sod2* muscle rescue),**p=0.003

(QC: Muscle Gal4+50mM rotenone vs Muscle *sod*2 OE+50mM rotenone), **p=0.0002 (Distance

crawled: Muscle Gal4 vs Muscle *ND-30*[RNAi]), ***p <0.0001; ns, not significant. Statistical analysis

based on one-way ANOVA with post-hoc Tukey's test for multiple and Student's t-test for pairwise

comparison.

TABLES

888   All tables contain raw data supporting the graphs in the figures and are housed in the supplemen-
889   tary section.

STAR METHODS

Experimental Model Details

Drosophila husbandry

Drosophila melanogaster was cultured on a standard cornmeal media containing molasses and

yeast prepared according to the Bloomington *Drosophila* Stock Center (BDSC, Bloomington, IN)

recipe. Fruit flies husbandry was performed according to standard practices [109]. As specified, larvae

were raised at 18°C, 25°C, or 29°C in humidity-controlled and light-controlled Percival DR-36VL in-

cubators (Geneva Scientific). Control,[RNAi] , and mutant animals were grown in media containing

0.5 mM curcumin (Sigma Aldrich), 0.5 mM N-acetyl cysteine amide (NACA) (Sigma Aldrich), or 50

µM rotenone (Sigma Aldrich) depending upon the experimental condition.

Drosophila stocks

904   *w*^{1118} was used as a non-transgenic wild-type stock. *UAS-IP₃-sponge* line (*UAS-IP₃-Sponge.m30*)

905   was obtained from Dr. Masayuki Koganezawa's lab. The GAL4 drivers used in this study were

906   *elav*^{C155}-Gal4, *Sca*-Gal4 and *BG57*-Gal4, *D42*-Gal4, and *OK371*-Gal4. Several *UAS-RNAi* and ge-

907   netic mutant lines were obtained from the Bloomington *Drosophila* stock center (Table 18).

Method Details

Immunohistochemistry

Wondering third instar larvae were dissected and fixed on a sylgard Petri plate in ice-cold HL-3 and fixed in 4% paraformaldehyde in PBS for 30 minutes or in Bouin's fixative for 2 minutes as described earlier [110]. Briefly, larvae were washed with PBS containing 0.2% Triton X-100 (PBST) for 30 min, blocked for an hour with 5% normal goat serum in PBST, and incubated overnight in primary antibodies at 4°C followed by washes and incubation in secondary antibodies. Monoclonal antibodies: anti-Dlg (4F3), anti-DGluRIIA (8B4D2), anti-CSP (ab49), anti-Synapsin (3C11), anti-Futsch (22C10), anti-Bruchpilot (nC82) and anti-α-Spectrin (3A9) were obtained from the Developmental Studies Hybridoma Bank (University of Iowa, USA) and were used at 1:30 dilution. Rabbit anti-GFP (Abcam) was used at 1:200 dilutions. Anti-GluRIII (1:100) ([111]; Table 18) was gifted by Aaron DiAntonio (Washington University, St. Louis, U.S.A.). Fluorophore coupled secondary antibodies Alexa Fluor 488, Alexa Fluor 568 or Alexa Fluor 647 (Molecular Probes, ThermoFisher Scientific) were used at 1:400 dilution. Alexa 488 or 647 and Rhodamine conjugated anti-HRP were used at 1:800 and 1:600 dilutions, respectively. The larval preparations were mounted in VECTASHIELD (Vector Laboratories, USA) and imaged with a laser scanning confocal microscope (LSM 710; Carl Zeiss). All the images were processed with Adobe Photoshop 7.0 (Adobe Systems, San Jose, CA).

Confocal imaging, quantification, and morphometric analysis

Samples were imaged using a Carl Zeiss scanning confocal microscope equipped with 63×/1.4 NA oil immersion objective using separate channels with four laser lines (405, 488, 561, and 637 nm) at room temperature. The stained NMJ boutons were counted using anti-Synapsin or anti-HRP co-stained with anti-Dlg on muscle 6/7 of A2 hemisegment, considering each Synapsin or HRP punctum to be a bouton. At least 8 NMJs were used for bouton number quantification. For fluorescence quantifications of GluRs, Dlg, Brp, α-Spectrin, HRP, and CSP, all genotypes were immunostained in the same tube with identical reagents, then mounted and imaged in the same session. Z-stacks were obtained using identical settings for all genotypes with z-axis spacing between 0.2-0.5 μm and optimized for detection without saturation of the signal.

ROS and rotenone incubation assay

Larvae were dissected in ice-cold calcium-free HL3 to label ROS in neurons. ROS levels were detected in mitochondria by incubating live preparation in 1X Schneider's media with MitoSOX™ Red (Molecular Probes, ThermoFisher Scientific) fluorogenic dye at 1:200 dilutions for 20-30

minutes. Briefly, larvae were washed with HL3, mounted in VECTASHIELD (Vector Laboratories,

USA.), and immediately imaged in a laser scanning confocal microscope (LSM 710; Carl Zeiss). To

study the effect of DMSO and rotenone (Sigma Aldrich) in BRP, larvae were dissected in HL3 and

incubated in 1X Schneider's media containing either 500 µM of rotenone or DMSO. After every 30

minutes, the old media was replaced with fresh media containing rotenone or DMSO. The above

preparations were fixed with 4% paraformaldehyde, stained with anti-nc82 antibodies, mounted and

imaged in a confocal microscope.

Electrophysiology and pharmacology

All dissections and recordings were performed in modified HL-3 saline [112] containing 70 mM NaCl, 5

mM KCl, 10 mM MgCl2, 10 mM NaHCO3, 115 mM sucrose, 4.2 mM trehalose, 5 mM HEPES, and

0.5 mM CaCl2 (unless otherwise noted), pH 7.2. Neuromuscular junction sharp electrode (electrode

resistance between 20-30 MΩ) recordings were performed on muscles 6/7 of abdominal segments

A2 and A3 in wandering third-instar larvae as described [109]. Recordings were performed on a Leica

microscope using a 10x objective and acquired using an Axoclamp 900A amplifier, Digidata 1440A

acquisition system, and pClamp 10.7 software (Molecular Devices). Electrophysiological sweeps

were digitized at 10 kHz and filtered at 1 kHz. Data were analyzed using Clampfit (Molecular Devic-

es) and MiniAnalysis (Synaptosoft) software. Miniature excitatory postsynaptic potentials (mEPSPs)

were recorded in the absence of any stimulation and motor axons were stimulated to elicit excitato-

ry postsynaptic potentials (EPSPs).

Larval crawling assay

Vials containing third instar larvae were analyzed for this assay. Vials were poured with 4 ml of 20%

sucrose solution and left for 10 min to let the larvae float on top. Floating third instar animals were

poured into a petri dish and washed gently twice with deionized water in a paintbrush. A minimum

of 10 larvae of each genotype were analyzed on a 2% agarose gel in a petri dish with gridline mark-

ings 1 cm on graph paper. The larvae were acclimatized in the petri dish before videotaping. The

average distance crawled (in centimeters) by larvae was calculated based on the average number

of gridlines passed in 30 sec [110].

Quantification And Statistical Analysis

Quantification of mitochondrial branch length

Controls and[RNAi] -depleted animals were analyzed using mitochondrial marker Mito-GFP, *D42* driver in cell bodies from larvae ventral nerve cord. All images were processed using image J software. Z-stacks of individual neurons were merged. The Mito-GFP signal was enhanced by adjusting brightness and contrast. The binary masks were created using Image>Adjust>Threshold., Method:Otsu, and Background:Dark. The branches were generated using Process>Binary>Skeletonised [113]. These skeletonized images were analyzed, and branch length of the individual cluster was manually calculated using image J tools.

Imaging quantifications

Maximum intensity projections were used for quantitative image analysis with the Image J software (National Institutes of Health) analysis toolkit. Boutons from muscle 4 or type Ib terminal boutons on the muscle 6/7 of A2 hemisegment from at least six NMJ synapses were used for quantification using Image J software. Student's t-test for pairwise and one-way ANOVA with post-hoc Tukey's test for multiple comparisons was used for statistical analysis, using GraphPad Prism Software. Specific p-value and tests are noted in the figures and figure legends and supplementary files and shown in graphs as follows: $*p<0.05$, $**p<0.001$, and $***p<0.0001$. The data are presented as mean ± s.e.m. For quantification of Futsch loops, third instar larval preparations were double immunostained with HRP, 22C10 and images were captured in Zeiss LSM710 confocal microscope. Only NMJs of muscles 6/7 of A2 hemisegment were used for quantification. The images were digitally magnified using image J software and the total number of HRP-positive boutons was manually counted in each image. Futsch positive loops, which are co-localized with HRP, were included in this analysis. Images with incomplete loops and diffused staining were not included in the count [114].

Electrophysiological analysis

Average mEPSP, EPSP, and quantal content were calculated for each genotype by dividing EPSP amplitude by mEPSP amplitude. Muscle input resistance (R_{in}) and resting membrane potential (V_{rest}) were monitored during each experiment. Recordings were rejected if the V_{rest} was above -60 mV, and R_{in} was less than 5 MΩ. Pharmacological agents were bath applied in recording saline at the final concentrations indicated in the text, figures, and tables. The agents included Xestospongin C (Abcam), Dantrolene (Tocris) and BAPTA-AM (Sigma Aldrich). Failure analysis was performed in HL-3 solution containing 0.1 mM CaCl2, which resulted in failures in about half of the stimulated responses in wild-type larvae. A total of 30 trials (stimulations) were performed at each NMJ in all genotypes. The failure rate was obtained by dividing the total number of failures by the total number

1007 of trials (100). High-frequency (10 Hz) recordings were performed at a Ca^{2+} concentration of 2 mM

1008 and paired-pulse recordings (10 Hz) were performed at a Ca^{2+} concentration of 0.4 mM and 1.5

1009 mM, respectively. Paired-pulse ratios were calculated as the EPSP amplitude of the second re-

1010 sponse divided by the first response (EPSP2/EPSP1).

1011

1012 **REFERENCES**

1013

1. Chamberlain, K.A., and Sheng, Z.H. (2019). Mechanisms for the maintenance and
regulation of axonal energy supply. Journal of neuroscience research *97*, 897-913.
10.1002/jnr.24411.

2. Hall, C.N., Klein-Flugge, M.C., Howarth, C., and Attwell, D. (2012). Oxidative
phosphorylation, not glycolysis, powers presynaptic and postsynaptic mechanisms
underlying brain information processing. J Neurosci *32*, 8940-8951.
10.1523/JNEUROSCI.0026-12.2012.

3. Neupane, P., Bhuju, S., Thapa, N., and Bhattarai, H.K. (2019). ATP Synthase: Structure,
Function and Inhibition. Biomol Concepts *10*, 1-10. 10.1515/bmc-2019-0001.

4. Brown, G.C. (1992). Control of respiration and ATP synthesis in mammalian mitochondria
and cells. Biochem J *284 (Pt 1)*, 1-13. 10.1042/bj2840001.

5. Faria-Pereira, A., and Morais, V.A. (2022). Synapses: The Brain's Energy-Demanding Sites.
International journal of molecular sciences *23*. 10.3390/ijms23073627.

6. Vandecasteele, G., Szabadkai, G., and Rizzuto, R. (2001). Mitochondrial calcium
homeostasis: mechanisms and molecules. IUBMB Life *52*, 213-219.
10.1080/15216540152846028.

7. Mattson, M.P., Gleichmann, M., and Cheng, A. (2008). Mitochondria in neuroplasticity and
neurological disorders. Neuron *60*, 748-766. 10.1016/j.neuron.2008.10.010.

8. Hamanaka, R.B., and Chandel, N.S. (2010). Mitochondrial reactive oxygen species regulate
cellular signaling and dictate biological outcomes. Trends Biochem Sci *35*, 505-513.
10.1016/j.tibs.2010.04.002.

9. Kastaniotis, A.J., Autio, K.J., Keratar, J.M., Monteuuis, G., Makela, A.M., Nair, R.R.,
Pietikainen, L.P., Shvetsova, A., Chen, Z., and Hiltunen, J.K. (2017). Mitochondrial fatty acid
synthesis, fatty acids and mitochondrial physiology. Biochim Biophys Acta Mol Cell Biol
Lipids *1862*, 39-48. 10.1016/j.bbalip.2016.08.011.

10. Banoth, B., and Cassel, S.L. (2018). Mitochondria in innate immune signaling. Transl Res
202, 52-68. 10.1016/j.trsl.2018.07.014.

11. Duarte, M., Sousa, R., and Videira, A. (1995). Inactivation of genes encoding subunits of the
peripheral and membrane arms of neurospora mitochondrial complex I and effects on
enzyme assembly. Genetics *139*, 1211-1221. 10.1093/genetics/139.3.1211.

12. Garcia, C.J., Khajeh, J., Coulanges, E., Chen, E.I., and Owusu-Ansah, E. (2017).
Regulation of Mitochondrial Complex I Biogenesis in Drosophila Flight Muscles. Cell reports
20, 264-278. 10.1016/j.celrep.2017.06.015.

13. Mallik, B., and Frank, C.A. (2022). Roles for Mitochondrial Complex I Subunits in Regulating
Synaptic Transmission and Growth. Front Neurosci *16*, 846425. 10.3389/fnins.2022.846425.

14. Sharma, L.K., Lu, J., and Bai, Y. (2009). Mitochondrial respiratory complex I: structure,
function and implication in human diseases. Curr Med Chem *16*, 1266-1277.
10.2174/092986709787846578.

15. Yang, J.L., Mukda, S., and Chen, S.D. (2018). Diverse roles of mitochondria in ischemic
stroke. Redox Biol *16*, 263-275. 10.1016/j.redox.2018.03.002.

16. Marin, S.E., Mesterman, R., Robinson, B., Rodenburg, R.J., Smeitink, J., and Tarnopolsky,
M.A. (2013). Leigh syndrome associated with mitochondrial complex I deficiency due to
novel mutations In NDUFV1 and NDUFS2. Gene *516*, 162-167.
10.1016/j.gene.2012.12.024.

17. Goto, Y. (2000). Mitochondrial encephalomyopathy. Neuropathology *20 Suppl*, S82-84.
10.1046/j.1440-1789.2000.00304.x.

18. Shoffner, J.M. (2000). Mitochondrial myopathy diagnosis. Neurol Clin *18*, 105-123.
10.1016/s0733-8619(05)70180-8.

19. Burman, J.L., Itsara, L.S., Kayser, E.B., Suthammarak, W., Wang, A.M., Kaeberlein, M.,
Sedensky, M.M., Morgan, P.G., and Pallanck, L.J. (2014). A Drosophila model of
mitochondrial disease caused by a complex I mutation that uncouples proton pumping from
electron transfer. Dis Model Mech *7*, 1165-1174. 10.1242/dmm.015321.

20. Schon, E.A., and Manfredi, G. (2003). Neuronal degeneration and mitochondrial
dysfunction. The Journal of clinical investigation *111*, 303-312. 10.1172/JCI17741.

21. He, L., He, T., Farrar, S., Ji, L., Liu, T., and Ma, X. (2017). Antioxidants Maintain Cellular
Redox Homeostasis by Elimination of Reactive Oxygen Species. Cell Physiol Biochem *44*,
532-553. 10.1159/000485089.

22. Folbergrova, J., and Kunz, W.S. (2012). Mitochondrial dysfunction in epilepsy.
Mitochondrion *12*, 35-40. 10.1016/j.mito.2011.04.004.

23. Lanza, I.R., and Nair, K.S. (2010). Mitochondrial function as a determinant of life span.
Pflugers Arch *459*, 277-289. 10.1007/s00424-009-0724-5.

24. Martin, L.J. (2012). Biology of mitochondria in neurodegenerative diseases. Prog Mol Biol
Transl Sci *107*, 355-415. 10.1016/B978-0-12-385883-2.00005-9.

25. Pilling, A.D., Horiuchi, D., Lively, C.M., and Saxton, W.M. (2006). Kinesin-1 and Dynein are
the primary motors for fast transport of mitochondria in Drosophila motor axons. Molecular
biology of the cell *17*, 2057-2068. 10.1091/mbc.E05-06-0526.

26. Misgeld, T., and Schwarz, T.L. (2017). Mitostasis in Neurons: Maintaining Mitochondria in an
Extended Cellular Architecture. Neuron *96*, 651-666. 10.1016/j.neuron.2017.09.055.

27. Sheng, Z.H. (2014). Mitochondrial trafficking and anchoring in neurons: New insight and
implications. J Cell Biol *204*, 1087-1098. 10.1083/jcb.201312123.

28. Sandoval, H., Yao, C.K., Chen, K., Jaiswal, M., Donti, T., Lin, Y.Q., Bayat, V., Xiong, B.,
Zhang, K., David, G., et al. (2014). Mitochondrial fusion but not fission regulates larval
growth and synaptic development through steroid hormone production. eLife *3*.
10.7554/eLife.03558.

29. Ge, Y., Shi, X., Boopathy, S., McDonald, J., Smith, A.W., and Chao, L.H. (2020). Two forms
of Opa1 cooperate to complete fusion of the mitochondrial inner-membrane. eLife *9*.
10.7554/eLife.50973.

30. Song, Z., Ghochani, M., McCaffery, J.M., Frey, T.G., and Chan, D.C. (2009). Mitofusins and
OPA1 mediate sequential steps in mitochondrial membrane fusion. Molecular biology of the
cell *20*, 3525-3532. 10.1091/mbc.E09-03-0252.

31. Smirnova, E., Griparic, L., Shurland, D.L., and van der Bliek, A.M. (2001). Dynamin-related
protein Drp1 is required for mitochondrial division in mammalian cells. Molecular biology of
the cell *12*, 2245-2256. 10.1091/mbc.12.8.2245.

32. Twig, G., Elorza, A., Molina, A.J., Mohamed, H., Wikstrom, J.D., Walzer, G., Stiles, L.,
Haigh, S.E., Katz, S., Las, G., et al. (2008). Fission and selective fusion govern
mitochondrial segregation and elimination by autophagy. EMBO J *27*, 433-446.
10.1038/sj.emboj.7601963.

33. Breuer, M.E., Koopman, W.J., Koene, S., Nooteboom, M., Rodenburg, R.J., Willems, P.H.,
and Smeitink, J.A. (2013). The role of mitochondrial OXPHOS dysfunction in the
development of neurologic diseases. Neurobiology of disease *51*, 27-34.
10.1016/j.nbd.2012.03.007.

34. Berger, I., Hershkovitz, E., Shaag, A., Edvardson, S., Saada, A., and Elpeleg, O. (2008).
Mitochondrial complex I deficiency caused by a deleterious NDUFA11 mutation. Ann Neurol
63, 405-408. 10.1002/ana.21332.

35. Formosa, L.E., Muellner-Wong, L., Reljic, B., Sharpe, A.J., Jackson, T.D., Beilharz, T.H.,
Stojanovski, D., Lazarou, M., Stroud, D.A., and Ryan, M.T. (2020). Dissecting the Roles of
Mitochondrial Complex I Intermediate Assembly Complex Factors in the Biogenesis of
Complex I. Cell reports *31*, 107541. 10.1016/j.celrep.2020.107541.

36. Guerrero-Castillo, S., Baertling, F., Kownatzki, D., Wessels, H.J., Arnold, S., Brandt, U., and
Nijtmans, L. (2017). The Assembly Pathway of Mitochondrial Respiratory Chain Complex I.
Cell metabolism *25*, 128-139. 10.1016/j.cmet.2016.09.002.

37. Stroud, D.A., Surgenor, E.E., Formosa, L.E., Reljic, B., Frazier, A.E., Dibley, M.G.,
Osellame, L.D., Stait, T., Beilharz, T.H., Thorburn, D.R., et al. (2016). Accessory subunits
are integral for assembly and function of human mitochondrial complex I. Nature *538*, 123-
126. 10.1038/nature19754.

38. Kauffman, M.E., Kauffman, M.K., Traore, K., Zhu, H., Trush, M.A., Jia, Z., and Li, Y.R. (2016). MitoSOX-Based Flow Cytometry for Detecting Mitochondrial ROS. React Oxyg Species (Apex) *2*, 361-370. 10.20455/ros.2016.865.

39. Robinson, K.M., Janes, M.S., Pehar, M., Monette, J.S., Ross, M.F., Hagen, T.M., Murphy, M.P., and Beckman, J.S. (2006). Selective fluorescent imaging of superoxide in vivo using ethidium-based probes. Proc Natl Acad Sci U S A *103*, 15038-15043. 10.1073/pnas.0601945103.

40. Little, A.C., Kovalenko, I., Goo, L.E., Hong, H.S., Kerk, S.A., Yates, J.A., Purohit, V., Lombard, D.B., Merajver, S.D., and Lyssiotis, C.A. (2020). High-content fluorescence imaging with the metabolic flux assay reveals insights into mitochondrial properties and functions. Commun Biol *3*, 271. 10.1038/s42003-020-0988-z.

41. Martelli, F., Hernandes, N.H., Zuo, Z., Wang, J., Wong, C.O., Karagas, N.E., Roessner, U., Rupasinghe, T., Robin, C., Venkatachalam, K., et al. (2022). Low doses of the organic insecticide spinosad trigger lysosomal defects, elevated ROS, lipid dysregulation, and neurodegeneration in flies. eLife *11*. 10.7554/eLife.73812.

42. Martelli, F., Zhongyuan, Z., Wang, J., Wong, C.O., Karagas, N.E., Roessner, U., Rupasinghe, T., Venkatachalam, K., Perry, T., Bellen, H.J., and Batterham, P. (2020). Low doses of the neonicotinoid insecticide imidacloprid induce ROS triggering neurological and metabolic impairments in Drosophila. Proc Natl Acad Sci U S A *117*, 25840-25850. 10.1073/pnas.2011828117.

43. Penugonda, S., Mare, S., Lutz, P., Banks, W.A., and Ercal, N. (2006). Potentiation of lead-induced cell death in PC12 cells by glutamate: protection by N-acetylcysteine amide (NACA), a novel thiol antioxidant. Toxicol Appl Pharmacol *216*, 197-205. 10.1016/j.taap.2006.05.002.

44. Missirlis, F., Hu, J., Kirby, K., Hilliker, A.J., Rouault, T.A., and Phillips, J.P. (2003). Compartment-specific protection of iron-sulfur proteins by superoxide dismutase. J Biol Chem *278*, 47365-47369. 10.1074/jbc.M307700200.

45. Oswald, M.C.W., Garnham, N., Sweeney, S.T., and Landgraf, M. (2018). Regulation of neuronal development and function by ROS. FEBS letters *592*, 679-691. 10.1002/1873-3468.12972.

46. Roos, J., Hummel, T., Ng, N., Klambt, C., and Davis, G.W. (2000). Drosophila Futsch regulates synaptic microtubule organization and is necessary for synaptic growth. Neuron *26*, 371-382.

47. Guo, X., Macleod, G.T., Wellington, A., Hu, F., Panchumarthi, S., Schoenfield, M., Marin, L., Charlton, M.P., Atwood, H.L., and Zinsmaier, K.E. (2005). The GTPase dMiro is required for axonal transport of mitochondria to Drosophila synapses. Neuron *47*, 379-393. 10.1016/j.neuron.2005.06.027.

48. Verstreken, P., Ly, C.V., Venken, K.J., Koh, T.W., Zhou, Y., and Bellen, H.J. (2005). Synaptic mitochondria are critical for mobilization of reserve pool vesicles at Drosophila neuromuscular junctions. Neuron *47*, 365-378. 10.1016/j.neuron.2005.06.018.

49. Milton, V.J., Jarrett, H.E., Gowers, K., Chalak, S., Briggs, L., Robinson, I.M., and Sweeney,
S.T. (2011). Oxidative stress induces overgrowth of the Drosophila neuromuscular junction.
Proc Natl Acad Sci U S A *108*, 17521-17526. 10.1073/pnas.1014511108.

50. Oswald, M.C., Brooks, P.S., Zwart, M.F., Mukherjee, A., West, R.J., Giachello, C.N.,
Morarach, K., Baines, R.A., Sweeney, S.T., and Landgraf, M. (2018). Reactive oxygen
species regulate activity-dependent neuronal plasticity in Drosophila. eLife *7*.
10.7554/eLife.39393.

51. Böhme, M.A., McCarthy, A.W., Grasskamp, A.T., Beuschel, C.B., Goel, P., Jusyte, M.,
Laber, D., Huang, S., Rey, U., Petzoldt, A.G., et al. (2019). Rapid active zone remodeling
consolidates presynaptic potentiation. Nat Commun *10*, 1085. 10.1038/s41467-019-08977-
6.

52. Goel, P., Dufour Bergeron, D., Bohme, M.A., Nunnelly, L., Lehmann, M., Buser, C., Walter,
A.M., Sigrist, S.J., and Dickman, D. (2019). Homeostatic scaling of active zone scaffolds
maintains global synaptic strength. J Cell Biol *218*, 1706-1724. 10.1083/jcb.201807165.

53. Gratz, S.J., Goel, P., Bruckner, J.J., Hernandez, R.X., Khateeb, K., Macleod, G.T., Dickman,
D., and O'Connor-Giles, K.M. (2019). Endogenous Tagging Reveals Differential Regulation
of Ca(2+) Channels at Single Active Zones during Presynaptic Homeostatic Potentiation and
Depression. J Neurosci *39*, 2416-2429. 10.1523/JNEUROSCI.3068-18.2019.

54. Hong, H., Zhao, K., Huang, S., Huang, S., Yao, A., Jiang, Y., Sigrist, S., Zhao, L., and
Zhang, Y.Q. (2020). Structural Remodeling of Active Zones Is Associated with Synaptic
Homeostasis. J Neurosci *40*, 2817-2827. 10.1523/JNEUROSCI.2002-19.2020.

55. Dannhauser, S., Mrestani, A., Gundelach, F., Pauli, M., Komma, F., Kollmannsberger, P.,
Sauer, M., Heckmann, M., and Paul, M.M. (2022). Endogenous tagging of Unc-13 reveals
nanoscale reorganization at active zones during presynaptic homeostatic potentiation.
Frontiers in cellular neuroscience *16*, 1074304. 10.3389/fncel.2022.1074304.

56. Ashrafi, G., de Juan-Sanz, J., Farrell, R.J., and Ryan, T.A. (2020). Molecular Tuning of the
Axonal Mitochondrial Ca(2+) Uniporter Ensures Metabolic Flexibility of Neurotransmission.
Neuron *105*, 678-687 e675. 10.1016/j.neuron.2019.11.020.

57. Datta, S., and Jaiswal, M. (2021). Mitochondrial calcium at the synapse. Mitochondrion *59*,
135-153. 10.1016/j.mito.2021.04.006.

58. Marland, J.R., Hasel, P., Bonnycastle, K., and Cousin, M.A. (2016). Mitochondrial Calcium
Uptake Modulates Synaptic Vesicle Endocytosis in Central Nerve Terminals. J Biol Chem
291, 2080-2086. 10.1074/jbc.M115.686956.

59. Paradis, M., Kucharowski, N., Edwards Faret, G., Maya Palacios, S.J., Meyer, C.,
Stumpges, B., Jamitzky, I., Kalinowski, J., Thiele, C., Bauer, R., et al. (2022). The ER
protein Creld regulates ER-mitochondria contact dynamics and respiratory complex 1
activity. Sci Adv *8*, eabo0155. 10.1126/sciadv.abo0155.

60. Benson, T.E., and Voigt, H.F. (1995). Neuron labeling by extracellular delivery of
horseradish peroxidase in vivo: a method for studying the local circuitry of projection and

interneurons at physiologically characterized sites. J Neurosci Methods *57*, 81-91. 10.1016/0165-0270(94)00131-y.

61. Budnik, V., Koh, Y.H., Guan, B., Hartmann, B., Hough, C., Woods, D., and Gorczyca, M. (1996). Regulation of synapse structure and function by the Drosophila tumor suppressor gene dlg. Neuron *17*, 627-640.

62. Pielage, J., Fetter, R.D., and Davis, G.W. (2006). A postsynaptic spectrin scaffold defines active zone size, spacing, and efficacy at the Drosophila neuromuscular junction. J Cell Biol *175*, 491-503.

63. Chen, K., and Featherstone, D.E. (2005). Discs-large (DLG) is clustered by presynaptic innervation and regulates postsynaptic glutamate receptor subunit composition in Drosophila. BMC Biol *3*, 1. 10.1186/1741-7007-3-1.

64. Accardi, M.V., Daniels, B.A., Brown, P.M., Fritschy, J.M., Tyagarajan, S.K., and Bowie, D. (2014). Mitochondrial reactive oxygen species regulate the strength of inhibitory GABA-mediated synaptic transmission. Nat Commun *5*, 3168. 10.1038/ncomms4168.

65. Doser, R.L., Amberg, G.C., and Hoerndli, F.J. (2020). Reactive Oxygen Species Modulate Activity-Dependent AMPA Receptor Transport in C. elegans. J Neurosci *40*, 7405-7420. 10.1523/JNEUROSCI.0902-20.2020.

66. Peng, J.J., Lin, S.H., Liu, Y.T., Lin, H.C., Li, T.N., and Yao, C.K. (2019). A circuit-dependent ROS feedback loop mediates glutamate excitotoxicity to sculpt the Drosophila motor system. eLife *8*. 10.7554/eLife.47372.

67. Wang, X., Winter, D., Ashrafi, G., Schlehe, J., Wong, Y.L., Selkoe, D., Rice, S., Steen, J., LaVoie, M.J., and Schwarz, T.L. (2011). PINK1 and Parkin target Miro for phosphorylation and degradation to arrest mitochondrial motility. Cell *147*, 893-906. 10.1016/j.cell.2011.10.018.

68. Wang, X., and Schwarz, T.L. (2009). The mechanism of Ca2+ -dependent regulation of kinesin-mediated mitochondrial motility. Cell *136*, 163-174. 10.1016/j.cell.2008.11.046.

69. Clark, I.E., Dodson, M.W., Jiang, C., Cao, J.H., Huh, J.R., Seol, J.H., Yoo, S.J., Hay, B.A., and Guo, M. (2006). Drosophila pink1 is required for mitochondrial function and interacts genetically with parkin. Nature *441*, 1162-1166. 10.1038/nature04779.

70. Park, J., Lee, S.B., Lee, S., Kim, Y., Song, S., Kim, S., Bae, E., Kim, J., Shong, M., Kim, J.M., and Chung, J. (2006). Mitochondrial dysfunction in Drosophila PINK1 mutants is complemented by parkin. Nature *441*, 1157-1161. 10.1038/nature04788.

71. Poole, A.C., Thomas, R.E., Andrews, L.A., McBride, H.M., Whitworth, A.J., and Pallanck, L.J. (2008). The PINK1/Parkin pathway regulates mitochondrial morphology. Proc Natl Acad Sci U S A *105*, 1638-1643. 10.1073/pnas.0709336105.

72. Yang, Y., Gehrke, S., Imai, Y., Huang, Z., Ouyang, Y., Wang, J.W., Yang, L., Beal, M.F., Vogel, H., and Lu, B. (2006). Mitochondrial pathology and muscle and dopaminergic neuron degeneration caused by inactivation of Drosophila Pink1 is rescued by Parkin. Proc Natl Acad Sci U S A *103*, 10793-10798. 10.1073/pnas.0602493103.

1237   73.   Jiang, H.C., Hsu, J.M., Yen, C.P., Chao, C.C., Chen, R.H., and Pan, C.L. (2015). Neural
1238   activity and CaMKII protect mitochondria from fragmentation in aging Caenorhabditis
1239   elegans neurons. Proc Natl Acad Sci U S A *112*, 8768-8773. 10.1073/pnas.1501831112.

1240   74.   Obashi, K., and Okabe, S. (2013). Regulation of mitochondrial dynamics and distribution by
1241   synapse position and neuronal activity in the axon. The European journal of neuroscience
1242   *38*, 2350-2363. 10.1111/ejn.12263.

1243   75.   Lin, M.T., and Beal, M.F. (2006). Mitochondrial dysfunction and oxidative stress in
1244   neurodegenerative diseases. Nature *443*, 787-795. 10.1038/nature05292.

1245   76.   Esterberg, R., Linbo, T., Pickett, S.B., Wu, P., Ou, H.C., Rubel, E.W., and Raible, D.W.
1246   (2016). Mitochondrial calcium uptake underlies ROS generation during aminoglycoside-
1247   induced hair cell death. The Journal of clinical investigation *126*, 3556-3566.
1248   10.1172/JCI84939.

1249   77.   Gorlach, A., Bertram, K., Hudecova, S., and Krizanova, O. (2015). Calcium and ROS: A
1250   mutual interplay. Redox Biol *6*, 260-271. 10.1016/j.redox.2015.08.010.

1251   78.   Yu, T., Robotham, J.L., and Yoon, Y. (2006). Increased production of reactive oxygen
1252   species in hyperglycemic conditions requires dynamic change of mitochondrial morphology.
1253   Proc Natl Acad Sci U S A *103*, 2653-2658. 10.1073/pnas.0511154103.

1254   79.   Jornayvaz, F.R., and Shulman, G.I. (2010). Regulation of mitochondrial biogenesis. Essays
1255   Biochem *47*, 69-84. 10.1042/bse0470069.

1256   80.   Shaw, J.M., and Winge, D.R. (2009). Shaping the mitochondrion: mitochondrial biogenesis,
1257   dynamics and dysfunction. Conference on Mitochondrial Assembly and Dynamics in Health
1258   and Disease. EMBO reports *10*, 1301-1305. 10.1038/embor.2009.247.

1259   81.   Esparza-Molto, P.B., Romero-Carraminana, I., Nunez de Arenas, C., Pereira, M.P., Blanco,
1260   N., Pardo, B., Bates, G.R., Sanchez-Castillo, C., Artuch, R., Murphy, M.P., et al. (2021).
1261   Generation of mitochondrial reactive oxygen species is controlled by ATPase inhibitory
1262   factor 1 and regulates cognition. PLoS Biol *19*, e3001252. 10.1371/journal.pbio.3001252.

1263   82.   Huddleston, A.T., Tang, W., Takeshima, H., Hamilton, S.L., and Klann, E. (2008).
1264   Superoxide-induced potentiation in the hippocampus requires activation of ryanodine
1265   receptor type 3 and ERK. Journal of neurophysiology *99*, 1565-1571.
1266   10.1152/jn.00659.2007.

1267   83.   Lee, K.Y., Chung, K., and Chung, J.M. (2010). Involvement of reactive oxygen species in
1268   long-term potentiation in the spinal cord dorsal horn. Journal of neurophysiology *103*, 382-
1269   391. 10.1152/jn.90906.2008.

1270   84.   Müller, M., Pym, E.C., Tong, A., and Davis, G.W. (2011). Rab3-GAP controls the
1271   progression of synaptic homeostasis at a late stage of vesicle release. Neuron *69*, 749-762.
1272   S0896-6273(11)00073-0 [pii]

10.1016/j.neuron.2011.01.025.

85.   Frank, C.A., Kennedy, M.J., Goold, C.P., Marek, K.W., and Davis, G.W. (2006).
Mechanisms underlying the rapid induction and sustained expression of synaptic
homeostasis. Neuron *52*, 663-677.

86.   Petersen, S.A., Fetter, R.D., Noordermeer, J.N., Goodman, C.S., and DiAntonio, A. (1997).
Genetic analysis of glutamate receptors in Drosophila reveals a retrograde signal regulating
presynaptic transmitter release. Neuron *19*, 1237-1248.

87.   Frank, C.A., Pielage, J., and Davis, G.W. (2009). A presynaptic homeostatic signaling
system composed of the Eph receptor, ephexin, Cdc42, and CaV2.1 calcium channels.
Neuron *61*, 556-569.

88.   Müller, M., and Davis, G.W. (2012). Transsynaptic control of presynaptic Ca(2)(+) influx
achieves homeostatic potentiation of neurotransmitter release. Curr Biol *22*, 1102-1108.
10.1016/j.cub.2012.04.018

S0960-9822(12)00410-1 [pii].

89.   Wang, T., Hauswirth, A.G., Tong, A., Dickman, D.K., and Davis, G.W. (2014). Endostatin Is
a Trans-Synaptic Signal for Homeostatic Synaptic Plasticity. Neuron *83*, 616-629.
10.1016/j.neuron.2014.07.003.

90.   Younger, M.A., Müller, M., Tong, A., Pym, E.C., and Davis, G.W. (2013). A presynaptic
ENaC channel drives homeostatic plasticity. Neuron *79*, 1183-1196.
10.1016/j.neuron.2013.06.048.

91.   Harris, N., Braiser, D.J., Dickman, D.K., Fetter, R.D., Tong, A., and Davis, G.W. (2015). The
Innate Immune Receptor PGRP-LC Controls Presynaptic Homeostatic Plasticity. Neuron *88*,
1157-1164. 10.1016/j.neuron.2015.10.049.

92.   Müller, M., Genç, Ö., and Davis, G.W. (2015). RIM-Binding Protein Links Synaptic
Homeostasis to the Stabilization and Replenishment of High Release Probability Vesicles.
Neuron *85*, 1056-1069. 10.1016/j.neuron.2015.01.024.

93.   Müller, M., Liu, K.S., Sigrist, S.J., and Davis, G.W. (2012). RIM Controls Homeostatic
Plasticity through Modulation of the Readily-Releasable Vesicle Pool. J Neurosci *32*, 16574-
16585. 10.1523/JNEUROSCI.0981-12.2012

32/47/16574 [pii].

94.   Weyhersmüller, A., Hallermann, S., Wagner, N., and Eilers, J. (2011). Rapid active zone
remodeling during synaptic plasticity. J Neurosci *31*, 6041-6052. 31/16/6041 [pii]

10.1523/JNEUROSCI.6698-10.2011.

95.   Goel, P., Dufour Bergeron, D., Bohme, M.A., Nunnelly, L., Lehmann, M., Buser, C., Walter,
A.M., Sigrist, S.J., and Dickman, D. (2019). Homeostatic scaling of active zone scaffolds
maintains global synaptic strength. J Cell Biol. 10.1083/jcb.201807165.

96.   Dhawan, S., Myers, P., Bailey, D.M.D., Ostrovsky, A.D., Evers, J.F., and Landgraf, M.
(2021). Reactive Oxygen Species Mediate Activity-Regulated Dendritic Plasticity Through

1311    NADPH Oxidase and Aquaporin Regulation. Frontiers in cellular neuroscience *15*, 641802.
1312    10.3389/fncel.2021.641802.

1313 97.  Frank, C.A., James, T.D., and Müller, M. (2020). Homeostatic control of Drosophila
1314    neuromuscular junction function. Synapse *74*, e22133. 10.1002/syn.22133.

1315 98.  Goold, C.P., and Davis, G.W. (2007). The BMP ligand Gbb gates the expression of synaptic
1316    homeostasis independent of synaptic growth control. Neuron *56*, 109-123.

1317 99.  Haghighi, A.P., McCabe, B.D., Fetter, R.D., Palmer, J.E., Hom, S., and Goodman, C.S.
1318    (2003). Retrograde control of synaptic transmission by postsynaptic CaMKII at the
1319    Drosophila neuromuscular junction. Neuron *39*, 255-267.

1320 100. McCabe, B.D., Marques, G., Haghighi, A.P., Fetter, R.D., Crotty, M.L., Haerry, T.E.,
1321    Goodman, C.S., and O'Connor, M.B. (2003). The BMP homolog Gbb provides a retrograde
1322    signal that regulates synaptic growth at the Drosophila neuromuscular junction. Neuron *39*,
1323    241-254.

1324 101. Kikuma, K., Li, X., Perry, S., Li, Q., Goel, P., Chen, C., Kim, D., Stavropoulos, N., and
1325    Dickman, D. (2019). Cul3 and insomniac are required for rapid ubiquitination of postsynaptic
1326    targets and retrograde homeostatic signaling. Nat Commun *10*, 2998. 10.1038/s41467-019-
1327    10992-6.

1328 102. Hauswirth, A.G., Ford, K.J., Wang, T., Fetter, R.D., Tong, A., and Davis, G.W. (2018). A
1329    postsynaptic PI3K-cII dependent signaling controller for presynaptic homeostatic plasticity.
1330    eLife *7*. 10.7554/eLife.31535.

1331 103. Orr, B.O., Fetter, R.D., and Davis, G.W. (2017). Retrograde semaphorin-plexin signalling
1332    drives homeostatic synaptic plasticity. Nature *550*, 109-113. 10.1038/nature24017.

1333 104. Goel, P., Li, X., and Dickman, D. (2017). Disparate Postsynaptic Induction Mechanisms
1334    Ultimately Converge to Drive the Retrograde Enhancement of Presynaptic Efficacy. Cell
1335    reports *21*, 2339-2347. 10.1016/j.celrep.2017.10.116.

1336 105. Penney, J., Tsurudome, K., Liao, E.H., Elazzouzi, F., Livingstone, M., Gonzalez, M.,
1337    Sonenberg, N., and Haghighi, A.P. (2012). TOR is required for the retrograde regulation of
1338    synaptic homeostasis at the Drosophila neuromuscular junction. Neuron *74*, 166-178.
1339    S0896-6273(12)00186-9 [pii]

1340 10.1016/j.neuron.2012.01.030.

1341 106. Pielage, J., Fetter, R.D., and Davis, G.W. (2005). Presynaptic spectrin is essential for
1342    synapse stabilization. Curr Biol *15*, 918-928.

1343 107. Pielage, J., Cheng, L., Fetter, R.D., Carlton, P.M., Sedat, J.W., and Davis, G.W. (2008). A
1344    presynaptic giant ankyrin stabilizes the NMJ through regulation of presynaptic microtubules
1345    and transsynaptic cell adhesion. Neuron *58*, 195-209. 10.1016/j.neuron.2008.02.017.

1346 108. Stephan, R., Goellner, B., Moreno, E., Frank, C.A., Hugenschmidt, T., Genoud, C., Aberle,
1347    H., and Pielage, J. (2015). Hierarchical microtubule organization controls axon caliber and
1348    transport and determines synaptic structure and stability. Dev Cell *33*, 5-21.
1349    10.1016/j.devcel.2015.02.003.

109. James, T.D., Zwiefelhofer, D.J., and Frank, C.A. (2019). Maintenance of homeostatic
plasticity at the Drosophila neuromuscular synapse requires continuous IP3-directed
signaling. eLife *8*. 10.7554/eLife.39643.

110. Raut, S., Mallik, B., Parichha, A., Amrutha, V., Sahi, C., and Kumar, V. (2017). RNAi-
Mediated Reverse Genetic Screen Identified Drosophila Chaperones Regulating Eye and
Neuromuscular   Junction   Morphology.   G3   (Bethesda)   *7*,   2023-2038.
10.1534/g3.117.041632.

111. Marrus, S.B., Portman, S.L., Allen, M.J., Moffat, K.G., and DiAntonio, A. (2004). Differential
localization of glutamate receptor subunits at the Drosophila neuromuscular junction. J
Neurosci *24*, 1406-1415.

112. Stewart, B.A., Atwood, H.L., Renger, J.J., Wang, J., and Wu, C.F. (1994). Improved stability
of Drosophila larval neuromuscular preparations in haemolymph-like physiological solutions.
J Comp Physiol A *175*, 179-191. 10.1007/BF00215114.

113. Terriente-Felix, A., Wilson, E.L., and Whitworth, A.J. (2020). Drosophila phosphatidylinositol-
4 kinase fwd promotes mitochondrial fission and can suppress Pink1/parkin phenotypes.
PLoS genetics *16*, e1008844. 10.1371/journal.pgen.1008844.

114. Choudhury, S.D., Mushtaq, Z., Reddy-Alla, S., Balakrishnan, S.S., Thakur, R.S., Krishnan,
K.S., Raghu, P., Ramaswami, M., and Kumar, V. (2016). sigma2-Adaptin Facilitates Basal
Synaptic Transmission and Is Required for Regenerating Endo-Exo Cycling Pool Under
High-Frequency   Nerve   Stimulation   in   Drosophila.   Genetics   *203*,   369-385.
10.1534/genetics.115.183863.

1371

| Gal4-control | *sod2* OE neuron | *ND-20L* neuron |

A, B, C

A′, B′, C′

ND-20L neuron+NACA | sod2 neuron rescue

D, E

D′, E′ merge

HRP
Futsch

F

A Gal4 control   B *sod2* OE neuron   C *ND-20L* RNAi neuron

D *ND-20L* RNAi neuron+NACA   E *sod2* neuron rescue

BRP-fire LUT

F

Normalized BRP Intensity/ μm² area of bouton

***   ***

H Gal4 control   I *ND-20L* RNAi neuron

H′ merge   I′ merge

BRP   GluRIII

G

BRP puncta density (per μm² area of bouton)

***   ***

3.0
2.5
2.0
1.5
1.0
0.5
0.0

Gal4 control sod2 OE ND-20L RNAi ND-20L+NACA sod2 rescue

Neuron

J Gal4 control   K *sod2* OE neuron   L *ND-20L* RNAi neuron   M *sod2* neuron rescue

*

N

mEPSP amplitude (mV)

***   ***   ns

1.5
1.0
0.5
0.0

1 2 3 4

O

EPSP amplitude (mV)

ns   ns   *

1 2 3 4

P

Quantal content

***   *   ns

1 2 3 4

1. Gal4 control
2. *sod2* OE neuron
3. *ND-20L* RNAi neuron
4. *sod2* neuron resue

Q Gal4 control   Mito-GFP HRP   S *ND-20L* RNAi   Mito-GFP HRP

R *sod2* OE   T *sod2* rescue

U

mito clusters/bouton

ns   ***

Gal4 control sod2 OE neuron ND-20L RNAi neuron sod2 neuron rescue

V Control normal ROS   W *ND-20L* loss elevated ROS   X *sod2* rescue normalized ROS

nROS   RRP   Bruchpilot   CaV2   GluR   Ca2+

ns

*

Neuron

Muscle

Evoked potentials normal   Evoked potentials normal   Evoked potentials small

A Glucose → Hexokinase (LDA or 2-Deoxy-D-glucose) → Glucose-6-Phosphate → Glycolysis → Pyruvate → TCA Cycle → ETS

2 ATP directly (Glycolysis)
2 ATP directly (TCA Cycle)

B Control+DMSO

C ns Control+LDA

D ND-20L RNAi+DMSO

E *** ND-20L RNAi+LDA

F Control+ 2-Deoxy-D-glucose

G ** ND-20L RNAi+ 2-Deoxy-D-glucose

H mEPSP amplitude (mV)
ns   ns   **

I mEPSP frequency (Hz)
ns   ns   ns

J EPSP amplitude (mV)
ns   ***   **

Control+DMSO
Control+LDA
ND-20L RNAi+DMSO
ND-20L RNAi+LDA
Control+2-Deoxy-D-glucose
ND-20L RNAi+2-Deoxy-D-glucose

K Control+DMSO
L ND-20L RNAi+DMSO
M Control+LDA
N ND-20L RNAi+LDA
O Control+2-Deoxy-D-glu
P ND-20L+2-Deoxy-D-glu

BRP-Fire-Lut

Q BRP relative Intensity (A.U.) μm² area of bouton
***   **   ***

Control+DMSO
ND-20L RNAi+DMSO
Control+LDA
ND-20L RNAi+LDA
Control+2-Deoxy-D-glucose
ND-20L RNAi+2-Deoxy-D-glucose

A Muscle-Gal4
B Muscle *ND-20L* RNAi
C Muscle UAS-*sod2* OE
D Muscle *ND-20L*+NACA
E Muscle *ND-20L*+*sod2* OE

GluRIII
BRP

F

# of missing GluRIII apposed puncta per bouton

*** ***

G Muscle-Gal4
H Muscle *ND-20L* RNAi
I Muscle *ND-20L*+NACA
J Muscle *ND-20L*+*sod2* OE

GluRIIA
BRP

K

# of missing GluRIIA apposed puncta per bouton

*** ***

Gal4 Control
ND-20L RNAi
UAS-*sod2* OE
ND-20L+NACA
ND-20L+*sod2* OE

A. Muscle Gal4 | Muscle *ND-20L* RNAi | *cat* muscle not rescue | *sod1* muscle not rescue | *sod2* muscle rescue

B. Muscle-Gal4+10% EtOH | Muscle *ND-20L* RNAi+10% EtOH | Muscle *ND-20L* RNAi+0.5mM curcumin | Muscle *ND-20L* RNAi+0.5mM NACA

C. *ND-30* mutant | *ND-30* mutant + 0.5mM NACA | *ND-30* mutant *sod2* muscle rescue

D. Muscle-Gal4+0.5% DMSO | Muscle-Gal4 (50µM rotenone) | Muscle *cat* OE (50µM rotenone) | Muscle *sod1* OE (50µM rotenone) | Muscle *sod2* OE (50µM rotenone)

E. mEPSP amplitude (mV)
F. EPSP amplitude (mV)
G. Quantal content

a. Muscle Gal4
b. Muscle *ND-20L* RNAi
c. *cat* muscle rescue
d. *sod1* muscle not rescue
e. *sod2* muscle not rescue
f. Muscle Gal4+10% EtOH
g. Muscle *ND-20L* RNAi+10% EtOH
h. Muscle *ND-20L* RNAi+0.5mM curcumin
i. Muscle *ND-20L* RNAi+0.5mM NACA
j. *ND-30* mutant
k. *ND-30* mutant+0.5mM NACA
l. *ND-30* mutant *sod2* muscle rescue
m. Muscle Gal4+0.5% DMSO
n. Muscle Gal4+50µM rotenone
o. Muscle *cat* OE+50µM rotenone
p. Muscle *sod1* OE+50µM rotenone
q. Muscle *sod2* OE+50µM rotenone

H. Distance crawled in cm

1. Muscle Gal4
2. Muscle *sod2* OE
3. Muscle *ND-30* RNAi
4. Muscle *ND-20L* RNAi
5. Muscle *sod2* rescue
6. Muscle *ND-20L* RNAi+NACA
7. Pan neuron Gal4
8. Pan neuron *ND-20L* RNAi
9. *ND-30* mutant
10. *ND-30* mutant *sod2* muscle rescue