Mitochondrial Complex I and ROS control synapse function through opposing pre- and postsynaptic mechanisms

Bhagaban Mallik[1], Sajad A. Bhat[2], Xinnan Wang[2], C. Andrew Frank[1,3]

[1]Department of Anatomy and Cell Biology, University of Iowa Carver College of Medicine, Iowa City, IA, 52242, USA

[2]Department of Neurosurgery, Stanford University School of Medicine, Stanford, CA, 94305, USA

[3]to whom correspondence should be addressed

andy-frank@uiowa.edu, @CAndrewFrank

ABSTRACT

Neurons require high amounts energy, and mitochondria help to fulfill this requirement. Dysfunctional mitochondria trigger problems in various neuronal tasks. Using the *Drosophila* neuromuscular junction (NMJ) as a model synapse, we previously reported that Mitochondrial Complex I (MCI) subunits were required for maintaining NMJ function and growth. Here we report tissue-specific adaptations at the NMJ when MCI is depleted. In *Drosophila* motor neurons, MCI depletion causes profound cytological defects and increased mitochondrial reactive oxygen species (ROS). But instead of diminishing synapse function, neuronal ROS triggers a homeostatic signaling process that maintains normal NMJ excitation. We identify molecules mediating this compensatory response. MCI depletion in muscles also enhances local ROS. But high levels of muscle ROS cause destructive responses: synapse degeneration, mitochondrial fragmentation, and impaired neurotransmission. In humans, mutations affecting MCI subunits cause severe neurological and neuromuscular diseases. The tissue-level effects that we describe in the *Drosophila* system are potentially relevant to forms of mitochondrial pathogenesis.

Keywords: homeostatic plasticity; mitochondria; Mitochondrial Complex I; *Drosophila*; *NDUFS7*; Mito-GFP; rotenone; NACA; *Sod2*; ROS

INTRODUCTION

Neurons have vast energy needs. These needs are primarily satisfied by healthy pools of mitochondria [1,2]. Mitochondria generate energy through the action of the ATP synthase complex in the electron transport chain [3,4]. They also perform complementary functions, including maintaining calcium homeostasis [5,6], promoting cell survival [7], triggering reactive oxygen species (ROS) signaling [8], stimulating lipid synthesis [9], and regulating innate immunity [10]. For energy-driven neurons, it is thought that the primary role of mitochondria is to provide ATP. It is less understood how other mitochondrial functions contribute to the regulation of normal neurophysiology. It is also not well understood how neural tissues or synaptic sites cope when they are challenged with a loss of mitochondria. Genetic models can help to address these puzzles.

Mitochondrial Complex I (MCI) (NADH ubiquinone oxidoreductase) is an essential part of the electron transport chain and ATP production, and it comprises dozens of distinct subunits. Much of our understanding about MCI derives from systemic analyses of its assembly. Studies have been performed on Complex I components from diverse organisms, including *Neurospora crassa* and *Drosophila melanogaster*. Those studies demonstrate that discrete MCI subunits are ancient; indeed, there are few differences between these MCI models from simple organisms and the corresponding human and bovine orthologs [11-14]. For *Drosophila melanogaster*, MCI consists of at least 42 distinct subunits; the 14 core MCI subunits are present, as are at least 28 accessory subunits [12].

In humans, MCI dysfunction has been linked to diseases such as Leigh syndrome, mitochondrial myopathy, and encephalomyopathy, as well as forms of stroke [15-18]. On a cellular level, MCI dysfunction can cause the demise of neurons and muscles; these phenotypes are typically attributed to defects in ATP production [19,20]. However, in addition to the ATP production defects, mutations affecting MCI subunit components are also associated with excess mitochondrial ROS. Normally, ROS accumulation can be neutralized by the cellular antioxidant system [21]. But if that system becomes overwhelmed, there can be consequences for cells and organ systems – including progressive neurodegeneration and seizures for the nervous system [22-24]. On the level of synapses, it is possible that MCI loss triggers severe molecular consequences, and it is also possible that excess ROS plays a role.

For a previous paper, we depleted MCI function at the *Drosophila* neuromuscular junction (NMJ). Our data suggested fundamental synaptic functions for MCI [13]. Here we expand upon that work, mostly taking advantage of RNAi-mediated depletion of the *Drosophila* homologs of human NDUFS7. We also scrutinize loss-of-function mutants of other MCI subunits and pharmacological inhibition of MCI. Our collective data show that MCI depletion causes *Drosophila* phenotypes remi-

niscent of mitochondrial diseases, such as progressive degeneration of muscle and presynaptic cy-
toskeleton, excess ROS production, loss of mitochondria and alteration in mitochondrial morpholo-
gy.

On single-tissue levels, we were surprised to find that there were opposite effects on syn-
apse activity in the presynaptic motor neurons vs. the postsynaptic muscles. MCI dysfunction in
Drosophila motor neurons causes profound cytological phenotypes, but there are no significant
functional phenotypes. This appears to be because neuronal mitochondrial ROS triggers an adap-
tive response, demonstrated visually by active zone enhancement. This ROS-driven enhancement
of active zones occurs through at least two processes, 1) regulation of calcium flux from intracellular
stores (ER) and mitochondria; and 2) use of glycolysis as an alternative energy source. By contrast,
postsynaptic depletion of MCI and the associated elevation of muscle ROS triggers a destructive
response: disruption of NMJ morphology and the Dlg-Spectrin scaffold that is critical for normal ac-
tive zone-receptor apposition. To our knowledge, these cellular and molecular mechanisms of MCI
deficiency have not previously been elucidated at synapse-specific or tissue-specific levels.

RESULTS

One RNAi transgene targets two *Drosophila* homologs of a core MCI subunit, NDUFS7
We previously reported impairments in neuromuscular junction (NMJ) synapse development and
function when Mitochondrial Complex I (MCI) was depleted [13]. We observed robust synaptic pheno-
types when driving a transgenic RNAi line against an MCI subunit, termed *UAS-ND-20L[RNAi]* (*ND-*
20L^HMJ23777) [13]. That RNAi line encodes a short hairpin that matches 21 consecutive nucleotides of
the *ND-20L* gene. Recent work indicates that *ND-20L* is only sparsely expressed in *Drosophila* tis-
sues [25-27], but the same short hairpin also matches 19 consecutive nucleotides of the *ND-20* gene,
and *ND-20* is ubiquitously expressed [25-27]. *ND-20* and *ND-20L* encode *Drosophila* homologs of the
core NDUFS7 MCI subunit. Because both genes are effectively "on-target" for the RNAi transgene,
we term the *UAS-ND-20L[RNAi]* transgene as *UAS-NDUFS7[RNAi]* for this study.

We used the GAL4/UAS system to verify that *UAS-NDUFS7[RNAi]* can diminish MCI func-
tion in relevant tissues. We drove its expression using neuronal or muscle GAL4 drivers. By quanti-
tative RT-PCR, message levels of both NDUFS7-encoding genes were significantly diminished in
those tissues in larvae (Fig. S1A-D). *ND-20L* message levels were down in both neurons and mus-
cle, but variable in neurons, likely due to the low endogenous expression of *ND-20L* (Fig. S1A-B).
ND-20 message levels were significantly knocked down by the transgene in both tissues (Fig. S1C-
D). We also tested if the transgene could diminish mitochondrial respiration by a Seahorse assay.

To acquire sufficient material for the assay, we collected mitochondria from adult muscles (see

Methods). Driving the *UAS-NDUFS7[RNAi]* transgene in adult muscle starkly reduced oxygen con-

sumption rate in muscle mitochondria across all timepoints tested (Fig. S1E-F). These assays con-

firmed that *UAS-NDUFS7[RNAi]* targets MCI.

Depletion of MCI affects mitochondrial integrity in multiple *Drosophila* synaptic tissues

We wanted to understand what was happening to mitochondria to affect synapse function when

MCI was depleted. We started by examining mitochondria by microscopy. To visualize them, we

expressed a *UAS-Mito-GFP* transgene [28] in *Drosophila* tissues. Concurrently, we used tissue-

specific GAL4 drivers alone (as controls) or GAL4 drivers + *UAS-NDUFS7[RNAi]* to deplete MCI

function [13]. With these tools, we made qualitative observations of mitochondrial morphology (Fig. 1),

and we quantified those observations in subsequent analyses.

In motor neurons, the Mito-GFP signal localized to the neuropil of the ventral nerve cord

(Fig.1: A-A′, B-B′). In control neurons, the neuropil mitochondria had a filamentous appearance. By

contrast, NDUFS7-depleted neurons had punctate and clustered mitochondria. (Fig. 1E-E′, F-F′).

We examined mitochondria in the motor axons that innervate proximal and distal NMJs (Fig. 1C-C′,

D-D′, G-G', H-H'). The proximal segment A2 axons had abundant mitochondria in all cases (Fig.

1C-C', G-G'). However, for the distal segment A5 axons, NDUFS7 depletion elicited an obvious de-

crease in mitochondria number (Fig. 1D-D,′ H-H,′). This A2 vs. A5 discrepancy was consistent with

prior work by others examining defects in mitochondrial trafficking dynamics: distal sites can show

phenotypes more prominently [29,30].

We hypothesized that fewer mitochondria in the A5 axon might correlate with a neurotrans-

mission defect at the NMJ. Yet by NMJ electrophysiology, we found no significant differences in the

evoked amplitude compared to the control NMJs in the distal segment A5 (Fig. 1I-K). These data

matched our prior examination of MCI at the A2 and A3 segments of the NMJ, where neuronal im-

pairment of MCI was not sufficient on its own to reduce evoked NMJ neurotransmission [13].

In muscle, we observed an array of phenotypes. As with neurons, there were clustered mi-

tochondria when *NDUFS7* gene function was depleted (Fig. 1L-M). Additionally, there was a tissue-

level phenotype: *NDUFS7*-depleted muscles were developed, but they looked disorganized and

fragmented, with oblong-shaped nuclei in the muscle syncytia (Fig. 1L-M). This phenotype could

explain why we previously observed that muscle impairment of MCI was sufficient to reduce evoked

NMJ neurotransmission [13].

To examine the mitochondria at presynaptic NMJ release sites, we used the motor neuron

GAL4 driver to label NMJ boutons with Mito-GFP. For image analysis, we marked the presynaptic

membrane boutons with anti-HRP immunostaining. Control NMJs contained abundant and large
clusters of mitochondria in synaptic boutons, but by comparison, *NDUFS7[RNAi]* boutons contained
small clusters and few mitochondria (Fig. 1N-P). We measured the mitochondrial volume in a 3-D
stack and compared it to the synaptic volume (Fig. 1Q-S). The Mito-GFP signal occupied a sizeable
proportion of the bouton volume in controls (~30%), but this value was significantly diminished in
NDUFS7-depleted animals (~10%) (Fig. 1S). Collectively, our data suggest that the depletion of
NDUFS7 by RNAi leads to abnormal mitochondrial clustering in the neuronal cell body and muscle
– as well as losses of distal axon and synaptic mitochondria.
142

Loss of MCI phenocopies loss of Mitofusin

The cell-level NDUFS7-depletion phenotypes were reminiscent of *Drosophila* mutants impairing mi-
tochondrial dynamics [29,31]. Therefore, we re-examined MCI-depleted mitochondria, this time addi-
tionally impairing genes known to mediate mitochondrial fusion and fission. Mitofusin 1 (Mfn1) and
Mitofusin 2 (Mfn2) are GTPases that regulate outer mitochondrial membrane fusion [32,33]. The *Dro-*
sophila gene encoding the Mitofusin homolog is called *marf*. Dynamin-related protein 1 is a GTPase
that regulates mitochondrial fission. In *Drosophila*, this factor is encoded by the gene *drp1* [34]. Previ-
ous work reported that defective fusion results in fragmented mitochondria, while defective fission
can lead to enlarged mitochondria [35]. We used RNAi-mediated knockdown constructs for each of
these genes.
As before, we observed that wild-type motor neurons had filamentous and oval mitochon-
dria, while NDUFS7-depleted neurons had fewer and smaller clustered mitochondria in the ventral
nerve cord (VNC) (Fig. 2A) and axons (Fig. 2B). Knockdown of the fusion gene *marf* phenocopied
NDUFS7 loss, revealing small mitochondria in motor neurons, while knockdown of the fission gene
drp1 yielded filamentous mitochondria (Fig. 2A). Simultaneously depleting motor neurons of *marf*
and *NDUFS7* by RNAi did not show any additive defect in mitochondrial appearance in the ventral
nerve cord (VNC) and axons (Figs. 2A, B). This result could mean that the genes share a common
process to regulate mitochondrial fusion. By contrast, depleting *drp1* and *NDUFS7* by RNAi simul-
taneously yielded punctate mitochondria. This result likely means that that the punctate *NDUFS7*
mitochondrial phenotypes (potential fusion phenotypes) are epistatic to *drp1* loss (Fig. 2A, B).
We measured mitochondrial branch length from skeletonized images of the mitochondria
(Skeletonize3D, ImageJ plugin, data in Tables 1, 2, 3). Control and *drp1* knockdown showed normal
mitochondrial branch length, but knockdown using *NDUFS7[RNAi]* or *marf[RNAi]* – or knockdowns
using combinations of each – exhibited short branch length (Fig. 2C, Table 1).

To quantify mitochondria in axons, we counted Mito-GFP positive puncta in distal A5 motor axons labeled by anti-GFP. Control axons and *drp1*-depleted axons contained abundant mitochondria (Fig. 2D, Table 1). By contrast, any gene manipulation or combination targeting *NDUFS7* or *marf* by RNAi resulted in diminished numbers of mitochondrial clusters (Fig. 2D, Table 1).

We extended the analysis to NMJ terminals. We counted Mito-GFP clusters in presynaptic boutons apposed by postsynaptic densities, labeled by anti-Discs Large 1 (Dlg1) (Figs. 2E-M, Table 4). The results matched our earlier observations (Figs. 1N-S). Control NMJs and *drp1*-depleted NMJs contained numerous mitochondrial clusters per bouton (Figs. 2E, M). However, NDUFS7-depleted boutons contained few mitochondria, and this was phenocopied by *marf[RNAi]* (Fig. 2E-M, Table 4). Collectively, these results suggest that *Drosophila* NDUFS7 (and hence MCI) contributes to normal mitochondrial fusion, likely in conjunction with the Mitofusin homolog, Marf.

Mitochondrial Reactive Oxygen Species Contribute to Synaptic Phenotypes
Several studies have demonstrated that Complex I loss results in high levels of mitochondrial reactive oxygen species (ROS) [36-40]. This means that excess ROS could be contributing to the cytological and mitochondrial fusion phenotypes that we have described.

We checked if we could observe mitochondrial ROS (superoxide) in living *Drosophila* tissue and if ROS levels corresponded to Complex I function (Fig. S1). We used a commercially available fluorescent mitochondrial superoxide indicator, MitoSOX (ThermoFisher, Methods) [41-43]. With MitoSOX, we observed mitochondrial superoxide in many tissues. There was a baseline level of ROS in controls (Fig. S2A, E, I, J, Table 5), and the level was greatly increased in NDUFS7-deficient motor neuron cell bodies and muscle (Fig. S2B, F, I; J, Table 5).

Next, we tested if ROS scavengers could reverse the high MitoSOX fluorescence levels in *NDUFS7*-deficient tissues. We fed a pharmacological scavenger, N-Acetyl Cysteine Amide (NACA) [44-46] to *Drosophila* larvae (Methods). We also used a transgene, *UAS-Sod2* [47], to express a superoxide dismutase enzyme. Both successfully diminished the high levels of mitochondrial ROS that resulted from *NDUFS7* depletion at the NMJ, and both worked in muscle and neurons (Fig. S2C, D, G-J, Table 5). We also tested a related idea: if genetic *Sod2* knockdown by RNAi could phenocopy loss of MCI by enhancing Mito-SOX levels in these same tissues. It did. Transgenic *UAS-Sod2[RNAi]* enhanced Mito-SOX signal in all tissues and sub compartments, including the ventral nerve cord (Fig. S3A-C, Table 6), the motor neuron axons (Fig. S3D-F, Table 6), the synaptic boutons (Fig. S3G-I, Table 6), and the muscle (Fig. S3J-L, Table 6).

Next, we tested if these same ROS scavengers could reverse mitochondrial phenotypes caused by loss of MCI. Co-expressing *UAS-Sod2* or rearing larvae with NACA suppressed the mi-

tochondrial morphology defects in the ventral nerve cord of *NDUFS7*-depleted animals; it also re-
stored axonal loss of mitochondria (Fig S4A-D, Table 2). To test an additional MCI manipulation, we
knocked down *Drosophila ND-30* (homologous to human *NDUFS3*) in motor neurons. As with
NDUFS7, depleting *ND-30* in motor neurons yielded punctate mitochondria in the ventral nerve
cord, but the addition of *UAS-Sod2* restored a wild-type, filamentous mitochondrial morphology (Fig.
S4A, C). Similarly, loss of *ND-30* gene function in neurons depleted A5 axonal mitochondria and
this phenotype was also reversed by *UAS-Sod2* transgenic expression (Fig. S4B, D, Table 2). But
in contrast to *UAS-Sod2*, neither *UAS-Sod1* nor *UAS-Catalase* worked to reverse mitochondrial
morphology defects (Fig. S5, Table 3). SOD1 (cytosol) and Catalase (peroxisome) localize to differ-
ent compartments than SOD2 (mitochondrial matrix). Therefore, these results could indicate that a
scavenger needs to access the proper compartment for rescue.

Because of the links between MCI and mitochondrial fusion, we considered whether a *marf*
loss of function could also yield high levels of neuronal ROS (Fig. S6). It did – both *marf* and
NDUFS7 loss-of-function conditions showed high levels of mitochondrial superoxide in motor neu-
ron cell bodies (Fig. S6A-D, A'-D', M, Table 5) in motor axons (Fig. S6E-H, E'-H', N, Table 5); and at
NMJ sites (Fig. S6I-L, I'-L', O, Table 5). These ROS phenotypes were not confined to genetic ma-
nipulations. We made similar observations when MCI was impaired pharmacologically by feeding
rotenone to developing larvae (Fig. S6, Table 5). In the case of rotenone, the amount of mitochon-
drial ROS in the tissues was high, but it was generally not increased as much as with the genetic
manipulations (Fig. S6M-O, Table 5).

Finally, because the distal A5 motor axons accumulated high levels of ROS when subjected
to these insults, we examined them for synaptic vesicle trafficking defects. We immunostained for
Cysteine String Protein (CSP), a DNAJ-like co-chaperone and synaptic vesicle-associated protein
(Cysteine string protein: CSP). *NDUFS7* depletion caused aberrant accumulation of CSP in the A5
motor axons; and this defect was suppressed by motor neuron transgene overexpression of *UAS-*
Sod2 or by feeding animals with NACA (Fig. S7A-L, Table 7). However, this phenotype was not
suppressed by motor neuron overexpression of *UAS-Sod1* or *UAS-Catalase* (Fig. S8, Table 8).

ROS scavengers did not reverse all mitochondrial abnormalities. Expressing *UAS-Sod2* in
the *UAS-NDUFS7[RNAi]* or *UAS-marf[RNAi]*-depletion backgrounds did not restore mitochondrial
clusters to motor neuron terminals (Fig. 2E-M, Table 2). For the remainder of the study, we used
scavengers as complementary tools to test which MCI-loss phenotypes were likely due to mito-
chondrial ROS.

Loss of MCI subunits impairs synaptic cytoskeletal stability

ROS can modulate the cytoskeleton, either through redox modification of cytoskeletal proteins or by altering pathways that regulate cytoskeletal organization [48]. To test whether the mitochondrial defects and abnormal accumulation of ROS were associated with the altered synaptic cytoskeleton, we labeled synaptic boutons with an anti-Futsch antibody (Fig. 3). Futsch is a *Drosophila* MAP1B homolog that associates with microtubules [49].

In motor neuron Gal4-control and *UAS-Sod2* overexpression larvae, Futsch organized in periodic loops, as expected from previous characterizations [49] (Fig. 3A-B, Table 7). But in NDUFS7-depleted larvae, the anti-Futsch staining showed a significant reduction in microtubule loops (Figure 3C, Table 7). These *NDUFS7* phenotypes were suppressed by motor neuron overexpression of *UAS-Sod2* or by raising animals on food containing 0.5 mM NACA (Fig. 3D-E, Table 7). However, they were not significantly restored by neuronal overexpression of *UAS-Sod1* or *UAS-Catalase* (Fig. S9, Table 8). These data indicate that loss of MCI regulates cytoskeletal architecture due to excessive accumulation of mitochondrial ROS in neurons.

Modest neurotransmission phenotypes after motor neuron-specific loss of MCI or Marf

Given the cytological phenotypes after neuronal MCI loss, it was puzzling that there seemed to be little to no electrophysiological consequence at NMJs ([13] and Fig. 1J). We probed this finding, this time depleting motor neurons of *marf* and/or *NDUFS7* gene function. We recorded spontaneous miniature postsynaptic potentials (mEPSP) and evoked excitatory postsynaptic potentials (EPSP).

Phenotypes were normal-to-mild (Fig. S10). For both *NDUFS7[RNAi]* and *marf[RNAi]*, there were small, but statistically significant decreases in mEPSP amplitude (Fig. S10A-C, G, Table 9). But for *NDUFS7[RNAi]*, evoked events (EPSP) and calculated quantal content (QC) were at control levels (Fig. S10A-B, H-I, Table 9). For neuronal *marf[RNAi]*, those measures were near-normal, with a slight decrease in EPSP amplitude (Fig. S10C, H, Table 9) and a slight increase in QC (Fig. S10I, Table 9). Finally, for a double *NDUFS7[RNAi]* + *marf[RNAi]* knockdown condition in neurons, we observed a small, but statistically significant decrease in EPSP amplitude (Fig. S10E, H, Table 9).

Synaptic phenotypes caused by mitochondrial dysfunction might be masked until synapses are challenged with extreme conditions, like high frequency stimulation [50,51]. Therefore, we challenged neuronally *NDUFS7*-depleted *Drosophila* NMJs in several ways (Fig. S11). First, we lowered recording saline $[Ca^{2+}]$ to 0.15 mM, which is roughly one order of magnitude lower than physiological calcium. In low calcium, the motor neuron-driven *NDUFS7[RNAi]* NMJs had slightly smaller evoked potentials compared to driver controls, but the numerical reduction was not statistically significant (Fig. S11A-C, Table 10). Next, we lowered extracellular $[Ca^{2+}]$ even further, to 0.1 mM,

which yielded a mix of successful EPSP firing events and failures. Failure analyses revealed an in-
crease in failure rate at the neuronally depleted *NDUFS7[RNAi]* NMJs compared to control animals
(Fig S11D, Table 10), demonstrating a decreased probability of release. Failure rates in 0.1 mM
calcium were restored to baseline levels when *NDUFS7*-depleted larvae were raised in a media
containing NACA or genetically expressing *Sod2* in motor neurons (Fig S11D, Table 10), indicating
that a sensitivity to low calcium could be related to ROS levels.

Finally, we checked if forms of short-term neuroplasticity were affected by *NDUFS7* loss in
motor neurons. For two different extracellular $[Ca^{2+}]$ conditions (0.4 to 1.5 mM), we did not observe
any significant changes in paired-pulse ratios (Figs. S11E-J, Table 10). Likewise, we did not note
depreciation of evoked neurotransmission over the course of high frequency stimulus trains in high
calcium (Figs. S11K-P). Collectively, these data suggest that there might be a small effect on NMJ
physiology due to defective mitochondrial fusion, and the defect could be sensitive to low levels of
extracellular calcium – but the aggregate data also indicate that neuronal mitochondrial defects
alone do not drastically affect NMJ neurotransmission.

Loss of MCI in neurons controls the level and distribution of the active zone to stabilize syn-
aptic strength
We wondered how synapse might evade severe dysfunction, despite loss of mitochondria in the
motor neurons. One possibility is that *NDUFS7* loss/MCI impairment could trigger a form of func-
tional homeostatic compensation of the NMJ. Another idea is that the mitochondrial ATP generated
is superfluous at the NMJ – and that any energy-intensive functions that mitochondria support could
be redundantly covered by glycolysis. These models are not mutually exclusive, and for any scenar-
io, mitochondrial ROS downstream of defective MCI could be a candidate signal. Recent findings
have demonstrated that ROS intermediates, mitochondrial distribution, and mitochondrial trafficking
all affect development of the *Drosophila* NMJ [48,52,53].

We imaged the presynaptic active zone apparatus in neuronally depleted *NDUFS7[RNAi]*
flies. Third-instar larval active zones showed a decrease in BRP puncta density per unit area in
NDUFS7-depeleted NMJs compared to control NMJs (Fig.4A-C, G, H-I, Table 11). But they also
showed a robust enhancement phenotype: in *NDUFS7*-depleted animals, we found a 40% increase
in active zone (BRP) immunofluorescence signal per unit area, compared to control (Fig. 4A-C, F,
H-I, Table 11), by laser scanning confocal microscopy. This result was intriguing because NMJ ac-
tive zone enhancements (or changes in active zone sub-structure) have been proposed by other
labs to be molecular correlates of forms of homeostatic plasticity and potentiation of neurotransmit-

ter release [54-58]. This raised the possibility that the NMJs evade severe dysfunction through a form
of synaptic homeostasis.

As an independent test, we impaired MCI pharmacologically. To do this, we raised larvae on
50 µM rotenone-spiked food; and we also incubated wild-type fillet preparations with 500 µM of ro-
tenone for extended time. For both cases, we observed significant increases in BRP protein at the
presynaptic active zones (Fig S12A-U, Table 12). For the extended incubation, the fillet prepara-
tions required sufficient rotenone incubation time (six hours) and an intact motor nerve to show the
active zone enhancement (Fig. S12U, Table 12). This result suggested that downstream of MCI de-
pletion, a compensatory delivery of active zone material required either substantial trafficking time
and/or fully intact neuroanatomy.

Next, we checked if the enhanced active zone signal by was triggered by excess mitochon-
drial ROS in motor neurons. Indeed, we found that the *NDUFS7*-depletion active zone enhance-
ments were fully reversed by ROS scavengers, either by raising larvae in food containing NACA or
by neuronally expressing *UAS-Sod2* (Fig. 4D-G, Table 11). However, they were not reversed by
overexpressing *UAS-Sod1* or *UAS-Catalase* (Fig. S13A-G, Table 13).

Finally, we assessed synapse function. As with our prior recordings, evoked postsynaptic
potentials at the NMJ were not significantly changed by *NDUFS7* depletion in motor neurons. But
interestingly, scavenging mitochondrial ROS in the *NDUFS7[RNAi]* neuronal depletion background
with *UAS-Sod2* unmasked a small deficit in NMJ excitation, compared to controls (Fig. 4J-P, Table
11). As before, neither *UAS-Sod1* overexpression nor *UAS-Catalase* overexpression unmasked this
deficit in NMJ excitation (Fig. S13N-P, Table 13).

These data could mean that mitochondrial ROS is helping to maintain synaptic activity. Neu-
ronal expression of *UAS-Sod2* did not restore mitochondrial clusters to the NMJ after *NDUFS7*
gene function depletion (Fig. 4Q-U, Table 11; like Fig. 2H), meaning that the synaptic sites were still
deficient in mitochondria. And as expected, neuronal expression of *UAS-Sod1* and *UAS-Catalase*
also failed to mitochondrial clusters to the NMJ (Fig. S13Q-W, Table 13). Collectively, our data sup-
port a model in which neuronal ROS (nROS) triggers active zone enhancement and functional
compensation when MCI is limiting (Fig. 4V-X, Table 11).

Neuronal MCI subunits stabilize synaptic strength in conjunction with intracellular calcium
signaling proteins
Recent work described a mechanism for local calcium uptake into mitochondria that drives ATP
production to maintain synaptic function [59]. The process is governed by the mitochondrial calcium

uniporter (MCU) and its accessory EF-hand MICU proteins [59]. Beyond this role for mitochondrial

calcium, there are also known roles for core synaptic functions like vesicle cycling [60,61].

To test if mitochondrial or neuronal calcium could be involved in maintaining synapse func-

tion at the NMJ, we acquired genetic reagents to examine depressed MCU function in conjunction

with depressed MCI. We also used pharmacological reagents to inhibit release of intracellular

sources of calcium, like those from the Ryanodine Receptor (RyR) and the IP_3 receptor (IP_3R) of the

endoplasmic reticulum (ER) [62], as well as a genetic reagent that we previously used at the NMJ to

deplete IP_3 signaling (*UAS-IP$_3$-sponge*) (Fig. 5A-D). For this set of experiments, we used the

NDUFS7 neuronal knockdown condition as a sensitized genetic background (Fig. 5E-T, Table 14).

We also used a pan-neuronal driver (*elaV-Gal4*) for ease *Drosophila* genetic stock construction in-

stead of a motor neuronal driver. We previously observed similar NMJ effects either for motor neu-

ronal or pan-neuronal depletion of MCI [13].

We observed no significant differences in EPSP amplitudes when we impaired *mcu* function

neuronally (Table 14). Similarly, we did not observe deficits in baseline synaptic activity by blocking

RyR and IP_3R, alone or in conjunction with *NDUFS7[RNAi]* (Fig. 5E-H, Q-T, Table 14). However,

when we concurrently impaired a combination of *NDUFS7*, *mcu*, and ER calcium store channels,

we observed marked decreases in evoked amplitude (Fig. 5K-L, S, Table 14). These results are

consistent with a model in which mitochondrial calcium uptake, MCU activation, and ER (store) cal-

cium efflux combine to stabilize synaptic strength.

If this idea were correct, then it should also be possible to chelate cytoplasmic calcium in a

neuronal *NDUFS7[RNAi]* background and reveal neurotransmission defects. Direct application of

the membrane-permeable chelator BAPTA-AM, followed by a wash to remove chelator residing in

the saline, had no significant effect on baseline neurotransmission parameters (DMSO carrier +

wash). But in the *NDUFS7[RNAi]* background, BAPTA-AM + wash significantly diminished evoked

potentials, compared to mock-treated (DMSO + wash) NMJs. (Fig. 5 M-T, Table 14).

To check if these effects on neurotransmission correlated with effects on active zone protein

accumulation, we also conducted anti-Brp immunostaining experiments (Fig. S14). As before, neu-

ronal knockdown of *NDUFS7* gene function triggered a marked, compensatory increase in presyn-

aptic active zone material that was readily apparent by confocal microscopy (Figs. S14A, C, I, Table

15). But this increase was reversed when combined with *mcu* gene function knockdown and phar-

macological blockade of store calcium release channels (Figs. S14G-I, Table 15). Together, our da-

ta indicate that loss of MCI subunits in neurons sensitizes synapses to decreases in intracellular

calcium.

A combination of mitochondria, glycolysis, and the TCA Cycle stabilizes NMJ function

Neuronal calcium handling and MCU play roles in NMJ stability that are uncovered by loss of MCI at the NMJ. Downstream of calcium handling, a logical hypothesis is that synaptic energy would play a role [59]. If this were the case, would mitochondria be the sole energy source? Alternatively, in the absence of full mitochondrial function, could glycolysis theoretically substitute, as a homeostatic (or redundant) means for staving off synapse dysfunction?

We tested these ideas by limiting glycolysis as an energy source in two ways: 1) swapping out sucrose and trehalose in our recording saline in favor of 2-deoxy-D-glucose (a non-glycolytic sugar; (Fig. 6A); 2) addition of lonidamine (LDA) to the saline to acutely inhibit hexokinase (Fig. 6A). Control recordings with these conditions showed little effect on baseline physiology (Fig. 6A, D, F, H-J, Table 16). However, when Mitochondrial Complex I was impaired neuronally through *NDUFS7[RNAi]* in combination with inhibition of glycolysis, there was a drop in evoked neurotransmission (Fig. 6C, E, G, H-J, Table 16). This correlated with a failure to increase active zone material after *NDUFS7* gene knockdown (Fig. 6K-Q, Table 16). These results match the idea that a combination of mitochondrial function or glycolysis can work to maintain normal levels of NMJ output.

We continued this line of investigation genetically. We acquired RNA interference-based transgenes to target five genes involved in *Drosophila* glycolysis or subsequent ATP generation in the Citric Acid (TCA) Cycle: *hexokinase A (hex-A)*, *hexokinase C (hex-C)*, *Citrate (Si) Synthase I*, *Isocitrate dehydrogenase* (*Idh*), and *Succinyl-coenzyme A synthetase α subunit 1* (*Scsα1*). We knocked down these genes neuronally, either alone or in combination with *NDUFS7[RNAi]* (Figs. S15 and S16). Neuronal impairment of *hex-C* had no effect on baseline neurotransmission, but *hex-A* impairment reduced it (Figs. S15A-E, Table 17). Impairment of the TCA Cycle enzymes on their own had little-to-no effect on baseline neurotransmission (Fig. S15A, F-H, Table 17). However, concurrent impairment of *NDUFS7* and most of these genes significantly blunted neurotransmission, with the exception being *hex-C* (Figs. S15I-N, Table 17). Collectively, the data suggest that MCI works in conjunction with – or redundantly to – alternative energy-generating pathways to support normal levels of neurotransmission (Fig. S15O-Q). In the case of the hexokinases, *Drosophila hex-A* seems more important for this process than *hex-C*.

We also quantified active zone material accumulation. The results mirrored the neurotransmission tests: as before, neuronal *NDUFS7[RNAi]* impairment elicited enhanced active zone material (Fig. S16A-B, M, Table 18). On their own, impairment of glycolysis or TCA Cycle genes had variable effects active zone material (Fig. S16C-G, M, Table 18). But concurrent neuronal impairment of *NDUFS7* and any of the glycolysis or TCA Cycle genes reversed active zone enhancement (Fig. S16H-M, Table 18).

MCI subunits in muscle are required for proper synapse development

We previously reported that impairments of MCI diminish *Drosophila* NMJ growth [13]. The roles of MCI in specific tissues for this developmental process were unclear. For the present study, we tested for tissue-specific roles of MCI in NMJ development. To visualize NMJ boutons, we co-stained larval fillets with anti-Horseradish Peroxidase (HRP) a presynaptic membrane marker, and anti-Discs Large (Dlg), a postsynaptic density marker [63,64].

On a coarse level, MCI loss in muscle (*BG57-Gal4 > UAS-NDUFS7[RNAi]*) caused a severe reduction in average bouton size, a decrease in bouton number, a notable decrease in Dlg expression, and a bouton "clustering" phenotype (Fig. 7 A-B), reminiscent of what we previously reported [13]. To quantify these observations, we measured bouton number, muscle area, and branch number per muscle in the third-instar larval NMJ synapses (Table 19). We found that *NDUFS7* muscle knockdown resulted in a significant reduction in all these parameters compared to controls (Fig 7J-L, U-V, Table 19).

In contrast to muscle knockdown, pan-neuronal or motor neuron-specific knockdown of *NDUFS7* showed slight NMJ overgrowth phenotypes (Fig. 7F-L, Table 19). As was the case with the active zone enhancement associated with neuronal *NDUFS7* knockdown (Fig. 4), this NMJ overgrowth could represent a developmental mechanism to stave off dysfunction caused by missing neuronal mitochondria. Taken together, our NMJ immunostaining results indicated to us that blunted NMJ growth due to MCI loss was likely due to muscle MCI dysfunction.

We tested if the NMJ undergrowth phenotypes could be due to the increased levels of mitochondrial reactive oxygen species we had observed in muscle (mROS) (Fig. S2F). If that idea were correct, then the undergrowth phenotypes should be reversed if mitochondrial mROS were scavenged. Consistent with this idea, bouton number and synaptic undergrowth phenotypes were fully restored to wild-type levels when the *NDUFS7[RNAi]* muscle knockdown animals also had *UAS-Sod2* transgenically expressed in the muscles (Fig. 7C, J-L, R, Table 19). They were also restored to wild-type levels when muscle *NDUFS7* knockdown animals were raised on food containing the antioxidant N-acetyl cysteine amide (NACA) (Fig. 7J-L, Table 19). By contrast, none of these NMJ growth parameters were restored to wild-type levels if scavengers *UAS-Sod1* or *UAS-Catalase* were misexpressed in the muscle (Fig. 7D-E, J-L, Table 19).

Loss of MCI and mROS in muscle disorganize NMJ postsynaptic densities

The muscle Dlg-Spectrin network functions as an organizing scaffold for synaptic assembly [64,65]. Dysregulation of this network can lead to an aberrant muscle subsynaptic reticulum (SSR) [64].

Therefore, one possible target of excess mROS in the absence of MCI function is Dlg. Dlg is the fly

homolog of PSD-95/SAP97/PSD-93, and it is a member of the membrane-associated guanylate ki-

nase (MAGUK) family of NMJ scaffolding proteins [64]. It is present both within presynaptic boutons

and in the portion of the SSR closest to the bouton.

Using the same antibodies detailed above (anti-Dlg and anti-HRP), we used a high magnifi-

cation to examine at the postsynaptic densities closely. By confocal immunofluorescence, Dlg area

was significantly reduced when *NDUFS7* was depleted postsynaptically by[RNAi] (Fig. 7M-U, Table

19). To quantify the relative Dlg area, we measured Dlg area with respect to HRP (Relative Dlg ar-

ea = Dlg area minus HRP area) in type 1b boutons at muscle 6/7 of the A2 hemi segment (Fig. 7U).

Compared with the control synapses, *NDUFS7* knockdown resulted in a significant reduction in the

relative Dlg area (Fig 7M-U, Table 19). Consistently, the relative α-Spectrin area was also reduced

when *NDUFS7* was depleted in muscle (Fig. S17A-F, Table 7).

Next, we scavenged mROS to check how that affected the postsynaptic densities. The rela-

tive Dlg and α-Spectrin areas were restored to wild-type levels when animals were grown in a me-

dia containing NACA (*NDUFS7[RNAi]/+; BG57-Gal4/+* with NACA) or genetically expressing *UAS-*

Sod2 in the muscle (*UAS-Sod2/NDUFS7[RNAi]; BG57-Gal4/BG57-Gal4*) (Fig.7R, U, Table 19; Fig.

S16D-F, Table 7). By contrast, Dlg levels were not restored while expressing other scavengers in

the muscle, encoded by *UAS-Sod1* and *UAS-Catalase* transgenes (Fig. 7S-U, Table 19). We con-

clude that depletion of MCI subunits in the muscle disables postsynaptic density formation via the

formation of mitochondrial reactive oxygen species intermediates.

The postsynaptic density (PSD-95/Dlg) has been shown to cluster glutamate receptors at

the SSR [66]. Our results raised the possibility that glutamate receptor clusters could be disrupted

when *NDUFS7* was depleted in the muscle (Fig. 8). To test this idea, we simultaneously

immunostained NMJs with antibodies against Brp (neuron, presynaptic active zone) and glutamate

receptor clusters (muscle). In controls, these pre- and postsynaptic structures were directly ap-

posed to one another (Fig. 8A, G). But when *NDUFS7* gene function was depleted, we observed

"missing" GluRIIA and GluRIII receptor clusters (Fig. 8 B, F, H, K, Table 7), i.e., Brp puncta without

apposed glutamate receptors. These lack of apposition phenotypes were fully reversed by raising

the larvae with NACA or genetically expressing *UAS-Sod2* in the muscle (Fig. 8 D-F, I-K, Table 7).

However, as with prior tests, these phenotypes were not reversed by *UAS-Sod1* or *UAS-Catalase*

overexpression (Fig. S18). Together, our data indicate that loss of MCI subunit in the muscle

(*NDUFS7*) disrupts several aspects of the postsynaptic density organization, and these disruptions

are likely due to the accumulation of mitochondrial ROS in the muscle.

Loss of MCI subunits in muscle diminishes evoked NMJ neurotransmission through postsynaptic mROS

Mitochondria and reactive oxygen species influence presynaptic vesicle release and plasticity at synapses. This has been shown in diverse model systems like flies, mice, and worms [52,53,67-69]. In our prior work, we identified an NMJ neurotransmission defect when MCI function is impaired [13], but based on our current study, that defect does not seem like it was dependent upon MCI's neuronal functions (Figs. 1, 4, 5, 6). Therefore, we turned to analyzing postsynaptic muscle MCI and mROS to check if these parameters influenced neurotransmission.

We performed sharp electrode electrophysiological recordings of miniature and evoked excitatory postsynaptic potentials (mEPSP and EPSP) at NMJ muscle 6, hemi segment A2. We also used average mEPSP and EPSP values to estimate quantal content (QC) for each NMJ. For the most part, mEPSP values remained steady (Figs. 9A-E, Table 20), with some exceptions. The starkest phenotypes came in terms of evoked amplitudes (Figs. 9A-D, F, Table 20), sometimes due to changes in QC (Fig. 9G) or combinatorial changes in both mEPSP (Fig. 9E) and QC (Fig. 9G).

Evoked synaptic vesicle release (EPSP) was significantly reduced when *NDUFS7* was depleted in muscles by RNAi (Fig. 9A, F, Table 20). That reduction was reversed after scavenging mitochondrial ROS in the muscle. Indeed, transgenic muscle-driven *Sod2* gene expression suppressed the *NDUFS7[RNAi]* phenotype, but expression of the *Catalase* and *Sod1* did not (Fig. 9A, F, Table 20). We also tested if feeding a ROS scavenger to developing larvae would reverse the same neurotransmission defect. Carrier feeding alone (10% EtOH) did not affect evoked neurotransmission, nor did it influence the neurotransmission loss caused by *NDUFS7[RNAi]* (Fig. 9B, F, Table 20). Feeding larvae 0.5 mM NACA successfully reversed the phenotype, but the nonspecific additive curcumin had no effect (Fig. 9B, F, Table 20).

We checked different MCI manipulations. First, we examined hemizygous *ND-30^{EY03664/Df}* genetic mutants. As with muscle-driven *NDUFS7[RNAi]*, the *ND-30* mutant NMJs had blunted evoked neurotransmission, but this defect was successfully reversed by ROS scavengers (Fig. 9C, Table 20). We also impaired MCI pharmacologically, by feeding larvae 50 µM rotenone (or 0.5% DMSO carrier control), similar to conditions we previously published [13]. As with the prior study, rotenone blunted neurotransmission (Fig. 9D, F, G, Table 20), but this effect was ameliorated by a genetic background overexpressing *UAS-Sod2* in the muscle (Fig. 9D, F, G, Table 20). By contrast, overexpressions of *UAS-Catalase* and *UAS-Sod1* were not effective at reversing the effect of rotenone (Fig. 9D, F, G, Table 20).

Finally, we performed behavioral experiments on *NDUFS7* and *ND-30[RNAi]* and mutant animals. Consistent with the electrophysiological recordings, MCI muscle-depleted and mutant ani-

mals showed severe defects in crawling ability. The crawling behavior was rescued by expressing

the *UAS-Sod2 transgene* in the muscle or raising the larvae in a media containing NACA (Fig. 9H,

Table 21). By contrast, neuronal depletion of *NDUFS7* in larvae did not show any significant crawl-

ing defects (Fig. 9H, Table 21). Together, our data suggest that excess mitochondrial ROS accumu-

lation in muscle (mROS) diminishes baseline synaptic physiology when MCI activity is lost, and it

also triggers aberrant crawling behavior.

DISCUSSION

We uncovered novel aspects of NMJ synapse biology controlled by Mitochondrial Complex I (MCI).

Impairment of MCI causes profound cytological phenotypes in synaptic tissues (Figs. 1-3) [13]. By ex-

amining mitochondria directly, we discovered shared phenotypes between MCI loss and loss of the

Drosophila Mitofusin, Marf (Fig. 2). Additionally, with MCI loss, we noted an enhancement of mito-

chondrial reactive oxygen species (ROS, Fig. S1), consistent with prior work [36-40].

Unexpectedly, these perturbations spur functionally opposite responses in presynaptic neu-

rons vs. postsynaptic muscles. In motor neurons, MCI loss and mitochondrial ROS appear to trigger

a compensatory response, where the underlying cytological problems are offset by an increase in

active zone material, resulting in normal levels of evoked excitation (Fig. 4). This process requires

known intracellular calcium signaling components (Fig. 5). It also appears to require energy stores

because loss of glycolysis – which may function as a supplemental energy source to mitochondria –

abrogates the presynaptic compensation (Fig. 6). By contrast, in the muscle, MCI loss and mito-

chondrial ROS trigger a destructive response, where there is a disassembly of the postsynaptic

density (Fig. 7). This disassembly correlates with mis-apposition of pre- and postsynaptic structures

(Fig. 8) and defective neurotransmission (Fig. 9).

Disruption of mitochondrial dynamics, ROS, and physiological responses in *Drosophila*

Energy is needed for normal levels of synaptic transmission [2]. Intuitively, a loss of synaptic mito-

chondria should blunt transmission because transmission requires energy. Consistently, several

labs have previously implicated mitochondrial dynamics in *Drosophila* synapse function, including

mitochondrial fission (Dynamin related protein 1, Drp1 [51]), fusion (Mitofusin/dMarf [31]), trafficking (Mi-

ro and Milton [50,70,71]), or quality control (Pink and Parkin [72-75]). Additionally, it has been established

from model organisms such as flies, worms, and mice that any misregulation in mitochondrial distri-

bution could affect synaptic activity [51,59,76,77]. Adding to that work, we uncovered synaptic transmis-

sion and developmental phenotypes after depletion of Mitochondrial Complex I (MCI) at the NMJ [13]

potentially related to defective mitochondrial fusion (this study). Additionally, ROS has been studied

in the context of mitochondrial dysfunction [78]. Excess ROS can trigger mitochondrial calcium uptake

and subsequently trigger apoptosis or degeneration of neurons or neural support cells [79,80]. MCI de-

ficiency elevates ROS levels, and this process can promote the fragmentation of mitochondria in

cells like fibroblasts [81].

Numerous papers using *Drosophila* as a model have examined potential connections be-

tween MCI, ROS, aging, and physiology. The consequences of MCI loss and the induction of ROS

are context dependent. They are not always in obvious agreement, either from tissue to tissue or

from study to study. For example, one analysis demonstrated that reverse electron transfer (RET)

at MCI increases ROS production, and this occurs with the normal aging process [82]. Consistently,

inhibition of RET decreases ROS and increases lifespan; this result would suggest that RET-

induced ROS hurts survival [82]. However, a different study reported the opposite finding under

stress-inducing conditions. For that scenario, inhibition of RET decreases ROS production and de-

creases lifespan – while activation of RET enhances ROS and extends lifespan. Those findings

would suggest that ROS is protective under some conditions [83]. Effects are not limited to Complex I

manipulation. *Drosophila* deficient for the Complex V subunit *bellweather* gene function demon-

strate enhanced mitochondrial ROS; downstream this ROS, they have broadly enhanced levels of

active zone material (BRP) in adult brains [84]. Consistently, genetically boosting mitochondrial func-

tion in adults has the opposite effect – attenuated BRP accumulation in the fly brain [84]. These find-

ings closely match our larval motor neuron observations.

On the molecular level there are signaling puzzles uncovered by *Drosophila* work. For ex-

ample, the unfolded protein response (UPR) is a key mediator when MCI function in lost. Neuronal

impairment of the NDUFS1 (MCI subunit) homolog ND-75 elicits a spectrum of movement and sei-

zure phenotypes, and it also activates the UPR on the cellular level [85]. All those *Drosophila* neu-

ronal phenotypes can be reversed when yeast ND1 is exogenously added to the system, which re-

stores NADH dehydrogenase function (while not restoring ATP production) [85]. Those findings sug-

gest a damaging role in adult neurons for the MCI-deficient ROS-UPR axis. By contrast, another

paper examined ND-75 disruption in adult muscle – and for this tissue, researchers found that ele-

vated ROS levels trigger the UPR activation ultimately to preserve function [86]. That suggests a pro-

tective in in muscles for the ROS-UPR axis [86].

Our study expands upon these results and supports the idea that MCI has roles at synaptic

sites through disrupted mitochondrial dynamics. Lack of MCI in larval motor neurons causes loss of

mitochondria at synaptic terminals (Fig. 1). This defect is linked to defective mitochondrial fusion

(Fig. 2), which is known to maintain mitochondrial integrity [87,88]. By contrast, lack of MCI in larval

muscle appears to disassemble the postsynaptic density, which hurts neurotransmission (Figs. 7-9).

A novel form of presynaptic homeostatic plasticity triggered by ROS?

ROS has been linked to short-term synaptic plasticity in *Drosophila* [53], as well as long-term potentia-

tion (LTP) in mammals [89-91]. Here, we uncovered a role for ROS in augmentation of active zone ma-

terial when MCI is impaired in neurons. This finding could be considered a form of homeostatic

plasticity: an increase in active zone components likely drives potentiated release to compensate

for defective baseline synaptic transmission, which would be expected after the loss of an energy

source like mitochondria.

Homeostatic augmentations of active zone material have been reported at the *Drosophila*

NMJ. For example, *rab3* mutants have an increase in active zone material to offset decreased syn-

apse growth [92]. Additionally, ROS has been shown to be an obligate signal in *Drosophila* to main-

tain fundamental properties in both pre- and postsynaptic compartments, including at the NMJ [53]. In

our study, we observe an increase in active zone intensity (Fig. 4), which could overlap with mech-

anisms uncovered in those prior studies. Alternatively, it could be consistent with a form of homeo-

static plasticity at the *Drosophila* NMJ called Presynaptic Homeostatic Potentiation (PHP). PHP is

initiated when the activity of postsynaptic glutamate receptors is impaired. This decreases quantal

size. The synapse detects the impairment, and muscle-to-nerve signaling drives an increase in pre-

synaptic glutamate release [93,94]. This happens in part through an increase in influx of calcium into

the neuron through Ca_V2 voltage-gated calcium channels [93,95-98]. PHP coincides with an increase in

the size of the readily releasable pool (RRP) of synaptic vesicles [99-102] and apparent increases in

active zone protein content or organization [54,56,103]. The general model is that these modifications

drive the neuron to release more glutamate, offsetting the initial synaptic challenge.

Could the excess mitochondrial ROS that is caused by the loss of Complex I be triggering

the same or overlapping homeostatic mechanisms? It is possible. Quenching presynaptic ROS by

expression of *Sod2* or by feeding flies NACA led to a reversion of active zone material back down to

the control levels (Fig. 4E). Under ideal conditions, control levels of active zone material would sup-

port normal neurotransmission (Fig. 4A, J). However, when combined with presynaptic *NDUFS7*

loss (already depleting synaptic mitochondria; Figs. 1O, 2G), there is a loss in neurotransmission

capacity (Fig. 4M).

The molecular underpinnings of this mitochondrial-loss-induced homeostatic plasticity are

unknown, but prior work offers ideas. One possibility is that *Sod2* changes the redox status of the

entire cell, which could potentially affect active zone components [89]. Another possibility is that ROS

could alter the release of calcium from intracellular stores such as ER; and in turn, this could induce calcium signaling to mitochondria, which would contribute to the formation of ATP and subsequent vesicle fusion [59,60]. Consistent with this idea, we observed a reduced synaptic strength at the terminals after simultaneous blockade of calcium release and import from ER stores to mitochondria through genetic or pharmacological manipulations (Fig. 5). The link would not have to be a direct one. Indeed, a recent study indicated that activity-driven mitochondrial calcium uptake does not depend on the ER as a source of calcium to maintain normal synaptic strength [59]. Future studies can refine conclusions about presynaptic calcium dynamics downstream of MCI disruption. This is possible through reagents in the *Drosophila* neurogenetic toolkit like compart-targeted genetic-encoded calcium indicators [104].

Loss of MCI in muscle and subsequent ROS accumulation diminish synaptic excitation

Little is known about what role postsynaptic ROS plays in regulating synaptic plasticity at the NMJ. It is unlikely that ROS is an abstract signal that triggers wholesale destruction. It likely has specific targets. There are clues from previous work. For example, ROS signaling plays direct roles in the activity-dependent structural plasticity of motor neurons and postsynaptic dendrites [53]. Additionally, postsynaptic ROS plays critical roles in dendritic development. In *Drosophila*, ROS appears to act as a developmental plasticity signal to regulate the size of the dendritic arbors. [105].

It is slightly surprising that the NMJ fails to compensate for a lack of MCI function in the muscle. This is because the NMJ employs multiple retrograde signals to stabilize function [106]. One hypothesis is that MCI deficiency and ROS interfere with important muscle-to-nerve signals normally required for maintaining NMJ setpoint function. Some of these signals are required to correct acute challenges (minutes) to NMJ function. Some of the best characterized are: Bone Morphogenetic Protein (BMP) signaling [107-109]; the Insomniac (adaptor)-Cul3 (ubiquitin ligase)-Peflin (substrate) signaling complex [110]; and the endosomal recycling molecules class II PI3 kinase (PI3K) and Rab11 [111]. Additionally, the field has uncovered instructive signaling molecules from the muscle that maintain robust neurotransmission, including: muscle-secreted Semaphorin-2b [112]; and Target of Rapamycin (TOR) [113,114]. Future studies can address whether these or similar targets are substrates of negative ROS regulation in the muscle.

The roles of ROS in muscle are less understood in terms of synapse regulation and function. In this study, we showed that excess ROS is sufficient to affect the organization of postsynaptic densities (Fig. 7), ionotropic glutamate receptor clusters (Fig. 8), and spectrin cytoskeleton (Fig. S11) in the muscle. The correlated uncoupling of pre- and postsynaptic structures seems to be responsible for neurotransmission defects (Fig. 9), and it likely coincides with broader structural insta-

bility [65,115]. These phenotypes are reminiscent of other *Drosophila* mutant phenotypes showing a

degenerative NMJ (e.g., [65,115-117]). Consistent with a model of degeneration, we found behavioral

defects in locomotion (Fig. 9).

Despite the profound defects that result from muscle depletion of MCI, we were able to re-

verse many of those structural and functional problems by culturing animals in an antioxidant-rich

media (0.5mM NACA) or through muscle expression of *UAS-Sod2*. This is interesting finding for

future work because it points to mitochondrial ROS as a signal of interest in the manifestation of

synapse dysfunction. In subsequent work, we will take advantage of the *Drosophila* neurogenetic

toolkit to identify specific targets.

Limitations and Future Directions: Linking Mitochondrial ROS and Synapse Function

There are fundamental questions that remain to be answered in future work. Many of these deal

with ROS. First, we have not identified the specific ROS molecule(s) responsible for the synaptic

phenotypes we have observed. Our data point to mitochondrial matrix ROS because *UAS-Sod2*

overexpression ameliorates MCI loss phenotypes, while *UAS-Catalase* and *UAS-Sod1* do not. Ad-

ditionally, *UAS-Sod2[RNAi]* phenocopies the ROS accumulation associated with MCI loss.

If the relevant ROS molecule is signaling from the mitochondrial matrix, this could mean su-

peroxide ($O_2\bullet-$) or hydrogen peroxide (H_2O_2) – or less likely, another species, like hydroxyl radicals

($\bullet OH$). The function of superoxide dismutase enzymes is to convert superoxide into hydrogen per-

oxide [118]. Because SOD2 is mitochondrial and it has a mollifying effect on MCI dysfunction in our

system, we would hypothesize that mitochondrial matrix superoxide ($O_2\bullet-$) is the likely culprit for the

synaptic phenotypes we observe. This idea would explain why other antioxidants like SOD1, and

Catalase do not help. SOD1 accumulates mainly in the cytoplasm [119,120], while Catalase accumu-

lates mainly in peroxisomes [121,122]. NACA, which is the amide form of the antioxidant N-

acetylcysteine, is membrane permeable; as such, it could accumulate in mitochondria [123]. We have

not affirmatively demonstrated sub-compartment mitochondrial localization of NACA in our system.

However, recent work by other groups is consistent with a reductive function of NACA for mito-

chondria [124,125].

Even though our data point to mitochondrial ROS, the mechanistic connection between ma-

trix $O_2\bullet-$ and the synapse is not clear. Additionally, we do not know the relevant subcellular site(s)

where ROS would need to accumulate to have its effects on the synapse. An intuitive idea is that

the affected mitochondria and ROS would be near the synapse itself. For the NMJ, this is possible

both in neurons and muscle. Yet our data do not neatly match that model. For example, six hours of

rotenone exposure to the NMJ can enhance active zone accumulation, but only if the motor nerve is

intact (Fig. S12). This finding suggests that the signaling downstream of mitochondrial ROS in neu-
rons needs both time and space to work.

Concerning the muscle, mitochondrial ROS seems to have a damaging effect directly on the
postsynaptic density. This effect hurts synaptic function. If the relevant site of mitochondrial ROS in
muscle happens to be closer to the NMJ than the relevant site in neurons, then the damaging ef-
fects from the muscle could partly a function of proximity. Determining if it matters whether mito-
chondrial ROS is physically near the NMJ could inform the possible signaling modalities that govern
mitochondrial regulation of synapse function.

RESOURCE AVAILABILITY

Lead Contact

The lead contact for this study is Dr. C. Andrew Frank (andy-frank@uiowa.edu).

Materials Availability

Dr. C. Andrew Frank's laboratory uses the fruit fly *Drosophila melanogaster* as a model organism.
Consistent with ethical conduct of research, the lab maintains *Drosophila* stocks that it generates
and publishes, as well as useful precursors and derivatives of those stocks. These stocks are avail-
able to researchers who make requests to Dr. Frank (andy-frank@uiowa.edu) – or to the appropri-
ate original source – and they are shipped promptly.

Data and Code Availability

Consistent with the Data Sharing Policy for NIH-funded research, Dr. C. Andrew Frank's laboratory
will share data related to this study. The lab will share final data through publication of summary fig-
ures or tables. In addition, a large amount of final data derives from performing electrophysiological
recordings or synapse imaging experiments and then performing subsequent analyses. Those ex-
periments yield raw data in electronic form. Therefore, in addition to publishing final data, the lab
will share raw data and related materials with researchers who make requests to Dr. Frank (andy-
frank@uiowa.edu). No original coding was required to execute this study.

Acknowledgments

We acknowledge the Developmental Studies Hybridoma Bank (University of Iowa, USA) for anti-
bodies used in this study and the Bloomington Drosophila Stock Center (Indiana University, Bloom-
ington, USA) for fly stocks. We thank members of the Frank lab for their helpful comments and dis-

708 cussions in this study. BM was supported in part by NIH/NINDS grants to CAF (NS085164,

709 NS130108, and NS136753). Collectively, the work was supported by those same NIH/NINDS

710 grants, as well as funds from the University of Iowa's Carver College of Medicine to CAF. Funding

711 supporting the work executed by SAB and XW was from an NIH/NINDS R01 grant NS128040 to

712 XW.

Author contributions

715 BM, SAB, XW, and CAF designed the research; BM and SAB performed the research; BM, SAB,

716 XW, and CAF analyzed the data; BM and CAF wrote the initial draft of the paper; BM, SAB, XW,

717 and CAF wrote and edited the revision draft of the paper.

Declarations of interests

720 The authors declare no competing financial interests.

SUPPLEMENTAL INFORMATION

Document S1:

• Supplemental Figures S1-S18 and legends.

Document S2:

• Tables 1-21: Summary data for graphs in all Figures and Supplemental Figures.
• Table 22: Detailed information about reagents and materials for items described in the
STAR METHODS section.

FIGURE LEGENDS

Figure 1: MCI-depleted flies harbor fewer mitochondria at neuromuscular junctions

Mitochondrial morphology and trafficking defects in the ventral nerve cord, distal axons, and boutons. RNAi lines and controls were crossed to a motor neuron driver (*D42-GAL4*) and a mitochondrial marker (*UAS-mitoGFP*). (A-A′), (B-B′), (C-C′) and (D-D′) represent control ventral nerve cord (VNC), a magnified section of VNC, proximal (A2) and distal (A5) axons, respectively. These tissues exhibit regular mitochondrial clusters in the soma and axons. (E-E′), (F-F′), (G-G′) and (H-H′) represent *NDUFS7* knocked down ventral nerve cord (VNC), a magnified section of VNC, proximal (A2) and distal (A5) axons, respectively. Mitochondria are abnormally clustered in *NDUFS7[RNAi]* in the ventral nerve cord and distal segments of A5 axons. *NDUFS7[RNAi]* yields fewer mitochondria in the distal segments when compared to the proximal segments. (I-J) Representative electrophysiological traces showing evoked potentials of *mitoGFP, D42-Gal4 × UAS-NDUFS7*[RNAi] larvae at the A5 hemi segment of muscle 6/7 synapse. Scale bars for EPSPs (mEPSP) are x=50 ms (1000 ms) and y= 10 mV (1 mV). Fewer mitochondria at the presynaptic A5 hemi segment did not affect evoked NMJ excitation. K. Quantification showing EPSP amplitude at NMJ 6/7 in control (*D42-Gal4/+*; EPSP: 19.99 mV ± 2.53, n=6) and RNAi-depleted animals (*NDUFS7[RNAi] /+;D42-Gal4/+* ;EPSP: 16.88 mV ± 1.21, n=9). (L-M) Representative images showing mitochondria morphology in control and RNAi-depleted animals in muscle. Mitochondria in *NDUFS7[RNAi]*-depleted larvae are clustered compared to the control larvae. (N-Q) *NDUFS7[RNAi]* contains almost no mitochondria in boutons when co-stained with pre- (HRP) or post-synaptic markers (Discs Large, Dlg). (A-H′, L-M, N-O) Scale bar: 10 µm. (P) Quantification showing the number of mitochondrial clusters at NMJ 6/7 in control (*D42-Gal4/+*; # clusters: 113.6 ± 11.97, n=7) and RNAi-depleted animals (*NDUFS7[RNAi]/+; D42-Gal4/+*; # clusters: 30.33 ± 4.02, n=6). (Q-R) 3D rendered image showing the volume (μm^3) of mitochondria at NMJ 6/7 in RNAi knockdown larvae (*NDUFS7[RNAi] /+; D42-Gal4/+ ;* 0.11 ± 0.01 μm^3, n=16) compared to the driver control animals (*D42-Gal4/+*; 0.29 ± 0.04 μm^3, n=14). (Q-R) Scale bar: 5 µm. (S) Quantification shows a significantly lower mitochondria volume in boutons at NMJ 6/7 in RNAi-depleted animals. ****p*<0.0001 and ****p*=0.0003 for mitochondrial clusters and volume, respectively. Statistical analyses based on Student's t-test. Error bars represent mean ± s.e.m.

Figure 2: Loss of *NDUFS7* in motor neurons phenocopies a *marf* depletion

Mitochondrial morphology and trafficking defects in the ventral nerve cord and distal axons. To label neuronal mitochondria, *UAS-[RNAi]* lines and controls were crossed to a motor neuron driver (*D42-Gal4*) and a mitochondrial marker (*UAS-mitoGFP*). (A) Ventral nerve cord (VNC): *UAS-mitoGFP* and *drp1[RNAi]* exhibit normal mitochondrial organization, *NDUFS7[RNAi]* and *marf*[RNAi] exhibit clustered mitochondria, *NDUFS7[RNAi] ; marf[RNAi]* and *ND-20[RNAi]; drp1[RNAi]* doubles exhibit clustered mitochondria in the soma. The fluorescent images were skeletonized to measure mitochondrial branch length. (B) Comparison of a proximal axonal segment in A2 and a distal segment in A5. Distal segments of A5 axons in *NDUFS7[RNAi]* and *marf[RNAi]* contain many fewer mitochondria than proximal segments. Knocking down *NDUFS7[RNAi]* and *marf[RNAi]* together does not show an additive effect. (A-B) Scale bar: 10 μm. (C-D) Histogram showing mitochondrial branch length (μm) and number (μm^2 area of bouton) in VNC and axons of the third instar larvae in the indicated genotypes. (E-L) Representative images of the A2 hemi segment of muscle 6/7 NMJs in (E) *UAS-mito-GFP, D42-Gal4/+*, (F) *UAS-Sod2/+; UAS-mito-GFP, D42-Gal4/+,* (G) *NDUFS7[RNAi] /+; UAS-mito-GFP, D42-Gal4/+*, (H) *NDUFS7[RNAi] /UAS-Sod2;UAS-mito-GFP,D42-Gal4/+*, (I) *UAS-mito-GFP,D42-Gal4/marf[RNAi]* , (J) *UAS-Sod2;UAS-mito-GFP,D42-Gal4/marf[RNAi]*, (K) *NDUFS7[RNAi]/+; UAS-mito-GFP,D42-Gal4/marf[RNAi]* and (L) *NDUFS7[RNAi] /UAS-Sod2;UAS-mito-GFP,D42-Gal4/marf[RNAi]* larvae immunostained with antibodies against HRP (magenta) and GFP (mito-GFP:green) to label neurons and mitochondria. *NDUFS7[RNAi]*- and *marf[RNAi]*-depleted animals harbor fewer mitochondria at the terminals as compared to control animals. Transgenic *UAS-Sod2* rescued mitochondrial clustering defects. (E-L) Scale bar: 5 μm. (M) Histograms showing quantification of mitochondrial clusters at the NMJs in the indicated genotypes. ***p <0.0001; ns, not significant. Statistical analysis based on one-way ANOVA followed by post-hoc Tukey's multiple-comparison test. Error bars represent mean ± s.e.m.

Figure 3: *NDUFS7* depletion in motor neurons affects synapse stability

Representative confocal images of NMJ synapses at muscle 6/7 of (A-A′) *D42-Gal4* control, (B-B′) *UAS-Sod2* overexpression (C-C′) *D42-Gal4*-driven *NDUFS7*[RNAi] (*NDUFS7[RNAi] /+; D42-Gal4/+*), (D-D′) *NDUFS7* knockdown with NACA rescue (*NDUFS7[RNAi] /+; D42-Gal4/+ +NACA*), (E-E′) *NDUFS7* knockdown with *UAS-Sod2* (*UAS-NDUFS7[RNAi]/UAS-Sod2; D42-Gal4/+*). Each condition was double immunolabeled with 22C10 (anti-Futsch, magenta) and anti-HRP (green) antibodies. The motor neuron-depleted *NDUFS7[RNAi]* larvae showed a decrease in the number of Futsch-positive loops as compared to the Gal4 control. Futsch-positive loops were significantly restored to the control number when *NDUFS7*[RNAi] was raised in media containing NACA or genetically expressing *UAS-Sod2* in the *UAS-NDUFS7[RNAi]* background. Scale bar: 10 μm. (F) Histograms showing the percentage of Futsch-positive loops in the indicated genotypes. *p=0.01 (Gal4 control vs *sod2* OE neuron), *p=0.0008 (Gal4 control vs *NDUFS7*[RNAi] neuron), **p=0.001 (*NDUFS7*[RNAi] neuron vs *NDUFS7*[RNAi] neuron + NACA) and **p=0.0005 (*NDUFS7*[RNAi] neuron vs *sod2* neuron rescue). Statistical analysis based on one-way ANOVA followed by post-hoc Tukey's multiple-comparison test. Error bars represent mean ± s.e.m.

Figure 4: Neuronal ROS (nROS) controls active zone material levels at NMJs

(A) Representative images of the A2 hemi segment of muscle 6/7 NMJs in *UAS-mito-GFP, D42-Gal4/+*, (B) *UAS-Sod2/+; UAS-mito-GFP, D42-Gal4/+* (C) *NDUFS7[RNAi]/+; UAS-mito-GFP, D42-Gal4/+*, (D) *NDUFS7[RNAi] /+; UAS-mito-GFP, D42-Gal4/+* with NACA and (E) *NDUFS7[RNAi]/UAS-Sod2; UAS-mito-GFP, D42-Gal4/+* larvae immunostained with antibodies against the active zone scaffold Bruchpilot (BRP:fire-LuT) to label the active zones. BRP levels are upregulated at the NMJs in *NDUFS7*[RNAi] depleted flies, while overexpression of ROS scavenger gene *Sod2* in the neuron or feeding the larvae with N-Acetyl L-cysteine amide (NACA) restores BRP to the control level. (A-E) Scale bar: 2.5 μm. (F-G) Histograms showing quantification of BRP intensity (F) and density (G) in μm^2 area of bouton at muscle 6/7 in the genotypes mentioned above. At least 8 NMJs of each genotype were used for quantification. ***p <0.0001. Error bars denote mean ± s.e.m. Statistical analysis based on one-way ANOVA followed by post-hoc Tukey's multiple-comparison test (H-H'-I-I'). Representative confocal images of muscle 6/7 NMJs in the (H-H') control (*D42-Gal4/+*) and (I-I') motor neuron Gal4 driven *NDUFS7[RNAi]* (*NDUFS7[RNAi] /+; D42-Gal4/+*) immunostained with antibodies against Bruchpilot (BRP: magenta) and GluRIII (green) to label a glutamate receptor subunit. (H-H'-I-I') Scale bar: 2.5 μm. There are no significant changes of GluRIII-BRP apposed clusters. At least 8 NMJs of each genotype were used for quantification (J-P). Representative traces, quantification of mEPSPs, EPSPs and quantal content in the indicated genotypes. Scale bars for EPSPs (mEPSP) are x=50 ms (1000 ms) and y= 10 mV (1 mV). EPSPs amplitudes were maintained in *NDUFS7[RNAi]*-depleted flies due to increased levels of active zone material (e.g., BRP); however, NMJs with *Sod2*-rescued *NDUFS7[RNAi]* in neurons showed diminished evoked release when compared with *NDUFS7[RNAi]*. Minimum 8 NMJs recordings of each genotype were used for quantification. *$p < 0.05$, ***$p <0.0001$; ns, not significant. Statistical analysis based on one-way ANOVA followed by post-hoc Tukey's multiple-comparison test. Error bars denote the standard error of the mean. (Q) Representative images of the A2 hemi segment of muscle 6/7 NMJs in *UAS-mito-GFP, D42-Gal4/+*, (R) *UAS-Sod2/+; UAS-mito-GFP, D42-Gal4/+* (S) *NDUFS7[RNAi] /+; UAS-mito-GFP, D42-Gal4/+*, and (T) *NDUFS7[RNAi]/UAS-Sod2; UAS-mito-GFP,D42-Gal4/+* larvae immunostained with antibodies against HRP (magenta) and GFP (mito-GFP:green) to label neurons and mitochondria. *NDUFS7*-depleted and *sod2*-rescued *NDUFS7*[RNAi] animals harbor fewer mitochondria at the terminals than control animals. (Q-T) Scale bar: 5 μm. (U) Histograms showing quantification of mitochondrial clusters in the above-indicated geno-types. (V-X) Schematic illustration showing ROS (magenta) levels, BRP (grey) and mitochondria (red) number in the indicated genotypes. At least 8 NMJs of each genotype were used for quantification. ***$p<0.0001$. Error bars represent mean ± s.e.m. Statistical analysis based on one-way ANOVA fol-lowed by post-hoc Tukey's multiple-comparison test.

Figure 5: Loss of an MCI subunit necessitates ER-mediated calcium release to maintain evoked neurotransmission at the NMJs

(A-D) Schematics illustrating the role of IP$_3$ receptor (IP$_3$R), Ryanodine receptor (RyR) in the endoplasmic reticulum, mitochondrial calcium uniporter Complex (MCU), Ca$_v$2 calcium channel and synaptic vesicles at the presynaptic nerve terminal. The IP$_3$ and Ryanodine receptors were blocked pharmacologically (red X symbols) by using the IP$_3$R antagonist Xestospongin C or neuronally expressing *UAS-IP$_3$-Sponge* and RyR antagonist Dantrolene, while *mcu[RNAi]* was used to block the Mitochondrial calcium uniporter complex (red X). (E-P) Representative traces of mEPSPs and EPSPs in (E) pan-neuronal Gal4 control (*elaV(C155)-Gal4/+*), (F) pan-neuronal Gal4 control (*elaV(C155)-Gal4/+*) with an acute application (10 minutes) of 20 μM Xestospongin C and 10 μM Dantrolene, (G) pan-neuronal Gal4 driven *NDUFS7*[RNAi] (*elaV(C155)-Gal4)/+;NDUFS7[RNAi] /+*), (H) pan-neuronal Gal4-driven *NDUFS7*[RNAi] (*elaV(C155)-Gal4/+;NDUFS7[RNAi] /+*) with 20 μM Xestospongin C and 10 μM Dantrolene, (I) pan-neuronal Gal4 driven UAS-*IP3-Sponge.m30* with an acute application of 10 μM Dantrolene (*elaV(C155)-Gal4)/+; NDUFS7[RNAi]/+ ;UAS-IP3-Sponge.m30/+*), (J) pan neuronal Gal4 driven *mcu*[RNAi] with 20 μM Xestospongin C and 10 μM Dantrolene (*elaV(C155)-Gal4)/+; mcu[RNAi]/+),* (K) pan-neuronal *mcu[RNAi]* + *NDUFS7*[RNAi] with 20 μM Xestospongin C and 10 μM Dantrolene and(*elaV(C155)-Gal4)/+; mcu[RNAi]/NDUFS7[RNAi]*) and (L) pan neuronal *UAS-IP3-Sponge.m30* + *mcu[RNAi]* + *NDUFS7*[RNAi] with 10 μM Dantrolene (*elaV(C155)-Gal4/+;NDUFS7[RNAi] /mcu[RNAi] ;UAS-IP$_3$-Sponge.m30/+*), (M) Pan neuronal Gal4 (*elaV(C155)-Gal4/+*) with DMSO, (N) pan neuronal Gal4 (*elaV(C155)-Gal4/+*) with 20 μM BAPTA-AM, (O) pan-neuronal Gal4 driven *NDUFS7*[RNAi] (*elaV(C155)-Gal4)/+;NDUFS7[RNAi] /+*) with DMSO and (P) pan neuronal Gal4 driven *NDUFS7*[RNAi] (*elaV(C155)-Gal4)/+;NDUFS7[RNAi] /+*) with 20 μM BAPTA-AM. Scale bars for EPSPs (mEPSP) are x=50 ms (1000 ms) and y= 10 mV (1 mV). Note that EPSPs amplitudes were reduced in pan-neuronal Gal4 driven *mcu[RNAi]* + *NDUFS7*[RNAi] with an acute exposure of 20 μM Xestospongin C, 10 μM Dantrolene or *UAS-IP$_3$-sponge.m30* + *mcu[RNAi]* + *NDUFS7*[RNAi] with 10 μM Dantrolene and pan-neuronal Gal4 driven *NDUFS7*[RNAi] with 20 μM BAPTA-AM. (Q-T) Histograms showing average mEPSPs, EPSPs amplitude, and quantal content in the indicated genotypes. A minimum of 8 NMJs recordings of each genotype were used for quantification. **$p < 0.05$ (EPSP and QC: *NDUFS7*[RNAi] neuron + DMSO, wash vs *NDUFS7*[RNAi] neuron + BAPTA-AM, wash), *$p < 0.05$, **$p=0.001$, ***$p <0.0001$; ns, not significant. Statistical analysis based on one-way ANOVA followed by post-hoc Tukey's multiple-comparison test. Error bars represent mean ± s.e.m.

Figure 6: Loss of an MCI subunit necessitates glycolysis to regulate levels of active zone materials and stabilize synaptic strength

(A) Schematic illustrations showing steps of glucose metabolism and ATP production during glycolysis and TCA cycle in the cell. (B-G) Representative traces of mEPSPs and EPSPs in (B) pan-neuronal Gal4 control (*elaV(C155)-Gal4/+* with DMSO), (C) pan-neuronal Gal4 control (*elaV(C155)-Gal4/+*) with an acute application (30 minutes) of 150 µM Lonidamine (LDA) (D) pan-neuronal Gal4 driven *NDUFS7*[RNAi] (*elaV(C155)-Gal4)/+; NDUFS7[RNAi] /+* with DMSO), (E) pan-neuronal Gal4 driven *NDUFS7*[RNAi] (*elaV(C155)-Gal4)/+; NDUFS7[RNAi]/+*) with 150 µM Lonidamine, (F) pan-neuronal Gal4 control (*elaV(C155)-Gal4/+*) and HL3 containing 2-Deoxy-D-glucose and (G) pan-neuronal Gal4 driven *NDUFS7*[RNAi] (*elaV(C155)-Gal4)/+; NDUFS7[RNAi] /+*) with HL3 containing 2-Deoxy-D-glucose. Scale bars for EPSPs (mEPSP) are x=50 ms (1000 ms) and y= 10 mV (1 mV). The EPSPs amplitudes were reduced in pan-neuronal Gal4-driven *NDUFS7*[RNAi] with an acute exposure of 150 µM Lonidamine (LDA) or pan-neuronal Gal4-driven *NDUFS7*[RNAi] in HL3 containing 2-Deoxy-D-glucose for 30 minutes. However, we saw a reduction in mEPSP amplitudes when incubated in HL3 containing 2-Deoxy-D-glucose for 30 minutes in pan-neuronal Gal4-driven *NDUFS7[RNAi]* -depleted larvae compared to the control animals. (H-J) Histograms showing average mEPSPs, EPSPs amplitude, and frequencies in the indicated genotypes. A minimum of 8 NMJ recordings of each genotype were used for quantification. ,**p=0.003 (mEPSP amplitude: *elaV(C155)-Gal4/+*2-Deoxy-D-glucose vs *elaV(C155)-Gal4)/+;NDUFS7[RNAi] / +*2-Deoxy-D-glucose), **p=0.002 (EPSP amplitude: *elaV(C155)-Gal4/+*2-Deoxy-D-glucose vs *elaV(C155)-Gal4)/+;NDUFS7[RNAi] /+*2-Deoxy-D-glucose),***p=0.0001 (EPSP amplitude: *elaV(C155)-Gal4/+*LDA vs *elaV(C155)-Gal4)/+;NDUFS7[RNAi] / +*LDA); ns, not significant. Statistical analysis is based on the Student's t-test for pairwise sample comparison. Error bars represent mean ± s.e.m. (K-P) Representative images of the A2 hemi segment of muscle 6/7 NMJs in the above-indicated genotypes immunostained with antibodies against the active zone scaffold bruchpilot (BRP:fire-LuT) to label the active zones. The BRP levels downregulated at the NMJs in pan-neuronal Gal4 driven *NDUFS7*[RNAi] either incubated with 150 µM LDA or in HL3 containing 2-Deoxy-D-glucose for 30 minutes. (K-P) Scale bar: 5 µm. (Q) Histograms showing quantification of BRP intensity in µm^2 area of bouton at muscle 6/7 in the genotypes mentioned above. At least 8 NMJs of each genotype were used for quantification.***p<0.0001 (BRP levels: *elaV(C155)-Gal4/+*DMSO vs *elaV(C155)-Gal4)/+;NDUFS7[RNAi]/+* +DMSO),**p=0.008 (BRP levels: *elaV(C155)-Gal4/+*LDA vs *elaV(C155)-Gal4)/+;NDUFS7[RNAi] /+*LDA), ***p=0.0005 (BRP levels: *elaV(C155)-Gal4/+*2-Deoxy-D-glucose vs *elaV(C155)-Gal4)/+;NDUFS7[RNAi] /+*2-Deoxy-D-glucose). Error bars denote mean ± s.e.m. Statistical analysis based on one-way ANOVA followed by post-hoc Tukey's multiple-comparison test.

Figure 7: *NDUFS7* in muscles is required to promote normal synapse growth

(A-I) Representative confocal images of NMJ synapses at muscle 6/7 of (A) Muscle-Gal4 control (*BG57-Gal4/+*), (B) Muscle Gal4 driven *NDUFS7*[RNAi] (*NDUFS7[RNAi] /+; BG57-Gal4/+*), (C) *sod2* muscle rescue (*UAS-Sod2/NDUFS7[RNAi] ; BG57-Gal4/BG57-Gal4*), (D) *Sod1* muscle rescue (*UAS-Sod1/NDUFS7[RNAi] ; BG57-Gal4/BG57-Gal4*), (E) Catalase muscle rescue (*UAS-Cat/NDUFS7[RNAi] ; BG57-Gal4/BG57-Gal4*), (F) Pan neuronal Gal4 control (*elaV(C155)-Gal4)/+*), (G) *elaV(C155)- Gal4)-Gal4* driven *NDUFS7*[RNAi] (*elaV(C155)-Gal4/+; NDUFS7[RNAi] /+*), (H) Motor neuron Gal4 control *(D42-Gal4/+)* and (I) *D42*-Gal4-driven *NDUFS7*[RNAi] (*NDUFS7[RNAi] /+; D42-Gal4/+*) double immunolabeled with Dlg (magenta) and HRP (green) antibodies. (A-I) Scale bar: 10 μm. The NMJ morphological defects in *NDUFS7*[RNAi] were restored upon co-expression of *UAS-Sod2 in* muscle; however, it did not rescue with *UAS-Sod1* or *UAS-Catalase* transgene. (J-L) Histograms show the number of boutons, muscle area, and average NMJ length at muscle 6/7 of A2 hemi segment in the indicated genotypes. *p <0.05, *p=0.006 (# of boutons: Muscle *NDUFS7*[RNAi] vs *sod2* muscle rescue), **p=0.0002 (# of boutons: Muscle *NDUFS7*[RNAi] vs Muscle *NDUFS7*[RNAi] +NACA), *p=0.009 (# of branches: Muscle Gal4 vs Muscle *sod2* OE), ***p=0.002 (# of branches: Muscle Gal4 vs Muscle *sod1* OE), *p=0.008 (# of branches: Muscle Gal4 vs Muscle *cat* OE), ***p=0.001 (Muscle Gal4 vs Muscle *NDUFS7*[RNAi]), ***p<0.0001; ns, not significant. Statistical analysis based on one-way ANOVA with post-hoc Tukey's test for multiple and Student's t-tests for pairwise comparison. Error bars represent mean ± s.e.m. (M-T) Representative confocal images of boutons at the third instar larval NMJ synapse in (M) Muscle-Gal4 control (*BG57-Gal4/+*), (N) Muscle Gal4 driven *UAS-Sod2* (*UAS-Sod2/+; BG57-Gal4/+*), (O) *UAS-Sod1* (*UAS-Sod1/+; BG57-Gal4/+*), (P) *UAS-Catalase* (*UAS-Catalase/+; BG57-Gal4/+*) (Q) Muscle *NDUFS7*[RNAi] (*NDUFS7[RNAi] /+; BG57-Gal4* (R) *sod2* muscle rescue (*UAS-Sod2/NDUFS7[RNAi] ; BG57-Gal4/BG57-Gal4*), (S) *sod1* muscle rescue (*UAS-Sod1/NDUFS7[RNAi] ; BG57-Gal4/BG57-Gal4*) and (T) Catalase muscle rescue (*UAS-cat/NDUFS7[RNAi] ; BG57-Gal4/BG57-Gal4*) animals double immunolabeled with anti-HRP (green) and anti-Dlg (magenta) antibodies. (M-T) Scale bar: 5 μm. Note that the gross morphology of SSRs and immunoreactivity of Dlg were reduced in *NDUFS7[RNAi]* animals. As mentioned, phenotypes were restored to wild-type level when *NDUFS7*[RNAi] -depleted flies were reared in NACA or by genetically expressing *Sod2 transgene* in muscle. (U-V) Histograms showing normalized synaptic fluorescence of Dlg and bouton area in the indicated genotypes. *p <0.01, **p <0.001, ***p<0.0001; ns, not significant. Error bars represent mean ± s.e.m. Statistical analysis based on one-way ANOVA with post-hoc Tukey's test for multiple and Student's t-tests for pairwise comparison.

Figure 8: *NDUFS7* subunit in muscle affects the organization of GluRs cluster in *Drosophila*.

Representative confocal images of boutons at third instar larval NMJ synapse in (A) Muscle-Gal4 control (*BG57-Gal4/+*), (B) Muscle *NDUFS7*[RNAi] (*NDUFS7[RNAi] /+; BG57-Gal4/+*), (C) Muscle Gal4 driven *UAS-Sod2* (*UAS-Sod2/+; BG57-Gal4/+*), (D) *NDUFS7[RNAi] /+; BG57-Gal4/+*NACA) and (E) *sod2* muscle rescue (*UAS-Sod2/NDUFS7[RNAi] ; BG57-Gal4/BG57-Gal4*) animals immunolabeled with active zone marker BRP (green) and anti-GluRIII (magenta) antibodies. Scale bar: 5 μm. Note that GluRIII apposed clusters with BRP are missing in the *NDUFS7*[RNAi]-depleted animals (marked in arrow) compared to control. These phenotypes were restored to normal when *NDUFS7*[RNAi]-depleted flies were reared in media containing NACA or genetically expressing *Sod2 transgene* in muscle. (F) Histograms showing quantification of the number of missing BRP-GluRIII apposed puncta per bouton in the indicated genotypes. (G-J) Similar phenotypes were observed when analyzed for BRP-GluRIIA apposed clusters in boutons. (K) Histograms showing quantification of the number of missing BRP-GluRIIA apposed puncta per bouton in the indicated genotypes. ***$p<0.0001$. Error bars represent mean ± s.e.m. Statistical analysis based on one-way ANOVA with post-hoc Tukey's test for multiple comparisons.

Figure 9: Loss of MCI subunits affects synaptic transmission via the formation of excess ROS in the muscle

(A) Representative traces of mEPSPs and EPSPs in muscle-Gal4 control (*BG57-Gal4/+*), muscle Gal4 driven *NDUFS7*[RNAi] (*NDUFS7[RNAi] /+; BG57-Gal4/+),* muscle *Catalase* rescue (2X muscle-Gal4>*UAS-Catalase/NDUFS7*[RNAi] : *UAS-Catalase/NDUFS7[RNAi] ; BG57-Gal4/BG57-Gal4*), muscle *Sod1* rescue (2X muscle-Gal4>*UAS-Sod1/NDUFS7*[RNAi] : *UAS-Sod1/NDUFS7[RNAi] ; BG57-Gal4/BG57-Gal4*), muscle *Sod2* rescue (2X muscle-Gal4>*UAS-Sod2/NDUFS7*[RNAi] : *UAS-Sod2/NDUFS7[RNAi] ; BG57-Gal4/BG57-Gal4*) animals. Note that EPSPs amplitudes were reduced in *NDUFS7[RNAi],* and the phenotype was restored to wild-type levels by expressing *Sod2 transgene* in the muscle but not with *Sod1* and *Catalase* transgenes. (B) Representative traces of mEPSPs and EPSPs in muscle Gal4 control (*BG57-Gal4/+*) larvae raised on 10% EtOH, *NDUFS7* muscle[RNAi] (*NDUFS7[RNAi] /+; BG57-Gal4/+*) raised on 10% EtOH, *NDUFS7* muscle[RNAi] (*NDUFS7[RNAi] /+; BG57-Gal4/+*) raised on 0.5mM curcumin, *NDUFS7* muscle[RNAi] (*NDUFS7[RNAi] /+; BG57-Gal4/+*) raised on 0.5mM NACA, (C) Representative traces of mEPSPs and EPSPs in *ND-30^{epgy}/Df* mutants, *ND-30^{epgy}/Df* mutants raised on 0.5 mM NACA and *UAS-Sod2* muscle rescued *ND-30^{epgy}/Df* mutant (*UAS-Sod2/+; ND-30 (Df), BG57-Gal4/ND-30^{epgy}*) animals. The EPSPs amplitudes were restored to wild type when muscle depleted *NDUFS7[RNAi]* larvae were raised in food containing NACA or by *UAS-Sod2* muscle overexpression in *NDUFS7[RNAi]* depleted animals. (D) Representative traces of mEPSPs and EPSPs in muscle-Gal4 control (*BG57-Gal4/+*) larvae raised on DMSO, muscle Gal4 (*BG57-Gal4/+*) larvae raised on 50μM rotenone (complex I inhibitor), *UAS-Catalase (UAS-Catalase/+; BG57-Gal4/)*, *UAS-Sod1(UAS-Sod1/+; BG57-Gal4/),* and *UAS-Sod2 (UAS-Sod2/+; BG57-Gal4/+)* muscle over-expression animals raised on 50μM rotenone. The EPSPs amplitudes were suppressed in rotenone raised larvae overexpressing *UAS-Sod2* in muscle *(UAS-Sod2/+; BG57-Gal4/)* due to its free radical scavenging activity. Scale bars for EPSPs (mEPSP) are x=50 ms (1000 ms) and y= 10 mV (1 mV). (E-G) Histograms showing average mEPSPs, EPSPs amplitude, and quantal content in the indicated genotypes. Minimum 8 NMJs recordings of each genotype were used for quantification. (H) Histogram representing crawling behavior (in cm) of the larvae in the indicated genotypes. Knocking down *NDUFS7*[RNAi] or *ND-30[RNAi]* in muscle and *ND-30^{epgy}/Df* mutants showed a severe defect in crawling behavior. The abnormal crawling behavior was rescued by expressing a *Sod2 transgene* in the muscle or rearing the larvae in a media containing NACA. Moreover, neuronally depleting *NDUFS7* did not show any notable change in crawling defects. Minimum 10 animals were analyzed for crawling behavioral analysis. *$p < 0.05$ (mEPSP amplitude: Muscle *NDUFS7*[RNAi] +10% EtOH vs Muscle *NDUFS7*[RNAi] + 0.5 mM NACA),**$p=0.006$ (mEPSP amplitude: *ND-30* mutant vs *N-30* mutant + 0.5

mM NACA), *p=0.0004 (mEPSP amplitude: Muscle-Gal4+0.5% DMSO vs. Muscle *Cat* OE + 50 μM ro-

tenone),*p=0.039 (mEPSP amplitude: Muscle Gal4+50mM rotenone vs Muscle *Sod2* OE+ 50 μM rote-

none), *p=0.001 (EPSP amplitude: Muscle *NDUFS7[RNAi]* vs *Sod2* muscle rescue),**p=0.003 (EPSP

amplitude: Muscle Gal4+10% EtOH vs Muscle *NDUFS7[RNAi]* +0.5mM curcumin), **p=0.0004 (EPSP

amplitude: Muscle *NDUFS7*[RNAi] +10% EtOH vs Muscle *NDUFS7*[RNAi] +0.5mM NACA), *p=0.004

(EPSP amplitude: Muscle Gal4+0.5% DMSO vs Muscle Gal4+50 μM rotenone), *p=0.015 (EPSP ampli-

tude: Muscle-Gal4+0.5% DMSO vs Muscle *cat* OE+50 μM rotenone),**p=0.015 (EPSP amplitude: Mus-

cle-Gal4+0.5% DMSO vs Muscle *sod1* OE+50 μM rotenone), **p=0.001 (QC*: ND-30* mutant vs *sod2*

muscle rescue),**p=0.003 (QC: Muscle Gal4+50 μM rotenone vs Muscle *Sod*2 OE+50 μM rotenone),

**p=0.0002 (Distance crawled: Muscle Gal4 vs Muscle *ND-30*[RNAi]), ***p <0.0001; ns, not significant.

Statistical analysis based on one-way ANOVA with post-hoc Tukey's test for multiple and Student's t-

test for pairwise comparison.

1000

1001 **TABLES**

1002 All tables contain raw data supporting the graphs in the figures and are housed in the supplemen-
1003 tary section.

1004

1005 **STAR METHODS**

1006

Experimental Model Details

***Drosophila* husbandry**

Drosophila melanogaster was cultured on a standard cornmeal media containing molasses and

yeast prepared according to the Bloomington *Drosophila* Stock Center (BDSC, Bloomington, IN)

recipe. Fruit fly husbandry was performed according to standard practices [126]. Both male and fe-

male larvae were used for this study. Larvae were raised at 25°C in humidity-controlled and light-

controlled Percival DR-36VL incubators (Geneva Scientific). For drug feeding experiments in larvae,

the control, RNAi, and mutant animals were grown throughout life (egg laying to analysis) in media

containing the relevant drug. Those drugs were 0.5 mM curcumin (Sigma Aldrich), 0.5 mM N-acetyl

cysteine amide (NACA) (Sigma Aldrich), or 50 μM rotenone (Sigma Aldrich).

1017

***Drosophila* stocks**

w^{1118} was used as a non-transgenic wild-type stock [127]. The GAL4 drivers used in this study were

elaV^{C155}*-Gal4* [128], *BG57-Gal4* [64], and *D42-Gal4* [129]. For the neuronal experiments, a switch from

D42-Gal4 (motor neuron) to *elaV^C155^-Gal4* (pan-neuronal) was made between Figures 4 and 5 for

practicality of genetic stock building with several markers in the latter experiments. The core neu-

ronal results remained unchanged.

Several *UAS-RNAi* and genetic mutant lines were obtained from the Bloomington *Drosophi-*

la stock center or directly from researchers who produced them (Table 18 includes the Bloomington

Line (BL) numbers). We verified the effectiveness of *UAS-ND-20L[RNAi]* (*ND-20L^HMJ23777^*, now

termed *UAS-NDUFS7[RNAi]*) in this study. Other mutant and transgenic lines have been character-

ized in prior studies. These include: *UAS-IP₃-sponge* (*UAS-IP₃-Sponge.m30* shared by Dr. Ma-

sayuki Koganezawa's lab) [126,130]; *UAS-Catalase* [131,132]; *UAS-Sod1* [133]; *UAS-Sod2* [47]; *UAS-*

Sod2[RNAi] [134]; *Df(3L)ED4288* [135]; *ND-30^epgy^* [136]; *UAS-ND-30[RNAi]* [12]; *UAS-mitoGFP* [137]; *UAS-*

marf[RNAi] [31]; *UAS-drp1[RNAi]* [138]; *UAS-CalX[RNAi]* [139]; *UAS-mcu[RNAi]* [140,141]; *UAS-hexokinase*

A[RNAi] [142]; *UAS-hexokinase C[RNAi]* [142]; *UAS-idh[RNAi]* [143]; *UAS-Cit.(Si)-Syn[RNAi]* [144]; *UAS-*

Scsα[RNAi] [145].

1034

Method Details

Immunohistochemistry

Wondering third instar larvae were dissected and fixed on a sylgard Petri plate in ice-cold HL3 and

fixed in 4% paraformaldehyde in PBS for 30 minutes or in Bouin's fixative for 2 minutes as de-

scribed earlier [146]. Briefly, larvae were washed with PBS containing 0.2% Triton X-100 (PBST) for

30 min, blocked for an hour with 5% normal goat serum in PBST, and incubated overnight in prima-

ry antibodies at 4°C followed by washes and incubation in secondary antibodies. Monoclonal anti-

bodies: anti-Dlg (4F3), anti-DGluRIIA (8B4D2), anti-CSP (ab49), anti-Synapsin (3C11), anti-Futsch

(22C10), anti-Bruchpilot (nC82) and anti-α-Spectrin (3A9) were obtained from the Developmental

Studies Hybridoma Bank (University of Iowa, USA) and were used at 1:30 dilution. Rabbit anti-GFP

(Abcam) was used at 1:200 dilutions. Anti-GluRIII (1:100) ([147]; Table 18) was gifted by Aaron

DiAntonio (Washington University, St. Louis, U.S.A.). Fluorophore coupled secondary antibodies

Alexa Fluor 488, Alexa Fluor 568 or Alexa Fluor 647 (Molecular Probes, ThermoFisher Scientific)

were used at 1:400 dilution. Alexa 488 or 647 and Rhodamine conjugated anti-HRP were used at

1:800 and 1:600 dilutions, respectively. The larval preparations were mounted in VECTASHIELD

(Vector Laboratories, USA) and imaged with a laser scanning confocal microscope (LSM 710; Carl

Zeiss). All the images were processed with Adobe Photoshop 7.0 (Adobe Systems, San Jose, CA).

1052

1053   **Confocal imaging, quantification, and morphometric analysis**

Samples were imaged using a Carl Zeiss scanning confocal microscope equipped with 63×/1.4 NA oil immersion objective using separate channels with four laser lines (405, 488, 561, and 637 nm) at room temperature. The stained NMJ boutons were counted using anti-Synapsin or anti-HRP co-stained with anti-Dlg on muscle 6/7 of A2 hemi segment, considering each Synapsin or HRP punctum to be a bouton. At least 8 NMJs were used for bouton number quantification. For fluorescence quantifications of GluRs, Dlg, Brp, α-Spectrin, HRP, and CSP, all genotypes were immunostained in the same tube with identical reagents, then mounted and imaged in the same session. Z-stacks were obtained using identical settings for all genotypes with z-axis spacing between 0.2-0.5 μm and optimized for detection without saturation of the signal.

ROS and rotenone incubation assay

Larvae were dissected in ice-cold calcium-free HL3 to label ROS in neurons. ROS levels were detected in mitochondria by incubating live preparation in 1X Schneider's media with MitoSOX Red (Molecular Probes, ThermoFisher Scientific) fluorogenic dye at 1:200 dilutions for 20-30 minutes. Briefly, larvae were washed with HL3, mounted in VECTASHIELD (Vector Laboratories, USA.), and immediately imaged in a laser scanning confocal microscope (LSM 710; Carl Zeiss). To study the effect of DMSO and rotenone (Sigma Aldrich) in BRP, larvae were dissected in HL3 and incubated in 1X Schneider's media containing either 500 μM of rotenone or DMSO. After every 30 minutes, the old media was replaced with fresh media containing rotenone or DMSO. The above preparations were fixed with 4% paraformaldehyde, stained with anti-nc82 antibodies, mounted and imaged in a confocal microscope.

Electrophysiology and pharmacology

All dissections and recordings were performed in modified HL3 saline [148] containing 70 mM NaCl, 5 mM KCl, 10 mM MgCl2, 10 mM NaHCO3, 115 mM sucrose, 4.2 mM trehalose, 5 mM HEPES, and 0.5 mM CaCl2 (unless otherwise noted), pH 7.2. Neuromuscular junction sharp electrode (electrode resistance between 20-30 MΩ) recordings were performed on muscles 6/7 of abdominal segments A2 and A3 in wandering third-instar larvae as described [126]. Recordings were performed on a Leica microscope using a 10x objective and acquired using an Axoclamp 900A amplifier, Digidata 1440A acquisition system, and pClamp 10.7 software (Molecular Devices). Electrophysiological sweeps were digitized at 10 kHz and filtered at 1 kHz. Data were analyzed using Clampfit (Molecular Devices) and MiniAnalysis (Synaptosoft) software. Miniature excitatory postsynaptic potentials (mEPSPs) were recorded in the absence of any stimulation and motor axons were stimulated to elicit excitatory postsynaptic potentials (EPSPs).

1088

Larval crawling assay

1089

Vials containing third instar larvae were analyzed for this assay. Vials were poured with 4 ml of 20%

sucrose solution and left for 10 min to let the larvae float on top. Floating third instar animals were

poured into a petri dish and washed gently twice with deionized water in a paintbrush. A minimum

of 10 larvae of each genotype were analyzed on a 2% agarose gel in a petri dish with gridline mark-

ings 1 cm on graph paper. The larvae were acclimatized in the petri dish before videotaping. The

average distance crawled (in centimeters) by larvae was calculated based on the average number

of gridlines passed in 30 sec [146].

1097

Real-time PCR (RT PCR)

1098

Total RNA was extracted from 10 larval muscle fillets or 20 larval brains for each experiment using

TRIzol (GIBCO) according to the manufacturer's instructions. Concentration and purity of total RNA

were measured using a NanoDrop. 1 µg of total RNA was then subjected to DNA digestion using

DNase I (Ambion). This was immediately followed by reverse transcription using the iScript Reverse

Transcription Supermix (1708841, Bio-Rad). qPCR was performed using the StepOnePlus instru-

ment (ThermoFisher Scientific) and SYBR Green Supermix (172-5270, Bio-Rad) by following the

manufacturer's instructions. The Step One Software analyzed the qPCR, and the relative expres-

sion level was presented as the ratio of the target gene to the internal standard gene, *rp49*. Each

sample was analyzed in triplicate. At least three independent biological repeats were obtained for

each experiment. The following primers were used for quantitative PCR (qPCR) analysis. For the

ND-20L gene, forward: 5' CATGCCGGTGTACGATTACC 3' and reverse: 5'

CGTCCCCAGTTTAGCAGGTC 3' primers are used. Similarly, for analyzing the *ND-20* transcript,

forward: 5' GAAGTGGCCCAAAATCTGCC 3' and reverse: 5' GAGCAGATCGTCCAGTCTGG 3'

primers were used respectively.

1113

Seahorse analysis for oxygen consumption rate (OCR) measurement

1114

The Seahorse analysis was carried out as previously described [149] with some modifications. Brief-

ly, mitochondria were isolated from the thoracic muscles of 20 adult *Drosophila melanogaster* using

a differential centrifugation method. Thoraces were homogenized in cold isolation buffer, followed

by a 300 × g spin to remove debris and a subsequent 3,000 × g spin to pellet mitochondria. The fi-

nal mitochondrial pellet was resuspended in mitochondrial isolation buffer, and protein concentra-

tion was quantified using a BCA assay. Isolated mitochondria from *Drosophila melanogaster* thorac-

ic muscles were resuspended in Agilent Seahorse XF assay medium supplemented with glucose

(10 mM), pyruvate (1 mM), and glutamine (2 mM), pH 7.4. For the Seahorse assay, ~5–10 µg of

isolated mitochondria were added to each well of a Seahorse XF Cell Culture Microplate. After cen-

trifugation (3,000 × g for 20 min at 4°C), the plate was then loaded into the Seahorse XFeMini Ana-

lyzer, and the mitochondrial respiration profile was measured. The OCR data were normalized to

protein concentration.

1127

Quantification And Statistical Analysis

Quantification of mitochondrial branch length

Controls and[RNAi] -depleted animals were analyzed using mitochondrial marker Mito-GFP, *D42*

driver in cell bodies from larvae ventral nerve cord. All images were processed using image J soft-

ware. Z-stacks of individual neurons were merged. The Mito-GFP signal was enhanced by adjusting

brightness and contrast. The binary masks were created using Image>Adjust>Threshold., Meth-

od:Otsu, and Background:Dark. The branches were generated using Process>Binary>Skeletonised

[150]. These skeletonized images were analyzed, and branch length of the individual cluster was

manually calculated using image J tools.

1137

Imaging quantifications

Maximum intensity projections were used for quantitative image analysis with the Image J software

(National Institutes of Health) analysis toolkit. Boutons from muscle 4 or type Ib terminal boutons on

the muscle 6/7 of A2 hemi segment from at least six NMJ synapses were used for quantification

using Image J software. Student's t-test for pairwise and one-way ANOVA with post-hoc Tukey's

test for multiple comparisons was used for statistical analysis, using GraphPad Prism Software.

Specific p-value and tests are noted in the figures and figure legends and supplementary files and

shown in graphs as follows: * $p<0.05$, ** $p<0.001$, and *** $p<0.0001$. The data are presented as

mean ± s.e.m. For quantification of Futsch loops, third instar larval preparations were double

immunostained with HRP, 22C10 and images were captured in Zeiss LSM710 confocal micro-

scope. Only NMJs of muscles 6/7 of A2 hemi segment were used for quantification. The images

were digitally magnified using image J software and the total number of HRP-positive boutons was

manually counted in each image. Futsch positive loops, which are co-localized with HRP, were in-

cluded in this analysis. Images with incomplete loops and diffused staining were not included in the

count [151].

1153

1154 **Electrophysiological analysis**

Average mEPSP, EPSP, and quantal content were calculated for each genotype by dividing EPSP amplitude by mEPSP amplitude. Muscle input resistance (R_{in}) and resting membrane potential (V_{rest}) were monitored during each experiment. Recordings were rejected if the V_{rest} was above -60 mV, and R_{in} was less than 5 MΩ. Pharmacological agents were bath applied in recording saline at the final concentrations indicated in the text, figures, and tables. The agents included Xestospongin C (Abcam), Dantrolene (Tocris) and BAPTA-AM (Sigma Aldrich). Failure analysis was performed in HL3 solution containing 0.1 mM CaCl2, which resulted in failures in about half of the stimulated responses in wild-type larvae. A total of 30 trials (stimulations) were performed at each NMJ in all genotypes. The failure rate was obtained by dividing the total number of failures by the total number of trials (100). High-frequency (10 Hz) recordings were performed at a calcium concentration of 2 mM and paired-pulse recordings (10 Hz) were performed at calcium concentrations of 0.4 mM and 1.5 mM, respectively. Paired-pulse ratios were calculated as the EPSP amplitude of the second response divided by the first response (EPSP2/EPSP1).

REFERENCES

1. Chamberlain, K.A., and Sheng, Z.H. (2019). Mechanisms for the maintenance and regulation of axonal energy supply. Journal of neuroscience research *97*, 897-913. 10.1002/jnr.24411.

2. Hall, C.N., Klein-Flugge, M.C., Howarth, C., and Attwell, D. (2012). Oxidative phosphorylation, not glycolysis, powers presynaptic and postsynaptic mechanisms underlying brain information processing. J Neurosci *32*, 8940-8951. 10.1523/JNEUROSCI.0026-12.2012.

3. Neupane, P., Bhuju, S., Thapa, N., and Bhattarai, H.K. (2019). ATP Synthase: Structure, Function and Inhibition. Biomol Concepts *10*, 1-10. 10.1515/bmc-2019-0001.

4. Brown, G.C. (1992). Control of respiration and ATP synthesis in mammalian mitochondria and cells. Biochem J *284 (Pt 1)*, 1-13. 10.1042/bj2840001.

5. Faria-Pereira, A., and Morais, V.A. (2022). Synapses: The Brain's Energy-Demanding Sites. International journal of molecular sciences *23*. 10.3390/ijms23073627.

6. Vandecasteele, G., Szabadkai, G., and Rizzuto, R. (2001). Mitochondrial calcium homeostasis: mechanisms and molecules. IUBMB Life *52*, 213-219. 10.1080/15216540152846028.

7. Mattson, M.P., Gleichmann, M., and Cheng, A. (2008). Mitochondria in neuroplasticity and neurological disorders. Neuron *60*, 748-766. 10.1016/j.neuron.2008.10.010.

8. Hamanaka, R.B., and Chandel, N.S. (2010). Mitochondrial reactive oxygen species regulate cellular signaling and dictate biological outcomes. Trends Biochem Sci *35*, 505-513. 10.1016/j.tibs.2010.04.002.

9.   Kastaniotis, A.J., Autio, K.J., Keratar, J.M., Monteuuis, G., Makela, A.M., Nair, R.R.,
Pietikainen, L.P., Shvetsova, A., Chen, Z., and Hiltunen, J.K. (2017). Mitochondrial fatty acid
synthesis, fatty acids and mitochondrial physiology. Biochim Biophys Acta Mol Cell Biol
Lipids *1862*, 39-48. 10.1016/j.bbalip.2016.08.011.

10.   Banoth, B., and Cassel, S.L. (2018). Mitochondria in innate immune signaling. Transl Res
202, 52-68. 10.1016/j.trsl.2018.07.014.

11.   Duarte, M., Sousa, R., and Videira, A. (1995). Inactivation of genes encoding subunits of the
peripheral and membrane arms of neurospora mitochondrial complex I and effects on
enzyme assembly. Genetics *139*, 1211-1221. 10.1093/genetics/139.3.1211.

12.   Garcia, C.J., Khajeh, J., Coulanges, E., Chen, E.I., and Owusu-Ansah, E. (2017).
Regulation of Mitochondrial Complex I Biogenesis in Drosophila Flight Muscles. Cell reports
20, 264-278. 10.1016/j.celrep.2017.06.015.

13.   Mallik, B., and Frank, C.A. (2022). Roles for Mitochondrial Complex I Subunits in Regulating
Synaptic Transmission and Growth. Front Neurosci *16*, 846425. 10.3389/fnins.2022.846425.

14.   Sharma, L.K., Lu, J., and Bai, Y. (2009). Mitochondrial respiratory complex I: structure,
function and implication in human diseases. Curr Med Chem *16*, 1266-1277.
10.2174/092986709787846578.

15.   Yang, J.L., Mukda, S., and Chen, S.D. (2018). Diverse roles of mitochondria in ischemic
stroke. Redox Biol *16*, 263-275. 10.1016/j.redox.2018.03.002.

16.   Marin, S.E., Mesterman, R., Robinson, B., Rodenburg, R.J., Smeitink, J., and Tarnopolsky,
M.A. (2013). Leigh syndrome associated with mitochondrial complex I deficiency due to
novel mutations In NDUFV1 and NDUFS2. Gene *516*, 162-167.
10.1016/j.gene.2012.12.024.

17.   Goto, Y. (2000). Mitochondrial encephalomyopathy. Neuropathology *20 Suppl*, S82-84.
10.1046/j.1440-1789.2000.00304.x.

18.   Shoffner, J.M. (2000). Mitochondrial myopathy diagnosis. Neurol Clin *18*, 105-123.
10.1016/s0733-8619(05)70180-8.

19.   Burman, J.L., Itsara, L.S., Kayser, E.B., Suthammarak, W., Wang, A.M., Kaeberlein, M.,
Sedensky, M.M., Morgan, P.G., and Pallanck, L.J. (2014). A Drosophila model of
mitochondrial disease caused by a complex I mutation that uncouples proton pumping from
electron transfer. Dis Model Mech *7*, 1165-1174. 10.1242/dmm.015321.

20.   Schon, E.A., and Manfredi, G. (2003). Neuronal degeneration and mitochondrial
dysfunction. The Journal of clinical investigation *111*, 303-312. 10.1172/JCI17741.

21.   He, L., He, T., Farrar, S., Ji, L., Liu, T., and Ma, X. (2017). Antioxidants Maintain Cellular
Redox Homeostasis by Elimination of Reactive Oxygen Species. Cell Physiol Biochem *44*,
532-553. 10.1159/000485089.

22.   Folbergrova, J., and Kunz, W.S. (2012). Mitochondrial dysfunction in epilepsy.
Mitochondrion *12*, 35-40. 10.1016/j.mito.2011.04.004.

23.   Lanza, I.R., and Nair, K.S. (2010). Mitochondrial function as a determinant of life span.
Pflugers Arch *459*, 277-289. 10.1007/s00424-009-0724-5.

24.   Martin, L.J. (2012). Biology of mitochondria in neurodegenerative diseases. Prog Mol Biol
Transl Sci *107*, 355-415. 10.1016/B978-0-12-385883-2.00005-9.

25.   Curcio, R., Frattaruolo, L., Marra, F., Pesole, G., Vozza, A., Cappello, A.R., Fiorillo, M.,
Lauria, G., Ahmed, A., Fiermonte, G., et al. (2024). Two functionally different mitochondrial
phosphate carriers support Drosophila melanogaster OXPHOS throughout distinct
developmental stages. Biochim Biophys Acta Mol Cell Res *1871*, 119615.
10.1016/j.bbamcr.2023.119615.

26.   Krause, S.A., Overend, G., Dow, J.A.T., and Leader, D.P. (2022). FlyAtlas 2 in 2022:
enhancements to the Drosophila melanogaster expression atlas. Nucleic Acids Res *50*,
D1010-D1015. 10.1093/nar/gkab971.

27.   Leader, D.P., Krause, S.A., Pandit, A., Davies, S.A., and Dow, J.A.T. (2018). FlyAtlas 2: a
new version of the Drosophila melanogaster expression atlas with RNA-Seq, miRNA-Seq
and sex-specific data. Nucleic Acids Res *46*, D809-D815. 10.1093/nar/gkx976.

28.   Pilling, A.D., Horiuchi, D., Lively, C.M., and Saxton, W.M. (2006). Kinesin-1 and Dynein are
the primary motors for fast transport of mitochondria in Drosophila motor axons. Molecular
biology of the cell *17*, 2057-2068. 10.1091/mbc.E05-06-0526.

29.   Misgeld, T., and Schwarz, T.L. (2017). Mitostasis in Neurons: Maintaining Mitochondria in an
Extended Cellular Architecture. Neuron *96*, 651-666. 10.1016/j.neuron.2017.09.055.

30.   Sheng, Z.H. (2014). Mitochondrial trafficking and anchoring in neurons: New insight and
implications. J Cell Biol *204*, 1087-1098. 10.1083/jcb.201312123.

31.   Sandoval, H., Yao, C.K., Chen, K., Jaiswal, M., Donti, T., Lin, Y.Q., Bayat, V., Xiong, B.,
Zhang, K., David, G., et al. (2014). Mitochondrial fusion but not fission regulates larval
growth and synaptic development through steroid hormone production. eLife *3*.
10.7554/eLife.03558.

32.   Ge, Y., Shi, X., Boopathy, S., McDonald, J., Smith, A.W., and Chao, L.H. (2020). Two forms
of Opa1 cooperate to complete fusion of the mitochondrial inner-membrane. eLife *9*.
10.7554/eLife.50973.

33.   Song, Z., Ghochani, M., McCaffery, J.M., Frey, T.G., and Chan, D.C. (2009). Mitofusins and
OPA1 mediate sequential steps in mitochondrial membrane fusion. Molecular biology of the
cell *20*, 3525-3532. 10.1091/mbc.E09-03-0252.

34.   Smirnova, E., Griparic, L., Shurland, D.L., and van der Bliek, A.M. (2001). Dynamin-related
protein Drp1 is required for mitochondrial division in mammalian cells. Molecular biology of
the cell *12*, 2245-2256. 10.1091/mbc.12.8.2245.

35.   Twig, G., Elorza, A., Molina, A.J., Mohamed, H., Wikstrom, J.D., Walzer, G., Stiles, L.,
Haigh, S.E., Katz, S., Las, G., et al. (2008). Fission and selective fusion govern
mitochondrial segregation and elimination by autophagy. EMBO J *27*, 433-446.
10.1038/sj.emboj.7601963.

36.   Breuer, M.E., Koopman, W.J., Koene, S., Nooteboom, M., Rodenburg, R.J., Willems, P.H.,
and Smeitink, J.A. (2013). The role of mitochondrial OXPHOS dysfunction in the
development of neurologic diseases. Neurobiology of disease *51*, 27-34.
10.1016/j.nbd.2012.03.007.

37.   Berger, I., Hershkovitz, E., Shaag, A., Edvardson, S., Saada, A., and Elpeleg, O. (2008).
Mitochondrial complex I deficiency caused by a deleterious NDUFA11 mutation. Ann Neurol
63, 405-408. 10.1002/ana.21332.

38.   Formosa, L.E., Muellner-Wong, L., Reljic, B., Sharpe, A.J., Jackson, T.D., Beilharz, T.H.,
Stojanovski, D., Lazarou, M., Stroud, D.A., and Ryan, M.T. (2020). Dissecting the Roles of
Mitochondrial Complex I Intermediate Assembly Complex Factors in the Biogenesis of
Complex I. Cell reports *31*, 107541. 10.1016/j.celrep.2020.107541.

39.   Guerrero-Castillo, S., Baertling, F., Kownatzki, D., Wessels, H.J., Arnold, S., Brandt, U., and
Nijtmans, L. (2017). The Assembly Pathway of Mitochondrial Respiratory Chain Complex I.
Cell metabolism *25*, 128-139. 10.1016/j.cmet.2016.09.002.

40.   Stroud, D.A., Surgenor, E.E., Formosa, L.E., Reljic, B., Frazier, A.E., Dibley, M.G.,
Osellame, L.D., Stait, T., Beilharz, T.H., Thorburn, D.R., et al. (2016). Accessory subunits
are integral for assembly and function of human mitochondrial complex I. Nature *538*, 123-
126. 10.1038/nature19754.

41.   Kauffman, M.E., Kauffman, M.K., Traore, K., Zhu, H., Trush, M.A., Jia, Z., and Li, Y.R.
(2016). MitoSOX-Based Flow Cytometry for Detecting Mitochondrial ROS. React Oxyg
Species (Apex) *2*, 361-370. 10.20455/ros.2016.865.

42.   Robinson, K.M., Janes, M.S., Pehar, M., Monette, J.S., Ross, M.F., Hagen, T.M., Murphy,
M.P., and Beckman, J.S. (2006). Selective fluorescent imaging of superoxide in vivo using
ethidium-based probes. Proc Natl Acad Sci U S A *103*, 15038-15043.
10.1073/pnas.0601945103.

43.   Little, A.C., Kovalenko, I., Goo, L.E., Hong, H.S., Kerk, S.A., Yates, J.A., Purohit, V.,
Lombard, D.B., Merajver, S.D., and Lyssiotis, C.A. (2020). High-content fluorescence
imaging with the metabolic flux assay reveals insights into mitochondrial properties and
functions. Commun Biol *3*, 271. 10.1038/s42003-020-0988-z.

44.   Martelli, F., Hernandes, N.H., Zuo, Z., Wang, J., Wong, C.O., Karagas, N.E., Roessner, U.,
Rupasinghe, T., Robin, C., Venkatachalam, K., et al. (2022). Low doses of the organic
insecticide spinosad trigger lysosomal defects, elevated ROS, lipid dysregulation, and
neurodegeneration in flies. eLife *11*. 10.7554/eLife.73812.

45.   Martelli, F., Zhongyuan, Z., Wang, J., Wong, C.O., Karagas, N.E., Roessner, U.,
Rupasinghe, T., Venkatachalam, K., Perry, T., Bellen, H.J., and Batterham, P. (2020). Low
doses of the neonicotinoid insecticide imidacloprid induce ROS triggering neurological and
metabolic impairments in Drosophila. Proc Natl Acad Sci U S A *117*, 25840-25850.
10.1073/pnas.2011828117.

46.   Penugonda, S., Mare, S., Lutz, P., Banks, W.A., and Ercal, N. (2006). Potentiation of lead-
induced cell death in PC12 cells by glutamate: protection by N-acetylcysteine amide

(NACA), a novel thiol antioxidant. Toxicol Appl Pharmacol *216*, 197-205.
10.1016/j.taap.2006.05.002.

47.   Missirlis, F., Hu, J., Kirby, K., Hilliker, A.J., Rouault, T.A., and Phillips, J.P. (2003).
Compartment-specific protection of iron-sulfur proteins by superoxide dismutase. J Biol
Chem *278*, 47365-47369. 10.1074/jbc.M307700200.

48.   Oswald, M.C.W., Garnham, N., Sweeney, S.T., and Landgraf, M. (2018). Regulation of
neuronal development and function by ROS. FEBS letters *592*, 679-691. 10.1002/1873-
3468.12972.

49.   Roos, J., Hummel, T., Ng, N., Klambt, C., and Davis, G.W. (2000). Drosophila Futsch
regulates synaptic microtubule organization and is necessary for synaptic growth. Neuron
26, 371-382.

50.   Guo, X., Macleod, G.T., Wellington, A., Hu, F., Panchumarthi, S., Schoenfield, M., Marin, L.,
Charlton, M.P., Atwood, H.L., and Zinsmaier, K.E. (2005). The GTPase dMiro is required for
axonal transport of mitochondria to Drosophila synapses. Neuron *47*, 379-393.
10.1016/j.neuron.2005.06.027.

51.   Verstreken, P., Ly, C.V., Venken, K.J., Koh, T.W., Zhou, Y., and Bellen, H.J. (2005).
Synaptic mitochondria are critical for mobilization of reserve pool vesicles at Drosophila
neuromuscular junctions. Neuron *47*, 365-378. 10.1016/j.neuron.2005.06.018.

52.   Milton, V.J., Jarrett, H.E., Gowers, K., Chalak, S., Briggs, L., Robinson, I.M., and Sweeney,
S.T. (2011). Oxidative stress induces overgrowth of the Drosophila neuromuscular junction.
Proc Natl Acad Sci U S A *108*, 17521-17526. 10.1073/pnas.1014511108.

53.   Oswald, M.C., Brooks, P.S., Zwart, M.F., Mukherjee, A., West, R.J., Giachello, C.N.,
Morarach, K., Baines, R.A., Sweeney, S.T., and Landgraf, M. (2018). Reactive oxygen
species regulate activity-dependent neuronal plasticity in Drosophila. eLife *7*.
10.7554/eLife.39393.

54.   Böhme, M.A., McCarthy, A.W., Grasskamp, A.T., Beuschel, C.B., Goel, P., Jusyte, M.,
Laber, D., Huang, S., Rey, U., Petzoldt, A.G., et al. (2019). Rapid active zone remodeling
consolidates presynaptic potentiation. Nat Commun *10*, 1085. 10.1038/s41467-019-08977-
6.

55.   Goel, P., Dufour Bergeron, D., Bohme, M.A., Nunnelly, L., Lehmann, M., Buser, C., Walter,
A.M., Sigrist, S.J., and Dickman, D. (2019). Homeostatic scaling of active zone scaffolds
maintains global synaptic strength. J Cell Biol *218*, 1706-1724. 10.1083/jcb.201807165.

56.   Gratz, S.J., Goel, P., Bruckner, J.J., Hernandez, R.X., Khateeb, K., Macleod, G.T., Dickman,
D., and O'Connor-Giles, K.M. (2019). Endogenous Tagging Reveals Differential Regulation
of Ca(2+) Channels at Single Active Zones during Presynaptic Homeostatic Potentiation and
Depression. J Neurosci *39*, 2416-2429. 10.1523/JNEUROSCI.3068-18.2019.

57.   Hong, H., Zhao, K., Huang, S., Huang, S., Yao, A., Jiang, Y., Sigrist, S., Zhao, L., and
Zhang, Y.Q. (2020). Structural Remodeling of Active Zones Is Associated with Synaptic
Homeostasis. J Neurosci *40*, 2817-2827. 10.1523/JNEUROSCI.2002-19.2020.

58. Dannhauser, S., Mrestani, A., Gundelach, F., Pauli, M., Komma, F., Kollmannsberger, P.,
Sauer, M., Heckmann, M., and Paul, M.M. (2022). Endogenous tagging of Unc-13 reveals
nanoscale reorganization at active zones during presynaptic homeostatic potentiation.
Frontiers in cellular neuroscience *16*, 1074304. 10.3389/fncel.2022.1074304.

59. Ashrafi, G., de Juan-Sanz, J., Farrell, R.J., and Ryan, T.A. (2020). Molecular Tuning of the
Axonal Mitochondrial Ca(2+) Uniporter Ensures Metabolic Flexibility of Neurotransmission.
Neuron *105*, 678-687 e675. 10.1016/j.neuron.2019.11.020.

60. Datta, S., and Jaiswal, M. (2021). Mitochondrial calcium at the synapse. Mitochondrion *59*,
135-153. 10.1016/j.mito.2021.04.006.

61. Marland, J.R., Hasel, P., Bonnycastle, K., and Cousin, M.A. (2016). Mitochondrial Calcium
Uptake Modulates Synaptic Vesicle Endocytosis in Central Nerve Terminals. J Biol Chem
291, 2080-2086. 10.1074/jbc.M115.686956.

62. Paradis, M., Kucharowski, N., Edwards Faret, G., Maya Palacios, S.J., Meyer, C.,
Stumpges, B., Jamitzky, I., Kalinowski, J., Thiele, C., Bauer, R., et al. (2022). The ER
protein Creld regulates ER-mitochondria contact dynamics and respiratory complex 1
activity. Sci Adv *8*, eabo0155. 10.1126/sciadv.abo0155.

63. Benson, T.E., and Voigt, H.F. (1995). Neuron labeling by extracellular delivery of
horseradish peroxidase in vivo: a method for studying the local circuitry of projection and
interneurons at physiologically characterized sites. J Neurosci Methods *57*, 81-91.
10.1016/0165-0270(94)00131-y.

64. Budnik, V., Koh, Y.H., Guan, B., Hartmann, B., Hough, C., Woods, D., and Gorczyca, M.
(1996). Regulation of synapse structure and function by the Drosophila tumor suppressor
gene dlg. Neuron *17*, 627-640.

65. Pielage, J., Fetter, R.D., and Davis, G.W. (2006). A postsynaptic spectrin scaffold defines
active zone size, spacing, and efficacy at the Drosophila neuromuscular junction. J Cell Biol
175, 491-503.

66. Chen, K., and Featherstone, D.E. (2005). Discs-large (DLG) is clustered by presynaptic
innervation and regulates postsynaptic glutamate receptor subunit composition in
Drosophila. BMC Biol *3*, 1. 10.1186/1741-7007-3-1.

67. Accardi, M.V., Daniels, B.A., Brown, P.M., Fritschy, J.M., Tyagarajan, S.K., and Bowie, D.
(2014). Mitochondrial reactive oxygen species regulate the strength of inhibitory GABA-
mediated synaptic transmission. Nat Commun *5*, 3168. 10.1038/ncomms4168.

68. Doser, R.L., Amberg, G.C., and Hoerndli, F.J. (2020). Reactive Oxygen Species Modulate
Activity-Dependent AMPA Receptor Transport in C. elegans. J Neurosci *40*, 7405-7420.
10.1523/JNEUROSCI.0902-20.2020.

69. Peng, J.J., Lin, S.H., Liu, Y.T., Lin, H.C., Li, T.N., and Yao, C.K. (2019). A circuit-dependent
ROS feedback loop mediates glutamate excitotoxicity to sculpt the Drosophila motor system.
eLife *8*. 10.7554/eLife.47372.

70. Wang, X., Winter, D., Ashrafi, G., Schlehe, J., Wong, Y.L., Selkoe, D., Rice, S., Steen, J., LaVoie, M.J., and Schwarz, T.L. (2011). PINK1 and Parkin target Miro for phosphorylation and degradation to arrest mitochondrial motility. Cell *147*, 893-906. 10.1016/j.cell.2011.10.018.

71. Wang, X., and Schwarz, T.L. (2009). The mechanism of Ca2+ -dependent regulation of kinesin-mediated mitochondrial motility. Cell *136*, 163-174. 10.1016/j.cell.2008.11.046.

72. Clark, I.E., Dodson, M.W., Jiang, C., Cao, J.H., Huh, J.R., Seol, J.H., Yoo, S.J., Hay, B.A., and Guo, M. (2006). Drosophila pink1 is required for mitochondrial function and interacts genetically with parkin. Nature *441*, 1162-1166. 10.1038/nature04779.

73. Park, J., Lee, S.B., Lee, S., Kim, Y., Song, S., Kim, S., Bae, E., Kim, J., Shong, M., Kim, J.M., and Chung, J. (2006). Mitochondrial dysfunction in Drosophila PINK1 mutants is complemented by parkin. Nature *441*, 1157-1161. 10.1038/nature04788.

74. Poole, A.C., Thomas, R.E., Andrews, L.A., McBride, H.M., Whitworth, A.J., and Pallanck, L.J. (2008). The PINK1/Parkin pathway regulates mitochondrial morphology. Proc Natl Acad Sci U S A *105*, 1638-1643. 10.1073/pnas.0709336105.

75. Yang, Y., Gehrke, S., Imai, Y., Huang, Z., Ouyang, Y., Wang, J.W., Yang, L., Beal, M.F., Vogel, H., and Lu, B. (2006). Mitochondrial pathology and muscle and dopaminergic neuron degeneration caused by inactivation of Drosophila Pink1 is rescued by Parkin. Proc Natl Acad Sci U S A *103*, 10793-10798. 10.1073/pnas.0602493103.

76. Jiang, H.C., Hsu, J.M., Yen, C.P., Chao, C.C., Chen, R.H., and Pan, C.L. (2015). Neural activity and CaMKII protect mitochondria from fragmentation in aging Caenorhabditis elegans neurons. Proc Natl Acad Sci U S A *112*, 8768-8773. 10.1073/pnas.1501831112.

77. Obashi, K., and Okabe, S. (2013). Regulation of mitochondrial dynamics and distribution by synapse position and neuronal activity in the axon. The European journal of neuroscience *38*, 2350-2363. 10.1111/ejn.12263.

78. Lin, M.T., and Beal, M.F. (2006). Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. Nature *443*, 787-795. 10.1038/nature05292.

79. Esterberg, R., Linbo, T., Pickett, S.B., Wu, P., Ou, H.C., Rubel, E.W., and Raible, D.W. (2016). Mitochondrial calcium uptake underlies ROS generation during aminoglycoside-induced hair cell death. The Journal of clinical investigation *126*, 3556-3566. 10.1172/JCI84939.

80. Gorlach, A., Bertram, K., Hudecova, S., and Krizanova, O. (2015). Calcium and ROS: A mutual interplay. Redox Biol *6*, 260-271. 10.1016/j.redox.2015.08.010.

81. Yu, T., Robotham, J.L., and Yoon, Y. (2006). Increased production of reactive oxygen species in hyperglycemic conditions requires dynamic change of mitochondrial morphology. Proc Natl Acad Sci U S A *103*, 2653-2658. 10.1073/pnas.0511154103.

82. Rimal, S., Tantray, I., Li, Y., Pal Khaket, T., Li, Y., Bhurtel, S., Li, W., Zeng, C., and Lu, B. (2023). Reverse electron transfer is activated during aging and contributes to aging and age-related disease. EMBO reports *24*, e55548. 10.15252/embr.202255548.

83. Graham, C., Stefanatos, R., Yek, A.E.H., Spriggs, R.V., Loh, S.H.Y., Uribe, A.H., Zhang, T.,
Martins, L.M., Maddocks, O.D.K., Scialo, F., and Sanz, A. (2022). Mitochondrial ROS
signalling requires uninterrupted electron flow and is lost during ageing in flies. Geroscience
44, 1961-1974. 10.1007/s11357-022-00555-x.

84. Fei, L., Liang, Y., Kintscher, U., and Sigrist, S.J. (2025). Coupling of mitochondrial state with
active zone plasticity in early brain aging. Redox Biol *79*, 103454.
10.1016/j.redox.2024.103454.

85. Granat, L., Knorr, D.Y., Ranson, D.C., Hamer, E.L., Chakrabarty, R.P., Mattedi, F., Fort-
Aznar, L., Hirth, F., Sweeney, S.T., Vagnoni, A., et al. (2023). Yeast NDI1 reconfigures
neuronal metabolism and prevents the unfolded protein response in mitochondrial complex I
deficiency. PLoS genetics *19*, e1010793. 10.1371/journal.pgen.1010793.

86. Owusu-Ansah, E., Song, W., and Perrimon, N. (2013). Muscle mitohormesis promotes
longevity via systemic repression of insulin signaling. Cell *155*, 699-712.
10.1016/j.cell.2013.09.021.

87. Jornayvaz, F.R., and Shulman, G.I. (2010). Regulation of mitochondrial biogenesis. Essays
Biochem *47*, 69-84. 10.1042/bse0470069.

88. Shaw, J.M., and Winge, D.R. (2009). Shaping the mitochondrion: mitochondrial biogenesis,
dynamics and dysfunction. Conference on Mitochondrial Assembly and Dynamics in Health
and Disease. EMBO reports *10*, 1301-1305. 10.1038/embor.2009.247.

89. Esparza-Molto, P.B., Romero-Carraminana, I., Nunez de Arenas, C., Pereira, M.P., Blanco,
N., Pardo, B., Bates, G.R., Sanchez-Castillo, C., Artuch, R., Murphy, M.P., et al. (2021).
Generation of mitochondrial reactive oxygen species is controlled by ATPase inhibitory
factor 1 and regulates cognition. PLoS Biol *19*, e3001252. 10.1371/journal.pbio.3001252.

90. Huddleston, A.T., Tang, W., Takeshima, H., Hamilton, S.L., and Klann, E. (2008).
Superoxide-induced potentiation in the hippocampus requires activation of ryanodine
receptor type 3 and ERK. Journal of neurophysiology *99*, 1565-1571.
10.1152/jn.00659.2007.

91. Lee, K.Y., Chung, K., and Chung, J.M. (2010). Involvement of reactive oxygen species in
long-term potentiation in the spinal cord dorsal horn. Journal of neurophysiology *103*, 382-
391. 10.1152/jn.90906.2008.

92. Müller, M., Pym, E.C., Tong, A., and Davis, G.W. (2011). Rab3-GAP controls the
progression of synaptic homeostasis at a late stage of vesicle release. Neuron *69*, 749-762.
S0896-6273(11)00073-0 [pii]

10.1016/j.neuron.2011.01.025.

93. Frank, C.A., Kennedy, M.J., Goold, C.P., Marek, K.W., and Davis, G.W. (2006).
Mechanisms underlying the rapid induction and sustained expression of synaptic
homeostasis. Neuron *52*, 663-677.

94. Petersen, S.A., Fetter, R.D., Noordermeer, J.N., Goodman, C.S., and DiAntonio, A. (1997).
Genetic analysis of glutamate receptors in Drosophila reveals a retrograde signal regulating
presynaptic transmitter release. Neuron *19*, 1237-1248.

95. Frank, C.A., Pielage, J., and Davis, G.W. (2009). A presynaptic homeostatic signaling
system composed of the Eph receptor, ephexin, Cdc42, and CaV2.1 calcium channels.
Neuron *61*, 556-569.

96. Müller, M., and Davis, G.W. (2012). Transsynaptic control of presynaptic Ca(2)(+) influx
achieves homeostatic potentiation of neurotransmitter release. Curr Biol *22*, 1102-1108.
10.1016/j.cub.2012.04.018

S0960-9822(12)00410-1 [pii].

97. Wang, T., Hauswirth, A.G., Tong, A., Dickman, D.K., and Davis, G.W. (2014). Endostatin Is
a Trans-Synaptic Signal for Homeostatic Synaptic Plasticity. Neuron *83*, 616-629.
10.1016/j.neuron.2014.07.003.

98. Younger, M.A., Müller, M., Tong, A., Pym, E.C., and Davis, G.W. (2013). A presynaptic
ENaC channel drives homeostatic plasticity. Neuron *79*, 1183-1196.
10.1016/j.neuron.2013.06.048.

99. Harris, N., Braiser, D.J., Dickman, D.K., Fetter, R.D., Tong, A., and Davis, G.W. (2015). The
Innate Immune Receptor PGRP-LC Controls Presynaptic Homeostatic Plasticity. Neuron *88*,
1157-1164. 10.1016/j.neuron.2015.10.049.

100. Müller, M., Genç, Ö., and Davis, G.W. (2015). RIM-Binding Protein Links Synaptic
Homeostasis to the Stabilization and Replenishment of High Release Probability Vesicles.
Neuron *85*, 1056-1069. 10.1016/j.neuron.2015.01.024.

101. Müller, M., Liu, K.S., Sigrist, S.J., and Davis, G.W. (2012). RIM Controls Homeostatic
Plasticity through Modulation of the Readily-Releasable Vesicle Pool. J Neurosci *32*, 16574-
16585. 10.1523/JNEUROSCI.0981-12.2012

32/47/16574 [pii].

102. Weyhersmüller, A., Hallermann, S., Wagner, N., and Eilers, J. (2011). Rapid active zone
remodeling during synaptic plasticity. J Neurosci *31*, 6041-6052. 31/16/6041 [pii]

10.1523/JNEUROSCI.6698-10.2011.

103. Goel, P., Dufour Bergeron, D., Bohme, M.A., Nunnelly, L., Lehmann, M., Buser, C., Walter,
A.M., Sigrist, S.J., and Dickman, D. (2019). Homeostatic scaling of active zone scaffolds
maintains global synaptic strength. J Cell Biol. 10.1083/jcb.201807165.

104. Oliva, M.K., Perez-Moreno, J.J., O'Shaughnessy, J., Wardill, T.J., and O'Kane, C.J. (2020).
Endoplasmic Reticulum Lumenal Indicators in Drosophila Reveal Effects of HSP-Related
Mutations on Endoplasmic Reticulum Calcium Dynamics. Front Neurosci *14*, 816.
10.3389/fnins.2020.00816.

105. Dhawan, S., Myers, P., Bailey, D.M.D., Ostrovsky, A.D., Evers, J.F., and Landgraf, M.
(2021). Reactive Oxygen Species Mediate Activity-Regulated Dendritic Plasticity Through
NADPH Oxidase and Aquaporin Regulation. Frontiers in cellular neuroscience *15*, 641802.
10.3389/fncel.2021.641802.

106. Frank, C.A., James, T.D., and Müller, M. (2020). Homeostatic control of Drosophila neuromuscular junction function. Synapse *74*, e22133. 10.1002/syn.22133.

107. Goold, C.P., and Davis, G.W. (2007). The BMP ligand Gbb gates the expression of synaptic homeostasis independent of synaptic growth control. Neuron *56*, 109-123.

108. Haghighi, A.P., McCabe, B.D., Fetter, R.D., Palmer, J.E., Hom, S., and Goodman, C.S. (2003). Retrograde control of synaptic transmission by postsynaptic CaMKII at the Drosophila neuromuscular junction. Neuron *39*, 255-267.

109. McCabe, B.D., Marques, G., Haghighi, A.P., Fetter, R.D., Crotty, M.L., Haerry, T.E., Goodman, C.S., and O'Connor, M.B. (2003). The BMP homolog Gbb provides a retrograde signal that regulates synaptic growth at the Drosophila neuromuscular junction. Neuron *39*, 241-254.

110. Kikuma, K., Li, X., Perry, S., Li, Q., Goel, P., Chen, C., Kim, D., Stavropoulos, N., and Dickman, D. (2019). Cul3 and insomniac are required for rapid ubiquitination of postsynaptic targets and retrograde homeostatic signaling. Nat Commun *10*, 2998. 10.1038/s41467-019-10992-6.

111. Hauswirth, A.G., Ford, K.J., Wang, T., Fetter, R.D., Tong, A., and Davis, G.W. (2018). A postsynaptic PI3K-cII dependent signaling controller for presynaptic homeostatic plasticity. eLife *7*. 10.7554/eLife.31535.

112. Orr, B.O., Fetter, R.D., and Davis, G.W. (2017). Retrograde semaphorin-plexin signalling drives homeostatic synaptic plasticity. Nature *550*, 109-113. 10.1038/nature24017.

113. Goel, P., Li, X., and Dickman, D. (2017). Disparate Postsynaptic Induction Mechanisms Ultimately Converge to Drive the Retrograde Enhancement of Presynaptic Efficacy. Cell reports *21*, 2339-2347. 10.1016/j.celrep.2017.10.116.

114. Penney, J., Tsurudome, K., Liao, E.H., Elazzouzi, F., Livingstone, M., Gonzalez, M., Sonenberg, N., and Haghighi, A.P. (2012). TOR is required for the retrograde regulation of synaptic homeostasis at the Drosophila neuromuscular junction. Neuron *74*, 166-178. S0896-6273(12)00186-9 [pii]

10.1016/j.neuron.2012.01.030.

115. Pielage, J., Fetter, R.D., and Davis, G.W. (2005). Presynaptic spectrin is essential for synapse stabilization. Curr Biol *15*, 918-928.

116. Pielage, J., Cheng, L., Fetter, R.D., Carlton, P.M., Sedat, J.W., and Davis, G.W. (2008). A presynaptic giant ankyrin stabilizes the NMJ through regulation of presynaptic microtubules and transsynaptic cell adhesion. Neuron *58*, 195-209. 10.1016/j.neuron.2008.02.017.

117. Stephan, R., Goellner, B., Moreno, E., Frank, C.A., Hugenschmidt, T., Genoud, C., Aberle, H., and Pielage, J. (2015). Hierarchical microtubule organization controls axon caliber and transport and determines synaptic structure and stability. Dev Cell *33*, 5-21. 10.1016/j.devcel.2015.02.003.

118. McCord, J.M., and Fridovich, I. (1969). Superoxide dismutase. An enzymic function for erythrocuprein (hemocuprein). J Biol Chem *244*, 6049-6055.

1541   119.   Okado-Matsumoto, A., and Fridovich, I. (2001). Subcellular distribution of superoxide
1542   dismutases (SOD) in rat liver: Cu,Zn-SOD in mitochondria. J Biol Chem *276*, 38388-38393.
1543   10.1074/jbc.M105395200.

1544   120.   Weisiger, R.A., and Fridovich, I. (1973). Superoxide dismutase. Organelle specificity. J Biol
1545   Chem *248*, 3582-3592.

1546   121.   Baudhuin, P., Beaufay, H., and De Duve, C. (1965). Combined biochemical and
1547   morphological study of particulate fractions from rat liver. Analysis of preparations enriched
1548   in lysosomes or in particles containing urate oxidase, D-amino acid oxidase, and catalase. J
1549   Cell Biol *26*, 219-243. 10.1083/jcb.26.1.219.

1550   122.   De Duve, C., and Baudhuin, P. (1966). Peroxisomes (microbodies and related particles).
1551   Physiol Rev *46*, 323-357. 10.1152/physrev.1966.46.2.323.

1552   123.   Grinberg, L., Fibach, E., Amer, J., and Atlas, D. (2005). N-acetylcysteine amide, a novel cell-
1553   permeating thiol, restores cellular glutathione and protects human red blood cells from
1554   oxidative stress. Free radical biology & medicine *38*, 136-145.
1555   10.1016/j.freeradbiomed.2004.09.025.

1556   124.   Patel, S.P., Sullivan, P.G., Pandya, J.D., Goldstein, G.A., VanRooyen, J.L., Yonutas, H.M.,
1557   Eldahan, K.C., Morehouse, J., Magnuson, D.S., and Rabchevsky, A.G. (2014). N-
1558   acetylcysteine amide preserves mitochondrial bioenergetics and improves functional
1559   recovery following spinal trauma. Exp Neurol *257*, 95-105. 10.1016/j.expneurol.2014.04.026.

1560   125.   Pandya, J.D., Readnower, R.D., Patel, S.P., Yonutas, H.M., Pauly, J.R., Goldstein, G.A.,
1561   Rabchevsky, A.G., and Sullivan, P.G. (2014). N-acetylcysteine amide confers
1562   neuroprotection, improves bioenergetics and behavioral outcome following TBI. Exp Neurol
1563   *257*, 106-113. 10.1016/j.expneurol.2014.04.020.

1564   126.   James, T.D., Zwiefelhofer, D.J., and Frank, C.A. (2019). Maintenance of homeostatic
1565   plasticity at the Drosophila neuromuscular synapse requires continuous IP3-directed
1566   signaling. eLife *8*. 10.7554/eLife.39643.

1567   127.   Hazelrigg, T., Levis, R., and Rubin, G.M. (1984). Transformation of white locus DNA in
1568   drosophila: dosage compensation, zeste interaction, and position effects. Cell *36*, 469-481.

1569   128.   Lin, D.M., and Goodman, C.S. (1994). Ectopic and increased expression of Fasciclin II alters
1570   motoneuron growth cone guidance. Neuron *13*, 507-523.

1571   129.   Yeh, E., Gustafson, K., and Boulianne, G.L. (1995). Green fluorescent protein as a vital
1572   marker and reporter of gene expression in Drosophila. Proc Natl Acad Sci U S A *92*, 7036-
1573   7040. 10.1073/pnas.92.15.7036.

1574   130.   Usui-Aoki, K., Matsumoto, K., Koganezawa, M., Kohatsu, S., Isono, K., Matsubayashi, H.,
1575   Yamamoto, M.T., Ueda, R., Takahashi, K., Saigo, K., et al. (2005). Targeted expression of
1576   Ip3 sponge and Ip3 dsRNA impaires sugar taste sensation in Drosophila. Journal of
1577   neurogenetics *19*, 123-141. 10.1080/01677060600569713.

131. Ha, E.M., Oh, C.T., Ryu, J.H., Bae, Y.S., Kang, S.W., Jang, I.H., Brey, P.T., and Lee, W.J. (2005). An antioxidant system required for host protection against gut infection in Drosophila. Dev Cell *8*, 125-132. 10.1016/j.devcel.2004.11.007.

132. Missirlis, F., Phillips, J.P., and Jackle, H. (2001). Cooperative action of antioxidant defense systems in Drosophila. Curr Biol *11*, 1272-1277. 10.1016/s0960-9822(01)00393-1.

133. Watson, M.R., Lagow, R.D., Xu, K., Zhang, B., and Bonini, N.M. (2008). A drosophila model for amyotrophic lateral sclerosis reveals motor neuron damage by human SOD1. J Biol Chem *283*, 24972-24981. 10.1074/jbc.M804817200.

134. Kirby, K., Hu, J., Hilliker, A.J., and Phillips, J.P. (2002). RNA interference-mediated silencing of Sod2 in Drosophila leads to early adult-onset mortality and elevated endogenous oxidative stress. Proc Natl Acad Sci U S A *99*, 16162-16167. 10.1073/pnas.252342899.

135. Ryder, E., Ashburner, M., Bautista-Llacer, R., Drummond, J., Webster, J., Johnson, G., Morley, T., Chan, Y.S., Blows, F., Coulson, D., et al. (2007). The DrosDel deletion collection: a Drosophila genomewide chromosomal deficiency resource. Genetics *177*, 615-629. 10.1534/genetics.107.076216.

136. Bellen, H.J., Levis, R.W., Liao, G., He, Y., Carlson, J.W., Tsang, G., Evans-Holm, M., Hiesinger, P.R., Schulze, K.L., Rubin, G.M., et al. (2004). The BDGP gene disruption project: single transposon insertions associated with 40% of Drosophila genes. Genetics *167*, 761-781.

137. Rikhy, R., Ramaswami, M., and Krishnan, K.S. (2003). A temperature-sensitive allele of Drosophila sesB reveals acute functions for the mitochondrial adenine nucleotide translocase in synaptic transmission and dynamin regulation. Genetics *165*, 1243-1253. 10.1093/genetics/165.3.1243.

138. Zhang, Y., Liu, X., Bai, J., Tian, X., Zhao, X., Liu, W., Duan, X., Shang, W., Fan, H.Y., and Tong, C. (2016). Mitoguardin Regulates Mitochondrial Fusion through MitoPLD and Is Required for Neuronal Homeostasis. Molecular cell *61*, 111-124. 10.1016/j.molcel.2015.11.017.

139. Paglione, M., Restivo, L., Zakhia, S., Llobet Rosell, A., Terenzio, M., and Neukomm, L.J. (2025). Local translatome sustains synaptic function in impaired Wallerian degeneration. EMBO reports *26*, 61-83. 10.1038/s44319-024-00301-8.

140. Lee, K.S., Huh, S., Lee, S., Wu, Z., Kim, A.K., Kang, H.Y., and Lu, B. (2018). Altered ER-mitochondria contact impacts mitochondria calcium homeostasis and contributes to neurodegeneration in vivo in disease models. Proc Natl Acad Sci U S A *115*, E8844-E8853. 10.1073/pnas.1721136115.

141. Rodriguez, L.R., Calap-Quintana, P., Lapena-Luzon, T., Pallardo, F.V., Schneuwly, S., Navarro, J.A., and Gonzalez-Cabo, P. (2020). Oxidative stress modulates rearrangement of endoplasmic reticulum-mitochondria contacts and calcium dysregulation in a Friedreich's ataxia model. Redox Biol *37*, 101762. 10.1016/j.redox.2020.101762.

142. Dus, M., Lai, J.S., Gunapala, K.M., Min, S., Tayler, T.D., Hergarden, A.C., Geraud, E.,
Joseph, C.M., and Suh, G.S. (2015). Nutrient Sensor in the Brain Directs the Action of the
Brain-Gut Axis in Drosophila. Neuron *87*, 139-151. 10.1016/j.neuron.2015.05.032.

143. Francois, C.M., Pihl, T., Dunoyer de Segonzac, M., Herault, C., and Hudry, B. (2023).
Metabolic regulation of proteome stability via N-terminal acetylation controls male germline
stem cell differentiation and reproduction. Nat Commun *14*, 6737. 10.1038/s41467-023-
42496-9.

144. Hudry, B., de Goeij, E., Mineo, A., Gaspar, P., Hadjieconomou, D., Studd, C., Mokochinski,
J.B., Kramer, H.B., Placais, P.Y., Preat, T., and Miguel-Aliaga, I. (2019). Sex Differences in
Intestinal Carbohydrate Metabolism Promote Food Intake and Sperm Maturation. Cell *178*,
901-918 e916. 10.1016/j.cell.2019.07.029.

145. Barry, W.E., and Thummel, C.S. (2016). The Drosophila HNF4 nuclear receptor promotes
glucose-stimulated insulin secretion and mitochondrial function in adults. eLife *5*.
10.7554/eLife.11183.

146. Raut, S., Mallik, B., Parichha, A., Amrutha, V., Sahi, C., and Kumar, V. (2017). RNAi-
Mediated Reverse Genetic Screen Identified Drosophila Chaperones Regulating Eye and
Neuromuscular   Junction   Morphology.   G3   (Bethesda)   *7*,   2023-2038.
10.1534/g3.117.041632.

147. Marrus, S.B., Portman, S.L., Allen, M.J., Moffat, K.G., and DiAntonio, A. (2004). Differential
localization of glutamate receptor subunits at the Drosophila neuromuscular junction. J
Neurosci *24*, 1406-1415.

148. Stewart, B.A., Atwood, H.L., Renger, J.J., Wang, J., and Wu, C.F. (1994). Improved stability
of Drosophila larval neuromuscular preparations in haemolymph-like physiological solutions.
J Comp Physiol A *175*, 179-191. 10.1007/BF00215114.

149. Groen, C.M., and Windebank, A.J. (2025). Isolation of Intact Mitochondria From Drosophila
melanogaster and Assessment of Mitochondrial Respiratory Capacity Using Seahorse
Analyzer. Bio Protoc *15*, e5180. 10.21769/BioProtoc.5180.

150. Terriente-Felix, A., Wilson, E.L., and Whitworth, A.J. (2020). Drosophila phosphatidylinositol-
4 kinase fwd promotes mitochondrial fission and can suppress Pink1/parkin phenotypes.
PLoS genetics *16*, e1008844. 10.1371/journal.pgen.1008844.

151. Choudhury, S.D., Mushtaq, Z., Reddy-Alla, S., Balakrishnan, S.S., Thakur, R.S., Krishnan,
K.S., Raghu, P., Ramaswami, M., and Kumar, V. (2016). sigma2-Adaptin Facilitates Basal
Synaptic Transmission and Is Required for Regenerating Endo-Exo Cycling Pool Under
High-Frequency   Nerve   Stimulation   in   Drosophila.   Genetics   *203*,   369-385.
10.1534/genetics.115.183863.

1651

Control

A — mito-GFP
A' merged — mito-GFP / BRP
B — mito-GFP
B' merged — mito-GFP / BRP
C merged A2 — mito-GFP / Futsch
C' — mito-GFP
D merged A5 — mito-GFP / Futsch
D' — mito-GFP

NDUFS7 RNAi

E — mito-GFP
E' merged — mito-GFP / BRP
F — mito-GFP
F' merged — mito-GFP / BRP
G merged A2 — mito-GFP / Futsch
G' — mito-GFP
H merged A5 — mito-GFP / Futsch
H' — mito-GFP

I
I Gal4 control at A5

J Neuron NDUFS7 RNAi at A5

K
EPSP amplitude (mV)
p=0.240
Gal4 control   Neuron NDUFS7 RNAi

Muscle
L Control — mito-GFP
M NDUFS7 RNAi — mito-GFP

Synapse
N Control — mito-GFP / HRP
O NDUFS7 RNAi

P
Mitochondria clusters NMJ 6/7
***

Synapse
Q Control — mito-GFP / Dlg
R NDUFS7 RNAi

S
Mitochondria vol. (mito/Dlg vol. μm³)
***
Control   Neuron NDUFS7 RNAi

	Gal4-control	sod2 OE neuron	NDUFS7 RNAi neuron
A	A'	B / B'	C / C'

	NDUFS7 neuron+NACA	sod2 neuron rescue	
D	D'	E	E' merged

HRP
Futsch

F

A Gal4 control **B** *sod2* OE neuron **C** *NDUFS7* RNAi neuron

D *NDUFS7* RNAi neuron+NACA **E** *sod2* neuron rescue

BRP-fire LUT

F

Normalized BRP Intensity/ μm² area of bouton

***   ***

H Gal4 control **I** *NDUFS7* RNAi neuron

H′ merged **I′** merged

BRP
GluRIII

G

BRP puncta density (per μm² area of bouton)

***   ***

3.0
2.5
2.0
1.5
1.0
0.5

Gal4 control
sod2 OE neuron
NDUFS7 RNAi
NDUFS7+NACA
sod2 rescue

Neuron

J Gal4 control **K** *sod2* OE neuron **L** *NDUFS7* RNAi neuron **M** *sod2* neuron rescue

*

N

mEPSP amplitude (mV)

***   ***
ns

1.0
0.8
0.6
0.4
0.2

1 2 3 4

O

EPSP amplitude (mV)

ns ns
ns *

45
40
35
30

1 2 3 4

P

Quantal content

***   *
ns

80
60
40
20

1 2 3 4

1. Gal4 control
2. *sod2* OE neuron
3. *NDUFS7* RNAi neuron
4. *sod2* neuron resue

mito-GFP, HRP

Q Gal4 control **R** *sod2* OE

S *NDUFS7* RNAi **T** *sod2* rescue

U

mito clusters/bouton

ns   ***

Gal4 control
sod2 OE neuron
NDUFS7 RNAi neuron
sod2 neuron rescue

V Control normal ROS

nROS

RRP

Bruchpilot

CaV2
Ca2+
GluR

Evoked potentials normal

W *NDUFS7* loss elevated ROS

ns

nROS

RRP

Bruchpilot

CaV2
Ca2+

Evoked potentials normal

X *sod*2 rescue normalized ROS

*

nROS

RRP

Bruchpilot

Neuron

CaV2
Ca2+
GluR

Muscle

Evoked potentials small

A

nROS
RyR
Ca²⁺
Ca²⁺
IP3R
MCU
Ca²⁺
Cav2
GluRs
Evoked potentials normal

ER
ns

B

nROS
RyR
Ca²⁺
Ca²⁺
IP3R
MCU
Ca²⁺
Cav2
GluRs
Evoked potentials normal

ER
ns

C

nROS
RyR
Ca²⁺
Ca²⁺
IP3R
MCU
Ca²⁺
Cav2
GluRs
Evoked potentials normal

ER
ns

D

nROS
RyR
Ca²⁺
Ca²⁺
IP3R
MCU
Ca²⁺
Cav2
GluRs
Evoked potentials small

ER
***

E Neuron Gal4 control

F Control+Xesto+Dant neuron

G *NDUFS7* RNAi neuron

H ns *NDUFS7* RNAi+Xesto +Dant neuron

I *IP3-Sponge.m30*+Dant +DMSO neuron

J *mcu* RNAi+Xesto +Dant neuron

K ** *mcu*+*NDUFS7* RNAi+ Xesto+Dant neuron

L *** *IP3-Sponge.m30*+*mcu*+ *NDUFS7* RNAi+Dant neuron

M Neuron-Gal4 +DMSO, wash

N ns Neuron-Gal4 +BAPTA-AM, wash

O *NDUFS7* RNAi neuron +DMSO, wash

P *** *NDUFS7* RNAi neuron +BAPTA-AM, wash

Q mEPSP amplitude (mV)
ns
*
ns   ns

R mEPSP Frequency (Hz)
ns
ns   ns

S EPSP amplitude (mV)
ns
** ***   ns   **

T Quantal content
ns
ns   **

elaV(C155)/++
elaV(C155)×Xesto+Dant
NDUFS7 RNAi×Xesto+Dant
IP3-Sponge.m30-Dant+DMSO
mcu RNAi+Xesto+Dant
NDUFS7 RNAi+Xesto+Dant
mcu+NDUFS7 RNAi+Xesto+Dant
IP3-Sponge.m30+mcu+NDUFS7 RNAi+Dant
Neuron-Gal4+DMSO, wash
NDUFS7 RNAi neuron+DMSO, wash
Neuron-Gal4+BAPTA-AM, wash
NDUFS7 RNAi neuron+BAPTA-AM, wash

A Glucose → Glucose-6-Phosphate → Pyruvate → TCA Cycle → ETS

Hexokinase LDA or 2-Deoxy-D-glucose

2 ATP directly

Glycolysis

2 ATP directly

B Control+DMSO

C ns Control+LDA

D NDUFS7 RNAi+DMSO

E *** NDUFS7 RNAi+LDA

F Control+ 2-Deoxy-D-glucose

G ** NDUFS7 RNAi+ 2-Deoxy-D-glucose

H mEPSP amplitude (mV)

ns   ns   **

I mEPSP frequency (Hz)

ns   ns   ns

J EPSP amplitude (mV)

ns   ***   **

Control+DMSO
Control+LDA
NDUFS7 RNAi+DMSO
NDUFS7 RNAi+LDA
Control+2-Deoxy-D-glucose
NDUFS7 RNAi+2-Deoxy-D-glucose

K Control+DMSO
L NDUFS7 RNAi+DMSO
M Control+LDA
N NDUFS7 RNAi+LDA
O Control+2-Deoxy-D-glu
P NDUFS7+2-Deoxy-D-glu

BRP-Fire-Lut

255
0

Q BRP relative Intensity (A.U) μm²/ area of bouton

***   **   ***

Control+DMSO
NDUFS7 RNAi+DMSO
Control+LDA
NDUFS7 RNAi+LDA
Control+2-Deoxy-D-glucose
NDUFS7 RNAi+2-Deoxy-D-glucose

J (Total number of boutons, A2 hemisegment 6/7)

K (Muscle area in μm²)

L (# of branches at A2 muscle 6/7)

1. Muscle Gal4
2. Muscle *sod2* OE
3. Muscle *sod1* OE
4. Muscle *cat* OE
5. Muscle *NDUFS7* RNAi
6. Muscle *NDUFS7* RNAi+NACA
7. *sod2* muscle rescue
8. *sod1* muscle not rescue
9. *cat* muscle not rescue
10. Pan neuron Gal4
11. Pan neuron *NDUFS7* RNAi
12. Motor neuron Gal4
13. Motor neuron *NDUFS7* RNAi

U (Relative Dlg area (Dlg-HRP per bouton)/μm²)

V (Bouton area μm²)

a. Muscle Gal4
b. Muscle *sod2* OE
c. Muscle *sod1* OE
d. Muscle *cat* OE
e. Muscle *NDUFS7* RNAi
f. Muscle *NDUFS7* RNAi+NACA
g. Muscle *sod2* rescue
h. Muscle *sod1* not rescue
i. Muscle *cat* not rescue

A Muscle Gal4 | Muscle *NDUFS7* RNAi | *cat* muscle not rescue | *sod1* muscle not rescue | *sod2* muscle rescue

B Muscle-Gal4+10% EtOH | Muscle *NDUFS7* RNAi+10% EtOH | Muscle *NDUFS7* RNAi+0.5mM curcumin | Muscle *NDUFS7* RNAi+0.5mM NACA

C *ND-30* mutant | *ND-30* mutant+0.5mM NACA | *ND-30* mutant *sod2* muscle rescue

D Muscle-Gal4+0.5% DMSO | Muscle-Gal4 (50μM rotenone) | Muscle *cat* OE (50μM rotenone) | Muscle *sod1* OE (50μM rotenone) | Muscle *sod2* OE (50μM rotenone)

E mEPSP amplitude (mV)

F EPSP amplitude (mV)

G Quantal content

a. Muscle Gal4
b. Muscle *NDUFS7* RNAi
c. *cat* muscle rescue
d. *sod1* muscle not rescue
e. *sod2* muscle not rescue
f. Muscle Gal4+10% EtOH
g. Muscle *NDUFS7* RNAi+10% EtOH
h. Muscle *NDUFS7* RNAi+0.5mM curcumin
i. Muscle *NDUFS7* RNAi+0.5mM NACA
j. *ND-30* mutant
k. *ND-30* mutant+0.5mM NACA
l. *ND-30* mutant *sod2* muscle rescue
m. Muscle Gal4+0.5% DMSO
n. Muscle Gal4+50μM rotenone
o. Muscle *cat* OE+50μM rotenone
p. Muscle *sod1* OE+50μM rotenone
q. Muscle *sod2* OE+50μM rotenone

H Distance crawled in cm

1. Muscle Gal4
2. Muscle *sod2* OE
3. Muscle *ND-30* RNAi
4. Muscle *NDUFS7* RNAi
5. Muscle *sod2* rescue
6. Muscle *NDUFS7* RNAi+NACA
7. Pan neuron Gal4
8. Pan neuron *NDUFS7* RNAi
9. *ND-30* mutant
10. *ND-30* mutant *sod2* muscle rescue