CORNICHON HOMOLOG 5-dependent ER export of membrane cargoes in phosphate-starved *Arabidopsis* root as revealed by membrane proteomic analysis

Tzu-Yin Liu,[1,2,3,†] Ming-Hsuan Tsai,[1,†] Jhih-Yi Wang,[1,†] Hui-Fang Lung,[1] Chang-Yi Chiu,[1] Hong-Xuan Chow,[1] Chiao-An Lu[3]

[1]Institute of Bioinformatics and Structural Biology, College of Life Sciences and Medicine, National Tsing Hua University, Hsinchu 30013, Taiwan.

[2]Department of Life Science, College of Life Sciences and Medicine, National Tsing Hua University, Hsinchu 30013, Taiwan.

[3]Interdisciplinary Program of Life Science and Medicine, College of Life Sciences and Medicine, National Tsing Hua University, Hsinchu 30013, Taiwan.

Author for correspondence: Tzu-Yin Liu

Tel: +886 3 574-2096; Fax: +886 3 571-5934

Email: tzliu@life.nthu.edu.tw

Institute of Bioinformatics and Structural Biology, College of Life Sciences and Medicine, National Tsing Hua University, Hsinchu 30013, Taiwan

[†]These authors have contributed equally to this work.

Running title: Membrane proteomics of Pi-starved *cnih5* root

Synopsis: *Arabidopsis thaliana* CORNICHON HOMOLOG 5 (*At*CNIH5) acts as a low phosphate (Pi)-responsive hub that controls efficient ER export of specific membrane cargoes beyond PHT1 Pi transporters to improve plant fitness under suboptimal Pi availability. Unlike the fungal and rice CNIHs interacting with their cognate cargoes through the C-terminal acidic motif, *At*CNIH5 employs a distinct cargo selection mechanism for *At*PHT1;1.

Key words: CORNICHON HOMOLOG, ER cargo receptor, coat protein complex II

32   (COPII), phosphate starvation, phosphate transporter, 4-hexylphenylazosulfonate (Azo)

Summary

- To find potential membrane cargoes of *Arabidopsis thaliana* CORNICHON HOMOLOG 5 (*At*CNIH5), a phosphate (Pi) deficiency-induced ER cargo receptor, we applied the UV-cleavable 4-hexylphenylazosulfonate (Azo)-solubilized microsomal protein extraction for iTRAQ-based proteomic analysis and identified 106 downregulated proteins in the Pi-limited roots of *cnih5*, including PHOSPHATE TRANSPORTER 1 proteins (PHT1s) and transmembrane enzymes responsible for the biosynthesis of very long-chain fatty acids and their derivative extracellular aliphatic compounds, and nucleotide sugar for cell walls.

- We confirmed the interaction of *At*CNIH5 with several downregulated transporters in *cnih5*, including *At*PHT1s, *At*OCT1, *At*URGT6, *At*DTX21, and *At*DTX35, using the yeast split-ubiquitin and the *in-planta* tripartite split-GFP assays.

- The C-terminal acidic residue of *At*CNIH5 is not required to interact with *At*PHT1;1, implying that *At*CNIH5 employs a distinct cargo selection mechanism for *At*PHT1;1.

- We showed the upregulation of *At*CNIH5 proteins in the Pi-limited root and in the Pi-replete root of the Pi overaccumulator *pho2*. As increasing *in-situ* *At*CNIH5 expression/activity enhances plant growth, we propose that *At*CNIH5 functions as a low Pi-responsive hub, facilitating the ER export of specific membrane cargoes and thereby improving plant fitness under suboptimal Pi availability and promoting agricultural sustainability.

Introduction

Phosphorus (P) is an essential macronutrient for plant growth and development
(Poirier & Bucher, 2002; Poirier *et al.*, 2022). Due to the low concentration of inorganic
phosphate (Pi) in soil—the primary form of P that plants absorb, applying Pi fertilizer has
become a common practice to improve crop productivity. However, the limited mobility
of Pi in soil prevents plants from acquiring most applied Pi fertilizer, leading to P runoff
into aquatic ecosystems. Therefore, developing plants with increased Pi uptake and high-
use efficiency is critical to minimizing the usage of Pi fertilizer, thus reducing
environmental pollution and achieving agricultural sustainability.

To adapt to the low Pi bioavailability, plants have many strategies to enhance external
Pi acquisition and internal Pi remobilization, which involve different Pi transporters at
distinct subcellular compartments. *PHOSPHATE TRANSPORTER 1* (*PHT1*) genes
encode the plasma membrane (PM)-localized Pi transporters responsible for the uptake
and the intercellular translocation of Pi. Even though Pi deprivation upregulates most
PHT1 genes at the transcript level, the biosynthesis, secretion, and recycling rate of PHT1
proteins (PHT1s) also determine their abundance at the PM (Nussaume *et al.*, 2011).
Multiple post-translational mechanisms have been elucidated to regulate the protein
expression and the PM localization of PHT1s. The plant-specific SEC12-related
PHOSPHATE TRANSPORTER TRAFFIC FACILITATOR1 (PHF1) contributes to the
accumulation of PHT1s at the PM by assisting the PHT1s transit through the endoplasmic
reticulum (ER) (González *et al.*, 2005; Bayle *et al.*, 2011; Chen *et al.*, 2015). The rice
casein kinase II (CK2) diminishes the interaction between *Os*PT8 and *Os*PHF1 by
phosphorylating *Os*PT8, resulting in the ER retention of *Os*PTs (Chen *et al.*, 2015). On
the contrary, the protein phosphatase type 2C *Os*PP95 acts antagonistically with *Os*CK2
(Yang *et al.*, 2020). In addition, under Pi sufficiency, the ubiquitin-conjugating E2
enzyme UBC24/PHOSPHATE2 (PHO2) at the ER or post-ER compartments mediates
the ubiquitination of the endomembrane-associated *At*PHT1s followed by the vacuolar
degradation (Huang *et al.*, 2013). The PM-localized ubiquitin E3 ligase NITROGEN
LIMITATION ADAPTATION (NLA) participates in the ubiquitination-mediated and
clathrin-dependent endocytosis of PHT1s when Pi is replete (Lin *et al.*, 2013; Park *et al.*,
2014).

The ER is the start site where newly synthesized PHT1s travel towards the PM. The

ER-to-Golgi transport of PHT1s relies on the coat protein complex II (COPII)-mediated

vesicles or carriers (Bayle *et al.*, 2011). However, how diverse membrane cargoes

associated with Pi starvation are transported in the late secretory pathway and whether

post-translational modification events regulate packaging of these cargoes into COPII

vesicles remains largely unknown. The formation of COPII vesicles is a multi-step

process that involves the recruitment of five core cytosolic factors, SAR1, SEC23, SEC24,

SEC13, and SEC31, to the ER exit sites (ERES) (Barlowe and Miller, 2013; Kurokawa

and Nakano, 2018). In particular, incorporation of multi-spanning transmembrane (TM)

proteins into COPII vesicles in the budding yeast *Saccharomyces cerevisiae* requires dual

interaction with the COPII coat subunit Sec24 and the ER cargo receptor Erv14 (Powers

and Barlowe, 1998; Harmel et al., 2012; Pagant et al., 2015). The function of Erv14 in

COPII coupling was presumably mediated by its role as a membrane chaperone that

exposes sorting signals of Erv14 clients to interact with Sec24 for efficient capture into

COPII vesicles (Dancourt and Barlowe, 2010; Pagant et al., 2015). Loss of the fungal

Erv14 compromised the targeting of several transporters destined for the late secretory

compartments, thus misregulating numerous cellular functions (Powers and Barlowe,

1998; Nakanishi et al., 2007; Pagant et al., 2015; Zimmermannová et al., 2019; Zheng et

al., 2023). The *Drosophila* Cornichon (Cni), like its yeast homolog Erv14, was shown to

recruit the epidermal growth factor (EGF)-like ligand Gurken into COPII vesicles (Roth

et al., 1995; Bökel *et al.*, 2006). Without Cni, the ER export and secretion of Gurken are

disrupted (Bökel *et al.*, 2006). Similarly, the plant CNIHs act as the ER cargo receptors

for various ion and metabolite transporters (Rosas-Santiago *et al.*, 2015; Wudick *et al.*,

2018; Wei *et al.*, 2023; Yáñez-Domínguez *et al.*, 2023).

Our recent side-by-side study uncovered that *Arabidopsis thaliana CORNICHON*

HOMOLOG 5 (*AtCNIH5*) is a Pi starvation-induced (PSI) gene and preferentially

expressed in the outer layers of the root above the apical meristem (Chiu *et al.*, 2024).

*At*CNIH5   localizes   to   the   ER   and   is   closely   associated   with   the

*At*SAR1A/*At*SEC16A/*At*SEC24A-labeled ERES (Chiu *et al.*, 2024). The *cnih5* mutant

exhibited a lower abundance of *At*PHT1s in the root microsomal fractions, with reduced

shoot Pi levels and shoot-to-root distribution of Pi under Pi sufficiency and decreased Pi

uptake under Pi deficiency (Chiu *et al.*, 2024). In the Pi-limited root of *cnih5*, the

fluorescent protein tagged-*At*PHT1;1 was in part retained intracellularly in the epidermis

within the transition/elongation zones and in the root hair (Chiu *et al.*, 2024). Consistently, the fluorescence intensities of *At*PHT1;1 fusion proteins at the PM of the root cells mentioned above also diminished in the Pi-limited *cnih5* relative to the wild-type (WT) control (Chiu *et al.*, 2024), supporting a potential role for *At*CNIH5 in the efficient transport of *At*PHT1s from the ER to the Golgi. Additional phenotypes, such as shorter root hair length and increased root hair and lateral root densities, were also observed for *cnih5* under Pi deficiency (Chiu *et al.*, 2024). Considering that *At*CNIH5 may serve as a cargo receptor for a broader range of membrane proteins and that inefficient ER export of cargoes in *cnih5* can cause their intracellular retention and eventual degradation due to the protein quality control, we inferred that the pleiotropy of *cnih5* is likely consequences of mistargeting and/or downregulation of membrane cargoes of *At*CNIH5 beyond *At*PHT1s.

To identify potential cargoes of *At*CNIH5 associated with Pi starvation, we employed isobaric tags for relative and absolute quantification (iTRAQ)-based membrane proteomics. Because membrane proteins are highly hydrophobic and have low abundance, isolation of microsomal proteins (MPs) enriched for PM, ER, Golgi, vacuolar membranes, and endosomal vesicles is challenging (Wang & Liang, 2012). Conventionally, MPs are solubilized with sodium dodecyl sulfate (SDS) or other anionic detergents. However, SDS impacts the liquid chromatography-tandem mass spectrometry (LC–MS/MS) performance, and the removal step of SDS before mass spectrometry analysis leads to an inevitable loss of MPs (Wang & Liang, 2012). A study using the rice leaf as a model system demonstrated that the microcentrifugation-based microsomal protein extraction (MME) method is effective for the downstream label-free quantitative proteome analysis, requiring neither a large amount of plant material nor a step of ultracentrifugation to enrich the PM proteins (Nguyen *et al.*, 2021). The authors also applied the UV-cleavable and mass spectrometry-compatible surfactant, 4-hexylphenylazosulfonate (Azo) (Brown *et al.*, 2019; Brown *et al.*, 2020), to avoid protein losses by omitting the SDS detergent wash step. Their results revealed several advantages of Azo over SDS in solubilizing the MPs, including a higher number and abundance of MPs (Nguyen *et al.*, 2021). In this study, we modified the Azo-solubilized MME method and isolated the root MPs from Pi-limited *Arabidopsis* WT and *cnih5* for iTRAQ-based proteomic analysis. We targeted our research to root MPs to reduce sample complexity,

thus increasing the possibility of identifying functionally relevant protein changes in
cnih5. In addition, the membrane proteome analysis of Pi-limited roots is valuable in
gaining insight into the physiological role of *At*CNIH5. It also provides the first database
of the membrane proteome for Pi-limited root biology research (Zhou *et al.*, 2022).

With this approach, we identified 4,317 proteins in the Pi-limited WT and *cnih5* roots,
in which 1231 proteins (28.5%) have at least one TM helix. Of the differentially expressed
proteins (DEPs) in *cnih5*, 372 and 106 proteins are upregulated and downregulated by a
1.25-fold change, respectively, including the downregulation of *At*PHT1;1–1;5,
corroborating our recent finding that *At*PHT1s are genuine cargoes of *At*CNIH5 (Chiu *et
al.*, 2024). Consistent with the primary role of *At*CNIH5 in the ER export of membrane
cargoes, the enrichment analyses of Gene Ontology (GO) and Kyoto Encyclopedia of
Genes and Genomes (KEGG) pathways suggested a close association of the
downregulated proteins with protein export and localization. We also found that the
transmembrane enzymes involved in the biosynthesis of nucleotide sugar for cell walls,
very long-chain fatty acids (VLCFAs), and their derivative extracellular aliphatic
compounds, such as cutin, suberin, and wax, were downregulated in the Pi-limited roots
of *cnih5*.

We used the yeast split-ubiquitin system (SUS) and the *in-planta* tripartite split-GFP
assay to confirm the interaction of the potential membrane cargoes with *At*CNIH5,
including *At*PHT1s and several transporters. Our immunoblot results demonstrated the
induction of *At*CNIH5 in the Pi-limited root and the upregulation of *At*CNIH5 in the Pi-
replete root of the Pi overaccumulator *pho2*, likely as a result of accommodating the
increased demand for ER export of *At*PHT1s. We also generated a series of C-terminally
truncated forms of *At*CNIH5 to find out which motif of *At*CNIH5 interacts with
*At*PHT1;1. We showed that the C-terminal acidic residue, aspartic acid, is not required to
interact with *At*PHT1;1. Instead, *At*CNIH5 1–26, containing the first transmembrane
domains (TMD) alone, is sufficient to interact with *At*PHT1;1. Therefore, unlike the
fungal and rice CNIHs interacting with their cognate cargoes through the C-terminal
acidic motif (Rosas-Santiago *et al.*, 2017), *At*CNIH5 employs a distinct mechanism to
select *At*PHT1;1 as its cargo. Importantly, introducing the *At*CNIH5 genomic-GFP
translation fusion to complement the *cnih5* mutant boosted plant growth under suboptimal
Pi supply, revealing that *At*CNIH5 acts as a hub to facilitate the ER export of specific

membrane cargoes associated with Pi starvation to improve plant fitness.

Materials and Methods

Plant Material and Growth Conditions

Seeds of the *A. thaliana* wild-type (Columbia, Col-0) and *cnih5* (SAIL_256_B03)
mutants used in this study were obtained from the Arabidopsis Biological Resource
Center. Seeds were surface-sterilized and germinated on agar plates with one-half
modified Hoagland's solution containing 1% sucrose and 0.7% Bacto agar and grown in
the growth chamber at 22 °C with a 16-h light/8-h dark cycle. Pi-sufficient (+P; 250 μM
Pi) or Pi-deficient (−P; 0 μM Pi) medium containing 1.2% agar was used for vertically
grown plants.

Tissue Homogenization and Microsomal Protein Preparation

The roots of 11-day-old seedlings with five days of Pi starvation were harvested.
Approximately 100 mg of root tissue was collected for each sample and stored at −80 °C
for subsequent homogenization. The frozen samples were homogenized by TissueLyser
II (QIAGEN). The method for microsomal protein extraction was modified from a
previous study (Abas & Luschnig, 2010), and all steps were performed on ice or 4 °C.
Briefly, each sample was resuspended in 100 μl of extraction buffer (EB) containing 50
mM of hydroxyethyl piperazine ethane sulfonic acid (HEPES; Melford, pH 7.5), 25%
(w/w 0.81 M) sucrose, 5% (v/v) glycerol, 10 mM of ethylenediaminetetraacetic acid
(EDTA; Amresco, pH 8), 5 mM of KCl, 1 mM of dithiothreitol (DTT; Sigma), and 2×
protease inhibitor cocktail (P9599, Sigma). An equivalent volume (100 μl) of 5%
polyvinylpolypyrrolidone (PVPP; Sigma) suspension solution was used for each sample.
The 5% PVPP was centrifuged at 2,000×g for 1 min at room temperature (RT). The
precipitate was activated for 2 hours in an equivalent volume of activated solution
containing 100 mM HEPES, 40% (w/w, 1.37 M) sucrose, 20 mM EDTA, and 10 mM
KCl. The activated PVPP was centrifuged at 5,000×g for 1 min at RT, and the supernatant
was removed. The activated PVPP pellet was mixed with the sample suspension and stood
for 5 min, followed by centrifugation thrice. The first centrifugation was processed at
600×g for 3 min. The pellet was then re-extracted with 50 μl EB and centrifuged at 600×g
for 3 min. The pellet was then re-extracted with 35 μl EB and centrifuged at 2,000×g for

30 s. After three times centrifugation, all the supernatants were combined and centrifuged

again at 600×g for 3 min. The collected supernatant was diluted with an equivalent

volume of ddH$_2$O to reach a final solution concentration containing 12–13% sucrose and

centrifuged at 21,000×g for 90 min. The pellet was resuspended in washing buffer

containing 10 mM HEPES, 5 mM EDTA, 1 mM phenylmethylsulfonyl fluoride (PMSF,

Sigma), and 1× protease inhibitor cocktail and centrifuged at 21,000×g for 45 min. Finally,

the pellet was stored at −80 °C for further analysis.

Protein Solubilization and In-Solution Digestion

Microsomal protein pellets were dissolved in 1% sodium 4-hexylphenylazosulfonate (Na-

Azo, Excenen Pharmatech) in 100 mM triethylammonium bicarbonate (TEABC, Sigma)

containing 2× protease inhibitor cocktail and 1 mM PMSF. The protein yield was

determined by BCA assay (Pierce). For each sample, an equal amount of protein was

reduced with 5 mM tris-(2-carboxyethyl)-phosphine (TCEP, Sigma) at 60 °C for 30 min,

followed by cysteine-blocking with 10 mM methyl methanethiosulfonate (MMTS, Sigma)

at 25 °C for 30 min. Samples were digested with sequencing-grade modified porcine

trypsin (Promega) at 37 °C with shaking for 1 hour and then illuminated by 6W UV light

302 nm (UVP UVM-57, Analytik Jena) at 4 °C for 40 min. The digested proteins were

then aliquot and stored at −20 °C until use.

Isobaric Tag for Relative and Absolute Quantitation (iTRAQ) Labeling and Peptide

Separation

After trypsin digestion, 0.5 μg of peptides was subjected to 1D–LC–MS/MS without pre-

fractionations to identify and quantify proteins as a quality control study. On the other

hand, 10 μg of peptides was labeled using an iTRAQ Reagent-4 plex Multiplex Kit

according to the manufacturer's instructions (Applied Biosystems) for 1 hour at RT. Two

WT replicates were labeled with iTRAQ-114 and -115, and two *cnih5* replicates were

labeled with iTRAQ-116 and -117, pooled and desalted by homemade C18-microcolumn

(SOURCE 15RPC, GE Healthcare). The eluted peptides were dried by vacuum

centrifugation. The desalted peptides were loaded onto a homemade strong cation

exchange (SCX) column (Luna SCX 5μm, 0.5 × 120 mm) at a 2 μl/min flow rate for 60

min. The peptides were then fractionated to 24 fractions by eluting with 0 to 100% HPLC

buffer B (1 M ammonium nitrate/25% acetonitrile/0.1% formic acid) using on-line two-
dimensional high-performance liquid chromatography (2D-HPLC; Dionex Ultimate
3,000, Thermo Fisher). Each SCX fraction was diluted in-line before trapping onto a
reverse-phase (RP) column (Zorbax 300SB-C18, 0.3 x 5 mm; Agilent Technologies). The
peptides were then separated on a column (Waters BEH 1.7 μm, 100 μm I.D. × 10 cm
with a 15 μm tip) using a multi-step gradient of HPLC buffer C (99.9% acetonitrile/0.1%
formic acid) for 60 min with a flow rate of 0.3 μl/min.
256
Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS) Analysis
The LC apparatus was coupled with a 2D linear ion trap mass spectrometer (LTQ-
Orbitrap ELITE; Thermo Fisher) operated using Xcalibur 2.2 software (Thermo Fisher).
The full-scan MS was performed in the Orbitrap over a range of 400 to 2,000 Da and a
resolution of 120,000 at m/z 400. For proteome analysis, the 10 data-dependent MS/MS
scan events (10 MS2 HCD) were followed by one MS scan for the 10 most abundant
precursor ions in the preview MS scan. The m/z values selected for MS/MS were
dynamically excluded for 80 sec with a relative mass window of 1 Da. The electrospray
voltage was set to 1.8 kV, and the capillary temperature was set to 200 °C. MS and MS2
automatic gain control were set at 1,000 ms (full scan) and 300 ms (MS2 for HCD), or 3
$\times 10^6$ ions (full scan) and 3×10^4 ions (MS2 for HCD) for maximum accumulated time
or ions.
269
LC-MS Data Processing, **Protein Identification and Quantification**
Proteome Discoverer software (version 2.3, Thermo Fisher Scientific) was used for data
analysis, incorporating the reporter ions quantifier node for iTRAQ quantification. The
MS/MS spectra were searched against the SwissProt database using the Mascot search
engine (Matrix Science, version 2.5). For peptide identification, a 10 ppm mass tolerance
was allowed for intact peptide masses and 0.2 Da for HCD fragment ions, with allowance
for two missed cleavages made from trypsin digestion. Variable modifications included
oxidized methionine, acetylation (protein N-terminal), and iTRAQ-4plex (peptide N-
terminal, K); methylthiolation (C) was considered a static modification. Peptide-spectrum
matches (PSMs) were filtered based on high confidence and Mascot search engine rank
1 of peptide identification, ensuring an overall false discovery rate (FDR) below 0.01.

Proteins with only a single peptide hit were removed. Quantitative protein data were
exported from Proteome Discoverer using Integration Methods of the Most Confident
Centroid with a 20 ppm tolerance. The data sets were analyzed by cross-comparison
between four samples (ratio 1, 116:114; ratio 2, 117:114; ratio 3, 116:115, and ratio 4,
117:115). Protein ratios in each comparison group were normalized using the median
ratio of each comparison group. Volcano plots of differentially expressed proteins were
generated using the ggplot2 (Wickham *et al.*, 2016) and ggrepel (Slowikowski *et al.*, 2024)
package in R version 4.3.1 (R Core Team, 2021). The code was written using RStudio
version 2023.06.2 (RStudio Team, 2020). A 1.25-fold cutoff was applied to define
differentially expressed proteins in each comparison group.

Protein Annotation and Interpretation
Transmembrane-related information was obtained through analysis using DeepTMHMM
(https://dtu.biolib.com/DeepTMHMM) (Hallgren *et al.*, 2022). Subcellular structure
localization was analyzed by SUBA5 (https://suba.live/) (Hooper *et al.*, 2016). For
functional annotation of proteins, Gene Ontology (GO) and Kyoto Encyclopedia of Genes
and Genomes (KEGG) enrichment analyses were determined by ShinyGO 0.80
(http://bioinformatics.sdstate.edu/go/) (Ge *et al.*, 2019), in which *p*-values were computed
using the hypergeometric test and multiple test correction was performed using the
Benjamini–Hochberg method based on an FDR cut-off of 0.05. Only pathways with a
minimum of 10 genes were selected. Similar pathways that share 95% of their genes and
50% of the words in their names were eliminated according to the setting of ShinyGO
0.80. Dot plots of GO enrichment analysis were generated using the ggplot2 (Wickham
et al., 2016) and stringr package (Wickham, 2023).

Microsomal protein extraction for immunoblot analysis
The MPs pellets were solubilized by 0.5% Azo in 25mM aqueous ammonium
bicarbonate, pH 7.0 (919233; Sigma Aldrich). Alternatively, the low-speed pellet (LSP)
was prepared as previously described (Yoshimoto *et al.*, 2004), and microsomal protein
was isolated using the Minute Plant Microsomal Membrane Extraction Kit (Invent, MM-
018) according to the manual instructions. A 15–30 µg of MPs from each sample was
loaded onto 4–12% Q-PAGE™ Bis-Tris Precast Gel (SMOBIO) or NuPAGE 4–12% Bis-

Tris Gels (Thermo Fisher Scientific). Gel was transferred to polyvinylidene difluoride

membranes (IPVH00010 Immobilon-P Membrane or ISEQ00010 Immobilon®-PSQ

Membrane; Merck). Membrane was blocked with 1–2% BSA in 1× PBS solution with

0.2% Tween 20 (PBST, pH 7.2) at RT for 1 h, and rinsed two times with distilled water

followed by hybridization with primary antibodies of *At*PHF1 (Huang *et al.*, 2013),

*At*PHO1 (Liu *et al.*, 2012), *At*NRT2.1 (AS12 2612; Agrisera), and *At*CNIH5 at RT for 2

h or at 4 °C overnight in blocking solution. Polyclonal rabbit *At*CNIH5 antibody was

raised against the region corresponding to the amino acid residues 96 to 114

(TELYNTNKWEQKKRVYKIG) by LTK BioLaboratories, Taiwan, and was affinity

purified. The membrane was washed four times with 1× PBST for 5 min followed by

hybridization with the horseradish peroxidase–conjugated secondary antibody

(GTX213110-01 or GTX213111-01; GeneTex) in blocking solution at RT for 1 h. After

four washes in 1× PBST for 5 min and a rinse with distilled water, chemiluminescent

substrates (WesternBright ECL or Sirius; Advansta) were applied for signal detection.

Construct design and yeast split-ubiquitin assay

For constructing the yeast expression clones, the coding sequence (CDS) of the gene of

interest (GOI), including *At*LRK10L-1.1, *At*GXM2, *At*SOS, *At*DTX21, *At*DTX35,

*At*URGT6, *At*OCT1, *At*SEC24A, *At*SEC24B, *At*SEC24C and the truncated *At*CNIH5

variants, was amplified by PCR, cloned into the pCR8/GW/TOPO vectors and sequenced.

The pCR8/GW/TOPO construct containing the CDS of *AtPHT1;1* was cloned as

previously reported (Huang *et al.*, 2013). The entry clones were recombined into

NX32_GW (ABRC CD3-1737) or XN22_GW (ABRC CD3-1735) through LR reactions,

resulting in N or C-terminally fused Nub fragments or into MetYC_GW (ABRC CD3-

1740) through LR reactions, resulting in *At*PHT1;1-PLVCub. The PLVCub-*At*CNIH5

construct was generated as recently reported (Chiu *et al.*, 2024). The Nub and PLVCub

expression clones were sequentially transformed into the yeast THY.AP4 strain (ABRC

CD3-808; (Obrdlik *et al.*, 2004). Fresh yeast colonies picked from yeast extract peptone

dextrose adenine (YPDA) medium or synthetic complete (SC) medium without leucine

(SC–L) were inoculated in the corresponding liquid medium and incubated overnight at

30 °C with shaking at 200 rpm. Overnight yeast culture was then diluted to $OD_{600} = 0.2$

and grew for another 4–6 h until OD_{600} reached 0.4–0.6. Yeast pellet was collected by

centrifugation at 2,000×g at 4 °C for 10 min, washed with ddH$_2$O, and resuspended in

Tris-EDTA/lithium acetate solution (TE/LiAc, 10 mM Tris-HCl [pH 8.0], 1 mM EDTA,

and 0.1 M lithium acetate). Yeast transformants carrying the PLVCub-*At*CNIH5 or

*At*PHT1;1-PLVCub were selected and maintained on SC–L for sequential transformation

with NubI (positive control), NubG (negative control), and GOI-NubG or NubG-GOI.

Yeast transformants carrying the Nub and the PLVCub constructs were selected and

maintained on SC medium without leucine and tryptophan (SC–LW). For the protein–

protein interaction testing, yeast cells co-expressing PLVCub-*At*CNIH5 or *At*PHT1;1-

CubPLV with GOI-NubG or NubG-GOI were grown on synthetic media lacking leucine,

tryptophan, adenine, and histidine (SC–LWAH) with 500 μM methionine or on SC–LW

containing 100 mg/L 5-bromo-4-chloro-3-indolyl-*β*-d-galactopyranoside (X-Gal) at 30

°C for 24–36 h. Primer sequences used are listed in Supporting Information Table S4.

Construct design for split-GFP assays in *N. benthamiana* leaves and *Arabidopsis*

transgenic plants

The *UBQ10:sXVE:* S10-*At*CNIH5 clone was obtained as recently reported (Chiu *et al.*,

2024). For the transient expression of *At*PHT1;1-S11, *At*SOS-S11, *At*DTX21-S11,

*At*DTX35-S11, *At*URGT6-S11, *At*OCT1-S11, *At*HAK5-S11, *At*VPE1-S11 and

*At*SULTR1;3-S11, the CDS or the corresponding fragment was amplified by PCR and

subcloned into *UBQ10:sXVE*: (MCS)-S11 (Addgene plasmid # 108179). For subcellular

localization analysis of the protein of interest, either *UBQ10:sXVE:* S11-GFP1–9

(Addgene plasmid # 125665) or *UBQ10:sXVE:* GFP1–10 (Addgene plasmid # 125663)

was co-expressed. For the tripartite split-GFP reconstitution, the constructs encoding

S10-tagged and S11-tagged protein fusions were co-expressed in the presence of

UBQ10:sXVE: GFP1–9 (Addgene plasmid # 108187). Co-expression with 3xHA-S11

(Addgene plasmid # 108238) was used as a negative control for protein–protein

interaction. The brightness of the images for the protein–protein interaction was adjusted

by a 20% increase and the contrast adjusted by a 20% decrease. Primer sequences used

are listed in Supporting Information Table S1. For the genomic GFP-*AtCNIH5*

complemented plants (*AtCNIH5pro*: GFP-*AtCNIH5/cnih5*), the construct design was

recently described (Chiu *et al.*, 2024).

***Agrobacterium tumefaciens*-mediated infiltration of *N. benthamiana* leaves**

Agrobacterium tumefaciens strain EHA105 harboring the expression constructs or p19

was cultured in Luria–Bertani (LB) medium with the appropriate antibiotics [gentamicin

(50 µg ml^{-1}), rifampicin (5 µg ml^{-1}) or kanamycin (50 µg ml^{-1})]. Cells were resuspended

in the infiltration medium containing 10 mM MgCl$_2$, 10 mM MES (pH 7.5), and 150 µM

acetosyringone, and co-infiltrated to the true leaves of 3–5-week-old tobacco plants. For

the β-estradiol-inducible split-GFP system, 36.7 µM β-estradiol was applied as previously

described (Liu *et al.*, 2018; Liu, 2021). Infiltrated leaves were collected 2 days post-

infiltration (dpi) for confocal imaging.

Confocal microscopy

Confocal microscopy images were acquired using Zeiss LSM 800 (Zeiss) with objectives

Plan-Apochromat 10×/0.45 M27, 20×/0.8 M27, and 40×/1.3 Oil DIC M27 in multi-track

mode with line switching and averaging of two reading. Excitation/emission wavelengths

were 488 nm/410–546 nm for GFP and 561 nm/656–700 nm for RFP and mCherry.

Chemical treatments

Acetosyringone (150 mM stock in DMSO; D134406; Sigma-Aldrich) was diluted to a

final concentration of 150 µM in ddH$_2$O. β-estradiol (36.7 mM stock in ethanol; E2758;

Sigma-Aldrich) was diluted to a final concentration of 36.7 µM in ddH$_2$O. For monitoring

the degradation of 1% Azo following the UV irradiation, 1% sodium 4-

hexylphenylazosulfonate (Na-Azo, Excenen Pharmatech) in 100 mM TEABC buffer was

1: 200 diluted and subjected to absorbance measurement.

Results

Azo-solubilized microcentrifugation-based microsomal protein extraction (MME)

of *Arabidopsis* roots

Because the application of Azo promoted fast and reproducible tryptic digestion for

quantitative proteomics (Brown *et al.*, 2020), we replaced the mass spectrometry-

incompatible surfactant SDS with Azo and established an efficient workflow of MPs

extraction for quantitative proteomic analysis of *Arabidopsis* roots (Fig. **1a**). To ensure

that the Azo wash step can be skipped prior to mass spectrometry analysis, we tested the

time-course effect of UV irradiation on Azo breakdown (Brown *et al.*, 2019). Within 40

min of UV illumination, 1% Azo in 100 mM triethylammonium bicarbonate (TEABC)

almost wholly degraded (Fig. **1b**). After tryptic digestion and Azo degradation, small

aliquots of four samples—two biological replicates each for WT and *cnih5*, were pre-

analyzed in a single one-dimensional (1D)–LC–MS/MS run. Summary of the label-free

proteomic analysis was shown in Supporting Information Fig. S1a. We used multiple

marker abundance profiling (MMAP) (Hooper *et al.*, 2016) to estimate the relative

organelle abundance in the MPs samples. The results suggested that the modified Azo-

solubilized MME method greatly removed proteins targeted to the plastid, nucleus, and

extracellular space while enriching proteins targeted to the mitochondria, ER, Golgi, PM,

and vacuole (Supporting Information Fig. S1b, Fig. S1c). To further evaluate this

modified method, we examined by immunoblot the expression of the well-established

root MPs, including the ER-resident *At*PHF1, the Golgi-localized *At*PHO1 (Arpat *et al.*,

2012; Liu *et al.*, 2012), and the PM-targeted *At*PHT1;1/2/3 and *At*NRT2.1 (Chopin *et al.*,

2007). As previously reported, *At*PHF1, *At*PHO1, and *At*PHT1;1/2/3 were upregulated

following Pi deprivation (Fig. **2a**). Interestingly, regardless of Pi availability, the protein

amounts of *At*PHF1 and *At*PHO1 were more abundant in *cnih5* than in WT, while

*At*PHT1;1/2/3 was downregulated in *cnih5* compared to the WT controls (Fig. **2a**).

*At*NRT2.1 expression declined by Pi starvation and showed no differences between WT

and *cnih5* (Fig. **2a**). Collectively, these results justified the suitability of our modified

Azo-solubilized MME method for enriching root MPs.

Upregulation of *At*CNIH5 in the root of the Pi overaccumulator *pho2*

As we attempted to determine the expression of *At*CNIH5 in the root MPs, a specific

polyclonal antibody against *At*CNIH5 was raised. This antibody recognized a band in the

WT at ~15 kDa that was absent in *cnih5* (Fig. **2b**). In agreement with the finding that low

Pi induces the gene expression of *AtCNIH5*, the protein amount of *At*CNIH5 was also

elevated in the Pi-limited root (Fig. **2b**), suggesting that *AtCNIH5* is upregulated by Pi

starvation at the transcript and protein level. Of note, under Pi sufficiency, *At*CNIH5 was

modestly increased in the root of the Pi overaccumulator *pho2* relative to the WT control

(Fig. **2b**), which was in line with the previous membrane proteomics data showing that

*At*CNIH5 significantly increased in the Pi-replete root of the overaccumulator *pho2*

relative to the WT (Huang *et al.*, 2013). The increase of *At*CNIH5 in the root of *pho2*

under Pi-sufficient conditions was also recapitulated by two other different MPs

extraction methods (Fig. **2b**). Taken together, these results suggested that *At*CNIH5 is

induced by Pi starvation to meet the increased demand for ER export of *At*PHT1s, which

is substantiated by our genetic evidence that loss of *AtCNIH5* alleviates the Pi toxicity of

pho2 (Chiu *et al.*, 2024).

Quantitative membrane proteomic analysis of MPs in *Arabidopsis cnih5* roots under

Pi limitation

Next, the peptides obtained after tryptic digestion were labeled with iTRAQ reagents,

combined, and separated by two-dimensional high-performance liquid chromatography

(2D–HPLC) into 24 fractions. Each fraction was diluted and trapped onto a reverse-phase

column for separation, followed by Orbitrap mass spectrometry. By combining the total

proteins of the four samples, we identified 32,812 peptides and 21,433 unique peptides

that matched with 4,317 proteins with confidence (criteria described in Material and

Method), resulting in an average sequence coverage of 17.8% (Supporting Information

Table S2). About 99.3% of the identified proteins were shared across the four samples

(Supporting Information Table S2), indicating that our proteomic data were unbiased, as

nearly all proteins within the detection range could be identified. The proteins containing

TM helices were predicted or annotated by DeepTMHMM and UniProtKB (Fig. **3a**)

(Krogh *et al.*, 2001; Hallgren *et al.*, 2022; The UniProt Consortium, 2023). 1231

identified proteins (28.5%) have at least one TM helix and are thus integral membrane or

TM proteins (Fig. **3a**). We classified the subcellular localizations of the 4,317 identified

proteins using SUBA5 (Hooper *et al.*, 2016), showing 7.8% of the identified proteins

localized to the ER, 10.1% to the Golgi, 4.4% to the vacuole, and 16.2% to the PM (Fig.

3b). About 37.3% (1609) and 21.2% (915) of the identified proteins represent MPs and

integral MPs, respectively (Fig. **3a**). In contrast, 19.3% of the 1231 TM proteins localized

to the ER, 15.7% to the Golgi, 9.7% to the vacuole, and 31.7% to the PM (Fig. **3c**). The

general and detailed descriptions of proteomics data, including the summary of proteomic

analysis, protein scores and mass, annotation, and quantification for the identified

proteins, are listed in Supporting Information Table S2. The 4,317 identified proteins

were subjected to GO enrichment analysis by Shiny GO 0.80 (Ge *et al.*, 2019). Only the

top significant GO terms (FDR adj. p-value < 0.05) were shown according to the ranking

of fold enrichment. In the molecular function (MF) category, the top terms are nucleoside-

triphosphatase (NTPase) activity, pyrophosphatase (PPase) activity, and mRNA binding;

in the biological process (BP) category, the top terms are intracellular transport, protein

transport, and establishment of protein localization; in the cellular component (CC)

category, plant-type vacuole, cell-cell junction, and plasmodesmata are enriched

(Supporting Information Fig. S2, Table S2), indicating that Pi depletion impacts NTP-

and PPi-dependent cellular events required for the endomembrane trafficking.

In comparison to the previous study by Lan et al. (Lan *et al.*, 2012), which examined

quantitative total protein profiling in Pi-deficient versus Pi-sufficient *Arabidopsis* roots,

we found that among 209 significantly up-regulated proteins in their datasets (Pi-

deficient/Pi-sufficient plants ratio of ≥ 1.29), 57 proteins (including 13 integral MPs)

overlapped with the 4,317 proteins we identified from the Pi-limited roots (Supporting

Information Table S2**)**. Although this overlap accounts for approximately 27.3% of the

PSI proteins reported in their research, we identified an additional 902 integral MPs,

emphasizing that enriching MPs for proteomics could greatly expand the inventory of

identified proteins implicated in Pi starvation responses (Supporting Information Table

S2).

As anticipated, several *At*PHT1 members were downregulated in the Pi-limited root of

cnih5 versus WT (Table **1**). Specifically, downregulation of *At*PHT1;2, *At*PHT1;3, or

*At*PHT1;4 was observed in at least three ratios (fold changes), with at least one ratio

showing FDR adj. p-value ≤ 0.05. While not statistically significant, the downregulation

of *At*PHT1;1 and *At*PHT1;5, along with the upregulation of *At*PHF1, displayed a

consistent trend across four ratios. In contrast, *At*PHT1;9 was upregulated in the Pi-

limited root of *cnih5* (Table **1**). These proteomic data were closely correlated with our

immunoblot results for *At*PHT1;1/2/3, *At*PHT1;4, and *At*PHF1 in the Pi-limited roots of

cnih5 (Fig. **2a**), confirming the reliability of our quantitative proteomic analysis.

Volcano plots summarized the results from the four comparison groups, highlighting

the differentially expressed *At*PHT1 members and *At*PHF1 in green (**Fig. 3d**). Due to the

variability in the ratios of quantified *At*PHT1s across two biological replicates, we

increased the stringency of our protein profile comparisons. We defined DEPs as those

with at least two ratios ≥ 1.25 or ≤ 0.75, without applying FDR adjusted p-value cutoffs,

and excluded candidates showing any opposite trends (≥ 1.25 or ≤ 0.75). As a result, we

identified 372 proteins that were upregulated and 106 proteins that were downregulated

in *cnih5* (Supporting Information Table S3).

The 372 upregulated proteins in the Pi-limited roots of *cnih5* were subjected to the

enrichment analysis of GO and KEGG pathways (FDR adj. *p*-value < 0.05). In the MF

category, the top terms include copper ion binding, antioxidant activity, and

oxidoreductase activity acting on the CH–OH group of donors NAD or NA; in the BP

category, the top terms are sulfur compound biosynthetic process, purine nucleotide

metabolic process, and purine-containing compound metabolic process; in the CC

category, polysomal ribosome, polysome, and cytosolic ribosome are enriched

(Supporting Information Fig. S3a, Table S3), likely reflecting compensation mechanisms

for loss of *At*CNIH5. In particular, when we focused on the 51 upregulated integral MPs

and subjected them to the GO enrichment analyses, we discovered significant

associations with transporter activities (Supporting Information Fig. S3b). The KEGG

pathway enrichment analyses of the 372 upregulated proteins identified the most relevant

pathways related to ascorbate and aldarate metabolism, sulfur metabolism, and fructose

and mannose metabolism (Supporting Information Fig. S3c). This suggests that the

upregulated proteins in the Pi-limited *cnih5* roots are associated with carbohydrate

metabolic pathways that help protect cells against oxidative stress. Additionally, the

KEGG analyses of the 51 upregulated integral MPs revealed interactions related to

SNARE-mediated vesicular transport (Supporting Information Fig. S3d). Overall, the

dysfunction of *At*CNIH5 leads to alterations in vesicular trafficking.

In the GO enrichment analysis of the 106 downregulated proteins, the most significant

terms within the MF category include 7S RNA binding, fatty acid elongase activity,

structural constituent of chromatin, and transporter activity; for the BP categories, the top

terms are protein transport, cellular macromolecule localization, and establishment of

protein localization; in the CC categories, the enriched terms are the AP-type membrane

coat adaptor complex, membrane coat, coated membrane, and cytoplasmic vesicle

membrane (Fig. **4a** and Supporting Information Table S3). These results can be attributed

to the downregulation of proteins involved in the Golgi-associated clathrin assembly and

the signal recognition particle (SRP) subunits that mediate co-translational membrane

protein targeting (Supporting Information Table S3). We thus surmised that the loss of

*At*CNIH5 leads to negative feedback regulation on post-Golgi membrane trafficking,

which reduces the futile coated-vesicle assembly, and on protein synthesis to prevent the

toxic accumulation of *At*CNIH5-dependent cargoes in the ER. The enriched KEGG

pathways also include fatty acid elongation, protein export, cutin, suberin, and wax

biosynthesis, and biosynthesis of nucleotide sugars (Fig. **4b**). Overall, the downregulated

proteins in the Pi-limited root of *cnih5* are associated with protein export and localization,

acyl-lipid metabolism, and nucleotide sugar biosynthesis.

Analysis of protein–protein interaction between *At*CNIH5 and its potential

membrane cargoes

Assuming that potential cargoes of *At*CNIH5 are downregulated integral MPs in *cnih5*,

we narrowed the number of candidates to 31 (Table **2**). This list is over-represented with

three hypersensitive response-like lesion-inducing protein (HRLI)-like proteins

(AT3G23175, AT3G23190, and AT4G14420), two *β*-ketoacyl-CoA synthases (*At*KCS2

and *At*KCS20), and two *At*PHT1 members (*At*PHT1;1 and *At*PHT1;3) (Table **2**).

Although *At*HRLI-like proteins have not been characterized, the *N. benthamiana*

*Nb*HRLI4 conferred salicylic acid-mediated resistance to turnip mosaic virus (TuMV)

(Wu *et al.*, 2021). *AtKCS* genes encode 3-ketoacyl CoA synthases catalyzing the rate-

limiting condensation step of VLCFAs biosynthesis (Evenson & Post-Beittenmiller,

1995). *AtKCS2* and *AtKCS20* are highly expressed in the root endodermis and redundant

in root suberin biosynthesis (Lee *et al.*, 2009). Interestingly, two alcohol-forming fatty

acyl-CoA reductases, *At*FAR3/CER4 and *At*FAR5, are also downregulated and over-

represented in the KEGG pathway of cutin, suberin and wax biosynthesis (Fig. **4b** and

Supporting Information Table S3), suggesting that biosynthesis of VLCFAs and their

derivative extracellular aliphatic compounds are misregulated in the Pi-limited roots of

cnih5. Nevertheless, since *At*HRLI-like proteins and *At*KCS are ER-localized integral

MPs, the downregulation of these proteins may be a secondary effect due to the

impairment of *At*CNIH5.

We then selected a subset of the potential membrane cargoes (highlighted in gray in

Table **2**) to test their interaction with *At*CNIH5 based on the two criteria: (1) a TM protein

located in post-ER compartments of the secretory pathway and (2) having four ratios $\leq$

0.75 or at least two ratios $\leq$ 0.75 with at least one ratio showing FDR adj. *p*-value $\leq$ 0.05.

With the yeast SUS, the nutrient selection and LacZ *β*-galactosidase assays suggested that

besides *At*PHT1;1 and *At*PHT1;4 (Chiu *et al.*, 2024), *At*CNIH5 interacted with *At*PHT1;2,

*At*PHT1;5, *At*PHT1;7 and *At*PHT1;9 (Fig. **5a**), suggesting that *At*CNIH5 recognizes

conserved motifs among several *At*PHT1 members. We also confirmed the interaction of

*At*CNIH5 with the PM-localized receptor-like kinase *At*LRK10L-1.1 (AT1G18390),

nucleotide exchange factor *At*SOS (AT1G26650), and organic cation/carnitine

transporter *At*OCT1 (AT1G73220), the Golgi-localized nucleotide sugar transporters

*At*URGT6 (AT1G34020) and glucuronoxylan 4-O-methyltransferase *At*GXM2

(AT4G09990), and the tonoplast (TP)-localized detoxification efflux carriers *At*DTX21

(AT1G33110) and *At*DTX35 (AT4G25640) (Fig. **5b)**. We obtained similar results when

the expression constructs for *At*SOS, *At*OCT1 *At*URGT6, *At*DTX21, and *At*DTX35 were

generated with different fusion orientations (Supporting Information Fig. S4). *AtOCT1* is

a PSI gene (Liu *et al.*, 2016). Its expression is restricted to the vasculature of the root and

at sites of lateral root formation (Lelandais-Brière *et al.*, 2007). Therefore, the interaction

of *At*CNIH5 and *At*OCT1 is likely to occur in the vascular tissues. *At*URGT6 and

*At*GXM2 are cell wall-modifying enzymes. *At*URGT6 is involved in the biosynthesis of

rhamnogalacturonan-I (RG-I), a component of pectin found in mucilage and various types

of plant cell walls (Saez-Aguayo *et al.*, 2020). *At*GXM2 is a methyltransferase that

catalyzes 4-O-methylation of glucuronic acid (GlcA) side chains present on the

hemicellulose xylan (Lee *et al.*, 2012; Urbanowicz *et al.*, 2012; Yuan *et al.*, 2014).

AtGXM2 is upregulated in response to Pi starvation (Liu *et al.*, 2016) and preferentially

expressed in the root hair (Salazar-Henao & Schmidt, 2016). *At*DTX21 and *At*DTX35

belong to the multidrug and toxic compound extrusion (MATE)-type transporter

superfamily. *At*DTX35 is also low Pi-responsive (Liu *et al.*, 2016) and functions as a

turgor-regulating chloride channel associated with root hair elongation (Zhang *et al.*,

2017). Defective *At*CNIH5 may modulate cell wall elasticity and root hair growth under

Pi limitation.

To explore other potential membrane cargoes of *At*CNIH5 that may be missing in our

iTRAQ-based proteomic data, we set out in parallel to find root PSI genes encoding

transporters that are co-regulated with *AtCNIH5*. We queried the RNA-seq dataset from

Arabidopsis seedlings subjected to Pi deprivation (P0: Pi-sufficiency; P1: one day of Pi

deprivation; P3: three days of Pi deprivation) (Liu *et al.*, 2016). Genes with expression

levels less than 1 RPKM were excluded. We filtered root PSI genes that displayed an expression pattern similar to that of *AtCNIH5* based on (1) the $\log_2FC \geq 0.5$ (P1/P0), (2) the $\log_2FC \geq 1$ (P3/P0), and (3) the difference of $\log_2FC \geq 0.5$ (P3/P0−P1/P0). Such cutoffs were applied to root PSI genes because the agriGO enrichment analysis (Tian *et al.*, 2017) yielded a satisfactory list for further investigation, including nearly all *AtPHT1s* and more than ten other genes encoding transporters (Supporting Information Table S4). Among the list, the PM-localized *At*OCT1, potassium transporter *At*HAK5 (AT4G13420), sulfate transporter *At*SULTR1;3 (AT1G22150), and the TP-localized Pi transporter *At*VPE1 (AT3G47420) were selected for testing interaction with *At*CNIH5 based on the tripartite split-GFP association in agro-infiltrated *N. benthamiana* leaves (Liu *et al.*, 2018; Liu, 2021). We also included *At*SOS, *At*URGT6, *At*DTX21, and *At*DTX35 for testing but not *At*LRK10L-1.1 or *At*GXM2 due to the same membrane topology of *At*CNIH5 and *At*GXM2 and the failure to find any agrobacteria transformants carrying the *At*LRK10L-1.1 construct. The S10-tagged *At*CNIH5 (S10-*At*CNIH5) and the S11-tagged fusion protein of interest correctly reached their subcellular destinations, except that *At*SOS-S11 and *At*SULTR1;3-S11 were retained in the ER (Fig. **6a**). Compared to the positive control *At*PHT1;1-S11 and the negative control 3xHA-S11 as recently reported (Chiu *et al.*, 2024), S10-*At*CNIH5 interacted with *At*OCT1-S11, *At*SOS-S11, *At*URGT6-S11, *At*DTX21-S11, and *At*DTX35-S11 *in planta*, yielding fluorescence signals as a result of GFP complementation, but failed to interact with *At*HAK5-S11, *At*SULTR1;3-S11, and *At*VPE1-S11 (Fig. **6b**). Taken together, our quantitative proteomics successfully identified specific membrane cargoes of *At*CNIH5, including *At*PHT1s.

The C-terminal acidic residue of *At*CNIH5 is not required to interact with *At*PHT1;1

As several *At*PHT1 members were identified as prominent cargoes of *At*CNIH5, we were curious to find out which motif of *At*CNIH5 interacts with *At*PHT1;1. *At*CNIHs have three putative TMD, with the N terminus facing the cytosol, the C terminus facing the ER lumen, and at least four putative functional motifs (Wudick *et al.*, 2018; Chiu *et al.*, 2024). The putative FLN motif in the second TMD of the yeast Erv14, known as a cargo-binding site (Pagant *et al.*, 2015), is present in *At*CNIH2–4, denoted as the YLC motif. The putative cytosolic IFRTL motif, located between the second and third TMD

of Erv14 and interacting with Sec24 (Powers & Barlowe, 2002), is also conserved in

*At*CNIH2–4 with several residues changed, denoted as the IFNNL motif. The C-terminal

acidic motif, which is only found in plants and fungi, is required for Erv14 to assist the

PM targeting of the yeast Nha1, Pdr12, and Qdr2 as well as for the rice *Os*CNIH1 to aid

the Golgi targeting of *Os*HKT1;3 (Rosas-Santiago *et al.*, 2017). Besides the cornichon

motif that is well conserved among all *At*CNIHs, *At*CNIH5 has the putative YLC and

IFNNL motifs with several residues changed and a shorter C-terminal putative acidic

motif with one aspartic acid residue (Supporting Information Fig. S5). We designed a

series of C-terminally truncated forms of *At*CNIH5. For unknown reasons, we failed to

obtain the clone expressing *At*CNIH5 1–102. Therefore, we only tested the interaction of

*At*PHT1;1 with *At*CNIH5 1–26, *At*CNIH5 1–49, *At*CNIH5 1–92, and *At*CNIH5 1–132 in

the yeast SUS (Fig. **7a**). *At*CNIH5 1–26, which contains only the first TMD, is sufficient

to interact with *At*PHT1;1. In comparison, *At*CNIH5 1–132, which has the last three

amino acids deleted (including the C-terminal aspartic acid), could still interact with

*At*PHT1;1 (Fig. **7b**). These results indicated that the putative C-terminal motif of

*At*CNIH5 is not necessary for interaction with *At*PHT1;1 and *At*CNIH5 employs a

different mechanism to interact with *At*PHT1;1 from those used by the fungi and rice

CNIHs for their cognate cargoes.

Increasing *in-situ* *At*CNIH5 expression/activity enables robust plant growth

Since the loss of *At*CNIH5 led to decreased abundance of *At*PHT1s (Table **2**),

inspecting whether overexpression of *At*CNIH5 can improve plant fitness under Pi

limitation was imperative. We found that the ectopic expression of RFP-*At*CNIH5, driven

by the cauliflower mosaic virus (CaMV) 35S promoter, resulted in distorted root

morphology (Supporting Information Fig. S6). This aberrant structure was likely caused

by deformation of the ER, a phenomenon previously associated with the overexpression

of Erv14 in budding yeast (Lagunas-Gomez *et al.*, 2023). Instead, the transgenic lines

expressing the genomic GFP-*At*CNIH5 fragment (*AtCNIH5pro*: GFP-*AtCNIH5*) in the

cnih5 background restored the expression of *At*PHT1;1/2/3 to the WT-like level and grew

bigger than WT and *cnih5* when Pi is replete (Chiu *et al.*, 2024). We then systematically

examined the growth of these *cnih5* complemented lines under different Pi regimes.

Under Pi sufficiency and modest Pi deficiency—three days of Pi deprivation (0 µM Pi)

or eleven days of suboptimal Pi supply (50 μM Pi), the complemented *cnih5* lines

exhibited higher shoot and/or root fresh weight than the WT and *cnih5* plants (Fig. **8**). In

contrast, under severe Pi deficiency—eleven days of Pi deprivation (0 μM Pi), none of

the complemented lines had a larger shoot size than WT (Fig. **8**). Although we could

detect the expression of GFP-*AtCNIH5* proteins in the Pi-limited root of *cnih5*

complemented lines, it is technically difficult to discern whether GFP-*AtCNIH5* proteins

were more abundant than the endogenous *At*CNIH5 in WT (Supporting Information Fig.

S7). Nonetheless, the higher biomass of *cnih5* complemented lines likely results from

more efficient assimilation of acquired Pi due to increased GFP-*AtCNIH5*

expression/activity, along with the coordinated upregulation of *At*CNIH5 membrane

cargoes beyond *At*PHT1s. We thus inferred that a subtle increase of *in-situ* rather than

ectopic expression/activity of GFP-*AtCNIH5* suffices to activate downstream effectors of

*At*CNIH5, thereby boosting plant growth.

Discussion

CNIHs are the eukaryotic conserved cargo receptors that cycle between the ER and

the early Golgi to select membrane proteins for incorporation into COPII vesicles. In this

study, we aimed to discover potential membrane cargoes of low Pi-induced *At*CNIH5 by

iTRAQ-based proteomic analysis and, in the meantime, to provide the first membrane

proteome for *Arabidopsis* roots subjected to Pi deprivation (Zhou *et al.*, 2022). Given that

our iTRAQ-based proteomic analysis did not include Pi-sufficient WT and *cnih5*

samples, we sought to evaluate whether the identified potential membrane cargoes of

*At*CNIH5 are indeed PSI proteins. To address this, we examined the overlap between the

proteins we identified in the Pi-limited roots and the root PSI genes (P3/P0, FC > 1.5; Liu

et al., 2016). We found minimal overlap, at only 5.8%, between the 4317 identified

proteins and the root PSI genes (Supporting Information Table. S2), which aligned with

the previous study indicating a weak correlation between the protein and transcript

changes following Pi deficiency in *Arabidopsis* root (Lan *et al.*, 2012). Still, of the

integral MPs in the Pi-limited roots of *cnih5*, 7.8% (4 out of 51) of the upregulated and

32.3% (10 out of 31) of the downregulated proteins were classified as root PSI genes

(Supporting Information Table. S5). These results suggested that PSI genes are enriched

5.7-fold among the downregulated integral MPs in the Pi-limited roots of *cnih5* when

compared to the 4317 identified proteins. Therefore, the *At*CNIH5-dependent membrane

cargoes we identified are mainly associated with Pi starvation.

Considering that only integral MPs at the post-ER compartments are potential cargoes

of *At*CNIH5, we tested their interaction with *At*CNIH5. When expressed in the yeast

heterologous expression system, *At*CNIH5 interacted not only with the two major Pi

transporters for Pi uptake, *At*PHT1;1 and *At*PHT1;4 (Chiu *et al.*, 2024), but also with

*At*PHT1;2, *At*PHT1;5, *At*PHT1;7, and *At*PHT1;9 (Fig. **5a**), supporting that *At*PHT1s are

the prominent membrane cargoes of *At*CNIH5 under Pi starvation. This was not

surprising because the amino acid sequences of *At*PHT1s share over 47.4% identity. It

has been shown that the yeast and rice CNIHs used the C-terminal acidic motif to interact

with their cognate cargoes (Rosas-Santiago *et al.*, 2017). However, a recent study

revealed that the cargo binding motif of the pumpkin *Cm*CNIH1 may reside in the N-

terminal rather than the C-terminal region (Wei *et al.*, 2023). This agreed with our results

for *At*CNIH5 (Fig. **7b**), indicating that the TMD1 of *At*CNIH5 contains a motif that

interacts with *At*PHT1s.

Besides *At*PHT1s, we identified other PM-localized membrane proteins

downregulated in the Pi-limited roots of *cnih5*. *At*LRK10L-1.2 is involved in ABA

signaling (Lim *et al.*, 2015). The animal Son of Sevenless (SOS) proteins encode the

guanine nucleotide exchange factors acting on the Ras subfamily of small GTPases

(Bowtell *et al.*, 1992). However, the physiological role of SOS in plants remains to be

studied. The PSI *At*OCT1 was also identified as a cargo of *At*CNIH5. Intriguingly, the

oct1 mutant exhibited enhanced root branching (Lelandais-Brière *et al.*, 2007),

phenocopying the higher lateral root density of *cnih5* (Chiu *et al.*, 2024). Although the

heterologous expression of *At*OCT1 in yeast could transport carnitine, it remains to be

determined which substrate *At*OCT1 transports that could negatively affect lateral root

development. Besides *At*URGT6 and *At*GXM2, various Golgi-localized TM proteins

involved in plant cell wall modification such as *At*QUL2 (AT2G03480) (Mouille *et al.*,

2007), *At*GAE1 (AT4G30440) (Mølhøj *et al.*, 2004), *At*GAUT3 (AT4G38270) (Sterling

et al., 2006), and *At*TMN9 (AT5G25100) (Hiroguchi *et al.*, 2021), are remarkably

downregulated in the Pi-limited *cnih5* roots (Table **2**). By contrast, the Golgi-localized

hydroxyproline-O-galactosyltransferases *At*GALT9 (AT1G53290), which is involved in

pectin metabolism (Zhang *et al.*, 2021), is most highly and significantly upregulated in

the Pi-limited *cnih5* roots (Fig. **3d**). Previous analyses of root-specific co-expression have

indicated that early PSR genes are involved in cell wall remodeling, highlighting the

importance of this process in response to Pi deficiency (Liao *et al.*, 2011; Lin *et al.*,

2011). It is likely that *At*CNIH5 plays a critical role in Pi starvation-triggered cell wall

composition changes. Nevertheless, the difference in cell wall structure between WT and

cnih5 awaits investigation. The TP-localized *At*DTX21 and *At*DTX35 were identified as

the potential cargoes of *At*CNIH5 (Fig. **5b** and **6b**). While *AtDTX35* is a PSI gene,

AtDTX21 is co-induced with the type III effectors secreted by *Pseudomonas syringae* and

may mediate the detoxification of chlorinated herbicide atrazine (de Torres-Zabala *et al.*,

2007; Ramel *et al.*, 2007). More studies are required to unravel the physiological role of

*At*DTX21 and *At*DTX35 in Pi starvation responses. Intriguingly, *At*CNIH5 failed to

interact with many other manifest PSI transporters such as *At*HAK5, *At*SULTR1;3, and

*At*VPE1 (Fig. **5b**). In fact, we identified the PSI transporters *At*VPE1 and its paralog

*At*VPE3 in our proteomic analysis, but their expression did not differ between WT and

cnih5 (Supporting Information Table S2), suggesting that *At*CNIH5 recognizes specific

PSI transporters through an unidentified mechanism.

In conclusion, an Azo-solubilized MME approach in conjunction with quantitative

proteomics was used to demonstrate the downregulation of *At*PHT1s and other membrane

proteins associated with Pi starvation in the roots of *cnih5* (Fig. **9**). Our analyses validated

the interaction between *At*CNIH5 and several membrane cargoes at post-ER

compartments, including *At*PHT1s, *At*OCT1, *At*URGT6, *At*DTX21, and *At*DTX35

transporters. We also showed that *At*CNIH5 increased in the Pi-depleted root as well as

in the Pi-replete root of *pho2*, likely due to a higher demand for the ER export of *At*PHT1s.

Moreover, introducing the genomic GFP-*AtCNIH5* fragment boosted plant growth across

a wide range of Pi supplies. Therefore, we propose that *At*CNIH5 acts as a Pi deficiency-

inducible hub to facilitate the ER-to-Golgi trafficking of specific membrane cargoes,

thereby improving plant fitness under suboptimal Pi availability.

Accession Numbers

Sequence data from this article can be found in the Arabidopsis Genome Initiative under

the following accession numbers: *At*CNIH5 (AT4G12090), *At*PHT1;1 (AT5G43350),

*At*PHT1;2 (AT5G43370), *At*PHT1;5 (AT2G32830), *At*PHT1;7 (AT3G54700), *At*PHT1;9

(AT1G76430), *At*LRK10L-1.1 (AT1G18390), *At*GXM2 (AT4G09990), *At*SOS

(AT1G26650)*, At*DTX21 (AT1G33110)*, At*DTX35 (AT4G25640)*, At*URGT6

(AT1G34020), *At*OCT1 (AT1G73220), *At*HAK5 (AT4G13420), *At*VPE1 (AT3G47420),

*At*SULTR1;3 (AT1G22150).

Acknowledgments

This work was supported by the Ministry of Science and Technology (MOST 108-2311-

B-007-003-MY3) and the National Science and Technology Council (NSTC 112-2313-

B-007-001-MY3). We acknowledge Ms. He-Ting Yang for cloning the CDS of *At*OCT1,

*At*HAK5, *At*VPE1, and *At*SULTR1;3, and Mr. Po-Ruey Huang for cloning the constructs

expressing the truncated *At*CNIH5 variants. We also thank Mr. Cheng-Da Tsai for testing

the interaction of PLVCub-*At*CNIH5 with NubG-*At*PHT1;2, NubG-*At*PHT1;5, NubG-

*At*PHT1;7 and NubG-*At*PHT1;9. We thank Dr. Tzvy-Jen Chiou at Academia Sinica,

Taiwan (R.O.C.), for sharing anti-PHO1 and anti-PHF1 antibodies and the technical

support from the confocal imaging core (sponsored by MOST 110-2731-M-007-001),

National Tsing Hua University.

Competing interests

None declared

Author contributions

T.-Y. L. conceived the project, designed the experiments, analyzed the experiments, and

wrote the article. M.-H. T., J.-Y. W., H.-F. L., C.-Y. C., H.-X. C. and C.-A. L. performed

and analyzed the experiments. T.-Y. L., M.-H. T. and J.-Y. W. contributed equally.

Data availability

All data supporting the findings of this study are available within the article and the

Supporting Information (Figs S1–S7, Tabs S1–S5).

References

Abas L, Luschnig C (2010) Maximum yields of microsomal-type membranes from small
amounts of plant material without requiring ultracentrifugation. Analytical
Biochemistry **401:** 217-227

Arpat AB, Magliano P, Wege S, Rouached H, Stefanovic A, Poirier Y (2012)

Functional expression of PHO1 to the Golgi and *trans*-Golgi network and its role
in export of inorganic phosphate. The Plant Journal **71:** 479-491

Barlowe CK, Miller EA (2013) Secretory Protein Biogenesis and Traffic in the Early
Secretory Pathway. Genetics **193:** 383-410

Bayle V, Arrighi J-F, Creff A, Nespoulous C, Vialaret J, Rossignol M, Gonzalez E,
Paz-Ares J, Nussaume L (2011) *Arabidopsis thaliana* High-Affinity Phosphate
Transporters Exhibit Multiple Levels of Posttranslational Regulation. The Plant
Cell **23:** 1523-1535

Bökel C, Dass S, Wilsch-Bräuninger M, Roth S (2006) *Drosophila* Cornichon acts as
cargo receptor for ER export of the TGFα-like growth factor Gurken.
Development **133:** 459-470

Bowtell D, Fu P, Simon M, Senior P (1992) Identification of murine homologues of the
Drosophila son of sevenless gene: potential activators of ras. Proceedings of the
National Academy of Sciences **89:** 6511-6515

Brown KA, Chen B, Guardado-Alvarez TM, Lin Z, Hwang L, Ayaz-Guner S, Jin S,
Ge Y (2019) A photocleavable surfactant for top-down proteomics. Nature
Methods **16:** 417-420

Brown KA, Tucholski T, Eken C, Knott S, Zhu Y, Jin S, Ge Y (2020) High-
Throughput Proteomics Enabled by a Photocleavable Surfactant. Angewandte
Chemie International Edition **59:** 8406-8410

Chen J, Wang Y, Wang F, Yang J, Gao M, Li C, Liu Y, Liu Y, Yamaji N, Ma JF,
Paz-Ares J, Nussaume L, Zhang S, Yi K, Wu Z, Wu P (2015) The Rice CK2
Kinase Regulates Trafficking of Phosphate Transporters in Response to
Phosphate Levels. The Plant Cell **27:** 711-723

Chiu C-Y, Tsai C-D, Wang J-Y, Tsai M-H, Kanno S, Lung H-F, Liu T-Y (2024)
Phosphate Starvation-Induced CORNICHON HOMOLOG 5 as Endoplasmic
Reticulum Cargo Receptor for PHT1 Transporters in *Arabidopsis*. bioRxiv**:**
2024.2006.2020.599911

Chopin F, Wirth J, Dorbe M-F, Lejay L, Krapp A, Gojon A, Daniel-Vedele F (2007)
The Arabidopsis nitrate transporter AtNRT2.1 is targeted to the root plasma
membrane. Plant Physiology and Biochemistry **45:** 630-635

Dancourt J, Barlowe C (2010) Protein Sorting Receptors in the Early Secretory
Pathway. Annual Review of Biochemistry **79:** 777-802

de Torres-Zabala M, Truman W, Bennett MH, Lafforgue G, Mansfield JW,
Rodriguez Egea P, Bögre L, Grant M (2007) *Pseudomonas syringae* pv. *tomato*
hijacks the *Arabidopsis* abscisic acid signalling pathway to cause disease. The
EMBO Journal **26:** 1434-1443

Evenson KJ, Post-Beittenmiller D (1995) Fatty Acid-Elongating Activity in Rapidly
Expanding Leek Epidermis. Plant Physiology **109:** 707-716

Ge SX, Jung D, Yao R (2019) ShinyGO: a graphical gene-set enrichment tool for
animals and plants. Bioinformatics **36:** 2628-2629

González E, Solano R, Rubio V, Leyva A, Paz-Ares J (2005) PHOSPHATE
TRANSPORTER TRAFFIC FACILITATOR1 Is a Plant-Specific SEC12-
Related Protein That Enables the Endoplasmic Reticulum Exit of a High-Affinity
Phosphate Transporter in *Arabidopsis*. The Plant Cell **17:** 3500-3512

Hallgren J, Tsirigos KD, Pedersen MD, Almagro Armenteros JJ, Marcatili P,
Nielsen H, Krogh A, Winther O (2022) DeepTMHMM predicts alpha and beta
transmembrane proteins using deep neural networks. bioRxiv**:**

2022.2004.2008.487609

Harmel N, Cokic B, Zolles G, Berkefeld H, Mauric V, Fakler B, Stein V, Klöcker N (2012) AMPA Receptors Commandeer an Ancient Cargo Exporter for Use as an Auxiliary Subunit for Signaling. PLOS ONE **7**: e30681

Hiroguchi A, Sakamoto S, Mitsuda N, Miwa K (2021) Golgi-localized membrane protein AtTMN1/EMP12 functions in the deposition of rhamnogalacturonan II and I for cell growth in Arabidopsis. Journal of Experimental Botany **72**: 3611-3629

Hooper CM, Castleden IR, Tanz SK, Aryamanesh N, Millar AH (2016) SUBA4: the interactive data analysis centre for Arabidopsis subcellular protein locations. Nucleic Acids Research **45**: D1064-D1074

Huang T-K, Han C-L, Lin S-I, Chen Y-J, Tsai Y-C, Chen Y-R, Chen J-W, Lin W-Y, Chen P-M, Liu T-Y, Chen Y-S, Sun C-M, Chiou T-J (2013) Identification of Downstream Components of Ubiquitin-Conjugating Enzyme PHOSPHATE2 by Quantitative Membrane Proteomics in *Arabidopsis* Roots. The Plant Cell **25**: 4044-4060

Krogh A, Larsson B, von Heijne G, Sonnhammer ELL (2001) Predicting transmembrane protein topology with a hidden markov model: application to complete genomes. Journal of Molecular Biology **305**: 567-580

Kurokawa K, Nakano A (2018) The ER exit sites are specialized ER zones for the transport of cargo proteins from the ER to the Golgi apparatus. The Journal of Biochemistry **165**: 109-114

Lagunas-Gomez D, Yañez-Dominguez C, Zavala-Padilla G, Barlowe C, Pantoja O (2023) The C-terminus of the cargo receptor Erv14 affects COPII vesicle formation and cargo delivery. Journal of Cell Science **136**

Lan P, Li W, Schmidt W (2012) Complementary Proteome and Transcriptome Profiling in Phosphate-deficient Arabidopsis Roots Reveals Multiple Levels of Gene Regulation. Molecular & Cellular Proteomics **11**: 1156-1166

Lee C, Teng Q, Zhong R, Yuan Y, Haghighat M, Ye Z-H (2012) Three Arabidopsis DUF579 Domain-Containing GXM Proteins are Methyltransferases Catalyzing 4-*O*-Methylation of Glucuronic Acid on Xylan. Plant and Cell Physiology **53**: 1934-1949

Lee S-B, Jung S-J, Go Y-S, Kim H-U, Kim J-K, Cho H-J, Park OK, Suh M-C (2009) Two Arabidopsis 3-ketoacyl CoA synthase genes, *KCS20* and *KCS2/DAISY*, are functionally redundant in cuticular wax and root suberin biosynthesis, but differentially controlled by osmotic stress. The Plant Journal **60**: 462-475

Lelandais-Brière C, Jovanovic M, Torres GAM, Perrin Y, Lemoine R, Corre-Menguy F, Hartmann C (2007) Disruption of *AtOCT1*, an organic cation transporter gene, affects root development and carnitine-related responses in Arabidopsis. The Plant Journal **51**: 154-164

Liao Y-Y, Buckhout TJ, Schmidt W (2011) Phosphate deficiency-induced cell wall remodeling. Plant Signaling & Behavior **6**: 700-702

Lim CW, Yang SH, Shin KH, Lee SC, Kim SH (2015) The AtLRK10L1.2, *Arabidopsis* ortholog of wheat LRK10, is involved in ABA-mediated signaling and drought resistance. Plant Cell Reports **34**: 447-455

Lin W-D, Liao Y-Y, Yang TJW, Pan C-Y, Buckhout TJ, Schmidt W (2011) Coexpression-Based Clustering of Arabidopsis Root Genes Predicts Functional Modules in Early Phosphate Deficiency Signaling. Plant Physiology **155**: 1383-

890   1402

891 **Lin W-Y, Huang T-K, Chiou T-J** (2013) NITROGEN LIMITATION ADAPTATION,
892   a Target of MicroRNA827, Mediates Degradation of Plasma Membrane–
893   Localized Phosphate Transporters to Maintain Phosphate Homeostasis in
894   *Arabidopsis*. The Plant Cell **25:** 4061-4074

895 **Liu T-Y** (2021) Using Tripartite Split-sfGFP for the Study of Membrane Protein–Protein
896   Interactions. *In* JJ Sanchez-Serrano, J Salinas, eds, Arabidopsis Protocols.
897   Springer US, New York, NY, pp 323-336

898 **Liu T-Y, Chou W-C, Chen W-Y, Chu C-Y, Dai C-Y, Wu P-Y** (2018) Detection of
899   membrane protein–protein interaction *in planta* based on dual-intein-coupled
900   tripartite split-GFP association. The Plant Journal **94:** 426-438

901 **Liu T-Y, Huang T-K, Tseng C-Y, Lai Y-S, Lin S-I, Lin W-Y, Chen J-W, Chiou T-J**
902   (2012) PHO2-Dependent Degradation of PHO1 Modulates Phosphate
903   Homeostasis in *Arabidopsis*. The Plant Cell **24:** 2168-2183

904 **Liu T-Y, Huang T-K, Yang S-Y, Hong Y-T, Huang S-M, Wang F-N, Chiang S-F,**
905   **Tsai S-Y, Lu W-C, Chiou T-J** (2016) Identification of plant vacuolar
906   transporters mediating phosphate storage. Nature Communications **7:** 11095

907 **Mølhøj M, Verma R, Reiter W-D** (2004) The Biosynthesis of d-Galacturonate in Plants.
908   Functional Cloning and Characterization of a Membrane-Anchored UDP-d-
909   Glucuronate 4-Epimerase from Arabidopsis. Plant Physiology **135:** 1221-1230

910 **Mouille G, Ralet M-C, Cavelier C, Eland C, Effroy D, Hématy K, McCartney L,**
911   **Truong HN, Gaudon V, Thibault J-F, Marchant A, Höfte H** (2007)
912   Homogalacturonan synthesis in *Arabidopsis thaliana* requires a Golgi-localized
913   protein with a putative methyltransferase domain. The Plant Journal **50:** 605-614

914 **Nakanishi H, Suda Y, Neiman AM** (2007) Erv14 family cargo receptors are necessary
915   for ER exit during sporulation in *Saccharomyces cerevisiae*. Journal of Cell
916   Science **120:** 908-916

917 **Nguyen TV, Gupta R, Annas D, Yoon J, Kim Y-J, Lee GH, Jang JW, Park KH,**
918   **Rakwal R, Jung K-H, Min CW, Kim ST** (2021) An Integrated Approach for the
919   Efficient Extraction and Solubilization of Rice Microsomal Membrane Proteins
920   for High-Throughput Proteomics. Frontiers in Plant Science **12**

921 **Nussaume L, Kanno S, Javot H, Marin E, Nakanishi TM, Thibaud M-C** (2011)
922   Phosphate Import in Plants: Focus on the PHT1 Transporters. Frontiers in Plant
923   Science **2**

924 **Obrdlik P, El-Bakkoury M, Hamacher T, Cappellaro C, Vilarino C, Fleischer C,**
925   **Ellerbrok H, Kamuzinzi R, Ledent V, Blaudez D, Sanders D, Revuelta JL,**
926   **Boles E, André B, Frommer WB** (2004) K$^+$ channel interactions detected by a
927   genetic system optimized for systematic studies of membrane protein interactions.
928   Proceedings of the National Academy of Sciences **101:** 12242-12247

929 **Pagant S, Wu A, Edwards S, Diehl F, Miller Elizabeth A** (2015) Sec24 Is a
930   Coincidence Detector that Simultaneously Binds Two Signals to Drive ER
931   Export. Current Biology **25:** 403-412

932 **Park BS, Seo JS, Chua N-H** (2014) NITROGEN LIMITATION ADAPTATION
933   Recruits PHOSPHATE2 to Target the Phosphate Transporter PT2 for
934   Degradation during the Regulation of *Arabidopsis* Phosphate Homeostasis. The
935   Plant Cell **26:** 454-464

936 **Poirier Y, Bucher M** (2002) Phosphate Transport and Homeostasis in Arabidopsis. The
937   Arabidopsis Book **2002**

Poirier Y, Jaskolowski A, Clúa J (2022) Phosphate acquisition and metabolism in
plants. Current Biology **32:** R623-R629
Powers J, Barlowe C (1998) Transport of Axl2p Depends on Erv14p, an ER–Vesicle
Protein Related to the *Drosophila cornichon* Gene Product. Journal of Cell
Biology **142:** 1209-1222
Powers J, Barlowe C (2002) Erv14p Directs a Transmembrane Secretory Protein into
COPII-coated Transport Vesicles. Molecular Biology of the Cell **13:** 880-891
R Core Team (2021) R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing
Ramel F, Sulmon C, Cabello-Hurtado F, Taconnat L, Martin-Magniette M-L,
Renou J-P, El Amrani A, Couée I, Gouesbet G (2007) Genome-wide
interacting effects of sucrose and herbicide-mediated stress in *Arabidopsis*
thaliana: novel insights into atrazine toxicity and sucrose-induced tolerance.
BMC Genomics **8:** 450
Rosas-Santiago P, Lagunas-Gómez D, Barkla BJ, Vera-Estrella R, Lalonde S, Jones
A, Frommer WB, Zimmermannova O, Sychrová H, Pantoja O (2015)
Identification of rice cornichon as a possible cargo receptor for the Golgi-
localized sodium transporter OsHKT1;3. Journal of Experimental Botany **66:**
2733-2748
Rosas-Santiago P, Lagunas-Gomez D, Yáñez-Domínguez C, Vera-Estrella R,
Zimmermannová O, Sychrová H, Pantoja O (2017) Plant and yeast cornichon
possess a conserved acidic motif required for correct targeting of plasma
membrane cargos. Biochimica et Biophysica Acta (BBA) - Molecular Cell
Research **1864:** 1809-1818
Roth S, Shira Neuman-Silberberg F, Barcelo G, Schüpbach T (1995) cornichon and
the EGF receptor signaling process are necessary for both anterior-posterior and
dorsal-ventral pattern formation in Drosophila. Cell **81:** 967-978
RStudio Team (2020) RStudio: Integrated Development Environment for R. RStudio,
PBC
Saez-Aguayo S, Parra-Rojas JP, Sepúlveda-Orellana P, Celiz-Balboa J, Arenas-
Morales V, Sallé C, Salinas-Grenet H, Largo-Gosens A, North HM, Ralet M-
C, Orellana A (2020) Transport of UDP-rhamnose by URGT2, URGT4, and
URGT6 modulates rhamnogalacturonan-I length. Plant Physiology **185:** 914-933
Salazar-Henao JE, Schmidt W (2016) An Inventory of Nutrient-Responsive Genes in
Arabidopsis Root Hairs. Frontiers in Plant Science **7**
Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,
Remmert M, Söding J, Thompson JD, Higgins DG (2011) Fast, scalable
generation of high-quality protein multiple sequence alignments using Clustal
Omega. Molecular Systems Biology **7:** 539
Spitzer M, Wildenhain J, Rappsilber J, Tyers M (2014) BoxPlotR: a web tool for
generation of box plots. Nature Methods **11:** 121-122
Sterling JD, Atmodjo MA, Inwood SE, Kumar Kolli VS, Quigley HF, Hahn MG,
Mohnen D (2006) Functional identification of an *Arabidopsis* pectin biosynthetic
homogalacturonan galacturonosyltransferase. Proceedings of the National
Academy of Sciences **103:** 5236-5241
The UniProt Consortium (2023) UniProt: the Universal Protein Knowledgebase in
2023. Nucleic Acids Research **51:** D523-D531
Tian T, Liu Y, Yan H, You Q, Yi X, Du Z, Xu W, Su Z (2017) agriGO v2.0: a GO

analysis toolkit for the agricultural community, 2017 update. Nucleic Acids Research **45:** W122-W129

Urbanowicz BR, Peña MJ, Ratnaparkhe S, Avci U, Backe J, Steet HF, Foston M, Li H, O'Neill MA, Ragauskas AJ, Darvill AG, Wyman C, Gilbert HJ, York WS (2012) 4-*O*-methylation of glucuronic acid in *Arabidopsis* glucuronoxylan is catalyzed by a domain of unknown function family 579 protein. Proceedings of the National Academy of Sciences **109:** 14253-14258

Wang X, Liang S (2012) Sample preparation for the analysis of membrane proteomes by mass spectrometry. Protein & Cell **3:** 661-668

Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ (2009) Jalview Version 2—a multiple sequence alignment editor and analysis workbench. Bioinformatics **25:** 1189-1191

Wei L, Liu L, Chen Z, Huang Y, Yang L, Wang P, Xue S, Bie Z (2023) CmCNIH1 improves salt tolerance by influencing the trafficking of CmHKT1;1 in pumpkin. The Plant Journal **114:** 1353-1368

Wickham H, Chang W, Henry L, Lin Pedersen T, Takahashi K, Wilke C, Woo K, Yutani H, Dunnington D, van den Brand T (2016) ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York

Wu X, Lai Y, Rao S, Lv L, Ji M, Han K, Weng J, Lu Y, Peng J, Lin L, Wu G, Chen J, Yan F, Zheng H (2021) Genome-Wide Identification Reveals That *Nicotiana benthamiana* Hypersensitive Response (HR)-Like Lesion Inducing Protein 4 (NbHRLI4) Mediates Cell Death and Salicylic Acid-Dependent Defense Responses to Turnip Mosaic Virus. Frontiers in Plant Science **12**

Wudick MM, Portes MT, Michard E, Rosas-Santiago P, Lizzio MA, Nunes CO, Campos C, Santa Cruz Damineli D, Carvalho JC, Lima PT, Pantoja O, Feijó JA (2018) CORNICHON sorting and regulation of GLR channels underlie pollen tube Ca^{2+} homeostasis. Science **360:** 533-536

Yáñez-Domínguez C, Lagunas-Gómez D, Torres-Cifuentes DM, Bezanilla M, Pantoja O (2023) A cornichon protein controls polar localization of the PINA auxin transporter in *Physcomitrium patens*. Development **150**

Yang Z, Yang J, Wang Y, Wang F, Mao W, He Q, Xu J, Wu Z, Mao C (2020) PROTEIN PHOSPHATASE95 Regulates Phosphate Homeostasis by Affecting Phosphate Transporter Trafficking in Rice. The Plant Cell **32:** 740-757

Yoshimoto K, Hanaoka H, Sato S, Kato T, Tabata S, Noda T, Ohsumi Y (2004) Processing of ATG8s, Ubiquitin-Like Proteins, and Their Deconjugation by ATG4s Are Essential for Plant Autophagy. The Plant Cell **16:** 2967-2983

Yuan Y, Teng Q, Lee C, Zhong R, Ye Z-H (2014) Modification of the degree of 4-*O*-methylation of secondary wall glucuronoxylan. Plant Science **219-220:** 42-50

Zhang H, Guo Z, Zhuang Y, Suo Y, Du J, Gao Z, Pan J, Li L, Wang T, Xiao L, Qin G, Jiao Y, Cai H, Li L (2021) MicroRNA775 regulates intrinsic leaf size and reduces cell wall pectin levels by targeting a galactosyltransferase gene in Arabidopsis. The Plant Cell **33:** 581-602

Zhang H, Zhao F-G, Tang R-J, Yu Y, Song J, Wang Y, Li L, Luan S (2017) Two tonoplast MATE proteins function as turgor-regulating chloride channels in *Arabidopsis*. Proceedings of the National Academy of Sciences **114:** E2036-E2045

Zheng J, Yao L, Zeng X, Wang B, Pan L (2023) *ERV14* receptor impacts mycelial growth via its interactions with cell wall synthase and transporters in *Aspergillus*

niger. Frontiers in Microbiology **14**

Zhou M, Zhu S, Mo X, Guo Q, Li Y, Tian J, Liang C (2022) Proteomic Analysis
Dissects Molecular Mechanisms Underlying Plant Responses to Phosphorus
Deficiency. Cells **11:** 651

Zimmermannová O, Felcmanová K, Rosas-Santiago P, Papoušková K, Pantoja O,
Sychrová H (2019) Erv14 cargo receptor participates in regulation of plasma-
membrane potential, intracellular pH and potassium homeostasis via its
interaction with K^+-specific transporters Trk1 and Tok1. Biochimica et
Biophysica Acta (BBA) - Molecular Cell Research **1866:** 1376-1388

1043

Supporting Information

Fig. S1 Multiple marker abundance profiling (MMAP) of the identified proteins in Azo-

solubilized microsomal protein extraction (MME) samples.

(a) Summary of label-free proteomic analysis of microsomal proteins in *Arabidopsis* WT

and *cnih5* Pi-limited roots.

(b) Relative number of high confidence marker (hcm) proteins for each compartment.

The relative number of distinct proteins for each compartment are displayed in the bar

graph as a percentage of all unique proteins in Arabidopsis (Standard) or all proteins in

the user sample (User).

(c) Relative protein abundance of high hcm proteins for each compartment. The

Normalized Gator Abundance Factor (NGAF) scores for each protein is summed per

subcellular compartment according to the hcm assignments. The abundance-scaling

factor for each compartment describes the skew between the standard observations in

Arabidopsis (Standard) and the estimated abundance observation in the user sample. The

NGAF scores for each protein in the user sample is multiplied by the compartmental

scaling factor and summed per subcellular compartment according to the hcm list. The

sums for each compartment are displayed in the bar graph as a percentage of the whole

protein abundance (Standard) or all proteins in the total user sample (User).

1062

Fig. S2 Gene ontology (GO) analysis of the 4,317 identified proteins in iTRAQ-based

proteomics of *Arabidopsis* WT and *cnih5* Pi-limited roots.

Dot plots of the top GO terms in enrichment analysis for categories of molecular function

(MF), biological process (BP), and cellular component (CC) ranked by fold enrichment.

Fold enrichment compares the sample frequency representing the number of genes

inputted under each GO term to the background frequency of total genes annotated to the

same term in *A. thaliana*. The size of the dot indicates the number of genes in each GO term. The more intense reddish color reflects a lower adjusted *p*-value shown by $-\log_{10}$(FDR).

Fig. S3 GO and KEGG pathway enrichment analyses of the 372 upregulated proteins and the 51 upregulated integral microsomal proteins (MPs) in the Pi-limited root of *cnih5*.
(a, b) Dot plots of the gene ontology (GO) analyses of the 372 upregulated proteins (a) and the 51 upregulated integral MPs (b) ranked by fold enrichment for categories of molecular function, biological process, and cellular component.
(c, d) Dot plots of the KEGG pathway analyses of the 372 upregulated proteins (c) and the 51 upregulated integral MPs (d) ranked by fold enrichment. Fold enrichment compares the sample frequency representing the number of genes inputted under each GO term or KEGG term to the background frequency of total genes annotated to the same term in *A. thaliana*. The size of the dot indicates the number of genes in each GO term or KEGG term. The more intense reddish color reflects a lower adjusted *p*-value shown by $-\log_{10}$ (FDR).

Fig. S4 Analysis of the interaction of PLVCub-*At*CNIH5 with NubG-*At*SOS, NubG-*At*OCT1, NubG-*At*URGT6, NubG-*At*DTX21, and NubG-*At*DTX35 using the yeast split-ubiquitin system (SUS). The co-expression of PLVCub-*At*CNIH5 with NubI and NubG are used as positive and negative controls, respectively. Yeast transformants were grown on synthetic medium without leucine and tryptophan (–LW; the left panel) or on synthetic medium lacking leucine, tryptophan, and adenine with 500 µM methionine (–LWAH; the middle panel) or on SC–LW containing 100 mg/L X-Gal (X-Gal; the right panel).

Fig. S5 Amino acid sequence alignment of *Sc*Erv14 and *At*CNIHs.
The sequence alignment was performed using the Clustal Omega algorithm (Sievers *et al.*, 2011) and visualized using Jalview version 2.11.4.0 (Waterhouse *et al.*, 2009). The degree of amino acid conservation among the CNIHs is represented by a light-to-dark blue color gradient. The transmembrane domains (TMD) are boxed by solid lines. The putative cornichon, YLC, IFNLL, and acidic motifs are boxed by dotted lines. The protein sequences were retrieved from UniprotKB (Uniprot ID: *Sc*Erv14, P53173; *At*CNIH1,

Q9C7D7; *At*CNIH2, Q3EDD7; *At*CNIH3, Q8GWT5; *At*CNIH4, Q84W04; *At*CNIH5, Q9SZ74).

Fig. S6 Expression of *35S:* RFP-*At*CNIH5 in *Arabidopsis*.

Expression of *35S*: RFP-*At*CNIH5 in 5-day-old *Arabidopsis* seedlings grown under Pi-deficient (–P, 0 μM KH_2PO_4, five days of starvation) conditions. Scale bars, 50 μm.

Fig. S7 Protein expression of the endogenous *At*CNIH5 in WT and GFP-*AtCNIH5* in the *AtCNIH5pro*: GFP-*AtCNIH5/cnih5* complementation (COM) plants under Pi deficiency. Immunoblot analysis of *At*CNIH5 and GFP-*At*CNIH5 in the root of *Arabidopsis* 11-day-old WT and COM seedlings under Pi-deficient (–P, 0 μM KH_2PO_4, seven days of starvation) conditions. The bands of *At*CNIH5 and GFP-*At*CNIH5 are indicated by solid arrowhead and open arrowhead, respectively.

Table S1. Oligonucleotides used for plasmid constructs.

Table S2. A list of the identified 4317 proteins in *Arabidopsis* Pi-limited WT and *cnih5* roots with GO enrichment analysis.

Detailed descriptions of our proteomic analysis, including the protein scores and mass, annotation, predicted transmembrane helices, quantification for the identified proteins, and a list of the 61 overlapped identified proteins between our datasets and the Pi starvation root proteome by (Lan *et al.*, 2012).

Table S3. A list of the 372 upregulated and 106 downregulated proteins in Pi-limited *cnih5* roots with GO enrichment analysis.

Detailed descriptions, including annotation, predicted transmembrane helices, and quantification for the proteins identified in proteomic analysis.

Table S4. A list of the 19 root Pi starvation-induced (PSI) genes encoding transporters.

Table S5. Enrichment of root Pi starvation-induced (PSI) genes in the iTRAQ-based proteomic analysis.

Figure legends

Figure 1. iTRAQ-based quantitative proteomic analysis of Pi-limited *Arabidopsis* roots.

(a) Schematic workflow of Azo-solubilized microsomal protein extraction for iTRAQ-based quantitative proteomic analysis of *Arabidopsis* Pi-limited roots.

(b) UV-Vis spectrum monitoring the degradation of 1% Azo as a function of UV irradiation time.

Figure 2. Immunoblot analysis of root microsomal proteins extracted from WT, *cnih5*, and *pho2* mutants.

(a, b) Protein expression of *At*PHF1, *At*PHO1, *At*PHT1;1/2/3, and *At*NRT2.1 (a) and *At*CNIH5 (b) in the root of 11-day-old seedlings of *Arabidopsis* WT, *cnih5*, and *pho2* under Pi-sufficient (+P) and Pi-deficient (–P, 0 μM KH_2PO_4, five days of starvation) conditions. The band of *At*CNIH5 is indicated by arrowhead. The relative protein expression level was normalized with the corresponding amido black staining and relative to the WT control. Error bars represent SE (n = 2–4). The applied extraction methods were shown as indicated.

Figure 3. Summary of iTRAQ-based proteomic analysis of microsomal proteins (MPs) in Pi-limited WT and *cnih5* roots.

(a) Venn diagram for comparing MPs and transmembrane (TM) proteins of the 4317 identified proteins. Subcellular localization was analyzed by SUBA5. TM protein prediction or annotation was performed by deepTMHMM or UniprotKB.

(b) Distribution of subcellular localizations of the identified 4317 proteins analyzed by SUBA5. Note that not all proteins are restricted to a single subcellular location.

(c) Distribution of subcellular localizations of the 915 integral MPs among the identified 1231 TM proteins analyzed by SUBA5. Note that not all proteins are restricted to a single subcellular location.

(d) Volcano plots of differentially expressed proteins (DEPs) in the Pi-limited root of *cnih5*. Results of the four comparison groups (four ratios of *cnih5* versus WT) were shown separately, with the $-\log_{10}$FDR adj. *p*-value plotted against the $\log_2$ fold-change (FC). The vertical dash lines denote ±1.25-FC ($\log_2$ FC ≈ 0.322 and $\log_2$FC ≈ −0.415), while the horizontal dash line denotes FDR adj. *p*-value = 0.05. Upregulated and

downregulated proteins are, respectively, indicated by red and blue dots. Only DEPs with ±2-FC are marked in red or blue. The *At*PHT1s and *At*PHF1 are marked in green. The proteins used for interaction testing are marked in purple.

Figure 4. GO and KEGG pathway enrichment analyses of the 106 downregulated proteins in Pi-limited *cnih5* roots.

(a) Dot plots of the GO enrichment analysis ranked by fold enrichment for categories of molecular function, biological process, and cellular component.

(b) Dot plots of the KEGG pathway enrichment analysis ranked by fold enrichment. Fold enrichment compares the sample frequency representing the number of genes inputted under each GO term or KEGG term to the background frequency of total genes annotated to the same term in *A. thaliana*. The size of the dot indicates the number of genes in each GO term or KEGG term. The more intense reddish color reflects a lower adj. *p*-value adjusted *p*-value shown by $-\log_{10}(\text{FDR})$.

Figure 5. Analysis of the interaction of *At*CNIH5 with its potential ER cargoes by the yeast split-ubiquitin system.

(a) Co-expression of PLVCub-*At*CNIH5 with NubG-*At*PHT1;2, NubG-*At*PHT1;5, NubG-*At*PHT1;7, and NubG-*At*PHT1;9.

(b) Co-expression of PLVCub-*At*CNIH5 with *At*LRK10L-1.1-NubG, *At*SOS-NubG, *At*OCT1-NubG, *At*URGT6-NubG, NubG-*At*GXM2, *At*DTX21-NubG, and *At*DTX35-NubG. The co-expression of PLVCub-*At*CNIH5 with NubI and NubG are used as positive and negative controls, respectively. Yeast transformants were grown on synthetic medium without leucine and tryptophan (–LW; the left panel) or on synthetic medium lacking leucine, tryptophan, adenine, and histidine with 500 μM methionine (–LWAH; middle panel) or on SC–LW containing 100 mg/L X-Gal (X-Gal; the right panel).

Figure 6. Analysis of the interaction of *At*CNIH5 with its potential ER cargoes by the tripartite split-GFP assay in *N. benthamiana* leaves.

(a, b) Subcellular localization (a) and protein–protein interaction analysis (b) of S10-*At*CNIH5, *At*PHT1;1-S11, 3xHA-S11, *At*SOS-S11, *At*OCT1-S11, *At*HAK5-S11, *At*SULTR1;3-S11, *At*URGT6-S11, *At*DTX21-S11, *At*DTX35-S11, and *At*VPE1-S11.

1197   The co-expression of S10-*At*CNIH5 with *At*PHT1;1-S11 and 3xHA-S11 are used as

1198   positive and negative controls, respectively. The percentage of laser power used for

1199   acquisition is shown. Scale bars, 5 μm.

1200

Figure 7. Analysis of the interaction of *At*PHT1;1 and the truncated *At*CNIH5 variants

by the yeast split ubiquitin assay.

(a) Schematic diagram of full-length (FL) and truncated *At*CNIH5 variants. The lengths

of the truncated variants are indicated by the number of residues. The transmembrane

domains and the putative cornichon, YLC, IFNLL, and acidic motifs are shown. The

black box, the transmembrane domain (TMD); the blue box, the cornichon motif; the

green box, the YLC motif; the yellow box, the IFNLL motif. The red box, the acidic

motif. The gray box shows the sequence alignment of the putative acidic motif of *Sc*Erv14

and *At*CNIHs. Noted that *At*CNIH5 has a shorter C-terminal tail carrying only one acidic

residue D134.

(b) Co-expression of *At*PHT1;1-PLVCub with NubG-*At*CNIH5, NubG-*At*CNIH5 1–26,

NubG-*At*CNIH5 1–49, NubG-*At*CNIH5 1–92, and NubG-*At*CNIH5 1–132. The co-

expression of *At*PHT1;1-PLVCub with NubI and NubG are used as positive and negative

controls, respectively. Yeast transformants were grown on synthetic medium without

leucine and tryptophan (–LW; the left panel) or on synthetic medium lacking leucine,

tryptophan, adenine, and histidine with 500 μM methionine (–LWAH; middle panel) or

on SC–LW containing 100 mg/L X-Gal (X-Gal; the right panel).

1218

Figure 8. Increasing *in-situ At*CNIH5 expression/activity boosts plant growth.

The shoot and root fresh weight (FW) of *Arabidopsis* 11-day-old WT, *cnih5* and

AtCNIH5pro: GFP-*AtCNIH5/cnih5* complementation (COM) lines under Pi-sufficient

(+P, 250 μM KH_2PO_4), modest Pi-deficient (–P, 0 μM KH_2PO_4, three days of starvation

or 50 μM KH_2PO_4, eleven days of starvation), and severe Pi-deficient (–P, 0 μM KH_2PO_4,

eleven days of starvation) conditions. Data are visualized using BoxPlotR (Spitzer *et al.*,

2014). Center lines show the medians; the central plus signs (+) show the means; box

limits indicate the 25th and 75th percentiles; whiskers extend to the 5th and 95th

percentiles, and data points are plotted as dots. Data significantly different from the other

groups are indicated by different lowercase letters (n, pools of seedings from independent

experiments; One-Way ANOVA, Duncan's test).

1230

Figure 9. *At*CNIH5-mediated ER-to-Golgi transport of membrane proteins associated

with Pi starvation responses in *Arabidopsis* root.

Under Pi starvation (−P), the transcripts of *CNIH5*, *PHT1* genes, *OCT1*, *GXM2*, and

DTX35 increase in *Arabidopsis* root cells (Liu *et al.*, 2016). As an ER cargo receptor,

CNIH5 assists the ER export of the newly synthesized PHT1s, LRK10L-1.1, SOS, OCT1,

URGT6, GXM2, DTX21, and DTX35 and their loading into COPII vesicles, thus their

corresponding subcellular destinations.

1238

1239   **Table 1.** Summary of *At*PHT1s and *At*PHF1 expression in Pi-limited WT and *cnih5* roots.

1240   The normalized ratios compared the protein expression in Col-0_1, Col-0_2, *cnih5*_1,

1241   and *cnih5*_2 samples. The FDR adj. *p*-values were in the parentheses.

1242

Table 2. List of the 31 downregulated microsomal transmembrane proteins in Pi-limited

cnih5 roots.

The normalized ratios compared the protein expression in Col-0_1, Col-0_2, *cnih5*_1,

and *cnih5*_2 samples. The FDR adj. *p*-values were in the parentheses. The candidates

highlighted in gray were selected for testing protein–protein interaction in this study.

Subcellular localization is predicted or annotated by SUBA5, UniprotKB or experimental

data.

1250

Figure 1. iTRAQ-based quantitative proteomic analysis of Pi-limited *Arabidopsis* roots.

(a) Schematic workflow of Azo-solubilized microsomal protein extraction for iTRAQ-based quantitative proteomic analysis of *Arabidopsis* Pi-limited roots.

(b) UV-Vis spectrum monitoring the degradation of 1% Azo as a function of UV irradiation time.

Figure 2. Immunoblot analysis of root microsomal proteins extracted from WT, *cnih5*, and *pho2* mutants.

(a and b) Protein expression of *At*PHF1, *At*PHO1, and *At*NRT2.1 (a) and *At*CNIH5 (b) in the root of 11-day-old seedlings of *Arabidopsis* WT, *cnih5*, and *pho2* under Pi-sufficient (+P) and Pi-deficient (–P, 0 μM KH$_2$PO$_4$, five days of starvation) conditions. The band of *At*CNIH5 is indicated by arrowhead. The relative protein expression level was normalized with the corresponding amido black staining and relative to the WT control. Error bars represent SE (n = 2–4). The applied extraction methods were shown as indicated.

Figure 3. Summary of iTRAQ-based proteomic analysis of microsomal proteins (MPs) in Pi-limited WT and *cnih5* roots.

(a) Venn diagram for comparing MPs and transmembrane (TM) proteins of the 4317 identified proteins. Subcellular localization was analyzed by SUBA5. TM protein prediction or annotation was performed by deepTMHMM or UniprotKB.

(b) Distribution of subcellular localizations of the identified 4317 proteins analyzed by SUBA5. Note that not all proteins are restricted to a single subcellular location.

(c) Distribution of subcellular localizations of the 915 integral MPs among the identified 1231 TM proteins analyzed by SUBA5. Note that not all proteins are restricted to a single subcellular location.

(d) Volcano plots of differentially expressed proteins (DEPs) in the Pi-limited root of *cnih5*. Results of the four comparison groups (four ratios of *cnih5* versus WT) were shown separately, with the −log$_{10}$FDR adj. *p*-value plotted against the log$_2$ fold-change (FC). The vertical dash lines denote ±1.25-FC (Log$_2$ FC ≈ 0.322 and Log$_2$FC ≈ −0.415), while the horizontal dash line denotes FDR adj. *p*-value = 0.05. Upregulated and downregulated proteins are, respectively, indicated by red and blue dots. Only DEPs with ±2-FC are specified in red or blue. The *At*PHT1s and *At*PHF1 are marked in green. The proteins used for interaction testing are marked in purple.

(a)

(b)

Figure 4. GO and KEGG pathway enrichment analyses of the 106 downregulated proteins in Pi-limited *cnih5* roots.

(a) Dot plots of the GO enrichment analysis ranked by fold enrichment for categories of molecular function, biological process, and cellular component.

(b) Dot plots of the KEGG pathway enrichment analysis ranked by fold enrichment. Fold enrichment compares the sample frequency representing the number of genes inputted under each GO term or KEGG term to the background frequency of total genes annotated to the same term in *A. thaliana*. The size of the dot indicates the number of genes in each GO term or KEGG term. The more intense reddish color reflects a lower adj. *p*-value adjusted *p*-value shown by $-\log_{10}(\text{FDR})$.

(a)

(b)

Figure 5. Analysis of the interaction of *At*CNIH5 with its potential ER cargoes by the yeast split-ubiquitin system.
(a) Co-expression of PLVCub-*At*CNIH5 with NubG-*At*PHT1;2, NubG-*At*PHT1;5, NubG-*At*PHT1;7, and NubG-*At*PHT1;9.
(b) Co-expression of PLVCub-*At*CNIH5 with *At*LRK10L-1.1-NubG, *At*SOS-NubG, *At*OCT1-NubG, *At*URGT6-NubG, NubG-*At*GXM2, *At*DTX21-NubG, and *At*DTX35- NubG. The co-expression of PLVCub-*At*CNIH5 with NubI and NubG are used as positive and negative controls, respectively. Yeast transformants were grown on synthetic medium without leucine and tryptophan (–LW; the left panel) or on synthetic medium lacking leucine, tryptophan, adenine, and histidine with 500 μM methionine (–LWAH; middle panel) or on SC–LW containing 100 mg/L X-Gal (X-Gal; the right panel).

Figure 6. Analysis of the interaction of *At*CNIH5 with its potential ER cargoes by the tripartite split-GFP assay in *N. benthamiana* leaves.

(a and b) Subcellular localization (a) and protein–protein interaction analysis (b) of S10-*At*CNIH5, *At*PHT1;1-S11, 3xHA-S11, *At*SOS-S11, *At*OCT1-S11, *At*HAK5-S11, *At*SULTR1;3-S11, *At*URGT6-S11, *At*DTX21-S11, *At*DTX35-S11, and *At*VPE1-S11. The co-expression of S10-*At*CNIH5 with *At*PHT1;1-S11 and 3xHA-S11 are used as positive and negative controls, respectively. The percentage of laser power used for acquisition is shown. Scale bars, 5 μm.

Figure 7. Analysis of the interaction of *At*PHT1;1 and the truncated *At*CNIH5 variants by the yeast split ubiquitin assay.

(a) Schematic diagram of full-length (FL) and truncated *At*CNIH5 variants. The lengths of the truncated variants are indicated by the number of residues. The transmembrane domains and the putative cornichon, YLC, IFNLL, and acidic motifs are shown. The black box, the transmembrane domain (TMD); the blue box, the cornichon motif; the green box, the YLC motif; the yellow box, the IFNLL motif. The red box, the acidic motif. The gray box shows the sequence alignment of the putative acidic motif of *Sc*Erv14 and *At*CNIHs. Noted that *At*CNIH5 has a shorter C-terminal tail carrying only one acidic residue D134.

(b) Co-expression of *At*PHT1;1-PLVCub with NubG-*At*CNIH5, NubG-*At*CNIH5 1–26, NubG-*At*CNIH5 1–49, NubG-*At*CNIH5 1–92, and NubG-*At*CNIH5 1–132. The co-expression of *At*PHT1;1-PLVCub with NubI and NubG are used as positive and negative controls, respectively. Yeast transformants were grown on synthetic medium without leucine and tryptophan (–LW; the left panel) or on synthetic medium lacking leucine, tryptophan, adenine, and histidine with 500 μM methionine (–LWAH; middle panel) or on SC–LW containing 100 mg/L X-Gal (X-Gal; the right panel).

Figure 8. Increasing *in-situ At*CNIH5 expression/activity boosts plant growth.

The shoot and root fresh weight (FW) of *Arabidopsis* 11-day-old WT, *cnih5* and *AtCNIH5pro*: GFP-*AtCNIH5/cnih5* complementation (COM) lines under Pi-sufficient (+P, 250 μM KH_2PO_4), modest Pi-deficient (–P, 0 μM KH_2PO_4, three days of starvation or 50 μM KH_2PO_4, eleven days of starvation), and severe Pi-deficient (–P, 0 μM KH_2PO_4, eleven days of starvation) conditions. Data are visualized using BoxPlotR . Center lines show the medians; the central plus signs (+) show the means; box limits indicate the 25th and 75th percentiles; whiskers extend to the 5th and 95th percentiles, and data points are plotted as dots. Data significantly different from the other groups are indicated by different lowercase letters (n, pools of seedings from independent experiments; One-Way ANOVA, Duncan's test).

Figure 9. *At*CNIH5-mediated ER-to-Golgi transport of membrane proteins associated with Pi starvation responses in *Arabidopsis* root.

Under Pi starvation (−P), the transcripts of *CNIH5*, *PHT1* genes, *OCT1*, *GXM2*, and *DTX35* increase in *Arabidopsis* root cells . As an ER cargo receptor, CNIH5 assists the ER export of the newly synthesized PHT1s, LRK10L-1.1, SOS, OCT1, URGT6, GXM2, DTX21, and DTX35 and their loading into COPII vesicles, thus their corresponding subcellular destinations.

Table 1. Summary of *At*PHT1s and *At*PHF1 expression in Pi-limited WT and *cnih5* roots.
The normalized ratios compared the protein expression in Col-0_1, Col-0_2, *cnih5*_1 and *cnih5*_2 samples. The FDR adj. *p*-values were in the parentheses.

Accession	Gene name	AGI code	Coverage [%]	Unique peptides	*cnih5*_1 /Col-0_1	*cnih5*_2 /Col-0_1	*cnih5*_1 /Col-0_2	*cnih5*_2 /Col-0_2
Q8VYM2	*AtPHT1;1*	AT5G43350	38	2	0.63 (0.41)	0.54 (0.28)	0.65 (0.31)	0.56 (0.22)
Q96243	*AtPHT1;2*	AT5G43370	30	1	0.61 (0.05)	0.97 (1.00)	0.78 (0.47)	1.24 (0.67)
O48639	*AtPHT1;3*	AT5G43360	10	2	0.58 (0.00)	0.65 (0.24)	0.56 (0.00)	0.74 (0.14)
Q96303	*AtPHT1;4*	AT2G38940	12	6	0.77 (0.40)	0.86 (0.89)	0.79 (0.06)	0.70 (0.00)
Q8GYF4	*AtPHT1;5*	AT2G32830	6	2	0.82 (0.78)	0.73 (0.58)	0.90 (0.87)	0.80 (0.42)
Q9S735	*AtPHT1;9*	AT1G76430	5	2	1.00 (0.99)	1.79 (0.20)	1.27 (0.79)	2.88 (0.00)
Q8GYE0	*AtPHF1*	AT3G52190	40	13	1.45 (0.05)	1.37 (0.36)	1.16 (0.48)	1.14 (0.62)

Coverage [%]: The percentage of the protein sequences covered by identified peptides.

Unique peptides: The total number of distinct peptide sequences unique to the protein group.

Table 2. List of the 31 downregulated microsomal transmembrane proteins in Pi-limited *cnih5* roots.
The normalized ratios compared the protein expression in Col-0_1, Col-0_2, *cnih5*_1 and *cnih5*_2 samples. The FDR adj. *p*-values were in the parentheses. The candidates highlighted in gray were selected for testing protein–protein interaction in this study. Subcellular localization is predicted or annotated by SUBA5, UniprotKB or experimental data.

AGI code	Accession	Gene name	*cnih5*_1/ Col-0_1	*cnih5*_2/ Col-0_1	*cnih5*_1/ Col-0_2	*cnih5*_2/ Col-0_2
		ER				
AT1G04220	Q5XEP9	3-ketoacyl-CoA synthase 2 (*AtKCS2*)	0.73 (0.57)	0.72 (0.66)	0.99 (0.99)	0.98 (1.00)
AT1G34130	Q9FX21	Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit STT3B (*AtSTT3B*)	0.75 (0.53)	0.65 (0.30)	0.91 (0.91)	0.77 (0.38)
AT1G51740	P59277	Syntaxin-81 (*AtSYP81*)	1.18 (0.97)	0.73 (0.88)	0.94 (0.98)	0.58 (0.52)
AT3G23175	Q8GYI3	HR-like lesion-inducing protein-like protein	0.68 (0.23)	0.69 (0.46)	0.94 (0.97)	0.96 (0.99)
AT3G23190	Q9LTC9	HR-like lesion-inducing protein-like protein	0.94 (0.99)	0.63 (0.20)	0.92 (0.92)	0.62 (0.00)
AT3G48570	Q9SMP2	Protein transport protein Sec61 subunit gamma-3 (*AtSEC61G3*)	0.72 (0.30)	0.69 (0.42)	0.71 (0.02)	0.72 (0.12)
AT4G14420	O23298	HR-like lesion-inducing protein-like protein	0.76 (0.53)	0.61 (0.07)	0.81 (0.30)	0.64 (0.00)
AT5G43760	Q9FG87	3-ketoacyl-CoA synthase 20 (*AtKCS20*)	1.03 (1.00)	0.60 (0.54)	1.23 (0.93)	0.72 (0.80)
		Golgi				
AT1G34020	Q9FDZ5	UDP-rhamnose/UDP-galactose transporter 6 (*AtURGT6*)	0.60 (0.49)	0.45 (0.18)	0.68 (0.69)	0.50 (0.27)
AT2G03480	Q3EC77	QUASIMODO2 LIKE 2 (*AtQUL2*)	0.78 (0.73)	0.64 (0.35)	0.87 (0.86)	0.68 (0.27)
AT2G40380	Q9SIY7	Prenylated Rab acceptor 1 (PRA1) family protein B2 (*AtPRA1B2*)	0.66 (0.26)	0.74 (0.72)	0.94 (0.97)	1.05 (0.99)
AT4G09990	Q9T0F7	Glucuronoxylan 4-O-methyltransferase 2 (*AtGXM2*)	0.71 (0.76)	0.73 (0.87)	0.58 (0.36)	0.60 (0.54)
AT4G30440	Q9M0B6	UDP-glucuronate 4-epimerase 1 (*AtGAE1*)	0.83 (0.93)	0.64 (0.58)	0.88 (0.93)	0.68 (0.38)
AT4G38270	Q0WQD2	Galacturonosyltransferase 3 (*AtGAUT3*)	0.87 (0.92)	0.62 (0.32)	0.95 (0.98)	0.69 (0.17)
AT5G25100	Q9C5N2	Transmembrane 9 superfamily member 9 (*AtTMN9*)	0.73 (0.69)	0.81 (0.89)	0.70 (0.33)	0.78 (0.62)
		Vacuole				
AT1G33110	Q8W488	Protein DETOXIFICATION 21 (*AtDTX21*)	0.52 (0.00)	0.76 (0.77)	0.48 (0.00)	0.71 (0.40)
AT4G17340	Q41975	Tonoplast intrinsic protein 2;2 (*AtTIP2-2*)	0.88 (0.95)	0.91 (0.97)	0.65 (0.34)	0.64 (0.40)
AT4G25640	F4JTB3	Protein DETOXIFICATION 35 (*AtDTX35*)	0.86 (0.89)	0.66 (0.31)	0.89 (0.83)	0.68 (0.02)
AT5G46090	Q9FNL3	DUF679 domain membrane protein 6 (*AtDMP6*)	0.74 (0.39)	0.64 (0.22)	0.86 (0.65)	0.74 (0.17)
		PM				
AT1G05570	Q9AUE0	Callose synthase 1 (*AtCALS1*)	0.74 (0.60)	0.73 (0.66)	0.82 (0.72)	0.80 (0.75)
AT1G17620	Q9LNP3	Late embryogenesis abundant (LEA) hydroxyproline-rich glycoprotein family	0.73 (0.54)	0.61 (0.23)	0.93 (0.96)	0.78 (0.68)
AT1G18390	P0C5E2	LEAF RUST 10 DISEASE-RESISTANCE LOCUS RECEPTOR-LIKE PROTEIN KINASE-like 1.2 (*AtLRK10L-1.2*)	0.74 (0.35)	0.72 (0.53)	0.75 (0.09)	0.73 (0.15)
AT1G26650	Q9LQY7	Son of sevenless protein (*AtSOS*)	0.61 (0.22)	0.70 (0.70)	0.61 (0.09)	0.70 (0.27)
AT1G73220	Q9CAT6	Organic cation/carnitine transporter 1 (*AtOCT1*)	0.89 (0.93)	0.86 (0.93)	0.73 (0.04)	0.69 (0.05)
AT2G27140	Q9ZVC6	HSP20-like chaperones superfamily protein	1.00 (0.99)	0.61 (0.13)	1.11 (0.87)	0.65 (0.01)
AT3G02600	Q8LFD1	Lipid phosphate phosphatase 3 (*AtLPP3*)	0.89 (0.92)	0.62 (0.16)	0.97 (0.98)	0.68 (0.02)
AT3G61270	Q9M2D7	Glycosyltransferase (*AtT20K12.170*)	0.84 (0.89)	0.67 (0.46)	0.80 (0.58)	0.64 (0.13)
AT4G01830	Q9SYI3	ATP binding cassette (ABC) transporter B family member 5 (*AtABCB5*)	0.75 (0.79)	0.68 (0.71)	0.99 (1.00)	1.03 (1.00)
AT5G20970	F4K6X6	HSP20-like chaperones superfamily protein	0.85 (0.86)	0.51 (0.00)	1.10 (0.89)	0.66 (0.01)
AT5G43350	Q8VYM2	Inorganic phosphate transporter 1;1 (*AtPHT1;1*)	0.63 (0.41)	0.54 (0.28)	0.65 (0.31)	0.56 (0.22)
AT5G43360	O48639	Inorganic phosphate transporter 1;3 (*AtPHT1;3*)	0.58 (0.00)	0.65 (0.24)	0.56 (0.00)	0.74 (0.14)