Transcriptional Dynamics Uncover the Role of BNIP3 in Mitophagy during Muscle Remodeling in *Drosophila*

(Running title: BNIP3-mediated mitophagy in fly muscle)

Hiroki Taoka[1,3,#], Tadayoshi Murakawa[2,#], Kohei Kawaguchi[1], Michiko Koizumi[1], Tatsuya Kaminishi[4], Yuriko Sakamaki[5], Kaori Tanaka[6], Akihito Harada[6,7], Keiichi Inoue[8], Tomotake Kanki[8], Yasuyuki Ohkawa[6], and Naonobu Fujita[1,2,*]

[1] Cell Biology Center, Institute of Integrated Research, Institute of Science Tokyo, 4259 S2-11 Nagatsuta, Midori-ku, Yokohama, Kanagawa 226-8501, Japan.

[2] School of Life Science and Technology, Institute of Science Tokyo, Yokohama, Kanagawa, Japan.

[3] Biological Science Research Laboratories, Kao Corporation, 2606, Akabane, Ichikai, Haga-gun, Tochigi 321-3497, Japan.

[4] Department of Genetics, Graduate School of Medicine, Osaka University, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan.

[5] Ochanomizu Research Facility (ORF), Bioscience Center, Research Infrastructure Management Center, Institute of Science Tokyo, 1-5-45, Yushima, Bunkyo-ku, Tokyo 113-8510, Japan.

[6] Division of Transcriptomics, Medical Institute of Bioregulation, Kyushu University, 3-1-1, Maidashi, Higashi-ku, Fukuoka 812-8582, Japan.

[7] Department of Multi-Omics, Graduate School of Medical Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-8582, Japan.

[8] Department of Cellular Physiology, Graduate School of Medical Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-8582, Japan.

[#] These authors equally contributed to this work.
[*] Corresponding author. Tel: +81 45 924 5131; E-mail: nafujita@cbc.iir.isct.ac.jp

Keywords: Muscle, *Drosophila*, Remodeling, BNIP3, mitochondria, RNA-seq, autophagy

Impact statement: A time-course RNA-seq analysis of muscle remodeling reveals transcriptional dynamics independent of autophagy and highlights the significance of BNIP3 in mitochondrial degradation *in vivo*.

Abbreviations used: APF, after puparium formation; ATG, autophagy-related; DIOM, dorsal internal oblique muscle; GO, gene ontology; LBWM, larval body wall muscle; LIR, LC3/Atg8-interacting regions; MER, minimal essential region; Mito-QC, mitophagy quality control; PCA, principal component analysis; TEM, transmission electron microscopy; TORC1, target of rapamycin complex 1; 3IL, third instar larvae.

Abstract

Differentiated muscle cells contain myofibrils and well-organized organelles, enabling powerful

contractions. Muscle cell reorganization occurs in response to various physiological stimuli;

however, the mechanisms behind this remodeling remain enigmatic due to the lack of a

genetically trackable system. Previously, we reported that a subset of larval muscle cells is

remodeled into adult abdominal muscle through an autophagy-dependent mechanism in

Drosophila. To unveil the underlying mechanisms of this remodeling, we performed a

comparative time-course RNA-seq analysis of isolated muscle cells with or without autophagy.

It revealed both transcriptional dynamics independent of autophagy and highlighted the

significance of BNIP3-mediated mitophagy in muscle remodeling. Mechanistically, we found

that BNIP3 recruits autophagic machinery to mitochondria through its LC3-interacting (LIR)

motif and minimal essential region (MER), which interact with Atg8a and Atg18a, respectively.

Loss of BNIP3 leads to a substantial accumulation of larval mitochondria, ultimately impairing

muscle remodeling. In summary, this study demonstrates that BNIP3-dependent mitophagy is

critical for orchestrating the dynamic process of muscle remodeling.

Introduction

Differentiated muscle cells are intricate assemblies of myofibrils and well-organized organelles, such as T-tubules and the sarcoplasmic reticulum. Coordinated sliding of actin and myosin filaments is regulated by the dynamics of calcium ions (Ca^{2+}) released from the sarcoplasmic reticulum, while mitochondria provide ATP as the energy source driving this process. Consequently, the structural alignment of organelles and myofibrils within muscle cells is essential for the precise orchestration of contraction (Kawaguchi and Fujita, 2024). Muscle cells are long-lived and continuously undergo atrophy and hypertrophy cycles in response to growth factors, mechanical loading, nutrient status, and aging (Sartori et al., 2021). Muscle organelles also undergo remodeling as the amount of myofibrils changes, a process mediated by the ubiquitin-proteasome system. However, the mechanisms behind these processes are still poorly understood due to the lack of a genetically tractable system for analyzing muscle remodeling.

Macroautophagy, hereafter referred to as autophagy, is essential for muscle function in both flies and mammals. Autophagy is an intracellular degradation system that delivers cytosolic materials to the lysosome (Morishita and Mizushima, 2019; Nakatogawa, 2020). A flat membrane sac elongates and encloses cytosolic content to form an autophagosome, a double membrane structure. The completed autophagosome fuse with lysosomes to degrade the enclosed cargo. To date, over 40 *ATG* genes have been identified as key regulators of autophagosome formation and cargo selectivity (Nakatogawa, 2020). In mammalian muscles, the loss of autophagy leads to reduced contractile function and abnormalities in myofibrils and organelles, including mitochondria and sarcoplasmic reticulum (Karsli-Uzunbas et al., 2014; Masiero et al., 2009; Yoshii et al., 2016). We have previously reported that a subset of larval body wall muscle cells is remodeled into adult dorsal internal oblique muscles (DIOM) through an autophagy-mediated mechanism (Fujita et al., 2017; Murakawa et al., 2020). The remodeled adult DIOMs function during eclosion, persist for approximately 12 hours, and are subsequently eliminated via programmed cell death (Kimura and Truman, 1990). To our knowledge, DIOM remodeling is the sole example of developmentally regulated muscle remodeling, making it an ideal model for studying muscle remodeling in general. Autophagy blockade results in severe phenotypes in DIOM (Fujita et al., 2017; Murakawa et al., 2020), yet the mechanisms by which autophagy loss induces defects remain unclear. Given that autophagy contributes to nutrient supply and transcriptional regulation (Morishita and Mizushima, 2019; Sánchez-Martín et al.,

2019),it is plausible that dysfunction in autophagy could also impact gene expression during

DIOM remodeling.

Autophagy can selectively target various substrates, including protein aggregates,

invading bacteria into the cytosol, and organelles (Lamark and Johansen, 2021; Vargas et al.,

2023). This selectivity is achieved through soluble and transmembrane cargo receptors, which

bridge the cargo and autophagy machinery. Most soluble cargo receptors target ubiquitinated

cargoes and harbor oligomerization domains, ubiquitin-binding domains, and LC3/Atg8-

interacting regions (LIR), which is a four-residue motif [W/F/Y] -X-X-[L/V/I] (Johansen and

Lamark, 2020). In contrast, transmembrane cargo receptors are anchored to target organelles

and recruit autophagy machinery. Soluble and transmembrane cargo receptors confer selectivity

through direct interaction with the ATG machinery. In most cases, in addition to the affinity

with Atg8/LC3 via the LIR motif, interaction with core machinery components, such as ULK1,

FIP200/Atg11, and Atg9, is required (Lamark and Johansen, 2021).

Mitophagy, selective autophagy for mitochondria, is physiologically crucial because

mitochondrial malfunction is harmful, resulting in energy insufficiency or excess reactive

oxygen species (ROS) in cells (Ney, 2015). In addition to damaged mitochondria, functional

mitochondria can be removed by mitophagy to meet distinct cellular demands. Multiple

mechanisms exist for mitophagy, including both soluble and transmembrane receptor-mediated

mechanisms (Onishi et al., 2021). In soluble receptor-mediated mitophagy, the PINK1/Parkin

axis labels mitochondria with ubiquitin. The ubiquitinated mitochondria are then recognized by

NDP52 and Optineurin, soluble cargo receptors, leading to their sequestration by autophagic

membranes (Narendra and Youle, 2024). On the other hand, transmembrane cargo receptors are

anchored to the outer mitochondrial membrane and recruit autophagy machinery via LIR and

other motifs interacting with the ATG machinery (Onishi et al., 2021). The concept of receptor-

mediated mitophagy was first established in yeast ATG32 studies (Kanki et al., 2009; Okamoto

et al., 2009), and then subsequently it was found that the basic mechanisms are also conserved

in multicellular organisms. The list of mitochondrial outer membrane-embedded cargo receptors

is growing: NIX/BNIP3L, BNIP3, FKBP8, FUNDC1, and BCL2L13 (Onishi et al., 2021).

Nevertheless, their relative contribution and molecular mechanisms still need to be fully

elucidated, especially under physiological conditions. So far, most mitophagy studies have been

performed in cultured cells with uncouplers, hypoxia, or genetic perturbations (Yamashita et al.,

2024).

NIX (also known as BNIP3L) and BNIP3 are homologous receptors for mitophagy in

mammals. NIX was initially discovered as a mitophagy receptor during reticulocyte maturation.

Seminal studies discovered that NIX is required to clear mitochondria from reticulocytes during

differentiation (Sandoval et al., 2008; Schweers et al., 2007). Following this, its paralog BNIP3

was also shown to regulate mitophagy. NIX and BNIP3 have an LIR motif near their N-

terminus and a transmembrane domain (TMD) for localization to the mitochondrial outer

membrane near their C-terminus. Thereby, it has been proposed that NIX and BINP3 tether the

Atg8/LC3-positive isolation membrane and mitochondria for mitophagy through these domains

(Li et al., 2022; Onishi et al., 2021; Schwarten et al., 2009). In addition, Prof. Ney's group in the

US identified a minimal essential region (MER), a short motif in NIX required for mitochondria

clearance in reticulocytes (Zhang et al., 2012). Recently, it has been reported that the MER

directly interacts with WIPIs, mammalian Atg18 orthologs, to recruit core autophagy machinery

in cultured cells overexpressing BNIP3 (Adriaenssens et al., 2024; Bunker et al., 2023). Despite

these findings, the roles of LIR-Atg8 and MER-Atg18 interactions in BNIP3-mediated

mitophagy have not been investigated under physiological conditions.

Drosophila BNIP3, the sole ortholog of NIX and BNIP3, contains LIR, MER, and

TMD, similar to its mammalian counterparts. In *Drosophila*, it has been reported that BNIP3

plays a role in programmed germline mitophagy required for mitochondrial DNA quality

control (Lieber et al., 2019; Palozzi et al., 2022). However, the mechanism is not well

understood. In addition, the contribution of BNIP3 to mitophagy in other physiological contexts,

such as cellular differentiation or remodeling programs, has not been explored yet. In this study,

we performed a comparative time-course RNA-seq analysis of isolated muscle cells, and

identified BNIP3-mediated mitophagy as a critical factor in developmentally programmed

muscle remodeling. We also explored the underlying mechanism by which BNIP3 tethers

mitochondria with the autophagic machinery, as well as the physiological significance of

BNIP3-mediated mitophagy *in vivo*.

Results

Transcriptional dynamics of DIOM remodeling during metamorphosis

During DIOM remodeling, autophagy occurs during atrophy, which lasts from 12 h to 2 d APF

(after puparium formation). Then, organelle and myofibril reformation accompanies

hypertrophy from 2 to 4 d APF (Fig. 1A). We previously reported the dynamics of several

organelles in DIOM remodeling (Fujita et al., 2017; Murakawa et al., 2020); however, the

overall mechanism is still enigmatic. To gain molecular insights into DIOM remodeling, we

performed time-course RNA-seq of isolated DIOMs from four animals at five time points (Fig.

1A). Dissected animals were fixed with methanol to avoid cell death induction (Wang et al.,

2021), and six DIOMs were isolated from each animal to prep total RNA (Fig. 1B). The

mRNAs were amplified, and the cDNA libraries were prepared using CEL-Seq2, a sensitive

protocol for single-cell RNA-seq (Hashimshony et al., 2016). Then, they are subjected to next-

generation sequencing. As a result, ~8,000 genes were counted in the DIOMs above background

(Supplementary File 1, ctrl). High similarities among four replicates at each time point show the

reproducibility of our protocol (Fig. 1-figure supplement 1A). We performed DESeq2 principal

component analysis (PCA) of all samples to reduce the dimensionality of the data (Fig. 1C and

Fig. 1-figure supplement 1B). In the PC1-PC2 plane, which explains 43% of expressed genes in

DIOMs, the transcriptome changed dynamically from 3IL to 4 d APF (Fig. 1C). Of note, there

was a substantial difference between before (3IL) and after (4 d APF) remodeling in both PC1-2

and PC3-4 (Fig. 1C), suggesting the functional importance of remodeling larval muscle to adult

abdominal muscle during this epoch.

To identify genes with similar temporal expression profiles, we used Fuzzy c-means to

categorize normalized read counts into twelve clusters (Fig. 1-figure supplement 1C), each with

379 to 937 genes (Fig. 1-figure supplement 1D). The twelve clusters represent unique temporal

expression dynamics, with each time point showing a distinct profile of clusters with high

expression (Fig. 1D and Fig. 1-figure supplement 1D). We also noticed that a time-dependent

shift in gene ontology (GO) terms is apparent among the clusters, reflecting the sequential

events during DIOM remodeling (Fig. 1F). The principal component coefficients (loadings) of

individual clusters to PC1 through 4 are shown in Fig. 1E. The direction and length of the

arrows indicate the contribution of each cluster to principal components (PC1 to 4). Briefly,

upon induction of muscle atrophy (1-2 d APF), the ubiquitin-proteasome system, developmental

signaling, autophagy, and histone modifications-related genes were upregulated. Then,

chromatin remodeling-related genes were elevated around 2 d APF. Myofibril, mitochondrion,

and ribosome biogenesis-related genes required for muscle hypertrophy were upregulated

around 3 d APF (Fig. 1F). The data illustrates the sequential events involved in DIOM

remodeling, aligning with previously observed morphological changes (Fujita et al., 2017;

Murakawa et al., 2020).

Transcriptional dynamics occur independently of autophagy during DIOM remodeling

Autophagy deficiency impacts both atrophy and myofibril regeneration during hypertrophy

(Fujita et al., 2017; Murakawa et al., 2020) (Fig. 2A). Given that autophagy is known to

influence gene expression through both direct and indirect mechanisms (Morishita and

Mizushima, 2019; Sánchez-Martín et al., 2019), we hypothesized that the loss of autophagy also

alters gene expression in DIOMs. To investigate this possibility, we performed a comparative

time-course RNA-seq of DIOM remodeling in *Atg18a* RNAi, *FIP200* RNAi, or *Stx17* RNAi

DIOMs and compared the results with control data (Fig. 2B). Atg18a and FIP200 are essential

factors for autophagosome formation, and Stx17 is an autophagosomal SNARE protein required

for autophagosome-lysosome fusion (Lőrincz and Juhász, 2020; Nakatogawa, 2020). Thus, their

knockdown blocks autophagy at early or late stages, respectively (Fig. 2B, bottom). Similar to

the experiment in Fig. 1, we performed RNA-seq at five time points from 3IL to 4 d APF (Fig.

2B) and confirmed high similarities among the four replicates in all 20 conditions (Fig. 2-figure

supplement 1A and Supplementary File 1). We confirmed that RNAi efficiently knocked down

targeted genes (Fig. 2-figure supplement 1B-D). In contrast to our prediction, the knockdown of

Atg18a, *FIP200*, or *Stx17* only had a slight impact on transcriptomic dynamics in DIOM

remodeling (Fig. 2C), with only minor changes detected (Fig. 2-figure supplement 2G).

Next, to examine whether loss of autophagy affects protein synthesis in DIOMs, we

used the quantification of DIOM volume changes as a surrogate measurement, and compared

controls with *FIP200* RNAi from 3IL to 4 d APF (Fig. 2D and E). As expected, muscle atrophy

observed until 2 d APF was affected by *FIP200* RNAi (Fig. 2A and D). Notably, DIOMs in the

FIP200 RNAi condition grew from 2 to 3 d APF at levels comparable to the control (Fig. 2E),

indicating that protein synthesis occurs independently of autophagy. Consistent with this

observation, transcription of both ribosomal RNAs and proteins, established indicators of the

target of rapamycin complex 1 (TORC1) activity (Saxton and Sabatini, 2017), did not drop but

instead were slightly elevated in conditions lacking autophagy (Fig. 2-figure supplement 2). The

synthesis of rRNA was upregulated in *ATG* RNAi between 2 to 3 d APF (Fig. 2-figure

supplement 2A). It was confirmed that the number of nuclei remained unchanged by the loss of

autophagy (Fig. 2-figure supplement 2B-C). The size of the nucleolus, which positively

correlates with TORC1 activity (Grewal et al., 2007), was increased by *FIP200* and *Stx17* RNAi

(Fig. 2-figure supplement 2D-F). Moreover, most ribosomal protein large subunit (RpLs) and

ribosomal protein small subunit (RpSs) mRNA levels were higher in *ATG* RNAi conditions

(Fig. 2-figure supplement 2G). These results suggest that TORC1 activity and protein synthesis

capacity are comparable between control and autophagy-deficient DIOMs. Altogether, we

conclude that gene expression occurs independently of autophagy in DIOMs.

Loss of BNIP3 leads to an accumulation of mitochondria in DIOMs

The above results suggest that the autophagic degradation of cytosolic components in itself is

critical for DIOM remodeling. We have previously reported that *FIP200* RNAi or *Stx17* RNAi

induce the accumulation of mitochondria or mitophagosomes, respectively (Fig. 3A and B)

(Fujita et al., 2017). In addition, mitophagosomes were often observed in wild-type DIOMs at 1

d APF, when autophagy is strongly induced (Fig. 3C) (Fujita et al., 2017). These results suggest

that mitochondria are major autophagy cargo, and impaired mitophagy contributes to the severe

loss of autophagy phenotype.

We tried to identify relevant mitophagy receptors directly via our RNA-seq approach.

The time course RNA-seq data (Fig. 1 and 2) indicated that *BNIP3* was robustly expressed in 1

d APF DIOMs among known mitophagy regulators (Fig. 3D). *Zonda*, *CG12511*, *Key*, and

Ref(2)P are *Drosophila* orthologs of FKBP8, FUNDC1, Optineurin, and p62, respectively.

BNIP3 is the sole fly ortholog of mammalian NIX and BNIP3. *BNIP3* mRNA levels were

relatively high at 3IL and 4 d APF when mitophagy was minimally induced. We noticed that

FBXL4 is expressed in 3IL and 4 d APF but not in 1 d APF DIOM (Fig. 3E). BNIP3 is known to

undergo ubiquitination and degradation through a mechanism mediated by FBXL4, a ubiquitin

E3 ligase (Niemi and Friedman, 2024). Previous data suggest that transcriptional and post-

translational mechanisms regulate BNIP3 protein levels. Consistent with expression levels,

BNIP3 RNAi led to mitochondrial accumulation in 4 d APF DIOMs, whereas RNAi targeting

other known mitophagy regulators did not (Fig. 3-figure supplement 1). We generated *BNIP3*

knockout (KO) flies lacking all exons using two gRNA to characterize BNIP3 further (Fig. 3F).

BNIP3 KO was compatible with viability and did not markedly impair mobility (Fig. 3-figure

supplement 2A-C). Similar to RNAi, mitochondria accumulated significantly in *BNIP3* KO

DIOMs at 4 d APF (Fig. 3G).

At the ultrastructural level, *BNIP3* KO induced the accumulation of mitochondria and

loss of myofibrils (Fig. 4A-C), consistent with our confocal data (Fig. 3G). To test the role of

BNIP3 in mitophagosome formation more directly, we performed TEM in the absence of Stx17

to block the autophagosome-lysosome fusion. As expected, mitophagosomes (pink-colored)

were frequently observed in the *Stx17* RNAi condition (Fig. 4D, D' and F). In stark contrast,

mitophagosomes were rare, and unengulfed mitochondria (green-colored) accumulated in the

combined condition of *Stx17* RNAi and *BNIP3* KO (Fig. 4E, E', and F). We also confirmed

there was no significant difference in the number of autophagosomes (blue-colored) between the

two conditions (Fig. 4G), indicating that BNIP3 is dispensable for autophagosome formation.

These data show that BNIP3 is critical for mitophagosome formation in DIOMs.

The LIR and MER motifs in BNIP3 are required for the clearance of mitochondria in

DIOMs

Drosophila BNIP3 harbors LIR, MER, and TMD domains, similar to mammalian NIX and

BNIP3 (Fig. 5A). Although the function of the MER domain has been unknown since it was

identified (Zhang et al., 2012), recent studies have shown that the MER of mammalian NIX and

BNIP3 directly interacts with WIPIs, the mammalian orthologs of Atg18 (Adriaenssens et al.,

2024; Bunker et al., 2023). AlphaFold 3 (Abramson et al., 2024) predicted that this interaction

is evolutionarily conserved in *Drosophila* in a manner analogous to their mammalian

counterparts (Fig. 5B) (Adriaenssens et al., 2024; Bunker et al., 2023). In this prediction,

residues 31–57 of *Drosophila* BNIP3 (slate blue and yellow) interact with the surface formed by

blades 2 (green) and 3 (pink) of the β-propeller structure of *Drosophila* Atg18a. Notably,

residues 42–53 (yellow), which correspond to the MER designated for NIX (Bunker et al.,

2023; Zhang et al., 2012), fold into an α-helix, docking into the groove formed by blades 2 and

3 and forming substantial hydrophobic interactions as well as defined polar contacts. We

confirmed that HA-tagged *Drosophila* Atg18a co-immunoprecipitated with GFP-tagged full-

length *Drosophila* BNIP3, and that this interaction was attenuated by the deletion of the MER

(residues 42-53) (Fig. 5C).

Next, to verify the importance of the LIR and MER motifs in BNIP3, we expressed a

series of BNIP3 mutants in *BNIP3* KO flies (Fig. 5A). ΔLIR (W16A/L19A) lacks critical

consensus residues in LIR due to double mutations (Johansen and Lamark, 2020). The MER

mutant (MER^{mut}, L49A) substituted residue L49 with an A, which is predicted to be deeply

buried at the interface with Atg18a (Fig. 5C). The corresponding L75A mutant in human NIX

has been shown to lose its affinity for WIPI2/Atg18 (Bunker et al., 2023). The ΔMER mutant

lacks an entire short α-helix (G42 to Q53) of the MER motif (Fig. 5C, shown yellow-orange).

The ΔLIR+ΔMER is a combined construct of ΔLIR and ΔMER. All BNIP3 constructs retain an

intact TMD at the C-terminus, ensuring their localization to the outer mitochondrial membrane

(Fig. 5-figure supplement 1A).

As shown in Fig. 5D, the expression of the full-length (Full) construct almost

completely suppressed the accumulation of mitochondria in BNIP3 KO at 4 d APF, confirming

the rescue of BNIP3 function using our construct-based approach. To our surprise, the BNIP3

ΔLIR construct rescued the *BNIP3* KO phenotype comparable to Full, indicating LIR on its own

may be redundant for mitophagy in DIOMs. In contrast, mitochondria accumulated in MER^{mut}

and ΔMER reconstituted DIOMs. Furthermore, the combination of ΔLIR and ΔMER resulted in

mitochondria accumulation nearly identical to that observed in BNIP3 KO flies (Fig. 5D and

D'). Consistent with these findings, *Atg101*-dependent autophagic degradation of GFP-BNIP3

was strongly suppressed by the ΔLIR+ΔMER mutations (Fig. 5E). Atg101 is an essential

subunit of the Atg1/ULK1 kinase complex (Guo et al., 2019; Hegedus et al., 2014; Noda and

Mizushima, 2016). From these results, we conclude that both MER and LIR motifs contribute to

BNIP3-mediated mitophagy, with some redundancy in their contributions to selectivity.

BNIP3-mediated mitophagy eliminates larval muscle mitochondria during muscle

remodeling

The data above suggest that a BNIP3-mediated mechanism degrades larval muscle-derived

mitochondria; however, in previous figures, we only observed the terminal phenotype of *BNIP3*

KO at 4 d APF. To observe the changes in the number of mitochondria and cell shape during

DIOM remodeling, we first performed a time-course experiment in control and *BNIP3* KO

conditions (Fig. 6A and Fig. 6-figure supplement 1A and B). There was no significant

difference in the number of mitochondria at 3IL; however, mitochondria started accumulating at

1 d APF (Fig. 6A and A'), suggesting that BNIP3-mediated mitophagy degrades larval muscle

mitochondria. Further, we tested the Mito-QC probe (Lee et al., 2018), a tandem GFP-mCherry

fusion protein targeted to the OMM, to compare mitophagy flux at 1d APF with and without

BNIP3. When the probe is delivered to lysosomes via autophagy, the GFP is quenched due to

the acidic environment of the lysosome. Robust mitophagy activity was detected in control

DIOMs at 1 d APF. In sharp contrast, *BNIP3* KO blocked mitophagy flux at a comparable level

with *FIP200* RNAi (Fig. 6B and B').

To show the degradation of larval muscle-derived mitochondria more directly, we

labeled larval muscle mitochondria specifically by using the temperature-sensitive mutant of

GAL80 (GAL80^{ts}), a repressor of GAL4 (Mcguire et al., 2004). At the restrictive temperature

(29˚C), the GAL80^{ts} mutant is inactive, allowing GAL4 to drive the expression of the UAS-

fused construct, UAS-Mito-GFP. Conversely, at the permissive temperature (18˚C), the

GAL80^{ts} is active, suppressing GAL4 activity. Using this system, Mito-GFP was expressed in

the larval muscle and subsequently suppressed throughout the entire pupal stage (Fig. 6C). Anti-

ATP5A antibodies were used to visualize total mitochondria. The amount of Mito-GFP-positive

mitochondria was comparable in control and *BNIP3* KO at 3IL muscle. However, at 4 d APF,

the Mito-GFP signal was nearly absent in the control, while a strong Mito-GFP signal was

observed in the *BNIP3* KO (Fig. 6D and D'), indicating that BNIP3-mediated mitophagy

degrades larval muscle mitochondria during muscle remodeling.

Discussion

The transcriptional dynamics associated with DIOM remodeling are largely independent of autophagy (Fig.2). Instead, our RNA-seq data suggest that it is regulated primarily by ecdysone signaling, with minimal influence from autophagy inhibition. However, a subset of genes exhibited altered expression levels in autophagy-deficient conditions (Fig. 2-figure supplement 2G). Of note, their expression levels were consistently upregulated or downregulated throughout all time points. Since no apparent phenotype was observed in *ATG* knockdown 3IL body wall muscles (Fujita et al., 2017), the observed expression changes are unlikely to underlie the loss-of-autophagy phenotype in DIOM remodeling. Furthermore, *FIP200* RNAi had minimal impact on the DIOM growth rate (Fig. 2 D and E), which reflects translational capacity. From these data, we conclude that the contribution of autophagy to gene expression during DIOM remodeling is minimal.

In the RNA-seq experiments (Fig. 2), *ATG* genes were exclusively depleted in muscle cells; thus, autophagy in non-muscle organs remained intact. It is plausible that essential nutrients for DIOM hypertrophy, such as amino acids, were supplied by other organs, including the fat body (Murakawa et al., 2022). Alternatively, the necessary amino acids might have been derived from myofibril breakdown via the ubiquitin-proteasome system (Quy et al., 2013).

Transcriptome changes before and after DIOM remodeling reflect adaptations to the shift from the anaerobic environment of larvae to the aerobic environment of adults, alongside alterations in nutritional pathways. Notable changes in energy production systems occurred between the 3IL stage (pre-remodeling) and 4 d APF (post-remodeling) (Fig. 1). Glycolysis-associated genes exhibited high expression levels at the 3IL stage, followed by a significant decline during the pupal and adult stages. For instance, the expression of *Lactate dehydrogenase* (*Ldh*), a key enzyme in switching between anaerobic and aerobic respiration (Liang et al., 2023; Semenza, 2014), was particularly noteworthy. Ldh catalyzes the conversion of pyruvate to lactate while regenerating NAD^+ from NADH under anaerobic conditions. Ldh levels were markedly high in larval muscles, suggesting reliance on anaerobic glycolysis. Conversely, its reduced expression in remodeled adult muscles reflects a metabolic shift toward aerobic pathways.

Genes related to serine-centered amino acid and lipid catabolism, both of which rely on oxygen to produce ATP, were significantly upregulated in adult muscles. For example, the

expression of *Alanine-glyoxylate aminotransferase* (*Agxt*), an enzyme that converts alanine to

pyruvate, was approximately 100 times higher in adult muscles compared to larvae. Similarly,

brummer (*bmm*), a triacylglycerol lipase, exhibited a fivefold increase, suggesting enhanced

reliance on fatty acids metabolized via β-oxidation and the TCA cycle. These findings highlight

the shift in energy sources from glycolysis in larval muscles to amino acids and lipids in adult

muscles, metabolized aerobically in mitochondria.

A recent study suggests that DIOMs are a primary source of pupal ecdysone (Zhang et

al., 2024). In line with this data, our time-course mRNA-seq analysis revealed high expression

of *Phm*, an enzyme essential for ecdysteroid biosynthesis, at 1 d APF (Supplementary File 1).

However, other Halloween genes required for ecdysteroid biosynthesis, such as *Nvd*, *Spo*, *Spok*,

Dib, *Sad*, and *Shd* (Kamiyama and Niwa, 2022), were not, or only minimally, expressed in

DIOMs during metamorphosis (Supplementary File 1). Consequently, whether DIOMs directly

secrete ecdysone remains unclear. Instead, DIOMs may contribute to a specific step in the

ecdysteroid biosynthesis pathway, potentially supplying intermediate 5β-ketotriol to other

tissues for further conversion.

Functional mitochondria are degraded during developmental muscle remodeling;

therefore, it is reasonable that a membrane-anchored mitophagy receptor, directly interacting

with the autophagic machinery, is selected over a ubiquitination-mediated mechanism involving

PINK1 and Parkin. The PINK1/Parkin axis is activated by mitochondrial membrane potential

depolarization (Narendra and Youle, 2024), which is not observed in remodeling muscles. The

protein level of BNIP3 appears to be regulated at transcriptional and post-translational levels

(Fig. 3E). It is known that BNIP3 is degraded via a mechanism mediated by Fbxl4, a ubiquitin

E3 ligase (Niemi and Friedman, 2024). The balance between synthesis and degradation likely

results in the highest BNIP3 protein levels at 1 d APF, a time when mitophagy is robustly

induced. However, BNIP3 protein levels have not been directly confirmed, as isolating a

sufficient number of DIOMs for western blotting is challenging. Generating a GFP knock-in

line for BNIP3 would allow for an imaging-based assay to measure its protein levels during

DIOM remodeling.

Overexpression of BNIP3 in larval muscle cells did not significantly induce mitophagy

(Fig. 5-figure supplement 2), suggesting that BNIP3 expression alone is insufficient to drive

mitophagy. BNIP3-mediated mitophagy likely requires not only BNIP3 expression but also

autophagy induction. We propose that autophagy induction in DIOMs at the early pupal stage is

regulated by ecdysone signaling, similar to other instances of developmental autophagy in

Drosophila (Murakawa et al., 2022). Understanding the mechanisms by which developmental

signals trigger autophagy remains a critical question for future research.

A key function of autophagy in DIOM remodeling is the degradation of mitochondria

from larval muscles. In the absence of BNIP3, mitochondria derived from the larval muscle

accumulate and cluster in the cell center, physically obstructing myofibril formation during

hypertrophy and restricting myofibrils to the cell periphery (Fig. 6E). Interestingly, unlike *ATG*

RNAi, *BNIP3* knockout muscles exhibited relatively normal but thinner myofibrils (Fig. 3G). In

contrast, *ATG* RNAi not only caused mitochondrial accumulation but also severely disrupted

myofibril organization, as indicated by misaligned sarcomeres (Fujita et al., 2017; Murakawa et

al., 2020). These findings suggest that autophagy contributes to DIOM remodeling beyond

mitochondrial degradation. Schnorrer's group in France has demonstrated that mechanical

tension is essential for sarcomere assembly and maturation (Mao et al., 2022; X. Zhang et al.,

2024). Given that *ATG* RNAi disrupts DIOM attachment (Murakawa et al., 2020; Ribeiro et al.,

2011), it is possible that reduced tension—caused by an unknown mechanism—leads to

disorganization of myofibrils in DIOMs when autophagy is impaired.

Contrary to the previous model, the LIR motif of BNIP3 was found to be dispensable

for mitophagy in DIOMs (Fig. 5D and E). Our findings align with reports stating that the

clearance of mitochondria in reticulocytes is independent of the LIR motif (Novak et al., 2010;

Zhang et al., 2012). Together, this suggests that the dependency on the LIR motif may differ

between *in vitro* and *in vivo* contexts. In studies of BNIP3 in cultured cells, such as HeLa cells,

BNIP3 is often overexpressed, and mitophagy is induced through chemical treatments (Bunker

et al., 2023; Yamashita et al., 2024). These experimental conditions may account for the

observed differences in LIR dependency. Understanding the factors that drive these contextual

variations in BNIP3-mediated mitophagy will require further investigation.

The MER motif in BNIP3 is required for mitophagy in *Drosophila*. Similar to its

mammalian counterpart, the MER motif of *Drosophila* BNIP3 is predicted to interact with the

groove between blades 2 and 3 of the β-propeller in Atg18a (Fig. 5B). Notably, this groove of

Atg18 overlaps with the binding site for Atg16 (Strong et al., 2021). We propose that the BNIP3

MER-Atg18a interaction functions at the early stages of mitophagosome formation, which is

later replaced by the Atg16-Atg18a interaction during the elongation phase. In this sequential

model, the BNIP3 LIR-Atg8a interaction would dominate at the elongation steps. Alternatively,

the MER-Atg18a and LIR-Atg8a interactions might serve redundant roles in determining

selectivity, as both Atg8 and Atg18 localize to the elongating autophagic membranes. This

redundancy could explain why the LIR motif is dispensable for BNIP3-mediated mitophagy

(Fig. 5D and E).

Martens's group (Austria) recently reported that WIPIs, the mammalian orthologs of

Atg18, bind to Atg13, a component of the FIP200/ULK1 complex (Adriaenssens et al., 2024).

They propose that the NIX/BNIP3-WIPI-Atg13 axis induces selective autophagy for

mitochondria, analogous to known selective autophagy receptors associated with FIP200

(Lamark and Johansen, 2021). It would be interesting to test whether an artificial interaction

motif with the Atg1 complex could compensate for the loss of function of ΔMER in DIOMs.

Our study provides new insights into the mechanisms underlying autophagy-

associated muscle remodeling. Transcriptional dynamics uncovered a sequence of events

involving muscle atrophy and hypertrophy, offering a valuable dataset for understanding the

process of muscle remodeling. In addition, we provided insights into the mechanism and

significance of BNIP3-mediated mitophagy *in vivo*. These findings are particularly important

given the evolutionary conservation of muscle remodeling mechanisms between *Drosophila* and

mammals (Piccirillo et al., 2014). Future studies should explore the broader implications of our

findings across various cellular, developmental, and organismal contexts.

Materials and methods

Reagents and antibodies

The following antibodies were used: Mouse monoclonal anti-ATP5A (1:300, ab14748, Abcam, Cambridge, UK), Rat monoclonal anti-HA (1:1000; 11867423001, Roche, Basel, Switzerland), Rabbit polyclonal anti-GFP (1:2000; 598, MBL, Nagoya, Japan), Rabbit polyclonal anti-Fibrillarin (1:300; ab5821, Abcam, Cambridge, UK), anti-rabbit IgG Alexa Fluor 594 conjugate (1:400; A11012, Thermo Fisher Scientific, Waltham, USA), anti-mouse IgG Alexa Fluor 488 conjugate (1:400; A11001, Thermo Fisher Scientific, Waltham, USA), anti-mouse IgG Alexa Fluor 594 conjugate (1:400; A11005, Thermo Fisher Scientific, Waltham, USA), HRP-conjugated AffiniPure Goat Anti-Rabbit IgG (1:10000; 111-035-144, Jackson ImmunoResearch, West Grobe, USA), HRP-conjugated AffiniPure Donkey Anti-Rat IgG (1:5000; 712-005-153, Jackson ImmunoResearch, West Grobe, USA), Bisbenzimide H33342 Fluorochrome Trihydrochloride DMSO Solution (1:10000; 04915-81, Nacalai Tesque, Kyoto, Japan), Alexa Fluor 633 Phalloidin (1:200; A22284, Thermo Fisher Scientific, Waltham, USA), and GFP-Trap Agarose (gta, Chromotek, Planegg-Martinsried, Germany).

Drosophila strains

Flies were reared under standard conditions at 25°C unless otherwise stated. The w^{1118} strain was used as a control for knockout strains. For muscle-targeted gene expression, Mef2-GAL4 was used. UAS-*LacZ* was used as a control for UAS transgenes. All genetic combinations were generated by standard crosses. Detailed genotypes are described in Supplementary Table 1. Genotypes of flies used in this study include the following: (1) y^1 w*; $P\{w^{+mC}$=GAL4-*Mef2.R*}3 (Bloomington *Drosophila* Stock Center, BL 27390; DMef2-GAL4), (2) w^{1118}; $P\{w^{+mC}$=UAS-lacZ.B}*Bg4-1-2* (BL 1776; UAS-LacZ), (3) w*; $P\{w^{+mC}$=UAS-Rheb.Pa}2 (BL 9688), (4) w^{1118}; $P\{w^{+mC}$=UAS-mito-HA-GFP.AP}2/CyO (BL 8442; Mito-GFP), (5) w^{1118}; $P\{y^{+t7.7}$ w^{+mC}=UAS-mito-QC}attP16 (BL 91640, Mito-QC), (6) w^{1118}; $P\{w^{+mC}$=UAS-Dcr-2.D}10 (BL 24651, UAS-Dcr2), (7) y^* w*; $P\{w^{+mC}$=UAS-tdTomato.mito}2 (Kyoto DGGR 117016, tdTomato-Mito), (8) w; UAS-IR-Atg18a^{KK100064} (VDRC 105366; Atg18a RNAi), (9) w; UAS-IR-Stx17^{KK100034} (VDRC 108825; Stx17 RNAi), (10) w; UAS-IR-BNIP3^{KK111246} (VDRC 107493; BNIP3 RNAi), (11) w; UAS-IR-Pink1^{KK101205} (VDRC 109614; Pink1 RNAi), (12) w; UAS-IR-Park^{KK107919} (VDRC 104363; Park RNAi), (13) w; UAS-IR-CG12511^{KK109618} (VDRC

101785; CG12511 RNAi), (14) *w*; *UAS-IR-Zonda^{KK107775}* (VDRC 110620; Zonda RNAi), (15) *y^1*

sc* v^1 sev^{21}; *P{y$^{+t7.7}$ v$^{+t1.8}$=TRiP.HMS01611}attP2/TM3, Sb^1* (TRiP, BL 36918; FIP200 RNAi),

(16) *y^1 v^1*; *P{y$^{+t7.7}$ v$^{+t1.8}$=TRiP.JF01937}attP2* (TRiP, BL 25896; Stx17 RNAi), (17) *y^1 sc* v^1*

sev^{21}; P{y$^{+t7.7}$ v$^{+t1.8}$=TRiP.GL00156}attP2 (TRiP, BL 35578; mTor RNAi), (18) *w*; *UAS-tub-*

GAL80^{ts} (from E. Kuranaga), (19) *w*; *UAS-mCD8:GFP*, (20) *w; DMef2-GAL4, UAS-Dcr2*

(Fujita et al., 2017), (21) *w*; *UAS-Dcr2*; *DMef2-GAL4, sqh-YFP:Mito^{BL7194}/TM6C Sb Tb*

(Murakawa et al., 2020). New genotypes generated during this study include the following: (22)

w*;; BNIP3 KO CRISPR{3xP3-RFP}/TM6B, Tb^1, (23) *w^{1118}, Atg101 KO CRISPR{3xP3-*

RFP}/FM7a (24) *w^{1118}*; *UASt-GFP-BNIP3_full^{attP40}*, (25) *w^{1118}*; *UASt-GFP-BNIP3_ΔLIR*

(W16A/L19A)^{attP40}, (26) *w^{1118}*; *UASt-GFP-BNIP3_MER mutant (L49A)^{attP40}*, (27) *w^{1118}*; *UASt-*

GFP-BNIP3_ΔMER (Δ42-53) ^{attP40} and (28) *w^{1118}*; *UASt-GFP-BNIP3_ ΔLIR+ΔMER^{attP40}*.

DNA engineering

Plasmid vectors were constructed using standard molecular biology techniques. The DNA

polymerase KOD One (TOYOBO, Osaka, Japan) and PrimeSTAR GXL Premix (Takara Bio,

Shiga, Japan) were used for PCR amplification. The DNA fragments were assembled using the

Gibson Assembly. Construction of the expression vectors of GFP-tagged BNIP3_full under

UAS control (pUASt-attB-GFP-BNIP3) was as follows: *Drosophila* BNIP3 CDS was amplified

from template RE48077 (DGRC Stock 9148; https://dgrc.bio.indiana.edu//stock/9148 ;

RRID:DGRC_9148) using the following primer sets: 5'-

ATGTCTACGACACCAAAATCGAG-3' and 5'-TCAGTCAATGACCACACGG-3'. This

fragment of BNIP3 CDS was ligated into the linearized pUAST-attB-GFP vector, which is

amplified with the following primer sets: 5'-

CGTGTGGTCATTGACTGAGCGGCCGCGGCTCGAGGGTACC-3' and 5'-

TTGGTGTCGTAGACATGGTGAAGGGGGCGGCCGCGGAG-3'. To generate plasmids with

mutations or deletions in the BNIP3 coding sequence (pUAST-attB-BNIP3_ΔLIR, pUAST-

attB-BNIP3_MER$^{mut (L49A)}$, and pUAST-attB-BNIP3_ΔMER), the plasmid pUAST-attB-GFP-

BNIP3_full was used as a template. For the construction of pUAST-attB-BNIP3_ΔLIR, these

primer sets: 5'- CTGCGATCGAAGCGAGCACAACAGCTGCGATG-3' and 5'-

CTCGCTTCGATCGCAGATTCGCCCAGCAAATC-3' were used to introduce the

W16A/L19A mutations. For the construction of pUAST-attB-BNIP3 MER$^{mut (L49A)}$, these primer

sets: 5'-AGACTTGCGCGCGAGGCCCAGCGCGAG-3' and 5'-

CTCGCGCGCAAGTCTCAGGTACTCCTC-3' were used to introduce the L49A mutation. For

the construction of pUAST-attB-BNIP3_ΔMER, these primer sets: 5'-

CAACAATCGCGAGTCGAACCAGTCG-3' and 5'-

GACTCGCGATTGTTGAATGGCAACGG-3' were used to delete G42 to Q53. To generate a

construct that harbors both ΔLIR and ΔMER (pUAST-attB-BNIP3_ΔLIR+ΔMER), the plasmid

pUAST-attB-GFP-BNIP3_ΔMER was used as a template. The same primer sets as pUAST-

attB-BNIP3_ ΔLIR construction were used for amplification. Each resultant linear DNA

fragment was ligated into a circular plasmid. Generation of the mammalian expression

constructs—pMRX-IRES-puro-GFP-BNIP3_full, pMRX-IRES-puro-GFP-BNIP3_ΔMER, and

pMRX-IRES-puro-3xHA-mCh-Atg18a—was performed by amplifying the corresponding

inserts from pUAST-attB-GFP-BNIP3_full, pUAST-attB-GFP-BNIP3_ΔMER, or LD38705

(DGRC Stock 2722; https://dgrc.bio.indiana.edu//stock/2722 ; RRID:DGRC_2722),

respectively. The amplified fragments were subsequently ligated into the linearized pMRX-

IRES-puro vector. All the resultant vectors were validated by DNA sequencing.

Generation of mutant and transgenic flies

To generate *Drosophila* mutants with a knockout of *BNIP3* (CG5059) or *Atg101* (CG7053), the

CRISPR/Cas9 system was used, following a modified method based on Kondo and Ueda

(Kondo and Ueda, 2013). Guide RNA (gRNA) sets for *BNIP3* KO and *Atg101* KO were

designed as follows: for *BNIP3* KO, upstream gRNA sequences, 5'-

CGTCAACACCAAAGATAACT[TGG]-3' and downstream gRNA sequences 5'-

CTTTCAGTCAATGACCACAC[GGG]-3'; for *Atg101 KO*, upstream gRNA sequences 5'-

GTCCACCTGACGACCCTCCA[TGG]-3' and downstream gRNA sequences 5'-

CTCGCAATGTGACGGGCTGT[CGG]-3'. Each gRNA was individually cloned into a plasmid

under the control of U6 promoter. A donor DNA cassette containing a 3x P3-RFP selection

marker, loxP sites, and two homology arms was cloned into the pUC57-Kan plasmid for DNA

repair. The gRNAs targeting *BNIP3* or *Atg101,* and the hs-Cas9 coding DNA plasmids, along

with the donor plasmid, were microinjected into embryos of the w^{1118} strain. F1 files were

screened by a selection marker of 3x P3-RFP and validated by genomic PCR and sequencing

(WellGenetics Inc.). To generate a series of UASt-GFP-BNIP3 transgenic flies, the vectors

described above were injected into embryos for phiC31-mediated insertion (WellGenetics Inc.).

RNA-seq analysis of DIOMs

For collecting DIOMs or their precursors, larvae or pupae were dissected and fixed with

methanol. Six DIOMs were isolated from each animal using forceps and a pipette under a

stereomicroscope (SZX16, EVIDENT, Tokyo, Japan). The collected muscle cells were lysed

with BLA buffer (ReliaPrep Tissue RNA Miniprep System, Promega) and stored at -80˚C.

Following sample preparation, the frozen samples were thawed, and total RNA was extracted

and eluted in RNase-free water according to the manufacturer's protocol (ReliaPrep Tissue

RNA Miniprep System, Promega). Subsequently, cDNA libraries were prepared using CEL-

Seq2, a single-cell RNA-seq protocol (Hashimshony et al., 2016). Briefly, reverse transcription

was performed using poly(T) primers with a T7 promoter to generate single-stranded DNA from

mRNA, which was then converted into double-stranded DNA through second-strand synthesis.

Amplified RNA was produced via in vitro transcription using the MEGAscript T7 kit (Thermo

Fisher Scientific, Waltham, MA) and subsequently reverse transcribed to construct the cDNA

library. Library sequencing was performed using the NovaSeq6000. Cell barcode and UMI in

Read1 were extracted by using UMI-tools with the following command "umi_tools extract -I

read1.fastq —read2-in=read2.fastq --bc-pattern=NNNNNNNNNNCCCCCCCCC". Adaptor

sequence and low-quality sequence were removed, read lengths below 20bp were discarded by

using Trim Galore, and reads were mapped to the GRCm38 reference using HISAT2. Read

counts for each gene were obtained by featureCounts, and UMI duplications were removed. The

expression levels of genes in each sample were determined from the UMI counts after a

normalization process using the DESeq2 package in R. The R programming language was used

for each type of analysis, utilizing the pheatmap package for cluster analysis, the prcomp

package for principal component analysis (PCA), and the Mfuzz package for fuzzy c-means

clustering analysis (Kumar and Futschik, 2007).

Quantification of rRNA in DIOMs

Total RNA was extracted as described in the "RNA-seq analysis of DIOMs" section. The

quality and composition of the extracted RNA were assessed using the Agilent 2100

Bioanalyzer system with the RNA 6000 Pico Kit (Agilent, Santa Clara, USA). Ribosomal RNA

(rRNA) content in the DIOMs was quantified by measuring the peak areas corresponding to 18S

and 28S rRNA within the total RNA profile.

Immunofluorescence staining

Muscle preparations were performed as previously described (Ribeiro et al., 2011). 3IL or pupae

were pinned on a Sylgard-covered petri dish in dissection buffer (5 mM HEPES, 128 mM NaCl,

2 mM KCl, 4 mM $MgCl_2$, 36 mM sucrose, pH 7.2). The animals were pinned flat and fixed (4%

PFA, 50 mM EGTA, PBS) at room temp for 20 min. Then, the samples were unpinned and

blocked (0.3% bovine serum albumin, 0.6% Triton X100, PBS) at room temp for 5 min,

incubated with primary antibody overnight at room temperature, washed with PBS, and

incubated with Alexa Fluor 488 or 594-conjugated secondary antibody (Thermo Fisher

Scientific, Waltham, MA) for 2 h at room temperature. The immunostained samples were

washed and mounted in FluorSave reagent (Merck Millipore, Darmstadt, Germany).

Confocal fluorescence microscopy

To image live pupal DIOMs, staged pupae were removed from their pupal cases and mounted

between a slide glass and a cover glass following the protocol described by Zitserman and

Roegiers (Zitserman and Roegiers, 2011). Imaging was performed through the dorsal cuticle.

Both live and fixed samples were observed using a confocal microscope (FV3000, EVIDENT,

Tokyo, Japan) equipped with either a 60× silicone/1.30 NA UPlanSApo or a 4×/0.16 objective

lens (EVIDENT, Tokyo, Japan). FLUOVIEW (EVIDENT, Tokyo, Japan) was used for image

acquisition, and the exported images were processed and analyzed with ImageJ (NIH, Bethesda,

USA).

Transmission electron microscopy

Staged pupae (1 d or 4 d APF) were removed from pupal cases, pinned on a Sylgard-covered

petri dish, and dissected directly in fixative (2% paraformaldehyde, 2.5% glutaraldehyde, 150

mM, 5 mM calcium chloride, sodium cacodylate, pH 7.4). They were then fixed for 2 h at room

temperature followed by overnight at 4˚C. The dissected fillets were washed with 0.1 M

phosphate buffer pH 7.4, post-fixed in 1% OsO_4 buffered with 0.1 M phosphate buffer for 2 h,

dehydrated in a graded series of ethanol, and embedded flat in Epon 812 (TAAB, Aldermaston,

UK). Ultrathin sections (70 nm thick) were collected on copper grids covered with Formvar

(Nisshin EM, Tokyo, Japan), double-stained with uranyl acetate and lead citrate, and then

observed using a JEM-1400Flash transmission electron microscope (JEOL, Tokyo, Japan).

Immunoblots and Immunoprecipitation

For immunoblots, five larval fillets were collected for each sample. The samples were lysed

directly in 1.5x sample loading buffer [100 mM Tris-HCl (pH 6.8), 3% SDS, 15% glycerol, and

0.0015% Bromophenol Blue]. Equal amounts of proteins per sample were subjected to SDS-

PAGE and transferred to Immobilon-PSQ PVDF transfer membrane (Merck Millipore,

Darmstadt, Germany) with Trans-Blot Turbo Transfer System (Bio-Rad, Hercules, CA, USA).

The membranes were blocked with TBS-T buffer [10 mM Tris-HCl (pH 8.0), 150 mM NaCl,

0.1% Tween 20, PBS] containing 5% skim milk for 1 h, and were then incubated with primary

antibodies in Can Get Signal Solution 1 (TOYOBO, Osaka, Japan) overnight at 4˚C.

Membranes were washed three times with TBST, incubated with HRP-conjugated secondary

antibodies in the blocking buffer for 2 h at room temperature, and washed five times with

TBST. Immunoreactive bands were detected using Clarity Western ECL Substrate (Bio-Rad,

Hercules, CA, USA) and the ChemiDoc MP Imaging System (Bio-Rad, Hercules, CA, USA).

The images were processed using ImageJ.

For immunoprecipitation, HEK293 cells were transfected with plasmids using

jetPRIME transfection reagent (Polyplus-transfecton, Illkirch, France) according to the

manufacturer's instructions. The following plasmids were used: pMRX-IRES-puro (-), pMRX-

IRES-puro-GFP-BNIP3_full (Full), pMRX-IRES-puro-GFP-BNIP3_ΔMER (MERΔ), and

pMRX-IRES-puro-3xHA-mCh-Atg18a. After 24 h of transfection, cells were washed with PBS

and lysed in lysis buffer [20 mM HEPES (pH 7.4), 1 mM EDTA, 150 mM NaCl, 0.1% Triton

X-100, 10% glycerol, 1 mM PMSF, and protease inhibitor cocktail (Nacalai Tesque, Kyoto,

Japan)]. Cell lysates were centrifuged at 17,000 x g for 1 min at 4˚C, and the supernatants were

incubated with GFP-Trap agarose beads (Chromotek, Planegg-Martinsried, Germany) for 1.5 h

at 4˚C with gentle rotation. After incubation, beads were washed with Wash Buffer [20 mM

HEPES (pH 7.4), 150 mM NaCl, 0.1% Triton X-100, 10% glycerol] to remove non-specific

proteins. Bound proteins were eluted by boiling in 2x sample buffer for 5 min, and subsequent

steps were performed as described above for immunoblots.

Structure prediction and visualization

The amino acid sequences of *Drosophila* BNIP3 (Q9VPD6) and Atg18a (Q9VSF0) were

retrieved from the UniProt Knowledgebase (https://www.uniprot.org/uniprotkb/) and submitted

to the AlphaFold Server (https://alphafoldserver.com/) to predict the structure of their complex.

The PyMOL Molecular Graphics System (Version 2.5.4; Schrödinger, LLC) was then used to

analyze the resulting five predictions and visualize the top-ranked structure.

Adult lifespan assay

The Adult lifespan assay was performed according to a previously reported protocol (Linford et

al., 2013). Briefly, age-synchronized adult flies were collected, sorted by sex, and housed in

groups of 30 per vial. Flies were maintained at 25˚C under a 12 h light/12 h dark cycle and

transferred to fresh food vials every 2-3 days without anesthesia. Mortality was recorded at each

transfer, and statistical significance was determined using the log-rank test.

Fly climbing assay

The climbing assay was conducted according to the protocol previously published (Gevedon et

al., 2019). Adult flies were gently transferred without anesthesia into a vertical climbing

chamber composed of two connected vials, marked at 5 cm from the bottom. After a brief

recovery period, flies were tapped to the bottom of the chamber, and the number of flies that

climbed above the 5 cm mark within 10 seconds was recorded. The climbing success rate was

calculated as the percentage of flies that reached or surpassed the mark. Each group was tested

in five trials, with a rest period of several minutes between trials. The median success rate

across trials was used as the representative value for each group.

Eclosion assay

Late-stage pupae were collected and maintained at 25˚C. The eclosion process was recorded,

and the time from initial rupture of the pupal case to the complete emergence of the adult fly

was measured. For each group, 7 to 10 individuals were analyzed, and the average elapsed time

was calculated.

Image analyses

For cell volume measurement (Fig. 2E), the DIOMs expressing GFP were imaged at 1 μm

intervals. The muscle cell volume was calculated by multiplying the area of each XY slice by

the interval length and summing the results across all slices. To measure nucleolus and nucleus

volume (Fig. 2-figure supplement 2E and F), DIOMs stained for DNA (H33342) and Fibrillarin

were imaged at 0.5 μm intervals. DAPI- and anti-Fibrillarin-stained areas were extracted by

binarization using ImageJ. This process was repeated for the z-stacks, and the nucleolus and

nucleus volume were determined by multiplying the area of each slice by the interval length

between slices. For mitochondrial area measurement in TEM images (Fig. 4C), each

mitochondrion and the corresponding DIOM region were manually segmented, and the total

mitochondrial area and DIOM area per image were calculated for each image using ImageJ. The

total mitochondrial area was then normalized to the DIOM area. For each genotype, at least 9

images from multiple animals were analyzed. Quantification of the number of autophagosomes,

mitophagosomes, or mitochondria in TEM images was conducted as follows (Fig. 4D and E):

Double-membrane-bound structures containing undigested cytoplasmic contents

(autophagosomes), autophagosomes enclosing mitochondria (mitophagosomes), and unengulfed

mitochondria were manually counted. At least 17 images derived from multiple animals were

analyzed for each genotype. Mitochondrial area in DIOMs was quantified in Fig. 5D' 6A' and

6D'. Cellular regions containing F-actin were manually segmented, and tdTomato (Fig. 5D') or

GFP (Fig. 6A' and 6D') channel intensity within these segmented regions was measured. For

Fig. 5D' and 6A', image preprocessing steps, including filtering, background subtraction, and

intensity binarization, were applied to the tdTomato (Fig. 5D') or GFP (Fig. 6A') channel using

ImageJ to reduce noise. The area of each detected object was measured, and the total

mitochondrial area was defined as the sum of these object areas. Mitochondrial area values were

then normalized to the corresponding cellular region area. For Fig. 6D', GFP intensity

quantification was performed as follows: because GFP signals in the 3IL BWM images were

weaker than those at 4 d APF, the GFP channel intensity was uniformly enhanced across all

images at a constant scale. All images were preprocessed using filtering and background

subtraction. The mean GFP intensity within each segmented area was then measured and

normalized, with the mean GFP intensity of the control for each developmental stage set to 1.

For Fig. 5D', at least 46 DIOMs from 5 animals were analyzed per genotype. For Fig. 6A' and

6D', at least five DIOMs per fly were analyzed, with three biological replicates per genotype.

For the Mito-QC assay, a representative DIOM was imaged for each genotype and analyzed by

ImageJ (Fig. 6B). Image preprocessing included filtering and background subtraction for both

GFP and mCherry channels. Dot plots were then generated, and Pearson's correlation

coefficients for GFP and mCherry intensities were calculated using the *Coloc2* plugin in ImageJ

(Fig. 6B and B').

Statistics

Each experiment was performed with at least three different cohorts of unique flies analyzed.

One exception was for TEM analyses, performed on two parallel replicates with multiple

animals each. All experiments were performed in parallel with controls. Raw data files

corresponding to all quantified results are provided in Supplementary File 2. Error bars show

the standard deviation for bar charts. When more than two genotypes were used in an

experiment, Sidak's test, Kruskal-Wallis test, or Dunnett's T3 multiple comparisons test was

used (GraphPad, Prism9 version 9.5.1). Mann-Whitney tests were used to compare two

means. $P<0.05$ was regarded as statistically significant (*$P<0.05$, **$P<0.001$, ***$P<0.0001$,

and ****$P<0.00001$).

Acknowledgments

We are grateful to JH. Lee (Univ. of Michigan), T. Yoshimori (Osaka Univ.), E. Kuranaga

(Kyoto Univ.), BDSC, VDRC, DGRC, and NIG-fly for reagents. We thank Y. Shimada and R.

Niwa (Univ. of Tsukuba) for the helpful discussion. We thank the Biomaterials Analysis

Division of the Institute of Science Tokyo, for the DNA sequencing. We are grateful to M.

Landekic (McGill Univ.) for English editing. This work was supported in part by Grant-in-Aid

for Transformative Research Areas (B) (grant number 21H05147, NF), Japan Science and

Technology Agency (JST) PRESTO (grant number JPMJPR18H8, NF), AMED PRIME (grant

number 24gm6410016h0001, NF), and MEXT Promotion of Development of a Joint Usage /

Research System Project: Pan-Omics DDRIC, MRCI for High Depth Omics, CURE:

JPMXP1323015486 for MIB, RIIT, and AMRC in Kyushu University.

Author contributions

NF conceived the project. HT, TM, TKanki, YO, and NF designed the experiments. HT, TM,

KK, MK, TKaminishi, YS, KT, AH, and NF performed the experiments. HT, TM, and NF

analyzed and interpreted data. YO contributed to the materials. NF took the lead in writing the

manuscript. All authors provided critical feedback and helped shape the research and

manuscript.

Competing financial interests

The authors declare no competing financial interests.

References

Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O, Willmore L,

Ballard AJ, Bambrick J, Bodenstein SW, Evans DA, Hung CC, O'Neill M, Reiman D,

Tunyasuvunakool K, Wu Z, Žemgulytė A, Arvaniti E, Beattie C, Bertolli O, Bridgland A,

Cherepanov A, Congreve M, Cowen-Rivers AI, Cowie A, Figurnov M, Fuchs FB,

Gladman H, Jain R, Khan YA, Low CMR, Perlin K, Potapenko A, Savy P, Singh S,

Stecula A, Thillaisundaram A, Tong C, Yakneen S, Zhong ED, Zielinski M, Žídek A,

Bapst V, Kohli P, Jaderberg M, Hassabis D, Jumper JM. 2024. Accurate structure

prediction of biomolecular interactions with AlphaFold 3. *Nature* **630**:493–500.

doi:10.1038/s41586-024-07487-w

Adriaenssens E, Schaar S, Cook ASI, Stuke JFM, Sawa-Makarska J, Nguyen TN, Ren X,

Schuschnig M, Romanov J, Khuu G, Lazarou M, Hummer G, Hurley JH, Martens S. 2024.

Reconstitution of BNIP3/NIX-mediated autophagy reveals two pathways and hierarchical

flexibility of the initiation machinery. *bioRxiv* doi:10.1101/2024.08.28.609967

Bunker EN, Le Guerroué F, Wang C, Strub M, Werner A, Tjandra N, Youle RJ. 2023. Nix

interacts with WIPI2 to induce mitophagy . *EMBO J* **42**. doi:10.15252/embj.2023113491

Fujita N., Huang W, Lin T-H, Groulx J-F, Jean S, Nguyen J, Kuchitsu Y, Koyama-Honda I,

Mizushima N, Fukuda M, Kiger AA. 2017. Genetic screen in Drosophila muscle identifies

autophagy-mediated T-tubule remodeling and a Rab2 role in autophagy. *Elife* **6**.

doi:10.7554/eLife.23367

Gevedon O, Bolus H, Lye SH, Schmitz K, Fuentes-González J, Hatchell K, Bley L, Pienaar J,

Loewen C, Chtarbanova S. 2019. In Vivo Forward Genetic Screen to Identify Novel

Neuroprotective Genes in Drosophila melanogaster. *J Vis Exp*. doi:10.3791/59720

Guo T, Nan Z, Miao C, Jin X, Yang W, Wang Z, Tu Y, Bao H, Lyu J, Zheng H, Deng Q, Guo

P, YongmeiXi X, Yang X, Ge W. 2019. The autophagy-related gene Atg101 in Drosophila

regulates both neuron and midgut homeostasis. *Journal of Biological Chemistry*

294:5666–5676. doi:10.1074/jbc.RA118.006069

Hashimshony T, Senderovich N, Avital G, Klochendler A, de Leeuw Y, Anavy L, Gennert D,

Li S, Livak KJ, Rozenblatt-Rosen O, Dor Y, Regev A, Yanai I. 2016. CEL-Seq2: Sensitive

highly-multiplexed single-cell RNA-Seq. *Genome Biol* **17**. doi:10.1186/s13059-016-0938-

8

Hegedus K, Nagy P, Gáspári Z, Juhász G. 2014. The putative HORMA domain protein Atg101

dimerizes and is required for starvation-induced and selective autophagy in Drosophila.

Biomed Res Int. doi:10.1155/2014/470482

Johansen T, Lamark T. 2020. Selective Autophagy: ATG8 Family Proteins, LIR Motifs and

Cargo Receptors. *J Mol Biol*. doi:10.1016/j.jmb.2019.07.016

Kamiyama T, Niwa R. 2022. Transcriptional Regulators of Ecdysteroid Biosynthetic Enzymes
and Their Roles in Insect Development. *Front Physiol*. doi:10.3389/fphys.2022.823418

Kanki T, Wang K, Cao Y, Baba M, Klionsky DJ. 2009. Atg32 Is a Mitochondrial Protein that
Confers Selectivity during Mitophagy. *Dev Cell* **17**:98–109.
doi:10.1016/j.devcel.2009.06.014

Karsli-Uzunbas G, Guo JY, Price S, Teng X, Laddha S V., Khor S, Kalaany NY, Jacks T, Chan
CS, Rabinowitz JD, White E. 2014. Autophagy is required for glucose homeostasis and
lung tumor maintenance. *Cancer Discov* **4**:915–927. doi:10.1158/2159-8290.CD-14-0363

Kawaguchi K, Fujita N. 2024. Shaping transverse-tubules: central mechanisms that play a role
in the cytosol zoning for muscle contraction. *J Biochem*. doi:10.1093/jb/mvad083

Kondo S, Ueda R. 2013. Highly Improved gene targeting by germline-specific Cas9 expression
in Drosophila. *Genetics* **195**:715–721. doi:10.1534/genetics.113.156737

Kumar L, Futschik M. 2007. Software Mfuzz: A software package for soft clustering of
microarray data. *Bioinformation* **2**:5–7. doi:10.6026/97320630002005

Lamark T, Johansen T. 2021. Mechanisms of Selective Autophagy. *Annual Review of Cell and
Developmental Biology Annu Rev Cell Dev Biol* **37**:143–69. doi:10.1146/annurev-cellbio-
120219

Lee JJ, Sanchez-Martinez A, Zarate AM, Benincá C, Mayor U, Clague MJ, Whitworth AJ.
2018. Basal mitophagy is widespread in Drosophila but minimally affected by loss of
Pink1 or parkin. *Journal of Cell Biology* **217**:1613–1622. doi:10.1083/jcb.201801044

Li Y, Zheng W, Lu Y, Zheng Y, Pan L, Wu X, Yuan Y, Shen Z, Ma S, Zhang Xingxian, Wu J,
Chen Z, Zhang Xiangnan. 2022. BNIP3L/NIX-mediated mitophagy: molecular
mechanisms and implications for human disease. *Cell Death Dis*. doi:10.1038/s41419-
021-04469-y

Liang M, Hody C, Yammine V, Soin R, Sun Y, Lin X, Tian X, Meurs R, Perdrau C, Delacourt
N, Oumalis M, Andris F, Conrard L, Kruys V, Gueydan C. 2023. eIF4EHP promotes Ldh
mRNA translation in and fruit fly adaptation to hypoxia. *EMBO Rep* **24**.
doi:10.15252/embr.202256460

Lieber T, Jeedigunta SP, Palozzi JM, Lehmann R, Hurd TR. 2019. Mitochondrial fragmentation
drives selective removal of deleterious mtDNA in the germline. *Nature* **570**:380–384.
doi:10.1038/s41586-019-1213-4

Linford NJ, Bilgir C, Ro J, Pletcher SD. 2013. Measurement of lifespan in Drosophila
melanogaster. *Journal of Visualized Experiments*. doi:10.3791/50068

Lőrincz P, Juhász G. 2020. Autophagosome-Lysosome Fusion. *J Mol Biol*.
doi:10.1016/j.jmb.2019.10.028

Mao Q, Acharya A, Rodríguez-Delarosa A, Marchiano F, Dehapiot B, Tanoury Z Al, Rao J,
Díaz-Cuadros M, Mansur A, Wagner E, Chardes C, Gupta V, Lenne PF, Habermann BH,
Theodoly O, Pourquié O, Schnorrer F. 2022. Tension-driven multi-scale self-organisation
in human iPSC-derived muscle fibers. *Elife* **11**. doi:10.7554/eLife.76649

Masiero E, Agatea L, Mammucari C, Blaauw B, Loro E, Komatsu M, Metzger D, Reggiani C,
Schiaffino S, Sandri M. 2009. Autophagy Is Required to Maintain Muscle Mass. *Cell*
Metab **10**:507–515. doi:10.1016/j.cmet.2009.10.008

Mcguire SE, Mao Z, Davis R. 2004. Spatiotemporal gene expression targeting with TARGET
and gene-swich systems in Drosophila. *Sci STKE* doi:10.1126/stke.2202004pl6

Morishita H, Mizushima N. 2019. Annual Review of Cell and Developmental Biology Diverse
Cellular Roles of Autophagy Autophagosome: a double-membrane structure enclosing
cytoplasmic materials. *Annu Rev Cell Dev Biol* **35**:453–75. doi:10.1146/annurev-cellbio-
100818

Murakawa T, Kiger AA, Sakamaki Y, Fukuda M, Fujita N. 2020. An autophagy-dependent
tubular lysosomal network synchronizes degradative activity required for muscle
remodeling. *J Cell Sci* **133**. doi:10.1242/jcs.248336

Murakawa T, Nakamura T, Kawaguchi K, Murayama F, Zhao N, Stasevich TJ, Kimura H,
Fujita N. 2022. A Drosophila toolkit for HA-tagged proteins unveils a block in autophagy
flux in the last instar larval fat body. *Development (Cambridge)* **149**.
doi:10.1242/dev.200243

Nakatogawa H. 2020. Mechanisms governing autophagosome biogenesis. *Nat Rev Mol Cell*
Biol. doi:10.1038/s41580-020-0241-0

Narendra DP, Youle RJ. 2024. The role of PINK1–Parkin in mitochondrial quality control. *Nat*
Cell Biol. doi:10.1038/s41556-024-01513-9

Ney PA. 2015. Mitochondrial autophagy: Origins, significance, and role of BNIP3 and NIX.
Biochim Biophys Acta Mol Cell Res. doi:10.1016/j.bbamcr.2015.02.022

Niemi NM, Friedman JR. 2024. Coordinating BNIP3/NIX-mediated mitophagy in space and
time. *Biochem Soc Trans*. doi:10.1042/BST20221364

Noda NN, Mizushima N. 2016. Atg101: Not Just an Accessory Subunit in the Autophagy-
initiation Complex. *Cell Structure and Function*. doi:10.1247/csf.15013

Novak I, Kirkin V, McEwan DG, Zhang J, Wild P, Rozenknop A, Rogov V, Löhr F, Popovic D,
Occhipinti A, Reichert AS, Terzic J, Dötsch V, Ney PA, Dikic I. 2010. Nix is a selective
autophagy receptor for mitochondrial clearance. *EMBO Rep* **11**:45–51.
doi:10.1038/embor.2009.256

Okamoto K, Kondo-Okamoto N, Ohsumi Y. 2009. Mitochondria-Anchored Receptor Atg32

Mediates Degradation of Mitochondria via Selective Autophagy. *Dev Cell* **17**:87–97.

doi:10.1016/j.devcel.2009.06.013

Onishi M, Yamano K, Sato M, Matsuda N, Okamoto K. 2021. Molecular mechanisms and

physiological functions of mitophagy. *EMBO J* **40**. doi:10.15252/embj.2020104705

Palozzi JM, Jeedigunta SP, Minenkova A V., Monteiro VL, Thompson ZS, Lieber T, Hurd TR.

2022. Mitochondrial DNA quality control in the female germline requires a unique

programmed mitophagy. *Cell Metab* **34**:1809-1823.e6. doi:10.1016/j.cmet.2022.10.005

Piccirillo R, Demontis F, Perrimon N, Goldberg AL. 2014. Mechanisms of muscle growth and

atrophy in mammals and Drosophila. *Developmental Dynamics* **243**:201–215.

doi:10.1002/dvdy.24036

Quy PN, Kuma A, Pierres P, Mizushima N. 2013. Proteasome-dependent activation of

mammalian target of rapamycin complex 1 (mTORC1) is essential for autophagy

suppression and muscle remodeling following denervation. *Journal of Biological*

Chemistry **288**:1125–1134. doi:10.1074/jbc.M112.399949

Ribeiro I, Yuan L, Tanentzapf G, Dowling JJ, Kiger A. 2011. Phosphoinositide regulation of

integrin trafficking required for muscle attachment and maintenance. *PLoS Genet* **7**.

doi:10.1371/journal.pgen.1001295

Sandoval H, Thiagarajan P, Dasgupta SK, Schumacher A, Prchal JT, Chen M, Wang J. 2008.

Essential role for Nix in autophagic maturation of erythroid cells. *Nature* **454**:232–235.

doi:10.1038/nature07006

Sanger JW, Wang J, Fan Y, White J, Sanger JM. 2010. Assembly and dynamics of myofibrils. *J*

Biomed Biotechnol. doi:10.1155/2010/858606

Sartori R, Romanello V, Sandri M. 2021. Mechanisms of muscle atrophy and hypertrophy:

implications in health and disease. *Nat Commun*. doi:10.1038/s41467-020-20123-1

Saxton RA, Sabatini DM. 2017. mTOR Signaling in Growth, Metabolism, and Disease. *Cell*.

doi:10.1016/j.cell.2017.02.004

Schwarten M, Mohrlüder J, Ma P, Stoldt M, Thielmann Y, Stangler T, Hersch N, Hoffmann B,

Merkel R, Willbold D. 2009. Nix directly binds to GABARAP: A possible crosstalk

between apoptosis and autophagy. *Autophagy* **5**:690–698. doi:10.4161/auto.5.5.8494

Schweers RL, Zhang J, Randall MS, Loyd MR, Li W, Dorsey FC, Kundu M, Opferman JT,

Cleveland JL, Miller JL, Ney PA. 2007. NIX is required for programmed mitochondrial

clearance during reticulocyte maturation. *Proc. Natl. Acad. Sci. U.S.A.* **104**:19500-19505.

doi:10.1073/pnas.0708818104

Semenza GL. 2014. Oxygen sensing, hypoxia-inducible factors, and disease pathophysiology.
Annual Review of Pathology: Mechanisms of Disease **9**:47–71. doi:10.1146/annurev-
pathol-012513-104720

Strong LM, Chang C, Riley JF, Boecker CA, Flower TG, Buffalo CZ, Ren X, Stavoe AK,
Holzbaur EL, Hurley JH. 2021. Structural basis for membrane recruitment of ATG16L1
by WIPI2 in autophagy. *Elife* doi:10.7554/eLife

Vargas JNS, Hamasaki M, Kawabata T, Youle RJ, Yoshimori T. 2023. The mechanisms and
roles of selective autophagy in mammals. *Nat Rev Mol Cell Biol*. doi:10.1038/s41580-022-
00542-2

Wang X, Yu L, Wu AR. 2021. The effect of methanol fixation on single-cell RNA sequencing
data. *BMC Genomics* **22**. doi:10.1186/s12864-021-07744-6

Yamashita SI, Sugiura Y, Matsuoka Y, Maeda R, Inoue K, Furukawa K, Fukuda T, Chan DC,
Kanki T. 2024. Mitophagy mediated by BNIP3 and NIX protects against ferroptosis by
downregulating mitochondrial reactive oxygen species. *Cell Death Differ* **31**:651–661.
doi:10.1038/s41418-024-01280-y

Yoshii SR, Kuma A, Akashi T, Hara T, Yamamoto A, Kurikawa Y, Itakura E, Tsukamoto S,
Shitara H, Eishi Y, Mizushima N. 2016. Systemic Analysis of Atg5-Null Mice Rescued
from Neonatal Lethality by Transgenic ATG5 Expression in Neurons. *Dev Cell* **39**:116–
130. doi:10.1016/j.devcel.2016.09.001

Zhang J, Loyd MR, Randall MS, Waddell MB, Kriwacki RW, Ney PA. 2012. A short linear
motif in BNIP3L (NIX) mediates mitochondrial clearance in reticulocytes. *Autophagy*
8:1325–1332. doi:10.4161/auto.20764

Zhang S, Wu S, Yao R, Wei X, Ohlstein B, Guo Z. 2024. Eclosion muscles secrete ecdysteroids
to initiate asymmetric intestinal stem cell division in Drosophila. *Dev Cell* **59**:125-
140.e12. doi:10.1016/j.devcel.2023.11.016

Zhang X, Avellaneda J, Spletter ML, Lemke SB, Mangeol P, Habermann BH, Schnorrer F.
2024. Mechanoresponsive regulation of myogenesis by the force-sensing transcriptional
regulator Tono. *Current Biology*. doi:10.1016/j.cub.2024.07.079

Zitserman D, Roegiers F. 2011. Live-cell Imaging of Sensory Organ Precursor Cells in Intact
Drosophila Pupae. *J Vis Exp*. doi:10.3791/2706

Figure legends

Figure 1. Time-course RNA-seq of the DIOM remodeling in wild-type *Drosophila*

(A) A timeline of DIOM remodeling. Samples were collected at five time points: third instar

larva (3IL) and 1 to 4 days after puparium formation (APF). (B) The scheme for sample

preparation of DIOMs. Red-colored rectangles indicate single DIOMs. (C) DESeq2 principal

component analysis (PCA) of all mRNA-seq libraries. The first four principal components are

shown. PC1-2, left; PC3-4, right. The dotted arrow in the PC1-2 plot represents the direction of

the transcription dynamics. (D) Heat map of fuzzy c-means cluster core expression profiles. All

genes were categorized into twelve clusters. (E) Factor loadings of each cluster to PC1-2 and

PC3-4. The circles show r = 0.8 (PC1-2) or r = 0.7 (PC3-4). (F) GO enrichment at each time

point during metamorphosis. The width of each line represents the expression level of the

clusters indicated.

Figure 2. Transcriptional dynamics during DIOM remodeling is independent of

autophagy

(A) A schematic of DIOM remodeling in wild-type and autophagy-deficient conditions.

Autophagy-dependent muscle atrophy starts at 12-14 h APF. (B) Genotypes and time points of

the comparative RNA-seq analysis (top) and a diagram of autophagosome formation (bottom).

FIP200 or *Atg18a* RNAi blocks autophagosome formation. *Stx17* RNAi blocks the

autophagosome-lysosome fusion. (C) DESeq2 PCA of all mRNA-seq libraries. PC1-2, left;

PC3-4, right. A total of 20 samples were analyzed. (D and E) DIOM volume changes in control

or *FIP200* RNAi from 3IL to 4 d APF. DIOMs were labeled with GFP. Projected images of XY

and XZ planes are shown (D). (E) Relative DIOM cell volume for each genotype normalized to

3IL (set to 1). N=5 (Mann-Whitney test).

Figure 3. Loss of BNIP3 results in accumulation of mitochondria in DIOMs

(A) A schematic of 4 d APF DIOMs in wild-type, *FIP200* RNAi, or *Stx17* RNAi. (B) TEM

image of *Stx17* RNAi DIOM at 4 d APF. White arrowheads indicate autophagosome structures.

(C) TEM image of wild-type DIOM at 1 d APF. White arrows indicate mitophagosome

membrane structure. (D) The expression level of mitophagy regulators in DIOMs at 1 d APF. N

= 4. Normalized counts in RNA-seq are shown. (E) The expression level of *BNIP3* and *Fbxl4* in

DIOMs. N = 4. (F) A diagram illustrating the *BNIP3* knockout strategy, where all exons were

deleted using two gRNAs and replaced with a 3xP3-RFP marker. (G) Loss of BNIP3 phenotype

on mitochondria and myofibrils in DIOM at 4 d APF. XY (top) and XZ (bottom) planes were

shown.

Figure 4. BNIP3 is required for mitophagosome formation

(A and B) TEM images of DIOM transverse sections at 4 d APF in wild-type (A) or *BNIP3* KO

(B). Mitochondria are shown in green. (C) The percentages of mitochondrial area per unit area

in the indicated genotypes. wild-type, N = 9; *BNIP3* KO, N = 12 (Mann-Whitney test). (D and

E) TEM images of DIOM transverse sections at 4 d APF in *Stx17* RNAi (D and D') or a

combination of *Stx17* RNAi and *BNIP3* KO (E and E'). Mitophagosomes, pink; Mitochondria,

green; autophagosomes, blue. (F) The number of mitophagosomes and mitochondria per unit

area in the indicated genotypes. (G) The number of total autophagosomes, including

mitophagosomes, per unit area in the indicated genotypes. *Stx17* RNAi, N = 20; *Stx17* RNAi

and *BNIP3* KO, N = 17 (Mann-Whitney test) (F and G).

Figure 5. The LIR and MER motifs are required for BNIP3-mediated mitochondrial

clearance

(A) Schematics of *Drosophila* BNIP3 and its mutants. (B) The structure of the BNIP3-Atg18a

complex predicted by AlphaFold 3. Top, overview: Atg18a is depicted in white, featuring a β-

propeller structure consisting of seven blades, with blades 2 and 3 highlighted in green and pink,

respectively. For BNIP3, only residues 29–74 are shown in blue for clarity, with the α-helix

spanning residues 42–53 highlighted in yellow. Bottom, close-up view: Amino acids positioned

to form intramolecular contacts through their side chains are labeled and represented as sticks,

with potential hydrogen bonds shown as dashed lines. (C) GFP pulldown experiment between

GFP-BNIP3_full or MERΔ construct and HA-mCh-Atg18a in HEK293 cells. (D and D') BNIP3

rescue experiment in DIOMs at 4 d APF. The indicated GFP-tagged BNIP3 constructs and

tdTomato-Mito were co-expressed in *BNIP3* KO flies using the GAL4/UAS system (D). Single-

channel images corresponding to the merged panels are presented in Fig. 5-figure supplement

1B. The percentages of mitochondrial area in DIOMs for the indicated genotypes. Empty (-), N

= 46; Full, N = 55; ΔLIR, N = 52; MER^{mut}, N = 48; ΔMER, N = 54; ΔLIR+ ΔMER, N = 51

(Kruskal-Wallis test) (D'). (E) The amount of GFP-BNIP3 in *WT* and *Atg101* KO muscles. The

GFP-BNIP3 constructs were expressed by Mef2-GAL4 in *WT* or *Atg101* KO flies. Larval fillets

were lysed and subjected to western blotting for GFP. After immunoblotting, the membrane was

stained with CBB to assess total protein loading.

Figure 6. BNIP3-mediated mitophagy eliminates larval muscle mitochondria during

muscle remodeling

(A and A') Time course microscopy of Mito-GFP and F-actin in control or *BNIP3* KO during

DIOM remodeling (A). Single-channel images corresponding to the merged panels are

presented in Fig. 6-figure supplement 1A. Mitochondrial area relative to total cell area. For 3IL,

control, N = 18; *BNIP3* KO, N = 17. For 1 d APF, control, N = 20; *BNIP3* KO, N = 18 (Mann-

Whitney test) (A'). (B and B') Mitophagy assay using Mito-QC in DIOMs at 1 d APF. Pixel

intensity correlation profiles and Pearson's correlation coefficients (R values) are shown.

Control, N = 6; *BNIP3* KO, N = 5. *FIP*200 RNAi, N = 5 (Sidak's test). (C, D, and D') Scheme

of the use of GAL80 temperature-sensitive mutants (GAL80^{ts}). The animals were raised at 29°C

(Restrictive) to induce Mito-GFP expression until mid-3IL, then shifted to 18°C (Permissive) to

block expression (C). Mito-GFP and ATP5A immunostaining (total mitochondria) signals in

muscles at 3IL or 4 d APF (D). Mito-GFP intensities in muscles at each time point normalized

to control (set to 1). For 3IL, control, N = 16; *BNIP3* KO, N = 15. For 1 d APF, control, N = 15;

BNIP3 KO, N = 15 (Mann-Whitney test) (D'). (E) A model of muscle remodeling with or

without BNIP3. Loss of BNIP3 leads to mitochondrial accumulation, which disrupts muscle

remodeling.

 Supplementary materials legends

Figure 1—figure supplement-1. RNA-seq of DIOM remodeling

(A) Heatmap of comparisons for each sample based on correlation coefficients. (B) The

contribution ratios (bar) and cumulative contribution ratios (red-colored line) of PC1 to 10 are

shown. (C) Fuzzy c-means clustering categorized expressed genes in DIOMs into twelve

groups. (D) Violin plot showing the expression level of each cluster, with the number of genes

in each cluster indicated at the bottom.

Figure 2—figure supplement-1. Expression levels of Atg18a, FIP200, and Stx17 during

DIOM remodeling

(A) Heatmap of comparisons for each sample based on correlation coefficients. (B-D)

Expression levels of *Atg18a* (B), *FIP200* (C), and *Stx17* (D) in DIOMs under the indicated

conditions. N = 4.

Figure 2—figure supplement-2. Effect of autophagy deficiency on transcription of

ribosomal RNA and proteins

(A) Time-course analysis of the amount of rRNA during DIOM remodeling in the indicated

genotypes. (B and C) Loss of autophagy on the number of nuclei in DIOMs at 3 d APF. Nuclei

in DIOMs were stained with Hoechst (B). N = 6 (Sidak's test) (C). (D-F) Loss of autophagy on

nucleolus volume in DIOMs at 3 d APF. DNA and fibrillarin were stained. XY and XZ planes

are shown (D). The nucleolus (E) and nucleus (F) volumes were calculated from Z-series

images. N = 12 (Dunnett's T3 multiple comparisons test). (G) A heatmap of the expression

levels of RpLs (left) and RpSs (right) relative to control at each time point.

1008

1009 **Figure 3—figure supplement-1. Loss of BNIP3 on mitochondria and myofibrils in DIOMs**

1010 YFP-Mito and Phalloidin-Alexa633 (F-actin) staining in DIOMs at 4 d APF in the indicated

1011 genotypes.

1012

1013 **Figure 3—figure supplement-2. Loss of BNIP3 on adult fly lifespan and mobility**

1014   (A) Survival rates of wild-type and *BNIP3* KO adult males. *WT*, N = 298; *BNIP3* KO, N = 297.

1015   (Log-rank test). (B) Climbing ability of wild-type and *BNIP3* KO adult males at 1 and 4 weeks

1016   old. N = 6 (Mann-Whitney test). (C) Elapsed time to eclosion in wild-type males and *BNIP3*

1017   KO males. WT, N = 10; *BNIP3* KO, N = 7 (Mann-Whitney test).

1018

Figure 5—figure supplement-1. GFP-BNIP3 constructs localize to the outer mitochondrial

membrane

(A) Localization of GFP-BNIP3 constructs and mitochondria matrix-targeted tdTomato

(tdTomato-Mito) in 3IL BWMs. (B) BNIP3 rescue experiment in DIOMs at 4 d APF. The

indicated GFP-tagged BNIP3 constructs and tdTomato-Mito were co-expressed in *BNIP3* KO

flies using the GAL4/UAS system. Individual fluorescence channels are displayed separately.

1025

1026   **Figure 5—figure supplement-2. Effect of GFP-BNIP3 overexpression on mitophagy flux in**

1027   **3IL BWMs**

1028   GFP-BNIP3 or mitochondria matrix-targeted GFP (Mito-GFP) were overexpressed together

1029   with tdTomato-Mito in 3IL BWMs. No clear difference in tdTomato-Mito signal was observed

1030   between samples with or without GFP-BNIP3 overexpression.

1031

Figure 6—figure supplement-1. The impact of BNIP3 knockout on the shape of the DIOM

during metamorphosis

(A) Time-course microscopy of mCD8-GFP in dorsal muscles imaged through the cuticle in

control and BNIP3 KO animals from 3IL to 4 d APF. (B) Time-course microscopy of Mito-GFP

and F-actin in control or *BNIP3* KO during DIOM remodeling. Individual fluorescence channels

are displayed separately.

1038

1039   **Supplementary Table 1**

1040   Detailed Drosophila genotypes, developmental stages, and experimental temperatures are

1041   provided.

1042

1043   **Supplementary File 1**

1044   Normalized counts from time-course RNA-seq of DIOM remodeling during metamorphosis.

1045

1046 **Supplementary File 2**

1047 Raw data files corresponding to all quantified results presented in the manuscript.

1048

1049 **Data availability**

1050 RNA-seq data have been deposited in GEO under accession number GSE293359.

Figure 1

Figure 2

Figure 3

wild-type

BNIP3 KO

Stx17 RNAi

Stx17 RNAi + BNIP3 KO

■ Mitophagosomes ■ Mitochondria ■ Autophagosomes

500 nm

Figure 4

Figure 5

Figure 6