scRNA+TCR-seq Reveals the Proportion and Characteristics of Dual TCR Treg Cells in Mouse Lymphoid and Non-lymphoid Tissues

Yuanyuan Xu 1#, Qipeng 1#, Xiaoping Lu 1, Long Ma 1, Jun Li 1, Xinsheng Yao 1*

1.Department of Immunology, Zunyi Medical University; Guizhou Provincial Research Center for Applied Immunomolecular Engineering; Guizhou Provincial Innovation Base for Graduate Education in Immunology; Guizhou Provincial Key Discipline of Immunology

#Co-first authors; * Corresponding author: immmunology@126.com

Abstract:

As an essential molecule for Treg cell function, TCR plays a critical role in various immune responses including cell development, immune tolerance, and self-antigen recognition. Current research has identified Treg cells in both lymphoid and non-lymphoid tissues as important for maintaining immune homeostasis and preventing autoimmune reactions. However, the TCR pairing types and CDR3 characteristics of these Treg cells are largely unknown. Through innovative analysis of scRNA+TCR-seq data, we discovered that 10.1% to 21.4% of dual TCR Treg cells exist in mouse lymphoid and non-lymphoid tissues. Dual TCR Treg cells in different tissues exhibit differential phenotypic and TCR repertoire characteristics. These findings provide key insights into the complex biological functions of Treg cell V(D)J recombination, tolerance selection, development, and migration.

Introduction

The discovery of CD4+CD25+ Treg cells in 1995, has since been linked to the occurrence and development of various human diseases. In addition to their strong regulatory role in self-reactivity and excessive inflammation, Treg cells play an important homeostatic role in the maintenance of tissue homeostasis, muscle damage repair, hair follicle regeneration, fat metabolism, among other complex cellular

activities[1][2]. The diverse negative regulatory functions of Treg cells suggest unique

characteristics for their TCR diversity, specificity, and memory responses. However,

the compositional characteristics of Treg cell TCR, their dependency on antigen

epitopes, activation, and maintenance mechanisms, remain largely unelucidated.

Some studies have garnered widespread attention, including:V(D)J recombination of

nTreg cells TCRs, self-tolerance selection, and migratory settlement with or without

lineage characterization;differences in antigen induction and activation mechanisms

for iTreg versus conventional helper and effector T cells;diversity and plasticity of

Treg cells in lymphoid and non-lymphoid tissues, such as TCR CDR3, and their

correspondence to regulatory functions.

Since their initial report in 1988[3], dual TCR expressing lymphocytes have been

widely supported by experimental evidence and play important roles in physiological

and pathological processes such as autoimmune tolerance, transplant rejection

reactions, and T cell tumors[4][5][6][7][8][9]. Due to technical limitations, comparing the

proportion, characteristics, origin, and mechanisms of single TCR and dual TCR T

cells in a single sample of more than a thousand T cells has been challenging for

immunologists. However, scRNA-seq combined with scTCR-seq can mark over 5,000

T cells from a single research sample at once, obtaining the pairing types of each T

cell's TCR beta chain and alpha chain, CDR3 sequence composition characteristics,

and complete mRNA expression data. This allows for the accurate analysis of TCR

CDR3 characteristics in the development, differentiation, maturation, and

response/tolerance processes of each T cell, as well as the corresponding expression

of regulatory and effector molecules. This provides unprecedented opportunities for

in-depth analysis of dual receptor T cells[8][9][10][11].

First, in the Treg study conducted by Oliver T et al.[12], which involved analyzing

single-cell RNA sequencing and T cell receptor (scRNA+TCR-seq) data of Foxp3+

Treg cells sorted from mouse blood, kidney, liver, LPL, and pancreas under

physiological conditions, we conducted a detailed comparative analysis of the

proportions of single TCR Treg and dual TCR Treg; the usage, diversity, clonality,

and overlap of CDR3 VJ; and the homogeneity and heterogeneity of characteristic

marker mRNA expression.Additionally, to further explore the differences in dual TCR Treg cells between lymphoid and non-lymphoid tissues in mice, we selected the scRNA+TCR-seq data shared by Annekathrin T et al.[13], which includes sequencing results of CD4+CD25+ Treg cells sorted from mouse spleen, inguinal lymph nodes (iLN), and mesenteric lymph nodes (mLN) under physiological conditions, as well as CD3+ T cells (including Treg and non-Treg) sorted from skin tissue, and performed comparative analysis again.

Results

1.A high proportion of dual TCR Treg cells in mouse lymphoid and non-lymphoid tissues:

The proportion of dual TCR Treg cells in mouse lymphoid tissues: spleen=21.4%; iLN=21.4%, mLN=21.0% (Figure 1D); The proportion of dual TCR Treg cells in mouse non-lymphoid tissues: blood=11.9%; kidney=11.2%; liver=13.6%, LPL=10.1%; pancreas=13.2%, The proportion of dual TCR T cells in mouse skin tissue: 18.6% (Figure 3D). Subsequently, a detailed analysis of the pairing of dual TCR T cells in the two sets of data was conducted, revealing that the TCR pairing types of dual TCR Treg include A+B1+B2, B+A1+A2 (the highest proportion), A1+A2+B1+B2 (the lowest proportion). There are significant differences in TCR pairing types between different tissues, with A1+A2+B1+B2 significantly increased in skin and kidney, and B+A1+A2 significantly higher in spleen and liver than in other groups (Figure 1D, Figure 3D).

2.Homogeneity and heterogeneity of TCR composition in mouse dual TCR Treg cells:

VJ usage(Figure 1E/F, Figure 3E/F):The overall VJ family distribution pattern of single TCR Treg and dual TCR Treg between different tissues is basically consistent. Compared with single TCR Treg, dual TCR Treg displays specific preferences in V and J usage, mainly including: TRBV10 and TRBV13-1 in the kidney and

liver,TRBV5 in LPL,TRBV19 in the skin;TRBJ2-3 in the liver, TRBJ2-1 in the

skin,etc. showing a significant advantage in usage; TRBV13-1 in peripheral blood,

kidney, and liver shows consistent advantage in usage; TRBJ1-2 in the spleen, iLN,

and mLN shows consistent advantage in usage, etc.

Diversity of CDR3(Figure 1G, Figure 3G):The 1/DS of single TCR Treg CDR3 is

significantly higher than that of dual TCR Treg; the CDR3 diversity of dual TCR Treg

in lymphoid tissues (spleen, iLN, mLN) is higher than that in non-lymphoid tissues;

the CDR3 diversity of dual TCR Treg in skin, LPL, and pancreas is very low.

Clonality of CDR3(Figure 1H, Figure 3H):The proportion of single TCR Treg with

clonal proliferation greater than or equal to 2 is higher than that of dual TCR Treg; the

proportion of dual TCR Treg with clonal proliferation greater than or equal to 2 in the

liver and skin is significantly higher than that in other tissues.

Overlap of CDR3(Figure 2A/B, Figure 4A/B):A certain proportion of overlap exists

between TRB CDR3 AA and TRA CDR3 AA in different tissues, with single TCR

Treg significantly higher than dual TCR Treg, and a lower proportion of TRB CDR3

AA compared to TRA CDR3 AA. This suggests that TRA allelic inclusive

recombination involves more CDR3 AA types in TCR pairing of Treg cells.

Meanwhile, it was found that single and dual TCR Treg in different tissues can also

share common CDR3 AA. This suggests that both single and dual TCR Treg can

respond to specific antigen epitopes.

All three tissues, the liver, kidney, and pancreas, show a significantly high

proportion of dual TCR Treg CDR3 AA overlap with peripheral blood, suggesting

their origin from the influx of peripheral blood. iLN, mLN, and skin all show a

significantly high proportion of dual TCR Treg CDR3 AA overlap with the spleen,

suggesting their origin from the influx of the spleen; Intriguingly, there is also a high

proportion of dual TCR Treg CDR3 AA overlap between the liver & kidney, kidney

& pancreas, and iLN & mLN, suggesting that dual TCR Treg cells in these tissue sites

may negatively regulate a shared set of antigens. Whereas the mechanism and

significance of the high percentage of overlap exhibited between iLN and skin TRA

CDR3 AA deserves further analysis.

3.Homogeneity of mouse dual TCR Treg cell phenotypes:

Comparative analysis of Treg cells with complete scRNA-seq and scTCR-seq results showed that the TCR pairing types and proportion were consistent with the results of direct analysis of scTCR-seq; the proportion of dual TCR Treg in the skin was higher than that of dual TCR non-Treg (Figure 2C, Figure 4C); in all tissues,the expression of signature molecules *Tnfrsf9, Stat3, Tnfrsf4, Cd27, Icos, Il2ra, Ikzf2, Lrrc32, Sh3rf1, Cd81, Baff, Smad2*, and *Foxo3* was identical between dual and single TCR Treg in all tissues, with slightly higher *Foxp3, Foxo1, and Ctla4* expression in dual TCR Treg than that in single TCR Treg(Figure 2D, Figure 4D).

4.Heterogeneity of mouse dual TCR Treg cell phenotypes:

Compared with iLN, mLN, and spleen, more characteristic mRNA molecules were highly expressed in dual TCR Tregs in skin (Figure 2E), such as *Rora, Dusp1, Junb* involved in immune responses and inflammatory diseases; *S100A6* involved in the proliferation, differentiation, and migration of immune cells; *Hopx* and *Nr4a1* involved in the development of immune cells, etc., suggesting that dual TCR Treg in the skin may be involved in more specific response negative regulations.

Between Blood & Liver; Kidney & Pancreas (Figure 2E), iLN & mLN & spleen (Figure 4E), the mRNA expression of cytokines, cytokine receptors, and transcription factors in dual TCR Treg shows higher homogeneity, suggesting that dual TCR Treg in these tissue sites may play convergent negative immune response regulations.

Differential mRNA expression analysis between dual TCR Treg and single TCR Treg in each tissue (Figure 2F, Figure 4F): showed significant high expressions of individual TRBV genes in different tissues, consistent with scTCR-seq results, suggesting specific dual TCR Treg lineage origins in different tissues; in addition to TRBV usage, each group's dual TCR Treg had significantly different characteristic molecule expressions from single TCR Treg, such as Blood: *H1f0* involved in immune cell proliferation and differentiation and *Hist1h2ap* related to immune cell activation were significantly increased; Kidney: *Fos* and *Cenpa* related to immune

cell proliferation and differentiation and *Ifit3* involved in the body's antiviral response

were significantly increased; Liver: *Cenpa* related to immune cell proliferation and

Birc5 involved in the body's anti-infection immune response were significantly

increased; LPL: *Flnb* related to cell adhesion and migration regulation and *Atp1a1*

related to cell proliferation, differentiation, and apoptosis regulation were

significantly increased; Pancreas: *Ccl5, Ccr2, Cxcr6* related to immune cell migration

and *Lbr* and *Id2* related to immune cell proliferation and differentiation, as well as

Fkbp4 involved in immune suppression signal transduction were significantly

increased; iLN: *Tgfbr1* involved in maintaining immune tolerance and regulating

inflammatory responses and *Traf3* involved in immune cell activation and cytokine

production were significantly increased; mLN: *Glcci1* involved in inflammation and

immune cell migration was significantly increased; skin: *Areg* involved in immune

cell proliferation, differentiation, migration, and immune suppression functions and

Ccr10 involved in T cell migration and positioning were significantly increased;

spleen: *Casc3* related to T cell activation and proliferation was significantly increased.

This suggests that dual TCR Treg and single TCR Treg in each specific tissue site

may be involved in different negative immune response regulations.

Consistent high expression of molecular mRNA between different lymphoid

tissues, such as: the presence of the *Nr4a1* gene related to T cell development,

activation, and proliferation in both iLN and mLN tissues; partial sharing of

molecular mRNA high expression between different non-lymphoid tissues, such as:

the expression of *Trbv13−1* and *Trbv5* genes in Blood, Kidney, and Liver tissues, and

the common expression of *Trbv1* in skin and Kidney, these results may suggest that

dual TCR Treg cells in some tissues have specific lineage origins; and the common

expression of the *Cenpa* gene in Kidney and Liver tissues may play a common

immune effect; similarly, single TCR Treg also has similar gene expressions, such as

the high expression of *Trbv1* in single TCR Treg cells in Blood and Liver tissues, and

the high expression of *Hist1h2ap* related to immune cell activation in skin and

Kidney.

Discussion

Currently, the proportion of dual TCR T cells is reported to be extremely variable across physiologic and pathologic states (0.1%-30%, etc.) due to differences in assay technology and methodology[4][5][6][7][8][9]. This study, for the first time in a large number of single Treg cell scRNA+TCR-seq studies, found that mouse lymphoid and non-lymphoid tissues both have a high proportion of dual TCR Treg cells, with lymphoid tissues being higher than non-lymphoid tissues, and the proportion of liver and pancreas dual TCR Treg cells in non-lymphoid tissues is relatively high.In addition, our analysis of dual TCR T cell pairing types revealed significant differences between tissues. These findings not only suggest that lymphoid tissues have a high proportion of dual TCR Treg cells but also provide a comparative baseline and research direction for tissue-specific Treg cells in different locations.

The lower diversity of dual TCR Treg CDR3, the tissue preference in V/J gene usage, and the high clonal proliferation frequency in some tissues reflect the limited breadth of the dual TCR Treg subset response. These cells may play a complementary role in specific physiological or pathological negative regulation, and the tissue-specific preference for V/J usage suggests that dual TCR Treg cells may have a characteristic lineage similar to that of γδ T cells. Current research indicates that γδ T cells in different tissues show differences in VJ subfamily usage due to directed migration after specific VJ recombination at different times[14]. Dual TCR Treg cells may have certain lineage characteristics in V(D)J recombination and self-tolerance selection, and migrate to settle in different tissues. The higher clonal proliferation frequency of dual TCR Treg cells in the liver may exert a stronger inhibitory response to certain specific antigenic epitopes. The relatively higher immune tolerance of liver transplantation compared to other organ transplants may be related to the high proportion and clonal proliferation of dual TCR Treg cells in the liver.

The majority of single/dual TCR Treg cells exhibit consistent expression of characteristic genes, and the high expression of certain genes in dual TCR Treg cells

(*Foxp3, Foxo1, Ctla4*) indicates that dual TCR Treg cells can function similarly to or

more potently than single TCR Treg cells. In addition, specific or shared Trbv genes

and mRNA molecule expression in dual TCR Treg cells has been identified in

lymphoid or non-lymphoid tissues. Recently, Malte et al. used scATAC-seq

technology to reveal organ-specific adaptation and conservation of tissue-resident

immune cells such as Treg and Th17, and identify key transcription factors for

multiple tissue-resident immune cells[15]. Overall, the homogeneity and heterogeneity

of these characteristic molecules within Treg cells suggests their potential induction in

response to different antigenic epitopes and corresponding negative regulatory

functions.

This study, based on single-cell RNA sequencing and T cell receptor

(scRNA+TCR-seq) data of Treg cells in lymphatic and non-lymphatic tissues of mice

under physiological conditions, reveals for the first time the phenomenon of a large

number of dual TCR Treg cells in mice. Moreover, the CDR3 regions of these dual

TCR Treg cells exhibit certain tissue-specific or lineage-specific characteristics in

terms of VJ usage, diversity, and clonality. Compared to single TCR Treg cells, dual

TCR Treg cells highly express immune-inhibitory related molecules, such as *Foxo3,*

Foxo1, *CD27*, *IL2RA*, *and Ikzf2*, etc. Dual TCR Treg cells in LPL and skin also have

more mRNA differentially expressed characteristic molecules. These results provide a

new Treg cell subset (dual TCR Treg) for studying phenotypic characteristics, TCR

composition characteristics, etc. This study delves into the complex biological effects

and mechanisms of Treg cells from the perspective of TCR mRNA expression,

providing a new perspective for Treg cell research and also offering technical

solutions for decoding the TCR pairing and characteristics of dual TCR T cells.

However, this study still has some shortcomings, such as the lack of protein data for

individual dual TCR Treg cells and research on human dual TCR Treg cells.

Subsequent studies can use dual receptor transgenic reporter mice[16], combined with

FCM, tetramers, and scCITE-seq technology, to provide a foundation for research at

the TCR protein expression level. We hope that more laboratories will participate in

the study of dual TCR Treg cells, especially to further clarify the diversity and

specificity of the immune negative regulatory role of Treg cells in human lymphatic

and non-lymphatic tissues.

Materials and Methods:

1. Research Subjects and Study Samples:(1) Blood,kidney,liver,LPL and panc

reas tissues were collected from 4 C57BL/6 mice and CD4+Foxp3+ Thy1.1+ T

reg cells were sorted for scRNA+TCR-seq analysis.(Source document:Oliver T

et al.The tissue-resident regulatory T cell pool is shaped by transient multi-tiss

ue migration and a conserved residency program.Immunity. 2024 Jul 9;57(7):158

6-1602.e10. Shared Data Link:https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc

=GSE266111)

(2) Spleen,iLN and mLN tissues from 4 mice were collected and sorted into C

D4+CD25+Treg cells,while skin tissues from 4 mice were collected and sorted

into CD3+ T cells(both Treg and non-Treg),These four tissue-sorted cells were

analyzed by scRNA+TCR-seq. (Source document:Nedwed et al.Using combined

single-cell gene expression,TCR sequencing and cell surface protein barcoding

to characterize and track CD4+ T cell clones from murin tissues.Front Immun

ol. 2023 Oct 12;14:1241283. Shared Data Link:https://www.ncbi.nlm.nih.gov/geo

/query/acc.cgi?acc=GSE240041)

2.The process of sharing raw data analysis： (1)Individual Treg cells (including

CD3+ T cells in skin tissues) were screened for sequencing of both the TCR alpha

chain (A) and TCR beta chain (B); (2) In a single T cell, two types of chains "can pair

and assemble into single TCR" T cells named "A+B"; Three or more types of chains

"can pair and assemble into dual TCR" T cells named

"A+B1+B2;B+A1+A2;A1+A2+B1+B2", (3) The proportions of total single TCR

Treg and dual TCR Treg were calculated for each mouse or each tissue.

3.Examples of single T-cell TCR pairing types: From the two data sets, single T

cell was selected to display four different TCR pairing types, VDJ gene family names

and CDR3 AA sequences:A+ B; A+B1+B2; B+A1+A2; and A1+A2+B1+B2.

4.Analysis of TCR pairing types and V-J usage: Statistically analyze the

proportions of five TCR T cell types in each sample, and use SPSS software to

perform statistical analysis on the proportions of the three dual TCR T cell types and

the single/dual TCR T cell ratios according to tissue origin; classify and calculate the

proportions of all corresponding sequences in the 'β-V' and 'β-J' columns of the single

and dual TCR T cell tables, and use SPSS software to conduct differential VJ usage

analysis between single/dual TCR T cells.

5.CDR3 diversity and clonality analysis: Use the inverse Simpson's index to assess

TCR Treg cell diversity and perform statistical analysis; define clonal expansion as

clone count$\geqslant$2, and analyze the proportion of clonal expansion of single/dual TCR

Treg cells in the two data sets.

6. CDR3 overlap analysis: The TRA CDR3 AA and TRB CDR3 AA of single TCR

Treg cells and dual TCR Treg cells from different tissue sources were compared for

overlap using the immunarch R package, and the Jacard index was used to analyze the

frequency of overlap of CDR3 AA between each tissues pair,; the first 10

high-frequency overlapping TRB CDR3 AA and TRA CDR3 AA sequences were

traced and analyzed between single TCR Treg cells and dual TCR Treg cells from

different tissue sources.

7. **Combined scRNA+TCR-seq analysis:** The scRNA-seq results of the two datasets

were quality-controlled to exclude data with nFeature >5,000, nCount <3,000

and >20,000, and percent.MT >5 (skin, spleen, iLN and mLN tissue datasets) or 10

(blood, kidney, liver, LPL and pancreas tissues). Subsequently, the "DoubletFinder"

R package was used to remove doublets from four tissue sources (including skin), and

doublets/negatives were eliminated from five tissue sources (including LPL) based on

HTO labels. T cells were then clustered and analyzed from different tissue sources

after quality control. The scTCR-seq results were analyzed in conjunction with the

scRNA-seq results, and only Treg cells meeting both criteria were included. Cells

containing scTCR-seq and scRNA-seq data were displayed in UMAP plots and

analyzed for the proportion of single TCR Treg cells and dual TCR Treg cells from

different tissue sources. In this analysis, skin tissue was divided into Treg and

non-Treg for comparison.

8.Cytokine, cytokine receptor, and transcription factor expression analysis:

Comparative analysis of the homogeneity of Treg cell signature molecules expressed

in single TCR Treg and dual TCR Treg; Comparative analysis of the heterogeneity of

the expression of the top 10 mRNA molecules of dual TCR Treg in different tissue

origins; Comparative analysis of the heterogeneity of the expression of single TCR

Treg and dual TCR Treg top 10 mRNA molecules in each tissue; Comparative

analysis of the heterogeneity of the expression of total single TCR Treg and total dual

TCR Treg top 10 mRNA molecules.

REFERENCES

[1] Sakaguchi S, Mikami N, Wing J B, et al. Regulatory T Cells and Human
Disease[J]. Annual Review of Immunology, 2020, 38: 541–566.
[2] Shevyrev D, Tereshchenko V. Treg Heterogeneity, Function, and Homeostasis[J].
Frontiers in Immunology, 2020, 10: 3100.
[3] Malissen M, Trucy J, Letourneur F, et al. A T cell clone expresses two T cell
receptor alpha genes but uses one alpha beta heterodimer for allorecognition and
self MHC-restricted antigen recognition[J]. Cell, 1988, 55(1): 49–59.
[4] Padovan E, Casorati G, Dellabona P, et al. Expression of Two T Cell Receptor α
Chains: Dual Receptor T Cells[J]. Science, American Association for the
Advancement of Science, 1993.
[5] Hinz T, Weidmann E, Kabelitz D. Dual TCR-Expressing T Lymphocytes in
Health and Disease[J]. International Archives of Allergy and Immunology, 2001,
125(1): 16–20.
[6] He X, Janeway C A, Levine M, et al. Dual receptor T cells extend the immune

repertoire for foreign antigens[J]. Nature Immunology, 2002, 3(2): 127–134.

[7] Nj S, Ba B. Dual TCR T Cells: Identity Crisis or Multitaskers?[J]. Journal of

immunology (Baltimore, Md. : 1950), J Immunol, 2019, 202(3).

[8] Zhu L, Peng Q, Li J, et al. scRNA-seq revealed the special TCR β & α V(D)J

allelic inclusion rearrangement and the high proportion dual (or more)

TCR-expressing cells[J]. Cell Death & Disease, Nature Publishing Group, 2023,

14(7).

[9] Yuanyuanxu, Qipeng, Qingqingma, et al. scRNA + TCR-seq revealed the dual

TCR pTh17 and Treg T cells involvement in autoimmune response in ankylosing

spondylitis[J]. International Immunopharmacology, 2024, 135: 112279.

[10] Papalexi E, Satija R. Single-cell RNA sequencing to explore immune cell

heterogeneity[J]. Nature Reviews. Immunology, 2018, 18(1): 35–45.

[11] Park J-E, Botting R A, Domínguez Conde C, et al. A cell atlas of human thymic

development defines T cell repertoire formation[J]. Science (New York, N.Y.),

2020, 367(6480): eaay3224.

[12] Burton O T, Bricard O, Tareen S, et al. The tissue-resident regulatory T cell pool

is shaped by transient multi-tissue migration and a conserved residency

program[J]. Immunity, 2024, 57(7): 1586-1602.e10.

[13] Nedwed A S, Helbich S S, Braband K L, et al. Using combined single-cell gene

expression, TCR sequencing and cell surface protein barcoding to characterize

and track CD4+ T cell clones from murine tissues[J]. Frontiers in Immunology,

2023, 14: 1241283.

[14] Papadopoulou M, Sanchez Sanchez G, Vermijlen D. Innate and adaptive γδ T

cells: How, when, and why[J]. Immunological Reviews, 2020, 298(1): 99–116.

[15] Simon M, Stüve P, Schmidleithner L, et al. Single-cell chromatin accessibility

and transposable element landscapes reveal shared features of tissue-residing

immune cells[J]. Immunity, 2024, 57(8): 1975-1993.e10.

[16] Yang L, Jama B, Wang H, et al. TCRα reporter mice reveal contribution of dual

TCRα expression to T cell repertoire and function[J]. Proceedings of the

National Academy of Sciences of the United States of America, 2020, 117(51):

32574–32583.

Acknowledgements：

We would like to express our gratitude to ***GEO*** and ***IMGT*** databases for providing the availability of the data. Additionally, We thank Oliver T et al. and Nedwed et al. for carrying out the innovative study of Treg in mice and for sharing all original data.

Funding：

This study was supported by the National Natural Science Foundation of China

(82471630&82160279) and the Guizhou Provincial Hundred-level Talent Fund
[No.(2018)5637].

Authorship Contributions：

Xinsheng Yao completed the experimental design and paper writing, Yuanyuanxu，

Qipeng, Xiaoping Lu, completed the data analysis and chart making.

Conflict of interest:

The authors declare no conflict of interests.

Figure 1. Characterizes single Treg TCR and dual TCR Treg in the Blood, Kldney,
Liver, LPL, and Pancreas of mice. **A,** Mouse source names, GEO accession number,
total number of paired TCR Treg cells, and number and proportion of TCR paired
types. **B,** Four different TCR pairing types, VDJ gene family names and CDR3 AA
sequences in single T cells. **C,** Proportion of TCR pairing types per mouse. **D,** Statistical
analysis of dual TCR Treg cells and the relative proportion of single /dual TCR Treg
Cells in Three TCR Pairing Types. **E,** Comparative analysis of single/dual TCR Treg
cell V gene usage variation. **F,** Comparative analysis of single/dual TCR Treg cell J
gene usage variation. **G,** Differential analysis of single/dual TCR Treg cell diversity
among tissues. **H,** Differential analysis of single/dual TCR Treg cell clone expansion
between tissues.

Figure 2. Comparative analysis of single Treg TCR and dual TCR Treg CDR3 overlap and gene mRNA expression in mice Blood, Kidney, Liver, LPL, and Pancreas.**A,**Clonal overlap analysis and jarcard index of TRA/B-chain CDR3 region in single /dual TCR Treg cells.**B,**Tracking results of the first 10 high-frequency overlapping sequences in the CDR3 region of the TRA/B chain of single /dual TCR Treg cells between tissues.**C,**Treg cell clustering results and proportion of Treg cells of the four TCR pairing types among tissues.**D,**Characterized mRNA expression in single/dual TCR Treg cells.**E,**Top 10 mRNA molecules highly expressed by dual

TCR Treg cells in each tissue.**F,**Top 10 mRNA molecules highly expressed by single/dual TCR Treg cells in each tissue and overall.

Figure 3. Characterizes single Treg TCR and dual TCR Treg in the iLN, mLN, skin, spleen of mice.**A,**Tissue source names, GEO accession number, total number of paired TCR T cells, and number and proportion of TCR paired types.**B,**Four different TCR pairing types, VDJ gene family names and CDR3 AA sequences in single T cells.**C,**Proportion of TCR pairing types per tissue.**D,**Statistical analysis of dual TCR T cells and the relative proportion of single /dual TCR T cells in three TCR pairing types.**E,**Comparative analysis of single/dual TCR T cell V gene usage variation.**F,**Comparative analysis of single/dual TCR T cell J gene usage variation.**G,**Differential analysis of single/dual TCR T cell diversity among

 tissues.**H,**Differential analysis of single/dual TCR T cell clone expansion between

 tissues.

Figure 4. Comparative analysis of single Treg TCR and dual TCR Treg CDR3 overlap and gene mRNA expression in mice iLN,mLN,skin,spleen.**A,**Clonal overlap analysis and jarcard index of TRA/B-chain CDR3 region in single /dual TCR T cells.**B,**Tracking results of the first 10 high-frequency overlapping sequences in the CDR3 region of the TRA/B chain of single /dual TCR T cells between tissues.**C,**Treg cell clustering results and proportion of Treg cells of the four TCR pairing types

422   among tissues.**D,**Characterized mRNA expression in single/dual TCR Treg

423   cells.**E,**Top 10 mRNA molecules highly expressed by dual TCR Treg cells in each

424   tissue.**F,**Top 10 mRNA molecules highly expressed by single/dual TCR Treg cells in

425   each tissue and overall.

A

sample	GEO accession		Total cells/paired TCR cells	Single TCR T cells(A+B)/proportion%	Dual TCR T cells/proportion%			
	scRNA-seq	scTCR-seq			Total	A+B1+B2	B+A1+A2	A1+A2+B1+B2
mouse1	GSM8240877	GSM8240878	2598/2072	1683/81.23%	389/18.77%	120/5.79%	218/10.52%	51/2.46%
mouse2	GSM8240880	GSM8240881	3198/2672	2202/82.41%	470/17.59%	115/4.30%	281/10.52%	74/2.77%
mouse3	GSM8240883	GSM8240884	3313/2626	2233/85.03%	393/14.97%	96/3.66%	225/8.57%	72/2.74%
mouse4	GSM8240886	GSM8240887	2140/1610	1398/86.83%	212/13.17%	63/3.91%	126/7.83%	23/1.43%