Characterization and modulation of human insulin degrading enzyme conformational dynamics to

control enzyme activity

Jordan M. Mancl[1], Wenguang G. Liang[1,5], Nicholas L. Bayhi[2], Hui Wei[3,6], William Budell[3], Joshua H.

Mendez[3], Tobin R. Sosnick[4], Bridget Carragher[3,7], Clinton S. Potter[3,7] and Wei-Jen Tang[1]*

1. Ben-May Institute for Cancer Research, The University of Chicago, 929 East 57th Street, Chicago, Illinois

60637, USA

2. Graduate Program in Biophysical Science, The University of Chicago, 929 East 57th street, Chicago,

Illinois 60637, USA

3. Simons Electron Microscopy Center, New York Structural Biology Center, 89 Convent Ave, New York,

New York, 10027, USA

4. Department of Biochemistry and Molecular Biology, The University of Chicago

5. Current affiliation - Bayer HealthCare Pharmaceuticals LLC, St Louis, Missouri, USA

6. Current affiliation - Rutgers New Jersey Medical School, Newark, New Jersey, USA

7. Current affiliation - Chan Zuckerberg Institute for Advanced Biological Imaging, Redwood City, California,

USA

* Corresponding author

Email: wtang@uchicago.edu

Author Contributions: W.-J.T., J.M.M., W.G.L. designed the project. For the molar ratio of IDE to insulin

of 2 to 1 cryo-EM analysis, W.G.L. performed protein purification, cryo-EM grid preparation, data acquisition

and processing assisted by H.W. and overseen by W.-J.T., B.C. and C.S.P. J.M.M. and W.G.L. built and

refined structural models. J.M.M. performed heterogeneity analysis of cryo-EM structures, MD simulations,

and mutant characterizations. For time-resolved CryoEM analysis, J.M.M. performed protein purification,

W.B. performed cryo-EM grid preparation, J.H.M performed data acquisition, and J.M.M. performed data

processing and model building. N.L.B performed the Upside simulations and analysis and initial all-atom

MD simulations under the guidance of T.R.S. J.M.M. W.-J. T. wrote the manuscript and all authors

contributed to manuscript finalization.

Acknowledgement: We would like to thank our colleagues at the University of Chicago: Adam Antoszewski

and Aaron Dinner for providing us with the data from their insulin unfolding simulations, as well as Rong

Shen for general guidance with the MD simulations, and Jessica Martinez Martinez for assistance with

cloning the IDE R668A construct. We would also like to thank James Byrnes at the LiX beamline at

Brookhaven National Laboratory for collecting SEC-SAXS data. Computer resources came, in part, from

an allocation on the Beagle3 computing cluster at the University of Chicago. This work was supported by

the NIH grant GM121964 to W.-J. Tang. The Simons Electron Microscopy Center is supported by a grant

from the Simons Foundation (SF349247). Use of the Advanced Photon Source was supported by the U.S.

Department of Energy, Office of Basic Energy Sciences, under contract No. DE-AC02-06CH11357. The

LiX beamline is supported by NIH grants P30GM133893 & S10 OD012331, DOE grants KP1605010 & DE-
SC0012704.

Competing Interest Statement: None.

Keywords (5 max): amyloid peptides, insulin degrading enzyme, proteostasis, diabetes, cryo-electron
microscopy

Abstract

Insulin degrading enzyme (IDE) is a dimeric M16A zinc metalloprotease that degrades amyloidogenic peptides diverse in shape and sequence, including insulin and amyloid-β, to prevent toxic amyloid fibril formation. IDE has a hollow catalytic chamber formed by two ~55 kDa N- and C- domains (IDE-N and IDE-C, respectively), in which peptides bind, unfold, and are repositioned for proteolysis. IDE is known to transition between a closed state, poised for catalysis, and an open state, able to release cleavage products and bind a new substrate. Here, we present six cryo-EM structures of the IDE dimer at 3.0-5.1 Å resolution, obtained in the presence of a sub-saturating concentration of insulin. Combining cryo-EM heterogeneity analysis with all-atom molecular dynamics (MD) simulations, we identified the structural basis and key residues for IDE conformational dynamics that were not previously revealed by IDE static structures. Notably R668 serves as a molecular latch mediating the open-close transition and facilitates key protein motions through charge-swapping interactions at the IDE-N/C interface. Our small-angle X-ray scattering analysis and enzymatic assays of an R668A mutant indicate a profound alteration of conformational dynamics and catalytic activity. By integrating coarse-grained MD simulations, our analysis reveals that IDE unfolds its substrates through the coordinated motion between IDE-N and IDE-C, as well as β-sheet formation between IDE and insulin. Additionally, our time-resolved cryo-EM analysis uncovers IDE allostery within the IDE dimer. Collectively, our findings demonstrate the strength of combining experimental and computational approaches to probe protein dynamics and pave the way for developing substrate-specific modulators of IDE activity.

Introduction

Protein homeostasis (a.k.a. proteostasis) is maintained by three primary mechanisms: chaperones,

ubiquitination/proteasome, and autophagy (Hipp *et al*, 2019; Lopez-Otin *et al*, 2013). Disruptions in

proteostasis can lead to the accumulation of amyloid fibrils and subsequent human diseases (Chiti &

Dobson, 2006; Chiti & Dobson, 2017; Greenwald & Riek, 2010). As a result, many proteases have evolved

to specifically target amyloidogenic peptides (Malito *et al*, 2008; Saido & Leissring, 2012). Among those,

the cryptidase family, which includes the M16 metalloproteases insulin degrading enzyme (IDE) use an

internal catalytic chamber, or "crypt", to capture and selectively degrade the monomeric form of amyloid

peptides to control the formation of amyloid fibrils (Liang *et al*, 2022; Malito *et al.*, 2008). IDE effectively

degrades various bioactive peptides, including amyloid-β (Aβ), a peptide associated with the progress of

Alzheimer's disease, and three blood glucose-regulating hormones: insulin, amylin, and glucagon (Tang,

2016). Consequently, defects in IDE alter the progression of type 2 diabetes mellitus and Alzheimer's

disease in animal models and are linked to these diseases in humans (Bertram *et al*, 2000; Bertram *et al*,

2007; Farris *et al*, 2003; Farris *et al*, 2004; Kim *et al*, 2007; Sladek *et al*, 2007; Tang, 2016; Tanzi & Bertram,

2005; Zeggini *et al*, 2007). IDE is a promising therapeutic target, as its inhibition improves glucose

tolerance, yet progress has been hampered by the contrary actions of its diverse substrate pool (Maianti *et*

al, 2014; Maianti *et al*, 2019; Tang, 2016). For example, IDE overexpression has been shown to reduce Aβ

loads, in mice, however, this reduction is accompanied by hyperglycemia from a similar reduction in insulin

levels (Leissring *et al*, 2003).

IDE-mediated substrate recognition and degradation is a complex process. IDE is capable of

degrading substrates with diverse sequences and structures, such as Aβ and the three glucose-regulating

hormones: insulin, amylin, and glucagon (Malito *et al.*, 2008; Tang, 2016). Despite this diversity, IDE

exhibits high selectivity—for example, it preferentially degrades insulin and atrial natriuretic peptide over

insulin-like growth factor-1 and brain natriuretic peptide, respectively (Malito *et al.*, 2008). The presence of

ATP, inositol phosphate, or specific mutations can markedly enhance the catalytic rate for small peptide

substrates like bradykinin, but does not significantly affect the degradation of larger substrates such as

insulin and Aβ (Im *et al*, 2007; McCord *et al*, 2013; Song *et al*, 2017; Song *et al*, 2004). Additionally, the

binding location of IDE inhibitors within the catalytic chamber can influence substrate preference, making

IDE more likely to degrade one substrate over another (Charton *et al*, 2014; Maianti *et al.*, 2019). This

complexity arises partly from the interplay between the unfoldase and protease activities of IDE, which are

essential for degrading diverse substrates (Malito *et al.*, 2008; Tang, 2016; Zhang *et al*, 2018). IDE relies

on its unique large catalytic chamber and associated conformational dynamics to unfold and degrade

amyloidogenic substrates. A deeper understanding of the molecular mechanisms by which IDE utilizes its

catalytic chamber for substrate unfolding and degradation is crucial for effectively harnessing this enzyme

in future therapeutic development.

IDE is a dimeric 110 kDa metalloprotease. Each protomer is comprised of four homologous

subdomains organized into two ~55 kDa domains, IDE-N and IDE-C, connected by a short flexible linker

(Li *et al*, 2006; Shen *et al*, 2006) (Fig. 1A). Our previous cryo-EM analysis revealed the major conformational

states of IDE. In the absence of substrate, each protomer was found to adopt an open (O) or partial open

(pO) state which are primarily differentiated by the displacement of IDE-N relative to IDE-C, the combination

of which led to three dimeric conformations (O/O, O/pO, pO/pO) (Zhang *et al.*, 2018). It was observed that

substrate binding induced both protomers of IDE to fully close, referred to as the partial closed (pC, pC/pC

for the dimer) state, as the cryo-EM closed state was found to be slightly more open than previously solved

crystal structures, likely influenced by the constraints of the crystal lattice (McCord *et al.*, 2013; Shen *et al.*,

2006). Only the O state permits substrate access to the catalytic chamber, yet the catalytic cleft is stabilized

only in the pC state with substrate bound, requiring IDE to undergo a substantial open-close transition

during catalysis, the specifics of which are unknown (Zhang *et al.*, 2018).

Biochemical and structural analyses indicate that many IDE substrates, such as insulin, must first

be unfolded prior to cleavage (Fig. 1B, C). The conformational dynamics of IDE are thought to play a central

role in substrate recognition and degradation, yet several mechanistic details underlying the unfoldase

activity of IDE remain unresolved (Fig. 1). We hypothesize that the unfoldase and protease activities of IDE

are regulated by the relative motions of the IDE-N and IDE-C domains, their interactions with substrates,

and allosteric communication between subunits within the IDE dimer. To investigate this, we solved cryo-

EM structures of IDE bound to insulin at a 2:1 IDE:insulin ratio, as well as structures captured following

rapid mixing of IDE with insulin under functionally relevant conditions. Utilizing newly developed

computational methods, we analyzed particle heterogeneity in these structures to infer the conformational

dynamics of IDE. By integrating these experimental data with all-atom and coarse-grained molecular

dynamics (MD) simulations, we identified and modulated specific molecular interactions to manipulate key

conformational dynamics of IDE, ultimately enhancing our understanding how to control its enzymatic

activity in vitro.

Results

Cryo-EM reveals new IDE structures in the presence of a sub-saturating concentration of insulin

We have previously used cryo-EM to solve the structure of IDE in the presence of a 5-fold molar

excess of insulin, resulting in a dominant pC/pC state with both substrate binding chambers yielding density

corresponding to unfolded insulin (Zhang *et al.*, 2018). To explore the mechanism of IDE-directed substrate

unfolding under more physiologically relevant conditions, we investigated IDE-insulin interactions with an

IDE:insulin molar ratio of 2:1. We collected a dataset of ~7,600 micrographs on a Titan Krios operated at

300 keV, from which 7.2 million particles were picked for processing in RELION. Following an established

workflow, and processing the data with C1 symmetry, we generated five structures: three previously

observed structures with improved resolution (pC/pC, O/O, and O/pO at 3.0 Å, 3.8 Å, and 4.1 Å resolution)

and two novel states (O/pC and pO/pC at 3.4 Å and 3.3 Å resolution, respectively) (Fig. 2A, Figure 2 –

figure supplement 1, Supplementary file 1). Interestingly, the pC/pC was still found to be dominant, despite

IDE and insulin being present at a 2:1 molar ratio, consistent with the allostery of IDE (McCord *et al.*, 2013;
Zhang *et al.*, 2018).

The individual protomers of IDE adopt the same three conformations: O, pO, and pC states,
reported previously but there are significant differences in the O and pO states (Fig. 2B) (Zhang *et al.*,
2018). While the pC/pC state is nearly identical to the two states reported previously (RMSD of ~0.6 Å; PDB
ID: 6BFC, 6B3Q) (Zhang *et al.*, 2018), the O and pO states appear more closed in the presence of insulin,
suggesting that the presence of insulin promotes the open to closed transition. Specifically, the pO and O
subunits are 3-5 Å more closed based on the distance between the centers-of-mass (COM) of the IDE D1
and D4 subdomains, while the dihedral angle formed by the COM of the D1-D2-D3-D4 subdomains is
reduced by 5-10° (Fig. 2B). Alignment of the new O/O and O/pO states to their previously solved
counterparts reveals little difference, with a global RMSD of 1.316 Å (O/O, PDB ID: 6B7Y) and 1.405 Å
(O/pO, PDB ID: 6BF8), respectively (Zhang *et al.*, 2018). The primary source of variation between the
different states remains the degree of opening between the two domains of each subunit, as a global
alignment of IDE-N and IDE-C across all structures reveal the domains primarily act as rigid bodies (RMSD
<1.4 Å). Consistent with previous observations, the door region, which contains the catalytic zinc binding
site, exhibited higher B-factors in the O states than in either the pO or pC state (Zhang *et al.*, 2018) (Figure
2 – figure supplement 2).

The cryo-EM structures presented here contain four subunits that have clear density present at the
catalytic cleft and exosite indicative of bound substrate, thus adopting the pC state. Previous structures
have modeled the same chain of insulin in both the exosite and catalytic cleft, but the best models we could
build into our density suggest that the chain bound to the exosite is not the same chain positioned for
cleavage at the catalytic cleft (Fig. 2C). The preponderance of data suggests it to be unlikely that insulin
adopts a preferred orientation within the catalytic chamber, rather the catalytic chamber can accommodate
insulin in four different orientations, with either the A or B chain bound to the exosite and catalytic cleft in
cis and *trans* orientations. This promiscuity of binding has been theorized as a reason for why there is no
chain preference for the initial insulin cleavage event (Manolopoulou *et al*, 2009). Crystal structures of
insulin are highly compact and globular, with little spatial separation between the N- and C- termini of each
chain, and the IDE cleavage sites are located within α-helices (Baker *et al*, 1988). In its crystallized state,
insulin cannot interact with both the exosite and catalytic cleft unless it unfolds, at least partially. Recent
work used MD simulations to study the conformational dynamics of insulin in solution and identified several
major "elements of disorder" to describe the partially unfolded intermediate structures they observed (Busto-
Moner *et al*, 2021). We docked these structures of insulin into the closed catalytic chamber of IDE and
found that the chamber was able to easily accommodate the positioning of either chain near the exosite
and catalytic cleft in *cis* and *trans* orientations (Figure 2 – figure supplement 3).

Heterogeneity analysis reveals 2 dominant components of structural variability with the IDE cryo-
EM particle population

Numerous structures of IDE exist with subunits adopting either an open or closed conformation,
yet we lack an understanding of how IDE transitions from an open to a closed state. The simplest
explanation of this transition, based on structural data, would be a direct, rigid body translation of IDE-N
relative to IDE-C, and this is the predominant model within the field (Shen *et al.*, 2006; Zhang *et al.*, 2018).
To better understand what this transition would look like, we measured and plotted the changes in the D1-
D4 COM distance versus the changes in the D1-D2-D3-D4 COM dihedral angle for all available cryo-EM
structures of IDE (Figure 3 – figure supplement 1). We found that in the absence of substrate, the transition
pathway produced a linear relationship between states. However, when our structures generated in the
presence of insulin were analyzed, the linear relationship no longer held (Figure 3 – figure supplement 1),
suggesting that the simple linear transition typically presumed for the open-close transition between distinct
IDE states does not accurately depict the complexity of IDE dynamics.

Recently, several approaches have been developed to understand the conformational
heterogeneity present within cryo-EM data. We employed multibody analysis in RELION and 3D variability
analysis (3DVA) in cryoSPARC to investigate the conformational heterogeneity within our particle
populations (Nakane *et al*, 2018; Punjani & Fleet, 2021). Of these, the range of motion predicted by
multibody analysis most closely matched the reported structures while 3DVA exhibited consistent results
across a smaller magnitude of transition (Figure 3 – figure supplements 1-3). Therefore, we focused on
multibody analysis below. Multibody analysis models the structural heterogeneity in our data as the result
of motions of independent, user-defined rigid bodies. As discussed above, the primary source of structural
variation in our structures is the positioning of IDE-N relative to IDE-C. We see little structural changes
within the domains, suggesting that IDE domains behave as three rigid bodies, IDE-N in two different
subunits within IDE dimer (IDE-N A, IDE-N B) and the dimeric IDE-C (IDE C/C). This is consistent with the
structural analysis of M16 family of metalloproteases (Liang *et al.*, 2022). This assertion is further supported
by our all-atom molecular dynamics (MD) simulations. While we observed small yet noticeable
conformational changes within IDE-N A, IDE-N B, and IDE-C/C, the overall RMSD of entire IDE dimer is
significantly higher than those three individual units (Fig. 3A). Multibody analysis defines the principal
components of variance along discrete eigenvectors, which we interpret as proxies for the dominant
components of molecular motion. We quantified the change in D1-D4 COM distance and D1-D2-D3-D4
COM dihedral of representative structures along the heterogeneity gradients of the top 9 eigenvectors for
each of our IDE structures, representing ~70-90% of the total structural heterogeneity per structure. The
results indicated that the particles comprising our cryo-EM structures sampled a significantly greater
conformational space than would be expected from analysis of the ensemble structures alone (Figure 3 -
figure supplement 2). Most intriguingly, the conformational changes displayed an unexpectedly high degree
of change in the D1-D2-D3-D4 COM dihedral angle, indicating a significant rotation of IDE-N relative to
IDE-C (Fig. 3B,C, Figure 3 – figure supplement 2).

Integration of the multibody results across all states of IDE suggest two dominant components of
structural variance. First is a translation-dominant conformational change wherein IDE-N swings toward or

away from IDE-C as if mediated by a hinge formed by the interdomain linker region of IDE (Fig. 3D, Figure
3 - video 1). This conformational change closely resembles our understanding of the IDE open-closed
transition inferred from analysis of the ensemble structures. The second component is a rotation-dominant
conformational change wherein IDE-N rotates orthogonal to the plane of the dimer as if it were being
screwed into or ground against IDE-C (Fig. 3E, Figure 3 – video 2). Importantly, these dominant
components of structural variance were also observed when the particle populations were analyzed with
3DVA in cryoSPARC, albeit over a smaller magnitude of conformational change (Figure 3 – figure
supplement 3). These dominant components also correlate well with the lowest frequency modes revealed
by normal mode analysis (Figure 3 – videos 3, 4). Interestingly, while the presence of insulin was found to
substantially influence the consensus reconstructions of IDE, we found that the presence or absence of
insulin had no significant effect on the principal components of structural variance for IDE, consistent with
the current understanding that enzyme conformational changes are "hardwired" into the structure (Bahar
et al, 2007; Bahar & Rader, 2005).

R668 plays a key role in the IDE open/close transition in all-atom MD simulations

Analysis of the particle heterogeneity comprising cryo-EM structures provides excellent information
about the conformational space sampled by the protein of interest. However, such analysis lacks a temporal
component, and thus only implies motions. To overcome this barrier and examine how the conformation of
IDE changes over time, we performed six replicate all-atom MD simulations with explicit solvent.
Simulations were initialized from the O/pO state (PDB: 7RZG), with the missing loops modeled in to
generate continuous peptide chains and run for one microsecond each. By starting with the O/pO structure,
we investigated the dynamics of both the open and closed subunits simultaneously. We observed that the
conformational space sampled by our simulations correlated well with the conformational space sampled
by our cryo-EM particle population, as revealed by multibody analysis (Figure 4 – figure supplement 1). In
five of our six simulations, the open subunit closed quickly, typically in <200 ns (Fig. 4A). Analysis of the
open-close transition revealed two key findings. First, the open subunits did not close to a singular
consensus structure. We define closing as a subunit reaching a similar D1-D4 COM distance as the pO or
pC states, but we observed distinct differences among the subunits indicative of IDE-N rotation relative to
IDE-C. These differences are in line with our second key finding, that the conformational changes
associated with the open-close transition did not follow the same general pathway among the simulations,
as we observed a high degree of variability upon closing (Fig. 4B).

Noticeably, R668 consistently stood out as one of the first IDE-C residues to form new interactions
with IDE-N, either D309 or E381 despite the path variation among our MD simulations (Fig. 4C). In most
simulations, formation of this interaction preceded a large decrease in the D1-D4 COM distance (Fig. 4D,
Figure 4 – figure supplement 2). Interestingly, these R668 interactions are not present in the ensemble
cryo-EM or crystal structures of IDE. We hypothesize that the biochemical properties and spatial localization
of R668 enable it to essentially reach out and grab onto the N-domain, initiating the formation of a complex
hydrogen bonding network that drives the open-close transition towards completion. Previously published

hydrogen-deuterium exchange mass spectrometry (HDX-MS) data indicates that R668 is stabilized under
conditions that promote IDE closing, suggesting that R668 may function as a molecular latch (Fig. 4E)
(Zhang *et al.*, 2018).

To further probe the importance of R668, we ran 6 additional simulations with IDE carrying a R668
to alanine (R668A) point mutation under identical conditions as described previously. These simulations
displayed significantly altered dynamics when compared to the wild-type simulations. Notably, the R668A
mutant was found to close more slowly, and many simulations did not reach the D1-D4 COM distance
typified by a closed state structure, instead stabilizing with a significantly larger D1-D4 COM distance than
we had previously observed in our wild type (WT) simulations or our experimental structures (Fig. 4F). We
also observed that the R668A mutant preferentially sampled a subset of conformational space rarely
explored in our WT simulations and displayed significantly greater variation in the D1-D2-D3-D4 dihedral
angle (Fig. 4G). As a result of this increased rotational exploration, several of our R668A simulations closed
to a conformation where IDE-N is offset and rotated relative to IDE-C when compared to the experimentally
determined structures (Figure 4 – figure supplement 3). This altered conformation results in a structure
where IDE-N and IDE-C are fully engaged, but the offset produces a deceivingly high D1-D4 COM distance.
With this in mind, we sought to better understand the conformational dynamics of IDE in the closed state.

R668A modulates IDE conformational dynamics *in vitro*

The R668A mutation was found to significantly alter the conformational dynamics of the IDE open-
close transition in our simulations (Fig. 4F, G), so we expressed and purified the R668A mutant to determine
if the changes predicted from our MD simulations resulted in altered enzymatic activity relative to WT.
Previous work has established that IDE adopts a dominant O/pO state in solution (Zhang *et al.*, 2018).
Consistent with our MD simulations, size-exclusion chromatography revealed that the R668A construct
eluted slightly earlier than WT IDE, suggesting that the R668A mutation induces a greater proportion of
molecules to adopt a more open conformation, possibly shifting to a dominant O/O state (Fig. 5A). Next,
we compared the enzymatic activity of the R668A and WT constructs using the fluorogenic substrate (7-
methoxycourmarin-4-yl)acetyl-RPPGFSAFK(2,4-dinitrophenyl)-OH (substrate V), a bradykinin mimetic
which has previously been used to characterize the enzymatic activity of IDE and related enzymes (King *et
al*, 2014; McCord *et al.*, 2013). We found that WT IDE has activity consistent with previously published data
(McCord *et al.*, 2013) and the R668A mutation produced a ~5-fold decrease in activity (Fig. 5B). ATP is
known to promote IDE degradation of small substrates, i.e. bradykinin, but not larger substrates, i.e. insulin
(33). Consistent with this observation, the addition of ATP increased the activity of both WT and R668A
constructs (Fig. 5B). We also tested if the R668A mutation would alter the degradation of larger, well-folded
substrates, using insulin in a direct competition assay. We found that, after accounting for the previously
observed 5x decrease in activity, insulin yielded an apparent K_i of ~8 nM for WT IDE, but this value was
reduced to ~52 nM for the R668A construct (Fig. 5C). When Aβ(1-40), an unstructured substrate, was
tested in the same competition assay, we observed an apparent K_i of ~688 nM for WT IDE, yet the R668A
mutant did not exhibit a dose-dependent response to the presence of Aβ and the data could not be fit to

any kinetic equation (Fig. 5D). We then performed size exclusion chromatography-coupled small-angle X-
ray scattering (SEC-SAXS) for both constructs to ascertain if the observed differences in enzymatic activity
could be explained by altered biophysical properties. The WT scattering profile produced an R_g of 49.4 +/-
0.9 Å, consistent with previous SEC-SAXS results (Fig. 5E, F) (Zhang *et al.*, 2018). The R668A scattering
profile produced an R_g of 54.7 +/- 0.4 Å, consistent with our SEC data that R668A mutant has slightly larger
hydrodynamic radius (Fig. 5A). Together, our data indicates that R668A mutation can profoundly affect the
conformational dynamics governing the open-closed transition, and that altered conformational dynamics
can alter enzymatic activity in a substrate-dependent manner.

287   Charge-swapping at the IDE-N/C interface mediates conformational dynamics in the closed state

The WT pO subunit remained closed throughout all simulations yet was found to initially relax to a
slightly more open state within the first 200 ns, roughly the same time frame it took the WT O subunit to
close (Fig. 6A) and, despite not opening, demonstrated a range of D1-D2-D3-D4 dihedral variation similar
to that of the O subunits (Fig. 6B). Interestingly, the conformational variation within our simulations
increased greatly after this relaxation point was reached (Fig. 6A). This relaxed conformational state
consistently displayed an altered IDE-N/C interface compared to the experimentally determined structures
that was replicated by the open WT subunits upon closing, leading to a nearly symmetric pO/pO state
(Figure 6 – figure supplement 1). Biochemically, this relaxed IDE-N/C interface makes more sense than the
interface observed experimentally. It has previously been demonstrated that the experimental constraints
of crystallographic structure determination force IDE to adopt a closed conformation and determination of
the open state can only be accomplished once those restraints are removed, i.e. cryo-EM (Zhang *et al.*,
2018). These results suggest that both ensemble methods of structure determination may impact the
conformation of IDE, however slightly. The IDE-N/C interface is littered with charged residues, yet few
interactions are observed in the ensemble structures (Fig. 6C), whereas our MD simulations reveal a
complex hydrogen bonding network (Fig. 6D). We observed an extensive network at the D2-D3 interface,
with sporadic patches of interactions between D1 and D4 (Figure 6 – figure supplement 2). Within the D2-
D3 network R668 again stands out as a notably key residue. In addition to forming hydrogen bonds with
D309 and E381, we observed R668 form π-π interactions with R311 (Fig. 6E). Arginine-mediated π-
interactions have been well characterized in other systems for their ability to stabilize interaction interfaces
and facilitate conformational changes (Armstrong *et al*, 2016; Vernon *et al*, 2018).

The IDE-N/C interactions we observe in our simulations are not restricted to discrete cognate pairs;
rather, we observed that residues within the network periodically swap among several interaction partners.
This "charge-swapping" phenomenon allows IDE-N to rotate or slide against IDE-C to adopt multiple
conformations and maintain favorable interacting contacts (Fig. 6F). While multiple, concurrent, instances
of charge-swapping were associated with rapid changes in the D1-D2-D3-D4 dihedral angle, singular
events of charge-swapping did not appear to be significant enough to stimulate large-scale conformational
change (Fig. 6G, H). Instead, substantial conformational changes appear to be caused by multiple charge-
swapping events occurring in conjunction. This may explain why we observed no noticeable change in the

conformational dynamics of the pO state subunits when comparing the WT and R668A constructs. Consurf
analysis reveals that most of the residues forming these interaction networks are highly conserved among
IDE homologs (Figure 6 – figure supplement 3) (Ben Chorin *et al*, 2020). This indicates that the revealed
rotational motion is likely evolutionarily conserved and offers a potential mechanism by which IDE unfolds
and repositions bound peptide substrates to degrade amyloid peptides.

Time-resolved cryo-EM suggests insulin-induced allostery in IDE

To begin understanding of how the interaction between IDE and insulin leads to insulin unfolding
and degradation, we first performed all-atom MD simulation of insulin-bound IDE in the closed state.
Unfortunately, we did not observe a noticeable structural change of insulin within our microsecond
simulations. Previously, we used time-resolved SAXS analysis to determine the time constant for the insulin
induced open to closed transition, tau (τ, a.k.a., monoexponentially decay) to be ~100 milliseconds (Zhang
et al., 2018). However, it has previously been determined that IDE cleaves about two insulin molecules
every second (Manolopoulou *et al.*, 2009). We employed spotiton to rapidly mix and freeze grids where IDE
has been exposed to insulin for 123 milliseconds prior to freezing (Dandey *et al*, 2020). We hoped that such
an experimental setup would permit the identification of novel insulin-bound states of IDE and offer insight
into the initial binding and unfolding of insulin. Previous work has established that, in the absence of
substrate, IDE adopts a dominant O/pO state in solution and the presence of a molar excess of substrate
shifts the particle population to a dominant pC/pC state (Zhang *et al.*, 2018). Surprisingly, after processing
the dataset in Relion, we observed a single conformational state within our rapidly-mixed dataset: the O/O
state (Fig. 7A, Figure 7 - figure supplement 1, supplementary file 2). We were unable to resolve the structure
beyond moderate resolution (~5.1 Å) and did not observe unambiguous density indicative of insulin binding
within the catalytic chamber, which is unsurprising given our previous observations of promiscuous binding
(Fig. 2C). Despite the relatively high particle number in our final reconstruction, we were unable to sub-
classify distinctly different conformations of IDE within the particle population as initial classes converged
back to the dominant O/O state upon subsequent refinement. This leads us to believe that the obtained
O/O state represents the average state of a particle population that exhibits a high degree of continuous
structural heterogeneity. The O/O state observed in our rapidly-mixed dataset is distinct from the previously
observed O/O states with D1-D4 center of mass distances of 55.1 Å and 46.8 Å matching the largest and
smallest, respectively, distances previously observed for individual O state subunits (Fig. 7B). The
discrepancy between the "openness" of the two subunits nearly doubles that observed in previous
structures, making this O/O state the most asymmetric O/O state yet observed (Fig. 7C). The observation
of a dominant O/O state in the presence of insulin under short timescales suggests that the binding of
insulin to the O subunit of the apo-dominant O/pO state can induce the pO state to open through an
unknown allosteric mechanism to reach the observed dominant O/O state (Fig. 7D). Positive allostery in
IDE has previously been identified in biochemical assays, yet we lack a detailed understanding of the
underlying mechanism (McCord *et al.*, 2013; Song *et al*, 2011).

Coarse-grained MD simulations reveal intermolecular cross-β sheet formation between insulin and
IDE in all four IDE domains

As cryo-EM proved unsuitable to explore the mechanism of insulin unfolding by IDE, we turned to
MD simulations. As stated above, there is a notable discrepancy in the time frame between the open/close
transition and insulin cleavage rate. This suggests that unfolding requires multiple cycles of the open-closed
transition and, thus it is not currently realistic to use all-atom MD simulations to probe the unfolding of insulin
by IDE. We then used a coarse-grained MD package, Upside, to probe the interaction between IDE and
insulin. Upside performs the dynamics simulation for just six atoms per residue with the sidechains being
represented by directional-dependent beads whose packing probabilities have the lowest free energy while
still including key structure details to compute realistic forces (Faruk *et al*, 2022; Faruk *et al*, 2023; Jumper
et al, 2018a, b). This allows rapid equilibration and sampling of the energy landscape when compared with
the conventional all-atom MD simulation, achieving ~10^4 acceleration for the calculation time. Consistent
with all-atom MD simulations, Upside simulations revealed the wide range of motions, including grid and
hinge motions found in our all-atom MD simulation (Figure 8 – figure supplement 1). We observed the
partial unfolding of insulin in our Upside simulations yet, importantly, the unfolding did not disrupt the inter-
or intra-molecular disulfide bonds (Figure 8 video 1).

As insulin partially unfolds, it readily adopts β-strand structures—a key characteristic of
amyloidogenic peptides—which can transiently form cross-β-sheets with the central β-sheet present in all
four domains of IDE (Figure 8A-B). Specifically, we observed that cross-β-sheet interactions occur at
distinct regions of insulin: the N-terminus of insulin chain A interacts with the IDE D2 domain near the
substrate N-terminal anchoring site (also known as the exosite); the middle portion of insulin chain A
interacts with the IDE D1 domain; and the C-terminal regions of insulin chain B interact with the D3 and D4
domains of IDE. While the interactions of insulin with IDE-C domain are novel, those with IDE D1/D2
domains are structurally similar to those previously observed (Guo *et al*, 2010; Manolopoulou *et al.*, 2009;
Ren *et al*, 2010; Shen *et al.*, 2006). The labile and transient β-strand-rich interactions have been shown to
mediate diverse protein-protein interactions, e.g., the assembly of intermediate filaments (Eibauer *et al*,
2024; Zhou *et al*, 2021).

This interaction provides molecular insight into how IDE recognizes and binds amyloidogenic
peptides with diverse sequences (Figure 8C). IDE is capable of forming cross-β interactions with many
regions of insulin, primarily mediated by backbone-to-backbone contacts that do not rely on specific amino
acid sequences. The structural arrangement of the four conserved domains of IDE enables it to interact
with multiple regions of its substrates, such as insulin, which are predisposed to forming β-strands (Figure
8C). We term this type of interaction "β-grabbing," and propose that it is a key mechanism by which IDE
unfolds insulin. This interaction effectively tethers the substrate to both the IDE-N and IDE-C domains,
facilitating the interdomain motions of IDE that promote substrate unfolding.

Discussion

Our results allow us to put forth a refined model describing how IDE recognizes amyloid peptides
that are diverse in size and shape (Fig. 9). The catalytic cycle of IDE starts with at least one subunit of
dimeric IDE adopting the open state, exposing the interior of the catalytic chamber to potential substrates
(Zhang *et al.*, 2018). The unconstrained motion of IDE-N likely facilitates substrate interaction in a variety
of initial orientations. IDE-N and IDE-C have negatively and positively charged surfaces, respectively, and
are able to attract peptide substrates with complementary charge profiles (Shen *et al.*, 2006; Zhang *et al.*,
2018). Such peptides have a high dipole moment and are often prone to aggregation. However, in the open
state, the door subdomain, which contains the key catalytic residues, is highly flexible, rendering the open
state catalytically incompetent (McCord *et al.*, 2013; Zhang *et al.*, 2018). Proper positioning and unfolding
of substrate is necessary to stabilize the cleft prior to catalysis. Substrate binding has been suggested to
enhance IDE closing through charge complementarity (Shen *et al.*, 2006). It is likely that the rotation of IDE-
N permits a range of closing pathways to accommodate non-optimal substrate binding geometry (Fig. 4). If
the closing geometry permits R668 to interact with E381 or D309, the closing reaction continues to
completion. If the closing geometry does not permit the R668-mediated interactions, this could be a signal
of improper substrate binding, allowing IDE-N to open and either release improper substrate or reposition
for another closing attempt. Once closed, IDE-N is capable of rotating against IDE-C mediated by charge-
swapping events at the IDE-N/C interface (Fig. 6). The combined motions associated with the open to
closed transition and rotation between IDE-N and IDE-C can lead to the distortion of the secondary structure
of bound substrate. Furthermore, the transient interaction of insulin with the catalytic chamber of IDE such
as β-grabbing via the β–strand pairing should further stress the integrity of bound substrate (Fig. 8).
Together, these forces should facilitate unfolding and proper positioning of substrate for cleavage. Our data
indicates that IDE is able to accommodate any possible orientation of insulin tethering within the catalytic
site (Fig. 2C), suggesting that IDE is able to cut both chains of insulin in rapid succession without requiring
significant repositioning (Manolopoulou *et al.*, 2009). IDE then opens to release cleavage products and the
cycle begins anew.

IDE utilizes both unfoldase and protease activities to degrade clinically-relevant peptides, including
three glucose-regulating hormones with contrary effects – insulin, amylin, and glucagon – and Aβ, the
accumulation of which is associated with the progression of Alzheimer's disease. This has made IDE an
attractive target for therapeutic intervention (Maianti *et al.*, 2019; Tang, 2016). One major challenge is how
to better control selectivity when the substrates IDE degrades are highly diverse in sequence and structure.
It remains unknown how IDE unfolds substrates prior to cleavage. Our studies characterize the
conformational dynamics of IDE and offer insight into the catalytic mechanisms, revealing that IDE-N rotates
against IDE-C. We envision that this motion works in concert with the open-close transition to reposition
and unfold substrates prior to cleavage. This rotation appears to be largely unconstrained in the open state
and may play a role in promoting substrate capture. In the closed state, we found the rotational motion to
be mediated by charge-swapping events at the IDE-N/C interface. These events are supplemented by
interactions between hydrophobic patches that provide a non-specific interaction interface and allow the
domains to slide against one another. We found that IDE-N/C interactions were mediated by the key residue

R668, and mutation of this residue altered the enzymatic activity and conformation of IDE, likely by impairing

the open-close transition. Our data suggests that, while the R668A mutation globally impaired IDE activity,

R668A exhibits significantly different kinetic profiles towards well-folded substrates (insulin) compared to

unstructured substrates (Aβ). This highlights the complexity of the IDE catalytic cycle as we envision that

IDE conformational dynamics can regulate solvent access to the catalytic chamber and substrate binding,

unfolding, and cleavage are all intimately linked.

Our work adds to the expanding body of literature that integrates computational and experimental

approaches (Arenz *et al*, 2016; Hirschi *et al*, 2021; Jiang & Doudna, 2017; Palermo *et al*, 2016; Roh *et al*,

2020; Saha *et al*, 2022). By employing multibody analysis, we identified the dominant sources of structural

heterogeneity within our cryo-EM particle population, uncovering that the conformational dynamics of IDE

involve a substantial rotational component that is not evident from ensemble structure analysis alone. We

showed that all-atom MD simulations can address limitations of cryo-EM single particle analysis when

investigating the role of residue R668. Because R668 forms a network of polar and electrostatic interactions

rather than exclusive partnerships, dynamic hydrogen bonding networks at the IDE-N/C interface—

particularly those involving R668—contribute to conformational heterogeneity, which further restricts the

achievable resolution in cryo-EM analysis. Furthermore, vitrification by freezing proteins in liquid ethane

can significantly alter the entropic contribution of R668 through conformation-dependent promiscuous

contacts. By combining cryo-EM and MD simulations, we gained insight into how flexible interactions

between IDE-N and IDE-C facilitate domain movements and highlighted the critical role of R668 in

mediating these interactions. We also used MD simulations to demonstrate altered open-closed dynamics

in the R668A mutant compared to the wild-type enzyme and validated these findings with experimental

enzymatic kinetic assays and SEC-SAXS experiments.

At its core, cryo-EM is an ensemble method of structure determination, averaging tens of thousands

of particles together to generate consensus structures. While methodologies have been well-established to

separate out discrete conformational classes, proteins that exhibit constant gradients of structural

heterogeneity remain problematic, although significant attention has been devoted to the issue in recent

years, with varying levels of success (Nakane *et al.*, 2018; Punjani & Fleet, 2021; Zhong *et al*, 2021).

Currently, the best approaches use various dimensionality reduction techniques to analyze the principal

components of particle heterogeneity. While a substantial innovation, it remains to be seen how the principal

components of structural heterogeneity, derived from a series of static snapshots of a particle at the time

of freezing, correlate with the real molecular motions of a protein in an aqueous environment. For many

systems, especially enzymes of clinical significance, proper characterization of the conformational

dynamics and transitional states of the protein are necessary for a full understanding of protein function

and lays the foundation for future therapeutic development. MD simulations can provide this information,

yet such studies are often met with skepticism by experimental researchers. By integrating MD simulations

with experimental techniques, the weaknesses of one technique can be compensated for by the strengths

of other techniques to provide comprehensive information about the system of interest. While this

integrative approach has proven beneficial across a wide range of systems, including proteorhodopsin, the

ribosome, and rotary ATPases, perhaps the best example of this approach is the CRISPR-Cas9 system,

where an exhaustive number of studies have combined cryo-EM, crystallography, and traditional

biophysical techniques with traditional and accelerated MD methods to drive the field forward at an

astounding pace (Arenz *et al.*, 2016; Hirschi *et al.*, 2021; Jiang & Doudna, 2017; Palermo *et al.*, 2016; Roh

et al., 2020; Saha *et al.*, 2022).

Our integrative structural analyses provide the structural basis for IDE conformational dynamics

and their roles in IDE catalytic cycle and the framework for future studies. We speculate that multi-cycles

of open-close transition and rotation driven by IDE conformational dynamics will be required for its

unfoldase activity. However, the kinetic timescale of this reaction is fast, estimated to be ~500 milliseconds

rendering investigation challenging experimentally while the timescale is long for all-atom MD simulation

(Manolopoulou *et al.*, 2009). Our cryo-EM analyses also shed light on IDE allostery (McCord *et al.*, 2013;

Song *et al.*, 2011; Zhang *et al.*, 2018) (Fig. 9). The IDE dimer primarily adopts the O/pO conformational

state (Zhang *et al.*, 2018). Through time-resolved cryo-EM analysis, our data suggest that, upon substrate

binding to the O state, the substrate-inaccessible pO state undergoes allosteric opening, allowing this

subunit to capture its substrate. Additionally, the prolonged incubation of IDE with sub-stoichiometric

amounts of insulin results in the predominance of the pC/pC state, suggestive of coordination in substrate

capture and open-closed transition. Collectively, these positive allosteric transitions would enhance the

capacity of IDE to capture and degrade its substrates more efficiently, though the underlying mechanisms

remain unclear. Future integrative structural studies are required to further elucidate and control the IDE

catalytic cycle, enabling substrate-specific modulation of IDE activity and advancing its therapeutic

potential.

Materials and Methods

<u>Expression and purification of human IDE:</u>

Cysteine-free IDE (referred to as IDE in this study) was expressed in *E. coli* BL21 (DE3) cells (at

25 °C for 20 h, 0.5mM IPTG induction using T7 medium). The R668A mutation was introduced into this

construct by site-directed mutagenesis with the following primer pair:

CCGGAAATTGTTAAGAGATGCCATATATGCTTCTTTG (forward)

CAAAGAAGCATATATGGCATCTCTTAACAACCGGGC (reverse) and verified by sequencing.

Recombinant IDE proteins were purified by Ni-NTA, source-Q, and Superdex 200 columns as previously

described (Zhang *et al.*, 2018). Protein was aliquoted and stored at -80° C.

<u>Cryo-EM data collection and analysis:</u>

Thawed IDE aliquots were further purified by Superdex 200 chromatography using buffer

containing 20 mM HEPES, pH 7.2, 150 mM NaCl, 10mM EDTA and then mixed with Fab$_{H11-E}$ at an equal

molar ratio. Fab$_{H11-E}$-IDE complex was purified by Superdex 200 chromatography in the presence of insulin

at molar ratio IDE:insulin=2:1. Insulin was purchased from SIGMA (91077C). All grids, 300 mesh carbon or

gold holey nanowire grids were plasma cleaned with O_2 and H_2 for 10 secs using a Solarus plasma cleaner

(Gatan) and plunged at 133 milli-second using Chameleon, Spotiton-based technology. All images were

acquired using a Titan Krios microscope (FEI) operated at 300KeV with a Gatan K3 direct electron detector

(Gatan) in counting mode. Images were automatically acquired using Leginon (Suloway *et al*, 2005).

Images were processed using software integrated into RELION3 (Zivanov *et al*, 2018). Frames were

aligned using MotionCor2 (Zheng *et al*, 2017) software with dose weighting, CTF was estimated using Gctf

(Zhang, 2016), particles were picked and extracted automatically using RELION. Particle stacks were

processed through several rounds of 2D and 3D classification. Selected classes were then processed for

high-resolution 3D refinement. The flowchart and detailed data processing is summarized in Figure 2 figure

supplement 1. Finally, the overall map was improved by particle polishing in RELION and sharpening. The

final resolution was estimated using Fourier Shell Correlation (FSC=0.143, gold-standard). The density

fitting and structure refinement was done using UCSF CHIMERA (Pettersen *et al*, 2004), COOT (Emsley &

Cowtan, 2004), REFMAC5 (Murshudov *et al*, 2011) and PHENIX (Adams *et al*, 2010). Time-resolved cryo-

EM grids by rapidly mixing IDE+Fab with insulin were prepared using the Spotiton device as previously

described (Budell *et al*, 2021; Dandey *et al.*, 2020). In brief, IDE+fab and insulin were thawed on ice and

diluted in 20 mM HEPES, pH 7.2, 150 mM NaCl, 10 mM EDTA to 25 µM and 127.5 µM, respectively. Self-

wicking, nanowire grids were plasma treated (Gatan Solarus) at 5 W in hydrogen and oxygen for 1-2

minutes (Wei *et al*, 2018). Each sample was aspirated into separate piezodispensers and applied

sequentially in 70-100% relative humidity at 22-25C onto plunging grids just before cryogen entry. The

maximum constant velocity of the plunging robot was adjusted to obtain the target sample mixing times of

123 milliseconds. The data was collected and processed as above, in Relion5, with Blush regularization

being employed during 3D refinement.

Multibody analysis:

We defined three rigid bodies as shown in figure 2A for the analysis. The size of the user-defined

bodies has been theorized to play a significant role in the success of multibody refinement, most likely due

to the fact that larger bodies will have a stronger signal-to-noise ratio (Nakane *et al.*, 2018). With a Fab

bound to each of IDE-N within IDE dimer, we assessed the impact of body size on multibody refinement by

examining how density corresponding to 2 F_v regions, 1 F_v region, or no F_v regions would affect the

multibody refinement. Fab has a variable region (F_v), and a constant region (F_c) and we chose to subtract

and mask out the density corresponding to the Fab F_c, as we speculated that any motion between the Fab

F_v and F_c domains would bias the results of the multibody refinement. The quality of the multibody results

was assessed based on the subsequent improvement in the Coulomb potential map quality and calculated

resolution. In the absence of any F_v density, the maps resulting from multibody refinement had the best

calculated resolution, yet the map quality was quite poor overall; much of the density appears globular and

featureless, particularly in the IDE-N regions (Figure 2 – figure supplement 4). Conversely, when F_v density

was present on both IDE-N bodies, the resolution of the multibody output maps (4.5 Å) was worse than the

resolution of the input map (4.3 Å). Thus, we found that the greatest improvement occurred when the F_v

density was present on only the exterior body of the pO subunit. In this case, multibody refinement improved

both the calculated resolution and density quality.

Following multibody refinement, the structural heterogeneity for each body was analyzed along six

principal component vectors, resulting in 18 vectors describing the primary components of structural

variation within the IDE dimer. We analyzed the variation along these vectors as a proxy for protein motion.

The results of the multibody analysis indicate the degree of structural variation that is explained by motion

along each of the primary component vectors. For all IDE states, we observed no single component vector

could explain the majority of structural variance (Supplementary files 3-8). Given the underlying

assumptions of multibody refinement, there is a question of whether any observed motions are biologically

relevant or simply the result of improper particle alignment. We reasoned that the greater the degree of

variation explained by motion along a specified component vector, the greater the likelihood was that the

motion would be biologically relevant. Therefore, we focus on the top 9 component vectors, which cover

~75% of total variance for each state, in greater detail.

Molecular dynamics simulations:

All-atom MD simulations were prepared from the O/pO structure of IDE (PDB: 7RZK) and

necessary CHARMM force field PARAM36 files using QwikMD (Ribeiro *et al*, 2016). Gaps in the structure

were modeled in using the AlphaFold model of IDE. A three-step minimization-annealing-equilibration

process was used to generate an equilibrated system. Simulations were performed under NPT (constant

numbers of particles N, pressure P, and temperature T) conditions at 310 K and 1 atm with periodic

boundary conditions in NAMD3.0 (Phillips *et al*, 2005). Explicit solvent was described with the TIP3P model

(Jorgensen *et al*, 1983). Custom TCL scripts in VMD were used to calculate the D1-D4 centers-of-mass

distance and D1-D2-D3-D4 centers-of-mass dihedral angle (Supplementary file 9). Residues 964-988 in

D4 that are mostly absence in all static structures of IDE were built using Alpha-fold, which is highly flexible

in our MD simulations. They were omitted from the analysis of MD simulations, as the inclusion of this

flexible loop significantly altered the center-of-mass independent of global conformational change.

Initial structure of insulin bound IDE, for Upside simulations, was generated using PYMOL that

started with IDE closed monomer (PDB code: 6BF8). Intact insulin (PDB code: 4INS) was fixed by mutation

of residue 30 of insulin chain B back to threonine and modeled into the catalytic chamber based on the

crystal structure of IDE in complex with the partially unfolded insulin (PDB code: 2WBY). The upside

simulation was run with 28 replicates at T_{Upside}=0.8 using Upside 1.0 force field parameters and our total

upside simulation time is approximately 28 μsec (Faruk *et al.*, 2022; Faruk *et al.*, 2023; Jumper *et al.*, 2018a,

b). We estimated the time step of Upside to be ~0.1 picosecond based on folding rates and chain motions.

Custom R, and Python scripts were used for data analysis and presentation (Supplementary files 10-12).

Enzymatic assays:

IDE constructs were exchanged into activity buffer comprised of 25 mM Tris-HCl pH 7.5, 150 mM

NaCl, and 10 mM $ZnCl_2$. MCA-RPPGFSAFK(Dnp) was purchased from Enzo Life Sciences

(BMLP2270001), resuspended in dimethyl sulfoxide at a concentration of 5 mM and diluted to 5 μM with

activity buffer. For activity measurements, 5 μM MCA-RPPGFSAFK(Dnp) was mixed with the desired IDE

construct at concentrations ranging from 1-100 nM in the presence or absence of 1 mM ATP in a 200 μl

reaction. Fluorescence was monitored every 30 sec for 30 minutes with excitation/emission wavelengths of

320/405 nm at 37° C. Initial velocity was calculated during the linear range. Insulin was purchased from MP

Biomedicals (#0219390010), resuspended in 0.01N HCl at 1 mM and diluted to desired concentration with

activity buffer. For competition assays, 5 nM IDE construct and 5 μM MCA-RPPGFSAFK(Dnp) was mixed

with insulin ranging from 0-100 μM. Fluorescence was monitored every 30 sec for 30 minutes with

excitation/emission wavelengths of 320/405 nm at 37° C. Initial velocity was calculated during the linear

range for each concentration of insulin and normalized relative to the velocity of the respective construct in

the absence of insulin to maintain consistency across plates, yielding values for relative inhibition which

were plotted vs [insulin] or [Aβ] and fit to the Michaelis-Menten equation in xmgrace to generate apparent

K_i values. All experiments were performed in triplicate.

Size-exclusion chromatography coupled small-angle X-ray scattering

SAXS/WAXS data collection was employed the Life Sciences X-ray Scattering (LIX) Beamline at

the National Synchrotron Light Source II (NSLS II) at Brookhaven National Laboratory in Upton, NY. Briefly,

60ul of samples in solution were pipetted into PCR tubes, placed into a Bio-Inert Agilent 1260 Infinity II

HPLC multisampler and measured using the isocratic SEC-SAXS format at the beamline (Yang *et al*, 2021).

50ul of sample was injected into the Phenomenex Biozen dSEC2 3 μm bead size with 200Å pore size

column was utilized at a flow rate of 0.35mL/min for 25 minutes. SAXS and WAXS data are collected

simultaneously on a Pilatus 1M (SAXS) and Pilatus 900K (WAXS) detectors with a 2 second exposure

(Yang *et al*, 2020). Data from both detectors is then scaled and merged. Intensity is normalized using the

water peak height at 2.0 Å $^{-1}$. Data processing and analysis was performed using py4xs and lixtools in

jupyter lab. Buffer frames 100-125 were averaged and used for subtraction of averaged frames under the

peak of interest.

Data Availability

Cryo-EM maps and refined models have been deposited to the EMDB and PDB, respectively with the

following accession numbers:

O/O state: EMDB-24760, PDB 7RZH

O/pO state: EMDB-24759, PDB 7RZG

pO/pC state: EMDB-24757, PDB 7RZE

O/pC state: EMDB-24758, PDB 7RZF

pC/pC state: EMDB-24761, PDB 7RZI

Time-resolved O/O state: EMDB-72393, PDB 9Y0H

Cryo-EM micrographs have been deposited to the Electron Microscopy Public Image Archive (EMPIAR)
under the following accession numbers:

IDE+insulin in a 2:1 ratio: EMPIAR-11610

Time-resolved IDE+insulin: EMPIAR-12968.

References

Adams PD, Afonine PV, Bunkoczi G, Chen VB, Davis IW, Echols N, Headd JJ, Hung LW, Kapral GJ, Grosse-
Kunstleve RW *et al* (2010) PHENIX: a comprehensive Python-based system for macromolecular structure
solution. *Acta Crystallogr D Biol Crystallogr* 66: 213-221
Arenz S, Bock LV, Graf M, Innis CA, Beckmann R, Grubmüller H, Vaiana AC, Wilson DN (2016) A combined
cryo-EM and molecular dynamics approach reveals the mechanism of ErmBL-mediated translation
arrest. *Nature Communications* 7: 12026
Armstrong CT, Mason PE, Anderson JLR, Dempsey CE (2016) Arginine side chain interactions and the role
of arginine as a gating charge carrier in voltage sensitive ion channels. *Scientific Reports* 6: 21759
Bahar I, Chennubhotla C, Tobi D (2007) Intrinsic dynamics of enzymes in the unbound state and, relation
to allosteric regulation. *Current opinion in structural biology* 17: 633-640
Bahar I, Rader AJ (2005) Coarse-grained normal mode analysis in structural biology. *Current opinion in
structural biology* 15: 586-592
Baker EN, Blundell TL, Cutfield JF, Cutfield SM, Dodson EJ, Dodson GG, Hodgkin DMC, Hubbard RE, Isaacs
NW, Reynolds CD *et al* (1988) The Structure of 2zn Pig Insulin Crystals at 1.5-a Resolution. *Philosophical
Transactions of the Royal Society B-Biological Sciences* 319: 369-456
Ben Chorin A, Masrati G, Kessel A, Narunsky A, Sprinzak J, Lahav S, Ashkenazy H, Ben-Tal N (2020)
ConSurf-DB: An accessible repository for the evolutionary conservation patterns of the majority of PDB
proteins. *Protein Science* 29: 258-267
Bertram L, Blacker D, Mullin K, Keeney D, Jones J, Basu S, Yhu S, McInnis MG, Go RC, Vekrellis K *et al*
(2000) Evidence for genetic linkage of Alzheimer's disease to chromosome 10q. *Science* 290: 2302-2303
Bertram L, McQueen MB, Mullin K, Blacker D, Tanzi RE (2007) Systematic meta-analyses of Alzheimer
disease genetic association studies: the AlzGene database. *Nat Genet* 39: 17-23
Budell WC, Allegri L, Dandey V, Potter CS, Carragher B (2021) Cryo-Electron Microscopic Grid
Preparation for Time-Resolved Studies using a Novel Robotic System, Spotiton. *Jove-Journal of Visualized
Experiments*
Busto-Moner L, Feng CJ, Antoszewski A, Tokmakoff A, Dinner AR (2021) Structural Ensemble of the
Insulin Monomer. *Biochemistry* 60: 3125-3136
Charton J, Gauriot M, Guo Q, Hennuyer N, Marechal X, Dumont J, Hamdane M, Pottiez V, Landry V,
Sperandio O *et al* (2014) Imidazole-derived 2-[N-carbamoylmethyl-alkylamino]acetic acids, substrate-
dependent modulators of insulin-degrading enzyme in amyloid-β hydrolysis. *Eur J Med Chem* 79: 184-
193

Chiti F, Dobson CM (2006) Protein misfolding, functional amyloid, and human disease. *Annual review of biochemistry* 75: 333-366

Chiti F, Dobson CM (2017) Protein Misfolding, Amyloid Formation, and Human Disease: A Summary of Progress Over the Last Decade. *Annual review of biochemistry* 86: 27-68

Dandey VP, Budell WC, Wei H, Bobe D, Maruthi K, Kopylov M, Eng ET, Kahn PA, Hinshaw JE, Kundu N *et al* (2020) Time-resolved cryo-EM using Spotiton. *Nature Methods* 17: 897-900

Eibauer M, Weber MS, Kronenberg-Tenga R, Beales CT, Boujemaa-Paterski R, Turgay Y, Sivagurunathan S, Kraxner J, Köster S, Goldman RD *et al* (2024) Vimentin filaments integrate low-complexity domains in a complex helical structure. *Nature Structural & Molecular Biology* 31

Emsley P, Cowtan K (2004) Coot: model-building tools for molecular graphics. *Acta Crystallographica Section D-Biological Crystallography* 60: 2126-2132

Farris W, Mansourian S, Chang Y, Lindsley L, Eckman EA, Frosch MP, Eckman CB, Tanzi RE, Selkoe DJ, Guenette S (2003) Insulin-degrading enzyme regulates the levels of insulin, amyloid beta-protein, and the beta-amyloid precursor protein intracellular domain in vivo. *Proc Natl Acad Sci USA* 100: 4162-4167

Farris W, Mansourian S, Leissring MA, Eckman EA, Bertram L, Eckman CB, Tanzi RE, Selkoe DJ (2004) Partial loss-of-function mutations in insulin-degrading enzyme that induce diabetes also impair degradation of amyloid beta-protein. *Am J Pathol* 164: 1425-1434

Faruk NF, Peng XD, Freed KF, Roux B, Sosnick TR (2022) Challenges and Advantages of Accounting for Backbone Flexibility in Prediction of Protein-Protein Complexes. *Journal of Chemical Theory and Computation* 18: 2016-2032

Faruk NF, Peng XD, Sosnick TR (2023) Factors That Control the Force Needed to Unfold a Membrane Protein in Silico Depend on the Mode of Denaturation. *International Journal of Molecular Sciences* 24

Greenwald J, Riek R (2010) Biology of amyloid: structure, function, and regulation. *Structure* 18: 1244-1260

Guo Q, Manolopoulou M, Bian Y, Schilling AB, Tang WJ (2010) Molecular basis for the recognition and cleavages of IGF-II, TGF-alpha, and amylin by human insulin-degrading enzyme. *J Mol Biol* 395: 430-443

Hipp MS, Kasturi P, Hartl FU (2019) The proteostasis network and its decline in ageing. *Nat Rev Mol Cell Biol* 20: 421-435

Hirschi S, Kalbermatter D, Ucurum Z, Lemmin T, Fotiadis D (2021) Cryo-EM structure and dynamics of the green-light absorbing proteorhodopsin. *Nature Communications* 12: 4107

Im H, Manolopoulou M, Malito E, Shen Y, Zhao J, Neant-Fery M, Sun CY, Meredith SC, Sisodia SS, Leissring MA *et al* (2007) Structure of substrate-free human insulin-degrading enzyme (IDE) and biophysical analysis of ATP-induced conformational switch of IDE. *J Biol Chem* 282: 25453-25463

Jiang FG, Doudna JA (2017) CRISPR-Cas9 Structures and Mechanisms. *Annual Review of Biophysics* 46: 505-529

Jorgensen WL, Chandrasekhar J, Madura JD, Impey RW, Klein ML (1983) Comparison of Simple Potential Functions for Simulating Liquid Water. *Journal of Chemical Physics* 79: 926-935

Jumper JM, Faruk NF, Freed KF, Sosnick TR (2018a) Accurate calculation of side chain packing and free energy with applications to protein molecular dynamics. *PLoS computational biology* 14: e1006342

Jumper JM, Faruk NF, Freed KF, Sosnick TR (2018b) Trajectory-based training enables protein simulations with accurate folding and Boltzmann ensembles in cpu-hours. *PLoS computational biology* 14: e1006578

Kim M, Hersh LB, Leissring MA, Ingelsson M, Matsui T, Farris W, Lu A, Hyman BT, Selkoe DJ, Bertram L *et al* (2007) Decreased catalytic activity of the insulin-degrading enzyme in chromosome 10-linked Alzheimer disease families. *J Biol Chem* 282: 7825-7832

King JV, Liang WG, Scherpelz KP, Schilling AB, Meredith SC, Tang W-J (2014) Molecular Basis of Substrate Recognition and Degradation by Human Presequence Protease. *Structure* 22: 996-1007

Leissring MA, Farris W, Chang AY, Walsh DM, Wu X, Sun X, Frosch MP, Selkoe DJ (2003) Enhanced
proteolysis of beta-amyloid in APP transgenic mice prevents plaque formation, secondary pathology,
and premature death. *Neuron* 40: 1087-1093
Li P, Kuo WL, Yousef M, Rosner MR, Tang WJ (2006) The C-terminal domain of human insulin degrading
enzyme is required for dimerization and substrate recognition. *Biochem Biophys Res Commun* 343:
1032-1037
Liang WG, Wijiya J, Wei H, Noble A, Mo S, Lee D, Mancl JM, King JL, Pan M, Liu C *et al* (2022) Structural
basis for the mechanisms of human presequence protease conformational switch and substrate
recognition. *Nature Communications* 13: 1833
Lopez-Otin C, Blasco MA, Partridge L, Serrano M, Kroemer G (2013) The hallmarks of aging. *Cell* 153:
1194-1217
Maianti JP, McFedries A, Foda ZH, Kleiner RE, Du XQ, Leissring MA, Tang W-J, Charron MJ, Seeliger MA,
Saghatelian A *et al* (2014) Anti-diabetic activity of insulin-degrading enzyme inhibitors mediated by
multiple hormones. *Nature* 511: 94-98
Maianti JP, Tan GA, Vetere A, Welsh AJ, Wagner BK, Seeliger MA, Liu DR (2019) Substrate-selective
inhibitors that reprogram the activity of insulin-degrading enzyme. *Nat Chem Biol* 15: 565-574
Malito E, Hulse RE, Tang WJ (2008) Amyloid β-degrading cryptidases: insulin degrading enzyme,
presequence peptidase, and neprilysin. *Cell Mol Life Sci* 65: 2574-2585
Manolopoulou M, Guo Q, Malito E, Schilling AB, Tang WJ (2009) Molecular basis of catalytic chamber-
assisted unfolding and cleavage of human insulin by human insulin-degrading enzyme. *J Biol Chem* 284:
14177-14188
McCord LA, Liang WG, Dowdell E, Kalas V, Hoey RJ, Koide A, Koide S, Tang W-J (2013) Conformational
states and recognition of amyloidogenic peptides of human insulin-degrading enzyme. *Proceedings of*
the National Academy of Sciences of the United States of America 110: 13827-13832
Murshudov GN, Skubák P, Lebedev AA, Pannu NS, Steiner RA, Nicholls RA, Winn MD, Long F, Vagin AA
(2011) REFMAC5 for the refinement of macromolecular crystal structures. *Acta Crystallographica*
Section D-Structural Biology 67: 355-367
Nakane T, Kimanius D, Lindahl E, Scheres SH (2018) Characterisation of molecular motions in cryo-EM
single-particle data by multi-body refinement in RELION. *Elife* 7: e36861
Palermo G, Miao YL, Walker RC, Jinek M, McCammon JA (2016) Striking Plasticity of CRISPR-Cas9 and Key
Role of Non-target DNA, as Revealed by Molecular Simulations. *Acs Central Science* 2: 756-763
Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE (2004) UCSF
chimera - A visualization system for exploratory research and analysis. *Journal of computational*
chemistry 25: 1605-1612
Phillips JC, Braun R, Wang W, Gumbart J, Tajkhorshid E, Villa E, Chipot C, Skeel RD, Kalé L, Schulten K
(2005) Scalable molecular dynamics with NAMD. *Journal of computational chemistry* 26: 1781-1802
Punjani A, Fleet DJ (2021) 3D variability analysis: Resolving continuous flexibility and discrete
heterogeneity from single particle cryo-EM. *Journal of Structural Biology* 213: 107702
Ren M, Guo Q, Guo L, Lenz M, Qian F, Koenen RR, Xu H, Schilling AB, Weber C, Ye RD *et al* (2010)
Polymerization of MIP-1 chemokine (CCL3 and CCL4) and clearance of MIP-1 by insulin-degrading
enzyme. *EMBO J* 29: 3952-3966
Ribeiro JV, Bernardi RC, Rudack T, Stone JE, Phillips JC, Freddolino PL, Schulten K (2016) QwikMD -
Integrative Molecular Dynamics Toolkit for Novices and Experts. *Scientific Reports* 6: 26536
Roh SH, Shekhar M, Pintilie G, Chipot C, Wilkens S, Singharoy A, Chiu W (2020) Cryo-EM and MD infer
water-mediated proton transport and autoinhibition mechanisms of V complex. *Science Advances* 6:
eabb9605
Saha A, Arantes PR, Palermo G (2022) Dynamics and mechanisms of CRISPR-Cas9 through the lens of
computational methods. *Current opinion in structural biology* 75: 102400

Saido T, Leissring MA (2012) Proteolytic degradation of amyloid beta-protein. *Cold Spring Harb Perspect*
Med 2: a006379
Shen Y, Joachimiak A, Rosner MR, Tang WJ (2006) Structures of human insulin-degrading enzyme reveal
a new substrate recognition mechanism. *Nature* 443: 870-874
Sladek R, Rocheleau G, Rung J, Dina C, Shen L, Serre D, Boutin P, Vincent D, Belisle A, Hadjadj S *et al*
(2007) A genome-wide association study identifies novel risk loci for type 2 diabetes. *Nature* 445: 881-
885
Song ES, Jang H, Guo HF, Juliano MA, Juliano L, Morris AJ, Galperin E, Rodgers DW, Hersh LB (2017)
Inositol phosphates and phosphoinositides activate insulin-degrading enzyme, while phosphoinositides
also mediate binding to endosomes. *Proceedings of the National Academy of Sciences of the United*
States of America 114: E2826-E2835
Song ES, Juliano MA, Juliano L, Fried MG, Wagner SL, Hersh LB (2004) ATP effects on insulin-degrading
enzyme are mediated primarily through its triphosphate moiety. *Journal of Biological Chemistry* 279:
54216-54220
Song ESE, Rodgers DWD, Hersh LBL (2011) Mixed dimers of insulin-degrading enzyme reveal a cis
activation mechanism. *The Journal of biological chemistry* 286: 13852-13858
Suloway C, Pulokas J, Fellmann D, Cheng A, Guerra F, Quispe J, Stagg S, Potter CS, Carragher B (2005)
Automated molecular microscopy: the new Leginon system. *J Struct Biol* 151: 41-60
Tang WJ (2016) Targeting Insulin-Degrading Enzyme to Treat Type 2 Diabetes Mellitus. *Trends Endocrinol*
Metab 27: 24-34
Tanzi RE, Bertram L (2005) Twenty years of the Alzheimer's disease amyloid hypothesis: a genetic
perspective. *Cell* 120: 545-555
Vernon RM, Chong PA, Tsang B, Kim TH, Bah A, Farber P, Lin H, Forman-Kay JD (2018) Pi-Pi contacts are
an overlooked protein feature relevant to phase separation. *Elife* 7: e31486
Wei H, Dandey VP, Zhang ZN, Raczkowski A, Rice WJ, Carragher B, Potter CS (2018) Optimizing "self-
wicking" nanowire grids. *Journal of Structural Biology* 202: 170-174
Yang L, Antonelli S, Chodankar S, Byrnes J, Lazo E, Qian K (2020) Solution scattering at the Life Science X-
ray Scattering (LiX) beamline. *J Synchrotron Radiat* 27: 804-812
Yang L, Lazo E, Byrnes J, Chodankar S, Antonelli S, Rakitin M (2021) Tools for supporting solution
scattering during the COVID-19 pandemic. *J Synchrotron Radiat* 28: 1237-1244
Zeggini E, Weedon MN, Lindgren CM, Frayling TM, Elliott KS, Lango H, Timpson NJ, Perry JRB, Rayner
NW, Freathy RM *et al* (2007) Replication of genome-wide association signals in UK samples reveals risk
loci for type 2 diabetes. *Science* 316: 1336-1341
Zhang K (2016) Gctf: Real-time CTF determination and correction. *Journal of Structural Biology* 193: 1-12
Zhang ZN, Liang WGG, Bailey LJ, Tan YZ, Wei H, Wang A, Farcasanu M, Woods VA, McCord LA, Lee D *et al*
(2018) Ensemble cryoEM elucidates the mechanism of insulin capture and degradation by human insulin
degrading enzyme. *Elife* 7: e33572
Zheng S, Polovcak E, Armache JP, Cheng Y, Agard D (2017) MotionCor2: anisotropic Correction of Beam-
induced Motion for Improved Single-particle Electron Cryomicroscopy. *Nature Methods* 14: 331-332
Zhong ED, Bepler T, Berger B, Davis JH (2021) CryoDRGN: reconstruction of heterogeneous cryo-EM
structures using neural networks. *Nature Methods* 18: 176-185
Zhou XM, Lin Y, Kato M, Mori E, Liszczak G, Sutherland L, Sysoev VO, Murray DT, Tycko R, McKnight SL
(2021) Transiently structured head domains control intermediate filament assembly. *Proceedings of the*
National Academy of Sciences of the United States of America 118
Zivanov J, Nakane T, Forsberg BO, Kimanius D, Hagen WJ, Lindahl E, Scheres SH (2018) New tools for
automated high-resolution cryo-EM structure determination in RELION-3. *Elife* 7: e42166

 Figures

Figure 1: Overview of structure and function of IDE. (A) Overall structure of IDE. IDE is comprised of
four structurally homologous domains: D1 (red, residues 43-285), D2 (orange, residues 286-515), D3 (blue,
residues 545-768), and D4 (cyan, residues 769-1,016). The domains are arranged in two roughly
hemispherical regions that enclose the catalytic chamber referred to as IDE-N (comprised of the D1 and
D2 domains) and IDE-C (comprising the D3 and D4 domains), which are joined by a linker region (residues
516-544). (B) Primary sequence and overall structure of insulin. Insulin cleavage by IDE as revealed by
mass spectrometry (Manolopoulou *et al.*, 2009). Cleavage sites are marked in arrows, which occur within
α-helical regions and requires the peptide to be unfolded prior to cleavage. (C) Overview of the IDE catalytic
cycle with key questions addressed in this study.

Figure 2: Cryo-EM structures. (A) Overview of the cryo-EM structures. See figure S1 for processing details. **(B)** Comparison of the open (O), partial open (pO), and partial closed (pC) subunit states present in our cryo-EM structures with domain organization. The distance between the D1 and D4 domain centers-of-mass (D1-D4 COM) along with the dihedral angle formed by the D1-D2-D3-D4 domain centers-of-mass (D1-D2-D3-D4 dihedral) described in Zhang et al. (21) and depicted in Figure 3 – figure supplement 1 were used as biologically important criteria to quantify observed conformations. **(C)** Insulin density and corresponding model in our cryo-EM structures. Both the A chain (magenta) and B chain (yellow) can fit the density in the exosite and catalytic cleft.

Figure 3: Conformational dynamics of IDE implied by structural heterogeneity. (A) All-atom MD
simulations analysis. The primary source of structural variance (RMSD) results from the IDE-N moving
against IDE-C as a rigid body. Rigid bodies were defined as colored for multibody refinement in RELION.
(B-C) Multibody analysis. The range of conformational variance described by the top principal component
vectors displays an unexpectedly high degree of rotational motion, as measured by the change in D1-D2-
D3-D4 dihedral angle across the gradient of structural heterogeneity for each vector, in both the absence
(B) and presence **(C)** of insulin compared to the expected open-close transition pathway predicted from a
linear interpolation of the experimentally determined structures of IDE (dashed line, Figure 3 – figure
supplement 1). Two dominant components of structural variance are revealed from multibody analysis: **(D)**
where IDE-N swings relative to IDE-N about the inter-domain linker, and **(E)** where IDE-N rotates against
IDE-C. Starting (orange) and ending (red) states of IDE-N shown with pathway depicted by arrows. IDE-C
shown as gray surface.

Figure 4: All-atom MD reveals a molecular basis for IDE conformational dynamics. **(A)** Measurements of the O subunit D1-D4 distance over the course of six separate microsecond long all-atom MD simulations of WT IDE. Of which, the open subunit closed in 5 of the 6 simulations. **(B)** Plot of the O subunit D1-D4 distance vs the D1-D2-D3-D4 COM dihedral angle over the course of the simulation of WT IDE. The open

subunits displayed a variety of closing pathways and did not close to a consensus structure. Starting

structure shown as black dot, pO structure shown as white dot. **(C)** R668 acts as a guidepost residue,

rapidly interacting with D309 or E381. Formation of this interaction is associated with rapid closing, as

measured by a decrease in D1-D4 distance **(D)**. **(E)** Hydrogen-deuterium exchange mass spectrometry

highlights the importance of R668 in mediating the open-close transition. In the presence of insulin (panel

1, red), Aβ (panel 2, red) , and BDM-44768 (panel 3, red), all of which promote IDE closing, the peptide

containing R668 shows reduced deuterium exchange relative to apo-IDE (black), yet in the presence of 6bk

(panel 4, red), which does not promote closing, there is no difference in the exchange rates for the R668

containing peptide relative to apo-IDE (black). Helix containing R668 colored by red (increase) – white (no

change) – blue (decrease) gradient depicting the degree of deuterium exchange relative to apo-IDE. **(F)**

Measurements of the O subunit D1-D4 distance over the course of six separate microsecond long all-atom

MD simulations of IDE R668A. **(G)** Plot of the O subunit D1-D4 distance vs the D1-D2-D3-D4 COM dihedral

angle over the course of the simulation of IDE R668A. The six separate microsecond long simulations

indicate that an R668A mutation significantly alters the closing dynamics of IDE **(F)** and increases the

rotational motion **(G)** relative to WT (panels **A** and **B** respectively). Starting structure shown as black dot,

pO structure shown as white dot.

Figure 5: R668A alters IDE activity *in vitro* (A) Elution profile of WT IDE (blue) compared to the R668A

mutant (orange) from a S200 SEC column. (**B**) Degradation of the fluorescent substrate MCA-

RPPGFSAFK(Dnp) by WT IDE and the R668A construct in the presence and absence of ATP. Data

represents the average initial velocities of three replicates performed at a protein concentration of 3.125

nM. Error bars (gray) represent the standard error. **(C)** Inhibition of MCA-RPPGFSAFK(Dnp) degradation

by WT IDE (circles, solid fit lines) and IDE R668A (squares, dashed fit lines) in the presence of varying

amounts of insulin. Data was fit to the Michaelis-Menten (black) and Hill equations (red). Relevant

parameters, Michaelis-Menten: WT: χ^2=0.001, V_{max}=0.951, K_i=8.3 nM; R668A: χ^2=0.005, V_{max}=0.892, K_i=52

nM; Hill: WT: χ^2=0.009, n=0.55, K_i=51 nM; R668A: χ^2=0.055, n=0.61, K_i=198 nM. Error bars represent

standard error, data points represent the average of three technical replicates. **(D)** Inhibition of MCA-

RPPGFSAFK(Dnp) degradation by WT IDE (circles, solid fit lines) and IDE R668A (squares, dashed fit

lines) in the presence of varying amounts of Aβ_{1-40}. WT data was fit to the Michaelis-Menten (black) and Hill

equations (red), R668A data could not be fit to either equation, indicating that the mutation confers

substrate-specific altered enzyme kinetics. Relevant WT parameters, Michaelis-Menten: χ^2=0.066,

V_{max}=0.823, K_i=353 nM; Hill: χ^2=0.08, n=0.72, K_i=688 nM. Error bars represent standard error, data points

represent the average of three technical replicates **(E)** SEC-SAXS profile of WT (black) and R668A (red)

constructs with Rg values calculated by both the Guinier and Porod methods along with Dmax derived from

the P(r) function **(F)**.

Figure 6 Structural basis of closed state conformational dynamics. (A) Measurements of the pO

subunit D1-D4 distance over the course of six separate microsecond long all-atom MD simulations of WT

IDE. **(B)** Plot of the pO D1-D4 distance vs the D1-D2-D3-D4 COM dihedral angle over the course of the

simulation of WT IDE. **(C)** IDE-N/C interface previously solved crystal structures (PDB:2G47 shown) shows

side chains are ill-positioned for interaction. **(D)** IDE-N/C interface formed upon open subunit closing in our

MD simulations reveals a complex hydrogen bonding network. **(E)** Heat map showing conformational

geometries that were preferentially sampled in our MD simulations by the open subunits upon closing.

Insets highlight how the IDE-N/C interface changes to permit interdomain motion. **(F)** Plot of the O subunit

D1-D2-D3-D4 dihedral angle during a subset of a single WT IDE MD simulation after the open-close

transition has been completed. Charge-swapping between residues at the IDE-N/C interface is associated

with changes in the D1-D2-D3-D4 dihedral. **(G)** For most of the simulation, D309 interacts with K483 (black),

however, this interaction is broken for ~100 ns, during which D309 instead interacts with R311 (blue) and

R668 (orange). **(H)** For most of the simulation, D426 interacts with K571 (black), yet this interaction is

periodically broken, and D426 instead interacts with K425 (green) and K899 (magenta). When these events

of charge-swapping coincide with D309 charge-swapping **(G)**, they are associated with a large change in

the D1-D2-D3-D4 dihedral angle **(F)**. When they occur alone, the effect on D1-D2-D3-D4 dihedral is smaller.

A

B

	Chain A	Chain B
D1-D4 COM	46.8 Å	55.1 Å
D1-D2-D3-D4 dihedral	-30.4°	-36.5°

C

9Y0H vs 7RZH

9Y0H vs 6B7Y

D

Conformational distribution of IDE by condition

Apo-IDE	IDE+Insulin (2:1)	IDE+Insulin (1:5) long	IDE+Insulin (1:5) short
	63% pC/pC		
72.8% O/pO	15.5% O/pO		
16% O/O	14.3% O/pC	100% pC/pC	100% O/O
11.2% pO/pO	3.6% O/O		
	3.6% pO/pC		

Figure 7: Time-resolved cryo-EM of IDE+insulin reveals a new O/O state. (A) 5.1 Å reconstruction of
IDE+Fab rapidly mixed with a 5x molar excess of insulin and vitrified with a mix-to-freeze time of 123
milliseconds. **(B)** Measurements (as in Fig. 2) of the distance between the D1-D4 centers-of-mass and
dihedral angle formed by the D1-D2-D3-D4 centers-of-mass as indicators of the "openness" of the structure

depicted in (A). **(C)** Alignment of the O/O state from our rapid-mixing dataset (domains colored as in Fig. 1)
compared to the previously solved O/O states (gray, as labeled). **(D)** Overview of the conformational states
adopted by IDE in all 4 available cryo-EM datasets with corresponding percentage of particles mapping to
each state (Zhang *et al.*, 2018)(this work).

Figure 8: Upside analysis of IDE-insulin interactions. (A) Transient cross-β-strand interactions between
insulin and IDE. Secondary structure (gray: coil, green: β-strand, yellow: α-helix) for each insulin residue
(Y-axis) over the course (X-axis) of a representative Upside simulation. Points indicate predicted hydrogen
bond interactions between IDE and insulin, colored by which domain of IDE that insulin residue is interacting
with (D1: red, D2: orange, D3: blue, D4: cyan). Stars below the X-axis indicate frames from which structures
were extracted for panel B. **(B)** Structure of IDE/insulin complex in Upside simulations at frames indicated
in (A). IDE domains colored as in (A), insulin shown in magenta (chain A) and yellow (chain B). Inset
highlights the intermolecular cross-β sheets formed between IDE and insulin within each of the four
homologous IDE domains. **(C)** Individual domains of IDE, colored as in (A), extracted from (B) and oriented
to demonstrate the structural conservation of intermolecular cross-β-sheet formation with bound insulin
(colored as in B).

Figure 9: Model for the catalytic cycle of IDE. The details are described in discussion. For simplicity,
only the scenario involving the addition of a second insulin molecule is illustrated, although this is not
required for the catalytic cycle. The mechanism by which IDE transitions between closed and open states
remains unknown and is therefore indicated with question marks.

901