Integration of head and body orientations in the macaque superior temporal sulcus is specific to

upright bodies

Yordanka Zafirova[1,2] & Rufin Vogels[1,2]

[1]Laboratorium voor Neuro- en Psychofysiologie, Department of Neurosciences, KU Leuven.

[2]Leuven Brain Institute, KU Leuven, Belgium.

Corresponding author: rufin.vogels@kuleuven.be

Abstract

The neural processing of faces and bodies is often studied separately, despite their natural

integration in perception. Unlike prior research on the neural selectivity for either head or body

orientation, we investigated their interaction in macaque superior temporal sulcus (STS) using a

monkey avatar with diverse head-body orientation angles. STS neurons showed selectivity for

specific combinations of head-body orientations. Anterior STS (aSTS) neurons enabled more reliable

decoding of head-body configuration angles compared to middle STS neurons. Decoding accuracy in

aSTS was lowest for head-body angle pairs differing only in sign (e.g. head-body orientation

difference of ±90° relative to the anatomical midline), and highest for aligned (0°) head-body

orientations versus those with maximum angular difference. Inverted bodies showed diminished

decoding of head-body orientation angle compared to upright bodies. These findings show that aSTS

integrates head and body orientation cues, revealing configuration-specific neural mechanisms, and

advance our understanding of social perception.

Introduction

Faces and bodies are typically studied separately, although both are parts of the same agent

(Freiwald *et al.*, 2016; Tsao and Livingstone, 2008; de Gelder *et al.*, 2010; Peelen and Downing, 2007).

Despite the distinct research traditions, behavioral and fMRI studies in humans suggest that face and

body processing interact (Hu *et al.*, 2020; Taubert *et al.*, 2022). Furthermore, face-selective units in

macaque mid Superior Temporal Sulcus (STS) face-selective patches respond to an object occluding a

face on top of a body (Arcaro *et al.*, 2020), which agrees with monkey (Fisher and Freiwald, 2015)

and human fMRI (Cox *et al.*, 2004) studies that showed that a blurred head on top of a body can

activate face-selective regions. Recently, we showed that anterior inferior temporal (IT) neurons

respond stronger to face-body configurations in which the face is located at its natural position on

top of the body than to unnatural face-body configurations, showing that the face and body interact

in a configuration-specific manner in IT (Zafirova *et al.*, 2024; Zafirova *et al.*, 2022). While these

studies demonstrate that face and body processing interact in the visual temporal cortex, the extent

and underlying mechanisms of this interaction remain unexplored.

The responses of most face-selective STS neurons are modulated by the 3D orientation of the

head (Freiwald and Tsao, 2010; Perrett *et al.*, 1992; Perrett *et al.*, 1984, 1985). Likewise, the

responses of most body-selective STS neurons are sensitive to body orientation (Bao *et al.*, 2020;

Kumar *et al.*, 2017). This might not be surprising because both head and body orientation are

important social cues, e.g. for the attention of a viewed agent or the direction of an agent's action. In

these studies, either face or body orientation, but not both, were manipulated. Here we address the

outstanding question of whether the orientation of the head and body interact in the STS: does the

selectivity for head orientation depend on the body orientation, and vice versa? Are STS neurons

selective for a particular combination of head and body orientation, suggesting a selectivity for both

head and body features in a particular pose, e.g. a selective response for a frontal body with a head

turned to the right? Our study was also motivated by human psychophysical studies that showed

that body and head orientation interact in social perception (Hietanen, 2002; Moors *et al.*, 2015;

Vrancken *et al.*, 2017), although it is unclear at which level – sensory or higher – the observed

interactions occurred.

Here, we manipulated independently the head and body orientation of a monkey avatar. We

employed both anatomically possible and impossible head-body orientations (e.g. a frontal body with

a backward-oriented head, i.e. 180° difference), allowing us to assess whether anatomically possible

and impossible head-body orientation angles are encoded differently by STS neurons.

Behavioral studies in humans (Griffin and Oswald, 2022) and monkeys (Matsuno and Fujita,

2018) provided evidence for greater body discrimination for upright compared to inverted bodies.

This body-inversion effect has been interpreted as providing evidence for the holistic encoding of

bodies, similar to that proposed for faces. Some studies suggested that the body-inversion effect is

driven by the head (for review, see (Griffin and Oswald, 2022)). However, other studies have

observed a behavioral body-inversion effect for headless bodies (for a meta-analysis, see (Griffin and

Oswald, 2022)). So far, no neural correlates of the body-inversion effect for whole bodies have been

described at the single-unit level. Here, we compared the responses and body-head interaction of

STS units between upright and inverted bodies, assessing whether these units show stronger

responses, or higher body-selectivity and body-head interactions for upright bodies.

We recorded the responses of units at two posterior-anterior levels of the ventral STS, in and

surrounding patches that were activated more by images of monkeys compared to objects in a fMRI

study with the same subjects ((Zafirova *et al.*, 2022); Figure 1A). Because it has been proposed that

face-body interactions are more prominent in anterior IT (Fisher and Freiwald, 2015; Hu *et al.*, 2020;

Zafirova *et al.*, 2022), we expected stronger head-body orientation interactions in the anterior STS.

Results

In experiment 1, we targeted two fMRI-defined patches that were defined by a higher

activation to images of monkeys compared to objects ((Zafirova *et al.*, 2022); Figure 1A). The

posterior patch was located in the ventral bank of the mid STS (mSTS) and the anterior one was

about 12 mm more anterior in the ventral bank of the STS (aSTS). In experiment 1, we recorded well-

isolated single units in both regions. In the Selectivity test (Methods), we compared the responses to

a set of monkeys, faces, headless bodies, and objects (see (Zafirova *et al.*, 2022)). We took neurons

(Methods) for further examination when their mean net response to the monkey, face, or body was

at least twice the mean response to the corresponding object control condition. Thus, we obtained

98 aSTS and 100 mSTS neurons, split approximately equally across the two subjects. In both regions,

the averaged responses of the selected neurons were strongest for the monkey images, followed

closely by the body and the face, and much weaker for the objects (Figure 1B). Some neurons

responded stronger to faces than bodies, or vice versa, and showed strong selectivity for different

body or face images (Figure 1B).

Figure 1. Recording locations and stimulus selectivity (A). Targeted patches (red circles) shown in the native space of M1 (left) and M2 (right) on 2 mm coronal slices. Brown vertical lines illustrate penetrations. Monkey patches were defined by the contrast monkeys > monkey control objects (M-O$_M$; p < 0.05 FEW). **(B).** Normalized net responses in the Selectivity test (Experiment 1) in aSTS (left) and mSTS (right). Rows represent neurons from M1 and M2 (separated by a dotted line). Columns correspond to the stimuli, color-coded per category with a bar and one example image. Below, bars in corresponding colors show the mean normalized net response to each stimulus (95% confidence intervals (1000 resamplings)). *n* indicates the number of neurons.

These neurons were tested in the Head-body Orientation test in which we manipulated

independently the orientation of the head and the body of a monkey avatar (Methods; Figure 2A).

The monkey avatar was either sitting (P1) or standing (P2). For each pose, we had 64 conditions in

which the orientation of the head and body was manipulated independently in steps of 45° (Figure

2A). To separate the effect of head/body location from head or body orientation, we presented all

images at two locations: monkey-centered (MC) and head-centered (HC; Figure 2B).

Figure 2. Monkey avatar stimuli for the Head-body orientation test (Experiment 1). (A). 64 combinations of head and body

orientations for the sitting (P1, left) and the standing pose (P2, right). Body orientation varies by row and head orientation

varies by column. Different colors group the 8 avatar orientations of four head-body angles. **(B).** Example stimuli of P1 and

P2 in the monkey-centered (MC, left) and the head-centered (HC, right) conditions, with zero angle between head and

body. **(C).** Headless bodies (B) and heads (H) for the zero and straight angle head-body configurations, showing the frontal,

back, and two lateral views for P1 and P2. Images are shown for MC (left) and HC (right) conditions and were positioned to

match their corresponding avatars in (A). The location of the red fixation target is indicated in **(B)** and **(C)**.

Responses for a monkey are on average less than the summed responses for the body and head (experiment 1)

For each pose, the head and body of the frontal, back, and two lateral orientations were also

presented in isolation at the same locations as in the avatar. The latter was done for head-body

configurations of zero and straight ($0°$ and $180°$) angles between the head and body orientation

(Figure 2B, 2C). This allowed us to compare the responses to the isolated body and head with those

to the monkeys having the same head and body at the same locations. For each case for which there

was a significant excitatory response to either the monkey or the corresponding head and body

compared to the baseline, we computed the Monkey-Sum Index (MSI) contrasting the net response

to the monkey and the sum of the net responses to its corresponding head and body (Methods). MSI

larger than 0 corresponds to a superadditive response to the monkey while MSI smaller than 0

indicates a subadditive response. For both subjects and regions, the median MSIs for P1 and P2 were

smaller than 0 (Figure 3A), which agrees with a single-unit study in anterior IT that employed images

of real macaques (Zafirova *et al.*, 2024). Surprisingly, we observed neurons that responded to either

the headless body or head but did not respond to the whole monkey that included the same body

and head (MSI close to -1; Figure 3A). The weak responses to the monkey were not limited to cases in

which the head-body configuration was anatomically impossible, i.e. a straight angle between the

head and the body. In fact, cases in which MSI was between -0.9 and -1 included 53% and 41% of

natural head-body configurations in aSTS and mSTS, respectively. In line with a previous study

(Zafirova *et al.*, 2024), only a minority of neurons demonstrated strong superadditive responses to

the monkey (MSI between 0.9 and 1: 4.3% and 6.2% in aSTS and mSTS, respectively). Figure 3E

illustrates the responses to heads, bodies, and monkeys of example units with different values of

MSI. These data show that heads and bodies interact at the level of the STS.

Figure 3. Neuronal responses in the Head-body orientation test (Experiment 1). (A). Distribution of MSI computed for the

responses for all orientation, pose (P1 and P2), and centering (MC (blue) and HC (red)) combinations in aSTS (upper panel)

and mSTS (lower panel). Data for M1 and M2, and medians (triangles) are shown in different hues. **(B).** Significant main

effects and interactions of the factors body and head orientation, for P1 and P2 in MC (blue) and HC (red) conditions in

corresponding colors) in aSTS (left) and mSTS (right). Pooled data of both subjects. Significant interactions in both MC and

HC are highlighted with darker blue and red (number of cases (n) in black). **(C).** Distribution of the correlation coefficients

between responses to the head-body configurations in MC and HC conditions in aSTS (left) and mSTS (right). Data from both

subjects and poses were pooled. Correlation coefficients for all cases are shown above the x-axis. Below the x-axis, the light

green bars represent the correlation coefficients for cases with a significant interaction (either MC or HC), while the dark

green bars indicate the coefficients for cases with a significant interaction in both MC and HC. N and medians (triangles) are

shown in corresponding colors. **(D).** Normalized net responses of example neurons to the 64 combinations of head and

body orientations in aSTS (left) and mSTS (right) for MC and HC conditions. Body orientation (ORT) varies by row and head

orientation varies by column, from 0° to 315°. **(E).** Net responses of example neurons to bodies, heads, and the

corresponding monkey configurations with MSI of 1, -1, or the median. Data for P1 and P2, MC and HC, and aSTS and mSTS

(smoothed with the Matlab function *interp1*).

Interaction of head and body orientation tuning (experiment 1)

The main question of this study was whether head and body *orientation tuning* interact.

Thus, we performed a two-way ANOVA on the responses (Methods) to the 64 head-body orientation

conditions, with factors the orientation of the head and body (Figure 2A). The large majority of

neurons showed a significant main effect ($p < 0.05$) of body orientation followed by a main effect of

head orientation (Figure 3B). Importantly, a large fraction of neurons showed a significant interaction

between head and body orientation. This was true for both centerings (MC and HC) and poses (P1

and P2) in each region (aSTS and mSTS, Figure 3B).

These results do not necessarily provide evidence for genuine head and body orientation

interactions because orientation changes could be confounded with location changes of the monkey

part within a centering condition (Figure 2B). To address this, we first assessed whether the tuning

for body/head orientation was location tolerant by correlating the responses to the 64 head-body

configurations between MC and HC. This was done for all neuron and pose combinations in which

there was a significant response in either centering condition (Split-plot ANOVA). The median

correlations were 0.68 and 0.57 in aSTS and mSTS, respectively (Figure 3C), indicating position

tolerance of the orientation selectivity in both regions. In line with its higher hierarchical stage and

thus expected higher location tolerance, aSTS showed significantly higher correlations than mSTS

(Wilcoxon rank sum test, p = 0.008; tested after pooling data of the poses per neuron; n = 75 aSTS,

and n = 77 mSTS neurons). The higher correlations for aSTS were not a result of differences in

response reliability between the two regions: the median Spearman-Brown corrected split-half

Pearson correlation (Methods) was 0.82 and 0.81 for the MC and HC aSTS neurons (n = 131 cases)

and 0.85 and 0.89 for the MC and HC mSTS neurons (n = 126 cases). The relatively higher reliability

for mSTS shows that the lower correlation of the head-body orientation selectivity between the two

centerings for mSTS neurons is not due to lower reliability. Furthermore, correcting the correlations

for reliability showed a lower median corrected correlation for mSTS (median: 0.72) compared to

aSTS (0.90).

To assess the position tolerance of the head-body orientation interaction per se, we

computed how many neurons for each pose show a significant interaction (two-way ANOVA; see

above) of head and body orientation for both centerings. In aSTS, 24% of the neurons responding for

both centerings of a pose (n=135) showed a significant interaction in both centerings, and this

proportion was similar for mSTS (26%; n=127; Figure 3B). Notably, the correlations between the

responses to the 64 head-body orientation conditions of the two centerings for the neuron and pose

combinations showing significant head-body interactions for the two centerings were similar to those

observed for the whole population (Figure 3C).

Next, we isolated the interaction of head and body orientation by computing the residuals of

the responses from a linear additive model in which the responses to a head-body configuration are

an addition of the mean responses to the body (averaging across the 8 head orientations) and head

(averaging across the 8 body orientations) orientations. The residuals were computed for both

centerings and then correlated across the centerings per neuron and pose combination. We

computed these correlations for those cases in which the 2-way ANOVA (see above) showed a

significant interaction for both centerings. The median correlation between the residuals of the two

centerings was 0.23 (quartiles: 0.08 – 0.36; n = 33) for mSTS, whereas it was 0.41 (0.21 – 0.55; n= 33;

median larger than 0: p = 9.35e-07) for aSTS, which was significantly greater than for mSTS (Wilcoxon

rank-sum test; p = 0.008). These data demonstrate a position-tolerance of the interaction of head

and body orientation in aSTS, but less so in mSTS.

Single neurons showed a variety of head-body orientation tunings (Figure 3D). Interestingly,

some neurons (e.g. Figure 3D) were tuned to a particular combination of a head and body

irrespective of centering.

<u>Decoding of orientation-invariant head-body configuration (experiment 1)</u>

Given the head-body orientation interactions in the STS, we examined whether we could

decode head-body configurations with different angles between head and body in an orientation-

invariant way from our sample of neurons using a linear classifier (linear kernel Support Vector

Machine (SVM)). We trained the classifier to distinguish two angles after pooling the responses of all

8 orientations of a head-body orientation angle configuration (Methods; examples in Figure 4A).

Because the training used responses to all orientations per angle and because the same head and

body orientations were present in both classes, we forced the classifier to employ the *combination* of

head and body orientation. We used 8-fold cross-validation with training and test trials evenly

distributed across orientations (Methods). We will report below the classification accuracies

obtained when training was performed using trials of one centering condition (MC) and testing for

the other centering condition (HC), requiring generalization across positions. The classification

accuracies for training on the HC and testing on the MC condition were highly similar (Figure S1A).

We employed for each region 60 neurons, 30 randomly selected for each monkey subject, as that

was the maximum number of neurons with responses in both centering conditions (MC and HC) per

subject and recording location (aSTS and mSTS). The classification was performed using raw firing

rates. We report the mean and standard deviation, across 100 resamplings, of the classification

accuracies for the test trials.

Figure 4. Decoding of head-body angles from neuronal responses in the Head-body orientation test (Experiment 1). (A).

Head-body configurations with zero and straight angles (black arrows in **(B)**) and leftward and rightward right angles of

head and body (grey arrows in **(B)**), of P1 and P2, at eight avatar orientations. **(B).** Mean decoding accuracies for the head-

body configurations with zero, straight, leftward, and rightward right angles versus other head-body angles as indicated, in

aSTS (top two panels) and mSTS (bottom two panels), for poses P1 and P2. N indicates the number of neurons. Results are

for MC training and HC testing. Responses to the 8 orientations of a configuration (rows in **(A)**) were pooled when training

the classifier. Error bars show standard deviations across resamplings. The shaded area indicates null distribution (stimulus

labels permutation). **(C).** Decoding accuracies for different head-body angle pairs in aSTS, for P1 (top) and P2 (bottom).

Decoded pairs are connected with lines, with decoding accuracies indicated by color. Training was performed after pooling

the 8 orientations of a head-body configuration defined by the angle of the head and body. Results are from MC training

and HC testing. Avatars of one orientation of the head-body configuration angles are shown.

Figure 4B shows the mean classification accuracies for pairs of head-body orientation angles,

plotted for each of 4 reference angles (0° (zero angle), 180° (straight angle), 90° (rightward right

angle), and -90° (leftward right angle)) versus other angles. For the zero and straight reference

angles, the classification accuracy increased with the absolute difference between the angles,

reaching a maximum of about 0.7 classification accuracy for zero versus straight angles. Classification

accuracies were smaller for mSTS compared to aSTS. Surprisingly, for the rightward (90°) and

leftward (-90°) right angles as references, classification scores were low (Figure 4B), which contrasts

with the higher scores for the same 180° orientation difference when the angles were zero and

straight (this difference was significant for both centerings and poses in aSTS; all p-values < 0.01).

Classification scores tended to be higher for 90° differences in angles (e.g. right versus straight) than

for the 180° difference (rightward versus leftward right angles) for the rightward or leftward right

reference angles.

The decoding accuracies showed the same pattern, although were higher, when training and

testing were performed with the same centering, requiring no generalization across locations (Figure

S1B and S1C).

We applied also a more stringent classification in which the trained and tested orientations

differed: we trained the classifier for the frontal, back, and two lateral orientations (0°, 180°, 90°, and

-90°, i.e. '+' orientations in Figure 5A), and tested with the oblique ones (45°, 135°, -135° and -45°, i.e.

'x' orientations in Figure 5A), and vice versa (Figure 5A, Figure S2A). Again, the trained and tested

responses were from different centerings. For this analysis, we employed the aSTS sample, as that

produced the highest decoding accuracies previously. For P1, the performance accuracies for the

aSTS sample were well above chance for the zero versus straight angle decoding, and, as before,

were lower for the leftward versus rightward right angle decoding (Figure 5B; for brevity, we will

refer to these angles as left and right angles, respectively, below). The same was true for P2 for two

of the four centering and orientation training-test combinations (Figure 5B; Figure S2B). However, for

the other two combinations, the performance accuracies for the zero versus straight angles were

barely above chance and were at a similar level as those for the right versus left angle decoding

(Figure 5B; Figure S2B). This interaction between generalization across orientation and centering for

P2 is likely related to the larger variations in the eccentricity of the head when changing the

orientation and centering of the standing avatar (see Figure S2). Nonetheless, when generalization

across orientation but not across centering was required, accuracies for P1 and P2 were similar for all

decodings (for the zero versus straight angle: classification accuracies for P1 ranging from 0.64 - 0.72

and for P2 ranging from 0.66 - 0.72; for the right versus left angle decoding: P1 0.49 - 0.54 and P2

0.48 - 0.56).

Figure 5. Decoding of head-body angles across orientations (Experiment 1). (A). Head-body configurations with zero and straight angles, and leftward and rightward right angles for P1 and P2, split into four orientations for training and testing. **(B).** Decoding accuracies for the head-body configurations with zero, straight, leftward, and rightward right angles in aSTS, for P1 (top) and P2 (bottom); *n* indicates the number of neurons. Results are for MC training and HC testing, using four orientations for training and testing. Panels one and three: classifier trained on '+' orientations and tested on 'x' orientations; panels two and four: classifier trained on 'x' orientations and tested on '+' orientations. Same conventions as Figure 4.

The marked difference in accuracy between the zero versus straight angle decoding and the right versus left angle decoding in aSTS was not trivially due to differences in mean response strength since there was no consistent effect of head-body orientation angle on mean response strength across poses (Friedman ANOVA; all p-values for both poses and centerings > 0.1).

The left and right angles have the same absolute angle between head and body (90°), only the sign of the angle differs with respect to the anatomical midline (90° versus -90°). The confusion

between these configurations suggests that this neuronal population is relatively insensitive to the

sign of the angle. This was confirmed by other decodings (Figure 4C; Figure S1C). We observed a

close-to-chance decoding of 45° versus -45° angles and 135° versus -135° angles, unlike the better

classification accuracy for the same 90° differences between angles of 0° and 90° or 180° and -90°,

the latter but not the former differing in the absolute angle between head and body. Other

configurations that differ in absolute angle also produced good decoding performance: 45° versus -

135° and 135° versus -45°, better than the left versus right angles, which differed only in sign (Figure

4C; Figure S1C).

One explanation of the poor decoding for the right versus left angles is that in those

conditions either the body or the face is at a lateral orientation. Neurons in anterior IT are less

sensitive to differences between mirror-symmetric images (Freiwald and Tsao, 2010; Rollenhagen

and Olson, 2000). If such neurons contribute to the computation of the head-body orientation angle,

poor decoding of right and left angles can be expected since these images are, to some extent,

mirror-symmetric. Therefore, we assessed the presence of mirror-symmetric tuning for the head

and/or body in the same sample of aSTS neurons used for the decoding. To assess this for the head,

we first averaged the responses, normalized by the maximum responses across the 64 stimuli, for the

8 monkey images having the same head (but a different body) orientation and then computed the

distance matrix using the 1-Pearson r distance metric. Similarly, to assess mirror-symmetric tuning

for body orientation, we averaged the responses to the monkeys having the same body orientation.

The distance matrices showed evidence for mirror-symmetric responses for the lateral head and

body orientations for P2 and face orientations of P1 (Figure S3). Mirror symmetry was less the case

for the body of P1, which fits the poorer image symmetry of the body in the case of the P1 lateral

views. Thus, the lower sensitivity for mirror-symmetric heads or bodies may have contributed to the

lower decoding of head-body angles that differed only in their sign. However, the difference in

decoding accuracy between the zero versus straight angle and the left versus right angle, especially

when training on the 'x' orientations and testing on '+' orientations (Figure 5), cannot be fully

attributed to the mirror symmetry of the avatar images.

<u>Decoding head-body orientation angle from the summed responses to body and face versus</u>

<u>from responses to the whole body (experiment 2)</u>

We obtained a second data set (experiment 2) from the same subjects in which we could test

the main findings of experiment 1 for a larger sample of neurons. Additionally, we evaluated whether

the angle between the head and the body could be computed from their individual orientations

when presented in isolation. We compared the responses to 16 head-body orientation configurations

to the responses to the heads and bodies shown in isolation with the same size and at the same

visual field location as in the monkey configuration. We recorded with V probes in mSTS and aSTS,

overlapping the monkey patches defined by fMRI (Figure 1A; Methods). The stimulus set included 16

monkey-centered images of the P1 avatar, crossing 4 head orientations and 4 body orientations (the

frontal, the back view, and the two lateral views), yielding 16 conditions. In addition, we included

images of the isolated heads, presented at the same position as in the avatar (16 conditions), and of

the 4 body orientations (Figure 6A).

Figure 6. Decoding of head-body angles in the Sum vs. Configuration test (Experiment 2). (A). Stimuli of Experiment 2: heads and bodies presented at four orientations (frontal, back, and the two lateral ones; B+H), along with all head-body configurations (M). Images were monkey-centered and the isolated heads and bodies were positioned to match their corresponding avatars. The red square indicates the fixation target. (B). Decoding accuracies for head-body configurations (M) and head-body sums (B+H) with zero, straight, leftward, and rightward right angles of head and body (trained pooling the four orientations (A)), for aSTS and mSTS, *n* indicates the number of units. Same conventions as Figure 4.

Spike sorting yielded 497 aSTS (M1 = 280; M2 = 217) and 1856 mSTS (M1= 804; M2 = 1052) units that responded selectively to the 16 monkey conditions (Methods). We decoded the head-body orientation angle for differences of 90 and 180° between angles of the head and body orientation (zero versus right angle, zero versus straight angle, …, etc), using the same procedure as in experiment 1. We found a highly similar pattern (n=430 units; equated across the two regions and subjects; randomly sampled per subject per resampling; Figure 6B) compared to that obtained with

the smaller sample of single units of experiment 1 (Figure 4B). For aSTS, the classification accuracy

for the zero versus straight angles was about 0.9 (Figure 6B). Again, classification accuracy for the

right versus left angles, differing only in their sign, was much lower than that obtained for zero versus

right angle, or straight versus left angle, differing in absolute angle (Figure 6B; $p < 0.01$). The

classification accuracy for mSTS units was again lower than for aSTS (Figure 6B), although no

generalization across centering conditions was required for this decoding. Importantly, in mSTS there

was no difference in decoding accuracy for the zero versus straight angle and right versus left angle.

The reliability of the responses of the units was similar for aSTS (median Spearman-Brown corrected

split-half Pearson correlation; M1: 0.70; M2: 0.69) and mSTS (median corrected r: M1: 0.72; M2:

0.71) and therefore cannot explain the difference in classification accuracy between the two regions.

Having replicated the main findings of experiment 1 (Figure 4B), we assessed whether the

simultaneous presentation of the head and body was necessary to decode the angle between head

and body orientation. To do this, we summed the net responses to the isolated head and body,

presented at the same locations as in the whole-monkey image, for individual trials, thus obtaining 8

head-body sums per orientation and angle (zero, straight, and the left and right angles) of head and

body orientation (Figure 6A). We decoded the angle using the face-body sums, employing the same

procedure and units as for the whole monkey. For aSTS, the classification accuracies for the sums

were markedly lower than those obtained for the monkey images (Figure 6B), and no difference in

classification accuracy between the zero versus straight angles and left versus right angles was

present for the sums (all p's > 0.4). The lower classification accuracy for the head-body sums is not

due to a lower reliability of the summed responses compared to the whole-body responses

(Spearman-Brown corrected split-half Pearson correlations; median whole-body r: M1: 0.7; M2: 0.69;

sums r: M1: 0.76; M2: 0.66). These data suggest that the difference in classification accuracy

between the zero versus straight angles and left versus right angles requires the simultaneous

presence of the head and body. For mSTS, the classification accuracies for the head-body sums were

similar to those obtained for the monkey images (Figure 6B), suggesting no genuine encoding of the

head-body angle in this region.

<u>Head- and body-selective aSTS units show similar decoding patterns of head-body orientation</u>

<u>angle (experiment 2)</u>

To assess whether the head/body selectivity matters for the decoding of the head-body

orientation angle, we subdivided our aSTS units into groups based on their head/body selectivity

(Methods; Figure S4). Head- and body-selective aSTS units showed, as a population, a similar

decoding bias for zero versus straight angles and both confused the left and right angles (Figure S4).

<u>Effect of body-inversion on responses and view selectivity (experiment 3)</u>

We determined whether body inversion impairs the decoding accuracy of the head-body

orientation angle. Inverting the body affects the global body configuration, keeping the local features

intact (except for an 180° change). First, we assessed for the first time whether STS neurons show a

body-inversion effect in response strength and/or capacity to discriminate between views of a pose.

For this, we obtained a new data set of aSTS and mSTS responses to the upright and inverted poses

of the avatar. We employed the same two poses as for the single units (Figure S5A). For each pose,

we presented 8 orientations (steps of 45°) of 4 head-body orientation angles (zero, straight, and left

and right angles) and this upright and inverted (Figure S5A). The images were monkey-centered, with

the heads shifted towards the fixation point, as in experiment 1 (Methods; Figure S5B for examples).

In addition, we included heads and bodies shown in isolation for a subset of these conditions

(Methods; Figure S4C).

We recorded with V-probes at the same mSTS and aSTS locations (3 anterior-posterior levels

in each region) as in experiment 2 (Figure 1A). First, we selected for each pose those units that

responded to the head, headless body, or monkey, either upright or inverted (Methods), having the

anatomically possible zero head-body angle (see Figure S5 for examples). Then, for each pose, we

selected the orientation ('preferred orientation') that produced the maximum response for the

stimuli corresponding to that orientation, irrespective of whether the stimuli were upright or

inverted. The latter ensured an unbiased selection of the preferred orientation with respect to the

upright versus inversion variable. We observed for each pose, in each region of each subject, a higher

mean response for the upright compared to the inverted preferred orientation of the whole monkey

(Wilcoxon signed rank tests on data pooled across monkeys: p's in Figure 7C; population PSTHs in

Figure S6). Selecting units that responded either to the upright or inverted heads showed less

consistent inversion effects for the head than observed for the monkey: upright heads showed across

subjects consistently and significantly greater responses than to inverted heads only for P2 in mSTS

but not for the other pose or region (Figure S7). The same held for the headless body for those units

that responded to either the upright or inverted bodies (Figure S7). Thus, the inversion effect was

consistently present for the head-body configurations, and less so for the face and body only.

Behaviorally, the body inversion effect corresponds to a higher discriminability of upright

bodies. To assess this for our STS neurons, we decoded the 8 whole-body orientations employing

units responding to the whole-body images with a zero body-head angle, either upright or inverted.

The decoding was performed separately for upright and inverted images. This was done for each

pose, drawing randomly in each of the 1000 resamplings 60 units per region (Methods). The mean

decoding accuracy was significantly higher (all p-values < 0.01; Methods) for upright compared to

inverted monkeys for both poses in aSTS (P1: upright versus inverted: 0.89 versus 0.82; P2: 0.89

versus 0.71, with standard deviations SD = 0.02) and mSTS (P1: 0.89 versus 0.79; P2: 0.93 versus 0.75;

SDs ranging between 0.01 and 0.02).

<u>Decoding of head-body orientation angle is impaired for inverted bodies (experiment 3)</u>

Next, we asked whether the decoding of the head-body orientation angle is affected by

inversion. For each pose, we selected units that responded to our stimulus set of 8 orientations of 4

angles for the upright or inverted avatar (Methods). Then, we decoded the head-body orientation

angle from the aSTS units using the same classification procedure as above. For both poses (n=460

responsive aSTS units; an equal number of units randomly sampled per subject per resampling), the

upright images showed again a higher classification performance for the zero versus straight angles

and the lowest for the left versus right angles (Figure 7B). The classification accuracy dropped

significantly (zero versus straight angle; all p-values < 0.01; n=100 resamplings; Methods) for the

inverted bodies (Figure 7B), and this drop in classification accuracy tended to be larger for the zero

versus straight angle decoding than for the left versus right angle decoding (the difference between

upright and inverted left versus right angle decoding not significant for the two poses; p = 0.2 and

0.18). Thus, the decoding of the head-body orientation angle is impaired when inverting the monkey

image.

Figure 7. Effect of head-body configuration inversion (Experiment 3) (A). Stimuli of the head-body configurations of P1 with zero and straight angles, and leftward and rightward right angles, at eight orientations, upright and inverted. **(B).** Decoding accuracies for the head-body configurations with zero, straight, leftward, and rightward right angles in aSTS, for poses P1 (top panels) and P2 (bottom panels), *n* indicates the number of units. Solid and dotted lines represent data for upright and inverted configurations, respectively. Same conventions as Figure 4. **(C).** Responses to the preferred orientation of upright or inverted head-body configurations for aSTS and mSTS and P1 and P2 (M1 (dark green); M2 (light green)), for the monkey-responsive units. Medians (diamonds) and the number of units per subject are shown in corresponding colors.

P values of Wilcoxon signed-rank tests (data pooled across subjects) testing the difference in response between upright and

inverted are shown (red: significant).

The stronger inversion effect for the zero versus straight angles decoding suggest that the

head-body angle signal in aSTS depends on the global head-body configuration and is not driven

merely by inversion-invariant local features. This aligns with the marked generalization across head-

body configuration orientation that we observed in experiment 1 for P1 (training using 4 orientations

and testing using the other 4); such across-orientation generalization would not be expected if local

features of individual images were employed by the classifier.

Discussion

We showed here that head and body orientations interact in ventral bank STS neurons, with

some neurons being tuned to a particular orientation of *both* the head and the body. Importantly,

the head-body orientation angle, and not only the orientations of the head and the body per se, can

be robustly decoded from a population of STS neurons. This decoding was greater for aSTS compared

to mSTS neurons. There is a marked bias in the decoding of head-body orientation angle from aSTS

with better decoding of absolute angle differences (e.g. 0 and 180° angles). Head-body orientation

angles that differ only in their sign with respect to the anatomical plane (e.g. 90° versus -90°angles)

can be decoded less from aSTS neurons. Also, we provide here the first demonstration of an

inversion effect for a body with a head in the macaque STS: the response strength, orientation

decoding, and head-body orientation angle decoding are less for inverted compared to upright

monkey avatars.

We could decode the orientation of the head-body configurations, which is an ecologically

important property. From the same population of neurons, we could also decode the angle between

the head and the body, irrespective of the orientation of the configuration (Figure 4A). This can be

viewed as a sensitivity for a particular head-body configuration that is defined by the orientation of

the head *and* body, and this independent of the overall orientation of the agent. Remarkably,

decoding of head-body configuration in the aSTS confused to a greater extent configurations in which

the head and body orientation angles differ only in their sign (e.g. head turned to the left versus to

the right with respect to the body). The latter is quite a robust effect in the aSTS: we observed it in

three different experiments. Images, especially of the standing pose, that differ only in the sign of the

head-body angle are also to some extent mirror-symmetric. Our population of single aSTS neurons

showed evidence of a lower sensitivity of mirror-symmetrical views for heads and bodies. Thus, one

factor that may have contributed to the lower decoding of head-body orientation angles that differ

only in sign is mirror symmetry. The highest decoding of the head-body angle occurred for the zero

versus straight angles. Basically, this is a classification of a frequently observed head-body

configuration, i.e. the same orientation of head and body, versus an anatomically impossible angle,

at least for primates. In fact, all well-decodable absolute differences in angles consisted of a possible

versus an impossible angle, while angles that differ only in their sign are either both anatomically

possible or both impossible. Thus, this anatomical possibility factor may explain the better decoding

for configurations that differ in their absolute angle. However, dissociating this factor from absolute

versus signed angle difference, or mirror-symmetry, is not trivial, since these 3 factors covary in

monkey configurations when manipulating the orientation of the head and body.

The decoding of whole-monkey orientation, which can be based on head and/or body

orientation, was present in mSTS and aSTS, which aligns with previous studies (Bao *et al.*, 2020;

Freiwald and Tsao, 2010; Kumar *et al.*, 2017). However, the decoding of head-body orientation angle,

especially when requiring location tolerance, was weaker in mSTS than in aSTS. This agrees with

previous work that suggested the presence of head-body integration in the anterior but less so in the

posterior temporal cortex (Fisher and Freiwald, 2015; Hu *et al.*, 2020; Zafirova *et al.*, 2022).

We observed no difference in response strength between anatomically possible and

impossible configurations. This differs from a previous study in which responses to head-body

configurations in which the body and head were misaligned were lower than those for a natural

head-body configuration (Zafirova *et al.*, 2024). However, that study targeted a patch that was

defined in fMRI by contrasting aligned versus misaligned head-body configurations, which only

partially overlapped the fMRI-defined monkey patch of the present study. Furthermore, the present

study examined head-body orientation interactions, while the previous one head-body relative

position interactions, which are fundamentally different. Nonetheless, the present study suggests

that mSTS and aSTS neurons respond equally strongly to anatomically possible and impossible head-

body configurations, at least when these are defined by the head-body orientation angle. However,

as a population, they can distinguish strikingly well the possible and impossible configurations since

these differ in their absolute angle (see above).

Our work demonstrates for the first time an inversion effect for bodies *with* heads in the

ventral bank of the STS. Previous studies reported a higher response to upright compared to inverted

bodies in the STS (Ashbridge *et al.*, 2000; Popivanov *et al.*, 2015), but the neurons in these studies

were searched with upright bodies. However, in our study, a search bias cannot explain the body-

inversion effect since we selected responsive units using both upright and inverted images. A body-

inversion effect has been observed in both humans and monkeys at the behavioral level (Griffin and

Oswald, 2022; Matsuno and Fujita, 2018). Some but not all human studies (see (Griffin and Oswald,

2022) for a meta-analysis) observed a stronger inversion effect for a body with than without a head.

Our data suggest also a more consistent inversion effect for head-body configurations compared to

headless bodies and heads in STS.

The body-inversion effect likely results from greater exposure to upright than inverted bodies

during development, although, as for the face-inversion effect, this remains to be proved (Kobylkov

and Vallortigara, 2024). Our monkeys were not exposed to the anatomically impossible head-body

orientations, but still, the responses to these configurations were similar to those to anatomically

possible configurations. This apparent discrepancy between the head-inversion effect and the

anatomically (im)possible head-body orientation configuration may result from both head and body

being in an infrequent orientation in the case of inversion while for the impossible head-body

orientation configuration either the head or the body, but not both, might be at an infrequently

exposed orientation.

In conclusion, we demonstrated that head and body orientations interact in the macaque

STS. The population of aSTS neurons can provide a reliable signal for the recognition of particular

head-body angles, especially between those that differ in their absolute angle with respect to the

anatomical midline. This signal is less reliable for inverted head-body configurations, showing a

strong inversion effect.

Materials and methods

<u>Subjects</u>

The experiments were performed with the two male rhesus monkeys (M1 and M2; *Macaca mulatta*, 6-7 years, 11-12 kg) which took part in our fMRI study (Zafirova et al., 2022). They were implanted with an MR-compatible plastic headpost for fixing the head during training, scanning, and recording, and a plastic custom-made recording chamber allowing a slightly oblique approach to the fMRI-defined monkey patches (for details, see Data Acquisition). All experimental procedures and animal care complied with national and European regulations and were approved by the Animal Ethical Committee at KU Leuven.

<u>Data acquisition</u>

Before the three electrophysiological experiments, reported here, we performed a functional MRI experiment in the same monkey subjects to determine the location of patches activated more by images of monkeys compared to objects (for details see (Zafirova *et al.*, 2022)). The fMRI localizer had a block design with a Brown-noise backgrounds-only condition (baseline) and the following 6 conditions, 20 images each: monkeys, monkey faces, monkey headless bodies, and three groups of balanced manmade objects for the monkeys, the faces and the bodies, respectively (centered on the Brown-noise backgrounds). We employed the contrast monkeys versus monkey control objects ($p <$.05, Family-wise error corrected) to define the monkey-selective regions (monkey patches) in each subject`s IT. In experiment 1, we targeted single units in the monkey patches in the ventral bank of the STS at two anterior-posterior levels (Figure 1A). In experiments 2 and 3 we recorded from a wider region, which overlapped with the two monkey patches and the recording locations of experiment 1. This was done in the context of another electrophysiological study in which we mapped the

responses to heads and body parts. In these two experiments, the penetrations were at 3 anterior-

posterior levels, spaced 1 mm apart in aSTS and mSTS. At each anterior-posterior level, we made

penetrations in 1 mm steps from the lip of the STS towards the fundus. This resulted in a total of 18

and 27 penetrations in aSTS and mSTS, respectively, for both subjects. We recorded from the left

hemisphere of M1 and the right hemisphere of M2 in all experiments.

Before the recording experiments, the subjects underwent an anatomical MRI (3.0 T full-

body Siemens scanner Magnetom Prisma Fit; magnetization-prepared rapid acquisition with gradient

echo sequence; 0.6 mm isotropic voxel resolution). We inserted into the recording chamber grid

(until the dura) long plastic capillaries filled with MRI opaque copper sulfate ($CuSO_4$) at

predetermined positions and filled the recording chamber with a Gadoteric Acid (Dotarem) solution

for visualization of the chamber and the grid. The activations were co-registered to the anatomical

scans   (Freesurfer   http://surfer.nmr.mgh.harvard.edu/;

FSLeyes, http://git.fmrib.ox.ac.uk/fsl/fsleyes/fsleyes/, SPM12, RRID: SCR_007037) and the co-

registration was verified by visual examination. The recording locations in the sagittal and coronal

planes were verified with structural MRIs that visualized the guide tube tracks. The dorsal/ventral

locations were based on observation of the white/gray matter transitions and silence associated with

the STS during the recordings.

The position of one eye was continuously tracked with an infrared video-based tracking

system (SR Research EyeLink; sampling rate 1 kHz). Stimuli were displayed on a VPixx LCD display

(1920 x 1080 resolution; 120 Hz) at a distance of 57 cm from the subjects' eyes. The onset and offset

of the stimulus were signaled utilizing a photodiode detecting luminance changes of a small square in

the corner of the display (not visible to the animal), placed in the same frame as the stimulus events.

A digital signal processing-based computer system developed in-house controlled stimulus

presentation, event timing, and juice delivery while sampling the photodiode signal, vertical, and

horizontal eye positions, spikes, and behavioral events.

Single-unit recordings in experiment 1 were performed with epoxylite-insulated tungsten

microelectrodes (FHC) lowered with a Narishige microdrive into the brain using a stainless-steel

guide tube that was fixed in a custom-made grid positioned within the recording chamber. After

amplification and filtering, spikes of a single unit were isolated online using a custom amplitude- and

time-based discriminator.

Spiking activity was recorded in experiments 2 and 3 with 24-channel V-probes (Plexon). The

V-probe was lowered into the brain through a stainless-steel guide tube with a Narishige microdrive.

The same grids were employed in experiment 1. After reaching the recording target, we waited for at

least 30 minutes before the tests were run to obtain stable recordings. During that waiting period,

the monkey could watch movies. We employed an INTAN RHD recording system to record spiking

activity. The signal of each channel was bandpass-filtered online between 500 and 7500 Hz. The

photo-diode signal from the display was fed into the INTAN recording system, allowing alignment of

stimulus events and the condition and trial data saved by the DSP-based computer system. We

employed Plexon Offline Sorter to sort the spiking activity of each channel into single units and small

clusters of (unsortable) units (multi-units). In this paper, we use the term 'units' for both single- and

multi-units, combining single and multi-units in the analyses of experiments 2 and 3.

<u>Stimuli and Experiments</u>

We report here three experiments designed to investigate the interaction between the

orientations of the head and the body. The first experiment included a Selectivity test to assess the

selectivity of the neurons recorded in the main Head-Body Orientation test. The Selectivity test

employed naturalistic greyscale static images centered on a uniform grey background. All subsequent

tests employed static images of realistic 3D monkey avatar in color,

(https://www.turbosquid.com/3d-models/monkey-fur-rigged-1493219) on the same uniform grey

background.

First, the avatar was positioned in two emotionally neutral natural monkey poses - a sitting

and a standing one (P1 and P2, respectively; Figure 2A). Using a 3D avatar instead of actual monkey

images allowed us to manipulate independently the orientation of the head and the body, for the

two poses. The head and body orientations were obtained by rotation around the vertical axis of the

avatar. The maximal vertical or horizontal extent of the images was set to 6°. Heads and headless

bodies of some of the configurations were presented in isolation at their original location which

allowed for direct comparison with the corresponding configurations. Details about the head and

body orientations and the centering of the configurations for each experiment are described below.

Experiment 1

Selectivity test

To assess the category selectivity of the neurons of experiment 1, we employed 60 stimuli

from a previous fMRI experiment (for the details about the stimuli, see (Zafirova *et al.*, 2024)). In the

Selectivity test, we presented 10 images of monkeys in diverse natural postures and orientations,

together with 10 images of headless monkey bodies and 10 images of monkey faces, all cropped

from different natural photographs of rhesus macaques. Additionally, we selected 10 control objects

for each of the groups (examples in Figure 1). The objects matched the face-, body-, and monkey

images as best as possible, as estimated by different shape-descriptive parameters (for details about

the images and the matching procedure, see (Zafirova *et al.*, 2024)). The low-level characteristics, like

luminance and contrast, were equated across the whole set (Matlab SHINE toolbox (Willenbockel *et*

al., 2010). The greyscale images were resized to fit a 6°x6° stimulus window and gamma-corrected.

Head-body Orientation test

In this test, we manipulated the orientations of the head and the body independently,

starting from the frontal view (0°) with eight 45° consecutive steps, generating 64 head-body

combinations for the sitting and the standing pose (P1 and P2; Figure 2A). In terms of angle between

the orientation of the head and the body, this resulted in eight head-body configurations with angles

0° (zero angle), 45°, 90° (rightward right angle), 135°, 180° (maximum possible absolute angle:

straight angle), -135°, -90° (leftward right angle), and -45° between the head and the body. Each of

these eight head-body configurations was spatially rotated with steps of 45° around its vertical axis,

that is, each configuration was presented in eight different orientations. Ultimately, we had eight

orientations of the head and the headless bodies (0° to -45° with steps of 45°), eight head-body

configurations with angles 0° to -45° with steps of 45° angle between the head and the body, and

eight different orientations of each of these configurations (rotated from 0° to -45° with steps of 45°

around their vertical axis). As a convention, we will further refer to the orientation of the head-body

configurations by the orientation of the body (torso). The head-body configurations with an angle

between the head and the body more than an absolute angle of 90° were anatomically impossible.

The configurations were centered on the head (HC condition) or the monkey avatar (MC

condition), for each of the two poses P1 and P2 (examples in Figure 2B). In the HC condition, the

maximum extent of the bodies was 2.8° and 3.3° from the fixation point for P1 and P2, respectively.

As we were interested in the possible head-body orientation interactions, and to ensure that the

orientation of the head was clearly visible, the images were shifted 1° down in the lower visual field

in the MC condition, so that the heads were closer to the fixation point. That way, the maximum

extent of the head was 1.6° and 2.6° from the fixation point for P1 and P2, respectively, in the MC

condition.

Additionally, heads and headless bodies from the configurations with zero and straight angles

(0° and 180°; anatomically possible and impossible) between the head and the body of P1 and P2

were presented in isolation at the same locations as in the head-body configurations. We did that for

four orientations of the two configurations: the frontal, the back, and the two lateral ones (0°, 180°,

90° and -90°). We had four orientations of the headless bodies for the MC condition and four

orientations of the head for the HC condition (one from each orientation; Figure 2C). These were

compared to the head-body configuration with the same body orientation for the MC condition and

to the head-body configuration with the same face orientation in HC conditions. We presented two

head orientations - one from the zero angle and one from the straight angle configurations for each

of the four body orientations in the MC condition, and two body orientations - one from the zero

angle and one from the straight angle - for each of the four head orientations in the HC condition.

Thus, we had four headless bodies and eight heads in the MC condition and four heads and eight

bodies in the HC condition, for P1 and P2 (Figure 2C), or altogether standalone heads and bodies

from the zero and straight angle configurations at four orientations, for each centering and pose. In

total, we had 152 images (64 + 12) for the MC and (64 + 12) for the HC condition) for each pose.

Experiment 2

Sum vs. Configuration test

In the second experiment, we presented only the sitting pose (P1) of the monkey avatar,

centered on the whole body. Additionally, we employed four out of the eight head-body

configurations: the zero angle (0°), the straight angle (180°), and the right and left angle

configurations (90° and -90°). These were chosen because experiment 1 showed that the 0°-180° pair

and the 90°- 90° pair showed the largest difference in the decoding of the head-body orientation

angle. Each of these configurations was presented at four orientations: the frontal, the back, and the

two lateral ones (0°, 180°, 90°, and -90°), determined by the viewpoint of the body (rotated with

steps of 90° around its vertical axis; Figure 6A).

As before, heads and headless bodies from the configurations were presented in isolation at

their original locations. We did that for the four orientations. Thus, we had four orientations of the

headless bodies. Here, we presented four head orientations from each of the four angles: the zero,

the straight, and the left and right ones, resulting in overall sixteen standalone heads and four

headless bodies (Figure 6A). Overall, the Sum vs. Configuration test employed 36 images (16

configurations and 20 images of heads and headless bodies).

Experiment 3

Upright vs. Inverted test

In the last experiment, we presented both the sitting and the standing pose (P1 and P2) of

the monkey avatar. We employed the same four head-body configurations as in experiment 2. Each

of these configurations was presented at the eight orientations (body rotated with steps of 45°

around its vertical axis; Suppl. Figure 4A). Crucially, we introduced one additional manipulation: all

images were presented either upright or flipped around their horizontal axis (inverted; Figure S5A).

All were centered on the monkey avatar, but as in the Head-body Orientation test, we wanted to

ensure that the orientation of the heads was visible. To that end, we shifted the upright images 1°

down in the lower visual field and the inverted images 1° up in the upper visual field, so that the

heads were closer to the fixation point (Figure 5B).

We presented also the heads and headless bodies from the zero and straight angle head-

body configurations of the two poses in isolation at their original locations. We did that for four

orientations: the frontal, the back, and the two lateral ones (0°, 180°, 90° and -90°). Thus, we had

four orientations of the headless bodies per pose. We presented two head orientations from each

configuration head-body angle, resulting in four headless bodies and eight heads for a pose (Figure

S5C). Overall, this test included 176 images (the upright and inverted configurations of the two poses

and their heads and bodies).

<u>Procedure</u>

In experiment 1 (Selectivity and Head-body Orientation test), we recorded well-isolated

single neurons, while single and multi-units were recorded with V-probes in experiment 2 (Sum vs.

Configuration test) and 3 (Upright vs. Inverted test). In each subject, we recorded in (experiments 1-

3) and surrounding (experiments 2-3; see Results) patches that were activated more by images of

monkeys compared to monkey control objects, in the anterior (aSTS) and mid ventral bank (aSTS) of

the superior temporal sulcus (Figure 1A). To control for potential order effects, we recorded first in

aSTS in one subject, and the mSTS in the other for each experiment.

In experiment 1, we recorded in the Head-body Orientation test well-isolated single neurons

that responded to monkeys, monkey faces *or* bodies in the preceding Selectivity test. All stimuli in all

experiments were presented for 250 ms, with at least 600 ms interstimulus interval, on a uniform

grey background. The subjects had to fixate a small (0.2°x0.2°) red target. They received juice

rewards for maintaining fixation in a 2°x2° window around the target for 300 ms before the stimulus,

during, and 300 ms after the stimulus. Trials in which the monkey failed to fixate continuously during

this period were not analyzed further. The juice rewards were administered at fixed intervals in

experiment 1 and after a completed trial in experiments 2 and 3. All stimuli of a test were presented

pseudorandomly in blocks, and the order of the stimuli within a block was random. A stimulus of an

aborted trial was presented again at a random moment during the same block.

<u>Data analyses</u>

All analyses were performed only on those units for which we had 5 unaborted trials per

stimulus for the Selectivity test and 8 unaborted trials per stimulus for all other experiments, and on

net firing rates (FRs) unless stated otherwise. Net FRs were computed by subtracting the mean FR

100 ms before the stimulus onset (baseline window), from the FR in a response window of 50 ms to

300 ms after stimulus onset.

<u>Unit selection</u>

Experiment 1

To determine the responsiveness of a neuron, we computed mean firing rates for each trial

in two windows: a 200 ms baseline window that started 200 ms before stimulus onset and a 300 ms

response window that started 50 ms after stimulus onset. We ran split-plot ANOVAs with within-trial

factor baseline vs. response window and between-trial factor stimulus. We considered as responsive

only those neurons for which (1) either the main effect of the repeated factor or the interaction of

the two factors was significant ($p < 0.05$), and (2) their response was excitatory. Neurons that were

irresponsive in the Selectivity test were excluded from the analyses of the Head-body Orientation

test (4 out of 102 recorded aSTS and 5 out of 105 recorded mSTS neurons). The tests of the two

poses were analyzed separately for responsiveness. Additionally, for each particular analysis, we

excluded neurons that were irresponsive to all conditions of that analysis (e.g. separately for each

pose and centering). These selections are described in the Results.

To compute the monkey-sum index (MSI; see below) we selected for each unit the monkey

and the corresponding isolated head and body images ('cases') for which there was a significant

response to either the monkey, the head, or the body. The significance was tested with a Wilcoxon

rank-sum test, comparing for each image the firing rate in baseline and response windows. The

response needed to be excitatory to be considered for further analyses.

Experiments 2 and 3

The responsive units of experiments 2 and 3 were selected using more stringent criteria since

these were not isolated online using traditional single-unit recording methodology in which the

electrode is moved for each unit to isolate it and responses are examined directly online. First, each

unit was tested with a Split-plot ANOVA with within-trial factor baseline vs. response window and

between-trial factor stimulus, as for experiment 1. Second, we employed a Kruskal–Wallis test ($p <$

0.05), performed on the mean raw firing rates in the response window per trial and for all stimuli of a

test, to exclude non-selective units. Third, only units with a z-score > 3 for at least one stimulus were

considered further. Z-scores for each stimulus were computed, as follows: first, we averaged the

mean firing rates in the baseline window (meanBaseStim) and the response window (meanRespStim)

for the eight trials of each stimulus. Then, we calculated the mean (meanBaseAll) and the standard

deviation of the baseline (stdBaseAll) across all stimuli. Finally, we subtracted the mean baseline

from the averaged response of each stimulus and divided this difference by the standard deviation of

the baseline: (meanRespStim-meanBaseAll)/stdBaseAll). The units in each analysis were subjected to

the three criteria (split-plot ANOVA, the Kruskal–Wallis test, and the z-scoring).

<u>*Reliability*</u>

We estimated the reliability of the responses of each responsive unit by randomly splitting

for each stimulus the 8 trials in two groups of 4 trials, and correlating the raw firing rate in the

response window between the halves across stimuli. The random splitting was performed 100 times.

The reliability corresponded to the mean of the 100 Spearman-Brown corrected Pearson correlation

coefficients between the two halves. In some analyses, we normalized the correlation between two

sets of responses (e.g. two centerings) by the reliability of the two responses by dividing the original

correlation by the geometric mean of the reliability of the two response sets.

Monkey-sum index MSI

For each orientation, pose, and centering combination in which the neuron responded

significantly either to the monkey (M), head (H), or body (B) (see unit selection above), we computed

a Monkey-sum index, MSI, using net FRs, as follows:

MSI = (Response M- (Response B+ Response H)) / (abs(Response M)+abs(Response B +

Response H)).

Interaction of head and body orientation tuning

For each responsive single-unit of experiment 1, we assessed statistically the interaction

between head and body orientation for each pose and centering with a two-way ANOVA with factors

of head and body orientation. We performed two ANOVAs, one using the raw spike counts per trial

and a second one using the following transformation of the raw spike counts per trial: square root

(spike count + 3/8). The latter serves to stabilize the variance across conditions for Poisson-like

distributions. Both produced similar results. We report the outcome of the ANOVAs using the square

root transformed data. We isolated the interaction of head and body orientation for each neuron by

computing the residuals of the square-root transformed responses from an additive model in which

the responses to a head-body configuration are the sum of the mean responses to the body

(computed by averaging responses across the 8 head orientations) and head (averaged across the 8

body orientations) orientations. The additive model response to the monkey configuration with a

head-body orientation (i, j) was modeled as:

Mean response head orientation (i) + mean response body orientation (j) – mean response to

the 64 stimuli.

The residuals were computed for both centerings and then correlated across the centerings

per neuron and pose combination.

Decoding analyses

We performed a series of decoding analyses (see Results) on the raw responses (unless

stated otherwise with custom Matlab code employing a linear kernel support vector machine (SVM)

classifier. For the binary classification (pairs of head-body orientation angles) we employed the

Matlab function *fitcsvm*. For multiclass classification, we employed the Matlab function *fitcecoc*. The

regularization parameter C was 1 (default) in all cases. Per stimulus, we produced 8 pseudo-

population vectors after permuting the order of the 8 trials per unit. We performed 8-fold cross-

validation, training on 7/8 of the pseudopopulation vectors, and the remaining ones were employed

for testing. The reported classification scores are means of the percent correct classifications for the

test vectors of 100 (or 1000; see below) resamplings of trials (experiments 1-3) and units

(experiments 2-3). For the main decodings, we ran the same decoding procedure after permuting the

stimulus labels, creating null distributions of chance performance. These were run 200 times, using

the same number of resamplings. We plot the range of the mean classification scores (mean of 100

resamplings) obtained after label permutation in the Figures.

Decoding of head-body orientation angle

We decoded the head-body angle for pairs of angles (e.g. zero versus the maximum possible

straight angle), for all possible orientations of the respective head-body configuration. To do this, we

created two classes of trials, each containing responses to the same angle (e.g. zero versus straight

angle). The head and body orientations per se were equal between the two classes. Since we had 8

trials of each of 8 (experiments 1 and 3) or 4 (experiment 2) head (and body) orientations per class,

there were 64 trials per class in experiments 1 and 3, and 32 trials per class in experiment 2. For each

orientation of a certain angle, we kept 7 trials for training, yielding in total 56 (7x8 orientations;

experiments 1 and 3) or 28 training trials (experiment 2) per class. Test trials consisted of one trial

per orientation, i.e. 8 trials. This was repeated eight times for each resampling, each time using a

different trial as a test trial, i.e. the training and testing trials were chosen pseudorandomly for each

class separately. This procedure was critical to avoid unbalancing of head and body orientation

between classes, which can produce artefactual results (e.g. classification scores well below chance

levels). The pairs of angles employed in the decoding are described in the Results.

We will illustrate our approach for the binary classification of two head-body angles for

experiment 1: zero versus straight angle ($0°$ versus $180°$; Figure 4A). We pooled 8 trials of the 8

orientations of each angle (e.g. zero angle configuration shown at a 0° (frontal) orientation, 45°, 90°

(right lateral), 135°, 180° (back view), -135°, -90° (left lateral) and -45°; Figure 4A), providing 64 trials

per angle. We employed an eightfold cross-validation approach, sampling in each cross-validation

step 7 trials of each orientation per angle, ensuring balanced sampling across orientations. The same

was done for the other angle (e.g. straight angle; Figure 4A). These 2 x 56 trials were then employed

to train the SVM classifier. Testing was performed using one trial per orientation per angle (in total 8

trials).

To assess the significance of the differences in classification scores between pairs of angles

(decoding of zero versus straight angle compared to decoding of the leftward versus rightward right

angles) we computed the difference in classification score between the two pairs for each resampling

and the percentile of 0 difference corresponded to the p-value. Note that the same set of neurons

was taken per resampling when comparing differences in classification accuracy between conditions.

In experiment 1, we performed two types of decoding: (1) cross-centering decoding, where

responses to monkey-centered images were used for training, and responses to head-centered

images for testing (and vice versa), and (2) within-centering decoding, where both training and

testing were within the same centering condition. We report the outcome of each. Additionally, we

conducted decodings using responses to four orientations for training (e.g., frontal, back, and the

two lateral views; '+' orientations) and responses to the other four orientations for testing (e.g. the

oblique views 45°, 135°, -135°, and -45°; 'x' orientations), and vice versa (Figure 5A).

For experiment 2, we performed also decoding of the summed responses to the isolated

bodies and heads. Thus, for each head-body configuration, we summed the response to the

corresponding head and body, using trials in which these stimuli were presented in isolation. Then,

the decoding was run on the summed responses, using the same procedure as described above for

the responses to monkey images, with two exceptions. First, as we employed summed head-body

responses, in this decoding analysis we used net firing rates (baseline subtracted) instead of raw

responses, to avoid inflating the response when computing the sums. Second, to prevent

unbalancing of head and body orientations across classes, testing trials were selected

pseudorandomly to ensure that head and body orientations were equally represented in the training

trials for both classes. The same balancing during cross-validation and the use of net firing rates was

also the case for the decoding of the angles for the monkey configurations (simultaneous presence of

head and body) in experiment 2.

Decoding of orientation

The main decodings in experiment 3 were conducted in line with the described default

procedure. Additionally, in experiment 3, we decoded the orientation of the monkey for the zero-

angle head-body configuration. This was done separately for the upright and inverted poses. We

employed 8-fold cross-validation, using 7 trials per orientation for training and one per orientation

for testing. In each of the 1000 resamplings, we randomly sampled 60 units from the population of

units (aSTS P1 n=332, P2 n=369; mSTS P1 n=1298, P2 n=1332; equated per monkey). This number of

units (equal to that of the experiment 1 decoding analyses) was used to avoid ceiling effects. The

mean of 10 classification accuracies (i.e., of 10 resamplings) was employed to obtain a distribution (n

= 100) of the differences in classification accuracy between upright and inverted images. The zero-

difference value fell outside the distribution of each of the two regions and poses. The reported

standard deviations of the classification accuracies are computed using also the means of 10

resamplings.

Acknowledgements: The authors thank I. Puttemans, A. Hermans, W. Depuydt, C. Ulens, S.

Verstraeten, J. Helin, S.T. Riyahi, M. De Paep and Y. Gürsoy for technical and administrative support.

This research was supported by Fonds Wetenschappelijk Onderzoek (FWO) Vlaanderen (G0E0220N),

KU Leuven grant C14/21/111, and the European Research Council (ERC) under the European Union's

Horizon 2020 research and innovation program (grant agreement 856495).

Author contributions: Conceptualization, R.V. and Y.Z.; methodology, R.V. and Y.Z.; investigation,

R.V. and Y.Z.; writing – original draft, R.V. and Y.Z.; writing – review & editing, Y.Z. and R.V.; funding

acquisition, R.V.; resources, R.V.; supervision, R.V.

Declaration of interests:

The authors declare no competing interests.

Data and code availability:

Data: https://osf.io/7t2e8/?view_only=d6b8c9e7715f40fdbc41a76c54da31dd

Codes: https://github.com/Yozafirova/head-body-orientation-interaction.git

References

Arcaro, M.J., Ponce, C., Livingstone, M., 2020. The neurons that mistook a hat for a face. *Elife* 9.
Ashbridge, E., Perrett, D.I., Oram, M.W., Jellema, T., 2000. Effect of image orientation and size on
object recognition: responses of single units in the macaque monkey temporal cortex. *Cogn*
Neuropsychol. 17**,** 13-34.
Bao, P., She, L., McGill, M., Tsao, D.Y., 2020. A map of object space in primate inferotemporal cortex.
Nature 583**,** 103-108.
Cox, D., Meyers, E., Sinha, P., 2004. Contextually evoked object-specific responses in human visual
cortex. *Science* 304**,** 115-117.
de Gelder, B., Van den Stock, J., Meeren, H.K., Sinke, C.B., Kret, M.E., Tamietto, M., 2010. Standing up
for the body. Recent progress in uncovering the networks involved in the perception of
bodies and bodily expressions. *Neurosci.Biobehav.Rev.* 34**,** 513-527.
Fisher, C., Freiwald, W.A., 2015. Whole-agent selectivity within the macaque face-processing system.
Proc.Natl.Acad.Sci.U.S.A 112**,** 14717-14722.
Freiwald, W., Duchaine, B., Yovel, G., 2016. Face Processing Systems: From Neurons to Real-World
Social Perception. *Annu.Rev.Neurosci.* 39**,** 325-346.
Freiwald, W.A., Tsao, D.Y., 2010. Functional compartmentalization and viewpoint generalization
within the macaque face-processing system. *Science* 330**,** 845-851.
Griffin, J.W., Oswald, F., 2022. A multilevel Bayesian meta-analysis of the body inversion effect:
Evaluating controversies over headless and sexualized bodies. *Psychonomic Bulletin & Review*
29**,** 1558-1593.
Hietanen, J.K., 2002. Social attention orienting integrates visual information from head and body
orientation. *Psychol Res* 66**,** 174-179.
Hu, Y., Baragchizadeh, A., O'Toole, A.J., 2020. Integrating faces and bodies: Psychological and neural
perspectives on whole person perception. *Neurosci Biobehav Rev* 112**,** 472-486.
Kobylkov, D., Vallortigara, G., 2024. Face detection mechanisms: Nature vs. nurture. *Front Neurosci*
18**,** 1404174.
Kumar, S., Popivanov, I.D., Vogels, R., 2017. Transformation of Visual Representations Across Ventral
Stream Body-selective Patches. *Cereb Cortex***,** 1-15.
Matsuno, T., Fujita, K., 2018. Body inversion effect in monkeys. *Plos One* 13.
Moors, P., Germeys, F., Pomianowska, I., Verfaillie, K., 2015. Perceiving where another person is
looking: the integration of head and body information in estimating another person's gaze.
Front Psychol 6**,** 909.
Peelen, M.V., Downing, P.E., 2007. The neural basis of visual body perception. *Nat.Rev.Neurosci.* 8**,**
636-648.
Perrett, D.I., Hietanen, J.K., Oram, M.W., Benson, P.J., 1992. Organization and functions of cells
responsive to faces in the temporal cortex. *Philos.Trans.R.Soc.Lond B Biol.Sci.* 335**,** 23-30.
Perrett, D.I., Smith, P.A., Potter, D.D., Mistlin, A.J., Head, A.S., Milner, A.D., Jeeves, M.A., 1984.
Neurones responsive to faces in the temporal cortex: studies of functional organization,
sensitivity to identity and relation to perception. *Hum.Neurobiol.* 3**,** 197-208.
Perrett, D.I., Smith, P.A., Potter, D.D., Mistlin, A.J., Head, A.S., Milner, A.D., Jeeves, M.A., 1985. Visual
cells in the temporal cortex sensitive to face view and gaze direction. *Proc.R.Soc.Lond B*
Biol.Sci. 223**,** 293-317.
Popivanov, I.D., Jastorff, J., Vanduffel, W., Vogels, R., 2015. Tolerance of macaque middle STS body
patch neurons to shape-preserving stimulus transformations. *J.Cogn Neurosci.* 27**,** 1001-
1016.
Rollenhagen, J.E., Olson, C.R., 2000. Mirror-image confusion in single neurons of the macaque
inferotemporal cortex. *Science* 287**,** 1506-1508.

Taubert, J., Ritchie, J.B., Ungerleider, L.G., Baker, C.I., 2022. One object, two networks? Assessing the
relationship between the face and body-selective regions in the primate visual system. *Brain*
Struct Funct 227**,** 1423-1438.
Tsao, D.Y., Livingstone, M.S., 2008. Mechanisms of face perception. *Annu.Rev.Neurosci.* 31**,** 411-437.
Vrancken, L., Germeys, F., Verfaillie, K., 2017. Holistic integration of gaze cues in visual face and body
perception: Evidence from the composite design. *J Vis* 17**,** 24.
Willenbockel, V., Sadr, J., Fiset, D., Horne, G.O., Gosselin, F., Tanaka, J.W., 2010. Controlling low-level
image properties: the SHINE toolbox. *Behav Res Methods* 42**,** 671-684.
Zafirova, Y., Bognar, A., Vogels, R., 2024. Configuration-sensitive face-body interactions in primate
visual cortex. *Prog Neurobiol* 232**,** 102545.
Zafirova, Y., Cui, D., Raman, R., Vogels, R., 2022. Keep the head in the right place: Face-body
interactions in inferior temporal cortex. *Neuroimage* 264**,** 119676.

Figure Legends

Figure 1. Recording locations and stimulus selectivity (A). Targeted patches (red circles) shown in

the native space of M1 (left) and M2 (right) on 2 mm coronal slices. Brown vertical lines illustrate

penetrations. Monkey patches were defined by the contrast monkeys > monkey control objects (M-

O_M; $p < 0.05$ FEW). **(B).** Normalized net responses in the Selectivity test (Experiment 1) in aSTS (left)

and mSTS (right). Rows represent neurons from M1 and M2 (separated by a dotted line). Columns

correspond to the stimuli, color-coded per category with a bar and one example image. Below, bars

in corresponding colors show the mean normalized net response to each stimulus (95% confidence

intervals (1000 resamplings)). *n* indicates the number of neurons.

Figure 2. Monkey avatar stimuli for the Head-body orientation test (Experiment 1). (A). 64

combinations of head and body orientations for the sitting (P1, left) and the standing pose (P2, right).

Body orientation varies by row and head orientation varies by column. Different colors group the 8

avatar orientations of four head-body angles. **(B).** Example stimuli of P1 and P2 in the monkey-

centered (MC, left) and the head-centered (HC, right) conditions, with zero angle between head and

body. **(C).** Headless bodies (B) and heads (H) for the zero and straight angle head-body

configurations, showing the frontal, back, and two lateral views for P1 and P2. Images are shown for

MC (left) and HC (right) conditions and were positioned to match their corresponding avatars in (A).

The location of the red fixation target is indicated in **(B)** and **(C)**.

Figure 3. Neuronal responses in the Head-body orientation test (Experiment 1). (A). Distribution of

MSI computed for the responses for all orientation, pose (P1 and P2), and centering (MC (blue) and

HC (red)) combinations in aSTS (upper panel) and mSTS (lower panel). Data for M1 and M2, and

medians (triangles) are shown in different hues. **(B).** Significant main effects and interactions of the

factors body and head orientation, for P1 and P2 in MC (blue) and HC (red) conditions in

corresponding colors) in aSTS (left) and mSTS (right). Pooled data of both subjects. Significant

interactions in both MC and HC are highlighted with darker blue and red (number of cases (n) in

black). **(C).** Distribution of the correlation coefficients between responses to the head-body

configurations in MC and HC conditions in aSTS (left) and mSTS (right). Data from both subjects and

poses were pooled. Correlation coefficients for all cases are shown above the x-axis. Below the x-axis,

the light green bars represent the correlation coefficients for cases with a significant interaction

(either MC or HC), while the dark green bars indicate the coefficients for cases with a significant

interaction in both MC and HC. N and medians (triangles) are shown in corresponding colors. **(D).**

Normalized net responses of example neurons to the 64 combinations of head and body orientations

in aSTS (left) and mSTS (right) for MC and HC conditions. Body orientation (ORT) varies by row and

head orientation varies by column, from 0° to 315°. **(E).** Net responses of example neurons to bodies,

heads, and the corresponding monkey configurations with MSI of 1, -1, or the median. Data for P1

and P2, MC and HC, and aSTS and mSTS (smoothed with the Matlab function *interp1*).

Figure 4. Decoding of head-body angles from neuronal responses in the Head-body orientation test

(Experiment 1). (A). Head-body configurations with zero and straight angles (black arrows in **(B)**) and

leftward and rightward right angles of head and body (grey arrows in **(B)**), of P1 and P2, at eight

avatar orientations. **(B).** Mean decoding accuracies for the head-body configurations with zero,

straight, leftward, and rightward right angles versus other head-body angles as indicated, in aSTS

(top two panels) and mSTS (bottom two panels), for poses P1 and P2. N indicates the number of

neurons. Results are for MC training and HC testing. Responses to the 8 orientations of a

configuration (rows in **(A)**) were pooled when training the classifier. Error bars show standard

deviations across resamplings. The shaded area indicates null distribution (stimulus labels

permutation). **(C).** Decoding accuracies for different head-body angle pairs in aSTS, for P1 (top) and

P2 (bottom). Decoded pairs are connected with lines, with decoding accuracies indicated by color.

Training was performed after pooling the 8 orientations of a head-body configuration defined by the

angle of the head and body. Results are from MC training and HC testing. Avatars of one orientation

of the head-body configuration angles are shown.

Figure 5. Decoding of head-body angles across orientations (Experiment 1). (A). Head-body

configurations with zero and straight angles, and leftward and rightward right angles for P1 and P2,

split into four orientations for training and testing. **(B).** Decoding accuracies for the head-body

configurations with zero, straight, leftward, and rightward right angles in aSTS, for P1 (top) and P2

(bottom); *n* indicates the number of neurons. Results are for MC training and HC testing, using four

orientations for training and testing. Panels one and three: classifier trained on '+' orientations and

tested on 'x' orientations; panels two and four: classifier trained on 'x' orientations and tested on '+'

orientations. Same conventions as Figure 4.

Figure 6. Decoding of head-body angles in the Sum vs. Configuration test (Experiment 2). (A).

Stimuli of Experiment 2: heads and bodies presented at four orientations (frontal, back, and the two

lateral ones; B+H), along with all head-body configurations (M). Images were monkey-centered and

the isolated heads and bodies were positioned to match their corresponding avatars. The red square

indicates the fixation target. **(B).** Decoding accuracies for head-body configurations (M) and head-

body sums (B+H) with zero, straight, leftward, and rightward right angles of head and body (trained

pooling the four orientations **(A)),** for aSTS and mSTS, *n* indicates the number of units. Same

conventions as Figure 4.

Figure 7. Effect of head-body configuration inversion (Experiment 3) (A). Stimuli of the head-body

configurations of P1 with zero and straight angles, and leftward and rightward right angles, at eight

orientations, upright and inverted. **(B).** Decoding accuracies for the head-body configurations with

zero, straight, leftward, and rightward right angles in aSTS, for poses P1 (top panels) and P2 (bottom

panels), *n* indicates the number of units. Solid and dotted lines represent data for upright and

inverted configurations, respectively. Same conventions as Figure 4. **(C).** Responses to the preferred

orientation of upright or inverted head-body configurations for aSTS and mSTS and P1 and P2 (M1

(dark green); M2 (light green)), for the monkey-responsive units. Medians (diamonds) and the

number of units per subject are shown in corresponding colors. P values of Wilcoxon signed-rank

tests (data pooled across subjects) testing the difference in response between upright and inverted

are shown (red: significant).