The RAB27A effector SYTL5 regulates mitophagy and mitochondrial metabolism

Ana Lapao[1,2*], Lauren Sophie Johnson[1,2,3*], Laura Trachsel-Moncho[1,2,3], Sakshi Singh[1,2], Matthew Yoke Wui Ng[1,2], Sigve Nakken[2,4,5], Eeva-Liisa Eskelinen[6], Anne Simonsen[1,2,3]

[1]Department of Molecular Medicine, Institute of Basic Medical Sciences, Faculty of Medicine, University of Oslo, 1112 Blindern, 0317 Oslo, Norway

[2]Centre for Cancer Cell Reprogramming, Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, 1171 Blindern, 0318 Oslo, Norway

[3]Department of Molecular Cell Biology, Institute for Cancer Research, Oslo University Hospital Montebello, 0379 Oslo, Norway

[4]Department of Tumour Biology, Institute for Cancer Research, Oslo University Hospital, Montebello, 0379 Oslo, Norway

[5]Department of Informatics, University of Oslo, Oslo, Norway

[6]Institute of Biomedicine, University of Turku, Turku, Finland

*The authors contributed equally

Correspondence: anne.simonsen@medisin.uio.no

Keywords

Mitochondrion, mitophagy, hypoxia, RAB27A, SYTL5, ACC

Abstract

SYTL5 is a member of the Synaptotagmin-Like (SYTL) protein family that differs from the Synaptotagmin family by having a unique N-terminal Synaptotagmin homology domain that directly interacts with the small GTPase RAB27A. Several SYTL protein family members have been implicated in plasma membrane transport and exocytosis, but the specific function of SYTL5 remains unknown. We here show that SYTL5 is a RAB27A effector and that both proteins localise to mitochondria and vesicles containing mitochondrial material. Mitochondrial recruitment of SYTL5 depends on its interaction with functional RAB27A. We demonstrate that SYTL5-RAB27A positive vesicles containing mitochondrial material and LAMP1 form during hypoxia and that SYTL5 and RAB27A promote mitophagy under hypoxia mimicking conditions, indicating a role for these proteins in mitophagy. Indeed, we find that SYTL5 interacts with proteins involved in vesicle-mediated transport and cellular response to stress and that its depletion compromises mitochondrial respiration and increases glucose uptake. Intriguingly, SYTL5 expression is significantly reduced in tumours of the adrenal gland, and correlates positively with survival for patients with adrenocortical carcinoma.

Introduction

The Synaptotagmin-Like (SYTL) protein family consists of 5 members (SYTL1-5), each containing an N-terminal Synaptotagmin homology domain (SHD) and two C-terminal C2 lipid-binding domains. The SHD domains of SYTL1-5 have been found to directly interact with the small GTPase protein RAB27[1,2]. The C2 domain is generally involved in phospholipid binding and interacts with cellular membranes[3], either in a calcium-dependent or calcium-independent manner[3,4]. Calcium was shown to be required for SYTL3 and SYTL5 phospholipid binding activity[1,5], whereas SYTL2 binding to phosphatidylserine (PS) was inhibited by calcium[6].

The SYTL protein family is generally involved in plasma membrane transport and exocytosis[2,4], mostly through their binding to RAB27 proteins. The main functions of RAB27 are related to vesicle budding, delivery, tethering and fusion with membranes[7] and its activity is regulated by a cyclic activation and inactivation state depending on it binding to guanosine-5'-triphosphate (GTP) or guanosine diphosphate (GDP), respectively[7]. RAB27A and RAB27B are the two RAB27 isoforms found in vertebrates[7] and their binding to effector proteins occurs

only when they are bound to GTP[7]. SYTL5 binds to the GTP-bound RAB27A form, indicating its possible role in membrane trafficking events[1].

Functional mitochondria are fundamental for normal cellular metabolism. Mitochondria are the primary source of adenosine triphosphate (ATP) obtained through oxidative phosphorylation (OXPHOS), and are important for cellular calcium homeostasis, lipid metabolism, reactive oxygen species (ROS) generation, and detoxification[8]. Mitochondria can integrate and generate signalling cues to adjust their metabolism and biogenesis to maintain cellular homeostasis[8,9]. Mitochondrial function and health are tightly regulated by several quality control processes involving targeting of parts of mitochondria to lysosomes for degradation, including macromitophagy[10,11] (selective degradation of mitochondria by autophagy), piecemeal mitophagy[12,13], mitochondrial-derived vesicles (MDVs)[14,15], and vesicles derived from the inner mitochondrial membrane (VDIMs)[16]. It has also been found that damaged mitochondria can be directly released to the extracellular space[17], either contained within extracellular vesicles[18] or by mitocytosis, where damaged mitochondria are expelled in migrasomes[19].

Exposure of mitochondria to various stressors can affect their function and lead to severe diseases, such as neurodegenerative disorders and cancer[20,21]. Cancer cells can trigger a change in mitochondrial metabolism to promote cell proliferation and survival by a process known as the Warburg effect, characterised by a switch from OXPHOS to ATP production through glycolysis even under normoxic conditions[22]. Adrenocortical carcinoma (ACC) is a rare type of cancer that develops in the adrenal gland cortex, having a very poor prognosis and limited treatment options[23]. Most patients are diagnosed at advanced cancer stages and present an excess in adrenocortical hormone production[23]. Mitochondria are known to be essential organelles in the synthesis of steroid hormones, including cortisol, since they contain specific enzymes that catalyse steroid synthesis from cholesterol delivered to the mitochondria[24].

Here we demonstrate that SYTL5 localises to mitochondria in a RAB27A-dependent manner and that both proteins regulate mitophagy and cellular metabolism. Cells lacking SYTL5 undergo a shift from mitochondrial oxygen consumption to glycolysis, which may explain the correlation between low SYTL5 expression and poor survival of ACC patients.

Results

SYTL5 localises to mitochondria and endolysosomal compartments

SYTL5 was identified as a putative candidate in a screen for lipid-binding proteins involved in the regulation of mitophagy[25]. To characterise the cellular localisation and function of SYTL5, we first generated U2OS cells with stable inducible expression of SYTL5-EGFP as there are no antibodies recognizing endogenous SYTL5. Intriguingly, live cell imaging analysis revealed that SYTL5-EGFP co-localised with filamentous structures positive for MitoTracker red **(Figure 1A)**, which labels active mitochondria. Moreover, several SYTL5-EGFP positive vesicles were observed, including some small and highly mobile SYTL5-EGFP vesicles moving along the mitochondrial network **(Supplementary video 1 and arrows in Figure 1B)**. SYTL5-EGFP staining at the plasma membrane was also seen, particularly in cells with higher expression levels **(Figure 1C)**, which may suggest a role for SYTL5 in secretion to the plasma membrane.

To determine the identity of the SYTL5 positive vesicles, U2OS SYTL5-EGFP cells were infected with a panel of lentiviral RAB GTPase constructs representing different cellular compartments **(Supplementary Figure 1A-F)** and analysed by live cell imaging. SYTL5-EGFP positive structures were observed to co-localise with vesicles positive for RAB4 and RAB11, two GTPases that mainly localise to recycling endosomes[26] and to some extent with RAB5, mostly localised to early endosomes[26] **(Supplementary Figure 1A-C)**. Co-localisation was also observed with RAB7, a marker of late endosomes, lysosomes and autophagosomes[26] and partially with RAB9, which associates with late endosomes and mediates the transport between late endosomes and the *trans*-Golgi network[26] **(Supplementary Figure 1D-E)**. SYTL5-EGFP showed little or no co-localisation with RAB6 positive structures, representing Golgi-derived vesicles **(Supplementary Figure 1F)**. In line with an endocytic identity of SYTL5-EGFP positive vesicles, they were also positive for lysosomal markers, including LysoTracker red, which labels acidic compartments in the cell **(Figure 1C)**, and the lysosomal marker LAMP1 **(Figure 1D)**.

Thus, based on live cell imaging analysis we conclude that SYTL5 localises to the mitochondrial network and to mitochondria-associated vesicles that partly overlap with endolysosomal compartments.

Mitochondrial localisation of SYTL5 requires both the RAB27-binding SHD domain and the lipid-binding C2 domains

To investigate whether the observed intracellular localisation of SYTL5-EGFP depends on its binding to RAB27 and/or lipids, we generated U2OS cells with stable expression of EGFP-3xFLAG-tagged wild type SYTL5 or SYTL5 lacking either the RAB27-binding SHD domain (SYTL5(ΔSHD)) or the two lipid-binding C2 domains (SYTL5(ΔC2AB)) **(Figure 1E)**.

To validate the SYTL5(ΔC2AB) mutant and determine the lipid-binding specificity of SYTL5, lysates from these cells were incubated with membranes containing various phosphoinositides and other lipids. Full-length SYTL5 was observed to bind to mono-phosphorylated phosphoinositides (PI(3)P, PI(4)P, PI(5)P), as well as PI(4,5)P2, PI(3,4,5)P3 and to some extent to phosphatidic acid (PA) and cardiolipin (CL) **(Figure 1F)**. As expected, upon removal of the C2 domains, specific binding to all lipid species was lost, while SYTL5 lacking the SHD domain retained the ability to bind to lipids **(Figure 1F)**.

The SYTL5(ΔSHD)-EGFP and SYTL5(ΔC2AB)-EGFP expressing cell lines were then incubated with MitoTracker red and analysed by live cell imaging. Indeed, the SYTL5 colocalisation to mitochondria was lost and both SYTL5(ΔSHD)-EGFP and SYTL5(ΔC2AB)-EGFP were dispersed in the cytosol **(Figure 1G-H),** indicating that both the RAB27-interacting SHD domain and the lipid binding C2 domains are required for mitochondrial localisation of SYTL5.

Mitochondrial localisation of SYTL5 requires RAB27A GTPase activity

Given that the SHD domain of SYTL5 has been shown to bind RAB27A[1] and our observations that the SHD domain is required for mitochondrial localisation of SYTL5, we asked whether RAB27A activity is required for SYTL5 recruitment to mitochondria. U2OS SYTL5-EGFP cells were infected with a mScarlet-RAB27A lentiviral construct to generate a stable cell line. Co-expression of SYTL5-EGFP and mScarlet-RAB27A resulted in a clear co-localisation of both proteins to filaments positive for MitoTracker DeepRed (DR) **(Figure 2A)**. The mitochondrial localisation of SYTL5-EGFP and mScarlet-RAB27A was confirmed by correlative light and EM (CLEM) analysis using the same cell line **(Figure 2B, zoom in 1)**. The mitochondrial localisation of SYTL5-EGFP was enhanced when co-expressed with mScarlet-RAB27A compared to cells expressing SYTL5-EGFP only **(Figure 1A-B)**, indicating that RAB27A might facilitate mitochondrial recruitment of SYTL5. Indeed, in U2OS cells with stable expression of mScarlet-RAB27A only, RAB27A strongly co-localised with MitoTracker green **(Supplementary Figure 2A)**, demonstrating its mitochondrial targeting. Moreover, SYTL5-EGFP and mScarlet-RAB27A were both detected in the mitochondrial fraction (containing

TIM23 and COXIV) of cells expressing SYTL5-EGFP or mScarlet-RAB27A **(Supplementary Figure 2B)**. Importantly, the mitochondrial localisation of SYTL5 and RAB27A is not specific to U2OS cells as HeLa cells with stable expression of SYTL5-EGFP or mScarlet-RAB27A show a similar expression pattern to U2OS cells (Figure 1A & C), as demonstrated by co-localisation of SYTL5-EGFP with MitoTracker Red and plasma membrane staining in cells having higher expression **(Supplementary Figure 2C)** and of mScarlet-RAB27A with MitoTracker Green **(Supplementary Figure 2D).** All further experiments were conducted in U2OS cells.

To further characterise mitochondrial recruitment of SYTL5 and RAB27A and elucidate their role at the mitochondrion, we used CRISPR/Cas9 to knock out (KO) RAB27A and SYTL5 in U2OS cells **(Supplementary Figure 3A-D)**. This double-KO (dKO) cell line was further transduced with lentiviral constructs to constitutively express SYTL5-EGFP, mScarlet-RAB27A, or both, and their mitochondrial localisation was analysed by live cell imaging **(Figure 2C-E)**. Intriguingly, mScarlet-RAB27A co-localised extensively with MitoTracker green and to small vesicles dispersed in the cytoplasm and located at or near mitochondria **(Figure 2C)**, indicating that SYTL5 is not required for RAB27A mitochondrial localisation. In contrast, SYTL5-EGFP was not recruited to mitochondria when expressed alone in dKO cells, although some SYTL5-EGFP vesicles were seen near mitochondrial structures **(Figure 2D)**. Upon co-expression of SYTL5-EGFP and mScarlet-RAB27A in dKO cells the mitochondrial localisation of SYTL5-EGFP was rescued **(Figure 2E),** indicating that mitochondrial recruitment of SYTL5 is dependent on RAB27A.

Since SYTL5 has previously been described as a RAB27A effector[1], we asked whether its mitochondrial recruitment depends on binding to the GTP-bound form of RAB27A. To address this, dKO cells expressing SYTL5-EGFP were rescued with the constitutively active (RAB27A-Q78L) or inactive (RAB27A-T23N) mutant forms of mScarlet-RAB27A. As expected, SYTL5-EGFP interacted specifically with RAB27A wild type (WT) and RAB27A-Q78L, as well as with endogenous RAB27A in control cells, while no interaction was detected with RAB27A-T23N, as assessed by GFP-pulldown **(Figure 2F)**. To our surprise, live cell imaging of the same cells revealed that, in contrast to mScarlet-RAB27A WT, neither the RAB27A Q78L nor the T23N mutant localised to the mitochondrial network when expressed together with SYTL5-EGFP in dKO cells. Also, SYTL5-EGFP failed to localise to mitochondria in cells expressing RAB27A Q78L, further demonstrating that mitochondrial recruitment of SYTL5 depends on mitochondrial RAB27A localisation **(Figure 2E).** Taken together, our data indicate that the

GTPase activity of RAB27A is required for its mitochondrial localisation, which facilitates further recruitment of SYTL5.

SYTL5 interacts with proteins involved in vesicle-mediated transport and cellular response to stress

As few molecular interactors of SYTL5 are known[1], we analysed the SYTL5 interactome by immunoprecipitating SYTL5-EGFP or an EGFP control, followed by mass spectrometry (MS) and protein-hit enrichment analysis. RAB27A was one of the most significant hits found, providing evidence that the IP was successful, together with several proteins known to participate in secretion such as PDCD6IP[28] **(Figure 3A)**. In total, 163 proteins were identified as significant SYTL5-EGFP interactors compared to the EGFP control **(Figure 3A, Table 1)** and were subjected to Gene Ontology (GO) term enrichment analysis. Considering GO enrichment within the subcellular compartment ontology (GO-CC), we discovered the following compartments as enriched (**Figure 3B**) (number of hits for each category in parentheses): cytoskeleton (43), intracellular vesicle (38), extracellular vesicle (36), plasma membrane-bound cell projection (33), endosome (18) and focal adhesion (17). Furthermore, for biological processes, encoded in the GO-BP subontology, we discovered the following categories as enriched **(Figure 3A and C):** vesicle-mediated transport (38), cellular response to stress (34), response to oxygen-containing compound (31), intracellular protein transport (22), secretion (21), autophagy (13), stress-activated MAPK cascade (10), cellular response to oxidative stress (9), reactive oxygen species metabolic process (7), regulation of mitochondrion organization (6), endosome organization (4) and protein insertion into mitochondrial membrane (3). Taken together, the SYTL5 interactome indicates a role for SYTL5 in mitochondrial processes and cellular response to stress, in line with its localisation to mitochondria, as well as a possible role in secretion and endocytosis, in line with its localisation to endocytic compartments and the plasma membrane **(Figure 1)**.

SYTL5-RAB27A positive vesicles contain mitochondrial components

Several proteins involved in autophagy were identified as specific SYTL5 interactors, including SQSTM1/p62, ATG4B, TBK1, and ATP6V1A **(Figure 3A)**, and given the mitochondrial localisation of SYTL5, we first investigated a possible role of SYTL5 in mitophagy. To induce mitophagy, dKO cells expressing mScarlet-RAB27A and SYTL5-EGFP were subjected to hypoxia

or the drugs DFP (deferiprone, an iron chelator) or DMOG (Dimethyloxalylglycine, a prolyl-4-hydroxylase inhibitor) that mimics hypoxia by activation of HIF1α. We noticed an increase in the number and size of vesicles positive for SYTL5-EGFP and mScarlet-RAB27A that also contained MitoTracker Deep Red (MitoTracker DR) in all conditions, compared to control cells **(Figure 4A).**

HIF-1α is a transcription factor that is a main driver of the Warburg effect[15] and an inducer of mitophagy via upregulation of the outer mitochondrial membrane proteins BNIP3 and BNIP3L, which bind to LC3. To identify the nature of the SYTL5/RAB27A/MitoTracker DR positive vesicular structures that increase with HIF-1α stabilisation, dKO cells expressing mScarlet-RAB27A and SYTL5-EGFP were immunostained with antibodies against various autophagy markers. Intriguingly, while observing strong co-localisation of SYTL5/RAB27A/MitoTracker DR positive vesicles with LAMP1 **(Figure 4B)**, the same structures did not co-localise with LC3 or the autophagy receptor p62 **(Supplementary Figure 4A).** Moreover, the mitochondrial localisation of SYTL5 and RAB27A was not affected in cells treated with the ULK1 inhibitor MRT68291 or with the class III phosphoinositide 3-kinase VPS34 inhibitor IN1 **(Supplementary Figure 4B)**, together indicating that SYTL5 and RAB27A are recruited to mitochondria independent of active autophagy.

SYTL5 and RAB27A regulate turnover of inner mitochondrial membrane proteins

To investigate whether SYTL5 and RAB27A are required for the turnover of mitochondrial proteins in hypoxic conditions, we took advantage of two mitophagy reporter U2OS cell lines, one expressing the inner mitochondrial membrane (IMM) reporter, pSu9-Halo-mGFP[29] and one expressing the mitochondrial targeting signal of the matrix protein NIPSNAP1 tagged with EGFP-mCherry[30] (hereafter referred to as IMLS cells).

The pSu9-Halo-mGFP cells allow a measure of mitophagy flux via the production of free Halo[TMR], which is stable in lysosomes when bound to the ligand. Activation of HIF1α by treatment with DFP or DMOG resulted in the appearance of a Halo Tag band due to cleaved Halo[TMR] **(Figure 4C-D)** consistent with an activation of mitophagy. The use of small interfering RNA (siRNA) to deplete SYTL5, RAB27A or both simultaneously in these cells, resulted in a significant reduction in the amount of free Halo[TMR] **(Figure 4C-D)** highlighting a reduction in mitophagy. This indicates that both SYTL5 and RAB27A are required for the turnover of mitochondrial inner membrane components in response to HIF1α activation.

Two independent siRNAs were used to deplete SYTL5 **(Supplementary Figure 5A)** in IMLS cells treated with DFP to induce Parkin-independent mitophagy and in IMLS cells with stable expression of Parkin treated with the uncoupler carbonyl cyanide m-chlorophenyl hydrazone (CCCP) to induce Parkin-dependent mitophagy. We performed imaging-based quantification of mCherry-only puncta (which represent mitochondria delivered to lysosomes as the EGFP signal is quenched in the acidic lysosome). Addition of the lysosomal V-ATPase inhibitor Bafilomycin A1 (BafA1) was used as a control to confirm that the mCherry-only puncta corresponded to lysosomal structures **(Supplementary Figure 5B-C)**. There was no significant difference in mCherry-only area per cell relative to the siControl in cells depleted of SYTL5, neither in cells treated with DFP **(Supplementary Figure 5B)** or with CCCP **(Supplementary Figure 5C)**. Moreover, IMLS turnover was not affected by depletion of RAB27A in these cells (**Supplementary Figure 5D**), together indicating that macro-mitophagy proceeds independently of SYTL5 and RAB27A. In line with this, the starvation-induced autophagic flux was not affected in cells lacking SYTL5 **(Supplementary Figure 5E)**.

Taken together, our results indicate that SYTL5 and RAB27A function to regulate the lysosomal turnover of mitochondrial inner membrane proteins in response to hypoxic conditions, in a macro-mitophagy independent manner.

SYTL5 regulates mitochondrial respiration

Our finding of mitochondrial material within SYTL5-RAB27A positive structures upon induction of mitochondrial stress through DFP and hypoxia **(Figure 4A)**, together with reduced turnover of the Fo-ATPase subunit 9 upon knockdown of SYTL5 and RAB27A **(Figure 4C-D)** and the identification of several proteins involved in the cellular response to stress **(Figure 3C)**, including mitochondrial proteins (e.g., DECR1 and TIMM13), in the SYTL5 interactome **(Figure 3A)** led us to investigate a potential role of SYTL5 and RAB27A in mitochondrial function and bioenergetics.

The cellular oxygen consumption rate (OCR) can be measured with the Seahorse analyser upon sequential addition of different inhibitors (oligomycin, CCCP, antimycin and rotenone) that block the function of specific electron transport chain (ETC) complexes[31]. SYTL5 KO cells had a significant decrease in both basal and ATP-linked respiration compared to WT U2OS cells **(Figure 5A-B),** indicating reduced mitochondrial OXPHOS activity. As for RAB27A, both basal respiration and ATP-linked OCR were significantly reduced in cells depleted of

RAB27A (RAB27A KO and RAB27A/SYTL5 dKO) relative to the control **(Figure 5A-B),** indicating an important function of these proteins in regulation of mitochondrial respiration.

To test whether the decreased respiration in cells depleted of SYTL5 and/or RAB27A is coupled to a shift to glycolysis, we performed Seahorse analysis of the extracellular acidification rate (ECAR) in cells upon the sequential addition of glucose (increases glycolytic capacity), oligomycin (inhibits mitochondrial ATP production), and 2-DG (a glycolysis inhibitor which causes an abrupt decrease in ECAR). Intriguingly, glycolysis was significantly increased in SYTL5 KO cells as compared to control **(Figure 5C-D)**, indicating a possible metabolic shift to compensate for the decrease in OCR observed when SYTL5 was depleted **(Figure 5A-B).** In contrast, ECAR was reduced in cells lacking RAB27A compared to the control, which may indicate an impairment in glucose uptake **(Figure 5C)** and therefore in glycolysis **(Figure 5D)**. After adding oligomycin the ECAR increased in all cell lines, but in RAB27A KO and dKO cell lines, the increase was reduced when compared to the control cell line **(Figure 5C-D).**

The clear reduction in mitochondrial respiration observed in cells lacking RAB27A and SYTL5 led us to investigate the abundance of various ETC proteins. Depletion of SYTL5 and/or RAB27A did not affect the abundance of ATP5A (complex V), UQCRC2 (complex III) or SDHB (complex II), but there was a significant increase in the level of MTCO1 (complex IV) in cells lacking RAB27A **(Figure 5E-F).** This is in line with the reduced turnover of Halo-tagged ATPase subunit Su9 observed in cells depleted of SYTL5 and/or RAB27A **(Figure 4C-D)**.

Taken together, our data show that both RAB27A and SYTL5 are important for normal mitochondrial respiration and ATP production while SYTL5 seems to regulate a metabolic switch to glycolysis.

Low SYTL5 expression is related to reduced survival for adrenocortical carcinoma patients
Since our data indicate that SYTL5 may be a regulator of the metabolic switch from oxidative phosphorylation to glycolysis (the Warburg effect) **(Figure 5A-D)** that is frequently associated with cancer cells, we decided to analyse publicly available expression datasets to explore a possible link between SYTL5 and cancer. Using bulk RNA expression data from healthy/normal tissue samples in the GTEx project, we found that SYTL5 is preferentially expressed in adrenal gland and various regions of the brain **(Figure 6A).** Adrenocortical carcinoma (ACC) is a rare type of aggressive cancer with poor prognosis and limited treatment options that develops in the adrenal cortex[23]. A comparison of gene expression levels[32] in normal adrenal gland

samples and ACC samples showed that SYTL5 mRNA expression is significantly reduced in tumour samples. **(Figure 6B).** Furthermore, patient survival analysis using data obtained from the TCGA-ACC cohort, indicates that low SYTL5 mRNA expression is associated with a significantly reduced survival of ACC patients **(Figure 6C)**, suggesting a possible tumour suppressor function for SYTL5 in ACC cancer progression. RAB27A expression levels were also reduced in tumor tissue compared to normal samples, although to a lesser extent than SYTL5 **(Figure 6D),** and overall patient survival in ACC patients did not show any relationship with RAB27A expression levels **(Figure 6E)**.

The adrenal gland cortex secretes several steroid hormones, such as mineralocorticoids (e.g. aldosterone), glucocorticoids (e.g. cortisol), and adrenal androgens (DHEA and DHEA-sulphate)[33], to the bloodstream. ACC tumours are characterised by excess adrenocortical hormone production in nearly 45–70% of patients, with hypercortisolism being the most common[23]. As mitochondria contain enzymes essential for steroid hormone biosynthesis from cholesterol[24], we asked whether SYTL5 may regulate cortisol production. SYTL5 or RAB27A were depleted by siRNA in the ACC cell line H295R, followed by quantification of cortisol secretion using a colorimetric assay. We did not observe a significant change in cortisol secretion upon SYTL5 or RAB27A knockdown (**Supplementary Figure 6**), indicating that the correlation between low SYTL5 expression levels and ACC patient survival likely is unrelated to effects on cortisol production.

Discussion

SYTL5 has been identified as a RAB27A effector protein with affinity for membranes through its C2 lipid-binding domains[1], but nothing is known about a potential role of SYTL5 in regulation of RAB27A-dependent membrane trafficking pathways. Using the osteosarcoma derived U2OS cell line, here we confirm that SYTL5 interacts with RAB27A, and show that it localises to mitochondria and small, highly mobile vesicles interacting with the mitochondrial network, as well as to endolysosomal and endocytic compartments, indicating a role of SYTL5 in cellular membrane trafficking events.

Most interestingly, mitochondrial recruitment of SYTL5 requires its RAB27-binding SHD domain and its lipid-binding C2 domains. Indeed, we demonstrate that SYTL5 is recruited to mitochondria in a RAB27A-dependent manner and that RAB27A itself is localised to mitochondria in a manner dependent on its GTPase activity, as both RAB27A GDP and GTP

mutants (T23N and Q78L, respectively) showed a dispersed cytoplasmic localisation. This is in line with a study showing that the GTPase activity of RAB27A is required for its localisation to melanosomes[34], but the mechanisms involved in mitochondrial recruitment of RAB27A are not clear. It is however interesting to note that ubiquitination of the RAB27A GTPase activating protein alpha (TBC1D10A) is reduced in the brain of Parkin KO mouse compared to controls[35], suggesting a possible connection of RAB27A with regulatory mechanisms that are linked with mitochondrial damage and dysfunction.

Given the presence of mitochondria-associated small and highly mobile SYTL5/RAB27A-positive vesicles that contain mitochondrial material and stain positive for LAMP1, we speculate that RAB27A and SYTL5 facilitate mitophagy-mediated lysosomal delivery of mitochondria or selected mitochondrial components. Indeed, SYTL5 and RAB27A positive vesicles containing mitochondrial material were observed in cells upon hypoxia or hypoxia-mimicking (DFP and DMOG) treatments. Intriguingly, in cells treated with DFP and DMOG we observed a significant decrease in lysosomal cleavage of the mitochondria inner membrane reporter pSu9-Halo-mGFP upon depletion of SYTL5 and/or RAB27A. Additionally, when blotting for various proteins of the electron transport chain complexes, we found that the level of the cytochrome c oxidase (complex IV) subunit MTCO1/COX1 was significantly increased in cells lacking RAB27A. In contrast, lysosomal targeting of a mitochondrial matrix reporter (IMLS) was not affected by depletion of SYTL5 and/or RAB27A, neither upon induction of Parkin-independent or Parkin-dependent mitophagy. In line with this, mitochondrial recruitment of SYTL5 and RAB27A was independent of the core autophagy machinery components ULK1 and VPS34 and the mitochondria-containing SYTL5/RAB27A-positive vesicles induced upon hypoxia did not stain positive for endogenous LC3B or p62, suggesting that SYTL5 and RAB27A-dependent clearance of mitochondrial components is independent of macroautophagy. The role of SYTL5 and RAB27A in the turnover of proteins involved in mitochondrial oxidative phosphorylation and ATP production is reminiscent of the recently described VDIM (Vesicles Derived from the Inner Mitochondrial membrane) pathway[16]. VDIMs are formed by IMM herniation through pores in the outer mitochondrial membrane, followed by their engulfment by lysosomes in proximity to mitochondria in a process that is independent of the core autophagy machinery. In line with our findings, VDIMs lack LC3 and p62 and VDIM formation was found to increase upon oxidative stress, leading to

selective degradation of IMM proteins (including COX4 and other proteins involved in OXPHOS) while sparing the remainder of the organelle.

Mitochondria are known stress sensors and hubs for cellular adaptation[8]. When analysing the interactome of SYTL5, we found that many candidates were related to the cellular response to stress and oxygen-containing compounds, as well as vesicle-mediated transport and secretion. In line with a role for SYTL5/RAB27A in the regulation of mitochondrial stress, we found that cells lacking SYTL5 and/or RAB27A displayed reduced OXPHOS activity and ATP-production. Interestingly, the depletion of SYTL5 triggered a shift to glycolysis as observed by an increase in the extracellular acidification rate (ECAR) due to cellular lactate production. A shift from OXPHOS to anaerobic glycolysis occurs in normal cells during hypoxic conditions and often in cancer cells even under normoxic conditions, commonly referred to as the Warburg effect[22].

The observed switch from OXPHOS to glycolysis in U2OS cells depleted of SYTL5 prompted us to look at cancer patient databases. We observed, using data from GTEx, that the adrenal gland is among the tissues with highest levels of SYTL5 gene expression, and that its expression is reduced in adrenocortical carcinoma samples (TCGA-ACC) when compared to normal adrenal gland tissue. Moreover, low SYTL5 expression is associated with reduced ACC patient survival. One characteristic of this type of cancer is an increase in steroid hormone production and secretion. Most steroid hormones are produced in mitochondria[24] but we did not observe any changes in the level of cortisol in ACC cells depleted of SYTL5. The mechanisms linking SYTL5 to the Warburg effect and ACC need further investigation, but it is interesting to note that RAB27A expression levels do not correlate with ACC survival.

To summarise, we show that SYTL5 is a RAB27A effector protein being recruited to mitochondria in a RAB27A-dependent manner. Upon hypoxia, RAB27A and SYTL5 promote lysosomal clearance of OXPHOS machinery components and both proteins are needed for mitochondrial respiration. We identify SYTL5 as a negative regulator of the Warburg effect and potentially tumorigenesis.

Materials and methods

Antibodies and dyes

Antibody	Source	Identifier
α-Tubulin	Sigma-Aldrich	#T5168
β-Actin	Cell signalling technology	#3700
COXIV	Cell signalling technology	#4850S
EGFP	Abcam; Clontech	#ab290; #632569
FLAG Tag	Cell signalling technology	#14793
GAPDH	Cell signalling technology	#5174
Halo Tag	Promega	#G9211
LAMP1	Santa Cruz Biotechnology	#sc-20011
OXPHOS	Abcam	#ab110413
p62	Progen	#GP62-C
RAB27A	Santa Cruz Biotechnology	#sc-74586
SYTL5	Sigma-Aldrich	#HPA026074
TIM23	BD biosciences	#611223
TOMM20	Santa Cruz Biotechnology	#sc-17764
LysoTracker red DND-99	ThermoFisher Scientific	#L7528
MitoTracker™ Deep Red FM	ThermoFisher Scientific	# M22426
MitoTracker™ Green FM	ThermoFisher Scientific	#M7514
MitoTracker™ Red CMXRos	ThermoFisher Scientific	#M7512

Cell culture, lentivirus production and stable cell line generation

Human bone osteosarcoma epithelial cells (U2OS FlpIn TRex cell line, kindly provided by Steve Blacklow, Harvard Medical School, US) were grown in complete medium of Dulbecco´s Modified Eagle Medium (DMEM; Lonza #12-604F and Gibco # 31966047) with 10% v/v foetal bovine serum (FBS; Sigma-Aldrich #F7524),100 U/ml Penicillin and 100 µg/ml Streptomycin (P/S; Thermo Fischer Scientific #15140122) at 37 °C with 5% CO_2.

For lentiviral particle production HeK-FT cells were co-transfected with 1.6 µg of pLVX lentiviral vector, pCMV-VSV-G and psPAX2 using X-tremeGENE™ 9 DNA Transfection Reagent

(Sigma #XTG9-RO). For virus infection, cells were seeded in complete medium and supplemented with 8 μg/ml polybrene (Santa Cruz Biotechnology #sc-134220). 2 μg/ml Puromycin (Sigma Aldrich #P7255) was added after 24 h post-infection for selection.

To generate U2OS stably expressing SYTL5-EGFP, U2OS FlpIn TRex cells and U2OS FlpIn TRex_mScarlet-RAB27A were co-transfected with 0.1 μg pcDNA™5/FRT/TO_SYTL5-EGFP and 0.9 μg pOG44 Flp recombinase expression vector using Lipofectamine™ 2000 (Thermo Fisher Scientific #11668019) according to the manufacturer´s instructions. To produce stable cell lines expressing pLVX-SV40-mScarlet-RAB27A(T23N/Q78L) and pLVX-CMV-SYTL5-EGFP-3xHA, U2OS dKO cells were infected with lentiviral particles followed by fluorescence-activated cell sorting (FACS) for double mScarlet/EGFP expressing cells. To generate stable cells expressing pLVX-CMV-SYTL5-EGFP-3xFLAG, pLVX-CMV-SYTL5 (ΔC2AB)-EGFP-3xFLAG and pLVX-CMV-SYTL5 (ΔSHD)-EGFP-3xFLAG, U2OS FlpIn TRex cells were infected with lentiviral particles. To generate U2OS stably expressing pSu9-Halo-mGFP, U2OS cells were infected with retroviral particles generated using pUMVC (addgene # 8449) and pMRX-IB-pSu9-HaloTag7-mGFP (addgene # 184905). Stable cell lines were maintained in complete medium with additional 5 μg/ml blasticidin S (Thermo Fisher Scientific #R21001) or 2 μg/ml puromycin (Sigma Aldrich #P7255). For starvation experiments, cells were washed with PBS and incubated for 4 h in Earle´s balanced salt solution (EBSS; Gibco #24010043).

NCI-H295R cells were obtained from ATCC (#CRL-2128) and maintained in DMEM: F12, HEPES medium (Gibco #11330032) supplemented with 2.5% Nu-Serum (Corning #355100), 1% ITS+Premix Universal culture supplement (Corning #354352) and 100 μg/ml P/S (Thermo Fischer Scientific #15140122) at 37 °C with 5% CO_2.

Hypoxia and drug treatments

After cells were seeded, they were transferred to a INVIVO$_2$ 200 Hypoxic workstation (Ruskinn). Hypoxia treatment was carried out for 24 hours with oxygen concentration set to 1%.

Drugs were added into the cell media. Treatment durations and drug concentrations are indicated in the figure legends. Drugs used were DFP (Sigma-Aldrich #379409), DMOG (Sigma-Aldrich #D3695), VPS34IN1 (Selleckchem # S7980) and MRT68291 (Selleckchem #S7949).

Molecular cloning

SYTL5 was amplified from WT U2OS cDNA (5´-ATGTCTAAGAACTCAGAGTTCATC-3´ and 5´-TTAGAGCCTACATTTTCCCATG-3´) and cloned into Zero Blunt TOPO vector using Zero Blunt™ TOPO™ PCR Cloning Kit (Thermo Fisher Scientific #450245) according to the manufacturer´s instructions. pcDNA™5/FRT/TO_SYTL5-EGFP was generated by Gibson Assembly using the TOPO-SYTL5 vector as a template for SYTL5 and cloned into a pcDNA™5/FRT Mammalian Expression Vector (Thermo Fisher Scientific #V601020) or into a pLVX-CMV lentiviral expression vector. Truncated SYTL5 (ΔSHD) and SYTL5 (ΔC2AB) were generated by Gibson assembly using the pcDNA™5/FRT/TO_SYTL5-EGFP construct as a template and cloned into a pLVX-CMV lentiviral expression vector.

RAB27A was amplified from WT U2OS cDNA and cloned into a pLVX-SV40 lentiviral expression vector. RAB27A mutations were performed using the site-directed mutagenesis QuikChange II kit (Agilent #200524) according to the manufacturer´s instructions using the following primer pairs:

Mutation	Forward primer	Reverse primer
RAB27A_T23N (c68a)	atattggtaaagtacactgttcttccctacaccagagtc	gactctggtgtagggaagaacagtgtactttaccaatat
RAB27A_Q78L (a233t)	tacgaaacctctccagccctgctgtgtcc	ggacacagcagggctggagaggtttcgta

CRISPR knockout cell line generation and validation

SYTL5 and RAB27A gene knockouts were performed as previously described by Ran *et al*., 2013**.** Briefly, gRNA target sequences were designed using Wellcome Sanger Institute Genome Editing (WGE) design tool available at https://wge.stemcell.sanger.ac.uk/find_crisprs.

Target gene	gRNA sequence	Genomic location
RAB27A	AGTGGCTCCATCCGGCCCAC	chr15:55230454
SYTL5 guide 1	CGACTAATACTTGCCAGAGC	chrX:38034292
SYTL5 guide 2	ATACGTAGAACAGCCTCCAC	chrX:38033693

DNA oligos were annealed and phosphorylated before cloning into pSpCas9(BB)-2A-Puro (PX459) V2.0 (addgene # 62988). The obtained CRISPR plasmids were transfected into U2OS Flp-In TRex cells using Lipofectamine™ 2000 according to the manufacturer´s instructions and

selection with 2 μg/ml puromycin (Sigma Aldrich #P7255) was applied for 24 h. Limiting dilutions were performed using a concentration of 15 cells/ml and seeded in complete medium in a 384 well plate. Single colonies were collected ~10 days after seeding and expanded for a further 3 weeks. For genotyping, gDNA from each single clone line was isolated using QIAamp DNA Mini Kit (QIAGEN #51304) and the targeted genomic region was amplified by PCR using flanking primers (RAB27A: 5´-ATAACCTCTCCCTTGACCTTGTATG-3´ and 5´-TAGATGCCTTTGGGATTTGTACTGA-3´; SYTL5: 5´-CAGGTCCCTTTCTTCTCGCA-3´ and 5´-TCAGTCAGCTGCAAGAGTGG-3´). The PCR product was cloned into Zero Blunt TOPO vector using Zero Blunt™ TOPO™ PCR Cloning Kit according to the manufacturer´s instructions. RAB27A clonal lines were validated by western blot, and SYTL5 deletions were detected by running PCR products in 1% w/v agarose gel (Thermo Fisher Scientific #R2801).

siRNA knockdown by reverse transfection

For siRNA knockdown, cells were incubated with OptiMEM (Thermo Fisher Scientific # 31985047) and Lipofectamine RNAiMAX (Thermo Fisher Scientific #13778150) for 5 min at room temperature (RT). Followed by addition of 20 nM siRNA diluted in OptiMEM and 15 min incubation at RT. Cells were fixed/harvested after 72 h of knockdown. The siRNA used for mRNA depletion are the following: siControl: 5´-UAACGACGCGACGACGUAAtt-3´; siSYTL5 #1: 5´-CUCUUAGAAGCAAAACGUAtt-3´; siSYTL5 #2: 5´-CAACAAGCGUAAGACCAAAtt-3´; siSYTL5 #3: 5'-GGUUUGUGCUUCAACCCAAtt-3'; siRAB27A #1: 5'-GGAAGACCAGUGUACUUUAtt-3´ and siRAB27A #2: 5´-CCAGUGUACUUUACCAAUAtt-3´.

RNA isolation, cDNA synthesis and RT-PCR

RNA was isolated using TRIzol reagent (Thermo Fisher Scientific #15596026) and cDNA was synthesised using SuperScript III Reverse Transcriptase (Thermo Fisher Scientific #18080085) according to the manufacturer´s instructions. qPCR analysis was performed using KAPA SYBR FAST qPCR Kit (Sigma-Aldrich #KK4601) and relative mRNA expression levels were normalised to the expression levels of TATA-binding protein (TBP) using the comparative ΔCt method. The primers used were the following:

SYTL5: 5´-GTGACAAAATCGCGCAGCTA-3´, 5´-GGACAACATCAGTGCCGAGA-3´

RAB27A: 5'-GGAGAGGTTTCGTAGCTTAACG-3', 5'-CCACACAGCACTATATCTGGGT-3'

TBP: 5´-CAGAAAGTTCATCCTCTGGGCT-3´, 5´-TATATTCGGCGTTTCGGGCA-3´

Immunofluorescent staining

Cells were fixed in 4% paraformaldehyde (PFA) (Sigma-Aldrich #158127) in PBS for at least 15min at 37°C, quenched for 10 min using 0.05 M NH4Cl in PBS, permeabilised in 0.05% saponin in PBS for 5 min and blocked in 1% BSA in PBS for 30 min at room temperature. Antibody staining was performed in a wet chamber for 1 h at room temperature and the antibodies were diluted in PBS containing 0.05% saponin. Nuclei staining was performed using Hoechst 33342 (Thermo Fisher Scientific #H1399) diluted in PBS at 1µg/ml.

Light microscopy

Cells expressing inducible fluorescent proteins were induced with 100 ng/ml doxycycline (Clontech #631311) for 24 h. For live cell confocal microscopy cells were incubated with indicated dyes prior to imaging. Imaging was carried out with either the Andor Dragonfly High Speed Confocal Microscope or the Nikon Crest V3 with a 60x oil immersion objective (NA 1.4). High content widefield imaging was conducted with the ImageXpress Micro Confocal (Molecular Devices) microscope with a 20x objective (NA 0.45). Fixed immunofluorescence confocal imaging was performed using the Andor Dragonfly High Speed Confocal Microscope or Nikon Crest V3 both with a 60x oil immersion objective (NA 1.4) or Zeiss LSM 800 microscope (Zen Black 524 2012 SP5 FP3, Zeiss) with a 63x oil immersion objective (NA 1.4).

Correlative light electron microscopy

U2OS cells co-expressing SYTL5-EGFP and mScarlet-RAB27A were seeded in gridded glass-bottom cell culture dishes (MatTek #P35G-1.5-14-CGRD). Cells growing in monolayer were fixed in warm (≈37°C) 3.7% PFA in 0.2 M HEPES (pH 7) and imaged using the Andor Dragonfly High Speed Confocal Microscope. After imaging, the samples were fixed using 2% v/v glutaraldehyde in 0.2 M HEPES, pH 7.4. After post-fixation in 2% v/v osmium tetraoxide and 3% v/v K4[Fe(CN)6], samples were embedded in Epon and 60 nm thickness sections were cut with a diamond knife. Sections were analysed using a JEM-1400 Plus Transmission Electron Microscope.

Western blot

Cells were harvested in RIPA buffer (50 mM TRIS pH 7.4, 150 mM NaCl (Merck #1064041000), 0,5% sodium deoxycholate (Sigma-Aldrich #30970), 1% NP-40 (Sigma-Aldrich #I3091), 1 mM EDTA (VWR #20302-293), 0,1% SDS (Sigma-Aldrich #L3771) containing 1x protease inhibitor (Merck #11697498001) and 1x phosphatase inhibitor (Merck #4906845001). 20 µg of lysate was loaded on SDS-PAGE gel (BioRad #5671095) and proteins transferred to a PVDF membrane (Merck #IPFL00010) which was blocked using 5x v/v casein blocking buffer in PBS for 1 h at room temperature. The obtained membranes were immunoblotted using primary and secondary antibodies with washing steps using PBST (1x Phosphate-Buffered Saline with 1% Tween20 (Sigma-Aldrich #P1379)). Membranes were analysed using the Odyssey CLx imaging system (Li-cor biosciences) and protein levels were quantified by densitometry with ImageStudio Lite software (Li-cor biosciences).

Macro-mitophagy analysis

siRNA knockdown was conducted as previously described in U2OS-IMLS cells seeded in complete media supplemented with 100 ng/ml doxycycline. Culture media was replaced 24 h after transfection and 1 mM DFP (Sigma-Aldrich #379409) treatment was added after 48 h. For U2OS-IMLS cells expressing PARKIN, 20 µM CCCP (Enzo Life Sciences #BML-CM124-0500) was used for 16 h and 10 µM Q-VD-OPh (Sigma-Aldrich #SML0063-1MG) pan-caspase inhibitor was included to support cell survival[11,36]. Control wells with 100 nM BafA1 (AH diagnostics #BML-CM110-0100) were dosed 2 h prior to fixation. After 72 h, cells were washed with PBS pH 7 and fixed in warm (≈37°C) 3.7% PFA in 0.2 M HEPES (pH 7) for 15 min at 37°C. After washing with PBS, cells were incubated with 2 µg/ml Hoechst and imaged.

Halo assay for mitophagy flux

siRNA knockdown was conducted as previously described in U2OS cells expressing the pSu9-Halo-mGFP reporter. 48 h after transfection culture media was changed to contain 100 nM tetramethylrhodamine (TMR)-conjugated Halo ligand (Promega #G8251). After 20 min incubation, cells were washed twice with PBS and the media replaced with either media containing no treatment, 1 mM DFP or 1 mM DMOG. 72 h after transfection cells were harvested and western blot conducted (as described earlier). Membranes were incubated with the Halo Tag antibody and actin antibody. Membranes were imaged using the Chemidoc MP system (Bio-Rad) and band intensity quantified in Image Lab 6.1 (Bio-Rad). Bands were

normalised to actin loading control. The percentage cleaved Halo was calculated by the formula (Free Halo / (Free Halo + Full Length)) x 100.

Oxygen consumption rate measurement by Seahorse XF analyser

U2OS cells were seeded in complete medium at the density of 4×10^4 cells/well into XF24 cell culture microplates (Agilent #100777-004). The sensor cartridge was hydrated for 24 h using XF calibrant in a 37°C non-CO_2 incubator. Before analysis, cell medium was replaced by warm (≈37°C) DMEM without Sodium Bicarbonate (pH 7.4) and placed for 1 h in a 37°C non-CO_2 incubator. Each measurement cycle consisted of a mixing time of 3 min and data acquisition period of 3 min (three data points). OCR results refer to the average rates during each measurement cycle. The final concentration for each mitochondrial inhibitor used in this assay was 1.5 µM Oligomycin A (SelleckChem #S1478), 1 µM CCCP, 0.5 µM Rotenone (Sigma Aldrich #R8875-1G) and 0.5 µM Antimycin (Sigma Aldrich #A8674). Four baseline measurements were performed before adding each compound and three response measurements were taken after each compound was added. Data analysis was performed using Seahorse Analytics software available at https://seahorseanalytics.agilent.com/ and OCR data was normalised to protein concentration in each well.

Extracellular acidification rate (ECAR) measurement by Seahorse XF analyser

U2OS cells were seeded in complete medium at the density of 4×10^4 cells/well into XF24 cell culture microplates. The sensor cartridge was hydrated for 24 h using XF calibrant in a 37°C non-CO_2 incubator. Before analysis, cell medium was replaced by warm (≈37°C) XF DMEM media (Seahorse Bioscience #102353-100) supplemented with 2 mM L-Glutamine (BioNordika #BE17-605E) and pH adjusted to 7.4. The final concentration for each reagent used in this assay was 11 mM Glucose**,** 1.3 µM Oligomycin A, 0.1 M 2-Deoxy-Glucose (2-DG; Sigma Aldrich #D8375-1G)[37]. Each measurement cycle consisted of a mixing time of 3 min and data acquisition period of 3 min (four data points). ECAR results refers to the average rates during each measurement cycle. Five baseline measurements were performed before adding each compound and four response measurements were taken after each compound was added. Data was normalised to protein concentration using Seahorse Analytics software available at https://seahorseanalytics.agilent.com/ and the glycolysis rate was calculated by subtracting the ECAR value after 2-DG treatment from the ECAR value after addition of glucose.

Purification of crude mitochondrial fraction

To purify crude mitochondrial fraction, cells expressing EGFP (control), mScarlet-RAB27A, SYTL5-EGFP and SYTL5(ΔC2AB)-EGFP were harvested from one 150 mm (80-90% confluency) dish, washed twice with ice cold homogenisation buffer (210 mM mannitol (Sigma Aldrich #M4125), 70 mM sucrose (Sigma Aldrich #S0389), 5 mM HEPES pH 7.12 at 25°C) and harvested in 1 mL homogenisation buffer supplemented (HBS) with 1 mM EGTA (Sigma Aldrich #E3889) and 1x protease inhibitor. The obtained cell lysate was further mechanically homogenised at 4°C using a cell homogeniser (Isobiotech) equipped with a 16 μm clearance ball by repeatedly passing the cell lysate for 10x repeats with the aid of 2 disposable syringes. The homogenised solution was further spun down at 1500xg for 3 min at 4°C to collect the TCL and the obtained supernatant collected and spun down at 13,000xg for 20 min. The obtained supernatant corresponds to the cytosolic fraction and the pellet was resuspended in 2 mL HBS and spun down at 13,000xg for 20 min. The obtained pellet corresponds to the enriched mitochondrial fraction and was resuspended in 900 μL HBS. 50 μL of each cellular fraction was resuspended in 30 μL 2x SDS-sample buffer and boiled at 100°C for 5 min and further analysed using SDS-PAGE.

GFP-TRAP and FLAG immunoprecipitation

Cells expressing EGFP were harvested from a 100 mm dish (80-90% confluency), washed with ice cold PBS and scraped using ice cold isolation buffer (1% NP-40, 2 mM EDTA, 136 mM NaCl, 20 mM Tris-HCl pH 7.4, 10% glycerol, 1x protease inhibitor and 1x phosphatase inhibitor). Cell lysate was spun down at 20,000xg for 10 min and the supernatant added to 25 μL equilibrated GFP-trap agarose beads (Chromotek #gta-20). Samples were rotated end-over-end for 1 h at 4°C. After 4x washing steps, the beads were collected by centrifugation at 2700xg for 1 min, resuspended in 30 μL 2x SDS-sample buffer and boiled at 100 °C for 5 min. The obtained supernatant was analysed by SDS-PAGE.

U2OS cells expressing pLVX-CMV-SYTL5-EGFP-3xFLAG, pLVX-CMV-SYTL5(ΔC2AB)-EGFP-3xFLAG and pLVX-CMV-SYTL5(ΔSHD)-EGFP-3xFLAG were harvested from a 150 mm dish (80-90% confluency), washed 2x with cold PBS and scraped using 1 ml of cold lysis buffer (50 mM Tris HCl, pH 7.4,150 mM NaCl, 1% Triton X-100, 5 mM $CaCl_2$ (Merck #1023821000, and 1x protease inhibitor. Cell lysates were spun down at 20,000xg for 10 min and the resulting

supernatants added to 40 μL of pre-washed anti-FLAG M2-Agarose Affinity Gel (Sigma Aldrich #FLAGIPT-1). Samples were rotated end-over-end for 2 h at 4°C. After 3x washing steps, the affinity gel was centrifuged at 5000xg for 30 seconds and the FLAG fusion proteins eluted using 3x FLAG peptide according to manufacturer's protocol. The eluted FLAG fusion proteins were diluted in 4 ml 1x TBST supplemented with 3% BSA and 5 mM $CaCl_2$.

I*n-vitro* protein-lipid binding assay

Eluted FLAG fusion proteins (SYTL5-EGFP-3xFLAG, SYTL5(ΔC2AB)-EGFP-3xFLAG and SYTL5(ΔSHD)-EGFP-3xFLAG were diluted in 1x TBST supplemented with 3% BSA and 5 mM $CaCl_2$. PIP strips (Echelon Biosciences #P-6001) and membrane lipid strips (Echelon Biosciences #P-6002) were blocked in 1x TBST supplemented with 3% BSA for 1 h at room temperature and incubated with the diluted FLAG fusion protein for 1 h at 4°C with gentle agitation. Lipid membranes were washed 3x with TBST and incubated with FLAG Tag primary antibody for 1 h at room temperature and detected using SuperSignal West Dura Extended Duration Substrate (Thermo Fisher Scientific #34076).

LC-MS/MS analysis of SYTL5-EGFP interactome

Cells expressing SYTL5-EGFP or EGFP (control) were harvested using NP-40 lysis buffer (10 mMTris Cl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 0.5% NP-40, 1x protease inhibitor. SYTL5-EGFP was immunoprecipitated using GFP-Trap agarose beads according to the manufacturer's protocol. The obtained beads were washed twice with 50 mM ammonium bicarbonate, reduced, alkylated and further digested by trypsin for overnight at 37 °C. Digested peptides were transferred to a new tube, acidified and the peptides were de-salted for MS analysis.

Peptide samples were dissolved in 10 μl 0.1% formic buffer and 3 μl was loaded for MS analysis. LC-MS/MS analysis of the resulting peptides was performed using an Easy nLC1000 liquid chromatography system (Thermo Electron) coupled to a QExactive HF Hybrid Quadrupole-Orbitrap mass spectrometer (Thermo Electron) with a nanoelectrospray ion source (EasySpray, Thermo Electron). The LC separation of peptides was performed using an EasySpray C18 analytical column (2 μm particle size, 100 Å, 75 μm inner diameter and 25 cm (Thermo Fisher Scientific). Peptides were separated over a 90min gradient from 2% to 30% (v/v) ACN in 0.1% (v/v) FA, after which the column was washed using 90% (v/v) ACN in 0.1% (v/v) FA for 20 min (flow rate 0.3 μL/min). All LC-MS/MS analyses were operated in data-

dependent mode where the most intense peptides were automatically selected for fragmentation by high-energy collision-induced dissociation.

Raw files from LC-MS/MS analyses were submitted to MaxQuant 1.6.17.0 software[38] for peptide/protein identification. Parameters were set as follow: Carbamidomethyl (C) was set as a fixed modification and PTY; protein N-acetylation and methionine oxidation as variable modifications. First search error window of 20 ppm and mains search error of 6 ppm. Trypsin without proline restriction enzyme option was used, with two allowed miscleavages. Minimal unique peptides were set to one, and FDR allowed was 0.01 (1%) for peptide and protein identification. The Uniprot human database was used. Generation of reversed sequences was selected to assign FDR rates. Further analysis was performed with Perseus[39], limma/voom[40] and Package R[41]. Volcano plots were plotted with EnhancedVolcano[42]. GO analysis was performed with shinyGO[43].

Bulk gene expression analysis – normal and tumor tissue

Bulk gene expression distribution in healthy/normal tissue samples was collected from GTEx v8 (PMID: 26484571). We utilised a combined RNA-seq expression dataset of TCGA and GTEx samples available through the Xena platform to investigate SYTL5 and RAB27A expression in normal and tumor tissue (PMID: 32444850). Here, data from a total of n = 77 samples provided normal adrenal gland expression (GTEx), and a total of n = 128 samples provided expression in adrenocortical carcinoma samples (TCGA-ACC). Survival data from samples in the TCGA-ACC cohort (one record per patient) was obtained using the TCGABiolinks R package[44]. Survival curve analysis was performed using GraphPad Prism 8.0.1 using Logrank test (Mantel-Cox).

Cortisol assay

To measure cortisol in cell culture supernates, siRNA knockdown by reverse transfection with siSYTL5 and siRAB27A was conducted as described earlier in NCI-H295R cells. Cells were harvested with TNTE Lysis Buffer. Cortisol from lysates was measured using the R&D Systems® Cortisol Immunoassay kit (# KGE008B) as described in the protocol provided from the company. Controls were performed using cells treated with 50 uM Forskolin as a positive control to stimulate cortisol secretion[45], or 10 uM mitotane to inhibit cortisol production[46].

Statistical analysis

Statistical analysis was performed with GraphPad Prism (8.0.1, 9.3.1 and 10.2.0 versions) and the tests used are indicated in the respective figure legends. All values come from distinct samples.

**** = p<0.0001, *** = p<0.001, ** = p<0.01, * = p< 0.05.

Acknowledgments

We thank Jenni Lane for assisting with EM sample preparation, Sachin Singh and Tuula Nyman at the OUS Proteomics Core Facility for assisting with mass spectrometry-based proteomic experiments, Ankush Sharma for help with analysis of mass spectrometry data, Anna Lång at ALM Core Facility Gaustad for assisting with widefield microscopy experiments, Santosh Phuyal for experimental assistance and Serhiy Pankiv for providing the lentiviral mScarlet-RAB expression constructs. We also thank Patricia González-Rodríguez for providing us with the U2OS pSu9-HaloTag7-mGFP cells.

This project was funded by the Research Council of Norway through its Centres of Excellence funding scheme (project number 262652) and FRIPRO grant (project number 249753), the Norwegian Cancer Society (project number 190251), Marie Skłodowska-Curie ETN grant under the European Union's Horizon 2020 Research and Innovation Programme (Grant Agreement No 765912 DRIVE) and by the University of Oslo Scientia Fellow program through the MSC scheme – Co-funding of Regional, National and International Programmes (COFUND). Mass spectrometry-based proteomic analyses were performed by the Proteomics Core Facility, Department of Immunology, University of Oslo/Oslo University Hospital, which is supported by the Core Facilities program of the South-Eastern Norway Regional Health Authority. This core facility is also a member of the National Network of Advanced Proteomics Infrastructure (NAPI), which is funded by the Research Council of Norway INFRASTRUKTUR-program (project number: 295910).

Author contributions

A.L., L.S.J., L.T.M. and A.S. designed and conceived the experimental work. A.L., L.S.J., L.T.M. and M.Y.W.N. performed the experiments and data analyses. S.S. carried out MS and GO term enrichment data analysis. S.N. provided GTEx and TCGA data analysis. E.L.E. performed and analysed CLEM data. A.L., L.S.J. and A.S. wrote the manuscript.

Author declarations

The authors have nothing to declare.

Figures

Figure 1. Lipid binding protein SYTL5 localises to mitochondria and endo-lysosomes

A. Live confocal microscopy imaging of U2OS stably expressing SYTL5-EGFP co-stained with 50 nM MitoTracker red. SYTL5-EGFP expression was induced for 24 h using 100 ng/ml doxycycline. MitoTracker red was added 30 min before imaging. Scale bars: 10 μm, 2 μm (insets).

B. Time-lapse video frames (Supplementary video 1) tracking SYTL5 vesicle movement along filaments positive for MitoTracker red (arrows). SYTL5-EGFP expression was induced for 24 h using 100 ng/ml doxycycline. Scale bars: 3 μm.

C. Live confocal microscopy imaging of U2OS stably expressing SYTL5-EGFP co-stained with 50 nM LysoTracker red. SYTL5-EGFP expression was induced for 24 h using 100 ng/ml doxycycline. LysoTracker red was added 30 min before imaging. Scale bars: 10 μm, 2 μm (insets).

D. Confocal imaging of U2OS stably expressing SYTL5-EGFP stained for endogenous LAMP1. Nuclei were stained using Hoechst. Scale bars: 10 μm, 2 μm (insets).

E. Overview of domain structure of SYTL5 full length, ΔSHD and ΔC2AB mutants.

F. 3xFLAG fusion proteins expressed in U2OS cells (SYTL5-EGFP-3xFLAG, SYTL5 (ΔSHD)-EGFP-3xFLAG and SYTL5 (ΔC2AB)-EGFP-3xFLAG) were immunoprecipitated and added to membranes spotted with lipids found in cell membranes: LPA (lipoprotein A); LPC (lysophosphatidylcholine); PI (phosphatidylinositol); PI(3)P (phosphatidylinositol 3-phosphate); PI(4)P (phosphatidylinositol 4-phosphate); PI(5)P (Phosphatidylinositol 5-phosphate); PE (phosphatidylethanolamine); PC (phosphatidylcholine); S1P (sphingosine-1-phosphate); PI(3,4)P2 (phosphatidylinositol 3,4-bisphosphate); PI(3,5)P2 (phosphatidylinositol 3,5-bisphosphate); PI(4,5)P2 (phosphatidylinositol 4,5-bisphosphate); PI(3,4,5)P3 (phosphatidylinositol 3,4,5 –triphosphate); PA (phosphatidic acid); PS (Phosphatidylserine); TAG (triglyceride); DAG (diglyceride); PG (phosphatidylglycerol); CL (cardiolipin); Cholesterol; SM (sphingomyelin); Sulfatide.

G. Live confocal microscopy imaging of U2OS stably expressing SYTL5 (ΔSHD)-EGFP-3xFLAG co-stained with 50 nM MitoTracker red added 30 min before imaging. Scale bars: 10 μm, 2 μm (insets).

H. Live confocal microscopy imaging of U2OS stably expressing SYTL5 (ΔC2AB)-EGFP-3xFLAG co-stained with 50 nM MitoTracker red added 30 min before imaging. Scale bars: 10 μm, 2 μm (insets).

Figure 2. Mitochondrial localisation of SYTL5 requires RAB27A GTPase activity

A. SYTL5-EGFP expression was induced for 24 h with 100 ng/ml doxycycline in U2OS cells with stable inducible expression of SYTL5-EGFP and constitutive expression of mScarlet-RAB27A. Cells were co-stained with MitoTracker DR for 30 min before live imaging. Arrows indicate mitochondrion filaments. Scale bars: 10 μm, 2 μm (insets).

B. CLEM analysis of the cells described in A. Before imaging, cells growing in monolayer were fixed in warm (≈37°C) 3.7% paraformaldehyde in 0.2 M HEPES (pH 7). After fixation, cells were imaged using a confocal microscope to acquire Z-stack of optical sections and DIC images to locate the cells of interest. The cells were finally fixed using 2% glutaraldehyde in 0.2 M HEPES (pH 7.4) for 120 min before sample preparation for TEM. Arrows in panel 1 indicate mitochondrion filaments. Scale bars: 15 μm (left), 5 μm (middle) and 2 μm (right).

C. U2OS dKO cells were rescued with mScarlet-RAB27A and co-stained with 50 nM MitoTracker green for 30 min before imaging. Scale bars: 10 μm, 2 μm (insets).

D. U2OS dKO cells were rescued with SYTL5-EGFP and co-stained with 50 nM MitoTracker red for 30 min before imaging. Scale bars: 10 μm, 2 μm (insets).

E. Live confocal microscopy imaging of U2OS dKO cells rescued with SYTL5-EGFP and mScarlet-RAB27A (upper panel); SYTL5-EGFP and mScarlet-RAB27A-T23N (middle panel) or SYTL5-EGFP and mScarlet-RAB27A-Q78L (lower panel). All cells were co-stained with 50 nM MitoTracker DR for 30 min before imaging. Scale bars: 10 μm, 2 μm (insets).

F. Lysates from U2OS cells stably expressing EGFP (control) and U2OS dKO cells expressing SYTL5-EGFP and/or mScarlet-RAB27A (wild-type, T23N or Q78L mutants) were immunoprecipitated using GFP-Trap beads and analysed by western blot using RAB27A and EGFP antibodies.

Figure 3. SYTL5 interacts with proteins involved in vesicle-mediated transport and cellular response to stress

A. SYTL5 interactome. SYTL5-EGFP expression in U2OS cells was induced for 24 h using 100 ng/ml doxycycline, followed by immunoprecipitation of SYTL5-EGFP using GFP-Trap beads and

identification of co-purified proteins by MS analysis. The list of co-purified proteins was compared with that obtained from cells expressing EGFP and only significant (p<0.05) SYTL5-EGFP specific protein hits are highlighted in colour. Protein hits falling into at least one of the biological processes; autophagy, regulation of mitochondrion organisation, protein insertion into mitochondrial membrane or cellular response to oxidative stress, are colour coded according to plot legend. All other significant hits are indicated in blue. Non-significant (p>0.05) are indicated in grey. Data are from 3 biological replicates.

B. GO cellular compartment term enrichment of significant protein hits co-purified with SYTL5-EGFP. Corresponding enrichment false discovery rate (FDR) value is represented inside each bar and bars are ordered from smallest to largest FDR (q-value) from top to bottom.

C. GO biological processes term enrichment of significant protein hits co-purified with SYTL5-EGFP.Corresponding enrichment FDR value is represented inside each bar and bars are ordered from smallest to largest FDR (q-value) from top to bottom. Proteins corresponding to each biological process category are listed in the table to the right.

Figure 4. Mitochondrial localisation of SYTL5 and RAB27A is not perturbed by autophagy inhibition or mitophagy stimulation

A. SYTL5/RAB27A dKO U2OS cells rescued with SYTL5-EGFP and mScarlet-RAB27A were untreated (upper panel) or treated with 1 mM DFP (2nd panel), hypoxia (1% O_2) (3rd panel) or 1 mM DMOG (4th panel) for 24 h. All cells were stained with 50 nM MitoTracker DR 30 min prior to live confocal microscopy imaging. Scale bars: 10 µm, 2 µm (insets).

B. SYTL5/RAB27A dKO U2OS cells rescued with SYTL5-EGFP and mScarlet-RAB27A were treated or not with 1 mM DFP for 24 h. Cells were stained with 50 nM MitoTracker DR prior to fixation. Fixed cells were stained with a LAMP1 antibody and imaged by confocal microscopy. Scale bars: 10 µm, 2 µm (insets). Arrowheads point to SYTL5/RAB27A-positive structures.

C. U2OS cells expressing pSu9-Halo-mGFP were transfected with 20 nM control siRNA, SYTL5 siRNA #1 or RAB27A siRNA #1 for 48 h before 20 min incubation with the TMR-conjugated Halo ligand followed by 24 h incubation with 1 mM DFP or 1 mM DMOG. Cell lysates were analysed by western blot for the Halo tag. Actin was used as a loading control.

D. Quantification of data in C. Error bars represent the mean with standard deviation between replicates (n=3). Significance was determined by two-way ANOVA followed by Tukey´s multiple comparison test. * = p> 0.05, ** =p> 0,01, *** = p>0.001.

Figure 5. SYTL5 and RAB27A regulate mitochondrial bioenergetics.

A. Mitochondrial oxygen consumption rate was analysed in U2OS control cells, SYTL5 KO, RAB27A KO and dKO cells using the Seahorse XF analyser. OCR was measured after sequential addition of oligomycin, CCCP and a combination of rotenone and antimycin-A into the assay medium. Error bars represent the mean with standard deviation between replicates (n=3).

B. The four basal OCR measurements per well (before oligomycin addition) were averaged to obtain the basal OCR value and the non-mitochondrial respiration was subtracted to determine the basal respiration associated to each condition. The ATP production was calculated by subtracting the proton leak from the maximal respiratory capacity. Error bars represent the mean with standard deviation between replicates (n=3). Significance was determined by two-way ANOVA followed by Bonferroni multiple comparison test, ** = p>0.01, * = p> 0.05.

C. The extracellular acidification rate (ECAR) was analysed in U2OS control cells, SYTL5 KO, RAB27A KO and dKO cells using the Seahorse XF analyser. ECAR was measured upon addition of glucose, oligomycin and 2-DG using control and SYTL5 depleted cells. Error bars represent the mean with standard deviation between replicates (n=3).

D. The glycolysis rate of each condition was calculated by subtracting the ECAR value after 2-DG treatment to the ECAR after addition of glucose (Glycolysis=ECAR after addition of glucose − ECAR after 2-DG treatment). Error bars represent the mean with standard deviation between replicates (n=3). Significance was determined by one-way ANOVA followed by Tukey multiple comparison test, * = p> 0.05.

E. Cell lysates from U2OS control, RAB27A KO, SYTL5 KO and dKO cells were analysed by western blot for the indicated electron transport chain complex-II, -III, -V and -V proteins. GAPDH was used as a loading control.

F. Quantification of data in E. Band intensities were quantified relative to GAPDH and normalised to control. Error bars represent the mean with standard deviation between replicates (n=4). Significance was determined by two-way ANOVA followed by Bonferroni multiple comparison test, ** = p>0.01.

Figure 6. Low SYTL5 expression is related to reduced survival for adrenocortical carcinoma patients.

A. SYTL5 expression levels in normal/healthy tissues, using data from the GTEx project (v8). Shown are the 10 tissue sites for which SYTL5 is most highly expressed (i.e. based on median expression levels). TPM = transcripts per million. Median is represented by a red solid line and quartiles by red dashed lines.

B. Comparison of SYTL5 gene expression levels in normal adrenal gland samples (GTEx, n = 128) and adrenocortical carcinoma samples (The Cancer Genome Atlas (TCGA) ACC cohort, n = 77). Median expression is represented by a red solid line and quartiles by red dashed lines. Statistical analysis was performed using the unpaired t-test with Welch´s correction. TPM= transcripts per million.

C. Kaplan-Meier plot for ACC patient survival related to SYTL5 expression levels. The events related to high SYTL5 expression (expression level above median) are indicated in red and the dotted lines represent 95% confidence interval. Events related to low SYTL5 expression (expression level below median) are indicated in blue and the dotted lines represent 95% confidence interval. Statistical analysis was performed using Logrank (Mantel-Cox) test.

D. Comparison of RAB27A gene expression levels in normal adrenal gland samples (GTEx, n = 128) and adrenocortical carcinoma samples (The Cancer Genome Atlas (TCGA) ACC cohort, n = 77). Median expression is represented by a red solid line and quartiles by red dashed lines. Statistical analysis was performed using the unpaired t-test with Welch´s correction.

E. Kaplan-Meier plot for ACC patient survival related to RAB27A expression levels. The events related to high RAB27A expression (expression level above median) are indicated in red and the dotted lines represent 95% confidence interval. Events related to low RAB27A expression (expression level below median) are indicated in blue and the dotted lines represent 95% confidence interval. Statistical analysis was performed using Logrank (Mantel-Cox) test.

Supplementary Figures

Figure S1

A-F. SYTL5-EGFP expression was induced for 24 h with 100 ng/ml doxycycline in U2OS cells with stable inducible expression of SYTL5-EGFP and constitutive expression of mScarlet-RAB4

(A), mScarlet-RAB11 (B), mScarlet-RAB5 (C), mScarlet-RAB7 (D), mScarlet-RAB9 (E) or mScarlet-RAB6 (F) followed by live confocal microscopy imaging analysis. Scale bars: 10 µm, 2 µm (insets).

Figure S2

A. Live confocal microscopy imaging of U2OS constitutively expressing mScarlet-RAB27A co-stained with MitoTracker green added 30 min before imaging. Scale bars: 10 µm, 2 µm (insets).

B. Total cell lysate (TCL), cytosol, and a crude mitochondria fraction from U2OS control cells and cells with stable expression of mScarlet-RAB27A or SYTL5-EGFP were analysed by western blotting for EGFP, RAB27A, tubulin, TIM23 and COXIV.

C. Live confocal microscopy imaging of HeLa constitutively expressing SYTL5-EGFP co-stained with MitoTracker Red added 30 min before imaging. Scale bars: 10 µm, 2 µm (insets).

D. Live confocal microscopy imaging of HeLa constitutively expressing mScarlet-RAB27A co-stained with MitoTracker green added 30 min before imaging. Scale bars: 10 µm, 2 µm (insets).

Figure S3

A. RAB27A KO validation by genotyping. The WT RAB27A gDNA sequence was aligned with the RAB27A KO clone 21(c21) gDNA sequence. CRISPR/Cas9 editing produced an indel (single nucleotide deletion indicated by *) in a genomic location shared between all RAB27A isoforms. gRNA targeting region is underlined in the WT RAB27A gDNA sequence.

B. RAB27A KO validation by western blot. Cell lysates from U2OS control cells and RAB27A KO cells (c21) were compared to cell lysates where RAB27A was depleted for 72 h using 2 independent siRNAs and probed with RAB27 and actin antibodies.

C. SYTL5 CRISPR/Cas9 exon deletion was performed using two gRNA simultaneously. These gRNA target the 5´UTR or intronic regions of SYTL5 (transcript NM_138780) that are shared by all SYTL5 isoforms. Genomic deletion validation and individual clone genotyping was performed using a primer pair that flanked the region to be edited (about 1307 bp length between these primer binding sites).

D. SYTL5 KO validation was performed with conventional PCR reaction. A PCR product of 1307 bp is expected for wild-type U2OS gDNA, while a PCR product of about 618 bp is expected if a clone has a deletion in the targeted region. This method was used to validate the SYTL5 KO

clone2 and for dKO RAB27A/SYTL5 clone57. The deleted sequence includes a coding exon from SYTL5, resulting in a non-functional SYTL5 protein.

Figure S4

A. SYTL5/RAB27A dKO U2OS cells rescued with SYTL5-EGFP and mScarlet-RAB27A were treated with 1 mM DFP for 24 h. Fixed cells were stained with a LAMP1 antibody in addition to either p62 (upper panel) or LC3 (lower panels) antibodies and imaged by confocal microscopy. Arrowheads point to SYTL5/RAB27A-positive structures. Scale bars: 10 μm, 2 μm (insets).

B. SYTL5/RAB27A dKO U2OS cells rescued with SYTL5-EGFP and mScarlet-RAB27A were treated with 1 μM MRT68921 for 2 h (upper panel) or 5 μM Vps34-IN1 for 2 h (lower panel). All cells were stained with 50 nM MitoTracker DR 30 min prior to live imaging

confocal microscopy. Scale bars: 10 μm, 2 μm (insets).

Figure S5

A. U2OS cells were transfected with 20 nM control siRNA or SYTL5 siRNA oligos for 72 h. Transfection efficiency was analysed by qPCR where the relative mRNA expression of SYTL5 was normalised to the expression levels of TBP and siControl. Error bars represent the standard deviation between replicates (n=3). Significance was determined by one-way ANOVA followed by Dunnett multiple comparison test, *** = p>0.001.

B. U2OS cells expressing the mitochondrial matrix reporter NIPSNAP1[1-53]-EGFP-mCherry[30] (referred to as IMLS cells) were transfected with 20 nM control siRNA or SYTL5 siRNA oligos for 48 h before 24 h incubation with 1 mM DFP in the absence or presence of 100 nM BafA1 for the last 2 h. After fixation, cell nuclei were stained with Hoechst and widefield images were obtained using a high-content imaging microscope. The area of red-only puncta per cell (>1000 cells analysed) was quantified using a Cell Profiler pipeline. The obtained results were normalised to the control siRNA and error bars represent the mean with standard deviation between replicates (n=3). Significance was determined by two-way ANOVA followed by Bonferroni multiple comparison test.

C. U2OS IMLS cells expressing PARKIN were transfected with 20 nM control siRNA or SYTL5 siRNA oligos for 48 h before a 16 h incubation with 20 μM CCCP to induce mitophagy, with treatment with 100 nM BafA1 or not for the last 2 h. The area of red-only puncta per cell

(>1000 cells analysed) was quantified using Cell Profiler from widefield images obtained using a high-content imaging microscope. The obtained results were normalised to the control siRNA and error bars represent the mean with standard deviation between replicates (n=3). Significance was determined by two-way ANOVA followed by Tukey´s multiple comparison test.

D. U2OS IMLS cells were transfected with 20 nM control siRNA or RAB27A siRNA oligos for 48 h before 24 h incubation with 1 mM DFP in the absence or presence of 100 nM BafA1 for the last 2 h. After fixation, cell nuclei were stained with Hoechst and widefield images were obtained using a high-content imaging microscope. The area of red-only puncta per cell (>1000 cells analysed) was quantified using a Cell Profiler pipeline. The obtained results were normalised to the siControl, and error bars represent the mean with standard deviation between replicates (n=3). Significance was determined by two-way ANOVA followed by Bonferroni multiple comparison test.

E. WT (control) or SYTL5 KO U2OS cells were starved in EBSS for 4 h or incubated with complete medium in the presence or absence of 100 nM BafA1 the last 2 h. Cell lysates were harvested and analysed by western blot for LC3B and actin proteins. LC3B-II band intensity was quantified relative to actin and normalised to control. Error bars represent the mean with standard deviation between replicates (n=3). Significance was determined by one-way ANOVA followed by Bonferroni multiple comparison test.

Figure S6

NCl-H295R cells were transfected with 20 nM control siRNA, SYTL5 siRNA or RAB27A siRNA oligos for 72 h. Cortisol secretion was measured and normalised to siControl. Error bars represent the mean with standard deviation between replicates (n=4).

Supplementary video 1. Live confocal microscopy imaging tracking SYTL5 (green) vesicle movement along filaments positive for MitoTracker Red (magenta).

Table 1. SYTL5 interactome

Proteins co-immunoprecipitated with SYTL5-EGFP vs EGFP expressing control

Gene ID	logFC	P.Value	Protein ID
SYTL5	8,190967	6,18E-06	Q8TDW5

RAB27A	4,904658	0,00013	P51159; O00194
DICER1	4,072889	3,36E-05	Q9UPY3
NRBP1	3,616261	0,000139	Q9UHY1
TUBGCP3	3,480067	2,23E-06	Q96CW5
UBXN4	3,364929	0,000445	Q92575
PPME1	2,860594	0,003682	Q9Y570
SUGT1	2,644558	3,96E-05	Q9Y2Z0
RRM2	2,58741	0,000772	P31350
S100A16	2,489595	0,019199	Q96FQ6
RBX1	2,283705	5,54E-05	P62877
CAPNS1	2,245698	0,0045	P04632
IPO5	2,152529	0,001141	O00410
SQSTM1	2,087749	0,019914	Q13501
CDK5RAP1	2,083531	0,022579	Q96SZ6
PSMD9	2,066147	0,000637	O00233
ANXA1	2,015019	0,008846	P04083
RIPK2	1,998221	0,012785	O43353
TUBB2A	1,937802	0,032975	Q13885
FBXL18	1,932709	0,000593	Q96ME1
ACACA	1,884369	0,00271	Q13085; O00763
TRAF2	1,879215	0,041615	Q12933
ATXN10	1,807462	0,001441	Q9UBB4
PFKFB3	1,806392	0,011617	Q16875
CASC5	1,787295	0,038936	Q8NG31
ACTR2	1,753581	0,00102	P61160
PDCD6IP	1,750369	0,001188	Q8WUM4
DNAJB2	1,746104	0,020493	P25686
MTAP	1,723933	0,037343	Q13126
NOSIP	1,722576	0,041459	Q9Y314
LARP4	1,673434	0,010333	Q71RC2
ENY2	1,672969	0,029441	Q9NPA8
ARMCX3	1,661646	0,020894	Q9UH62
TIMM13	1,652287	0,016672	Q9Y5L4
ZC3HAV1L	1,646732	0,000559	Q96H79
CLU	1,641816	0,018032	P10909
STAT3	1,630662	0,017936	P40763
MMP2	1,596861	0,01583	P08253
ARHGAP28	1,594874	0,029771	Q9P2N2
CASP2	1,54764	0,006055	P42575
NECAP1	1,532475	0,001098	Q8NC96; Q9NVZ3
CHORDC1	1,530842	0,006654	Q9UHD1
AHNAK2	1,518123	0,003188	Q8IVF2
P4HA2	1,507659	0,01657	O15460
SLAIN2	1,504025	0,038526	Q9P270
TTLL12	1,496256	0,016402	Q14166
CIAO1	1,47458	0,002001	O76071

PSMD13	1,469132	0,004425	Q9UNM6
PAPSS2	1,468267	0,024743	O95340
PKP3	1,462143	0,000241	Q9Y446
MAPKAPK3	1,43686	0,036615	Q16644
LAMC2	1,424905	0,014389	Q13753
TMED10	1,420889	0,033461	P49755
GGA2	1,401072	0,036446	Q9UJY4
SMYD5	1,380067	0,000962	Q6GMV2
MYD88	1,379573	0,0036	Q99836
PTPN23	1,372845	0,042911	Q9H3S7
PRMT3	1,367264	0,035439	O60678
SRR	1,335156	0,018756	Q9GZT4
METTL2B	1,318423	0,002052	Q6P1Q9
CEP55	1,303196	0,042401	Q53EZ4
KNTC1	1,293798	0,013151	P50748
MMADHC	1,287966	0,007304	Q9H3L0
RPS6KA4	1,278372	0,026028	O75676
TCEAL4	1,277767	0,041367	Q96EI5; Q9H3H9
KRT18	1,273972	0,000277	P05783
EIF3D	1,249965	0,001692	O15371
MAP3K6	1,235349	0,038163	O95382
FAM83D	1,235204	0,001544	Q9H4H8
FLCN	1,223376	0,025555	Q8NFG4
NARFL	1,207308	0,005387	Q9H6Q4; Q9UHQ1
TK1	1,191836	0,022913	P04183
TSN	1,182404	0,012781	Q15631
MOGS	1,181984	0,013095	Q13724
TRIP13	1,179669	0,007178	Q15645
SH3GL1	1,148087	0,001254	Q99961; Q99962
TGFB1	1,146935	0,009311	P01137
TBK1	1,146181	0,005314	Q9UHD2
UBAP2L	1,131475	0,034553	Q14157
HMMR	1,124677	0,038667	O75330
MAP2K4	1,123533	0,004105	P45985
LRRC40	1,116807	0,03566	Q9H9A6
TAB2	1,114102	0,012624	Q9NYJ8
IPO7	1,106622	0,002335	O95373
DECR1	1,096331	0,014198	Q16698
PRKAR1A	1,090646	0,049639	P10644
ATP6V1A	1,082598	0,006145	P38606
ATG4B	1,062937	0,01514	Q9Y4P1
CIAPIN1	1,056349	0,026294	Q6FI81
STIM1	1,036432	0,036684	Q13586
TSNAX	1,033898	0,012445	Q99598
PDLIM2	1,014087	0,008808	Q96JY6
KIF11	1,012826	0,007811	P52732

NCAPD2	1,005509	0,004454	Q15021
FLYWCH2	1,002772	0,036116	Q96CP2
FLNC	0,997402	0,031493	Q14315
ZCCHC11	0,995037	0,011036	Q5TAX3
ARF5	0,98277	0,002189	P84085; P61204; P18085; P84077
ARHGAP29	0,982689	0,020558	Q52LW3
INPP4A	0,977505	0,046529	Q96PE3
IKBKG	0,964638	0,017171	Q9Y6K9
CACYBP	0,96422	0,007507	Q9HB71
SEC23B	0,951773	0,047922	Q15437
CARHSP1	0,950642	0,016008	Q9Y2V2
STRN	0,93495	0,004936	O43815
KIAA0930	0,920553	0,044996	Q6ICG6
DPYSL3	0,910877	0,006555	Q14195
ACAA1	0,902484	0,014043	P09110
PPIA	0,901801	0,005015	P62937
JAK1	0,898744	0,029572	P23458
PXN	0,89135	0,032021	P49023
RIPK1	0,880408	0,02096	Q13546
HACL1	0,873144	0,005027	Q9UJ83
C16orf13	0,872785	0,043785	Q96S19
STAU2	0,871501	0,007129	Q9NUL3
CTTN	0,868965	0,002283	Q14247
NEDD1	0,865712	0,00298	Q8NHV4
RELA	0,848508	0,008652	Q04206
MOCS2	0,844002	0,02455	O96007
SESTD1	0,843323	0,022617	Q86VW0
CNN2	0,84227	0,005121	Q99439
TNS3	0,824331	0,019403	Q68CZ2; Q63HR2
MYH9	0,820463	0,020254	P35579; P35749
FAM120B	0,81995	0,012614	Q96EK7
PHLDB1	0,816857	0,002876	Q86UU1
LAMB3	0,811098	0,007705	Q13751
ERCC6L	0,809206	0,022191	Q2NKX8
RASA4	0,79333	0,044026	O43374; C9J798
SH3KBP1	0,78816	0,014872	Q96B97
ACTB	0,783173	0,00407	P60709; Q6S8J3; A5A3E0; P0CG38; Q562R1; P0CG39; Q9BYX7
CNN3	0,781781	0,003728	Q15417
DNMBP	0,770568	0,030522	Q6XZF7
INF2	0,769802	0,005056	Q27J81
NFKB2	0,762598	0,02469	Q00653
DOK1	0,758782	0,022414	Q99704
MVP	0,758137	0,025696	Q14764
SEPT9	0,750806	0,015745	Q9UHD8
SFN	0,747282	0,038485	P31947

UFSP2	0,736462	0,032911	Q9NUQ7
GNA13	0,729323	0,01046	Q14344
VPS37B	0,721343	0,006336	Q9H9H4
MCMBP	0,716545	0,008801	Q9BTE3
DPYSL2	0,707331	0,02086	Q16555
RNPEP	0,705101	0,032267	Q9H4A4
UBA6	0,691729	0,015468	A0AVT1
GSTK1	0,687332	0,046851	Q9Y2Q3
GMPS	0,683068	0,014906	P49915
POLR2B	0,676992	0,027717	P30876
DBNL	0,675429	0,006534	Q9UJU6
PLCG1	0,669824	0,049137	P19174
HSPA4	0,665847	0,027702	P34932
LIMD1	0,661786	0,018739	Q9UGP4
CAT	0,653596	0,036557	P04040
ANXA2	0,652983	0,016136	P07355; A6NMY6
CBR1	0,64346	0,032038	P16152; O75828
PRDX2	0,643131	0,022495	P32119
UBA1	0,634137	0,022102	P22314
NCAPG	0,632896	0,021943	Q9BPX3
MAT2A	0,622619	0,013543	P31153; Q00266
GRB2	0,619474	0,025687	P62993
KIF1B	0,612282	0,046537	O60333; Q12756; O43896
KIF5B	0,611182	0,018915	P33176; O60282; Q12840
MAT2B	0,602416	0,016771	Q9NZL9

References

Kuroda, T. S., Fukuda, M., Ariga, H. & Mikoshiba, K. Synaptotagmin-like protein 5: a novel Rab27A effector with C-terminal tandem C2 domains. *Biochemical and biophysical research communications* **293**, 899-906 (2002). https://doi.org/10.1016/s0006-291x(02)00320-0

Kuroda, T. S., Fukuda, M., Ariga, H. & Mikoshiba, K. The Slp homology domain of synaptotagmin-like proteins 1-4 and Slac2 functions as a novel Rab27A binding domain. *The Journal of biological chemistry* **277**, 9212-9218 (2002). https://doi.org/10.1074/jbc.M112414200

Nalefski, E. A. & Falke, J. J. The C2 domain calcium-binding motif: structural and functional diversity. *Protein Sci* **5**, 2375-2390 (1996). https://doi.org/10.1002/pro.5560051201

Gálvez-Santisteban, M. *et al.* Synaptotagmin-like proteins control the formation of a single apical membrane domain in epithelial cells. *Nature cell biology* **14**, 838-849 (2012). https://doi.org/10.1038/ncb2541

Fukuda, M. The C2A domain of synaptotagmin-like protein 3 (Slp3) is an atypical calcium-dependent phospholipid-binding machine: comparison with the C2A domain of synaptotagmin I. *The Biochemical journal* **366**, 681-687 (2002). https://doi.org/10.1042/bj20020484

Yu, M. *et al.* Exophilin4/Slp2-a targets glucagon granules to the plasma membrane through unique Ca2+-inhibitory phospholipid-binding activity of the

C2A domain. *Molecular biology of the cell* **18**, 688-696 (2007). https://doi.org/10.1091/mbc.e06-10-0914

Fukuda, M. Rab27 effectors, pleiotropic regulators in secretory pathways. *Traffic* **14**, 949-963 (2013). https://doi.org/10.1111/tra.12083

Montava-Garriga, L. & Ganley, I. G. Outstanding Questions in Mitophagy: What We Do and Do Not Know. *J Mol Biol* **432**, 206-230 (2020). https://doi.org/10.1016/j.jmb.2019.06.032

Martínez-Reyes, I. & Chandel, N. S. Mitochondrial TCA cycle metabolites control physiology and disease. *Nature communications* **11**, 102 (2020). https://doi.org/10.1038/s41467-019-13668-3

Allen, G. F., Toth, R., James, J. & Ganley, I. G. Loss of iron triggers PINK1/Parkin-independent mitophagy. *EMBO reports* **14**, 1127-1135 (2013). https://doi.org/10.1038/embor.2013.168

Lazarou, M. *et al.* The ubiquitin kinase PINK1 recruits autophagy receptors to induce mitophagy. *Nature* **524**, 309-314 (2015). https://doi.org/10.1038/nature14893

Le Guerroué, F. *et al.* Autophagosomal Content Profiling Reveals an LC3C-Dependent Piecemeal Mitophagy Pathway. *Molecular cell* **68**, 786-796.e786 (2017). https://doi.org/10.1016/j.molcel.2017.10.029

Abudu, Y. P. *et al.* SAMM50 acts with p62 in piecemeal basal- and OXPHOS-induced mitophagy of SAM and MICOS components. *The Journal of cell biology* **220** (2021). https://doi.org/10.1083/jcb.202009092

Neuspiel, M. *et al.* Cargo-selected transport from the mitochondria to peroxisomes is mediated by vesicular carriers. *Curr Biol* **18**, 102-108 (2008). https://doi.org/10.1016/j.cub.2007.12.038

Soubannier, V. *et al.* A vesicular transport pathway shuttles cargo from mitochondria to lysosomes. *Curr Biol* **22**, 135-141 (2012). https://doi.org/10.1016/j.cub.2011.11.057

Prashar, A. *et al.* Lysosomes drive the piecemeal removal of mitochondrial inner membrane. *Nature* **632**, 1110-1117 (2024). https://doi.org/10.1038/s41586-024-07835-w

Choong, C. J. *et al.* Alternative mitochondrial quality control mediated by extracellular release. *Autophagy* **17**, 2962-2974 (2021). https://doi.org/10.1080/15548627.2020.1848130

Phinney, D. G. *et al.* Mesenchymal stem cells use extracellular vesicles to outsource mitophagy and shuttle microRNAs. *Nature communications* **6**, 8472 (2015). https://doi.org/10.1038/ncomms9472

Jiao, H. *et al.* Mitocytosis, a migrasome-mediated mitochondrial quality-control process. *Cell* **184**, 2896-2910.e2813 (2021). https://doi.org/10.1016/j.cell.2021.04.027

Youle, R. J. & Narendra, D. P. Mechanisms of mitophagy. *Nature reviews. Molecular cell biology* **12**, 9-14 (2011). https://doi.org/10.1038/nrm3028

Ding, W. X. & Yin, X. M. Mitophagy: mechanisms, pathophysiological roles, and analysis. *Biol Chem* **393**, 547-564 (2012). https://doi.org/10.1515/hsz-2012-0119

Liberti, M. V. & Locasale, J. W. The Warburg Effect: How Does it Benefit Cancer Cells? *Trends Biochem Sci* **41**, 211-218 (2016). https://doi.org/10.1016/j.tibs.2015.12.001

Else, T. *et al.* Adrenocortical carcinoma. *Endocr Rev* **35**, 282-326 (2014). https://doi.org/10.1210/er.2013-1029

Miller, W. L. Steroid hormone synthesis in mitochondria. *Mol Cell Endocrinol* **379**, 62-73 (2013). https://doi.org/10.1016/j.mce.2013.04.014

Munson, M. J. *et al.* GAK and PRKCD are positive regulators of PRKN-independent mitophagy. *Nature communications* **12**, 6101 (2021). https://doi.org/10.1038/s41467-021-26331-7

Wandinger-Ness, A. & Zerial, M. Rab proteins and the compartmentalization of the endosomal system. *Cold Spring Harbor perspectives in biology* **6**, a022616 (2014). https://doi.org/10.1101/cshperspect.a022616

Marchler-Bauer, A. *et al.* CDD: NCBI's conserved domain database. *Nucleic Acids Res* **43**, D222-226 (2015). https://doi.org/10.1093/nar/gku1221

Baietti, M. F. *et al.* Syndecan–syntenin–ALIX regulates the biogenesis of exosomes. *Nature cell biology* **14**, 677-685 (2012). https://doi.org/10.1038/ncb2502

Yim, W. W.-Y., Yamamoto, H. & Mizushima, N. A pulse-chasable reporter processing assay for mammalian autophagic flux with HaloTag. *eLife* **11**, e78923 (2022). https://doi.org/10.7554/eLife.78923

Princely Abudu, Y. *et al.* NIPSNAP1 and NIPSNAP2 Act as "Eat Me" Signals for Mitophagy. *Developmental cell* **49**, 509-525.e512 (2019). https://doi.org/10.1016/j.devcel.2019.03.013

Hill, B. G. *et al.* Integration of cellular bioenergetics with mitochondrial quality control and autophagy. *Biol Chem* **393**, 1485-1512 (2012). https://doi.org/10.1515/hsz-2012-0198

Tang, Z. *et al.* GEPIA: a web server for cancer and normal gene expression profiling and interactive analyses. *Nucleic Acids Res* **45**, W98-w102 (2017). https://doi.org/10.1093/nar/gkx247

Wang, T. & Rainey, W. E. Human adrenocortical carcinoma cell lines. *Mol Cell Endocrinol* **351**, 58-65 (2012). https://doi.org/10.1016/j.mce.2011.08.041

Ishida, M., Arai, S. P., Ohbayashi, N. & Fukuda, M. The GTPase-deficient Rab27A(Q78L) mutant inhibits melanosome transport in melanocytes through trapping of Rab27A effector protein Slac2-a/melanophilin in their cytosol: development of a novel melanosome-targetinG tag. *The Journal of biological chemistry* **289**, 11059-11067 (2014). https://doi.org/10.1074/jbc.M114.552281

Key, J. *et al.* Ubiquitylome profiling of Parkin-null brain reveals dysregulation of calcium homeostasis factors ATP1A2, Hippocalcin and GNA11, reflected by altered firing of noradrenergic neurons. *Neurobiol Dis* **127**, 114-130 (2019). https://doi.org/10.1016/j.nbd.2019.02.008

Caserta, T. M., Smith, A. N., Gultice, A. D., Reedy, M. A. & Brown, T. L. Q-VD-OPh, a broad spectrum caspase inhibitor with potent antiapoptotic properties. *Apoptosis* **8**, 345-352 (2003). https://doi.org/10.1023/a:1024116916932

TeSlaa, T. & Teitell, M. A. Techniques to monitor glycolysis. *Methods Enzymol* **542**, 91-114 (2014). https://doi.org/10.1016/b978-0-12-416618-9.00005-4

Cox, J. & Mann, M. MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat Biotechnol* **26**, 1367-1372 (2008). https://doi.org/10.1038/nbt.1511

Tyanova, S. *et al.* The Perseus computational platform for comprehensive analysis of (prote)omics data. *Nat Methods* **13**, 731-740 (2016). https://doi.org/10.1038/nmeth.3901

Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* **43**, e47 (2015). https://doi.org/10.1093/nar/gkv007

Dessau, R. B. & Pipper, C. B. ["R"--project for statistical computing]. *Ugeskr Laeger* **170**, 328-330 (2008).

Kevin Blighe, S. R., Myles Lewis. *EnhancedVolcano: publication-ready volcano plots with enhanced colouring and labeling*, <https://github.com/kevinblighe/EnhancedVolcano> (2021).

Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for animals and plants. *Bioinformatics* **36**, 2628-2629 (2020). https://doi.org/10.1093/bioinformatics/btz931

Colaprico, A. *et al.* TCGAbiolinks: an R/Bioconductor package for integrative analysis of TCGA data. *Nucleic Acids Res* **44**, e71 (2016). https://doi.org/10.1093/nar/gkv1507

Maglich, J. M., Kuhn, M., Chapin, R. E. & Pletcher, M. T. More than just hormones: H295R cells as predictors of reproductive toxicity. *Reprod Toxicol* **45**, 77-86 (2014). https://doi.org/10.1016/j.reprotox.2013.12.009

Lin, C. W., Chang, Y. H. & Pu, H. F. Mitotane exhibits dual effects on steroidogenic enzymes gene transcription under basal and cAMP-stimulating microenvironments in NCI-H295 cells. *Toxicology* **298**, 14-23 (2012). https://doi.org/10.1016/j.tox.2012.04.007

Figure 1

A
SYTL5-EGFP MitoTracker Red

B
SYTL5-EGFP MitoTracker Red

C
SYTL5-EGFP LysoTracker Red

D
SYTL5-EGFP LAMP1

E

SHD 7 123 C2A 408 511 C2B 574 678 730
1 C2

SYTL5 full-length
SYTL5 (ΔSHD)
SYTL5 (ΔC2AB)

F

PIP strip

Left column	Full-length	ΔSHD	ΔC2AB	Control	Right column
LPA					S1P
LPC					PI(3,4)P2
PI					PI(3,5)P2
PI(3)P					PI(4,5)P2
PI(4)P					PI(3,4,5)P3
PI(5)P					PA
PE					PS
PC					Blank

Lipid membrane strip

Left column	Full-length	ΔSHD	ΔC2AB	Control	Right column
TAG					PI
DAG					PI(4)P
PA					PI(4,5)P2
PS					PI(3,4,5)P3
PE					Cholesterol
PC					SM
PG					Sulfatide
CL					Blank

G
SYTL5(ΔSHD)-EGFP MitoTracker Red

H
SYTL5(ΔC2AB)-EGFP MitoTracker Red

Figure 2

Figure 3

Figure 4

A

RAB27A SYTL5 dKO rescue with SYTL5-EGFP & mScarlet-RAB27A

B

RAB27A SYTL5 dKO rescue with SYTL5-EGFP & mScarlet-RAB27A

C

D

Figure 5

Figure 6

A

B

C

D

E

Supplementary Figure 1

Supplementary Figure 2

A ■ MitoTracker Green ■ mScarlet-RAB27A

B

C ■ SYTL5-EGFP ■ MitoTracker Red

HeLa

D ■ MitoTracker Green ■ mScarlet-RAB27A

HeLa

Supplementary Figure 3

A

RAB27A frameshift mutation
NM_183235 (2nd coding exon):

```
        WT  GTGTACAGAGCCAGTGGGCCGGATGGAGCCACTGGCAGAGGCCAGAGAATCCACCTGCAGTTATGGGACACAGCAGGGCAGGAGAG
            |||||||||||||||*||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
RAB27A KO c21 GTGTACAGAGCCAGT-GGCCGGATGGAGCCACTGGCAGAGGCCAGAGAATCCACCTGCAGTTATGGGACACAGCAGGGCAGGAGAG
```

B

C

```
GGAGGCTGTTCTACGTATAGCCTGAAAGAATACTTAACTACCATACGCAATTCTGCAGAGACCTTGTAAAAATCACACTTTACACCAAACAACTGAGTGATTCTGA
ATTGGTTGGGGGAATCTTAGTTGTAGATATCAATTCACCTTCTTGAAGATTCAACCACTGCTACTCAGAGCTGCTGCTGAAATACCATGTCTAAGAACTCAGAGTT
CATCAATCTGTCATTTTTATTAGATCATGAGAAGGAAATGATCCTGGGCGTCCTAAAGAGAGATGAATATTTGAAAAAGTGGAGGACAAGAGAATAAGGTAGTATT
CTTTTTATTTTTTCAGTCCTCCTTGACTGCTTTCTGTTTGTTTGAAGCAATTTAGATCAGAAATGGGCCAAGTGCAACCACAATGGGGACAGATGTCAGTTAGGAA
CTTCATGAGTTCTCATCAGTAGCCCAAATATCGGAAAACTTTGGCATTTCACCATTTAGCCAGACATGAAAATGGAATTTAGCTAATAAAGTGGTATTGATGAGAAC
ACTTAATTTGCTAATAAAGTCATAATTTTTAGTTAAGAGGAAGGACTACCTAAGTTTGAGTACCAGCTCTGGCAA
```

RAB27A/SYTL5 dKO deletion

```
GAGGCTGTTCTACGTATAGCCTGAAAGAATACTTAACTACCATACGCAATTCTGCAGAGACCTTGTAAAAATCACACTTTACACCAAACAACTGAGTGATTCTGA
ATTGGTTGGGGGAATCTTAGTTGTAGATATCAATTCACCTTCTTGAAGATTCAACCACTGCTACTCAGAGCTGCTGCTGAAATACCATGTCTAAGAACTCAGAGT
TCATCAATCTGTCATTTTTATTAGATCATGAGAAGGAAATGATCCTGGGCGTCCTAAAGAGAGATGAATATTTGAAAAAGTGGAGGACAAGAGAATAAGGTAGTA
TTCTTTTTATTTTTTCAGTCCTCCTTGACTGCTTTCTGTTTGTTTGAAGCAATTTAGATCAGAAATGGGCCAAGTGCAACCACAATGGGGACAGATGTCAGTTAGG
AACTTCATGAGTTCTCATCAGTAGCCCAAATATCGGAAAACTTTGGCATTTCACCATTTAGCCAGACATGAAAATGGAATTTAGCTAATAAAGTGGTATTGATGAGA
ACACTTAATTTGCTAATAAAGTCATAATTTTTAGTTAAGAGGAAGGACTACCTAAGTTTGAGTACCAGCT
```

D

Forward primer CAGGTCCCTTTCTTCTCGCA
Reverse primer TCAGTCAGCTGCAAGAGTGG

Supplementary Figure 4

A

RAB27A SYTL5 dKO rescue with SYTL5-EGFP & mScarlet-RAB27A

B

RAB27A SYTL5 dKO rescue with SYTL5-EGFP & mScarlet-RAB27A

Supplementary Figure 5

Supplementary Figure 6