Predicting microbial transcriptome using genome sequence

Guohao Fu[1, 5], Yujing Yan[2, 5], Ye Chen[3 *], Bin Shao[4, 5 *]

[1] Xuteli School, Beijing Institute of Technology, Beijing 100081, P. R. China

[2] Department of Organismic and Evolutionary Biology, Harvard University Herbaria, Harvard University,

Cambridge MA 02138, USA

[3] Key Laboratory of Quantitative Synthetic Biology, Shenzhen Institute of Synthetic Biology, Shenzhen

Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen, 518055, China

[4] Advanced Research Institute of Multidisciplinary Sciences, Beijing Institute of Technology, Beijing

100081, P. R. China

[5] These authors contribute equally to this work.

*Correspondence should be addressed to Ye Chen (chenye.h.ai@gmail.com), Bin Shao

(shaobinlx@gmail.com)

Abstract: We present TXpredict, a transformer-based framework for predicting microbial transcriptomes

using annotated genome sequences. By leveraging information learned from a large protein language

model, TXpredict achieves an average Spearman correlation of 0.53 and 0.62 in predicting gene

expression for new bacterial and fungal genomes. We further extend this framework to predict

transcriptomes for 2,685 additional microbial genomes spanning 1,744 genera, 82% of which remain

uncharacterized at the transcriptional level. Our analysis highlights conserved and divergent

transcriptional programs across understudied genera, providing a powerful resource for uncovering

microbial adaptation strategies and metabolic potential across the tree of life.

For a genome-sequenced microbe, transcriptome profiling provides critical insights into its adaptation, survival, and pathogenicity, and bridges the gap between raw genomic data and functional biology[1–6]. Despite its importance, only a small fraction of sequenced microbes has been characterized at the transcriptional level. A major challenge is that most microbes remain uncultivated due to the limited knowledge of their growth requirements or their inhabitation of extreme environments[7]. Recent advances in predicting physicochemical requirements have begun to address these challenges, representing an important first step[8–10]. Moreover, even when cultivation is successful, RNA-seq protocols often require organism-specific optimizations, such as cell lysis and rRNA depletion, which can involve extensive trial and error for novel organisms. These technical issues hinder large-scale, cross-species studies of microbial gene expression across the tree of life.

Current machine learning approaches for predicting gene expression have primarily focused on analyzing untranslated regions (UTRs)[11–16] or large genomic windows surrounding the gene of interest [17,18]. While these methods show promise, their application to non-model organisms remains limited, as trained models often fail to generalize across species or even within the same species[19,20]. Additionally, previous studies have demonstrated evolutionary constraints on protein synthesis, such as conserved enzyme stoichiometry in metabolic pathways[21–23]. Yet these constraints have not been translated into predictive models that are broadly applicable to diverse microbial species.

Here we introduce TXpredict, a novel framework for predicting microbial transcriptomes using only annotated genome sequences. Our approach assumes that gene expression is shaped by evolutionary constraints on protein sequences, and it leverages a protein language model, ESM2[24], to extract predictive features. Using transcriptome data from 22 bacterial, 9 archaeal and 9 fungal species obtained from the NCBI GEO database[25], we trained a transformer-based model that achieves a Spearman correlation up to 0.70 in leave-one-genome-out evaluations. We then validated the model's performance in the non-model organism *Pseudomonas sp.* WBC-3 by conducting a series of RNA-seq experiments. Additionally, we trained the model to account for condition-dependent gene expression using data from 13 bacterial species across 4.6k unique conditions and demonstrated that it can identify sequence determinants underlying these variations. Lastly, we applied the trained TXpredict models to predict transcriptomes for 2,685 additional genomes, the majority of them have never been transcriptionally characterized. This resulted in the creation of TXpredictDB, a database containing expression profiles for 9.8 million genes. The models developed in this study are publicly available: https://github.com/lingxusb/TXpredict

We first collected microbial RNA-seq datasets from the NCBI GEO datasets and selected the dataset with

the highest number of samples for each microbial species. Read counts were normalized with the same

pipeline (Methods). Taken together, our dataset includes 22 bacterial, 9 archaeal and 9 fungal species

and a total of 24.8M gene expression measurements (Supplementary Table 1). For each microbial

species, gene expression was averaged across all samples to construct a standard expression profile.

Protein embeddings for each gene were computed using the ESM2-650M model and combined with

basic protein statistics to serve as the input features for TXpredict (Fig. 1a, Methods). For model

evaluation, we employed a leave-one-genome-out cross-validation strategy. For bacterial species, the

model was trained on genes from 21 species and tested on the held-out species (Fig. 1b). We followed

the same approach for archaeal and fungal species. Model performance was evaluated using Spearman

correlation coefficients between predicted and measured expression values for the held-out genome.

We found our model achieved a mean Spearman correlation coefficient of 0.53 for the bacterial species

(Fig. 1c), and performed well for non-model organisms, with correlation coefficients of 0.64, 0.62 and

0.62 for *B. hinzii, B. thetaiotaomicron*, and *C. beijerinckii*. It is worth noting that transcriptome

predictions for these bacteria were based solely on the computational analysis of annotated genome

sequences, providing an alternative to labor-intensive RNA-seq experiments. The model's performance

in predicting gene expression variation decreased modestly as the variation calculations depend on the

diversity of experimental conditions (Supplementary Fig. 1). Despite having a limited training set for

fungal species, our TXpredict model obtained a mean correlation of 0.62 and reached a correlation

coefficient up to 0.70 for non-model organisms such as *O. oligospora* (Fig. 1c). For archaea, TXpredict

showed a mean Spearman correlation of 0.44, partly due to the limited availability of training data (Fig.

1b). The model demonstrated broad applicability to extremophiles from distinct natural niches (ranging

from Dead Sea to volcanic springs), further highlighting the robustness of our method.

In our initial evaluation, TXpredict performed poorly on the archaeon *N. maritimus*, with a Spearman

correlation of 0.23 in the leave-one-genome-out evaluation (Supplementary Fig. 2). Upon closer

examination of this dataset, we found that fewer than 40% of the reads were successfully mapped to

reference genome, demonstrating the potential of our model to detect potential problems in RNA-seq

data. In addition, we examined whether protein sequence similarity affects model performance. For

bacterium *B. hinzii*, we used BLASTP[26] to compute sequence similarity between its proteins and those in

the other bacterial strains. The model was trained on other species and tested on rare proteins from *B.*

hinzii that were not present in the training data, showing a correlation of 0.46 for these rare proteins

(Supplementary Fig. 3). While this is lower than the correlation for all proteins, it suggests that the

model captures expression-related features beyond sequence similarity.

We next explored TXpredict's cross-taxa performance by training separate models on bacterial,

archaeal, or fungal species and then testing them on other domains (Fig. 1d). Interestingly, we still found

our model showed moderate performance on cross-taxa predictions, suggesting that they captured

general principles of gene expression regulation. The archaeal model achieved a higher mean

correlation for eukaryotic species than for archaea, indicating that the gene expression is more

constrained in eukaryotes. At the species level, models trained on other taxa generally performed worse

than those trained on the same taxon (Fig. 1e and Supplementary Fig. 4).

To validate the prediction of the TXpredict model, we conducted new RNA-seq experiments on

Pseudomonas sp. WBC-3 and compared the predicted expression levels with the measured values. The

Spearman correlation between measured and predicted expression was 0.40 for all proteins and 0.50

for proteins with GO term annotations (Fig. 1f). Notably, our model showed high correlations (0.55) for

certain GO categories including catalytic activity, and also for operons like the flagellar genes (Fig. 1g and

Supplementary Fig. 5).

We next extended our framework to model condition-dependent gene expression (Fig. 1h). We

collected RNA-seq profiles from 13 bacterial strains across 4.6k unique conditions from iModulonDB,

resulting in a total of 12.0M measurements (Supplementary Table 2, Fig. 1i). The model integrated one-

hot encoded 5'UTR sequences with protein embeddings as input features. For each gene, we calculated

the correlation between measured and predicted expression values across all conditions. Our TXpredict

model achieved an average correlation coefficient of 0.52 across all bacterial strains (Fig. 1i). For *E. coli*,

the distributions of Pearson's and Spearman's correlation coefficients show similar patterns, with 41%

and 38% of genes having correlation coefficients above 0.5 (Fig. 1j).

To further interpret the model's predictions, we calculated input gradients[17] to identify sequence motifs

contributing to gene expression variation across conditions (Supplementary Fig. 6). In *E. coli*, we

observed a peak in the input × gradient overlap with the -10 box of the sigmaS-dependent promoter of

the *yadV* gene[27]. Also input gradients increased after the transcriptional start site, suggesting potential

contributions from transcribed sequences (Supplementary Fig. 5). In a non-model organism *P. putida*,

we identified a peak overlapping with a Zur binding motif for the *PP_0120* gene, consistent with its role

as a Zur iModulon gene[1]. These findings illustrate our model's ability to capture condition-specific gene

expression changes in an interpretable manner.

Lastly, we applied the trained TXpredict model to predict transcriptomes for new genomes including

2,169 bacterial, 423 archaeal and 93 fungal species (Fig. 2a and b). Collectively, these new genomes span

1,744 genera, of which 82% have no prior RNA-seq data according to publicly available data in the NCBI

GEO database. To explore the relationship between predicted gene expression and protein function, we

used DeepFRI[28] to predict Gene Ontology (GO) terms for the new genes (Supplementary Fig. 7). Among

the GO terms with the highest predicted expression levels were structural constituents of ribosome

(GO:0003735) (Fig. 2e, Supplementary Fig. 7). In contrast, GO terms with the lowest predicted

expression included proteins with DNA-binding activity (GO:0003677), consistent with their function as

transcriptional regulators. Collectively, our model predicted the expression of up to 9.8M genes across

these genomes, which were compiled into a database named TXpredictDB. The predicted gene

expressions in different domains followed similar distributions (Fig. 2f). When visualizing protein

embeddings of all proteins, we observed a clear separation of bacterial, archaeal and fungal genes (Fig.

2g, Supplementary Fig. 8). Small clusters of ribosomal genes located at the periphery of the plot and

showed high predicted expression, while transmembrane transporter proteins formed distinct low-

expression clusters.

We focused on genera without prior RNA-seq data and grouped genes based on their annotated Gene

Ontology (GO) terms (Supplementary Fig. 9). There was a positive correlation between the predicted

expression levels of tRNA ligases and ribosomal proteins across genera (Fig. 2h, Spearman correlation

0.52). *Actinoplanes* had the lowest predicted expression for ribosomal proteins, in accord with their

known slow growth rate[29]. The genus *Candidatus* includes a symbiotic bacterium found in carpenter ants

(*Blochmanniella*) and showed low expression of both tRNA ligases and ribosomal proteins. This pattern

likely reflects its reduced need for protein turnover and genome stasis[30], in contrast to free-living

bacteria. We further visualized the embeddings of tRNA ligases to explore their sequence variations (Fig.

2i). tRNA ligases from *Candidatus* formed distinct low-expression clusters, whereas those from

Acidiferrobacter were distributed across multiple clusters despite similarly low expression levels,

suggesting divergent evolutionary constraints associated with their ecological niche.

We also clustered protein sequences using MMseq2[31] and predicted gene expression within each high-

similarity cluster using the TXpredict model. Interestingly, several clusters displayed non-unimodal

expression distributions, with conserved amino acid differences separating low- and high-expression

variants (Fig. 2j). For example, in cluster of the PTS IIA component homologs, low-expression variants

shared a V116K substitution (Supplementary Fig. 10), which could reduce translation efficiency and

mRNA stability due to the presence of constitutive lysine residues[32,33].

We further analyzed the predicted gene expression profiles of biosynthetic gene clusters (BGCs) across

TXpredictDB, which were identified using antiSMASH[34] (n = 11,400). Although BGCs are typically lowly

expressed, distinct patterns were observed across different cluster types. For example, the proteusin

BGC showed high expression of core biosynthetic genes, whereas the ectoine BGC exhibited elevated

expression of regulatory genes (Supplementary Fig. 11). These reference expression profiles in non-

model organisms may offer valuable insights into the functional organization of BGCs.

In summary, we present TXpredict, a transcriptome prediction model that can generalize to a wide

range of microbes. For a microbial genome with 4.6k genes, transcriptome prediction can be completed

in approximately 22 minutes using a Google Colab notebook (Methods), demonstrating the model's

efficiency and accessibility. Our approach is not without limitations: while it effectively leverages protein

embeddings for expression prediction, it does not account for other regulatory elements or non-coding

RNAs. Moreover, our model's performance is constrained by the limited training profiles, especially for

archaeal species. Lastly, we provide a single reference transcriptome per genome in TXpredictDB, which

does not capture gene expression dynamics. Despite these limitations, TXpredict has the potential to

complement experimental approaches and advance the study of the vast number of sequenced yet

poorly characterized microbes. For example, identifying conserved expressed genes in pathogens may

enable the discovery of novel drug targets[35]. Furthermore, predicting gene expression across genomes

can provide insight into how genome plasticity contributes to bacterial pathogenicity[36]. Extending

transcriptome prediction to unculturable microbes also provides critical insights into microbial

adaptation to extreme environments and the evolution of gene regulation across prokaryotes[37,38]. These

efforts could further benefit from integrating single-cell RNA-seq data [5,39,40] and computational tools for

mRNA synthesis and stability prediction[41–43], paving the way for a deeper understanding of microbial

transcriptional landscapes.

Figures

Figure 1. Prediction of transcriptome based on the genome sequence

a) Schematic of TXpredict for transcriptome prediction. **b)** Scatter plots showing predicted and

measured gene expression in leave-one-genome-out evaluations for one species from bacteria, archaea,

and fungi. Spearman correlation coefficients are reported in the top-left corner of each plot. **c)**

Spearman correlation coefficients between predicted and measured gene expression for all species.

TXpredict models were separately trained for bacterial, archaeal, and fungal species. Results from five

rounds of model training and testing are reported (n = 5). Dashed lines indicate the mean correlations

for each domain. **d)** Cross-domain prediction performance of TXpredict models. Models were trained on

all species from one domain and tested on species from the other domains. Mean Spearman

correlations across all test species are shown. Diagonal values represent leave-one-genome-out

performance within each domain. **e)** Predictions of archaeal and fungal models on all bacterial species.

Blue dots indicate leave-one-genome-out evaluations for bacteria. Predictions for the same species

were connected. **f)** Model predictions versus measured gene expression in new RNA-seq experiment for

Pseudomonas sp. WBC-3. Each dot represents one functional gene and the Spearman correlation is

shown. **g)** Predicted and measured gene expression for genes with different GO annotations in

Pseudomonas sp. WBC-3. **h)** Schematic of TXpredict for condition-specific gene expression prediction. **i)**

Mean correlations for models trained separately on different bacterial species. Correlations from 5-fold

cross-validation tests are shown on the left (n = 5). Gene and condition counts for each species are

shown on the right. **j)** Distribution of Spearman and Pearson correlation coefficients for gene expression

across conditions in *E. coli* (n = 498).

Figure 2. Application of TXpredict to new genomes

a) Transcriptome prediction for new genomes. **b)** Genome sizes and gene counts for the new genomes

(n = 2,685). **c)** Number of genera in the new genomes (light colors) compared to training genomes (dark

colors). **d)** Polygenetic tree of newly added genomes. Each dot represents a phylum. The dot size is

proportional to the number of genera it contains and color indicates the proportion of genera with

existing RNA-seq data based on the NCBI GEO database. **e)** Top 20 GO terms of the genes in the new

genomes calculated by the DeepFRI model[28] (left) and their predicted gene expression levels (right). **f)**

Distributions of predicted gene expression levels across new bacterial (blue), archaeal (orange) and

fungal (green) genomes. **g)** t-SNE plots of protein embeddings generated by the ESM2 model[24] (n =

9.8M). Each dot represents a gene in the new genomes, colored by taxa (left), predicted gene expression

(middle), and GO terms (right). GO terms are colored as in panel (e). **h)** Predicted expression of tRNA

ligases and ribosomal proteins across understudied genera (n = 40), colored by phylum. **i)** t-SNE plots of

tRNA ligase embeddings across understudied genera, colored by phylum (left) and predicted expression

level (right). **j)** Histogram of predicted expression levels for proteins in a high-similarity cluster. Inset

shows the t-SNE plot of embeddings for these proteins, colored by predicted expression.

Methods

Training dataset

We collected RNA-seq data from the NCBI GEO database (https://www.ncbi.nlm.nih.gov/geo/). For each

microbe species, we selected the dataset with the highest number of samples and used either raw count

data or RPKM (Reads Per Kilobase of transcript per Million mapped reads) values when available. For

raw count data, we calculated the RPKM for each gene, followed by log (1 + p) and z-score

normalization. For RPKM values, we applied log (1 + p) transformation and z-score normalization. In

total, we gathered data from 22 bacterial species (2,649 samples), 9 archaeal species (169 samples) and

9 fungal species (2,078 samples, Supplementary Table 1). For each species, we constructed a standard

gene expression profile by averaging the normalized expression values across all available samples.

To model gene expression across conditions, we also collected RNA-seq data from iModulonDB

(https://imodulondb.org/index.html). We downloaded the processed datasets for each microbial

species, resulting in a dataset that includes 13 bacterial species and a total of 4,619 conditions. For each

microbial species, we only kept the genes with mean normalized gene expressions greater than -3 and a

standard deviation less than 1 across conditions.

Protein embeddings for all annotated genes were calculated using the ESM2-650M model

(*esm.pretrained.esm2_t33_650M_UR50D()*), following the instructions provided at

https://github.com/facebookresearch/esm. Proteins longer than 1,500 amino acids (AAs) were excluded

from this analysis. In addition to the protein embeddings, we also included basic protein sequence

statistics in the model input: the normalized protein length (calculated by dividing each protein length

by the longest protein length within each organism) and the amino acid composition (the proportion of

each of the 20 amino acids). The final input dimension of the model is 1301.

Transcriptome prediction model

We developed a Transformer-based model to predict gene expression levels from sequence features.

The transformer encoder utilizes a multi-head self-attention mechanism (num_heads=4) with a single

encoder layer. The encoder processes input features through a linear transformation into a hidden

dimension of 128, followed by positional encoding. We implemented sinusoidal positional encoding to

capture sequential information. The output of the transformer layer was then fed into two fully

connected layers (dimension 512 and 128) with ReLU activations. The final linear layer produces a single

output value representing the predicted gene expression level.

Model training and evaluation

We performed separate cross-species evaluations for bacteria and archaea to account for potential

domain-specific expression patterns. For each domain, we employed a cross-validation strategy where

models were trained on data from all but one microbial species and tested on the held-out species. This

approach was repeated for each species in our dataset to assess generalization capability. The models

were trained using the Adam optimizer with a learning rate of 0.0002 for 5 epoch with a batch size of 32.

To ensure robust evaluation, we performed 5 independent training runs with different random seeds

(42, 123, 456, 789, 1011). The mean squared error (MSE) loss function was used for optimization. Model

performance was evaluated using both Spearman and Pearson correlation coefficients between

predicted and measured expression levels.

New bacterial, archaeal and fungal genomes were downloaded from NCBI Genbank

(https://ftp.ncbi.nlm.nih.gov/genomes/genbank/). To maximize genus-level representation for

transcriptome prediction, we selected 2,169 bacterial, 423 archaeal, and 93 fungal genomes spanning a

total of 1,744 genera. Gene expression for these genomes was predicted using TXpredict models,

forming the basis of the TXpredictDB. To determine whether a genus has been transcriptional

characterized, we searched the GEO database using the following query: "*genus_name [Organism] AND*

("Expression profiling by array"[Filter])". We annotated taxonomy of these genomes by querying the

GTDB database[44]. For model training and inference, we used 3090 Ti GPU (24GB) and the software

packages PyTorch (version 2.1.1) and transformer (version 4.28.1).

Conditional gene expression model

The conditional gene expression model shares the same transformer encoder architecture with the

transcriptome prediction model, and trained separately for each species. The UTR branch processes

one-hot encoded sequences from -100 to +100 bp relative to the transcription start site. The

transformer layer has 2 attention heads and a feedforward dimension of 128. The protein embedding

branch utilizes 8 attention heads with a hidden dimension of 256. The outputs from both branches are

concatenated together and processed through a fully connected layer (dimension: 4096) with ReLU

activation. The output dimension of the model is the number of conditions for each bacterial species.

For each bacterial species, we trained the model for 10 epochs using the Adam optimizer with a learning

rate of 0.0002 and a batch size of 32. Model performance was evaluated using 5-fold cross-validation.

Pearson correlation coefficients were calculated between predicted and measured expression levels

across all conditions for genes in the test set.

Analysis of protein sequences

To visualize the high-dimensional embeddings in a two-dimensional space, we implemented t-SNE (t-

Distributed Stochastic Neighbor Embedding) using the *opentsne* package. We initialized the embedding

using PCA and constructed a uniform affinity matrix with 10 nearest neighbors. The optimization process

consisted of three phases: an early exaggeration phase of 125 iterations with an exaggeration factor of

12 and momentum of 0.5, followed by a gradual exaggeration annealing phase where the exaggeration

factor was linearly decreased from 12 to 1 over 125 iterations with momentum of 0.8. The final

optimization phase ran for 2000 iterations without exaggeration. Function prediction for these proteins

was carried out using the default deepFRI model[28], as available on GitHub

(https://github.com/flatironinstitute/DeepFRI). We used a score cutoff of 0.5 which was reported to be

significant in the original publication. To cluster the protein sequences in TXpredictDB, we used the

MMseqs2[31] package with default settings and a minimum sequence identity threshold of 0.9.

Calculation of input gradient

To investigate the contribution of individual nucleotides in the UTR sequence to gene expression

prediction across conditions, we implemented a gradient-based analysis. Using a sequence-only model

variant where the protein embedding branch was masked, we calculated the gradient of the model

output with respect to the input DNA sequence. For each target gene and condition, we performed a

forward pass with the one-hot encoded UTR sequence while setting all other inputs to zero. We then

computed the gradients through backpropagation and multiplied them element-wise with the input

sequence (Gradient × Input method)[17]. The absolute values of these products were summed across the

four nucleotide channels (A, T, C, G) to obtain single-base resolution importance scores. For visualization

purposes, we aggregated these scores into 5-bp bins by summing the scores within each bin.

RNA-seq experiment of the *Pseudomonas* sp. WBC-3 strain

For the RNA-seq experiment, *Pseudomonas* sp. WBC-3 was cultured under three different conditions to:

PNP1 (minimal medium with 0.01% glycerol), PNP2 (minimal medium with 0.04% PNP), and PNP3

(minimal medium with 0.01% glycerol and 0.04% PNP). Each condition was prepared by diluting a 2×

stock of yeast extract minimal medium supplemented with either 0.02% glycerol, 0.08% PNP, or both,

resulting in the desired final concentrations. WBC-3 strain was first grown overnight in LB medium, then

diluted 1:100 into 4 mL of each experimental medium and incubated overnight at 30°C with shaking at

200 rpm. Cultures were harvested at OD600 between 0.5 and 1.0 by centrifugation for 2 minutes,

ensuring that the total collected biomass did not exceed $1×10^9$ cells. Total RNA was extracted using the

RNAprep Pure Bacteria Kit (TIANGEN, DP430), followed by DNase I treatment (TIANGEN, RT411) to

remove genomic DNA contamination. RNA quality and concentration were confirmed to meet

sequencing requirements. Raw sequencing reads were aligned and processed using standard RNA-seq

pipelines. Read counts were quantified using StringTie with the following command: stringtie -p 8 -G

GCF_035758525.1_ASM3575852v1_genomic.gtf -l sample_id -o sample.gtf sample.bam, using the

Pseudomonas sp. WBC-3 reference genome (GCF_035758525.1).

Code availability

Our trained model and inference codes are available from GitHub:

https://github.com/lingxusb/TXpredict

References

1.   Lim, H. G. *et al.* Machine-learning from Pseudomonas putida KT2440 transcriptomes reveals its
transcriptional regulatory network. *Metab Eng* **72**, 297–310 (2022).

2.   Patel, A. *et al.* Proteome allocation is linked to transcriptional regulation through a modularized
transcriptome. *Nat Commun* **15**, 5234 (2024).

3.   Ryan, D. *et al.* An expanded transcriptome atlas for Bacteroides thetaiotaomicron reveals a small
RNA that modulates tetracycline sensitivity. *Nat Microbiol* **9**, 1130–1144 (2024).

4.   Shin, J., Rychel, K. & Palsson, B. O. Systems biology of competency in Vibrio
natriegens is revealed by applying novel data analytics to the transcriptome. *Cell Rep* **42**,
(2023).

5.   Ma, P. *et al.* Bacterial droplet-based single-cell RNA-seq reveals antibiotic-associated
heterogeneous cellular states. *Cell* **186**, 877–891 (2023).

6.   Ryu, M.-H. *et al.* Control of nitrogen fixation in bacteria that associate with cereals. *Nat Microbiol*
5, 314–330 (2020).

7.   G, L. K., D, S. A., Joshua, L., Junqi, Y. & Lonnie, C. Phylogenetically Novel Uncultured Microbial
Cells Dominate Earth Microbiomes. *mSystems* **3**, 10.1128/msystems.00055-18 (2018).

8.   Barnum, T. P. *et al.* Predicting microbial growth conditions from amino acid composition. *bioRxiv*
2023–2024 (2024).

9.   Ramoneda, J. *et al.* Building a genome-based understanding of bacterial pH preferences. *Sci Adv*
9, eadf8998 (2024).

10. Price, M. N., Deutschbauer, A. M. & Arkin, A. P. Filling gaps in bacterial catabolic pathways with
computation and high-throughput genetics. *PLoS Genet* **18**, e1010156- (2022).

11. Zrimec, J. *et al.* Deep learning suggests that gene expression is encoded in all parts of a co-
evolving interacting gene regulatory structure. *Nat Commun* **11**, 6141 (2020).

12. Peleke, F. F., Zumkeller, S. M., Gültas, M., Schmitt, A. & Szymański, J. Deep learning the cis-
regulatory code for gene expression in selected model plants. *Nat Commun* **15**, 3488 (2024).

13. Dudnyk, K., Cai, D., Shi, C., Xu, J. & Zhou, J. Sequence basis of transcription initiation in the human
genome. *Science (1979)* **384**, eadj0116 (2024).

14. LaFleur, T. L., Hossain, A. & Salis, H. M. Automated model-predictive design of synthetic
promoters to control transcriptional profiles in bacteria. *Nat Commun* **13**, 5159 (2022).

15. Urtecho, G., Tripp, A. D., Insigne, K. D., Kim, H. & Kosuri, S. Systematic Dissection of Sequence
Elements Controlling σ70 Promoters Using a Genomically Encoded Multiplexed Reporter Assay in
Escherichia coli. *Biochemistry* **58**, 1539–1551 (2019).

16. Vaishnav, E. D. *et al.* The evolution, evolvability and engineering of gene regulatory DNA. *Nature*
(2022) doi:10.1038/s41586-022-04506-6.

17. Avsec, Ž. *et al.* Effective gene expression prediction from sequence by integrating long-range
interactions. *Nat Methods* **18**, 1196–1203 (2021).

18. Kelley, D. R. *et al.* Sequential regulatory activity prediction across chromosomes with
convolutional neural networks. *Genome Res* **28**, 739–750 (2018).

19. Huang, C. *et al.* Personal transcriptome variation is poorly explained by current genomic deep
learning models. *Nat Genet* **55**, 2056–2059 (2023).

20. Sasse, A. *et al.* Benchmarking of deep neural networks for predicting personal gene expression
from DNA sequence highlights shortcomings. *Nat Genet* **55**, 2060–2064 (2023).

21. Li, G.-W., Burkhardt, D., Gross, C. & Weissman, J. S. Quantifying Absolute Protein Synthesis Rates
Reveals Principles Underlying Allocation of Cellular Resources. *Cell* **157**, 624–635 (2014).

22. Lalanne, J.-B. *et al.* Evolutionary Convergence of Pathway-Specific Enzyme Expression
Stoichiometry. *Cell* **173**, 749-761.e38 (2018).

23. Klumpp, S., Scott, M., Pedersen, S. & Hwa, T. Molecular crowding limits translation and cell
growth. *Proceedings of the National Academy of Sciences* **110**, 16754–16759 (2013).

24. Lin, Z. *et al.* Evolutionary-scale prediction of atomic-level protein structure with a language
model. *Science (1979)* **379**, 1123–1130 (2023).

25. Clough, E. & Barrett, T. The gene expression omnibus database. *Statistical Genomics: Methods
and Protocols* 93–110 (2016).

26. Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421 (2009).

27. Raina, S., Missiakas, D., Baird, L., Kumar, S. & Georgopoulos, C. Identification and transcriptional
analysis of the Escherichia coli htrE operon which is homologous to pap and related pilin operons.
J Bacteriol **175**, 5009–5021 (1993).

28. Gligorijević, V. *et al.* Structure-based protein function prediction using graph convolutional
networks. *Nat Commun* **12**, 3168 (2021).

29. Couch, J. N. Actinoplanes, a new genus of the Actinomycetales. *Journal of the Elisha Mitchell
Scientific Society* **66**, 87–92 (1950).

30. Degnan, P. H., Lazarus, A. B. & Wernegreen, J. J. Genome sequence of Blochmannia
pennsylvanicus indicates parallel evolutionary trends among bacterial mutualists of insects.
Genome Res **15**, 1023–1033 (2005).

31. Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence searching for the
analysis of massive data sets. *Nat Biotechnol* **35**, 1026–1028 (2017).

32. Radhakrishnan, A. & Green, R. Connections underlying translation and mRNA stability. *J Mol Biol*
428, 3558–3564 (2016).

33. Meydan, S. & Guydosh, N. R. Disome and Trisome Profiling Reveal Genome-wide Targets of
Ribosome Quality Control. *Mol Cell* **79**, 588-602.e6 (2020).

34.   Blin, K. *et al.* antiSMASH 7.0: new and improved predictions for detection, regulation, chemical
structures and visualisation. *Nucleic Acids Res* **51**, W46–W50 (2023).

35.   Wang, C., Hong, Z., Rongfen, Z., Clement Kin-Ming, T. & and Deng, S. Genomic Characterization
and Antimicrobial Resistance of ESBL-Producing, Escherichia coli Isolates in Suzhou, China. *Infect
Drug Resist* **18**, 1049–1057 (2025).

36.   Krishnakant Kushwaha, S. *et al.* Comprehensive blueprint of Salmonella genomic plasticity
identifies hotspots for pathogenicity genes. *PLoS Biol* **22**, e3002746- (2024).

37.   Shu, W.-S. & Huang, L.-N. Microbial diversity in extreme environments. *Nat Rev Microbiol* **20**,
219–235 (2022).

38.   Caro-Astorga, J. *et al.* Polyextremophile engineering: a review of organisms that push the limits of
life. *Front Microbiol* **15**, (2024).

39.   Imdahl, F., Vafadarnejad, E., Homberger, C., Saliba, A.-E. & Vogel, J. Single-cell RNA-sequencing
reports growth-condition-specific global transcriptomes of individual bacteria. *Nat Microbiol* **5**,
1202–1206 (2020).

40.   Kuchina, A. *et al.* Microbial single-cell RNA sequencing by split-pool barcoding. *Science (1979)*
371, eaba5257 (2021).

41.   LaFleur, T. L., Hossain, A. & Salis, H. M. Automated model-predictive design of synthetic
promoters to control transcriptional profiles in bacteria. *Nat Commun* **13**, 5159 (2022).

42.   Cetnar, D. P., Hossain, A., Vezeau, G. E. & Salis, H. M. Predicting synthetic mRNA stability using
massively parallel kinetic measurements, biophysical modeling, and machine learning. *Nat
Commun* **15**, 9601 (2024).

43.   Xia, Y. *et al.* Language model generates cis-regulatory elements across prokaryotes. *bioRxiv*
2011–2024 (2024).

44.   Parks, D. H. *et al.* GTDB: an ongoing census of bacterial and archaeal diversity through a
phylogenetically consistent, rank normalized and complete genome-based taxonomy. *Nucleic
Acids Res* **50**, D785–D794 (2022).

45.   Jeckel, H. *et al.* Simultaneous spatiotemporal transcriptomics and microscopy of Bacillus subtilis
swarm development reveal cooperation across generations. *Nat Microbiol* **8**, 2378–2391 (2023).

46.   Rachael, N., Swathi, P., S, P. F. & Peter, B. Metabolic changes associated with polysaccharide
utilization reduce susceptibility to some β-lactams in Bacteroides thetaiotaomicron. *mSphere* **9**,
e00103-24 (2024).

47.   Kamat, A. *et al.* Widespread prevalence of a methylation-dependent switch to activate an
essential DNA damage response in bacteria. *PLoS Biol* **22**, e3002540- (2024).

48.   Lakey, B. D. *et al.* The essential Rhodobacter sphaeroides CenKR two-component system
regulates cell division and envelope biosynthesis. *PLoS Genet* **18**, e1010270- (2022).

49. A, P. D. *et al.* Exogenous butyrate inhibits butyrogenic metabolism and alters virulence
phenotypes in Clostridioides difficile. *mBio* **15**, e02535-23 (2024).

50. D, S. M. *et al.* Arabinose-Induced Catabolite Repression as a Mechanism for Pentose Hierarchy
Control in Clostridium acetobutylicum ATCC 824. *mSystems* **3**, 10.1128/msystems.00064-18
(2018).

51. Rahimi, S. *et al.* Co-culturing Bacillus subtilis and wastewater microbial community in a bio-
electrochemical system enhances denitrification and butyrate formation. *Chemical Engineering*
Journal **397**, 125437 (2020).

52. Pennings, J. L. A. *et al.* Regulation of Clostridium tetani Neurotoxin Expression by Culture
Conditions. *Toxins (Basel)* **14**, 31 (2022).

53. Espah Borujeni, A., Zhang, J., Doosthosseini, H., Nielsen, A. A. K. & Voigt, C. A. Genetic circuit
characterization by inferring RNA polymerase movement and ribosome usage. *Nat Commun* **11**,
5001 (2020).

54. Wilson, E. H. *et al.* A Computational Framework for Identifying Promoter Sequences in Nonmodel
Organisms Using RNA-seq Data Sets. *ACS Synth Biol* **10**, 1394–1405 (2021).

55. Young, A. T. *et al.* Multi-omic regulatory networks capture downstream effects of kinase
inhibition in Mycobacterium tuberculosis. *NPJ Syst Biol Appl* **7**, 8 (2021).

56. Xavier, C. *et al.* Multisystem Analysis of Mycobacterium tuberculosis Reveals Kinase-Dependent
Remodeling of the Pathogen-Environment Interface. *mBio* **9**, 10.1128/mbio.02333-17 (2018).

57. Wu, M.-L. *et al.* Developmental transcriptome of resting cell formation in Mycobacterium
smegmatis. *BMC Genomics* **17**, 837 (2016).

58. Nobori, T. *et al.* Transcriptome landscape of a bacterial pathogen under plant immunity.
Proceedings of the National Academy of Sciences **115**, E3055–E3064 (2018).

59. M, E. J. *et al.* Integrative genomic, virulence, and transcriptomic analysis of emergent
Streptococcus dysgalactiae subspecies equisimilis (SDSE) emm type stG62647 isolates causing
human infections. *mBio* **15**, e02578-24 (2024).

60. Kachroo, P. *et al.* Integrated analysis of population genomics, transcriptomics and virulence
provides novel insights into Streptococcus pyogenes pathogenesis. *Nat Genet* **51**, 548–559
(2019).

61. Martim, D. B. *et al.* Resolving the metabolism of monolignols and other lignin-related aromatic
compounds in Xanthomonas citri. *Nat Commun* **15**, 7994 (2024).

62. M, V. J. *et al.* Genome-Scale Transcription-Translation Mapping Reveals Features of Zymomonas
mobilis Transcription Units and Promoters. *mSystems* **5**, 10.1128/msystems.00250-20 (2020).

63. Sakrikar, S. & Schmid, A. K. An archaeal histone-like protein regulates gene expression in
response to salt stress. *Nucleic Acids Res* **49**, 12732–12743 (2021).

64. Hackley, R. K. *et al.* TbsP and TrmB jointly regulate gapII to influence cell development
phenotypes in the archaeon Haloferax volcanii. *Mol Microbiol* **121**, 742–766 (2024).

65. D, T. B., Cem, M., B, N. T., Q, L. S. & F, H. J. Growth Kinetics, Carbon Isotope Fractionation, and
Gene Expression in the Hyperthermophile Methanocaldococcus jannaschii during Hydrogen-
Limited Growth and Interspecies Hydrogen Transfer. *Appl Environ Microbiol* **85**, e00180-19
(2019).

66. Payne, D. *et al.* Alternative sources of molybdenum for Methanococcus maripaludis and their
implication for the evolution of molybdoenzymes. *Commun Biol* **7**, 1337 (2024).

67. Spietz, R. L. *et al.* Investigating Abiotic and Biotic Mechanisms of Pyrite Reduction. *Front*
Microbiol **13**, (2022).

68. Yang, Y. *et al.* A novel RHH family transcription factor aCcr1 and its viral homologs dictate cell
cycle progression in archaea. *Nucleic Acids Res* **51**, 1707–1723 (2023).

69. Topçuoğlu, B. D., Meydan, C., Orellana, R. & Holden, J. F. Formate hydrogenlyase and formate
secretion ameliorate H2 inhibition in the hyperthermophilic archaeon Thermococcus
paralvinellae. *Environ Microbiol* **20**, 949–957 (2018).

70. G, D. E., E, S. J., Alix, A., Patricia, O. & A, H. D. Balancing Positive and Negative Selection: In Vivo
Evolution of Candida lusitaniae MRR1. *mBio* **12**, 10.1128/mbio.03328-20 (2021).

71. Kang, Y. S. *et al.* Leveraging a new data resource to define the response of Cryptococcus
neoformans to environmental signals. *Genetics* **229**, iyae178 (2025).

72. Schnepf, V., Vlot, A. C., Kugler, K. & Hückelhoven, R. Barley susceptibility factor RACB modulates
transcript levels of signalling protein genes in compatible interaction with Blumeria graminis f.sp.
hordei. *Mol Plant Pathol* **19**, 393–404 (2018).

73. Hunsaker, E. W., Yu, C.-H. A. & Franz, K. J. Copper Availability Influences the Transcriptomic
Response of Candida albicans to Fluconazole Stress. *G3 Genes|Genomes|Genetics* **11**, jkab065
(2021).

74. D, M. N. *et al.* Independent component analysis reveals 49 independently modulated gene sets
within the global transcriptional regulatory architecture of multidrug-resistant Acinetobacter
baumannii. *mSystems* **9**, e00606-23 (2024).

75. Lamoureux, C. R. *et al.* A multi-scale expression and regulation knowledge base for Escherichia
coli. *Nucleic Acids Res* **51**, 10176–10193 (2023).

76. Jonathan, J.-S. *et al.* Reconstructing the transcriptional regulatory network of probiotic L. reuteri
is enabled by transcriptomics and machine learning. *mSystems* **9**, e01257-23 (2024).

77. Reo, Y. *et al.* Machine Learning of All Mycobacterium tuberculosis H37Rv RNA-seq Data Reveals a
Structured Interplay between Metabolism, Stress Response, and Infection. *mSphere* **7**, e00033-22
(2022).

78. Rajput, A. *et al.* Advanced transcriptomic analysis reveals the role of efflux pumps and media
composition in antibiotic responses of Pseudomonas aeruginosa. *Nucleic Acids Res* **50**, 9675–
9688 (2022).

79. Heera, B., Kevin, R., R, L. C., V, S. A. & O, P. B. Machine learning uncovers the Pseudomonas
syringae transcriptome in microbial communities and during infection. *mSystems* **8**, e00437-23
(2023).

80. Saugat, P. *et al.* Coordination of CcpA and CodY Regulators in Staphylococcus aureus USA300
Strains. *mSystems* **7**, e00480-22 (2022).

81. Yuan, Y. *et al.* Machine learning reveals the transcriptional regulatory network and circadian
dynamics of Synechococcus elongatus PCC 7942. *Proceedings of the National Academy of*
Sciences **121**, e2410492121 (2024).

82. Yujiro, H. *et al.* Elucidation of independently modulated genes in Streptococcus pyogenes reveals
carbon sources that control its expression of hemolytic toxins. *mSystems* **8**, e00247-23 (2023).

Supplementary Table

Supplementary Table S1. Datasets used in the transcriptome prediction model

Bacteria				
Species	**GSE number**	**Reference genome**	**Gene#**	**Condition#**
Bacillus subtilis	GSE224332[45]	NZ_CP020102.1	4220	317
Bacteroides thetaiotaomicron	GSE251676[46]	GCF_000011065.1	4618	64
Bordetella hinzii	GSE181542	CP052845.1	4531	228
Caulobacter vibrioides	GSE246782[47]	NC_011916.1	3882	34
Cereibacter sphaeroides	GSE186600[48]	GCA_000012905.2	4232	17
Chlamydia trachomatis	GSE262652	GCA_000068585.1	857	30
Clostridioides difficile	GSE249810[49]	GCA_000009205.5	3560	24
Clostridium acetobutylicum	GSE107804[50]	GCA_000008765.1	3755	48
Clostridium beijerinckii	GSE150480[51]	GCF_000833105.2	5316	16
Clostridium butyricum	GSE150480[51]	GCF_001456065.2	3848	16
Clostridium tetani	GSE182408[52]	NC_004557.1	2562	39
Escherichia coli	GSE152664[53]	NC_010473.1	3260	9
Erythrobacter litoralis	GSE126532	GCA_001719165.1	3003	49
Methylotuvimicrobium buryatense	GSE162089[54]	NZ_CP035467.1	3927	56
Mycobacterium tuberculosis	GSE110508[55,56]	GCA_000195955.2	3957	87
Mycolicibacterium smegmatis	GSE69983[57]	GCA_000015005.1	6700	30
Pseudomonas aeruginosa	GSE251671	GCA_000014625.1	5659	858
Pseudomonas syringae	GSE103441[58]	GCA_000007805.1	5586	114
Streptococcus dysgalactiae	GSE272047[59]	GCA_029234115.1	1939	114
Streptococcus pyogenes	GSE113055[60]	CP000056.2	1608	442
Xanthomonas citri	GSE252662[61]	AE008923.1	4294	38
Zymomonas mobilis	GSE139939[62]	CP023715.1	1759	19
Summary			83,073	2,649

508 **Supplementary Table S1 - continue. Datasets used in the transcriptome prediction model**

Archaea				
Species	**GSE number**	**Reference genome**	**Gene#**	**Condition#**
Haloarcula hispanica	GSE227031	GCF_000223905.1	3776	32
Halobacterium salinarum	GSE182493[63]	GCF_000006805.1	2325	24
Haloferax volcanii	GSE248564[64]	GCF_000025685.1	3873	39
Methanocaldococcus jannaschii	GSE112986[65]	GCF_000091665.1	1724	13
Methanococcus maripaludis	GSE261517[66]	GCF_000011585.1	1730	12
Methanosarcina barkeri	GSE168895[67]	GCF_000970025.1	3338	12
Sulfolobus islandicus	GSE220819[68]	GCF_000189555.1	2682	18
Thermococcus kodakarensis	GSE186817	NC_006624.1	2214	4
Thermococcus paralvinellae	GSE101078[69]	GCF_000517445.1	1995	15
Summary			25,448	169

509 **Supplementary Table S1 - continue. Datasets used in the transcriptome prediction model**

Fungi				
Species	**GSE number**	**Reference genome**	**Gene#**	**Condition#**
Clavispora lusitaniae	GSE162151[70]	ATCC42720_w_CBS_6936_MT	5851	54
Cryptococcus neoformans	GSE226255[71]	CP022321.1	6718	720
Saccharomyces cerevisiae	GSE234431	GCA_000146045.2	5498	897
Coccidioides posadasii	GSE279406	NC_089407.1	8103	144
Blumeria graminis	GSE69215[72]	GCA_000151065.3	6329	48
Lipomyces starkeyi	GSE276000	GCA_001661325.1	7928	36
Neurospora crassa	GSE181566	GCF_000182925.2	9532	80
Candida albicans	GSE159545[73]	GCF_000182965.3	5918	75
Orbilia oligospora	GSE245014	GCF_000225545.1	11195	24
Summary			67,072	2,078

Supplementary Table S2. Datasets used in the conditional gene expression model (https://imodulondb.org/)

Species	Reference	Reference genome	Gene#	Condition#
Acinetobacter baumannii	Menon, et al.[74]	CP008706.1	1788	139
Bacillus subtilis	Sastry, et al.	NC_000964.3	1969	265
Escherichia coli	Lamoureux, et al.[75]	U00096.3	2490	1035
Limosilactobacillus reuteri	Josephs-Spaulding, et al.[76]	GCA_020785475.1	1159	117
Mycobacterium tuberculosis	Yoo, et al.[77]	AL123456.3	3214	647
Pseudomonas aeruginosa	Rajput, et al.[78]	NC_002516.2	3981	411
Pseudomonas putida	Lim, et al.[1]	AE015451.2	4379	321
Pseudomonas syringae	Bajpe, et al.[79]	AE016853.1	4296	202
Staphylococcus aureus	Poudel, et al.[80]	NC_010079.1	1736	385
Synechococcus elongatus	Yuan, et al.[81]	NC_007604.1	1522	300
Salmonella enterica	Yuan, et al.[81]	AE006468.2	1808	533
Streptococcus pyogenes	Hirose, et al.[82]	CP008776.1	1173	116
Vibrio natriegens	Shin, et al.[4]	GCA_001456255.1	1981	148
Summary			31,496	4,619

Supplementary Figures

Supplementary Figure 1: Prediction of gene expression variation. Spearman correlations between

predicted and measured gene expression variations in leave-one-genome-out experiments for bacterial

(a), archaeal **(b)**, and fungal **(c)** species. Results from five rounds of model training and evaluation are

shown (n = 5). **d)** Number of conditions and corresponding mean Spearman correlations for expression

variation prediction in bacterial species (n = 22).

Supplementary Figure 2: **Detection of issues related to RNA-seq dataset. a)** Spearman correlation

coefficients between predicted and measured gene expression for all archaeal species in the initial

evaluation. Results from five rounds of model training and testing are shown (n = 5). **b)** Proportion of

mapped/unmapped reads in the *N. maritimus* dataset (GSE103699).

Supplementary Figure 3: Prediction of gene expression for rare proteins. a) Histogram showing the

distribution of overlap ratios between *B. hinzii* proteins and proteins in the training sequences. Overlap

ratios were calculated as the aligned sequence length divided by the query length. **b)** Predicted versus

measured gene expression for proteins with low homology to the training set (overlap ratio < 0.01). **c)**

Predicted and measured gene expressions for proteins with close homologs in the training dataset

(overlap > 0.99).

Supplementary Figure 4: Cross-domain prediction performance of TXpredict. TXpredict was trained on
all species from a single domain and evaluated on species from the two other domains. Results from
three domain-specific models are shown: bacteria (blue), archaea (orange), and fungi (green).

Supplementary Figure 5: Performance of TXpredict on *Pseudomonas sp.* WBC-3. a) Correlation of

measured gene expression values of *Pseudomonas sp. WBC-3* across three experimental conditions. **b)**

Predicted versus measured gene expression under each condition. **c)** Scatter plot comparing predicted

and mean measured gene expression across three conditions for all genes. **d)** Predicted and measured

gene expression levels for genes in the *fli* operon.

Supplementary Figure 6: **Sequence determinant of condition-dependent gene expression. a)**

Calculation of input gradients for identifying sequence determinants of condition-dependent gene

expression. **b)** Two examples of input × gradients and their correlations with known sequence motifs.

Supplementary Figure 7: Top GO terms and the predicted gene expressions in TXpredictDB. Top 20

Gene Ontology (GO) terms for genes from newly added bacterial, archaeal, and fungal genomes, as

annotated by the DeepFRI model[28]. Predicted gene expression levels for these genes are shown

alongside their associated GO terms.

<u>**Supplementary Figure 8:**</u> **Prediction of gene expression variation in newly added genomes. a)**

Predicted gene expression variation for genes associated with the top 20 GO terms in the newly added

genomes. **b)** Predicted gene expression variation visualized in t-SNE plots of protein embeddings (n =

9.8M).

Supplementary Figure 9: Prediction of gene expression for understudied genera. Gene expression

values for the top 40 GO terms and 40 genera are shown. Only genus–GO term combinations with more

than 10 associated genes are included, with mean gene expression levels presented.

Supplementary Figure 10: **Amino acid frequency differences between low and high expression genes**

within a protein cluster. TXpredict was used to predict gene expression for proteins in cluster

ERS451416_00727, identified using MMseq2[31] at a 90% sequence identity threshold. Differences in

amino acid frequencies between lowly and highly expressed genes within the cluster are shown.

Supplementary Figure 11: **Predicted gene expression across biosynthetic gene clusters (BGCs). a)**

Predicted gene expression levels for core biosynthetic, additional biosynthetic, regulatory, and transport

genes across different BGC types. **b)** Number of regions and predicted expression levels for the top 20

BGC types. **c)** Histogram of mean expression levels of BGCs (n = 11,400).