Family life and cadmium ingestion independently shape offspring microbiomes in a subsocial insect

Marie-Charlotte Cheutin[*#1], Romain Honorio[#1,2], Joël Meunier[1]

[1] Institut de Recherche sur la Biologie de l'Insecte, UMR 7261, CNRS, University of Tours, Tours, France
[2] Laboratoire d'Ethologie Expérimentale et Comparée UR 4443 (LEEC), Université Sorbonne Paris Nord, 93430 Villetaneuse, France

*Corresponding author: mccheutin@gmail.com
# These authors contributed equally to this work

ABSTRACT
Symbiotic microbes are essential for host health and fitness. In family-living species, these microbes are often acquired through vertical transmission from parents and horizontal transmission from siblings. However, it is unclear how environmental stressors, such as chemical pollution, influence these contributions to the juvenile microbiome. Here, we tested the independent and interactive effects of social environment and cadmium ingestion - a highly toxic and common heavy metal pollutant - on the microbiome of juvenile European earwigs. We reared 900 juveniles either alone, with siblings or with siblings plus the mother. We exposed them to cadmium-enriched food at 0, 25 or 100mg.L^{-1}, and analysed their microbiome composition and diversity at the end of the family life period. Our results showed that both social environment and cadmium exosure shaped the phylogenetic beta-diversity of the juvenile microbiome, with no evidence of an interaction between these factors. In contrast, alpha-diversity remained unaffected by either factor or their interaction. Notably, several specific bacterial taxa, including putatively pathogenic (Serratia) and mutualistic (Lactobacillus) symbionts, were more abundant in juveniles reared with family members than in those reared in isolation, reflecting classical patterns observed in social species. Overall, our findings suggest that while both social environment and cadmium shape the microbiome of earwig juveniles, family life neither amplifies nor mitigates the effects of chemical exposure. This highlights the robustness of microbial sharing within families, even under strong environmental stress.

Keywords: chemical pollutant; facultative social life; Parental care; Microbial heritability; Social microbiome

Introduction

Most animals harbour microorganisms that are essential to their health and fitness. These microorganisms play crucial roles in hosts' biological processes, such as aiding in nutrient acquisition (Berlanga & Guerrero, 2016; Calderón-Cortés et al., 2012), regulating the immune system (Horak et al., 2020), supporting development (Girard et al., 2023; Y. Wang & Rozen, 2017) and providing protection against pathogens (Muhammad et al., 2019; Shukla et al., 2018). The acquisition of these microorganisms, whether from the environment or from parents (*e.g.,* transovarial pathway), is thus essential for the host, especially for juveniles that are born or hatched with an underdeveloped microbiome (Videvall et al., 2019) or that have lost a large part of their microbiome during development, such as through moulting in insects (Cheutin, Boucicot, et al., 2024; Manthey et al., 2022; Schmidt & Engel, 2021).

In group-living species, high nest fidelity and high frequency of social interactions between group members are often key factors facilitating the acquisition of core microorganisms by juveniles (Onchuru et al., 2018). In family groups, this acquisition occur through two main pathways. The first pathway is vertical transmission through parental care - a widespread phenomenon across animal taxa (Trumbo, 2012). Parental care typically allows the transfer of microorganisms and community of microorganisms from parents to offspring through behaviours such as trophallaxis and coprophagy (Gurevich et al., 2020; Hosokawa et al., 2012). The second pathway involves horizontal transmission between juveniles. This may occur through active and cooperative behaviours among siblings, such as stomodeal (mouth-to-mouth) or proctodeal (mouth-to-anus) trophallaxis, that facilitate direct microbial transfer between individuals, or through passive processes involving the excretion and consumption of feces by family members (Lombardo, 2008; Onchuru et al., 2018; Powell et al., 2014). Together these vertical and horizontal pathways are important drivers of the juvenile microbiome (Onchuru et al., 2018). However, the relative contributions of each pathway during family life, and the influence of external factors on these contributions, remain largely unexplored.

Multiple external factors can directly or indirectly shape the vertical and horizontal transfers of microorganisms within families, whether by directly modifying the composition of the microbiome of donor individuals and/or the ability of parents and siblings to perform these transfers (Bright & Bulgheresi, 2010; Hosokawa & Fukatsu, 2020; Sachs et al., 2011). Among these factors, human-induced chemical pollution, particularly heavy metals, has gained increasing attention due to its widespread impact on ecosystems. Heavy metals accumulate in soil, water, and air as a result of agricultural activities, urbanization, and industrial processes (Johnson & Munshi-South, 2017; Seto et al., 2012). They are known to disrupt microbial communities across different environments (Mogren & Trumble, 2010; Sun et al., 2020; S. Zhang et al., 2015). For example, heavy metals (such as cadmium, lead or mercury) disrupt soil microbial communities by reducing bacterial richness and suppressing beneficial taxa (such as *Pseudomonas* and *Bacillus*) which are essential for plant health and nutrient cycling (Z. Cheng et al., 2023; Q. Wang et al., 2022). Similarly, in animal hosts, heavy metal exposure disrupts gut microbial composition, leading to shifts in community structure and metabolic functions. For instance, on earthworms, cadmium exposure affects key taxa (such as *Verrucomicrobia* and *Firmicutes*) which play important roles in digestion and detoxification processes (Wu et al., 2020).

Among heavy metals, cadmium is one of the most biotoxic and mobile elements in ecosystems (Kubier et al., 2019; Luo et al., 2020; Mulligan et al., 2001). Cadmium alters the behaviour, cognition, development, survival or fecundity of many organisms ranging from mammals to insects (*e.g.,* Arito et al., 1981; Gustin et al., 2018; Honorio et al., 2021). Research on cadmium's effects on animal microbiomes has expanded only recently, particularly regarding its influence on insect-associated microbial communities. Studies on honeybees (*Apis mellifera*) and pygmy grasshoppers (*Tetrix tenuicornis*) have shown that cadmium exposure alters their gut microbiomes, increasing the abundance of opportunistic pathogens such as *Aeromonas* and *Chryseobacterium*, possibly reducing host immune responses and fitness (Li et al., 2021; Rothman et al., 2019). In contrast, some gut microbes appear to confer resistance to cadmium toxicity. For instance, in mice, specific

strains of *Lactobacillus plantarum* can sequester cadmium in the intestine, acting as a protective
barrier against its toxic effects when administered as probiotics (Zhai et al., 2014, 2016). Despite
these insights, the broader implications of cadmium-induced microbiome disruptions remain poorly
understood, particularly in terms of microbial transmission between family members. If cadmium
alters the gut microbial composition of a host, it may thus disrupt both vertical and horizontal
transfer of symbiotic bacteria, potentially leading to shifts in community composition, reduced
microbial richness, and consequently major impairments in host digestion, immunity, and
development (Teyssier et al., 2018, 2020).
In this study, we investigated how the social environment and cadmium ingestion influence
the whole microbiome (encompassing gut, internal non-gut, and surface bacteria) of European
earwig juveniles (*Forficula auricularia*). The European earwig is a hemimetabolous insect, meaning
that juveniles (called nymphs) and adults share similar morphology and ecology. Nymphs undergo
five moults before reaching adulthood (Honorio et al., 2024; Tourneur et al., 2020). Mothers live
with their young nymphs in family groups that persist for about two weeks after egg hatching. During
this period, mothers frequently interact with their nymphs (Kölliker, 2007; Kölliker & Vancassel,
2007), engaging in grooming, trophallaxis and coprophagy (Falk et al., 2014), which are behaviours
typically known to facilitate microbial transfer among social insects – unlike solitary species that
primarily acquire their microbiome from a internal route or from their food environment (Onchuru et
al., 2018; Y. Wang & Rozen, 2017). However, family life is facultative in the European earwig, as
nymphs quickly become mobile after hatching, can forage for themselves, and can reach adulthood
without parents and siblings (Falk et al., 2014; Meunier & Kölliker, 2012; Van Meyel & Meunier,
2022). Recent studies indicate that the earwig microbiome is predominantly composed of
Proteobacteria and Firmicutes, with dominant bacterial families such as Lachnospiraceae,
Enterobacteriaceae or Pseudomonadaceae varying among individuals (Cheutin, Leclerc, et al.,
2024; Van Meyel et al., 2021). This microbial composition can be altered by social and chemical
stresses, including the absence of mothers during early life stages and the consumption of the
antibiotic rifampicin (Cheutin, Boucicot, et al., 2024; Greer et al., 2020; Van Meyel et al., 2021). In
contrast, cadmium exposure appears to have minimal effects on the behaviour, survival and
development of both adults and juveniles in the European earwig (Honorio, Depierrefixe, et al.,
2023; Honorio, Moreau, et al., 2023), which stands in contrast to the typical negative effects
observed in most animal species (*e.g.,* Arito et al., 1981; Gustin et al., 2018; Honorio et al., 2021).
Here, we investigated whether cadmium exposure can alter the microbiome of the European
earwig, and whether the presence of siblings during family life may mitigate these potential
alterations, thereby explaining the apparent absence of the typical cadmium-induced phenotypic
effects observed in other species.
We reared 900 earwig juveniles either isolated (alone), with siblings or in a family group
(with siblings and the mother) and exposed them to one of three concentrations of cadmium (0, 25
and 100 mg.L^{-1}) for the entire duration of family life, *i.e.* 14 days (Meunier, 2024). We then extracted
the whole microbiome from a subset of 90 nymphs (10 per combination of social environment and
cadmium concentration) and used 16S rRNA metabarcoding to analyse their alpha- and beta-
diversity, as well as identify potential treatment-specific bacterial taxa. If the frequent grooming,
trophallaxis and allo-coprophagy behaviours expressed among siblings and between mother and
nymphs mediate microbial (horizontal and vertical, respectively) transfers to juveniles, we predict
greater microbial (alpha and/or beta) diversity in nymphs living in group (sibling and family groups)
compared to isolated nymphs. Additionally, if mothers possess microbial taxa that are not
environmentally accessible to the nymphs, we expect that some microbial taxa will be specific to
nymphs reared with their mother. If cadmium alter the microbiome composition of the hosts, we
predict a change in microbial diversity in nymphs treated with cadmium. This change could be
manifested as a reduced diversity due to the elimination/selection of certain taxa, or as an
increased diversity related with a "dysbiosis" by the proliferation of non-core members (*i.e.,* Anna
Karenina principle, Zaneveld et al., 2017). Finally, if cadmium ingestion alters the social
transmission of the microbiome to the nymphs, we predict that the nymphs' microbiota will be
shaped by an interaction between both parameters: for instance, the diversity of the nymphs'

microbiome would depend on the presence or absence of siblings and mothers only in the absence
of cadmium exposure (or vice versa).

Material and methods

Laboratory rearing and experimental social environment

This study involved the offspring of 30 females collected in a peach orchard near Valence,
France (Lat 44.9772790, Long 4.9286990). All these individuals belonged to *Forficula auricularia*
Linneaus, 1758, also called *Forficula auricularia* clade A (González-Miguéns et al., 2020). Just after
collection, these females were reared under standard laboratory conditions until oviposition and
egg hatching (Meunier et al., 2012). In brief, we maintained the group of females with males for
four months to allow uncontrolled mating (Sandrin et al., 2015) and then isolated each mated
female in individual containers to allow egg production. This isolation occurred at the end of
October, approximately 50 days before the typical oviposition period for this species. From the
isolation date, each female received 50 mg of a standard laboratory-made diet (*i.e.,* mostly
composed of pollen, cat food, carrots, agar (see details in Kramer et al., (2015) without preservative
acids – not useful in this study), covered by one of three concentrations of cadmium (see details
below). The food (including cadmium) was changed twice a week until oviposition. At that time, we
removed the food as mothers typically stop their foraging activity to care for their eggs (which
typically lasts for about 50 days; Kölliker, 2007).
One day after egg hatching, we randomly assigned 30 nymphs per family (mean clutch size per
family ± SD = 51.6 ± 12.7 nymphs) to three experimental groups: family group (10 nymphs), sibling
group (10 nymphs), and isolated nymphs (10 nymphs). The family group consisted of 10 nymphs
reared with their mother, the sibling group consisted of 10 nymphs reared together without their
mother, and the isolated group consisted of single nymphs reared in isolation. This setup allowed
us to compare the microbiome of isolated nymphs with that of nymphs maintained in groups, either
with siblings only or with siblings and their mother. We then provided each experimental group with
50 mg of the standard laboratory-made diet described above, mixed with one of the three
concentrations of cadmium. Each mother and her nymphs received the same cadmium treatment
throughout the experiment. We maintained these experimental groups in Petri dishes lined with
moistened sand at 20-18°C under a 12:12h light:dark cycle. To standardise the space available per
individual, we maintained the family and sibling groups in large Petri dishes (diameter 9 cm, height
2 cm), and the isolated nymphs in small Petri dishes (diameter 3.6 cm, height 1 cm). Fourteen days
later, which marks the end of family life under natural conditions (Meunier et al., 2012), one nymph
per experimental group and cadmium concentration was transferred to an Eppendorf tube and
stored at -80°C until the extraction of the genomic material. Hence, a total of 30 nymphs were
obtained for each concentration of cadmium, comprising 10 that were raised in a family group, 10
that were raised with siblings only, and 10 that were raised in isolation. This experiment ended at
this point.

Cadmium exposure

We exposed the 30 mothers and their nymphs to cadmium using the protocol and
concentrations presented in Honorio, Moreau, et al., (2023). In particular, we applied 10 µL drops
of each cadmium solution directly onto the 50 mg of food provided to individuals. This quantity and
application method ensured that the food piece was homogeneously covered by the solution. We
used cadmium concentrations of either 0, 25, or 100 mg.L^{-1} (corresponding to 0, 5, 20 mg.kg^{-1} of
food, respectively), which were obtained by diluting cadmium chloride powder (Sigma-Aldrich
#202908, CAS number: 10108-64-2, purity: 99.99%) in MilliQ water. We chose cadmium exposure
via ingestion because this is the main source of cadmium exposure in nature (H. Zhang & Reynolds,
2019). We chose these concentrations of aqueous solutions to cover both those found in the food
chain and soil (Butt et al., 2018; FitzGerald & Roth, 2015; H. Zhang & Reynolds, 2019) and those

known to induce behavioural and microbial changes after ingestion in other species (Dési et al.,
1998; Rothman et al., 2019; Su et al., 2021). A previous study confirmed that this method of
exposure ensures the uptake of cadmium by earwig females (Honorio et al, *submitted*; Honorio,
Moreau, et al., 2023).

Genomic extraction and 16S amplification

The 90 nymphs were individually isolated in 2mL Eppendorf tube and immediately stored at -
80°C until the extraction of both external and internal DNA material (*i.e.,* whole individual body)
using the NucleoMag® Tissue extraction kit (Macherey-Nagel™, Düren, Germany). We then
amplified the hypervariable region V3-V4 of the 16S rRNA gene using the primers 343F (5′ -
ACGGRAGGCAGCAG - 3′) and 784R (5′- TACCAGGGTATCTAATC – 3′) (Muyzer et al., 1993)
coupled with platform-specific Illumina linkers. PCR reactions were processed using the Taq
Polymerase Phusion® High-Fidelity PCR Master Mix with GC buffer following the manufacturer's
instructions (Qiagen, Hilder, Germany). Each PCR reaction involved an initial step of 30 sec
denaturation at 98°C, followed by 22 amplification cycles (10 sec denaturation at 98°C; 30 sec
annealing at 61.5°C; 30 sec extension at 72°C), and ended with a final step of 10 min extension at
72°C. Each reaction was checked for the probe specificity and amplification success with an
electrophoresis migration. Extraction and amplification steps involved several blank controls to
confirm that samples were not contaminated by environmental microorganisms. As blanks were all
negative, we did not send them for sequencing (Figure S1). Samples were amplified in duplicate
and equally pooled for a final product of 30 µL which was further sequenced using 2x250bp Illumina
MiSeq technology at the Bio-Environnement platform (University of Perpignan, France). Finally,
two of the 90 nymphs used for genomic extraction were removed from the study as one did not
amplify (isolated nymphs) and one did not respect the experimental condition (the sibling group
was only constituted by two nymphs instead of 10). Both samples were exposed to 100 mg.L^{-1} Cd.

Bioinformatic process

We used the DADA2 algorithm v1.24.0 (Callahan et al., 2016) in R software v4.2.0 (R Core
Team, 2024) to process sequences. Libraries were first trimmed for primers and reads were
truncated at 240 base pairs (both forward and reverse), filtered with their quality profiles, cleaned,
dereplicated and inferred into Amplicon Sequence Variants (ASVs) (Glassman & Martiny, 2018).
The resulting chimeras were removed and taxonomic assignment using the SILVA reference
database (release 138) (Quast et al., 2012). Sequences were aligned using the MAFFT program
(Katoh, 2002) and entered into a phyloseq object using the *phyloseq* package v1.40.0 (McMurdie
& Holmes, 2013). We inferred the phylogenetic tree associated with these sequences using the
FastTree 2 tool (Price et al., 2010) and the *Phangorn* package v2.8.1 (Schliep, 2011). We removed
2 211 sequences that belonged to 65 ASVs that were assigned as mitochondrial sequences or not
assigned at least at the Phylum level. We also filtered the sequence using a standard method that
(1) preserves rare sequences while filtering out statistically random sequences that may introduce
noise from exaggerated inter-individual variability (*i.e.,* unique sequences present in only few
individuals), and (2) avoids relying on subjective or arbitrary occurrence and abundance thresholds
(Magurran & Henderson, 2003). Specifically, we calculated an index of dispersion (*i.e.,* the variance
to mean ratio, VMR) for each ASV. We then tested whether these indices followed a Poisson
distribution, checking if they fall within the 2.5 and 97.5% confidence limits of the Chi^2 distribution
(Krebs, 1999) and we preserved the ASVs with index values outside these confidence limits (Figure
S2). This removal represents 1.5% of the sequences in the initial dataset, for an occurrence that
did not exceed three individuals out of 90.

Microbial diversity and indicators

Microbial diversity and indicators

We tested the effects of social environment and cadmium ingestion on the alpha- and beta-
diversity of the resulting microbiome of each host nymph. To calculate alpha-diversity, we first
rarefied all samples to 11 637 sequences corresponding to the threshold at which rarefaction
curves reach their stationary phases (Cameron et al., 2021; Figure S3). The rarefaction involved
the removal of one sample of family nymphs (100 mg.L^{-1} of Cd), which contained fewer reads than
the threshold. We then calculated the observed richness (*i.e.,* ASV or bacterial genus counts), the
Shannon entropy that integrates the abundance of each ASV (or genus) present in each samples,
and their respective equivalents the Faith and Allen indices that integrate the phylogenetic
relationships between ASVs (or genus). For beta-diversity, we calculated the Jaccard distance
between samples with ASVs (or genus) presence/absence and its quantitative equivalent Bray-
Curtis. As for alpha-diversity, we also used their phylogenetic equivalents with the Unifrac metrics
for qualitative (unweighted) and quantitative (weighted) distances (Chao et al., 2010; Yang et al.,
2021). The dissimilarities between samples were projected into a Principal Coordinates Analysis
(PCoA), which we plotted on the first two axes. The significant microbial genera correlated along
these axes were projected through the PCoA using the function *envfit()* from the package *vegan*
v2.6-4 (Oksanen et al., 2022). Alpha- and beta-diversity were also calculated at genus level by
merging all ASVs in their belonging genus using the function *tax_glom()* from the *phyloseq* package
(McMurdie & Holmes, 2013).

Statistics

All statistical analyses were performed using R v4.3.4 (R Core Team, 2024). First, we analysed
the alpha-diversity of nymph microbiome using four linear mixed models (LMER) with the *lme4*
v1.1-31 (Bates et al., 2015) and analyses of variances (Anova) were run using *car* version
packages v3.1-1 (Fox & Weisberg, 2019). In these models, we entered either the observed
richness, Shannon entropy, Faith or Allen indices as a response variable, and social environment
(isolated, sibling or family nymphs) and cadmium concentration (discrete; 0, 25 or 100 mg.L^{-1}) as
explanatory variables. We also included the interaction between these two factors to test whether
cadmium ingestion affects the social transmission, or wether being in a specific social context
modulates the effect of the cadmium ingestion on the microbiome. To control for the same origin
of each social environment, we also entered the mother identity as a random effect in each model.
When necessary, we conducted post-hoc pairwise comparisons using the estimated marginal
means via the *emmeans* package v1.8.0 (Lenth, 2022). The R^2 corresponding to each pairwise
comparison was calculated with the *MuMIn* package v1.47.1 (Bartoń, 2022). Second, we analysed
the beta-diversity of nymph microbiome using four permutational analyses of variances
(PERMANOVAs). In these models, we entered either Jaccard, Bray-Curtis, unweighted and
weighted Unifrac metrics as response variables, and social environment, cadmium concentration
and their interaction as explanatory variables. We specified the origin of the mother as strata to
control for the effects of this variable. We conducted post-hoc pairwise comparisons using the
pairwiseAdonis package v0.4.1 (Martinez Arbizu, 2017). Overall, we ran these eight models on
diversity indices calculated on both ASVs and Genus levels. All models were simplified after AICc
comparisons by removing the non-significant interactions. Finally, to identify bacterial indicators of
both the social environment and the cadmium concentration, we used the *DESeq2* package v1.36-
0 (Love et al., 2014). Each raw counts were clustered into bacterial genera, normalized by its
geometric mean, and then compared to a negative binomial distribution. The estimated coefficient
(Log2FoldChange) for each genus in each sample group was transformed into a Z-statistic,
compared to a normal distribution, and then corrected for multiple comparisons.

We checked all model assumptions using the *DHARMa* package (Hartig, 2022) and the
homoscedasticity of the variances between groups of each variable with the function *betadisper*
from the *vegan* package v2.6-4 (Oksanen et al., 2022). To fulfil model assumptions, we log-
transformed alpha proxies (*i.e.,* Observed richness, Shannon, Faith and Allen indices) as response

variable. All p-values were adjusted for multiple testing with False Discovery Rate method (FDR)
and Benjamini-Hochberg correction (Haynes, 2013).

Results

Overall microbial composition of the nymphs

Overall, the microbiome of the tested nymphs included 2 075 719 sequences (98.52%
abundance of the initial dataset) representing 590 different ASVs (31.45% richness of the initial
dataset; Figure S2). While Proteobacteria largely dominated the microbiome (78.7%), the
Bacteroidota represented 15.2%, and the Firmicutes 5.43%. Finally, the microbiome also contained
other bacterial phyla at a low level, such as Actinobacteriota, Bdellovibrionota and Patescibacteria
(all < 1 %). The dominant genera (Figure 1) were *Buttiauxella* (Proteobacterales,
Enterobacteriacae) (32.1 % of all reads), *Chryseobacterium* (Bacteroidota, Weeksellaceae)
(11.5%), *Acinetobacter* (Proteobacteria, Moraxellaceae) (6.33%), *Serratia* (Proteobacteria,
Yersiniaceae) (6.31%) and *Pseudomonas* (Proteobacteria, Pseudomonadaceae) (5.34%).

**Figure 1 - Microbiome composition of nymphs from isolated, sibling and
family treatments.** Barplots represent the individual composition for the top 20
microbial genera. Taxa belonging to Proteobacteria are in blue, Firmicutes in
red/purple, and Bacteroidota in green. The gray contrasted horizontal bar
represent the cadmium concentration for exposed nymphs. N indicates the
number of nymphs used per social environment.

Effects of social environment and cadmium on microbial diversity

Social environment and cadmium ingestion had independent effects on the unweighted Unifrac metric (Figure 2B), but no effect on alpha diversity (Figure 2A, Figure S4) neither the other beta diversity metrics (Figure S4). The social environment affected the unweighted Unifrac dissimilarities at both ASV level (R^2 = 0.063, F = 2.90, P_{adj} = 0.016; Figure S4) and Genus level (R^2 = 0.070, F = 3.28, P_{adj} = 0.016). This effect showed that nymphs exhibited microbiome structures that were specific of their social environment (Table 1, Figure 2B): isolated vs family treatments (ASV R^2 = 0.066, F = 3.10, P_{adj} = 0.001; Genus R^2 = 0.075, F = 4.61, P_{adj} = 0.001), isolated vs sibling treatments (ASV R^2 = 0.039, F = 2.25, P_{adj} = 0.018; Genus R^2 = 0.047, F = 2.73, P_{adj} = 0.022), and family vs sibling treatments (ASV R^2 = 0.038, F = 2.25, P_{adj} = 0.014; Genus R^2 = 0.038, F = 2.23, P_{adj} = 0.029). These results were consistent with the first two axes of the PCoA associated with the unweighted Unifrac distances where microbial structures of isolated nymphs were seemingly projected in the opposite direction of other samples along the first axis (Figure 2C). Microbial taxa such as bacteria belonging to the Phylum Bdellovibrionota and bacterial genera *Niabella* (Bacteroidota, Chitinophagaceae), *Rurimicrobium* (Bacteroidota, Chitinophagaceae), *Pedobacter* (Bacteroidota, Sphingobacteriaceae) or *Alkanindiges* (Proteobacteria, Moraxellaceae) were significantly correlated along the PCoA axes (P < 0.01) and tended to project in the direction of isolated nymph's microbiomes along the first PCoA axis (Figure 2B).

Figure 2 – Phylogenetic alpha and beta-diversity of the nymph microbiomes.
Differences in alpha-diversity where **(A)** the Faith (phylogenetic diversity) and **(B)** the unweighted Unifrac distances of the ASVs composition between samples were tested according to the social environment and the cadmium concentration. Boxplots show the median (middle bar) and the interquartile range (box) with whiskers representing the 1.5-fold the interquartile range. Non-significant results (P_{adj} > 0.05) are labelled by 'ns'. Dissimilarities are visualized by the first two axes of a Principal Coordinates Analyses (PCoA), where each nymph is coloured according to their social environment and shaped by the cadmium concentration. The microbial genera and phyla that are significantly correlated (P < 0.01) are projected in black and red, respectively.

Additionally, cadmium ingestion marginally affected unweighted Unifrac dissimilarities at ASV level ($R^2 = 0.041$, F = 1.88, $P_{adj} = 0.053$). Nymphs exposed to 100 mg.L^{-1} cadmium had a different microbial structure than nymphs exposed to 0 mg.L^{-1} ($R^2 = 0.035$, F = 2.04, $P_{adj} = 0.023$) and 25 mg.L^{-1} ($R^2 = 0.038$, F = 2.21, $P_{adj} = 0.018$), whereas there was no difference between 0 mg.L^{-1} and 25 mg.L^{-1} ($R^2 = 0.022$, F = 1.29, $P_{adj} = 0.188$). Note, however, that these last results must be taken with caution as dispersion (*i.e.,* homogeneity) of microbiomes regarding the unweighted Unifrac distances differed between the three cadmium concentrations with highest distance from the centroid in the concentration 25 mg.L^{-1} ($P_{adj} = 0.026$). Finally, we did not detect any effect of the interaction between social environment and cadmium ingestion, either for alpha diversity (all $P_{adj} \geq 0.123$) or beta diversity (all $P_{adj} \geq 0.392$; Table 1).

Table 1 - Effects of social environment, cadmium ingestion, and their interaction on the alpha-diversity and beta-diversity of nymph microbiomes at both ASV and Genus levels. Statistical values from linear models with random effects (LMERs) or PERMANOVAs conducted on each index. P-values were corrected for multiple testing according to False Discovery Rate (FDR) correction. The models were simplified by removing the interaction as it was not significant for both alpha (all $P_{adj} \geq 0.136$) and beta diversity (all $P_{adj} \geq 0.169$). Significant results ($P_{adj} \leq 0.05$) are indicated in bold.

Index	Social environment			Cadmium ingestion		
Alpha-diversity (LMERs)						
	LR X^2_2	P	P_{adj}	LR X^2_2	P	P_{adj}
- ASV level						
Observed richness	0.587	0.746	0.892	1.231	0.540	0.720
Shannon	0.141	0.932	0.962	0.371	0.831	0.892
Faith	7.581	0.023	0.123	2.129	0.345	0.552
Allen	2.609	0.271	0.551	0.358	0.836	0.892
- Genus level						
Observed richness	4.054	0.132	0.453	3.550	0.170	0.453
Shannon	0.370	0.831	0.892	0.075	0.963	0.963
Faith	**9.268**	**0.010**	**0.080**	2.243	0.326	0.551
Allen	2.956	0.228	0.551	0.653	0.721	0.887
Beta-diversity (PERMANOVAs)						
	R^2	P	P_{adj}	R^2	P	P_{adj}
- ASV level						
Jaccard	0.026	0.161	0.453	0.027	0.389	0.556
Bray-Curtis	0.029	0.156	0.453	0.031	0.363	0.553
Unweighted unifrac	**0.063**	**0.001**	**0.016**	**0.041**	**0.005**	**0.053**
Weighted unifrac	0.026	0.285	0.551	0.015	0.572	0.732
- Genus level						
Jaccard	0.032	0.093	0.392	0.025	0.290	0.551
Bray-Curtis	0.035	0.098	0.392	0.027	0.304	0.551
Unweighted unifrac	**0.070**	**0.001**	**0.016**	0.040	0.019	0.122
Weighted unifrac	0.026	0.327	0.551	0.012	0.507	0.705

Effect of social environment and cadmium on bacterial genera

Out of a total of 84 bacterial genera that formed the nymph's microbiome, 18 and 7 were statistical indicators of the social environment and the cadmium concentration, respectively. Of these, 13 and 5 bacterial genera showed a |log2fold change| > 2 (*i.e.,* for which the abundance is at least 4 times different), respectively. Among these bacteria, *Janthinobacterium* (Proteobacteria, Oxalopacteraceae), *Kosakonia* (Proteobacteria, Enterobacteriaceae), *Pedobacter* (Bacteroidota, Sphingobacteriaceae), *Rurimicrobium* (Bacteroidota, Chitinophagaceae) were more abundant in isolated nymphs than in sibling or family nymphs (Figure 3; Table S1). In contrast, the bacterial genera *Duganella* (Proteobacteria, Oxalobacteraceae), *Cellvibrio* (Proteobacteria, Cellvibrionaceae) and *Serratia* (Proteobacteria, Yersiniaceae) dominated the sibling nymphs. *Rhodococcus* (Actinobacteria, Nocardiaceae), *Lactobacillus* (Firmicutes, Lactobacillaceae), *Paracoccus* (Proteobacteria, Rhodobacteraceae), *Tsukamurella* (Actinobacteria, Tsukamurellaceae), *Erwinia* (Proteobateria, Erwiniaceae) and *Sphingobium* (Proteobacteria, Sphingomonadaceae) were the most abundant bacteria in family nymphs (Figure 3; Table S1).

Regarding cadmium ingestion, we detected a striking increase in *Duganella* (Proteobacteria, Oxalobacteraceae), *Hafnia-Obesumbacterium* (Proteobacteria, Hafniaceae), *Erwinia* (Proteobateria, Erwiniaceae), *Nubsella*, *Pedobacter* (Bacteroidota, Sphingobacteriaceae) and *Rurimicrobium* (Bacteroidota, Chitinophagaceae) in nymphs exposed to 25 mg.L^{-1} of Cd compared to nymphs exposed to 0 or 100 mg.L^{-1} of Cd (Figure 4; Table S1). None of these bacteria were represented as discriminant of both parameters (*i.e.,* social environment and cadmium ingestion), suggesting that the effect of the social environment does not influence (or is not influenced) by the cadmium ingestion.

Figure 3 - Abundance of each bacterial genus indicators related to isolated, sibling or family nymphs. Indicators are delineated using a DESeq2 approach (P < 0.05) for which the significant contrast showed a |log2fold change| > 2. The prevalence for each genus is labelled in the bar of the corresponding social environment. Different letters indicate P < 0.05.

Discussion

Understanding the factors that drive changes in communities of mutualistic and commensal
microorganisms in animal hosts is crucial for enhancing our knowledge of host fitness, population
dynamics and ecosystem functioning (Martiny et al., 2023). This is particularly true in the context
of anthropogenic environmental changes, which are increasingly known to both directly and
indirectly shape host-symbiont relationships (Cheutin et al., 2021; Maraci et al., 2022; Teyssier et
al., 2018). In this study, we tested the simultaneous effects of the social environment and exposure
to a chemical pollutant on the diversity and nature of the microbial communities present in earwig
juveniles. Our results first show that the social environment affected the phylogenetic diversity of
the microbial communities (*i.e.,* unweighted Unifrac distances) at both the bacterial sequence (*i.e.,*
ASV) and genus levels. Our results then show that cadmium ingestion influenced the phylogenetic
qualitative composition of the microbiome at the ASV level. Notably, we found that changes in the
microbiome induced by the social environment and cadmium ingestion were associated with the
turnover of specific bacterial taxa. However, there was no effect of the social environment and
cadmium ingestion on the other indices of alpha-diversity and beta-diversity of the nymph
microbiome. Finally, as we found no interaction between cadmium and social environment, our
results suggest that cadmium ingestion did not affect the social transmission of the microbiome to
earwig nymphs.
Our first finding is that nymphs exhibited social environment-dependent microbial communities
according to the unweighted Unifrac metric. This indicates that nymphs reared alone, with siblings,
or with a full family group (siblings plus mother) harbour distinct microbiome profiles, each primarily
shaped by the rare or low abundance of bacteria from phylogenetically distant taxa. These results
provide two key insights. First, the presence of specific taxa in the isolated nymphs suggest that
living in a group may not only enhance access to new bacterial sources, as reported in other social
mammals and insects (Hosokawa et al., 2007; Raulo et al., 2021; Tung et al., 2015; Y. Wang &
Rozen, 2017), but also help to prevent colonization by opportunistic environmental taxa. This is
examplified by the phylum Bdellovibrionota (including the two genera *Peredibacter* and
Bdellovibiro), which was more abundant in isolated nymphs compared to those reared with siblings
or within a full family group (Figure 2B). This phylum is typically abundant in the soil and is
occasionally found in the gut microbiome of certain insects (Davidov et al., 2006). The proliferation
of Bdellovibrionota in isolated nymphs might thus stem from their initial acquisition from the
environment, combined with the absence of competitive bacterial strains that are typically
transmitted by family members (Figueiredo & Kramer, 2020). Further testing is needed to validate
this hypothesis. The second key finding is that the composition of the group plays an important role
in shaping the nymph's microbiome. Some bacterial taxa were more abundant in nymphs reared
in a whole-family group than in nymphs reared with their siblings only. This suggests a maternal
transmission of bacteria to nymphs during earwig family life. Interestingly, many of the bacteria

putatively transmitted by earwig mothers have known beneficial functions related to digestion and
immunity in other (social) insects. For example, the Proteobacteria *Sphingobium* or *Rhodococcus*
present a large variety of catabolic functions such as aromatic hydrocarbons metabolization or
cellulose digestion in cockroaches, termites or beetles (Biswas & Paul, 2021; Delalibera et al.,
2005; Wenzel et al., 2002). Similarly, the bacterium *Lactobacillus* can protect against the negative
effects of pathogens, pesticides and cadmium, as well as help for the fermentation of carbohydrate
plants and cereals in bees, honeybees or grasshoppers (Lazzeri et al., 2020; Li et al., 2021; Nowak
et al., 2021; Vásquez et al., 2012). To date, the role of the gut microbiome in the fitness of the
European earwig remains unclear (Cheutin, Leclerc, et al., 2024; Van Meyel et al., 2021), and
future experimental and transcriptomic studies are needed to investigate the potential function of
these bacteria in this species.

We found that the social environment affected the nymph microbiome not only through
differences in rare/low-abundant phylogenetically distant taxa, but also via an over-representation
of 13 bacterial taxa (4 specific to the isolated treatment, 3 to the sibling treatment, and 7 to the full
family treatment). In isolated nymphs, two of these four taxa (*Kosakonia* and *Janthinobacterium)*
are insect mutualists and typically enhance host protection against pathogens. For instance,
Janthinobacterium has antifungal functions in many insects (Haack et al., 2016; Magoga et al.,
2023; Rodrigues et al., 2006), while *Kosakonia* can inhibit the growth of the entomopathogenic
strains belonging to the bacteria *Serratia* (Grimont et al., 1979; Muratoğlu et al., 2011) and render
mosquitoes and tsetse flies hosts refractory to parasite infections such as *Trypanosoma* and
Plasmodium (Weiss et al., 2019). The overabundance of these bacteria in isolated nymphs might
be an adaptive strategy, enabling them to mount an effective defense against pathogens in the
absence of social immunity provided by siblings and mothers (Diehl et al., 2015; Diehl & Meunier,
2018). This hypothesis would also be consistent with our results showing that *Serratia* (an
opportunistic entomopathogen) is less abundant in isolated than group-living nymphs (see above).

Independent of the social environment, we found that cadmium exposure also played an
important role in shaping the diversity of the nymph microbiome. First, cadmium ingestion induced
unstable community states consistent with the Anna Karenina principle (Zaneveld et al., 2017),
whereby microbiomes became more heterogeneous and compositionally distinct in nymphs
exposed to the highest (100mg.L^{-1}) compared to intermediate (25 mg.L^{-1}) or control (0 mg.L^{-1})
doses, as revealed by both the Faith and unweighted Unifrac metrics. This effect is in line with the
typical effects of cadmium on microbial communities reported in a taxonomically wide range of
animals (Li et al., 2021; Rothman et al., 2019; Trevors et al., 1986; S. Zhang et al., 2015; Y. Zhang
et al., 2016). Second, cadmium ingestion at the intermediate concentration was associated with
the presence or absence of five bacterial taxa. Of these, only one (*Hafnia-Obesumbacterium*) has
been reported in animal microbiomes (S. Zhang et al., 2023). This bacterium is a pathogen in many
species, where its abundance often increases under stressful conditions (Janda & Abbott, 2006).
For example, *Hafnia-Obesumbacterium* is dominant in the gut microbiomes of tadpole *Rana*
chensinensis and crayfish *Procambarus clarkii*, respectively exposed to lead pollution (S. Zhang et
al., 2023) and acidic conditions (Z. Wang et al., 2024). As such, our data suggest that this genus
is the only bacterium sensitive to cadmium ingestion in juveniles of the European earwig. The other
four bacterial taxa specific to the intermediate dose of cadmium probably reflect the good
development of these bacteria on medium contaminated with low levels of cadmium and their
subsequent passive environmental uptake by earwig nymphs via contact with cadmium-
contaminated food. This is consistent with previous studies indicating that the growth of *Pedobacter*
and *Duganella* is typically favoured in cadmium-enriched soils (X. Cheng et al., 2022; Stout et al.,
2010; H. Wang et al., 2023). However, while our findings align with the existing literature, they must
be considered with caution, as they are based on metabarcoding approach (Djemiel et al., 2022).
Given the qualitative nature of our results, they may be biased due to their compositional
framework. Quantitative and transcriptomic studies are now needed to validate these findings and
further elucidate the functional role of the microbiome under environmental stress (Gloor et al.,
2017).

Overall, this study shows that the access to a social environment, the composition of family group and exposure to chemical pollutants can all influence the composition and diversity of the microbiome in juvenile European earwigs. Interestingly, we show that these effects do not interact with each other, *i.e.* varying the access to a social environment or changing the composition of that nymph microbiome, and vice versa. As anthropogenic environmental changes continue to accelerate globally (Johnson & Munshi-South, 2017; Seto et al., 2012), these results demonstrate that the ability to naturally switch between solitary and social life (as found in the European earwig nymphs, Meunier, 2024) does not necessarily confer an advantage in mitigating the negative effects of chemical contaminants (at least) on the host microbiome. Future studies are now needed to better understand the role of the earwig microbiome on its fitness and population dynamics (Cheutin, Leclerc, et al., 2024; Van Meyel et al., 2021), and to determine whether the reported effects of social environment and cadmium intake on its diversity can alter these parameters.

Data availability

The data set and the scripts are available on Zenodo with the url https://zenodo.org/doi/10.5281/zenodo.13380552. The reads for each sample were deposited in the NCBI Sequence Read Archive (SRA) under the BioProject accession no. PRJNA1152783.

Acknowledgements

We thank Charlotte Lécureuil for her comments on this manuscript. We also thank all members of the EARWIG group at IRBI for their help with animal rearing and setup installation, and the Bio-environment platform of the University of Perpignan (France) for providing the MiSeq Illumina. We thank Armand Guillermin and Stéphanie Drusch of the INRAE unité expérimentale Recherche Intégrée in Gotheron for giving us access to their orchards for earwig field sampling. The graphical abstract was made with BioRender application. Finally, we also thank two anonymous reviewers for their insightful comments in a previous version of this manuscript.

Author's contribution

Marie-Charlotte Cheutin: Data curation; Formal analysis; Investigation; Methodology; Writing - original draft; and Writing - review & editing. – **Romain Honorio**: Conceptualization; Methodology; Writing - original draft; and Writing - review & editing. **Joël Meunier**: Conceptualization; Funding acquisition; Supervision; Writing - review & editing

Funding information

This work was funded by the Région Centre-Val de Loire (Project DisruptCare; n°2021 00149474) and by a research grant from the Agence Nationale de la Recherche (ANR-20-CE02-0002).

Conflict of interest

The authors declare that this study had no conflicts of interest.

Ethics approval statements

Our investigation complies with the current European Directive 2010/63/EU that does not acquire ethical approval on invertebrates. All animals were handled with care until necessary sacrifices.

Reference

Arito, H., Sudo, A., & Suzuki, Y. (1981). Aggressive behavior of the rat induced by repeated administration of cadmium. *Toxicology Letters*, *7*(6), 457–461. https://doi.org/10.1016/0378-4274(81)90093-X

Bartoń, K. (2022). *MuMIn: Multi-Model Inference*. https://CRAN.R-project.org/package=MuMIn

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/jss.v067.i01

Berlanga, M., & Guerrero, R. (2016). The holobiont concept: The case of xylophagous termites and cockroaches. *Symbiosis*, *68*(1–3), 49–60. https://doi.org/10.1007/s13199-016-0388-9

Biswas, S., & Paul, D. (2021). Isolation and characterization of *Rhodococcus qingshengii*, a cellulolytic bacteria from *Cnaphalocrocis medinalis* (Lepidoptera: Crambidae) gut. *Environmental and Experimental Biology*, *19*(4). https://doi.org/10.22364/eeb.19.19

Bright, M., & Bulgheresi, S. (2010). A complex journey: Transmission of microbial symbionts. *Nature Reviews Microbiology*, *8*(3), 218–230. https://doi.org/10.1038/nrmicro2262

Butt, A., Qurat-ul-Ain, Rehman, K., Khan, M. X., & Hesselberg, T. (2018). Bioaccumulation of cadmium, lead, and zinc in agriculture-based insect food chains. *Environmental Monitoring and Assessment*, *190*(12), 698. https://doi.org/10.1007/s10661-018-7051-2

Calderón-Cortés, N., Quesada, M., Watanabe, H., Cano-Camacho, H., & Oyama, K. (2012). Endogenous plant cell wall digestion: A key mechanism in insect evolution. *Annual Review of Ecology, Evolution, and Systematics*, *43*(1), 45–71. https://doi.org/10.1146/annurev-ecolsys-110411-160312

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*, *13*(7), 581–583. https://doi.org/10.1038/nmeth.3869

Cameron, E. S., Schmidt, P. J., Tremblay, B. J.-M., Emelko, M. B., & Müller, K. M. (2021). Enhancing diversity analysis by repeatedly rarefying next generation sequencing data describing microbial communities. *Scientific Reports*, *11*(1), 22302. https://doi.org/10.1038/s41598-021-01636-1

Chao, A., Chiu, C.-H., & Jost, L. (2010). Phylogenetic diversity measures based on Hill numbers. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*(1558), 3599–3609. https://doi.org/10.1098/rstb.2010.0272

Cheng, X., Cao, X., Tan, C., Liu, L., Bai, J., Liang, Y., & Cai, R. (2022). Effects of four endophytic bacteria on cadmium speciation and remediation efficiency of *Sedum plumbizincicola* in farmland soil. *Environmental Science and Pollution Research*, *29*(59), 89557–89569. https://doi.org/10.1007/s11356-022-21711-y

Cheng, Z., Shi, J., He, Y., Chen, Y., Wang, Y., Yang, X., Wang, T., Wu, L., & Xu, J. (2023). Enhanced soil function and health by soybean root microbial communities during *in situ* remediation of Cd-contaminated soil with the application of soil amendments. *mSystems*, *8*(3), e01049-22. https://doi.org/10.1128/msystems.01049-22

Cheutin, M.-C., Boucicot, M., & Meunier, J. (2024). Microbiome turnover during offspring development varies with maternal care, but not moult, in a hemimetabolous insect. *Peer Community Journal*, *4*, e67. https://doi.org/10.24072/pcjournal.446

Cheutin, M.-C., Leclerc, B., & Meunier, J. (2024). With or without you: Gut microbiota does not predict aggregation behavior in European earwig females. *Behavioral Ecology*, *35*(3), arae022. https://doi.org/10.1093/beheco/arae022

Cheutin, M.-C., Villéger, S., Hicks, C. C., Robinson, J. P. W., Graham, N. A. J., Marconnet, C., Restrepo, C. X. O., Bettarel, Y., Bouvier, T., & Auguet, J.-C. (2021). Microbial shift in the enteric bacteriome of coral reef fish following climate-driven regime shifts. *Microorganisms*, *9*(8), 1711. https://doi.org/10.3390/microorganisms9081711

Davidov, Y., Friedjung, A., & Jurkevitch, E. (2006). Structure analysis of a soil community of predatory bacteria using culture-dependent and culture-independent methods reveals a hitherto undetected diversity of *Bdellovibrio* -and-like organisms. *Environmental Microbiology*, *8*(9), 1667–1673. https://doi.org/10.1111/j.1462-2920.2006.01052.x

Delalibera, I., Handelsman, J., & Raffa, K. F. (2005). Contrasts in cellulolytic activities of gut microorganisms between the wood borer, *Saperda vestita* (Coleoptera: Cerambycidae), and the bark beetles, *Ips pini* and *Dendroctonus frontalis* (Coleoptera: Curculionidae). *Environmental Entomology*, *34*(3), 541–547. https://doi.org/10.1603/0046-225X-34.3.541

Dési, I., Nagymajtényi, L., & Schulz, H. (1998). Behavioural and neurotoxicological changes caused by cadmium treatment of rats during development. *Journal of Applied Toxicology*, *18*(1), 63–70. https://doi.org/10.1002/(SICI)1099-1263(199801/02)18:1<63::AID-JAT475>3.0.CO;2-Z

Diehl, J. M. C., Körner, M., Pietsch, M., & Meunier, J. (2015). Feces production as a form of social immunity in an insect with facultative maternal care. *BMC Evolutionary Biology*, *15*(1), 40. https://doi.org/10.1186/s12862-015-0330-4

Diehl, J. M. C., & Meunier, J. (2018). Surrounding pathogens shape maternal egg care but not egg production in the European earwig. *Behavioral Ecology*, *29*(1), 128–136. https://doi.org/10.1093/beheco/arx140

Djemiel, C., Maron, P.-A., Terrat, S., Dequiedt, S., Cottin, A., & Ranjard, L. (2022). Inferring microbiota functions from taxonomic genes: A review. *GigaScience*, *11*, giab090. https://doi.org/10.1093/gigascience/giab090

Falk, J., Wong, J. W. Y., Kölliker, M., & Meunier, J. (2014). Sibling cooperation in earwig families provides insights into the early evolution of social life. *The American Naturalist*, *183*(4), 547–557. https://doi.org/10.1086/675364

Figueiredo, A. R. T., & Kramer, J. (2020). Cooperation and conflict within the microbiota and their effects on animal hosts. *Frontiers in Ecology and Evolution*, *8*, 132. https://doi.org/10.3389/fevo.2020.00132

FitzGerald, R. E., & Roth, N. (2015). Cadmium in mineral fertilizers – human and environmental risk update. https://doi.org/10.13140/RG.2.2.26451.17442

Fox, J., & Weisberg, S. (2019). An R companion to applied regression (Third). Sage. https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Girard, M., Luis, P., Valiente Moro, C., & Minard, G. (2023). Crosstalk between the microbiota and insect postembryonic development. *Trends in Microbiology*, *31*(2), 181–196. https://doi.org/10.1016/j.tim.2022.08.013

Glassman, S. I., & Martiny, J. B. H. (2018). Broadscale ecological patterns are robust to use of Exact Sequence Variants versus Operational Taxonomic Units. *mSphere*, *3*(4), e00148-18. https://doi.org/10.1128/mSphere.00148-18

Gloor, G. B., Macklaim, J. M., Pawlowsky-Glahn, V., & Egozcue, J. J. (2017). Microbiome datasets are compositional: And this is not optional. *Frontiers in Microbiology*, *8*, 2224. https://doi.org/10.3389/fmicb.2017.02224

González-Miguéns, R., Muñoz-Nozal, E., Jiménez-Ruiz, Y., Mas-Peinado, P., Ghanavi, H. R., & García-París, M. (2020). Speciation patterns in the *Forficula auricularia* species complex: Cryptic and not so cryptic taxa across the western Palaearctic region. *Zoological Journal of the Linnean Society*, *190*(3), 788–823. https://doi.org/10.1093/zoolinnean/zlaa070

Greer, J. A., Swei, A., Vredenburg, V. T., & Zink, A. G. (2020). Parental care alters the egg microbiome of maritime earwigs. *Microbial Ecology*, *80*(4), 920–934. https://doi.org/10.1007/s00248-020-01558-x

Grimont, P. A. D., Grimont, F., & Lysenko, O. (1979). Species and biotype identification of *Serratia* strains associated with insects. *Current Microbiology*, *2*(3), 139–142. https://doi.org/10.1007/BF02605870

Gurevich, Y., Lewin-Epstein, O., & Hadany, L. (2020). The evolution of paternal care: A role for microbes? *Philosophical Transactions of the Royal Society B: Biological Sciences*, *375*(1808), 20190599. https://doi.org/10.1098/rstb.2019.0599

Gustin, K., Tofail, F., Vahter, M., & Kippler, M. (2018). Cadmium exposure and cognitive abilities and behavior at 10 years of age: A prospective cohort study. *Environment International*, *113*, 259–268. https://doi.org/10.1016/j.envint.2018.02.020

Haack, F. S., Poehlein, A., Kröger, C., Voigt, C. A., Piepenbring, M., Bode, H. B., Daniel, R., Schäfer, W., & Streit, W. R. (2016). Molecular keys to the *Janthinobacterium* and *Duganella* spp. Interaction with the plant pathogen *Fusarium graminearum*. *Frontiers in Microbiology*, *7*. https://doi.org/10.3389/fmicb.2016.01668

Hartig, F. (2022). DHARMa: Residual diagnostics for hierarchical (multi-Level / mixed) regression models. *R Package Version 0.4.6*. https://CRAN.R-project.org/package=DHARMa

Haynes, W. (2013). Benjamini–Hochberg Method. In W. Dubitzky, O. Wolkenhauer, K.-H. Cho, & H. Yokota (Eds.), *Encyclopedia of Systems Biology* (pp. 78–78). Springer New York. https://doi.org/10.1007/978-1-4419-9863-7_1215

Honorio, R., Cheutin, M. –C., Pasquier, L., De Wever, S., Perdereau, E., Villalta, I., Lécureuil, C., & Meunier, J. (2024). The European earwig: A model species for studying the (early) evolution of social life. *Insectes Sociaux*. https://doi.org/10.1007/s00040-024-00985-0

Honorio, R., Depierrefixe, P., Devers, S., Rouelle, M., Meunier, J., & Lécureuil, C. (2023). Effects of cadmium ingestion on reproduction and maternal egg care in the European earwig. *Animal Behaviour*, *195*, 1–8. https://doi.org/10.1016/j.anbehav.2022.10.008

Honorio, R., Jacquier, L., Doums, C., & Molet, M. (2021). Disentangling the roles of social and individual effects on cadmium tolerance in the ant *Temnothorax nylanderi*. *Biological Journal of the Linnean Society*, *134*(4), 823–834. https://doi.org/10.1093/biolinnean/blab116

Honorio, R., Moreau, S., Lécureuil, C., & Meunier, J. (2023). Cadmium does not affect post-hatching maternal care or early offspring development in earwigs. *Behavioral Ecology*, *34*(3), 399–407. https://doi.org/10.1093/beheco/arad011

Horak, R. D., Leonard, S. P., & Moran, N. A. (2020). Symbionts shape host innate immunity in honeybees. *Proceedings of the Royal Society B: Biological Sciences*, *287*(1933), 20201184. https://doi.org/10.1098/rspb.2020.1184

Hosokawa, T., & Fukatsu, T. (2020). Relevance of microbial symbiosis to insect behavior. *Current Opinion in Insect Science*, *39*, 91–100. https://doi.org/10.1016/j.cois.2020.03.004

Hosokawa, T., Hironaka, M., Mukai, H., Inadomi, K., Suzuki, N., & Fukatsu, T. (2012). Mothers never miss the moment: A fine-tuned mechanism for vertical symbiont transmission in a subsocial insect. *Animal Behaviour*, *83*(1), 293–300. https://doi.org/10.1016/j.anbehav.2011.11.006

Hosokawa, T., Kikuchi, Y., & Fukatsu, T. (2007). How many symbionts are provided by mothers, acquired by offspring, and needed for successful vertical transmission in an obligate insect–bacterium mutualism? *Molecular Ecology*, *16*(24), 5316–5325. https://doi.org/10.1111/j.1365-294X.2007.03592.x

Janda, J. M., & Abbott, S. L. (2006). The Genus *Hafnia*: From Soup to Nuts. *Clinical Microbiology Reviews*, *19*(1), 12–28. https://doi.org/10.1128/CMR.19.1.12-28.2006

Johnson, M. T. J., & Munshi-South, J. (2017). Evolution of life in urban environments. *Science*, *358*(6363), eaam8327. https://doi.org/10.1126/science.aam8327

Katoh, K. (2002). MAFFT: A novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research*, *30*(14), 3059–3066. https://doi.org/10.1093/nar/gkf436

Kölliker, M. (2007). Benefits and costs of earwig (*Forficula auricularia*) family life. *Behavioral Ecology and Sociobiology*, *61*(9), 1489–1497. https://doi.org/10.1007/s00265-007-0381-7

Kölliker, M., & Vancassel, M. (2007). Maternal attendance and the maintenance of family groups in common earwigs (*Forficula auricularia*): A field experiment. *Ecological Entomology*, *32*(1), 24–27. https://doi.org/10.1111/j.1365-2311.2006.00831.x

Kramer, J., Thesing, J., & Meunier, J. (2015). Negative association between parental care and sibling cooperation in earwigs: A new perspective on the early evolution of family life? *Journal of Evolutionary Biology*, *28*(7), 1299–1308. https://doi.org/10.1111/jeb.12655

Krebs, C. (1999). *Ecological methodology* (2nd ed.). Harper and Row.

Kubier, A., Wilkin, R. T., & Pichler, T. (2019). Cadmium in soils and groundwater: A review. *Applied*
Geochemistry, *108*, 104388. https://doi.org/10.1016/j.apgeochem.2019.104388

Lazzeri, A. M., Mangia, N. P., Mura, M. E., Floris, I., Satta, A., & Ruiu, L. (2020). Potential of novel
food-borne *Lactobacillus* isolates against the honeybee pathogen *Paenibacillus larvae*.
Biocontrol Science and Technology, *30*(9), 897–908.
https://doi.org/10.1080/09583157.2020.1769556

Lenth, R. (2022). *emmeans: Estimated Marginal Means, aka Least-Squares Means*.
https://CRAN.R-project.org/package=emmean

Li, X.-D., Xin, L., Rong, W.-T., Liu, X.-Y., Deng, W.-A., Qin, Y.-C., & Li, X.-L. (2021). Effect of heavy
metals pollution on the composition and diversity of the intestinal microbial community of
a pygmy grasshopper (*Eucriotettix oculatus*). *Ecotoxicology and Environmental Safety*,
223, 112582. https://doi.org/10.1016/j.ecoenv.2021.112582

Lombardo, M. P. (2008). Access to mutualistic endosymbiotic microbes: An underappreciated
benefit of group living. *Behavioral Ecology and Sociobiology*, *62*(4), 479–497.
https://doi.org/10.1007/s00265-007-0428-9

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550.
https://doi.org/10.1186/s13059-014-0550-8

Luo, W., Wang, D., Xu, Z., Liao, G., Chen, D., Huang, X., Wang, Y., Yang, S., Zhao, L., Huang, H.,
Li, Y., Wei, W., Long, Y., & Du, Z. (2020). Effects of cadmium pollution on the safety of rice
and fish in a rice-fish coculture system. *Environment International*, *143*, 105898.
https://doi.org/10.1016/j.envint.2020.105898

Magoga, G., Piombo, C., Locatelli, D. P., Limonta, L., & Montagna, M. (2023). The microbiota of
Idaea inquinata developing on dry herbs. *Entomologia Experimentalis et Applicata*,
171(10), 796–805. https://doi.org/10.1111/eea.13347

Magurran, A. E., & Henderson, P. A. (2003). Explaining the excess of rare species in natural
species abundance distributions. *Nature*, *422*(6933), 714–716.
https://doi.org/10.1038/nature01547

Manthey, C., Johnston, P. R., Nakagawa, S., & Rolff, J. (2022). Complete metamorphosis and
microbiota turnover in insects. *Molecular Ecology*, mec.16673.
https://doi.org/10.1111/mec.16673

Maraci, Ö., Corsini, M., Antonatou-Papaioannou, A., Jünemann, S., Sudyka, J., Di Lecce, I.,
Caspers, B. A., & Szulkin, M. (2022). Changes to the gut microbiota of a wild juvenile
passerine in a multidimensional urban mosaic. *Scientific Reports*, *12*(1), 6872.
https://doi.org/10.1038/s41598-022-10734-7

Martinez Arbizu, P. (2017). *pairwiseAdonis: Pairwise Multilevel Comparison using Adonis*.

Martiny, J. B. H., Martiny, A. C., Brodie, E., Chase, A. B., Rodríguez-Verdugo, A., Treseder, K. K.,
& Allison, S. D. (2023). Investigating the eco-evolutionary response of microbiomes to
environmental change. *Ecology Letters*, *26*(S1). https://doi.org/10.1111/ele.14209

McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R package for reproducible interactive analysis
and graphics of microbiome census data. *PLoS ONE*, *8*(4), e61217.
https://doi.org/10.1371/journal.pone.0061217

Meunier, J. (2024). The biology and social life of Earwigs (Dermaptera). *Annual Review of*
Entomology, *69*(1), annurev-ento-013023-015632. https://doi.org/10.1146/annurev-ento-
013023-015632

Meunier, J., & Kölliker, M. (2012). Parental antagonism and parent–offspring co-adaptation interact
to shape family life. *Proceedings of the Royal Society B: Biological Sciences*, *279*(1744),
3981–3988. https://doi.org/10.1098/rspb.2012.1416

Meunier, J., Wong, J. W. Y., Gómez, Y., Kuttler, S., Röllin, L., Stucki, D., & Kölliker, M. (2012). One
clutch or two clutches? Fitness correlates of coexisting alternative female life-histories in
the European earwig. *Evolutionary Ecology*, *26*(3), 669–682.
https://doi.org/10.1007/s10682-011-9510-x

Mogren, C. L., & Trumble, J. T. (2010). The impacts of metals and metalloids on insect behavior.
Entomologia Experimentalis et Applicata, *135*(1), 1–17. https://doi.org/10.1111/j.1570-
7458.2010.00967.x

Muhammad, A., Habineza, P., Ji, T., Hou, Y., & Shi, Z. (2019). Intestinal microbiota confer
protection by priming the immune system of red palm weevil *Rhynchophorus ferrugineus*

Olivier (Coleoptera: Dryophthoridae). *Frontiers in Physiology*, *10*, 1303. https://doi.org/10.3389/fphys.2019.01303

Mulligan, C. N., Yong, R. N., & Gibbs, B. F. (2001). Remediation technologies for metal-contaminated soils and groundwater: An evaluation. *Engineering Geology*, *60*(1–4), 193–207. https://doi.org/10.1016/S0013-7952(00)00101-0

Muratoğlu, H., Sezen, K., & DemiRbağ, Z. (2011). Determination and pathogenicity of the bacterial flora associated with the spruce bark beetle, *Ips typographus* (L.) (Coleoptera: Curculionidae: Scolytinae). *Turkish Journal of Biology*. https://doi.org/10.3906/biy-0902-12

Muyzer, G., de Waal, E. C., & Uitterlinden, A. G. (1993). Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Applied and Environmental Microbiology*, *59*(3), 695–700. https://doi.org/10.1128/aem.59.3.695-700.1993

Nowak, A., Szczuka, D., Górczyńska, A., Motyl, I., & Kręgiel, D. (2021). Characterization of *Apis mellifera* gastrointestinal microbiota and lactic acid bacteria for honeybee protection—A review. *Cells*, *10*(3), 701. https://doi.org/10.3390/cells10030701

Oksanen, J., Simpson, G., Blanchet, F., Kindt, R., Legendre, P., Minchin, P., O'Hara, R., Solymos, P., Stevens, M., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B., Borcard, D., Carvalho, G., Chirico, M., De Caceres, M., Durand, S., … Weedon, J. (2022). *vegan: Community ecology package*. https://CRAN.R-project.org/package=vegan

Onchuru, T. O., Javier Martinez, A., Ingham, C. S., & Kaltenpoth, M. (2018). Transmission of mutualistic bacteria in social and gregarious insects. *Current Opinion in Insect Science*, *28*, 50–58. https://doi.org/10.1016/j.cois.2018.05.002

Powell, J. E., Martinson, V. G., Urban-Mead, K., & Moran, N. A. (2014). Routes of acquisition of the gut microbiota of the honey bee *Apis mellifera*. *Applied and Environmental Microbiology*, *80*(23), 7378–7387. https://doi.org/10.1128/AEM.01861-14

Price, M. N., Dehal, P. S., & Arkin, A. P. (2010). FastTree 2 – Approximately maximum-likelihood trees for large alignments. *PLoS ONE*, *5*(3), e9490. https://doi.org/10.1371/journal.pone.0009490

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., & Glöckner, F. O. (2012). The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Research*, *41*(D1), D590–D596. https://doi.org/10.1093/nar/gks1219

R Core Team. (2024). *R: A language and environment for statistical computing* (Version 4.3.4) [Computer software]. https://www.R-project.org/

Raulo, A., Allen, B. E., Troitsky, T., Husby, A., Firth, J. A., Coulson, T., & Knowles, S. C. L. (2021). Social networks strongly predict the gut microbiota of wild mice. *The ISME Journal*, *15*(9), 2601–2613. https://doi.org/10.1038/s41396-021-00949-3

Rodrigues, J. L. M., Silva-Stenico, M. E., De Souza, A. N., Lopes, J. R. S., & Tsai, S. M. (2006). *In situ* probing of *Xylella fastidiosa* in honeydew of a xylem sap-feeding insect using 16S rRNA-targeted fluorescent oligonucleotides. *Environmental Microbiology*, *8*(4), 747–754. https://doi.org/10.1111/j.1462-2920.2005.00958.x

Rothman, J. A., Leger, L., Kirkwood, J. S., & McFrederick, Q. S. (2019). Cadmium and selenate exposure affects the honey bee microbiome and metabolome, and bee-associated bacteria show potential for bioaccumulation. *Applied and Environmental Microbiology*, *85*(21), e01411-19. https://doi.org/10.1128/AEM.01411-19

Sachs, J. L., Skophammer, R. G., & Regus, J. U. (2011). Evolutionary transitions in bacterial symbiosis. *Proceedings of the National Academy of Sciences*, *108*(supplement_2), 10800–10807. https://doi.org/10.1073/pnas.1100304108

Sandrin, L., Meunier, J., Raveh, S., Walser, J.-C., & Kölliker, M. (2015). Multiple paternity and mating group size in the European earwig, *Forficula auricularia*. *Ecological Entomology*, *40*(2), 159–166. https://doi.org/10.1111/een.12171

Schliep, K. P. (2011). phangorn: Phylogenetic analysis in R. *Bioinformatics*, *27*(4), 592–593. https://doi.org/10.1093/bioinformatics/btq706

Schmidt, K., & Engel, P. (2021). Mechanisms underlying gut microbiota–host interactions in insects. *Journal of Experimental Biology*, *224*(2), jeb207696. https://doi.org/10.1242/jeb.207696

Seto, K. C., Güneralp, B., & Hutyra, L. R. (2012). Global forecasts of urban expansion to 2030 and
direct impacts on biodiversity and carbon pools. *Proceedings of the National Academy of*
Sciences, *109*(40), 16083–16088. https://doi.org/10.1073/pnas.1211658109

Shukla, S. P., Plata, C., Reichelt, M., Steiger, S., Heckel, D. G., Kaltenpoth, M., Vilcinskas, A., &
Vogel, H. (2018). Microbiome-assisted carrion preservation aids larval development in a
burying beetle. *Proceedings of the National Academy of Sciences*, *115*(44), 11274–11279.
https://doi.org/10.1073/pnas.1812808115

Stout, L. M., Dodova, E. N., Tyson, J. F., & Nüsslein, K. (2010). Phytoprotective influence of
bacteria on growth and cadmium accumulation in the aquatic plant *Lemna minor*. *Water*
Research, *44*(17), 4970–4979. https://doi.org/10.1016/j.watres.2010.07.073

Su, H., Wu, J., Zhang, Z., Ye, Z., Chen, Y., & Yang, Y. (2021). Effects of cadmium stress at different
concentrations on the reproductive behaviors of beet armyworm *Spodoptera exigua*
(Hübner). *Ecotoxicology*, *30*(3), 402–410. https://doi.org/10.1007/s10646-021-02365-x

Sun, Y., Tang, Y., Xu, X., Hu, K., Zhang, Z., Zhang, Y., Yi, Z., Zhu, Q., Xu, R., Zhang, Y., Liu, Z., &
Liu, X. (2020). Lead exposure results in defective behavior as well as alteration of gut
microbiota composition in flies and their offsprings. *International Journal of Developmental*
Neuroscience, *80*(8), 699–708. https://doi.org/10.1002/jdn.10067

Teyssier, A., Matthysen, E., Hudin, N. S., De Neve, L., White, J., & Lens, L. (2020). Diet contributes
to urban-induced alterations in gut microbiota: Experimental evidence from a wild
passerine. *Proceedings of the Royal Society B: Biological Sciences*, *287*(1920),
20192182. https://doi.org/10.1098/rspb.2019.2182

Teyssier, A., Rouffaer, L. O., Saleh Hudin, N., Strubbe, D., Matthysen, E., Lens, L., & White, J.
(2018). Inside the guts of the city: Urban-induced alterations of the gut microbiota in a wild
passerine. *Science of The Total Environment*, *612*, 1276–1286.
https://doi.org/10.1016/j.scitotenv.2017.09.035

Tourneur, Cole, C., & Meunier, J. (2020). The first of five moults of *Forficula auricularia* L.
(Dermaptera: Forficulidae). *The Canadian Entomologist*, *152*(6), 783–789.
https://doi.org/10.4039/tce.2020.57

Trevors, J. T., Stratton, G. W., & Gadd, G. M. (1986). Cadmium transport, resistance, and toxicity
in bacteria, algae, and fungi. *Canadian Journal of Microbiology*, *32*(6), 447–464.
https://doi.org/10.1139/m86-085

Trumbo, S. (2012). Patterns of parental care in invertebrates. In N. Royle, P. Smiseth, & M. Kölliker,
The evolution of parental care (1st ed., pp. 81–93). Oxford University Press.

Tung, J., Barreiro, L. B., Burns, M. B., Grenier, J.-C., Lynch, J., Grieneisen, L. E., Altmann, J.,
Alberts, S. C., Blekhman, R., & Archie, E. A. (2015). Social networks predict gut
microbiome composition in wild baboons. *eLife*, *4*, e05224.
https://doi.org/10.7554/eLife.05224

Van Meyel, S., Devers, S., Dupont, S., Dedeine, F., & Meunier, J. (2021). Alteration of gut
microbiota with a broad-spectrum antibiotic does not impair maternal care in the European
earwig. *Journal of Evolutionary Biology*, *34*(7), 1034–1045.
https://doi.org/10.1111/jeb.13791

Van Meyel, S., & Meunier, J. (2022). Costs and benefits of isolation from siblings during family life
in adult earwigs. *Animal Behaviour*, *193*, 91–99.
https://doi.org/10.1016/j.anbehav.2022.09.003

Vásquez, A., Forsgren, E., Fries, I., Paxton, R. J., Flaberg, E., Szekely, L., & Olofsson, T. C. (2012).
Symbionts as major modulators of insect health: Lactic acid bacteria and honeybees.
PLoS ONE, *7*(3), e33188. https://doi.org/10.1371/journal.pone.0033188

Videvall, E., Song, S. J., Bensch, H. M., Strandh, M., Engelbrecht, A., Serfontein, N., Hellgren, O.,
Olivier, A., Cloete, S., Knight, R., & Cornwallis, C. K. (2019). Major shifts in gut microbiota
during development and its relationship to growth in ostriches. *Molecular Ecology*, *28*(10),
2653–2667. https://doi.org/10.1111/mec.15087

Wang, H., Du, X., Zhang, Z., Feng, F., & Zhang, J. (2023). Rhizosphere interface microbiome
reassembly by arbuscular mycorrhizal fungi weakens cadmium migration dynamics. *iMeta*,
2(4), e133. https://doi.org/10.1002/imt2.133

Wang, Q., Zhou, Q., Huang, L., Xu, S., Fu, Y., Hou, D., Feng, Y., & Yang, X. (2022). Cadmium
phytoextraction through *Brassica juncea L.* under different consortia of plant growth-

promoting bacteria from different ecological niches. *Ecotoxicology and Environmental*
Safety, *237*, 113541. https://doi.org/10.1016/j.ecoenv.2022.113541

Wang, Y., & Rozen, D. E. (2017). Gut microbiota colonization and transmission in the burying
beetle *Nicrophorus vespilloides* throughout development. *Applied and Environmental*
Microbiology, *83*(9), e03250-16. https://doi.org/10.1128/AEM.03250-16

Wang, Z., Li, J., Zhao, P., Yu, Z., Yang, L., Ding, X., Lv, H., Yi, S., Sheng, Q., Zhang, L., Zhou, F.,
& Wang, H. (2024). Integrated microbiome and metabolome analyses reveal the effects of
low pH on intestinal health and homeostasis of crayfish (*Procambarus clarkii*). *Aquatic*
Toxicology, *270*, 106903. https://doi.org/10.1016/j.aquatox.2024.106903

Weiss, B. L., Maltz, M. A., Vigneron, A., Wu, Y., Walter, K. S., O'Neill, M. B., Wang, J., & Aksoy, S.
(2019). Colonization of the tsetse fly midgut with commensal *Kosakonia cowanii* Zambiae
inhibits trypanosome infection establishment. *PLOS Pathogens*, *15*(2), e1007470.
https://doi.org/10.1371/journal.ppat.1007470

Wenzel, M., Schonig, I., Berchtold, M., Kampfer, P., & Konig, H. (2002). Aerobic and facultatively
anaerobic cellulolytic bacteria from the gut of the termite *Zootermopsis angusticollis*.
Journal of Applied Microbiology, *92*(1), 32–40. https://doi.org/10.1046/j.1365-
2672.2002.01502.x

Wu, Y., Chen, C., Wang, G., Xiong, B., Zhou, W., Xue, F., Qi, W., Qiu, C., & Liu, Z. (2020).
Mechanism underlying earthworm on the remediation of cadmium-contaminated soil.
Science of The Total Environment, *728*, 138904.
https://doi.org/10.1016/j.scitotenv.2020.138904

Yang, Z., Xu, F., Li, H., & He, Y. (2021). Beyond samples: A metric revealing more connections of
gut microbiota between individuals. *Computational and Structural Biotechnology Journal*,
19, 3930–3937. https://doi.org/10.1016/j.csbj.2021.07.009

Zaneveld, J. R., McMinds, R., & Vega Thurber, R. (2017). Stress and stability: Applying the Anna
Karenina principle to animal microbiomes. *Nature Microbiology*, *2*(9), 17121.
https://doi.org/10.1038/nmicrobiol.2017.121

Zhai, Q., Tian, F., Zhao, J., Zhang, H., Narbad, A., & Chen, W. (2016). Oral administration of
probiotics inhibits absorption of the heavy metal cadmium by protecting the intestinal
barrier. *Applied and Environmental Microbiology*, *82*(14), 4429–4440.
https://doi.org/10.1128/AEM.00695-16

Zhai, Q., Wang, G., Zhao, J., Liu, X., Narbad, A., Chen, Y. Q., Zhang, H., Tian, F., & Chen, W.
(2014). Protective effects of *Lactobacillus plantarum* CCFM8610 against chronic cadmium
toxicity in mice indicate routes of protection besides intestinal sequestration. *Applied and*
Environmental Microbiology, *80*(13), 4063–4071. https://doi.org/10.1128/AEM.00762-14

Zhang, H., & Reynolds, M. (2019). Cadmium exposure in living organisms: A short review. *Science*
of The Total Environment, *678*, 761–767. https://doi.org/10.1016/j.scitotenv.2019.04.395

Zhang, S., Chen, A., Deng, H., Jiang, L., Liu, X., & Chai, L. (2023). Intestinal response of *Rana*
chensinensis larvae exposed to Cr and Pb, alone and in combination. *Ecotoxicology and*
Environmental Safety, *255*, 114774. https://doi.org/10.1016/j.ecoenv.2023.114774

Zhang, S., Jin, Y., Zeng, Z., Liu, Z., & Fu, Z. (2015). Subchronic exposure of mice to cadmium
perturbs their hepatic energy metabolism and gut microbiome. *Chemical Research in*
Toxicology, *28*(10), 2000–2009. https://doi.org/10.1021/acs.chemrestox.5b00237

Zhang, Y., Yu, X., Sun, S., Li, Q., Xie, Y., Li, Q., Zhao, Y., Pei, J., Zhang, W., Xue, P., Zhou, Z., &
Zhang, Y. (2016). Cadmium modulates hematopoietic stem and progenitor cells and skews
toward myelopoiesis in mice. *Toxicology and Applied Pharmacology*, *313*, 24–34.
https://doi.org/10.1016/j.taap.2016.10.016

909