Nde1 Promotes Lis1 Binding to Full-Length Autoinhibited Human Dynein-1

Jun Yang[1,4], Yuanchang Zhao [2,3,4], Pengxin Chai [1,4], Ahmet Yildiz[2,3,*], Kai Zhang[1,*]

[1]Department of Molecular Biophysics and Biochemistry, Yale University, New Haven, CT, 06511, USA

[2]Physics Department, University of California, Berkeley, CA 94709, USA

[3]Department of Molecular and Cell Biology, University of California, Berkeley, CA 94709, USA

[4]These authors contributed equally

*Correspondence: Email: yildiz@berkeley.edu (A.Y.), jack.zhang@yale.edu (K.Z.)

Abstract

Cytoplasmic dynein-1 (dynein) is the primary motor for the retrograde transport of intracellular cargoes along microtubules. The activation of the dynein transport machinery requires the opening of its autoinhibited Phi conformation by Lis1 and Nde1/Ndel1, but the underlying mechanism remains unclear. Using biochemical reconstitution and cryo-electron microscopy, we show that Nde1 significantly enhances Lis1 binding to autoinhibited dynein and facilitates the opening of Phi. We discover a key intermediate step in the dynein activation pathway where a single Lis1 dimer binds between the Phi-like (Phi^L) motor rings of dynein. In this "Phi^L-Lis1", Lis1 interacts with one of the motor domains through its canonical interaction sites at the AAA+ ring and stalk and binds to the newly identified AAA5, AAA6, and linker regions of the other motor domain. Mutagenesis and motility assays confirm the critical role of the Phi^L-Lis1 interface. This intermediate state is instantly and efficiently formed in the presence of Nde1, but Nde1 is not part of the Phi^L-Lis1. These findings provide key insights into the mechanism of how Nde1 promotes the Lis1-mediated opening of Phi dynein.

Introduction

Cytoplasmic dynein is a motor protein essential for transporting nearly all intracellular cargoes toward the minus end of microtubules (MTs) in most eukaryotes. Dynein's cargoes include organelles, proteins, RNA, viruses, and vesicles[1]. Additionally, dynein plays key roles in other cellular processes such as mitosis and organelle positioning[1,2]. Mutations affecting dynein or its regulatory proteins have been linked to neurodevelopmental and neurodegenerative diseases, including spinal muscular atrophy (SMA), ALS, Huntington's disease, lissencephaly, and microcephaly[3-9].

The 1.4 MDa dynein complex contains pairs of six subunits. The largest subunit is the dynein heavy chain (DHC), which contains the N-terminal tail domain and C-terminal motor domain. The tail domain facilitates dimer formation, recruits the dimers of intermediate chain (DIC), light intermediate chain (DLIC), and interacts with dynactin, cargo adaptors and other regulatory proteins[10-13]. The N-terminus of DIC (DIC-N) recruits three pairs of light chains, Robl, LC8, and Tctex[1,14,15]. DIC segments binding to LC8 and Tctex, known as IC-LC tower, play a crucial role in assembling the dynein-dynactin-adaptor (DDA) complex[11]. The dynein motor domain belongs to AAA+ (ATPases Associated with diverse cellular Activities) protein families and consists of six AAA subdomains (AAA1-6)[16,17]. AAA1 is the primary subdomain that hydrolyzes ATP to power dynein motility along MTs. The dynein motor domain attaches to MTs via the coiled-coil stalk and the MT-binding domain (MTBD) and connects to the tail through the linker domain located at the surface of the AAA+ ring[17-20].

Dynein alternates between two key conformational states: the autoinhibited Phi (similar to the Greek letter "φ") and the open conformations[14,21]. Phi dynein adopts a compact structure that limits its interactions with MTs and dynactin, which serves to minimize unnecessary ATP hydrolysis when motor protein is not engaged in active transport[14,22]. Opening of the Phi enables dynein to assemble with dynactin and the cargo adaptor, allowing processive movement along MTs[10,14], a process supported by dynein regulators Nde1/Ndel1 and Lis1[3,15,23-30]. However, the underlying mechanism promoting Phi to open transition is unclear.

Lis1, the first gene identified in relation to a neuronal migration disease, plays a crucial role in dynein-related function[3,11,31-39]. Lis1 possesses an N-terminal LisH domain that facilitates dimerization and a C-terminal WD-40 β-propeller domain that binds to Nde1/Ndel1, dynein, and other proteins[3]. Both domains are important at different stages of dynein activation[40-42]. The LisH domain interacts with dynactin p150 and DIC-N, thus promoting the recruitment of dynactin and adaptors at the later stage of dynein activation[11]. WD-40 domains of Lis1 can directly bind to the dynein motor domain at the AAA3-AAA4 sites (site-ring) and stalk coiled-coil (site-stalk)[11,38,43]. Functional studies proposed that Lis1 facilitates the formation of highly processive DDA complex by favoring the release of Phi dynein and stabilizing open dynein[34-37,44]. Lis1 binding is

thought to be incompatible with Phi dynein based on the steric clash when docking a
Lis1 β-propeller to the motor domains of Phi dynein[3,35,37,44]. However, there is no direct
biochemical or structural evidence of whether Lis1 can open Phi dynein and
intermediary states that facilitate opening of Phi dynein remain unclear. A recent
structural study on tail-truncated yeast dynein motor domains reported a "Chi"
conformation, in which two Lis1 dimers are wedged between two AAA+ rings[37],
suggesting that two Lis1 dimers between the motor domains are required to crack open
Phi dynein. Furthermore, it is well-known that the AAA+ ring undergoes substantial
conformational changes under different nucleotide-binding states, which can potentially
regulate dynein–Lis1 binding. A more recent work has demonstrated that specific
nucleotide "codes" at the three variable nucleotide-binding sites (AAA1, 3, and 4)
govern the stoichiometry of dynein–Lis1 interactions by tuning their binding affinity at
two distinct locations[45]. However, the intermediate structure for full-length human dynein
alone bound to Lis1 is still lacking and whether dynein forms a Chi conformation at the
initial state of its activation remains unclear.

Nde1 and its paralog Ndel1[12,23], are critical for all dynein-mediated functions in cell
division[28,46], cargo trafficking[47,48] and neuronal migration[49,50]. Nde1/Ndel1 is predicted to
be composed of the N-terminal coiled-coil domain and the C-terminal unstructured
region[51-53]. The coiled-coil domain interacts with DIC-N, which overlaps with the DIC
binding site of the p150 subunit of dynactin[52,54,55]. It also interacts with the WD-40
domains of Lis1, and its binding site on WD-40 domain overlaps with the Lis1 binding
site of dynein[26,52]. The multi-protein interaction modes of Nde1/Ndel1, along with its
overlapping binding sites with other proteins, make the functional interpretation of
Nde1/Ndel1 elusive. It has been proposed that Nde1 tethers Lis1 to dynein[12,26,56-58] and
promotes Lis1-mediated activation of dynein[11,26] (**Fig. 1a**). According to this model, Phi
dynein first adopts an 'open' conformation. Lis1 stabilizes the open dynein by preventing
it from transitioning back to Phi, thus favoring the assembly of the DDA complex[11,34,35,44].
Consistent with this model, overexpression of Lis1 can rescue the deletion of
Nde1/Ndel1 in cells[58,59] and Nde1/Ndel1-mediated recruitment of Lis1 to dynein
enhances DDA assembly in vitro[26]. An alternative model suggests that Ndel1 negatively
regulates dynein activation by competing with p150 for DIC-N binding and by
sequestering Lis1 away from dynein[52]. Consequently, how Lis1 and Nde1/Ndel1 form a
complex with dynein and promote the opening of Phi dynein is not well understood.

To understand the mechanism by which Nde1/Ndel1 and Lis1 prime dynein for the DDA
assembly, we investigated how human Lis1 and Nde1 affect the conformational states
of full-length human dynein using biochemical reconstitution and electron microscopy.
We showed that Nde1 promotes the formation of the dynein-Lis1 complex. Using
negative-stain EM, we provide direct evidence that Lis1 or Nde1 alone has little effect
on the equilibrium between the Phi and open conformations of dynein, but Lis1 and

Nde1 together significantly bias the equilibrium toward the open conformation,
demonstrating that Nde1 acts like a molecular chaperone to promote Lis1-mediated
opening of dynein. Cryo-EM imaging of dynein-Lis1 in the presence or absence of Nde1
captures a new intermediate and rate-limiting state during the dynein activation process,
characterized by a single Lis1 dimer binding between the motor domains of Phi dynein.
Lis1 binding to Phi dynein causes the rotation between two motor domains to relieve the
steric clash, forming a Phi-like dynein and Lis1 complex, named "Phi^L-Lis1". While Lis1
binds to one of the motor domains through its canonical interaction sites, it also forms
new interfaces between the other motor domain at the AAA5, AAA6, and linker regions.
Mutagenesis at the novel interfaces together with single molecule motility assays
supported the critical role of this Phi^L-Lis1 during the dynein activation process.
Collectively, our findings shed light on the roles of Nde1 and Lis1 in the dynein
activation pathway.

Results

Nde1 promotes Lis1 binding to Phi dynein and cooperatively releases dynein
autoinhibition

To determine whether Nde1 promotes Lis1 binding to dynein, we performed mass
photometry (MP) assays to assess the Lis1 and Nde1 binding to full-length human
dynein under different conditions. In the absence of Nde1, Lis1 alone exhibited an
increased binding to dynein in a time-dependent manner, with the formation of a 37%
1:1 dynein:Lis1 (DL) complex within 60 minutes. The inclusion of Nde1 significantly
enhanced the dynein-Lis1 binding (69% of complex formation) in less than a minute,
consistent with Nde1/Ndel1-mediated tethering of Lis1 to dynein[12,26,57] (**Fig. 1b**). To
further test if Nde1 preferentially recruits Lis1 to Phi or open conformation, we
performed these assays using the Phi mutant of dynein[14] that only forms open
conformation. Lis1 was readily bound to open dynein, but the addition of Nde1 did not
further enhance Lis1 binding to dynein (**Extended Data Fig.1**), suggesting that Nde1 is
required for Lis1 recruitment to Phi dynein. These results are also consistent with the
previous observation that the Nde1 addition does not further enhance Lis1-mediated
activation of DDA complexes assembled with the Phi dynein mutant[26]. Interestingly, the
formation of the dynein-Lis1-Nde1 tripartite complex was not observed (**Fig. 1b and**
Extended Data Fig.1), regardless of different nucleotide conditions (**Extended Data**
Fig. 2). Although previous reports indicated that Nde1 can interact with DIC-N in single
molecule imaging[12,26] and pull-down assays[26,52], we also did not detect the dynein-Nde1
complex (**Fig. 1b**, and **Extended Data Fig. 2**), suggesting that Nde1 rapidly dissociates
from the complex after handing off Lis1 to dynein[12].

To determine whether Nde1-mediated Lis1 recruitment to dynein shifts the equilibrium
between Phi and open conformations, we used the negative stain EM imaging[14] to

quantify the ratio of Phi dynein in the presence and absence of Lis1 and Nde1.
Specifically, we used freshly prepared dynein with ~50% of motors forming the Phi and
then incubated dynein with Lis1 and Nde1 (**Fig. 1c and Extended Data Fig. 3**) in the
presence of ATP. We found that Lis1 or Nde1 alone does not change the Phi ratio
compared with the control (**Fig. 1d-e**). However, the Phi ratio decreased 44% when we
incubated dynein with both Lis1 and Nde1 at a ratio of 1:2:2 (**Fig. 1d-e**). Collectively,
our results demonstrate that Nde1 specifically promotes Lis1 binding to Phi dynein and
facilitates opening of this autoinhibited conformation.

A novel Phi^L-Lis1 structure

Our MP results suggest that there is a rate-limiting step of Lis1 binding to Phi dynein,
and this step can be significantly accelerated by the addition of Nde1 (**Fig. 1b**). We
used cryo-EM to capture the potential intermediate states to reveal the structural basis
of this process. We focused on the particles that form the autoinhibited dynein (**Fig. 2a,
b and Extended Data Fig. 4, 5**). In the absence of Nde1, we unexpectedly observed a
novel structure in which a Lis1 dimer is wedged between the two stacked motor rings of
Phi dynein. The dynein in this complex shows a compact conformation, similar to but
not the same as the previously reported Phi structure[14] and we referred this complex as
the "Phi^L-Lis1" (**Fig. 2a, c and Supplementary Video 1**). Despite excess Lis1, nearly
half of Phi dynein does not bind to Lis1 (42.7% Phi vs. 57.3% Phi^L-Lis1) (**Fig. 2a**).
Notably, we did not observe the "Chi" (two Lis1s bound to dynein), suggesting that the
"Chi" conformation may be specific to yeast dynein or may form when truncated dynein
containing only the motor domains is used instead of full-length motor[37].

In the Phi^L-Lis1 structure, Lis1 appears on the same side as the IC-LC tower of dynein
(front side), independent of the presence of Nde1 (**Fig. 2a, b**). Based on this structural
observation, along with previous evidence indicating that DIC-N can bind to both Lis1[11]
and Nde1[12], we speculate that the transient dynein-Lis1-Nde1 forms only on the IC-LC
tower side. Although we do not observe clear cryo-EM densities for the DIC-N, it is
possible that the DIC-N can weakly interact with Lis1, thus recruiting Lis1 to the front
side of dynein.

Previous results suggest Lis1 can affect dynein's mechanochemical cycle and
nucleotide state[11,60,61]. The high-resolution structure of the motor domain enabled us to
identify and compare the nucleotide states of the motor domains in Phi and Phi^L-Lis1
structures. Specifically, we found that AAA1 pockets of both motor domains exhibit the
same ADP-Mg^{2+} density (**Extended Data Fig.6a, b**), accompanied by a flexible sensor-I
loop, indicating the intermediate state of Pi releasing (**Extended Data Fig. 6c, d**).
Similarly, AAA3 pockets show clear ADP binding (**Extended Data Fig.6a, b**), indicating
that Lis1 binding does not influence the nucleotide states of the motor domains of
autoinhibited dynein.

We next premixed equimolar Nde1 and Lis1, then added dynein to achieve a final 1:2:2
ratio of dynein:Lis1:Nde1 (**Fig. 2b**). Strikingly, we obtained a similar Phi^L-Lis1 structure,
but no longer observed the Phi dynein alone in the presence of Nde1 (**Fig. 2a, b and**
Supplementary Video 1). This is consistent with the MP analysis showing that Nde1
instantly promotes Lis1 binding to dynein (**Fig. 1b**) and forming Phi^L-Lis1. These results
suggest that Phi^L-Lis1 formation is a rate-limiting intermediate state before dynein
opening. Remarkably, none of the 3D classes shows clear Nde1 density in Phi^L-Lis1,
suggesting that Nde1 is not part of this complex.

We also analyzed individual motor domains of open dynein and did not observe an
increase in the propensity of the dynein-Lis1 complex compared to open dynein alone in
the presence of Nde1 (**Extended Data Fig. 7**). Consistent with MP of the Phi dynein
mutant[14] (**Extended Data Fig. 1**), these results show that Nde1 promotes Lis1 binding
to Phi dynein to form Phi^L-Lis1, whereas Lis1 can readily bind to open dynein and does
not require Nde1.

Lis1 induces a relative rotation in the Phi conformation to accommodate its
binding

Consistent with previous reports[35,37,44], our structural analysis reveals a severe steric
clash between Lis1 and Phi motor domain A (**Fig. 3a**) and this needs to be relieved to
accommodate Lis1's binding in Phi^L-Lis1 (**Fig. 3b**). Comparing Phi to Phi^L-Lis1 reveals a
relative rotation between the two motor domains, which results in the groove on the
front side of dynein becoming larger than the corresponding groove on the back side
(**Fig. 3c, d**). The enlarged groove on the front side allows Lis1 to fit between Phi^L motor
domains. However, docking of Lis1 to the back side with a smaller groove shows severe
clashes, explaining why there is only one Lis1 present in Phi^L-Lis1 (**Fig. 3e**). The
rotation between the motor domains in Phi^L-Lis1 also causes a slight anticlockwise twist
in the neck region (**Fig. 3d**). This twist likely promotes the unwinding of the tail,
generating a trend toward an open conformation of dynein. We concluded that Lis1
binding induces a rotation of Phi dynein motor domains to avoid steric clash with Lis1
(**Fig. 3 and Supplementary Video 1**).

Novel interactions identified in Phi^L-Lis1

In Phi^L-Lis1, WD-40 domains of Lis1 interact with both motor domains of dynein (MD-A
and MD-B). While Lis1 interacts with MD-B through its canonical interaction sites at the
AAA3, AAA4, and AAA5 regions[38,43,62], we observed previously uncharacterized
interaction sites of Lis1 with the linker, AAA6, and AAA5 regions of MD-A (**Fig. 4a,**
Extended Data Fig. 8 and Supplementary Video 1). Sequence alignment of Lis1
homologs shows that MD-A and Lis1 interface is highly conserved among higher
eukaryotes but less conserved in yeast (**Extended Data Fig. 9**), suggesting different
regulatory roles of Lis1 between higher eukaryotes and yeast. Structural comparison

indicates that two motor domains in Phi^L-Lis1 adopt an almost identical conformation.
The root-mean-squared-displacement of alpha carbon atoms (Cα-RMSD) of the two
motor domains was 0.513 Å (**Fig. 4b**). Additionally, the MD-B bound with Lis1$_{ring}$ and
Lis1$_{stalk}$ in our results also shows no significant difference from the structure of Lis1
bound to the human dynein motor domain[38] (Cα-RMSD: 0.868 Å) (**Fig. 4c**), suggesting
Lis1 binding does not induce structural changes within an individual motor domain.

The interactions between MD-A and Lis1$_{ring}$ are notably compact (**Fig. 4a and**
Extended Data Fig. 8). The WD-40 domain of Lis1 interacts with dynein MD-A at
regions distributed across the linker, AAA6, and AAA5 region (**Fig. 4d-f**). Specifically, at
the linker-Lis1 binding site, the side chains of M329 and E300 of Lis1 engage in
hydrophobic and polar interactions with the side chains of V1563, P1562 and H1559 of
dynein. Additionally, K303 and S304, located on the flexible loop of the Lis1 WD-40
surface, interact with R1621, D1556, and E1622 of linker region (**Fig. 4d, Extended**
Data Fig. 8a).

Within the AAA6 interaction region, the interface is characterized by polar interactions
involving N203, D205, and Q222 of the Lis1 WD-40 domain and K4089, N4085, Q4117,
and S4115 of MD-A. The side chain of K4089 forms a salt bridge with the side chain of
D205 and establishes a hydrogen bond with the oxygen atom in the main chain of N203.
Q222 and D205 of Lis1 also form polar interactions with S4115 and Q4117 and N4085
residues of dynein (**Fig. 4e, Extended Data Fig. 8b**).

The AAA5-Lis1$_{ring}$ WD-40 interface shows a more compact interaction (**Fig. 4f**). This
interface is mainly composed of residues Q3636, S3613, T3612, D3616, D3617, A3618,
and K3621 of dynein MD-A and Y225, C226, G190, H191, M172, and D192 of the Lis1
WD-40 domain (**Fig. 4f and Extended Data Fig. 8b**). Notably, the side chain of Y225 of
Lis1 forms a polar interaction with the side chain of Q3636. Additionally, the side chain
of D3616 and the main chain of D3617 form hydrogen bonds with the main chain of
C226. The side chain of A3618 forms hydrophobic interaction with the main chain of
G190, while residues M172 and H191 of Lis1 form hydrophobic and hydrophilic
interactions with the side chain of K3621. Additionally, novel interactions are formed
between MD-A and MD-B in the Phi^L-Lis1, compared with canonical Phi (**Extended**
Data Fig. 10), suggesting that the dynein Phi^L-Lis1 is a stable conformation.

The Phi^L-Lis1 interface regulates Nde1-dependent dynein activation

To evaluate whether the new interaction sites we detected between Lis1 and dynein
MD-A in Phi^L-Lis1 are critical for activation of dynein, we introduced three sets of Lis1
mutations targeting the interfaces that interact with the linker, AAA6, and AAA5 of
dynein. Key residues of Lis1$_{ring}$ at each interface were mutated to charged residues or
alanine to disrupt these interactions (Lis1^{linker}: E300K, K303E, S304R and M329A;
Lis1^{AAA6}: N203K, D205K and Q222A; Lis1^{AAA5}: M172K, D192K, Y225A, C226D). Similar

to wild-type Lis1 (WT Lis1), these Lis1 mutants formed homodimers and interacted with
Nde1 (**Fig. 4g**). However, the binding efficiency of Nde1 was reduced for the Lis1^{AAA5}
mutant (**Fig. 4g**), suggesting that the AAA5-Lis1$_{ring}$ interface may share a region
involved in Nde1 binding to Lis1 (**Extended Data Fig. 11**). The Lis1 mutants also bound
to dynein, and the dynein binding efficiency of Lis1 was increased with Nde1 (**Fig. 4h**).
Notably, we detected a mass population corresponding to dynein bound to one Nde1
and two Lis1^{AAA6} mutants, indicating that mutations to the AAA6 interaction site of Lis1
prevent dissociation of Nde1 from the dynein-Lis1 complex (**Fig. 4h**).

To determine how these mutations affect activation of dynein motility, we assayed
single molecule motility of complexes assembled with wild type dynein, dynactin, and
the BicDR1 adaptor (DDR) on surface-immobilized MTs in vitro in the presence and
absence of Lis1 and Nde1. Consistent with our previous observations[26], Lis1 enhanced
the run frequency of DDR about 3-fold, and Nde1 and Lis1 together increased the run
frequency 15-fold (**Fig. 4i, j and Supplementary Video 2**). In the absence of Nde1,
Lis1 mutants activated dynein motility at similar levels of WT Lis1. In the presence of
Nde1, Lis1^{linker} and Lis1^{AAA5} triggered activation of DDR motility similar to WT Lis1 (**Fig.**
4i, j). In comparison, Nde1 failed to enhance Lis1^{AAA6}-mediated dynein motility,
suggesting that Lis1^{AAA6} cannot form the stable Phi^{L}-Lis1 complex and open the Phi
conformation (**Fig. 4i, j**). Together with MP results, we demonstrate that mutations in
the Lis1^{AAA6} interface disrupt Nde1-mediated opening of the Phi conformation by Lis1,
highlighting the importance of the Phi^{L}-Lis1 structure in the dynein activation pathway.

Discussion

In this study, we investigated the structure and mechanism of how Lis1 and Nde1
rescue dynein from autoinhibition prior to the assembly of active dynein transport
machinery. Using negative stain EM imaging, we directly showed that Nde1 and Lis1
cooperatively promote the opening of Phi dynein, whereas neither Nde1 nor Lis1 alone
exhibited a significant effect on Phi opening (**Fig. 1c-e**). Lis1 can readily bind to open
dynein (**Extended Data Fig. 1**) and facilitate the assembly of DDA complexes and Nde1
addition does not further enhance DDA motility[26]. Despite that Lis1 alone can bind to the
Phi dynein and potentially open this autoinhibited conformation[45], we show that Nde1
facilitates Lis1 to dock onto Phi dynein more efficiently and promote its switch it to the
open state before DDA assembly in this work. Similar to molecular chaperones, Nde1
rapidly dissociates from dynein after handing off Lis1, promoting dynein-Lis1 complex
formation but not existing in the final complex. Because Nde1 has an overlapping
binding site with the p150 subunit of dynactin on DIC-N[12], dissociation of Nde1 from
DIC-N after tethering Lis1 to dynein may enable efficient recruitment of dynactin to
dynein-Lis1 complexes.

Our structural and functional studies of dynein-Lis1 complexes revealed a key
intermediate step on Lis1 and Nde1 mediated opening of dynein. Although a dynein
dimer contains Lis1 binding sites on each motor domain, our MP assays showed that
Phi dynein recruits a single Lis1. Using cryo-EM imaging, we revealed a Phi^L-Lis1
structure in which the two AAA+ rings of Phi dynein rotate slightly backward to
accommodate Lis1 binding to the front side. This rotational motion reduces the spacing
between the AAA+ rings, thereby preventing Lis1 binding to the back side. Preferential
binding of Lis1 to the front side may be due to the IC-LC tower, which is located at the
front side of the Phi motor.

The Phi^L-Lis1 is fundamentally distinct from the previously reported Chi of yeast dynein
monomers[37]. The Chi is stabilized by two Lis1, one on each side, and adopts a more
open and extended conformation compared to the Phi^L-Lis1 motor domains (**Extended
Data Fig. 12**). In comparison, our study utilized full-length, wild-type human dynein and
we could not detect Chi dynein even when we used this conformation as a reference
during cryo-EM image processing. It is possible that isolated dynein motor domains may
prefer to recruit two Lis1s and form more extended Chi. In comparison, full-length
dynein readily forms the compact Phi and structural constraints imposed by the tail
domain may restrict the relative movement of the Phi motor domains. Lis1 binding to the
front side of Phi^L dynein reduces the spacing on the back side, thereby preventing the
formation of Chi. Most interaction sites located at Lis1$_{ring}$ surface of Chi-Lis1 are also
present in that of Phi^L-Lis1 (**Extended Data Fig. 9**). In comparison, Phi^L-Lis1 exhibits
more compact interactions (**Fig 4 and Extended Data Fig. 8,10**). Consistent with
Phi^L-Lis1, we did not detect complexes with one dynein and two Lis1s in MP,
suggesting that Phi^L-Lis1, rather than Chi, is the stable intermediate of full-length human
dynein.

The mutagenesis of the interactions between Lis1 and the AAA6 subdomain of MD-A in
Phi^L-Lis1 disrupts Nde1's ability to promote Lis1-mediated dynein activation, confirming
that the Phi^L-Lis1 is a key intermediate in the dynein activation pathway. However,
these mutations did not disrupt the mutant Lis1's ability to increase the run frequency of
dynein several-fold on its own. This is because WT dynein can be either in open or Phi
conformations with near equal probability in our conditions. Our model predicts that
mutant Lis1 can still bind and enhance DDA assembly of open dynein without Nde1.
However, it cannot further enhance dynein motility synergistically with Nde1 because
this mutant is deficient in forming Phi^L-Lis1.

Based on our results and previous observations, we propose a mechanism underlying
dynein activation by Lis1 and Nde1. In the absence of Nde1, Lis1 alone can bind to both
DIC-N and dynein motor domains[11], inducing a conformational change from canonical
Phi to Phi^L. However, the efficiency of this process is low and Lis1 cannot open Phi^L
dynein on its own (**Fig.1c-e, 2a**). In the presence of Nde1, Lis1-Nde1 is initially recruited

to DIC-N positioned at the front side of Phi dynein, facilitating more efficient binding of
Lis1 to the front side of Phi. The local enhancement of Lis1 near the Phi motor by Nde1
facilitates more efficient binding of Lis1 to Phi dynein (**Fig. 5 step-i, -ii**). Nde1 and Lis1
form a transient Phi^L-Lis1-Nde1 complex (**Fig. 5 step-i**). Nde1 dissociates
spontaneously, leading to Phi^L-Lis1 formation (**Fig. 5 step-ii**). Lis1 binding induces a
slight backward rotation of the two motor rings in Phi^L-Lis1 (**Fig. 3c**), suggesting an
intermediate state prior to an open state. Additionally, a slight twist in the neck region,
caused by motors rotation and likely inducing an unwinding trend in the tail, may also
contribute to dynein opening (**Fig. 3d**). Subsequently, Phi^L-Lis1 transitions to open
dynein-Lis1 with the assistance of Nde1 (**Fig. 5 step-iii**). The binding of Lis1 to dynein
facilitates DDA assembly and activates dynein motility by recruiting the p150 subunit of
dynactin to dynein through its LisH domain[11] (**Fig. 5, step-iv**). Future studies are
required to understand how Nde1 hands off Lis1 to dynein and why it rapidly dissociates
from dynein. In addition to its tethering role and facilitating the formation of Phi^L-Lis1, it
remains to be determined whether Nde1 has additional roles in helping Lis1 convert
Phi^L-Lis1 to the open dynein.

Methods

Cloning and expression

The plasmid encoding full-length human dynein[63] was generously provided by Andrew Carter (His-ZZ-TEV-SNAPf DHC1_IC2C_LIC2_Tctex1_Robl1_LC8, Addgene plasmid #111903). The His-ZZ-TEV-SNAPf tag is fused to the N-terminus of the dynein heavy chain. Human Lis1 and Nde1 (residues N-terminal 1-190 residues, which functions similarly to full-length Nde1 in the single motility assay[26]), and the mouse BIDCR1 gene, were each cloned individually into the pOmniBac backbone. The constructs featured a ZZ-TEV tag at the N-terminus and a SNAPf tag at the C-terminus. Lis1 mutants containing point mutations were generated using purchased DNA fragments (IDT) containing the mutations and inserted into the plasmid backbone. The mutations were verified by Oxford Nanopore full-plasmid sequencing. The constructs used in this study are listed in **Supplementary Table 1**.

These proteins were all expressed in insect sf9 cells, as describe previously[14,19,26] with slight modifications. Briefly, Bacmid DNA isolated from the from DH10MultiBac competent cells (Geneva Biotech) were transfected into the in sf9 insect cells with the Cellfectin® II (Gibco) reagent. Protein expression in sf9 cells was accomplished by infecting them with P2 virus at a cell density of 2.5 million cells/mL. For dynein expression, 28 mL of P2 virus was added into a 1.4 L culture of sf9 cells. For Lis1, Nde1, and BicDR1, 7 mL P2 virus was used to infect the 0.7 L sf9 cells. Cells were harvested after 75 hours by centrifugation at 1000 rcf for 15 minutes at 4°C. The cell pellets were flash-frozen in liquid nitrogen and stored at -80°C.

Protein purification

Purification for full-length human dynein was previous described[19]. Briefly, the cell pellets from a 1.4 L cell culture were resuspended in 100 mL lysis buffer (50 mM HEPES pH 7.2, 100 mM NaCl, 1 mM DTT, 0.1 mM ATP, 10% glycerol) containing 2 tablets of Complete EDTA-free protease inhibitor (Roche) and 2 mM PMSF. The suspension was homogenized using a Dounce with a tight plunger for 15~25 strokes, followed by clarification through centrifugation at 65,000 rpm with a Ti70 rotor (Beckman) for 1 hour at 4°C. The supernatant was then incubated with 3 mL IgG Sepharose 6 fast flow resin (Cytiva) for 3~4 hours on a roller at 4°C, followed by washed with 200 mL lysis buffer and 200 mL TEV buffer (50 mM Tris–HCl pH 7.4, 150 mM K-acetate, 2 mM Mg-acetate, 1 mM EGTA, 10% glycerol, 0.1 mM ATP, 1 mM DTT). Afterward, the resins were incubated with TEV buffer supplemented with 400 ug TEV protease overnight at 4°C. The supernatant was collected and concentrated with a 100 kDa MWCO Amicon concentrator, then loaded into a TSKgel G4000 column pre-equilibrated with the GF150 buffer (25 mM HEPES pH 7.2, 150 mM KCl, 1 mM $MgCl_2$, 5 mM DTT, 0.1 mM ATP). Peak fractions were collected and concentrated to 2~3 mg/mL for Cryo-EM grid

preparation. The quality of the sample was evaluated with the SDS-PAGE gels and the
negative-stain EM.

The purification of Lis1, Nde1, and BicDR1 from a 0.7 L cell culture followed a similar
protocol to that of dynein, with a few modifications. Specifically, 50 mL of lysis buffer
was used to resuspend the cell pellets, and ATP was omitted from the GF150 buffer.
And Superose 6 Column (Cytiva) was used for size exclusion chromatography. The
concentrated proteins were aliquoted, flash-frozen in liquid nitrogen, and stored at -80°C.
The quality of the proteins was assessed using SDS-PAGE gels.

Dynactin was isolated from pig brains through a series of purification steps, including
SP Sepharose Fast Flow and MonoQ ion exchange chromatography (Cytiva), followed
by size exclusion chromatography using a TSKgel G4000SWXL column (Tosoh), as
described by previous protocol[64].

MT reconstitution

MTs were reconstituted using porcine tubulin, which was either purchased from
Cytoskeleton or purified in-house in MT buffer (25 mM MES, 70 mM NaCl, 1 mM MgCl2,
1 mM EGTA, and 1 mM DTT, pH 6.5). The tubulin was concentrated to 10 mg/mL at
4°C, then flash-frozen and stored at -80°C. To polymerize the MTs, the tubulin was
diluted to 5 mg/mL in MT buffer supplemented with 3 mM GTP. The tubulin mixture was
incubated on ice for 5 minutes and then transferred to a 37°C incubator for 1 hour. After
incubation, the MTs were pelleted at 20,000 rcf for 8 minutes at room temperature and
resuspended in MT buffer supplemented with 5 µM paclitaxel before being stored at
room temperature.

MP assay

High-precision coverslips (Azer Scientific) were cleaned by alternating washes with
isopropanol and water three times in a bath sonicator, then air-dried. The gasket was
cleaned similarly, without sonication, and air-dried before being placed onto a clean
coverslip. A total of 14 µL of filtered mass photometry buffer (30 mM HEPES pH 7.4, 5
mM $MgSO_4$, 1 mM EGTA, and 10% glycerol) was added to a well for autofocus. The
protein sample was then applied to the well and diluted to a concentration of 5–20 nM in
the buffer. Protein contrast data were collected using a TwoMP mass photometer
(Refeyn 2) with two technical replicates. The instrument was calibrated with a standard
mix of conalbumin, aldolase, and thyroglobulin. MP profiles were analyzed by fitting to
multiple Gaussian peaks, with the mean, standard deviation, and percentages
calculated using DiscoverMP software (Refeyn). The data for parameters of a multi-
Gaussian fit of MP measurements is summarized in **Supplementary Table 2**.

Negative-stain EM and data quantification

Freshly purified dynein was diluted in GF150 buffer to a final concentration of 14.3 nM in
the presence of 0.1 mM ATP and subsequently evaluated using negative-stain electron
microscopy (EM). A 4 μL aliquot of the sample was applied to glow-discharged carbon
film grids (Electron Microscopy Sciences) and stained with 2% uranyl acetate. The grids
were then imaged using a 120 kV Talos L120C electron microscope. Micrographs were
manually acquired at a magnification of 45,000x. More than 40 micrographs were
collected per experiment for each condition to obtain sufficient particles for statistical
analysis.

Samples for statistical analysis of the Phi ratio were prepared with the following molar
ratios: dynein: Lis1(dimer) at 1:0, 1:1, and 1:2; dynein: Nde1(dimer) at 1:0, 1:1 and 1:2;
dynein : Lis1 : Nde1 at 1:0:0, 1:1:1 and 1:2:2. The mixtures were incubated on ice for 90
minutes and subsequently subjected to negative staining. Each experimental group was
accompanied by its respective control. The assay was repeated independently three or
more times, using different batches of freshly purified protein.

Micrographs were processed using cryoSPARC, including blob picking, micrograph
extraction, and 2D classification. Briefly, micrographs from each experimental group
were merged, and particle picking was performed using templates of Phi and open
dynein. The particle diameter was set to 750 Å, and the distance cutoff for dynein
particles was 400 Å to optimize particle selection. All particles were extracted, followed
by three rounds of 2D classification. Phi and open dynein particles were identified from
the 2D classification and traced back to the corresponding micrographs for each
condition, where Phi and open particles were quantified. To calculate the normalized
fraction of Phi, the total number of Phi and paired open dynein particles was determined
for micrographs under a given condition. The ratio of Phi particles in the dynein-alone
condition (N(Phi) / N(Phi + open)) was defined as proportion A, serving as the control for
each group. The ratio of Phi particles in each experimental condition, excluding the
dynein-alone control, was defined as proportion B (N(Phi) / N(Phi + open)). The
normalized fraction of Phi was calculated as B/A, and GraphPad Prism was used to plot
the normalized fraction of Phi.

Cryo-EM sample preparation

For the dynein, Lis1, and Nde1 sample, Lis1 and Nde1 were incubated at a 1:1 molar
ratio for 30 minutes on ice. Freshly purified dynein, at a concentration of 2 mg/mL, was
then added to the Lis1-Nde1 complex at a 1:2:2 molar ratio and incubated on ice for 5
minutes, with 5 mM ATP added immediately prior to freezing. For the dynein and Lis1
complex, dynein (2 mg/mL) was incubated with Lis1 for 90 minutes on ice, and 5 mM
ATP was added just before vitrification.

For vitrification, 3.5 μL of the prepared sample was applied to glow-discharged
Quantifoil holey carbon grids (R2/1, 300 mesh gold), which were treated for 45 seconds

at 25 mA using a GloQube Glow Discharge system (Quorum Technologies). The grids
were blotted for 2.5 to 4.5 seconds at 4°C and 100% humidity, then vitrified by plunging
into liquid ethane using a Vitrobot Mark IV (Thermo Fisher Scientific).

Cryo-EM data collection

Data were collected at the Yale ScienceHill Cryo-EM facility using a Glacios microscope
(Thermo Fisher Scientific) operated at 300 keV and equipped with a K3 detector. Data
collection was facilitated by SerialEM software, targeting a defocus range of -1.2 µm to -
2.6 µm. Four exposures per hole were recorded as movies, comprising 40 frames each,
with a total electron dose of 40 e$^-$/Å². A total of 7,128 movies were collected for the
Dynein-Lis1-Nde1 condition, while 16,558 movies were collected for the Dynein-Lis1
condition.

Cryo-EM data processing

Cryo-EM movies were pre-processed using CryoSPARC Live, which included patch
motion correction and patch CTF estimation. The processing workflows are illustrated in
Extended Data Fig. 4 and 5. The statistics are summarized in **Table 1**.

For the dynein-Lis1 condition dataset, particles were picked using the blob picker,
extracted with a box size of 512 pixels, and downscaled to 128 pixels with a pixel size of
3.456 Å. In total, 7,254,515 particles were extracted. The iterative 2D classification was
performed to filter the particles, resulting in the selection of 204,005 high-quality
particles for ab initio reconstruction. Initial maps for the dynein single motor domain and
the Phi^L-Lis1 motor domains were identified. The map of the Phi^L-Lis1 motor domains
was subsequently used for heterogeneous refinement of all original particles. The
original particles were divided into four subsets, each subjected to heterogeneous
refinement (4 classes). Three rounds of heterogeneous refinement were performed,
updating the reference each time, ultimately identifying the Phi and Phi^L-Lis1 motor
domains. The Phi and Phi^L-Lis1 motor domains were merged separately and extracted
from micrographs using a box size of 512 pixels, which was then binned to 384 pixels,
resulting in a pixel size of 1.1573 Å. Two rounds of heterogeneous refinement were
conducted to exclude junk particles. High-quality subsets were selected for
homogeneous refinement, followed by two rounds of CTF refinement and local
refinement. The Phi^L-Lis1 motor domains achieved a resolution of 2.86 Å, and the Phi
motor domains reached a resolution of 2.71 Å, exhibiting C2 symmetry.

To reconstitute the tails of the Phi^L-Lis1 and Phi motor domains, the particles were
recentered at the tail and then extracted from micrographs using a box size of 512
pixels, binned to 256 pixels, yielding a pixel size of 1.736 Å/px. Following this,
heterogeneous refinement was applied to filter the particles, and high-quality subsets
were selected for homogeneous refinement. The overall tail resolutions reached 4.21 Å

for the Phi tail and 4.05 Å for the Phi^L-Lis1 tail. Four masks were devised to cover the
tail segments, which were divided into the NDD, left, right, and neck regions. Maps of
local refinement using these masks were integrated, referring to the consensus map of
the tail. The full-length map was assembled by stitching together the tail and motor
domains in ChimeraX, corresponding to Phi and Phi^L-Lis1, respectively.

For the Lis1-Nde1-dynein condition dataset, the process closely mirrored that of the
dynein-Lis1 condition described above. Briefly, the blob picker identified 2,393,279
particles. Following extraction from the micrographs and iterative 2D classification,
204,005 particles were selected for initial map generation through ab initio
reconstruction. An initial map for the Phi^L-Lis1 motor domains were obtained, which
were then subjected to iterative heterogeneous refinement using all particles with the
Phi^L-Lis1 initial map as a reference. However, the Phi motor domains did not appear in
the heterogeneous refinement, even when the verified Phi map from this study was
utilized as a reference. Subsequently, the Phi^L-Lis1 domain map was re-extracted from
the micrographs and underwent two rounds each of heterogeneous refinement, CTF
refinement, and local refinement, achieving a resolution of 2.88 Å. The consensus map
of the tail reached a resolution of 6.22 Å after recentering and re-extracting the tail
region. Masks and local refinement were employed to enhance the local resolution of
the tail. The full-length Phi^L-Lis1 structure was reconstituted by integrating the
composite tail map and motor map in ChimeraX.

Model building and refinement

For model building, previously reported structures 9BLY[19], 9BLZ[19], and 8FDT[38] were
utilized as the initial models for the full-length and motor domains of Phi and Phi^L-Lis1.
The individual domains, including the tail, single motor and Lis1 dimer, were extracted
from the 9BLY, 9BLZ, and 8FDT, and rigid-body fitting into the Cryo-EM maps were
performed using UCSF ChimeraX. The models were then manually constructed in
COOT[65,66], and followed by real-space refinement in Phenix[67]. The quality of the refined
models was assessed using the MolProbity integrated into Phenix, with the statistics
reported in **Table 1**.

Single-molecular motility assay

Fluorescent imaging was conducted using a custom-built, multicolor objective-type TIRF
microscope based on a Nikon Ti-E microscope body. It was equipped with a 100X
magnification, 1.49 N.A. apochromatic oil-immersion objective (Nikon) and a Perfect
Focus System. Fluorescence signals were captured by an electron-multiplied charge-
coupled device camera (Andor, Ixon EM+, 512 × 512 pixels), with an effective pixel size
of 160 nm after magnification. Probes such as Alexa488/GFP/mNeonGreen, LD555,
and LD655 were excited by 488 nm, 561 nm, and 633 nm laser beams (Coherent),
coupled to a single-mode fiber (Oz Optics), and their emissions were filtered using

525/40, 585/40, and 697/75 bandpass filters (Semrock), respectively. The entire system
was controlled via MicroManager 1.4 software.

Biotin-PEG treated flow chambers were treated with 5 mg/ml streptavidin for 2 minutes,
followed by washing with MB buffer (30 mM HEPES pH 7.0, 5 mM $MgSO_4$, 1 mM EGTA,
1 mg/ml casein, 0.5% pluronic acid, 0.5 mM DTT, and 1 µM Taxol). Biotinylated MTs
were then added to the chamber for 2 minutes and washed again with MB buffer.
Proteins were prepared by diluting them to the desired concentrations in MB buffer. For
DDRNL complex assembly, a mixture of 10 nM dynein, 150 nM dynactin, 50 nM BicDR1,
200nM Lis1 and 10nM Nde1 was incubated on ice for 15 minutes, then diluted tenfold
into imaging buffer (MB buffer containing 0.1 mg/ml glucose oxidase, 0.02 mg/ml
catalase, 0.8% D-glucose, and 2 mM ATP) and introduced to the flow chamber. Motility
was observed and recorded for 40 seconds.

Data analysis for single-molecular motility assay

Single-molecule motility of the DDR complex was captured for 200 frames per imaging
area and analyzed as kymographs made in FIJI. Run frequency was determined by
counting the number of processive BicDR1 molecules on each MT, then dividing this
number by the MT length and the total data collection time, with a custom MATLAB
script. The p-values for the two-tailed Student's t-test were determined in Excel.

Data availability

Cryo-EM Density maps and models have been deposited in the Electron Microscopy
Data Bank and Protein Data Bank as follows: In the dynein and Lis1 condition: PDB-
XXX/EMD-XXX for full length Phi and PDB-XXX/EMD-XXX for motor domains of Phi;
PDB-XXX/EMD-XXX for full length Phi^L-Lis1 and PDB-XXX/EMD-XXX for motor
domains of Phi^L-Lis1. In the dynein, Lis1 and Nde1 condition: PDB-XXX/EMD-XXX for
full-length Phi^L-Lis1 and XXX/EMD-XXX for motor domains of Phi^L-Lis1.

Acknowledgement

We are very grateful to members of the Zhang and Yildiz laboratories for their valuable
discussions. This work was funded by the NIH/NIGMS (GM136414 to A.Y., and
GM142959 to K.Z.) and in part by a Collaboration Development Award Program (to K.Z.)
from the Pittsburgh Center for HIV Protein Interactions (U54AI170791). The Cryo-EM
data were collected at the Yale ScienceHill Cryo-EM facility. We would like to thank
Jianfeng Lin and Kaifeng Zhou for assistance with the data collection. Yale Cryo-EM
Resource is funded in part by the NIH grant S10OD023603 awarded to Frederick J.
Sigworth.

Author contributions

K.Z. and A.Z. designed the study. J.Y. expressed and purified dynein, Lis1 and Nde1
proteins for EM. J.Y. and P.C. prepared the cryo-EM samples, collected and processed
the data, and built the PDB models. P.C. and J.Y. processed negative stain EM data
and quantified the particle numbers. Y.Z. performed Lis1 mutagenesis, protein
preparation, TIRF and mass photometry assays. J.Y., P.C., Y.Z., K.Z., and A.Y.
analyzed the data and prepared figures. J.Y., Y.Z., P.C., K.Z., and A.Y. wrote the
manuscript with input from all authors.

Competing Interests

The authors declare no competing interests.

Figures

Fig. 1. Nde1 promotes Lis1 binding to Phi dynein and cooperatively releases dynein autoinhibition. a, Schematic of dynein activation. Nde1 tethers Lis1 to dynein and promotes Lis1-mediated formation of the active dynein-dynactin-adaptor-Lis1 (DDAL) complex. Intermediate states between the association of Lis1-Nde1 and opening of dynein are unknown. **b**, MP shows that Lis1 alone slowly binds to dynein in

tens of minutes, whereas Nde1 promotes more rapid and efficient binding of Lis1 to
dynein. Dynein, Lis1 and Nde1 were included at a 1:2:2 ratio. Only one Lis1 dimer is
tethered to one dynein. Solid curves represent a fit to multiple Gaussians to predict the
average mass and percentage of each population. **c**, A representative image of dynein
motors captured using negative stain electron microscopy. Without Lis1 and Nde1,
dynein is distributed almost equally between Phi and open conformations. The
percentage of Phi was quantified after incubating dynein with Lis1, Nde1, or both
proteins for 90 min. The percentage of Phi (**d**) and the relative change of Phi (**e**) in the
presence and absence of Lis1 and Nde1 (mean ± s.d.; from left to right, n=14, 6, 6, 3, 4,
4, 4 from three or more independent experiments). P values are calculated from a two-
tailed *t* test. The control (Ctrl) of panel e represents dynein alone from the three groups
of panel d (Ctrl) normalized to 100%.

Fig. 2. The structure of Phi^L-Lis1 complex. a, (Top) Full length human dynein was incubated with Lis1 with a molar ratio of 1:2 on ice for 90 min before flash frozen. (Bottom) Cryo-EM maps of Phi and Phi^L-Lis1 complex structures in front view and percentages of particles with these two conformations (open dynein excluded). **b,** (Top) Dynein was incubated with Lis1 and Nde1 with a molar ratio of 1:2:2 on ice for 5 min before flash frozen. (Bottom) The structure of Phi^L-Lis1 obtained under this condition is shown in front view and side view. All particles were classified into the Phi^L-Lis1 conformation and Phi conformation could not be detected. **c,** Model of the Phi^L-Lis1 motor domains is shown in three views. One Lis1 dimer is clamped between MD-A and MD-B of Phi^L-Lis1 on the front side, while the back side of the motor domains is vacant,

609   with no Lis1 bound. The top view reveals that the linker regions (purple) of MD-A and -B

610   interact with each other.

Fig. 3. Lis1 induces a transition in the Phi conformation to accommodate its binding. a, Docking of a Lis1 monomer between dynein motor rings shows a clash with the rigid canonical Phi MD-A. **b**, A Lis1 dimer bound to dynein motors on the front side is compatible with the coordinated Phi^L structure. **c**, (Top) The superimposition of the Phi and Phi^L-Lis1 motor domains. Phi and Phi^L-Lis1 are colored with grey and rainbow respectively. (Bottom) Comparison of the motor domains between Phi^L-Lis1 and canonical Phi. Lis1 is hidden for clarity. Vectors present interatomic distance of pairwise Cα atoms between the Phi to Phi^L-Lis1 structures. Lis1 binding induces a slight rotation of the dynein motor domains from the Lis1-bound (front) side toward the back, resulting in a larger groove at the front side compared to the smaller groove at the back. **d**, Overlay of the motor domain maps of Phi and Phi^L-Lis1 with MD-A aligned at a lower contour level. The rotation of Phi^L-Lis1 motor domain induces rotation in the neck region, showing a slight unwinding trend, which contributes to the formation of a smaller groove on the back side. **e**, Docked Lis1 on the back side of Phi^L-Lis1 causes a significant clash with MD-B.

628
Fig. 4. Novel interactions identified between Lis1$_{ring}$ and MD-A in Phi^L-Lis1. **a**,
Cryo-EM density map highlighting the subdomains involved in the interface between
Lis1 and the dynein motor. Lis1 and subdomains of MD-A and MD-B are colored
separately. Novel interfaces between MD-A and Lis1$_{ring}$ are marked by dash rectangle
and enlarged in panels d, e, and f. **b**, Comparison of MD-A (grey) and -B (rainbow) of
Phi^L-Lis1. **c**, Comparison of MD-B (rainbow) with the Lis1-bound structure of the motor
domain of human dynein (PDB: 8FDT, grey). Representative interactions located at
linker-Lis1$_{ring}$ (**d**), AAA6-Lis1$_{ring}$ (**e**), and AAA5-Lis1$_{ring}$ (**f**) interface are shown with stick

mode, colored according to their respective subdomains. **g**, MP profiles illustrate the
interaction of Nde1 (N) with WT Lis1 (L) and Lis1 mutants (Lis1[AAA5], Lis1[AAA6], and
Lis1[linker]). Nde1 interacts with one (NL) or two (NL$_2$) Lis1 dimers. Lis1[AAA5] exhibits a
reduced binding percentage to Nde1. **h**. MP shows the binding of Lis1 mutants to
dynein with or without Nde1. Dynein, Lis1 and Nde1 were incubated for 2 minutes at
1:2:2 ratio (D: dynein only, DL: one dynein and one Lis1, DL$_2$N: one dynein, two Lis1,
and one Nde1). **i**, Representative kymographs show the motility of WT DDR complexes
with or without Nde1 and Lis1. **j**, Run frequency of WT DDR with or without Nde1 and
Lis1 (mean ± s.d.; n = 20 MTs for each condition; statistics from two independent
experiments). Results were normalized to the -Lis1, -Nde1 condition.

Fig. 5. Model for the initial stage of dynein activation by Nde1 and Lis1. Step-i, Nde1 tethers Lis1 to Phi dynein and forms a transient Phi^L-Lis1-Nde1 complex. **Step-ii,** Nde1 dissociates spontaneously from dynein, leading to the formation of Phi^L-Lis1. Without Nde1, Lis1 binding to Phi dynein and formation of Phi^L-Lis1 becomes a less efficient and slow process. The binding of Lis1 induces a slight backward rotation of the two motor rings and a twist in the neck region, likely leading an unwinding trend in the tail and contributing to dynein opening. **Step-iii**, Phi^L-Lis1 transitions to open dynein-Lis1 assisted by Nde1. **Step-iv**, Lis1 promotes processive DDA complex assembly by interacting with p150 and DIC-N through its LisH domain. The dashed line indicates the possibility that Lis1 in the Phi^L-Lis1 complex promotes dynactin recruitment and DDA assembly without switching to the open conformation.

Description	In dynein and Lis1 condition				In Nde1, Lis1 and dynein condition	
	Composite map of full-length Phi	Composite map of full-length Phi^L-Lis1	Phi motor	Phi^L-Lis1 motor	Composite map of full-length Phi^L-Lis1	Phi^L-Lis1 motor
PDB code	xxx	xxx	xxx	xxx	xxx	xxx
Data Collection and Processing						
Facility	Yale ScienceHill-Cryo-EM facility				Yale ScienceHill-Cryo-EM facility	
Microscope	Glacios				Glacios	
Voltage (kV)	200				200	
Camera	K3				K3	
Magnification	45k				45k	
Pixel Size (Å)	0.432 (super resolution)				0.432 (super resolution)	
Total Electron Exposure (e-/A^2)	40				40	
Defocus Range (μm)	1.5-2.7				1.5-2.7	
Symmetry Imposed	C1	C1	C2	C1	C1	C1
Num of mics	16,558				7,128	
Initial Particles	7,254,515				2,393,279	
Final Particles	103,097	127,963	160,539	215,049	61,684	127,611
Refinement						
Initial models	9BLY ab initio	9BLY 8FDT ab initio	9BLZ ab initio	8FDT ab initio	9BLY 8FDT ab initio	8FDT ab initio
Map pixel size	1.7360	1.7360	1.1573	1.1573	1.736	1.157
Map Resolution (Å) (FSC 0.143)	4.50	4.50	2.71	2.86	5.0	2.88
Map sharpening B-factor (Å^2)	/	/	72.3	69.6	/	51.7
Model Composition						
Non-hydrogen atoms	89,388	94,497	47418	21,370	94,497	52,508
Protein residues	11,061	11,704	5784	6514	11,704	6,515
Ligands	MG:4 ATP:2 ADP:6	MG:4 ATP:2 ADP:6	MG:4 ATP:2 ADP:6	MG:4 ATP:2 ADP:6	MG:4 ATP:2 ADP:6	MG:4 ATP:2 ADP:6
Model vs. Data						
FSC Map to Model (Å) (FSC 0.5)	3.5	3.8	3.0	3.1	3.5	3.4
Correlation coefficient (mask)	0.60	0.59	0.86	0.87	0.51	0.84
B factors (A^2)						
Protein	117.05	112.65	14.13	38.69	260.48	52.33
Ligand	75.90	79.64	1.76	14.44	69.27	32.47
R.m.s deviation						
Bond length (Å)	0.006	0.004	0.003	0.003	0.002	0.002
Bond angles (_)	0.683	0.693	0.548	0.527	0.544	0.548
Validation						
Molprobity score	1.87	1.88	1.40	1.41	1.75	1.45
Clashscore	13.26	13.07	7.31	7.33	10.62	8.06
Rotamer outliers (%)	0.00	0.00	0.00	0.00	0.00	0.02
Ramachandran plot						
Outliers (%)	0.09	0.07	0.00	0.00	0.05	0.00
Allowed (%)	3.52	3.74	1.50	2.05	3.23	2.05
Favored (%)	96.39	96.20	98.50	97.95	96.72	97.95
Rama-Z (whole)	0.62	0.53	1.60	1.72	0.92	1.86

660 **Table 1. Cryo-EM data collection, refinement, and validation statistics.**

Construct	Vector	Source	Figures
SNAPf-DYNH1C1-IC2C-LIC2-Robl1-Tctex1-LC8	pOmniBac-pIDC	Schlager et al., 2014	F1, 2, 3, 4, EDF2, 3, 4, 5, 6, 8, 9, 12
SNAPf-DYNH1C1$^{R1567E, K1610E}$-IC2CLIC2-Robl1-Tctex1-LC8	pOmniBac-pIDC	Zhang et al., 2017	EDF1, 7, 13
BicDR1-SNAPf	pOmniBac	Urnavicius et al. 2018	F4, EDF13
Lis1-SNAPf	pOmniBac	Elshenawy et al., 2020	F1, 2, 3, 4, EDF1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13
Lis1$^{E300K, K303E, S304R, M329A}$-SNAPf	pOmniBac	This study	F4, EDF11
Lis1$^{N203K, D205K, Q222A}$-SNAPf	pOmniBac	This study	F4, EDF11
Lis1$^{M172K, D192K, Y225A, C226D}$-SNAPf	pOmniBac	This study	F4, EDF11
Nde1$^{1-190}$-SNAPf	pOmniBac	Zhao et al., 2023	F1, 4, EDF1, 2, 7, 11, 13

Supplementary Table 1. The list of protein constructs used in this study. Dynein
chains were codon-optimized for *Spodoptera frugiperda* (Sf9) expression and inserted
into the pOmniBac backbone. Nde1, Lis1, and BicDR1 constructs were cloned into the
pOmniBac backbone. Constructs were tagged with an N-terminal 6xHis-ZZ-TEV site for
affinity purification and TEV protease cleavage during protein purification. The SNAPf
tag was inserted for labeling the proteins with fluorescent dyes (F: Figure, EDF:
Extended Data Figure).

Fig.	Sample	Complex	Expected (kDa)	Measured (kDa)	%
F 1b	WT Dyn	WT Dyn	1376	1326 ± 49	100[*]
	WT Dyn + Lis1 0 min	WT Dyn	1376	1350 ± 65	100[*]
	WT Dyn + Lis1 10 min	WT Dyn	1376	1344 ± 55	80[*]
		WT Dyn + Lis1	1546	1521 ± 129	20[*]
	WT Dyn + Lis1 30 min	WT Dyn	1376	1364 ± 60	65[*]
		WT Dyn + Lis1	1546	1544 ± 68	35[*]
	WT Dyn + Lis1 60 min	WT Dyn	1376	1359 ± 48	63[*]
		WT Dyn + Lis1	1546	1528 ± 40	37[*]
	WT Dyn + Lis1 + Nde1 0 min	WT Dyn	1376	1356 ± 64	31[*]
		WT Dyn + Lis1	1546	1525 ± 70	69[*]
	WT Dyn + Nde1	WT Dyn	1376	1330 ± 83	100[*]
F 4g	WT Lis1	WT Lis1	133	137 ± 17	70
	Lis1^{AAA5}	Lis1^{AAA5}	133	141 ± 26	87
	Lis1^{AAA6}	Lis1^{AAA6}	133	141 ± 26	87
	Lis1^{Linker}	Lis1^{Linker}	133	142 ± 31	90
	Nde1$^{1-190}$ + WT Lis1	Nde1$^{1-190}$	84	85 ± 18	31
		WT Lis1	133	142 ± 15	29
		1 Nde1$^{1-190}$ + 1 WT Lis1	217	231 ± 20	28
		1 Nde1$^{1-190}$ + 2 WT Lis1	340	371 ± 44	12
	Nde1$^{1-190}$ + Lis1^{AAA5}	Nde1$^{1-190}$	84	90 ± 17	37
		Lis1^{AAA5}	133	142 ± 13	53
		1 Nde1$^{1-190}$ + 1 Lis1^{AAA5}	217	229 ± 48	10
	Nde1$^{1-190}$ + Lis1^{AAA6}	Nde1$^{1-190}$	84	82 ± 17	25
		Lis1^{AAA6}	133	141 ± 15	30
		1 Nde1$^{1-190}$ + 1 Lis1^{AAA6}	217	233 ± 30	29
		1 Nde1$^{1-190}$ + 2 Lis1^{AAA6}	340	373 ± 16	9
	Nde1$^{1-190}$ + Lis1^{Linker}	Nde1$^{1-190}$	84	88 ± 16	24
		Lis1^{Linker}	133	143 ± 14	26
		1 Nde1$^{1-190}$ + 1 Lis1^{Linker}	217	229 ± 18	30
		1 Nde1$^{1-190}$ + 2 Lis1^{Linker}	340	372 ± 15	12
F 4h	WT Dyn	WT Dyn	1376	1367 ± 93	100[*]
	WT Dyn + WT Lis1	WT Dyn	1376	1387 ± 52	82[*]
		WT Dyn + WT Lis1	1509	1530 ± 39	18[*]
	WT Dyn + WT Lis1 + Nde1$^{1-190}$	WT Dyn + WT Lis1	1509	1514 ± 67	100[*]
	WT Dyn + Lis1^{AAA5}	WT Dyn	1376	1367 ± 94	100[*]
	WT Dyn + Lis1^{AAA5} + Nde1$^{1-190}$	WT Dyn	1376	1361 ± 57	57[*]
		WT Dyn + Lis1^{AAA5}	1509	1581 ± 76	43[*]
	WT Dyn + Lis1^{AAA6}	WT Dyn	1376	1356 ± 60	100[*]
	WT Dyn + Lis1^{AAA6} + Nde1$^{1-190}$	WT Dyn	1376	1356 ± 75	77[*]
		WT Dyn+2*Lis1^{AAA6} +	1716	1680 ± 77	23[*]

			1376	1349 ± 56	
		Nde1[1-190]			
	WT Dyn + Lis1[Linker]	WT Dyn	1376	1349 ± 56	100[*]
	WT Dyn + Lis1[Linker] + Nde1[1-190]	WT Dyn + Lis1[Linker]	1509	1464 ± 75	100[*]
EDF 1	mtDyn	mtDyn	1376	1362 ± 46	100[*]
	mtDyn + Lis1 + Nde1	mtDyn	1376	1386 ± 77	32[*]
		mtDyn + Lis1	1546	1525 ± 74	68[*]
	mtDyn + Lis1	mtDyn	1376	1411 ± 73	50[*]
		mtDyn + Lis1	1546	1540 ± 57	50[*]
EDF 2a	WT Dyn	WT Dyn	1376	1396 ± 241	100[*]
	WT Dyn + Lis1	WT Dyn	1376	1414 ± 85	100[*]
	WT Dyn + Nde1	WT Dyn	1376	1371 ± 68	100[*]
	WT Dyn + Lis1 + Nde1	WT Dyn	1376	1415 ± 85	30[*]
		WT Dyn + Lis1	1546	1580 ± 63	70[*]
EDF 2b	WT Dyn	WT Dyn	1376	1347 ± 278	100[*]
	WT Dyn + Lis1	WT Dyn	1376	1395 ± 167	100[*]
	WT Dyn + Nde1	WT Dyn	1376	1376 ± 78	100[*]
	WT Dyn + Lis1 + Nde1	WT Dyn	1376	1391 ± 65	48[*]
		WT Dyn + Lis1	1546	1585 ± 133	52[*]
EDF 2c	WT Dyn	WT Dyn	1376	1381 ± 61	100[*]
	WT Dyn + Lis1	WT Dyn	1376	1371 ± 136	100[*]
	WT Dyn + Nde1	WT Dyn	1376	1377 ± 113	100[*]
	WT Dyn + Lis1 + Nde1	WT Dyn	1376	1409 ± 75	45[*]
		WT Dyn + Lis1	1546	1584 ± 47	55[*]
EDF 2d	WT Dyn	WT Dyn	1376	1411 ± 108	100[*]
	WT Dyn + Lis1	WT Dyn	1376	1377 ± 124	100[*]
	WT Dyn + Nde1	WT Dyn	1376	1366 ± 130	100[*]
	WT Dyn + Lis1 + Nde1	WT Dyn	1376	1383 ± 56	28[*]
		WT Dyn + Lis1	1546	1561 ± 51	72[*]
EDF 2e	WT Dyn	WT Dyn	1376	1396 ± 304	100[*]
	WT Dyn + Lis1	WT Dyn	1376	1403 ± 76	100[*]
	WT Dyn + Nde1	WT Dyn	1376	1379 ± 54	100[*]
	WT Dyn + Lis1 + Nde1	WT Dyn	1376	1402 ± 53	27[*]
		WT Dyn + Lis1	1546	1581 ± 46	73[*]

Supplementary Table 2. The parameters of a multi-Gaussian fit of MP measurements. Dynein (Dyn), Lis1, and Nde1 were mixed at 1:2:2 ratio if present and diluted into 5-20 nM. Measured mass and percentage represent the center (mean ± s.d.) and the percent area of the corresponding Gaussian peak ([*]percentages are normalized so that the sum of all dynein included peaks are 100%). Expected mass corresponds to the dimeric forms of Lis1, Nde1, and Dyn (F: Figure, EDF: Extended Data Figure). Lis1 used in F 1b, and EDF 1 contains a ZZ tag, and its expected MW is 170 kDa.

Supplementary Video Legends

Supplementary Video 1. Full-length human dynein in Phi^L conformation, bound to a
Lis1 dimer and displaying the newly identified interface with Lis1.

Supplementary Video 2. Single-molecule motility recordings of WT DDR complexes, in
the presence or absence of Nde1, WT Lis1, and Lis1 mutants. The fluorescence signal
originates from BicDR1-mNeonGreen.

Extended Data Figures

Extended Data Fig. 1. MP analysis of Nde1's effect on Lis1 binding to open dynein.
MP shows that Nde1 does not promote increased Lis1 binding to open dynein
compared to the Lis1-alone condition, indicating that Lis1 can efficiently bind to open
dynein and Nde1 does not enhance Lis1's interaction with open dynein.

Extended Data Fig. 2. MP analysis of nucleotide conditions on Nde1 and Lis1 binding to WT dynein. MP shows that under apo buffer (**a**), 0.1 mM ATP (**b**), ADP (**c**), ATP.vi (**d**), and AMPPNP (**e**) conditions, Nde1 promotes Lis1 binding to dynein, forming a 1:1 dynein-Lis1 (DL) complex. The nucleotide condition does not affect Nde1's ability to tether Lis1 to dynein. Importantly, the formation of dynein-Nde1, dynein-Lis1-Nde1

696   and 1:2 dynein-Lis1 complexes were not observed. In the Lis1-alone condition, no
697   significant DL complex was formed immediately.

Extended Data Fig. 3. Workflow for negative-stain EM data processing. a, Representative micrographs for dynein alone (42 micrographs), dynein-Lis1 at 1:1 (41 micrographs), and dynein-Lis1 at 1:2 (47 micrographs) molar ratios from batch #1. **b**, Particle picking from representative micrographs in each dataset using a template matching approach based on Phi and open dynein models (particle diameter: 750 Å, distance cutoff: 400 Å). **c**, Three rounds of 2D classification were performed after extracting all particles (box size: 960 Å × 960 Å), yielding class averages of Phi dynein, open dynein motors, single motors, and junk particles. **d**, Final classified 2D averages showing Phi dynein, two-motor open dynein, and single-motor open dynein. The particle numbers for each group were counted, and single motors were considered as open dynein by dividing the total number of particles by two.

710
Extended Data Fig. 4. Cryo-EM data processing for the dynein-Lis1 dataset. a, A
representative cryo-EM micrograph and the flowchart of cryo-EM data processing. **b**,

Fourier Shell Correlation (FSC) curves showing the final resolution estimates for the
motor domains of the Phi (2.71 Å) and Phi^L-Lis1(2.86 Å) datasets.

Extended Data Fig. 5. Cryo-EM data processing for the Nde1-Lis1-dynein dataset.
a, A representative cryo-EM micrograph and the flowchart of cryo-EM data processing.
b, Fourier Shell Correlation (FSC) curve showing the final resolution estimate for the
motor domains of the Phi^L-Lis1 (2.88 Å) dataset.

721
Extended Data Fig. 6. Comparison of local resolution, nucleotide binding in AAA1,
AAA3, AAA4, and sensor-I loop conformation in MD-A of the Phi and Phi[L]-Lis1.
Local resolution, and nucleotide binding states in MD-A at AAA1, AAA3, and AAA4 of
the Phi (**a**) and Phi[L]-Lis1 (**b**). MD-A and -B share the same nucleotide binding in AAA1,
AAA3, and AAA4 across both the Phi and Phi[L]-Lis1. The sensor-I loop adopts almost
the same conformation in MD-A (or -B) of both Phi (**c**) and Phi[L]-Lis1 (**d**), indicating that
Lis1 binding does not affect phosphate release. The color scheme is the same with Fig.
4.

Extended Data Fig. 7. Comparison of Nde1's effect on Lis1 binding to the open

dynein. The particle numbers for open dynein-Lis1 and open dynein alone were

quantified in both the dynein-Lis1 and dynein-Lis1-Nde1 datasets. These results

indicate that Nde1 does not promote Lis1 binding to the open dynein motor.

a Linker-Lis1$_{ring}$

b AAA5-Lis1$_{ring}$
      AAA6-Lis1$_{ring}$

Extended Data Fig. 8. Density quality at the dynein MD-A and Lis1 interface in Phi^{L}-Lis1. a, Flexible density at the linker-Lis1$_{ring}$ interface, indicating dynamic interactions in this region. **b**, Well-defined density at the AAA5-Lis1$_{ring}$ and AAA6-Lis1$_{ring}$ regions, showing compact and stable interactions. The color scheme for the motor domains is consistent with Fig. 4.

Extended Data Fig. 9. Sequence alignment of Lis1 homologs among multiple
species. Sequence alignment of Lis1 proteins among *Homo sapiens* (Hs Lis1),
Xenopus laevis (*Xl* Lis1), *Gallus gallus* (*Gg* Lis1), *Danio rerio* (*Dr* Lis1), and
Saccharomyces cerevisiae (*Sc* Lis1). The secondary structure elements are placed on
the top of the alignment. Strictly conserved residues are highlighted in shaded red

boxes, while less-conserved residues are shown in open red boxes. In these open red
boxes, red font indicates residues with similar polarity and high conservation, whereas
black font represents residues with low similarity. The colored triangles represent key
residues involved in interactions at the linker-Lis1$_{ring}$ (purple), AAA5-Lis1$_{ring}$ (green), and
AAA6-Lis1$_{ring}$ (orange) interfaces in Phi^{L}-Lis1. The black triangle represents reported
interactions at MD-A and Lis1 interface of modeled human Chi-Lis1 based on the yeast
Chi-Lis1[37]. *S. cerevisiae* Lis1 shows more variation compared to the vertebrate species,
suggesting greater evolutionary divergence.

**Extended Data Fig. 10. Comparison of motor domain A-B interfaces in Phi and
Phi[L]-Lis1. a**. Representative residues located on MD-A that are involved in the MD-A
and MD-B interface of Phi, showing interactions at the linker-linker, linker-AAA4, AAA4-
linker, AAA4-AAA4, AAA5-AAA5, and stalk-stalk interfaces. **b**, Representative residues

located on MD-A of the Phi^L-Lis1, involved in the MD-A and $Lis1_{ring}$ interface (including
linker-$Lis1_{ring}$, AAA6-$Lis1_{ring}$, and AAA5-$Lis1_{ring}$), and the motor domain A-B interface
(including linker-linker, AAA3-linker, linker-AAA4, AAA5-AAA5, and stalk-stalk
interfaces). Residues in both panels (a) and (b) are displayed in sphere mode. **c**,
Detailed view of the motor domain A-B interface in Phi^L-Lis1, showing key residues
involved in interactions at the linker-linker, AAA3-linker, AAA5-AAA5, and linker-AAA4
interfaces. **d**, Detailed view of motor domain A-B interface of Phi, showing key residues
at the linker-linker, linker-AAA4, AAA4-linker, AAA4-AAA4, and AAA5-AAA5 interfaces.

Extended Data Fig. 11. Nde1-Lis1 interface predicted by Alphafold. a, Predicted structure of the Nde1-Lis1 complex. Interactions involved in the Phi^L MD-A and Lis1 interface are shown on the Lis1 surface. **b**, D192 and M172 involved in AAA5-Lis1 interface also show contact with R115 of Nde1. **c**, D192K and M172K mutation of Lis1^{AAA5} show charge repulsion with R115 of Nde1. The prediction supports an overlap of the interfaces between the AAA5-Lis1 and Nde1-Lis1.

Extended Data Fig. 12. Comparison of yeast Chi-Lis1 and human Phi^L-Lis1 motor domains. a, The structure of yeast Chi-Lis1 (PDB:8DZZ)[37], showing two tail-truncated yeast dynein motor domains (grey) bound to two Lis1 dimers (colored, Chi-Lis1 1:2). **b**, Residues of MD-A that interact with Lis1$_{ring}$ are located in AAA6-Lis1$_{ring}$ and AAA5-Lis1$_{ring}$ regions and highlighted with dashed rectangle. Representative residues of MD-A involved in the canonical Lis1$_{ring}$ binding sites are located in Lis1$_{ring}$-AAA3, Lis1$_{ring}$-AAA4, Lis1$_{ring}$-AAA5 and Lis1$_{ring}$-stalk region. Interactions between MD-A and MD-B are in stalk-stalk region. Residues are displayed in sphere mode and are colored according to the subdomains in Fig. 4. **c**, Superimposition of the human Phi^L-Lis1 and yeast Chi-Lis1 structures, showing that Chi-Lis1 adopts a more expanded conformation, with larger grooves on both the front and back sides compared to the more compact Phi^L-Lis1

788   structure. Lis1 is hidden for clarity. Vectors represent interatomic distances of pairwise

789   Cα atoms between the Phi^L-Lis1 and Chi-Lis1 structures.

Reference

1. Reck-Peterson, S.L., Redwine, W.B., Vale, R.D. & Carter, A.P. The cytoplasmic dynein transport machinery and its many cargoes. *Nat Rev Mol Cell Biol* **19**, 382-398 (2018).
2. McNally, F.J. Mechanisms of spindle positioning. *J Cell Biol* **200**, 131-40 (2013).
3. Markus, S.M., Marzo, M.G. & McKenney, R.J. New insights into the mechanism of dynein motor regulation by lissencephaly-1. *Elife* **9**(2020).
4. Willemsen, M.H. et al. Mutations in DYNC1H1 cause severe intellectual disability with neuronal migration defects. *J Med Genet* **49**, 179-83 (2012).
5. Scoto, M. et al. Novel mutations expand the clinical spectrum of DYNC1H1-associated spinal muscular atrophy. *Neurology* **84**, 668-679 (2015).
6. Poirier, K. et al. Mutations in TUBG1, DYNC1H1, KIF5C and KIF2A cause malformations of cortical development and microcephaly. *Nat Genet* **45**, 639-47 (2013).
7. Guven, A., Gunduz, A., Bozoglu, T.M., Yalcinkaya, C. & Tolun, A. Novel NDE1 homozygous mutation resulting in microhydranencephaly and not microlyssencephaly. *Neurogenetics* **13**, 189-94 (2012).
8. Lipka, J., Kuijpers, M., Jaworski, J. & Hoogenraad, C.C. Mutations in cytoplasmic dynein and its regulators cause malformations of cortical development and neurodegenerative diseases. *Biochemical Society Transactions* **41**, 1605-1612 (2013).
9. Eschbach, J. & Dupuis, L. Cytoplasmic dynein in neurodegeneration. *Pharmacol Ther* **130**, 348-63 (2011).
10. Urnavicius, L. et al. Cryo-EM shows how dynactin recruits two dyneins for faster movement. *Nature* **554**, 202-206 (2018).
11. Singh, K. et al. Molecular mechanism of dynein-dynactin complex assembly by LIS1. *Science* **383**, eadk8544 (2024).
12. Okada, K. et al. Conserved roles for the dynein intermediate chain and Ndel1 in assembly and activation of dynein. *Nature Communications* **14**(2023).
13. Chaaban, S. & Carter, A.P. Structure of dynein-dynactin on microtubules shows tandem adaptor binding. *Nature* (2022).
14. Zhang, K. et al. Cryo-EM Reveals How Human Cytoplasmic Dynein Is Auto-inhibited and Activated. *Cell* **169**, 1303-1314 e18 (2017).
15. Canty, J.T. & Yildiz, A. Activation and Regulation of Cytoplasmic Dynein. *Trends in Biochemical Sciences* **45**, 440-453 (2020).
16. Neuwald, A.F., Aravind, L., Spouge, J.L. & Koonin, E.V. AAA+: A class of chaperone-like ATPases associated with the assembly, operation, and disassembly of protein complexes. *Genome Res* **9**, 27-43 (1999).
17. Carter, A.P., Cho, C., Jin, L. & Vale, R.D. Crystal Structure of the Dynein Motor Domain. *Science* **331**, 1159-1165 (2011).
18. Canty, J.T., Tan, R., Kusakci, E., Fernandes, J. & Yildiz, A. Structure and Mechanics of Dynein Motors. *Annu Rev Biophys* **50**, 549-574 (2021).
19. Chai, P. et al. Cryo-EM Reveals the Mechanochemical Cycle of Reactive Full-length Human Dynein-1. *BioRixv* (2024).

20. Cianfrocco, M.A., DeSantis, M.E., Leschziner, A.E. & Reck-Peterson, S.L.
Mechanism and regulation of cytoplasmic dynein. *Annu Rev Cell Dev Biol* **31**, 83-
108 (2015).
21. Amos, L.A. Brain dynein crossbridges microtubules into bundles. *J Cell Sci* **93**
(Pt 1), 19-28 (1989).
22. Torisawa, T. et al. Autoinhibition and cooperative activation mechanisms of
cytoplasmic dynein. *Nature Cell Biology* **16**, 1118-+ (2014).
23. Garrott, S.R., Gillies, J.P. & DeSantis, M.E. Nde1 and Ndel1: Outstanding
Mysteries in Dynein-Mediated Transport. *Frontiers in Cell and Developmental*
Biology **10**(2022).
24. Bradshaw, N.J., Hennah, W. & Soares, D.C. NDE1 and NDEL1: twin
neurodevelopmental proteins with similar 'nature' but different 'nurture'. *Biomol*
Concepts **4**, 447-64 (2013).
25. Monda, J.K. & Cheeseman, I.M. Nde1 promotes diverse dynein functions through
differential interactions and exhibits an isoform-specific proteasome association.
Molecular Biology of the Cell **29**, 2336-2345 (2018).
26. Zhao, Y., Oten, S. & Yildiz, A. Nde1 promotes Lis1-mediated activation of dynein.
Nat Commun **14**, 7221 (2023).
27. Shu, T.Z. et al. Ndel1 operates in a common pathway with LIS1 and cytoplasmic
dynein to regulate cortical neuronal positioning. *Neuron* **44**, 263-277 (2004).
28. Stehman, S.A., Chen, Y., McKenney, R.J. & Vallee, R.B. NudE and NudEL are
required for mitotic progression and are involved in dynein recruitment to
kinetochores. *Journal of Cell Biology* **178**, 583-594 (2007).
29. Wang, S.S. et al. Nudel/NudE and Lis1 promote dynein and dynactin interaction
in the context of spindle morphogenesis. *Molecular Biology of the Cell* **24**, 3522-
3533 (2013).
30. McKenney, R.J., Vershinin, M., Kunwar, A., Vallee, R.B. & Gross, S.P. LIS1 and
NudE Induce a Persistent Dynein Force-Producing State. *Cell* **141**, 304-314
(2010).
31. Reiner, O. & Sapir, T. LIS1 functions in normal development and disease.
Current Opinion in Neurobiology **23**, 951-956 (2013).
32. Huang, J., Roberts, A.J., Leschziner, A.E. & Reck-Peterson, S.L. Lis1 Acts as a
"Clutch" between the ATPase and Microtubule-Binding Domains of the Dynein
Motor. *Cell* **150**, 975-986 (2012).
33. Qiu, R.D., Zhang, J. & Xiang, X. LIS1 regulates cargo-adapter-mediated
activation of dynein by overcoming its autoinhibition in vivo. *Journal of Cell*
Biology **218**, 3630-3646 (2019).
34. Elshenawy, M.M. et al. Lis1 activates dynein motility by modulating its pairing
with dynactin. *Nat Cell Biology* **22**, 570-578 (2020).
35. Htet, Z.M. et al. LIS1 promotes the formation of activated cytoplasmic dynein-1
complexes. *Nature Cell Biology* **22**, 518-+ (2020).
36. McKenney, R.J. LIS1 cracks open dynein. *Nature Cell Biology* **22**, 515-517
(2020).
37. Karasmanis, E.P. et al. Lis1 relieves cytoplasmic dynein-1 autoinhibition by
acting as a molecular wedge. *Nat Struct Mol Biol* **30**, 1357-1364 (2023).

38. Ton, W.D. et al. Microtubule-binding-induced allostery triggers LIS1 dissociation from dynein prior to cargo transport. *Nature Structural & Molecular Biology* (2023).

39. Kusakci, E. et al. Lis1 slows force-induced detachment of cytoplasmic dynein from microtubules. *Nature Chemical Biology* **20**(2024).

40. Neer, E.J., Schmidt, C.J., Nambudripad, R. & Smith, T.F. The ancient regulatory-protein family of WD-repeat proteins. *Nature* **371**, 297-300 (1994).

41. Emes, R.D. & Ponting, C.P. A new sequence motif linking lissencephaly, Treacher Collins and oral-facial-digital type 1 syndromes, microtubule dynamics and cell migration. *Hum Mol Genet* **10**, 2813-20 (2001).

42. Mateja, A., Cierpicki, T., Paduch, M., Derewenda, Z.S. & Otlewski, J. The dimerization mechanism of LIS1 and its implication for proteins containing the LisH motif. *Journal of Molecular Biology* **357**, 621-631 (2006).

43. Reimer, J.M., DeSantis, M.E., Reck-Peterson, S.L. & Leschziner, A.E. Structures of human dynein in complex with the lissencephaly 1 protein, LIS1. *Elife* **12**(2023).

44. Marzo, M.G., Griswold, J.M. & Markus, S.M. Pac1/LIS1 stabilizes an uninhibited conformation of dynein to coordinate its localization and activity. *Nature Cell Biology* **22**, 559-+ (2020).

45. Geohring, I.C. et al. A nucleotide code governs Lis1's ability to relieve dynein autoinhibition. *bioRxiv* (2024).

46. Wynne, C.L. & Vallee, R.B. Cdk1 phosphorylation of the dynein adapter Nde1 controls cargo binding from G2 to anaphase. *Journal of Cell Biology* **217**, 3019-3029 (2018).

47. Lam, C., Vergnolle, M.A., Thorpe, L., Woodman, P.G. & Allan, V.J. Functional interplay between LIS1, NDE1 and NDEL1 in dynein-dependent organelle positioning. *J Cell Sci* **123**, 202-12 (2010).

48. Zhang, Y.F. et al. Nde1 is a Rab9 effector for loading late endosomes to cytoplasmic dynein motor complex. *Structure* **30**, 386-+ (2022).

49. Doobin, D.J., Helmer, P., Carabalona, A., Bertipaglia, C. & Vallee, R.B. The Role of Nde1 phosphorylation in interkinetic nuclear migration and neural migration during cortical development. *Mol Biol Cell* **35**, ar129 (2024).

50. Pei, Z. et al. The Expression and Roles of Nde1 and Ndel1 in the Adult Mammalian Central Nervous System. *Neuroscience* **271**, 119-136 (2014).

51. Derewenda, U. et al. The structure of the coiled-coil domain of Ndel1 and the basis of its interaction with Lis1, the causal protein of Miller-Dieker lissencephaly. *Structure* **15**, 1467-81 (2007).

52. Garrott, S.R. et al. Ndel1 disfavors dynein-dynactin-adaptor complex formation in two distinct ways. *J Biol Chem* **299**, 104735 (2023).

53. Ye, F. et al. DISC1 Regulates Neurogenesis via Modulating Kinetochore Attachment of Ndel1/Nde1 during Mitosis. *Neuron* **96**, 1204 (2017).

54. McKenney, R.J., Weil, S.J., Scherer, J. & Vallee, R.B. Mutually Exclusive Cytoplasmic Dynein Regulation by NudE-Lis1 and Dynactin. *Journal of Biological Chemistry* **286**, 39615-39622 (2011).

55.   Nyarko, A., Song, Y.J. & Barbar, E. Intrinsic Disorder in Dynein Intermediate
Chain Modulates Its Interactions with NudE and Dynactin. *Journal of Biological*
Chemistry **287**, 24884-24893 (2012).
56.   Moon, H.M. et al. LIS1 controls mitosis and mitotic spindle organization via the
LIS1-NDEL1-dynein complex. *Hum Mol Genet* **23**, 449-66 (2014).
57.   Zylkiewicz, E. et al. The N-terminal coiled-coil of Ndel1 is a regulated scaffold
that recruits LIS1 to dynein. *J Cell Biol* **192**, 433-45 (2011).
58.   Wang, S.S. & Zheng, Y.X. Identification of a novel dynein binding domain in
nudel essential for spindle pole organization in Xenopus egg extract. *Journal of*
Biological Chemistry **286**, 587-593 (2011).
59.   Efimov, V.P. Roles of NUDE and NUDF Proteins of Aspergillus nidulans: Insights
from Intracellular Localization and Overexpression Effects. *Molecular Biology of*
the Cell **14**, 871-888 (2003).
60.   Toropova, K. et al. Lis1 regulates dynein by sterically blocking its
mechanochemical cycle. *Elife* **3**(2014).
61.   Cianfrocco, M.A. et al. Lis1 has Two Distinct Modes of Regulating Dynein's
Mechanochemical Cycle. *Biophysical Journal* **112**, 43a-43a (2017).
62.   Gillies, J.P. et al. Structural basis for cytoplasmic dynein-1 regulation by Lis1.
Elife **11**(2022).
63.   Schlager, M.A., Hoang, H.T., Urnavicius, L., Bullock, S.L. & Carter, A.P. In vitro
reconstitution of a highly processive recombinant human dynein complex. *EMBO*
J **33**, 1855-68 (2014).
64.   Urnavicius, L. et al. The structure of the dynactin complex and its interaction with
dynein. *Science* **347**, 1441-1446 (2015).
65.   Casanal, A., Lohkamp, B. & Emsley, P. Current developments in Coot for
macromolecular model building of Electron Cryo-microscopy and
Crystallographic Data. *Protein Sci* **29**, 1069-1078 (2020).
66.   Kidmose, R.T. et al. Namdinator - automatic molecular dynamics flexible fitting of
structural models into cryo-EM and crystallography experimental maps. *IUCrJ* **6**,
526-531 (2019).
67.   Afonine, P.V. et al. Real-space refinement in PHENIX for cryo-EM and
crystallography. *Acta Crystallogr D Struct Biol* **74**, 531-544 (2018).

955