Title: ProteoPlotter: an executable proteomics visualization tool compatible with Perseus

Authors: Esther Olabisi-Adeniyi[1], Jason A. McAlister[1], Daniela Ferretti[2], Juergen Cox[2], Jennifer

Geddes-McAlister[1,*]

Affiliations: [1]Molecular and Cellular Biology, University of Guelph, Guelph, Ontario, Canada,

N1G 2W1. [2]Max Planck Institute of Biochemistry, Martinsried, Bayern, Germany, 82152

*Corresponding Author: Dr. Jennifer Geddes McAlister, jgeddesm@uoguelph.ca

Keywords: Proteomics, Data visualization, Perseus, *Klebsiella pneumoniae*, 1D annotation

enrichment, heat maps, Venn diagrams, volcano plots, UpSet plots, dynamic range

Abstract

Mass spectrometry-based proteomics experiments produce complex datasets requiring

robust statistical testing and effective visualization tools to ensure meaningful conclusions are

drawn. The publicly-available proteomics data analysis platform, Perseus, is extensively used to

perform such tasks, but opportunities to enhance visualization tools and promote accessibility of

the data exist. In this study, we developed ProteoPlotter, a user-friendly, executable tool to

complement Perseus for visualization of proteomics datasets. ProteoPlotter built on the Shiny

framework for R programming and enables illustration of multi-dimensional proteomics data.

ProteoPlotter provides mapping of one-dimensional enrichment analyses, enhanced adaptability

of volcano plots through incorporation of Gene Ontology terminology, visualization of 95%

confidence intervals in principal component analysis plots using data ellipses, and customizable

features. ProteoPlotter is designed for intuitive use by biological and computational researchers

alike, providing descriptive instructions (i.e., Help Guide) for preparing and uploading Perseus

output files. Herein, we demonstrate the application of ProteoPlotter towards microbial proteome

remodeling under altered nutrient conditions and highlight the diversity of visualizations enabled

with the platform for improved biological characterization. Through its comprehensive data

visualization capabilities, linked to the power of Perseus data handling and statistical analyses,

ProteoPlotter facilitates a deeper understanding of proteomics data to drive new biological

discoveries.

Introduction

Mass spectrometry (MS)-based proteomics allows for the comprehensive exploration of an organism's proteome (i.e., composition of proteins within a biological system). Insights derived from powerful MS-based proteomics technology continue to bring advancements and innovation to diverse fields within the life and biomedical sciences [1–4]. During proteomics experiments, MS instruments analyze protein mixtures across tens to thousands of samples through label-based and label-free approaches for detection, abundance, modifications, and interactions [5]. To derive meaningful biological insights from the high-resolution data acquired, powerful computational platforms and search engines, such as MaxQuant, MSFragger, and DIA-NN [6–8], have been developed, applying comprehensive qualitative and quantitative analyses to identify and quantify peptides and proteins. The resulting data allows researchers to investigate the proteome, and apply statistical testing, to define and interpret important biological findings.

To facilitate the analysis of proteomics data and provide strategies for visualization and interpretation, software platforms for data visualization have been developed. These platforms range in utility and accessibility based on user expertise, experience, and expectations. For example, RStudio is a computational environment for the R programming language that offers several libraries for the statistical analysis of MS-based proteomics data but requires a working knowledge of R[9–11]. As a result, many researchers opt for computational programs which provide an intuitive graphical user interface (GUI) offering advanced features (e.g., statistical testing, normalization), with minimal requirement for computational expertise. This approach increases accessibility for biological researchers and facilitates robust approaches for proteomics data analyses. For example, Perseus, a powerful and versatile software that is freely available for the statistical analysis and visualization of proteomics datasets, is a widely-used and accessible

platform [12]. The Perseus GUI boasts an array of options for analyzing data through an interactive

workflow that enables the user to track, preserve, and revisit a record of their activities on the

platform. Some visualization options in Perseus include, but are not limited to, volcano plots,

hierarchical clustering, and principal component analysis (PCA) plots.

To further harness the capabilities of the Perseus software, we introduce ProteoPlotter, an

executable platform designed to utilize exported files from Perseus in generating multi-

dimensional visualizations. Following user-enabled analysis in Perseus, such as filtering,

normalization, computation of 1D annotation enrichment scores, and/or two-sample t-test, a raw

tabular output can be exported from Perseus. ProteoPlotter utilizes these tabular datasets to

transform the analyses into enhanced graphical representations. With its user-friendly GUI

integrated within the Shiny web platform[13], ProteoPlotter is designed to facilitate data visualization

regardless of the user's computational expertise to generate 1D annotation enrichment heat maps,

interactive volcano plots, PCA plots with confidence intervales, dynamic range plots, Venn

diagrams, and UpSet plots, within minimal effort. Here, we demonstrate application of

ProteoPlotter to visualize proteome remodeling of the bacterial pathogen, *Klebsiella pneumoniae*,

under altered iron conditions, providing new visualization of our biological findings [14]. With

ProteoPlotter, researchers can create multi-dimensional figures for MS data, effectively

summarizing and conveying information for accessible and comprehensive data visualization and

interpretation.

Materials and Methods

Software development and availability

ProteoPlotter is an interactive software designed to complement Perseus in the downstream

analysis of quantitative proteomics data. ProteoPlotter was developed within the Shiny framework,

based on the R programming language (version 4.3.2), incorporating several R libraries in its

architecture, including ggplot2 (graphs), Shiny (GUI and web deployment), and Tidyverse (data

structure design)[9,10,13,15,16]. The executable file to launch ProteoPlotter locally, with no RStudio or

R software requirements, is available at https://github.com/JGM-Lab-UoG/. ProteoPlotter's source

code, which can be initiated within RStudio, is also available within the GitHub repository.

ProteoPlotter Input

ProteoPlotter is specifically designed for all calculations to be performed within Perseus,

taking full advantage of the Perseus data analysis workflow and archiving features. ProteoPlotter

uses data contained within specific columns of the exported matrices to generate figures. The

software accepts up to four Perseus matrices with specific plot-dependent column requirements,

enabling users to generate the following data visualizations: (i) 1D annotation enrichment heat

maps, (ii) multi-dimensional volcano plots, (iii) PCA plots with ellipses indicating 95% confidence

intervals, (iv) dynamic range plots, (v) Venn diagrams of up to five groups, and (vi) UpSet plots

(Table 1). All figures generated with ProteoPlotter are downloadable in high quality Portable

Network Graphics (PNG) format. Additionally, the software features a 'Guide' tab with usage

instructions. Helpful prompts also appear in the console as needed to assist users with

troubleshooting.

Table 1. Perseus matrices and column requirements for upload to ProteoPlotter.

Matrix uploaded to ProteoPlotter	Column headers in Perseus					
	LFQ intensity \<samples\>	Majority protein IDs	Student t-test columns*	GO / Keywords columns+	x and y coordinates	1D annotation columns−
1D Annotation matrix						✓
Volcano, PCA and Dynamic Range matrix	✓	✓	✓ (Optional: PCA and Dynamic Range plots)	Optional		
Threshold Curve matrix					✓	
Venn Diagram and UpSet matrix	✓ (Mean intensity values)					

*Student's t-test columns include: Significant, p-value, q-value, difference, and additional auto-generated columns.
+GO columns can include any or all of the following: Cellular Component, Molecular Function, Biological Process, and Keywords.
~1D annotation columns include the following: column, type, name, size, score, p-value, Benjamini-Hochberg False Discovery Rate [17].

1D Gene Ontology Annotation Enrichment Heat Maps

Within Perseus, the user can perform a functional annotation of identified proteins, and subsequently, compute 1D annotation scores to investigate the enrichment of these annotation terms within samples[18]. Computed results are exported from Perseus as a matrix and uploaded to ProteoPlotter to generate 1-D enrichment heat maps. The matrix contains 1D annotation enrichment scores for included GO terms. Each 1D score is defined between -1 to 1 such that a value close to 1 correlates to the colour red on the heat map and indicates that the functional annotation term is enriched among proteins within one group of the comparative pair under consideration, while a value approaching -1 correlates to the colour blue, meaning that the annotation term is enriched among proteins within the other group of the comparative pair. A value near 0 is depicted as the colour white on the heat map and suggests non-preferential enrichment between the two groups. Within the '1D Annotation Heat Map' tab in ProteoPlotter, the user

uploads the Perseus matrix and up to four distinct heat maps are generated for (i) GO – cellular

component (GOCC), (ii) GO – molecular function (GOMF), (iii) GO – biological process (GOBP),

and (iv) UniProt Keywords. On each heat map, annotation terms are displayed along rows with

columns indicating comparative pairs. Customizable options include: order of the groups, false

discovery rate setting, score minimum/maximum values, and x-axis labels.

Volcano plot

ProteoPlotter's volcano plot feature allows the user to illustrate multiple layers of

information from the dataset. For example, average protein intensity values and GO term

annotations can be layered upon the plot (Figure 1). ProteoPlotter accepts up to two input matrices

for the volcano plot, but the 'Threshold Curve Matrix' is optional; the 'Matrix with t-test' is

required. The 'Matrix with t-test' input needs to contain the imputed protein intensities, Student's

t-test results, and majority protein identifications. When generating volcano plots for multiple

comparative pairs, a separate t-test matrix should be generated for each pair. Subsequently, each

matrix containing t-test results for a specific comparative pair is exported individually, rather than

consolidating all t-test results into a single matrix. Furthermore, any of these t-test matrices can

also serve as inputs for generating dynamic range or PCA plots given that the structure provides

the required information regardless of the t-test performance.

Figure 1: ProteoPlotter volcano plots. The size of each point (corresponding to a protein) varies

by average protein abundance across replicates. The GO term annotation layer highlights the

significantly different proteins annotated with a selected term.

On the volcano plot, negative $\log_{10}$ p-values and Student's t-test differences from the

matrix are plotted against each other. For depiction of functional annotations on the figure, the

user must ensure that at least one functional annotation column (e.g., GOBP, GOCC, GOMF,

Keywords) is provided. The user can also generate the False Discovery Rate (FDR) curves matrix

in Perseus to layer FDR threshold curves on the volcano plot[17]. To achieve this, the user first

creates a volcano plot in Perseus with the same S_0 and FDR values that will be applied within

ProteoPlotter. Subsequently, the FDR curves matrix containing the x and y coordinates is exported

and uploaded. Additional volcano plot options within ProteoPlotter include hovering over points

to disapply their respective GO terms within a chosen GO category, filtering GO terms to retain

only those with 1D annotation scores (requires uploading the 1D annotation enrichment matrix),

adjustment of axes limits, adjustment of FDR and S_0 thresholds, and selection of colours from

multiple colour palette libraries.

Dynamic range and principal component analysis plots

As previously stated, the t-test matrix generated for the volcano plot can be used as an input

file to generate dynamic range plots for all groups within the uploaded matrix, as well as a PCA

plot. Alternatively, if volcano plots are not being generated, the user may export the matrix

containing imputed protein intensities without performing t-tests from Perseus. Once the required

matrix is uploaded, the 'Generate Plot' button within the 'Dynamic Range Plot' tab or the 'PCA

Plot' tab can be used to generate the figures. The dynamic range plots illustrate the average protein

abundance ($\log_{10}$) across replicates against their descending-order ranking, depicting the

distribution of protein abundance across measured proteomes. With ProteoPlotter's traditional

PCA function, users can investigate clustering patterns among group replicates and the variance

driving separation among groups. Moreover, the software also offers the option to overlay data

ellipses onto the PCA plot to illustrate the multivariate t-distribution of the data at a 95%

confidence interval[19,20]. Note that this data ellipse feature is only applicable to groups with four or

more replicates.

Venn diagram and UpSet plots

ProteoPlotter generates Venn diagrams and UpSet plots from average protein intensity

values computed in Perseus. To produce a Venn diagram, the user uploads the matrix within the

'Venn Diagram' tab, then selects the 'Generate Plot' button. The Venn diagram allows users to

assess unique and shared proteins across up to five groups, and users can select colours for each

group within the Venn diagram. In contrast, the UpSet plot in ProteoPlotter largely overcomes this

limitation of group number by enabling users to visualize up to 25 groups using the same matrix

uploaded for the Venn diagram. With the matrix already uploaded, the user selects the 'Generate

Plot' button within the 'UpSet Plot' tab to produce the figure.

Analysis of Sample Dataset

A dataset from a previous proteomics study on the bacterial pathogen, *Klebsiella*

pneumoniae, was selected to demonstrate the multi-dimensional visualization features available

within ProteoPlotter[14]. The dataset was obtained from the PRIDE database (Project PXD015623)[21].

In the study, the impact of iron limitation on bacterial growth, survival, and proteome

differentiation of *K. pneumoniae* was analyzed. The bacterium was cultured in low iron media,

and media supplemented with $Fe_2(SO_4)_3$ at 10 µM and 100 µM in four biological replicates for

each condition. Cellular proteome samples were obtained from each treated group using tandem

mass spectrometry techniques (LC-MS/MS) as previously described[14]. Raw mass spectrometry

files were processed with MaxQuant (version 1.6.0.26), and the tab-delimited output from

MaxQuant, proteingroups.txt, was imported into Perseus (version 2.1.3.0) (Figure 2).

Figure 2: Perseus workflow for cellular proteome analysis of *K. pneumoniae* under altered iron conditions. Teal workflow: before GO annotation; Red workflow: after GO annotation.

The workflow in Perseus performed the following analyses, including removal of proteins that matched to the reverse database, contaminants, and proteins only identified with modified peptides (matrices 1-3). Intensities were converted to a $\log_2$ scale (matrix 4), and LFQ intensity columns were categorically annotated as follows: LIM, Fe10 µM, and Fe100 µM (matrix 5). Only proteins with a minimum of three valid values within at least one group were retained to filter the total number of proteins identified (matrix 6). Next, we averaged intensity values to derive the matrix for the Venn diagram and UpSet plot, applying a minimum of three valid values per group

as the threshold (matrix 6A). The resulting matrix was reannotated with group names and exported

(matrix 6B). Subsequently, missing values were imputed with numbers drawn from a normal

distribution with a downshift by 1.8 standard deviations and a width of 0.3 standard deviations

(matrix 7). GO and UniProt Keywords annotations were collected from UniProtKB using the

UniProt ID mapper feature[22,23]. The annotations were added to the matrix outside of the Perseus

platform and the annotated matrix was re-uploaded to the Perseus workflow (matrix 'GO

annotated).

For statistical testing, Student's t-test (two-sample test) were computed to identify proteins

with significant differences in abundance (p-value ≤ 0.05) between LIM and Fe10 µM; LIM and

FE100 µM; Fe10 µM and Fe100 µM (matrix 9-11). Permutation-based FDR was applied for

multiple hypothesis testing correction with 250 randomizations, FDR = 0.05, and $S_0 = 1$. To use

the ProteoPlotter volcano plot feature, we exported the matrix containing the Student's t-test

results for LIM vs. Fe10 µM (matrix 9). We used the same matrix to create both the PCA and

dynamic range plots. Additionally, threshold curves for the volcano plots were generated by

exporting the curve matrix produced by the volcano plot feature in Perseus (matrix 9A). For the

1D annotation heat maps, we computed 1D annotation scores suing the 't-test difference' columns

of all three comparative pairs present within matrix 11 (FDR = 0.05). We also filtered the resulting

matrix (matrix 12) to exclude rows with non-annotation terms. The final matrix was exported to

ProteoPlotter to generate the 1D annotation heat maps (matrix 13). All matrices were exported in

the .txt format.

Results

Quantitative characterization of the K. pneumoniae cellular proteome under altered iron

conditions using ProteoPlotter

We applied ProteoPlotter's volcano plot feature for exploring the proteins that were significantly differentially abundant between LIM and FE10 µM conditions (Figure 3A). A significant difference was observed for 31 proteins; 16 proteins were significantly more abundant under iron-limited (LIM) conditions while 15 proteins were significantly more abundant under iron-replete (10 µM $Fe_2(SO_4)_3$) conditions. The S_0 and FDR values, which are adjustable when creating the plot, are displayed in purple at the base of the plot for the user's information. Users can choose to include or exclude the threshold curves.

ProteoPlotter also allows the user to highlight proteins IDs, enabling rapid identification of specific significantly different proteins. Next, we overlaid select UniProt Keywords onto the already labeled volcano plot, thereby allowing us to visualize and investigate the functional properties of significantly different proteins (Figure 3B). For example, we illustrate the feature by labeling GO annotations associated with the TonB box, a protein involved in ferric-siderophore receptor function to provide a mechanical linkage between the cytoplasmic and outer membrane, promoting energy transduction from the proton motive force to the receptor[24–27]. Other GO categories can also be illustrated in ProteoPlotter. In addition, we visualized the average protein intensity for each significantly altered protein. Through this figure, we observed that most of the significantly differentially abundant proteins in the iron-limited samples exhibited relatively moderate intensity levels (Figure 3C). In contrast, the Fe10 µM samples demonstrated a consistent distribution of both low and high intensity values.

Figure 3: Significantly different protein abundance profiles under iron-limited (LIM) vs. iron-replete (Fe10 µM) conditions using ProteoPlotter's volcano plot feature. A) Protein ID selection layer labeling select proteins. B) GO annotation layer displaying the significantly different proteins associated with TonB box, a GO term under the Keywords category. C) Volcano plot layer with average intensity values for significantly different proteins in each treatment group. Student's t-test p-value < 0.05; FDR = 0.05; $S_0 = 1$.

Using ProteoPlotter, we generated a 1D annotation heat map based on UniProt Keywords to gain another perspective into the functional enrichment of the cellular proteome under all three tested conditions (Figure 4). Several categories showed positive enrichment in replete (Fe10 µM

and Fe100 µM) conditions compared to limited conditions, including rRNA-binding, ribosomal,

and ribonucleoprotein categories. We also observed a positive enrichment of iron transport and

TonB GO terms under limited conditions relative to Fe100 µM replete conditions. More iron-

associated terms, such as iron-sulfur and 4Fe-4S clusters were positively enriched under 10 µM

replete conditions relative to 100 µM replete conditions. The patterns observed in the heat maps

suggest that changes in iron availability alters the cellular environment in *K. pneumoniae*, leading

to an increase in iron-associated protein production under low iron conditions.

Figure 4: 1D annotation enrichment heat map for UniProt Keywords for *K. pneumoniae*

under altered iron conditions. Student's t-test p-value < 0.05; FDR = 0.05; -1.0 < Score < 1.0.

Qualitative characterization of the K. pneumoniae cellular proteome under altered iron conditions using ProteoPlotter

The Venn diagram in ProteoPlotter illustrates the shared and unique number of proteins identified among groups, as counts, percentages or both (Figure 5A). The majority of identified proteins are defined within a core proteome of 1855 proteins present across all three tested conditions. Notably, 33 proteins were unique to Fe100 µM and 36 proteins were unique to Fe10 µM conditions, whereas the iron-limited conditions demonstrated unique identification of 52 proteins. This information is also depicted using the UpSet plot feature (Figure 5B).

Figure 5: Qualitative analysis Venn and UpSet plots in ProteoPlotter. A) Venn diagram comparing the number of proteins identified in the cellular proteome cultured under limited (LIM) and replete (Fe10 µM and Fe100 µM) iron conditions. B) UpSet plot demonstrating protein identification patterns across the conditions.

Next, we generated PCA and dynamic range plots using imputed LFQ intensities, to explore the distribution of protein abundance values. In the PCA, treatment groups clustered, with

the largest variability observed between the iron-limited group and both iron-replete groups

(Principal Component 1 = 22.78%; Figure 6A). The second principal component accounted for

16.64% of the variance, reflecting the reproducibility among replicates. We also added data

ellipses to the PCA plot to explore the t-distribution of samples within each treatment group. This

step revealed an overlap between both replete conditions and an isolated ellipse for the iron-limited

samples (95% CI; Figure 6B). Lastly, dynamic range plots were generated to assess the varying

abundance of proteins identified in the cellular proteome under iron-limited and iron-replete

conditions. The protein abundance within each group exhibited a dynamic range of distribution,

with proteins showing high, moderate, and low intensities, all spanning across a similar range of

8 to 11 ($\log_{10}$ LFQ intensity) (Figure 7).

Figure 6: Qualitative analysis PCA plots in ProteoPlotter. A) PCA of the cellular proteome

identifying clusters by group. B) Data ellipses visualizing the multivariate t-distribution of each

treatment group (95% CI).

Figure 7: Qualitative analysis dynamic range plots in ProteoPlotter. Average protein abundance plotted against protein ranks (highest to lowest) to illustrate the dynamic range of *K. pneumoniae* cellular proteome under iron-limited and iron-replete conditions.

Discussion

ProteoPlotter provides new visualization tools and capabilities, and application to a previously published proteomics dataset of *K. pneumoniae* results in a demonstrated ability to decipher insights into how iron availability affects the bacteria's cellular proteome. We selected proteins highlighted in the volcano plot and identified them with the UniProtKB database to illustrate our ability to use ProteoPlotter for investigation of biologically relevant proteins. For instance, CirA (A6TBP0), an outer membrane pore protein requiring TonB, was present at

significantly higher abundance under iron-limited conditions [14,27]. CirA associated with

siderophore secretion, which is demonstrated by *K. pneumoniae* to improve iron uptake under low-

iron availability[28,29]. Gram-negative bacteria, such as *Klebsiella* spp., use siderophore-release

mechanisms under iron-limited conditions as the molecule's high affinity for metals improves iron

uptake[30,31]. Therefore, an increase in CirA's abundance likely reflects the activation of

siderophore-release mechanisms triggered by low-iron conditions. Another protein, Fe(II)-2OG-

dependent oxygenase (A6T915) was also more abundant under iron-limited conditions, which is

consistent with previous findings that show a correlation between increased Fe(II)-2OG-dependent

oxygenase activity in *Saccharomyces cerevisiae* and low iron conditions[32].

ProteoPlotter's multidimensional approach to data visualization enhances data analysis and

interpretation of results. For instance, overlaying GO terms onto the volcano plot revealed that

CirA protein is associated with the TonB box annotation term. We also observed a positive

enrichment of TonB under iron-limited conditions in the 1D heat map, which is likely correlated

with the significantly higher abundance of CirA, a protein that is dependent on TonB. Analyzing

this dataset from multiple perspectives through the volcano plot and heat map enables the user to

corroborate associations between a protein and defined biological functions. To further investigate

GO terms associated with significantly different proteins, ProteoPlotter users can focus on key

terms by applying the filter that retains only those terms enriched in the 1D annotation dataset.

During proteomics analysis, it is crucial for proteins to be detectable across varying

abundances, particularly for low abundant proteins[33]. Through the dynamic range plot in

ProteoPlotter, we observed a well distributed spectrum of protein abundances, including those at

lower levels, across five orders of magnitude. Researchers can apply this tool before proceeding

with in-depth analysis and visualization. Additional analysis, with the Venn diagram and UpSet

plot features indicates that the cellular proteome under iron-limited conditions contained more

unique proteins relative to the proteome within iron-replete environments. Given the significantly

higher abundance of siderophore-associated proteins like CirA under iron-limited conditions, it is

likely that iron-limited samples produce more condition-specific proteins to facilitate efficient

siderophore-release and iron uptake. Dimensionality reduction with PCA in ProteoPlotter also

revealed a clear distinction between the replicates in iron-limited environments and those in iron-

replete environments. The clusters for both 10 µM and 100 µM samples were positioned closely,

suggesting a similarity between the proteomic profiles and a separation from the iron-limited

cluster. T-distribution ellipses of the groups further demonstrate that the iron-limited proteome of

distinct from both iron-replete proteomes. From the overlapping ellipses, we can infer that both

iron-replete samples maintain similarity to one another whereas the proteomic profile under

limited conditions remains distinct (95% CI).

Conclusion

ProteoPlotter serves as a valuable supplementary visualization tool for proteomics data

analyzed in Perseus to offer new modes of extracting biological insights from complex data. The

program enabled us to successfully illustrate, and extract details embedded within our sample data.

ProteoPlotter integrates multiple data visualization techniques into a user-friendly, executable

platform, allowing for a harmonized exploration of data in tandem with Perseus. Presenting results

graphically and exploring traditional visualization methods in a multidimensional manner

empowers researchers to gain a clearer understanding of their data, thereby improving data

interpretation.

Conflict of Interest

J.C. is developer of the Perseus software.

Data availability statement

The executable file to launch ProteoPlotter locally, with no RStudio or R software requirements, is available at https://github.com/JGM-Lab-UoG/. ProteoPlotter's source code, which can be initiated within RStudio, is also available within the GitHub repository. Proteomics data is available through PRIDE database (Project PXD015623).

Acknowledgements

The authors thank members of the Geddes-McAlister lab for testing of ProteoPlotter and insightful discussions and feedback throughout the development process.

Funding

Funding is provided to J.G.M. through the Natural Sciences and Engineering Research Council of Canada (Discovery Grant), New Frontiers Research Fund: Exploration, and the Canada Research Chairs Program.

Contributions

J.A.M. & J.G.-M. conceptualized the design and development of ProteoPlotter. E.O.-A. designed and developed ProteoPlotter. D.F. & J.C. provided guidance and input into the design and development of ProteoPlotter. E.O.-A. & J.A.M. prepared the first draft of the manuscript. E.O.-

380   A. & J.G.-M. prepared figures. J.A.M. & J.G.-M. prepared the final version of the manuscript. All

381   authors have reviewed and approve the submitted manuscript.

382   **References**

383   (1)   Meissner, F.; Geddes-McAlister, J.; Mann, M.; Bantscheff, M. The Emerging Role of Mass

384   Spectrometry Based Proteomics in Drug Discovery. *Nat Rev Drug Discov* **2022**.

385   (2)   Geddes-McAlister, J.; Prudhomme, N.; Dutierrez Gongora, D.; Cossar, D.; McLean, M. D.

386   The Emerging Role of Mass Spectrometry-Based Proteomics in Molecular Pharming. *Curr*

387   *Opin Chem Biol* **2022**, *68*, 102133.

388   (3)   Geddes-McAlister, J.; Roux-Dalvai, F.; Droit, A. Proteomics, Bioinformatics, and

389   Infectious Diseases. In *Genetics and Evolution of Infectious Diseases*; Elsevier, 2024; pp

390   465–492. https://doi.org/10.1016/B978-0-443-28818-0.00024-0.

391   (4)   Wenk, D.; Zuo, C.; Kislinger, T.; Sepiashvili, L. Recent Developments in Mass-

392   Spectrometry-Based Targeted Proteomics of Clinical Cancer Biomarkers. *Clin Proteomics*

393   **2024**, *21* (1), 6. https://doi.org/10.1186/s12014-024-09452-1.

394   (5)   Aebersold, R.; Mann, M. Mass-Spectrometric Exploration of Proteome Structure and

395   Function. *Nature* **2016**, *537* (7620), 347–355. https://doi.org/10.1038/nature19949.

396   (6)   Cox, J.; Mann, M. MaxQuant Enables High Peptide Identification Rates, Individualized

397   p.p.b.-Range Mass Accuracies and Proteome-Wide Protein Quantification. *Nat Biotechnol*

398   **2008**, *26* (12), 1367–1372. https://doi.org/10.1038/nbt.1511.

399   (7)   Kong, A. T.; Leprevost, F. V; Avtonomov, D. M.; Mellacheruvu, D.; Nesvizhskii, A. I.

400   MSFragger: Ultrafast and Comprehensive Peptide Identification in Mass Spectrometry–

401   Based Proteomics. *Nat Methods* **2017**, *14* (5), 513–520. https://doi.org/10.1038/nmeth.4256.

402   (8)   Demichev, V.; Messner, C. B.; Vernardis, S. I.; Lilley, K. S.; Ralser, M. DIA-NN: Neural

403   Networks and Interference Correction Enable Deep Proteome Coverage in High

404   Throughput. *Nat Methods* **2020**, *17* (1), 41–44. https://doi.org/10.1038/s41592-019-0638-x.

405   (9)   R Core Team. R: A Language and Environment for Statistical Computing. Foundation for

406   Statistical Computing 2024. https://www.R-project.org/ (accessed 2024-10-06).

407   (10)   RStudio Team. RStudio: Integrated Development for R. RStudio 2020.

408   http://www.rstudio.com/ (accessed 2024-10-06).

409   (11)   Gatto, L.; Gibb, S.; Rainer, J. MSnbase, Efficient and Elegant R-Based Processing and

410   Visualization of Raw Mass Spectrometry Data. *J Proteome Res* **2021**, *20* (1), 1063–1069.

411   https://doi.org/10.1021/acs.jproteome.0c00313.

412   (12)   Tyanova, S.; Temu, T.; Sinitcyn, P.; Carlson, A.; Hein, M. Y.; Geiger, T.; Mann, M.; Cox,

413   J. The Perseus Computational Platform for Comprehensive Analysis of (Prote) Omics Data.

414   *Nat Methods* **2016**, *13* (9), 731–740.

415   (13)   Chang W., et al, Shiny: Web Application Framework for R. 2024. https://CRAN.R-

416   project.org/package=shiny.

417   (14)   Muselius, B.; Sukumaran, A.; Yeung, J.; Geddes-McAlister, J. Iron Limitation in *Klebsiella*

418   *pneumoniae* Defines New Roles for Lon Protease in Homeostasis and Degradation by

419   Quantitative Proteomics. *Front Microbiol* **2020**, *11*, 546.

420   https://doi.org/10.3389/fmicb.2020.00546.

421   (15)   Wickham, H.; Averick, M.; Bryan, J.; Chang, W.; McGowan, L.; François, R.; Grolemund,

422   G.; Hayes, A.; Henry, L.; Hester, J.; Kuhn, M.; Pedersen, T.; Miller, E.; Bache, S.; Müller,

423   K.; Ooms, J.; Robinson, D.; Seidel, D.; Spinu, V.; Takahashi, K.; Vaughan, D.; Wilke, C.;

424   Woo, K.; Yutani, H. Welcome to the Tidyverse. *J Open Source Softw* **2019**, *4* (43), 1686.

425   https://doi.org/10.21105/joss.01686.

426   (16)   Wickham, H. Ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York

427   2016.

(17) Benjamini, Y.; Hochberg, Y. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B*

(Methodological) **1995**, *57* (1), 289–300. https://doi.org/10.1111/j.2517-

6161.1995.tb02031.x.

(18) Cox, J.; Mann, M. 1D and 2D Annotation Enrichment: A Statistical Method Integrating

Quantitative Proteomics with Complementary High-Throughput Data. *BMC Bioinformatics*

2012, *13*, S12. https://doi.org/10.1186/1471-2105-13-S16-S12.

(19) Tang, Y.; Horikoshi, M.; Li, W. Ggfortify: Unified Interface to Visualize Statistical Results

of Popular R Packages. *R J* **2016**, *8* (2), 474. https://doi.org/10.32614/RJ-2016-060.

(20) Brereton, R. G. The *t* -distribution and Its Relationship to the Normal Distribution. *J*

Chemom **2015**, *29* (9), 481–483. https://doi.org/10.1002/cem.2713.

(21) Perez-Riverol, Y.; Csordas, A.; Bai, J.; Bernal-Llinares, M.; Hewapathirana, S.; Kundu, D.

J.; Inuganti, A.; Griss, J.; Mayer, G.; Eisenacher, M.; Pérez, E.; Uszkoreit, J.; Pfeuffer, J.;

Sachsenberg, T.; Yilmaz, Ş.; Tiwary, S.; Cox, J.; Audain, E.; Walzer, M.; Jarnuczak, A. F.;

Ternent, T.; Brazma, A.; Vizcaíno, J. A. The PRIDE Database and Related Tools and

Resources in 2019: Improving Support for Quantification Data. *Nucleic Acids Res* **2019**.

https://doi.org/10.1093/nar/gky1106.

(22) UniProt: The Universal Protein Knowledgebase in 2021. *Nucleic Acids Res* **2021**, *49* (D1),

D480–D489. https://doi.org/10.1093/nar/gkaa1100.

(23) Zaru, R.; Orchard, S. UniProt Tools: BLAST, Align, Peptide Search, and ID Mapping. *Curr*

Protoc **2023**, *3* (3). https://doi.org/10.1002/cpz1.697.

449 (24) Mossialos, D.; Amoutzias, G. Role of Siderophores in Cystic Fibrosis Pathogenesis: Foes

450   or Friends? *International Journal of Medical Microbiology* **2009**, *299* (2), 87–98.

451   https://doi.org/10.1016/j.ijmm.2008.06.008.

452 (25) Kumar, A.; Chakravorty, S.; Yang, T.; Russo, T. A.; Newton, S. M.; Klebba, P. E.

453   Siderophore-Mediated Iron Acquisition by *Klebsiella pneumoniae*. *J Bacteriol* **2024**, *206*

454   (5). https://doi.org/10.1128/jb.00024-24.

455 (26) Hsieh, P.; Lin, T.; Lee, C.; Tsai, S.; Wang, J. Serum-Induced Iron-Acquisition Systems and

456   TonB Contribute to Virulence in *Klebsiella pneumoniae* Causing Primary Pyogenic Liver

457   Abscess. *J Infect Dis* **2008**, *197* (12), 1717–1727. https://doi.org/10.1086/588383.

458 (27) Noinaj, N.; Guillier, M.; Barnard, Travis J.; Buchanan, S. K. TonB-Dependent Transporters:

459   Regulation, Structure, and Function. *Annu Rev Microbiol* **2010**, *64* (1), 43–60.

460   https://doi.org/10.1146/annurev.micro.112408.134247.

461 (28) Lan, P.; Lu, Y.; Jiang, Y.; Wu, X.; Yu, Y.; Zhou, J. Catecholate Siderophore Receptor CirA

462   Impacts Cefiderocol Susceptibility in *Klebsiella pneumoniae*. *Int J Antimicrob Agents* **2022**,

463   *60* (4), 106646. https://doi.org/10.1016/j.ijantimicag.2022.106646.

464 (29) Zhang, Z.; Du, W.; Wang, M.; Li, Y.; Su, S.; Wu, T.; Kang, Y.; Shan, X.; Shi, Q.; Zhu, G.

465   Contribution of the Colicin Receptor CirA to Biofilm Formation, Antibotic Resistance, and

466   Pathogenicity of *Salmonella* Enteritidis. *J Basic Microbiol* **2020**, *60* (1), 72–81.

467   https://doi.org/10.1002/jobm.201900418.

468 (30) Holden, V. I.; Bachman, M. A. Diverging Roles of Bacterial Siderophores during Infection.

469   *Metallomics* **2015**, *7* (6), 986–995. https://doi.org/10.1039/C4MT00333K.

470   (31)   Crumbliss, A. L.; Harrington, J. M. Iron Sequestration by Small Molecules:

471   Thermodynamic and Kinetic Studies of Natural Siderophores and Synthetic Model

472   Compounds; 2009; pp 179–250. https://doi.org/10.1016/S0898-8838(09)00204-9.

473   (32)   Deepa, A.; Naveena, K.; Anindya, R. DNA Repair Activity of Fe(II)/2OG-Dependent

474   Dioxygenases Affected by Low Iron Level in Saccharomyces Cerevisiae. *FEMS Yeast Res*

475   **2018**, *18* (2). https://doi.org/10.1093/femsyr/foy014.

476   (33)   Schessner, J. P.; Voytik, E.; Bludau, I. A Practical Guide to Interpreting and Generating

477   Bottom-up Proteomics Data Visualizations. *Proteomics* **2022**, *22* (8).

478   https://doi.org/10.1002/pmic.202100103.