Theory of non-dilute binding and surface phase separation applied to membrane-binding proteins

Xueping Zhao[1,†]**, Daxiao Sun**[2,3†]**, Giacomo Bartolucci**[4]**, Anthony A. Hyman**[2,5]**, Alf Honigmann**[3,5,6*]**, Christoph A. Weber**[7*]

[1]Department of Mathematical Sciences, University of Nottingham Ningbo China; [2]Max Planck Institute of Molecular Cell Biology and Genetics, Dresden, Germany; [3]Technische Universität Dresden, Biotechnologisches Zentrum, Center for Molecular and Cellular Bioengineering (CMCB), Dresden, Germany; [4]Department of Physics, Universitat de Barcelona, Barcelona, Spain; [5]Cluster of Excellence Physics of Life, TU Dresden, Dresden, Germany; [6]Center for Systems Biology Dresden, Dresden, Germany; [7]Faculty of Mathematics, Natural Sciences, and Materials Engineering: Institute of Physics, University of Augsburg, Augsburg, Germany; [8]Max Planck Institute for the Physics of Complex Systems, Dresden, Germany

*For correspondence:
christoph.weber@physik.
uni-augsburg.de;
alf.honigmann@tu-dresden.de

[†]These authors contributed equally to this work

Abstract   Surface binding and surface phase separation of cytosolic scaffold proteins on lipid membranes are involved in many cellular processes such as cell signaling, cell adhesion, and cortex regulation. However, the interplay between surface binding and surface phase separation is poorly understood. In this work, we study this interplay by deriving a general thermodynamic model and applying it to in vitro reconstitution experiments of membrane-binding proteins involved in tight junction initiation. Our theory extends the classical surface binding isotherm to account for non-dilute and heterogeneous conditions where components can phase separate. Using this framework we demonstrate how surface phase separation is governed by the interaction strength among membrane-bound scaffold proteins and their binding affinity to the membrane surface. Comparing the theory to reconstitution experiments, we show that tuning the oligomerization state of the adhesion receptors in the membrane controls surface phase transition and patterning of the scaffold protein ZO1. These findings suggest a fundamental role of the interplay between non-dilute surface binding and surface phase separation in the formation of the tight junction. More broadly, our work highlights non-dilute surface binding and surface phase separation as a common organizational principle for membrane-associated structures in living cells.

Introduction

Liquid-liquid phase separation (LLPS) has emerged as a fundamental principle governing the spatial organization of cellular components into dynamic, membrane-less compartments termed biomolecular condensates *Brangwynne et al.* (*2011*); *Mahen and Venkitaraman* (*2012*); *Gall* (*2003*); *Brangwynne et al.* (*2009*); *Fritsch et al.* (*2021*); *JR and R.* (*2009*); *CJ and R.* (*2012*). Such condensates create distinct microenvironments that influence a wide range of cellular functions, including signaling *Su et al.* (*2016*); *Wang et al.* (*2021*); *Xiao et al.* (*2022*), transcription *Boija et al.* (*2018*); *Lu et al.* (*2020*), and stress responses *Riback et al.* (*2017*).

While LLPS has been extensively studied in the cytoplasm, protein phase separation is also involved in many biological processes happening at membrane surfaces such as signaling *Li et al.* (*2012*); *CASE et al.* (*2019*); *Zeng et al.* (*2019*); *Li et al.* (*2022*); *Zeng et al.* (*2018*); *Su et al.* (*2016*); *Zheng et al.* (*2022*), synapse formation *Zeng et al.* (*2018*), cell adhesion *Beutel et al.* (*2019*); *Pombo-García et al.* (*2024*); *Wang et al.* (*2021*) and cortex regulation *CASE et al.* (*2019*); *Wiegand and Hyman* (*2020*). Protein condensation at membranes is often induced by binding of multivalent scaffold proteins present in the cytoplasm to integral membrane proteins or lipids on the surface of the membrane. Membrane binding of scaffold proteins can result in condensation far below the concentration required for phase separation in the bulk, providing spatial-temporal control over the condensation process. Therefore, membrane binding and surface phase separation are tightly linked processes *GrandPre et al.* (*2024*).

How membrane binding and surface phase separation influence each other is not well understood. Classical surface binding theories, such as the Langmuir isotherm *Langmuir* (*1918*), assume dilute concentrations where binding sites are independent and non-interacting. These assumptions are not valid for phase separating systems, where surface densities are *non-dilute* and intermolecular interactions among membrane-bound molecules become important. More recently, non-dilute binding and phase separation have been incorporated into theoretical models *Zhao et al.* (*2021*, 2024); *Liese et al.* (*2023*). However, these models still rely on simplifying assumptions such as uniform binding sites, which is not matching the reality of biological membranes.

Binding of proteins to biological membranes is typically tightly controlled in space and time by either increasing local densities of binding sites such as in the formation of cell adhesion complexes *Otani et al.* (*2019*); *Zhao et al.* (*2018b*), or by increasing binding affinities for example by phosphorylation *Li et al.* (*2012*). To accurately describe such phenomena, theoretical models must incorporate both non-homogeneous and non-dilute surface densities of proteins at the membrane. While a recent thermodynamic model has linked binding heterogeneity to prewetting phenomena *Rouches et al.* (*2021*), developing a theory that integrates these heterogeneous, non-dilute conditions and captures the potential for surface-bound phase transitions remains crucial for a comprehensive understanding of membrane-associated assemblies.

In this work, we present a theoretical framework for the interplay between surface binding and surface phase separation that includes heterogeneous and non-dilute membrane surface densities. We show how the classical binding isotherm must be fundamentally modified when molecules bound to membrane receptors interact strongly and undergo surface condensation. We derived a generalized binding isotherm which extends classical adsorption models by accounting for the non-dilute nature of surface-bound complexes and their interactions. Using this framework, we explored the behaviors of non-dilute surface binding under varying bulk and surface compositions, binding affinities, and interaction strengths between membrane bound complexes. To illustrate the applicability of our theory, we focus on in vitro reconstituted membrane systems that model components of tight junctions. We explain how oligomerization states influence membrane organization. More generally, our theory reveals how non-dilute surface binding and membrane-associated phase separation interact to influence membrane organization, shedding light on a fundamental physicochemical mechanism through which cells regulate membrane associated assembly processes.

Model for protein binding to receptors on a membrane

In the first part of this work, we discuss a theoretical framework that describes binding of soluble scaffold proteins to lipid membranes via interactions with receptor proteins embedded in a membrane. In addition, the scaffold proteins can undergo (surface) phase separation, which results in formation of protein-dense and dilute regions on the membrane (Fig. 1a,b). The theoretical framework builds upon previous works on the interplay of surface-associated phase transition (surface phase separation, prewetting) and surface binding *Zhao et al.* (*2021*, 2024). However, previous models do not involve a receptor component, reducing the stable coexistence of domains poor

Figure 1. Schematics of the bulk-membrane coupled system. a Schematic representation of scaffold proteins binding to the receptors anchored on a lipid bilayer. Each receptor (magenta) is tagged to a lipid (orange) in the supported lipid bilayers (SLBs). Scaffold proteins P (green) can bind to receptors R, forming a receptor-protein complex PR. We introduce two more effective components in the membrane: the lipids with a receptor without any protein bound but a water layer on top, and a lipid patch without a receptor but with the water layer on top. These components are termed "effective" because they also include water molecules that form a molecular layer adjacent to the membrane. For clarity, we do not depict these water molecules in the schematics, thereby providing a clearer view of the surface binding structure. We note that the height of the effective binding layer is denoted as h, which is also the height of protein-receptor complex units. In addition, h_R represents the height of free receptors. Most importantly, the effective components are confined to the membrane and can only move diffusely parallel to it, or unbind and dissolve in the bulk. **b** shows the mathematical notations in our theoretical model. The volume fraction of proteins in the bulk is ϕ; ϕ_{PR}, ϕ_R are the area fraction of membrane-bound complexes and free receptors. Moreover, $\phi|_{z=0}$ is the volume fraction of protein at the membrane surface, which is denoted as S. S_o indicates the boundary surface of the bulk region V opposite to the membrane surface S. The boundary surfaces of the bulk region V, excluding S and S_o, are denoted by ∂V; The membrane surface's boundary is represented by ∂S.

---

and rich in bound protein to a specific value of the bulk protein concentration where the phase transition occurs. However, this behavior is not consistent with in vitro experiments where phase coexistence exists over a large range of bulk protein concentrations. In this light, we propose an extended but still minimal model in terms of the number of membrane-associated components: free lipid, free receptor, and receptor bound to a protein from the bulk (Fig. 1a).

The proteins in bulk can bind to receptors in the membrane, while membrane lipids are not accessible for protein binding. Note that the bulk does not contain any receptors. In our theory, we focus on the classical case of supported lipid bilayers (SLBs); however, we note that the theory can be straightforwardly extended to model real cell membranes even more realistically, e.g. accounting for more components and the effects of curvature. Since receptors are confined to the membrane, bound proteins can only diffuse within a layer adjacent to the membrane surface. This layer also contains solvent molecules, which are exchanged with the bulk when proteins bind. In the three-dimensional bulk V, a binary mixture of protein and solvent molecules is present. The quasi-two-dimensional domain related to the membrane is denoted as S; see Fig. 1b for an illustration. Moreover, S_o indicates the boundary surface of bulk region V opposite to membrane surface S. The boundary surfaces of the bulk region V, excluding S and S_o, are denoted by ∂V. The membrane (one-dimensional) boundary is represented by ∂S.

To describe the setting outlined above, we propose a spatially coarse-grained model for bulk-membrane coupled systems. To this end, we introduce three effective components on a surface S: lipid with receptors bound to protein PR, lipid with receptors but without protein-bound R, and lipid with only the water layer on top L. Proteins in the bulk are denoted by P. For simplicity, we consider an incompressible mixture where the molecular volume of solvent v and protein v_P in bulk are constants. Similarly, the molecular areas of lipid v_L, receptor v_R and protein-receptor complex v_{PR} are constants. The thermodynamics of such an incompressible mixture is characterized by the bulk volume fraction of the proteins $\phi(\pmb{x}, t)$, and the area fraction of the receptor-protein complex PR that is confined in the membrane $\phi_{PR}(\pmb{x}, t)$, and the area fraction, $\phi_R(\pmb{x}, t)$, of free receptor R on the membrane. The solvent in the bulk is then given by $(1 - \phi)$, and the free lipid in the membrane

is given by $(1 - \phi_R - \phi_{PR})$. All volume fraction fields, in general, depend on position $\boldsymbol{x} = (x, y, z)$ and time t. The membrane is located at $z = 0$. We denote $\phi|_{z=0}$ as the volume fraction of protein at the boundary surface $S = \{(x, y, z = 0) \in V\}$. All area fraction fields depend on position $\boldsymbol{x} = (x, y)$.

The binding reaction between proteins in the bulk and receptors tagged to lipids in the membrane is

$$\alpha P + \beta R \rightleftharpoons PR, \tag{1}$$

where the effective component PR is composed of β receptors and α proteins, thus α and β are stoichiometric coefficients of the binding process; see Fig. 1a. In other words, our model can account for many proteins binding a receptor ($\alpha > \beta$), or a protein binding many receptors ($\alpha < \beta$). The molecular areas of the membrane components are denoted as v_{PR} for receptor-protein complex, v_R for the free receptor, and v_L for the free lipid. For binding to be volume conserving and thereby also ensure incompressibility in the layer adjacent the membrane, we have

$$\alpha v_P + \beta (h_R v_R) = h v_{PR}, \tag{2}$$

where v_P is the molecular volume of protein in the bulk. Moreover, h is the height of the effective membrane-bound layer and h_R denotes the height of the receptor protruding the membrane (see Fig. 1 for an illustration). When measuring experimentally the molecular volume of the receptor $\tilde{v}_R$, the height of the receptor can be estimated via $h_R = \tilde{v}_R / v_R$. The height of the effective membrane-bound layer h can then be determined via Eq. (2). Consistently, the molecular areas obey $v_{PR} = \beta v_R$. Note that $\alpha v_P / v$ are the number of solvent molecules removed from the effective bound layer when α proteins bind β receptors. Here, v denotes the molecular volume of the solvent in the bulk. In general, the binding scheme (Eq. (1)) should contain the solvent molecules removed from the layer increasing solvent amount in the bulk. However, the related changes in bulk volume and composition due to this release is negligible since the bound layer is microscopic.

Non-equilibrium thermodynamics with non-dilute surface binding

The conservation laws for membrane and bulk are

$$\partial_t \phi = -\nabla \cdot \boldsymbol{j}, \qquad\qquad x \in V, \tag{3a}$$

$$\partial_t \phi_{PR} = -\nabla_{||} \cdot \boldsymbol{j}_{PR} + r, \qquad\qquad x \in S, \tag{3b}$$

$$\partial_t \phi_R = -\nabla_{||} \cdot \boldsymbol{j}_R - r, \qquad\qquad x \in S, \tag{3c}$$

where $\boldsymbol{j}$, $\boldsymbol{j}_{PR}$ and $\boldsymbol{j}_R$ are the diffusive fluxes of bulk protein P, complex PR and free receptor R. Moreover, r denotes the binding rate between bulk and membrane. We define $\nabla = (\partial_x, \partial_y, \partial_z)$ as the gradient operator in the three-dimensional bulk and $\nabla_{||} = (\partial_x, \partial_y)$ on the two-dimensional membrane surface.

Binding relates the protein flux $\boldsymbol{n} \cdot \boldsymbol{j}$ normal to the membrane S to the binding rate r:

$$\boldsymbol{n} \cdot \boldsymbol{j} = \alpha \frac{v_P}{v_{PR}} r, \quad x \in S, \tag{3d}$$

where α is the stoichiometric coefficient of the binding protein. Please remember that v_P is the protein molecular volume and v_{PR} is the molecular area of the bound complex. Thus, v_P / v_{PR} provides the length scale related binding from three to two dimensions. The remaining bulk boundaries are considered to be impermeable for all components with the diffusive flux vanishing: $\boldsymbol{n} \cdot \boldsymbol{j} = 0$, $x \in \partial V, S_o$.

The total free energy F of the system is composed of different contributions: the free energy density $f_b(\phi)$ in the bulk V, the free energy density $f_m(\phi_{PR}, \phi_R)$ at the membrane surface S, and coupling between bulk and membrane $J(\phi|_{z=0}, \phi_{PR}, \phi_R)$. In specific, the total free energy reads:

$$F[\phi, \phi_{PR}, \phi_R] = \int_V d^3x \left[f_b(\phi) + \frac{1}{2}\kappa |\nabla \phi|^2 \right] + \int_S d^2x \left[f_m(\phi_{PR}, \phi_R) + J(\phi|_{z=0}, \phi_{PR}, \phi_R) \right.$$
$$\left. + \frac{1}{2}\kappa_R |\nabla_{||} \phi_R|^2 + \frac{1}{2}\kappa_{PR} |\nabla_{||} \phi_{PR}|^2 + \kappa_{PR\text{-}R} \nabla_{||} \phi_R \cdot \nabla_{||} \phi_{PR} \right]. \tag{4}$$

The free energy penalties due to concentration gradients are characterized by κ, κ_{PR}, κ_R, and $\kappa_{PR\text{-}R}$ in bulk and membrane, respectively.

In the following, we derive the kinetic equations that can relax toward thermodynamic equilibrium. In an isothermal system, the rate of change of the system entropy $S(t)$ can be expressed in terms of the integrated change of the free energy densities f_m, f_b in membrane S and bulk V respectively, the coupling free energies J, as well as the reactive free energy fluxes $\boldsymbol{j}_{r1}$ and $\boldsymbol{j}_{r2}$ through the boundaries of bulk and membrane, respectively *Zhao et al.* (*2024*):

$$
T\dot{S} = -\int_V d^3x\,\partial_t\left[f_b(\phi) + \frac{1}{2}\kappa|\nabla\phi|^2\right] - \int_S d^2x\,\partial_t\left[f_m(\phi_{PR},\phi_R) + J(\phi|_{z=0},\phi_{PR},\phi_R)\right]
$$
$$
-\int_S d^2x\,\partial_t\left[\frac{1}{2}\kappa_R|\nabla_\parallel\phi_R|^2 + \frac{1}{2}\kappa_{PR}|\nabla_\parallel\phi_{PR}|^2 + \kappa_{PR\text{-}R}|\nabla_\parallel\phi_{PR}|^2\right]
$$
$$
-\int_{\partial V} d^2x\,\boldsymbol{n}\cdot\boldsymbol{j}_{r1} - \int_{\partial S} dx\,\boldsymbol{t}\cdot\boldsymbol{j}_{r2}\,. \tag{5}
$$

Using the conservation laws Eqs. (3)(a-c) and boundary conditions (3d), defining the reactive free energy fluxes $\boldsymbol{j}_{r1} = \boldsymbol{j}\mu/v_P$ and $\boldsymbol{j}_{r2} = \boldsymbol{j}_{PR}\mu_{PR}/v_{PR} + \boldsymbol{j}_R\mu_R/v_R$, we obtain for the entropy production rate, where we could define a pair of thermodynamic fluxes and forces. Using Onsager principle, we get the constitutive relation between fluxes and forces. Substituting the well-defined fluxes into the continuity equation (3)(a,b,c), we obtain the governing equations of the system as follows

$$
\partial_t\phi = \nabla\cdot(\Lambda_P\nabla\mu), \qquad\qquad x\in V\,, \tag{6a}
$$
$$
\partial_t\phi_{PR} = \nabla\cdot(\Lambda_{PR}\nabla\mu_{PR}) + r\,, \qquad\qquad x\in S\,, \tag{6b}
$$
$$
\partial_t\phi_R = \nabla\cdot(\Lambda_R\nabla\mu_R) - r\,, \qquad\qquad x\in S\,, \tag{6c}
$$

with boundary conditions:

$$
\boldsymbol{n}\cdot\nabla\phi = -\kappa^{-1}\frac{\partial J}{\partial\phi|_{z=0}}\,, \qquad\qquad x\in S\,, \tag{7a}
$$
$$
\boldsymbol{n}\cdot\nabla\phi = 0\,, \qquad\qquad x\in S_o\,, \tag{7b}
$$
$$
-\boldsymbol{n}\cdot(\Lambda_P\nabla\mu) = \alpha\,\frac{v_P}{v_{PR}}\,r\,, \qquad\qquad x\in S\,, \tag{7c}
$$
$$
-\boldsymbol{n}\cdot(\Lambda_P\nabla\mu) = 0\,, \qquad\qquad x\in S_o\,. \tag{7d}
$$

We note that it is possible to have more dissipative fluxes on the boundaries S when the system evolves approaching the equilibrium state *Zhao et al.* (*2024*), e.g., $(\partial J/\partial\phi + \boldsymbol{n}\cdot\kappa\nabla\phi)\Lambda_k = \partial_t\phi|_{z=0}$, where Λ_k is the corresponding kinetic coefficient. Here, we only focus on the effects of binding rate r and thus set other fluxes to be zero for all times during the kinetics. The boundary condition $\partial J/\partial\phi + \boldsymbol{n}\cdot\kappa\nabla\phi = 0$ in Eq. (7)a characterizes the interactions with the membrane surface and thereby also determine wetting behavior of bulk condensates. The second condition specifies that the opposite side of membrane S_o are neutral, i.e., the free energy is not affected by increasing or decreasing the bulk volume fractions adjacent to S_o. Moreover, we did not consider the cross-coupling interaction in the membrane.

We consider the standard mobility scalings,

$$
\Lambda_P = \Lambda_{P,0}\,\phi\,(1-\phi)\,, \tag{8a}
$$
$$
\Lambda_{PR} = \Lambda_{PR,0}\,\phi_{PR}\,(1-\phi_{PR}-\phi_R)\,, \tag{8b}
$$
$$
\Lambda_R = \Lambda_{R,0}\,\phi_R\,(1-\phi_R-\phi_{PR})\,, \tag{8c}
$$

and consider for simplicity the mobility prefactors $\Lambda_{P,0}$, $\Lambda_{PR,0}$, and $\Lambda_{R,0}$ as constants.

The binding rate r obeys detailed balance of the rates *Weber et al.* (*2019*). However, by Onsager linear response thermodynamics *Zhao et al.* (*2024*), we can only obtain the binding rate to linear order in the thermodynamic force $\Delta\mu = \mu_{PR} - \beta\,\mu_R - \alpha\mu$. To account for a general non-linear binding

181 rate that also obeys detailed balance of the rates, we follow Ref. *Van Kampen* (*1973*) and write the
182 binding rate

$$r(\phi_{PR}, \phi_R, \phi) = -\Lambda \left[\exp \left((\alpha \mu + \beta \mu_R) / k_\mathrm{B} T \right) - \exp \left(\mu_{PR} / k_\mathrm{B} T \right) \right] , \tag{9}$$

183 where Λ is the binding rate coefficient. At thermodynamic equilibrium (Eqs. (13)), the binding rate
184 $r = 0$.

Thermodynamic equilibrium with non-dilute surface binding

186 When each diffusive flux and the binding rate defined above vanish to zero, the system is at ther-
187 modynamic equilibrium. At thermodynamic equilibrium, the total free energy F is minimal. There
188 are constraints for the minimization since the total number of proteins N and the total number of
189 receptors at the membrane N_R are conserved. The minimum of the free energy fulfils

$$0 = \delta \left\{ F - \lambda \left[\int_V d^3x\, \phi / v_P + \alpha \int_S d^2x\, \phi_{PR} / v_{PR} - N \right] - \lambda_R \left[\int_S d^2x \left(\phi_R / v_R + \beta\, \phi_{PR} / v_{PR} \right) - N_R \right] \right\} , \tag{10}$$

190 where λ and λ_R denote Lagrange multipliers corresponding to the two respective conservation
191 laws. Using Eq. (4), the variation of the total free energy is given as

$$\begin{aligned}
\delta F = &\int_V d^3x \left(\frac{\partial f}{\partial \phi} - \kappa \nabla^2 \phi \right) \delta\phi + \int_S d^2x \left(\frac{\partial f_m \left(\phi_{PR}, \phi_R \right)}{\partial \phi_{PR}} - \kappa_{PR} \nabla_{||}^2 \phi_{PR} - \kappa_{PR\text{-}R} \nabla_{||}^2 \phi_R + \frac{\partial J}{\partial \phi_{PR}} \right) \delta\phi_{PR} \\
&+ \int_S d^2x \left(\frac{\partial f_m \left(\phi_{PR}, \phi_R \right)}{\partial \phi_R} - \kappa_R \nabla_{||}^2 \phi_R - \kappa_{PR\text{-}R} \nabla_{||}^2 \phi_{PR} + \frac{\partial J}{\partial \phi_R} \right) \delta\phi_R + \int_S d^2x \left(\frac{\partial J}{\partial \phi |_{z=0}} \right) \delta\phi \\
&+ \int_{S, \partial V} d^2x \left(\kappa\, \boldsymbol{n} \cdot \nabla \phi \right) \delta\phi + \int_{\partial S} dx\, \boldsymbol{t} \cdot \left(\kappa_{PR} \nabla_{||} \phi_{PR} + \kappa_{PR\text{-}R} \nabla_{||} \phi_R \right) \delta\phi_{PR} \\
&+ \int_{\partial S} dx\, \boldsymbol{t} \cdot \left(\kappa_R \nabla_{||} \phi_R + \kappa_{PR\text{-}R} \nabla_{||} \phi_{PR} \right) \delta\phi_R .
\end{aligned} \tag{11}$$

192 We thus obtain the exchange chemical potentials of the proteins in the bulk μ, the protein-receptor
193 complex μ_{PR} and the free receptor μ_R in the membrane:

$$\mu \equiv v_P \frac{\delta F}{\delta \phi} = v_P \left(\frac{\partial f}{\partial \phi} - \kappa \nabla^2 \phi \right) , \tag{12a}$$

$$\mu_{PR} \equiv v_{PR} \frac{\delta F}{\delta \phi_{PR}} = v_{PR} \left(\frac{\partial f_m}{\partial \phi_{PR}} + \frac{\partial J}{\partial \phi_{PR}} - \kappa_{PR} \nabla_{||}^2 \phi_{PR} - \kappa_{PR\text{-}R} \nabla_{||}^2 \phi_R \right) , \tag{12b}$$

$$\mu_R \equiv v_R \frac{\delta F}{\delta \phi_R} = v_R \left(\frac{\partial f_m}{\partial \phi_R} + \frac{\partial J}{\partial \phi_R} - \kappa_R \nabla_{||}^2 \phi_R - \kappa_{PR\text{-}R} \nabla_{||}^2 \phi_{PR} \right) . \tag{12c}$$

194 Using Eqs. (12), we find that $\mu_{PR} = \alpha \lambda + \beta \lambda_R$, with $\lambda = \mu$ and $\lambda_R = \mu_R$. As a result, binding equilibrium
195 of the system composed of membrane and bulk is given as:

$$\alpha \mu + \beta \mu_R = \mu_{PR} . \tag{13a}$$

Here, the stoichiometric coefficients α and β reflect that effective component PR is composed of α
proteins and β receptors.

Once the protein binds to the membrane-bound receptor and its local concentration exceeds
a certain threshold, surface phase separation may occur, driven by the attractive interaction be-
tween protein–receptor complexes. When two phases, denoted as I and II, coexist in the mem-
brane, phase equilibrium holds:

$$\mu_{PR}^{\mathrm{I}} = \mu_{PR}^{\mathrm{II}}, \tag{13b}$$

$$\mu_R^{\mathrm{I}} = \mu_R^{\mathrm{II}}, \tag{13c}$$

$$f^{\mathrm{I}} - f^{\mathrm{II}} = \frac{\mu_{PR}^{\mathrm{I}}}{v_{PR}} \left(\phi_{PR}^{\mathrm{I}} - \phi_{PR}^{\mathrm{II}} \right) + \frac{\mu_R^{\mathrm{I}}}{v_R} \left(\phi_R^{\mathrm{I}} - \phi_R^{\mathrm{II}} \right) . \tag{13d}$$

The total amount of receptors in the membrane is conserved:

$$A\,\phi_R^0 = A^{\mathrm{I}}\left(\phi_{PR}^{\mathrm{I}} + \phi_R^{\mathrm{I}}\right) + \left(A - A^{\mathrm{I}}\right)\left(\phi_{PR}^{\mathrm{II}} + \phi_R^{\mathrm{II}}\right),$$ (13e)

where A^{I} is the surface condensate area in the membrane, $A = L^2$ is the total area of the membrane, and ϕ_R^0 is the initial area fraction of the receptor in the membrane.

We use the boundary conditions at the membrane S, the opposite boundary surface S_o, and the rest of the boundaries enclosing the bulk, ∂V, at thermodynamic equilibrium in Eq. (7). Boundary conditions (7) and equilibrium conditions (13) are composite of non-dilute binding equilibrium conditions. In the following section, we will discuss the non-dilute surface binding in detail based on these equilibrium conditions.

Interplay between surface binding and surface phase separation

Non-dilute surface binding

In this section, we derive the general binding isotherm for equilibrium surface binding under non-dilute conditions. Unlike the classical Langmuir model *Langmuir* (*1918*), which assumes idealized dilute conditions and no intermolecular interactions, here we incorporate activity coefficients to account for deviations from ideality, such as interactions between different kinds of molecules. This approach leads to a generalized form of the binding isotherm that reduces to the classical Langmuir model under appropriate limiting assumptions.

To capture non-dilute behavior, we express the chemical potentials in terms of standard chemical potentials and activity coefficients:

$$\mu = \mu_0 + k_{\mathrm{B}}T \ln\left(\gamma(\phi)\phi\right),$$ (14a)

$$\mu_R = \mu_{R,0} + k_{\mathrm{B}}T \ln\left(\gamma_R(\phi_R,\phi_{PR})\phi_R\right),$$ (14b)

$$\mu_{PR} = \mu_{PR,0} + k_{\mathrm{B}}T \ln\left(\gamma_{PR}(\phi_R,\phi_{PR})\phi_{PR}\right).$$ (14c)

where the terms $\gamma(\phi)$, $\gamma_R(\phi_R,\phi_{PR})$, and $\gamma_{PR}(\phi_R,\phi_{PR})$ are activity coefficients capturing non-ideal effects, such as intermolecular interactions or internal energies. The constants μ_0, $\mu_{R,0}$, and $\mu_{PR,0}$ are the corresponding standard chemical potentials that include the internal free energies of the components.

Substituting (14) into the binding equilibrium condition (13a), we rearrange to find,

$$\alpha \ln\phi + \beta \ln\phi_R = \ln\phi_{PR} + \Omega,$$ (15)

or equivalently,

$$\phi_{PR} = (\phi_R)^\beta\,(\phi)^\alpha\,e^{-\Omega},$$ (16a)

where

$$\Omega = \frac{\Delta\mu_0}{k_{\mathrm{B}}T} - \ln\left(\frac{(\gamma)^\alpha\,(\gamma_R)^\beta}{\gamma_{PR}}\right),$$ (16b)

$$\Delta\mu_0 = \mu_{PR,0} - \alpha\mu_0 - \beta\mu_{R,0},$$ (16c)

Equation (16a) represents a general non-dilute binding relation, where the effects of interactions at non-dilute conditions are accounted for by activity coefficients γ, γ_R and γ_{PR}. Moreover, $\Delta\mu_0$ denotes the differences of standard chemical potential.

To determine the surface coverage, we must also consider the conservation of the total amount of receptor on the membrane. Defining ϕ_R^0 as the total receptor coverage, in a spatially homogeneous system we have :

$$\phi_R^0 = \phi_{PR} + \phi_R,$$ (16d)

Figure 2. Non-dilute surface binding behaviors with varying surface and bulk compositions. (a) Binding fraction curves ($\theta = \phi_{PR}/\phi_R^0$) are shown for three different initial receptor area fractions: $\phi_R^0 = 1$ (solid line), $\phi_R^0 = 0.65$ (dashed line) and $\phi_R^0 = 10^{-5}$ (dotted line), under the assumption of no surface phase separation ($\chi_{PR\text{-}L} = 2$). (b) With a stronger interaction strength ($\chi_{PR\text{-}L} = 3.2$) and a fixed initial receptor area fraction ($\phi_R^0 = 0.65$), the homogeneous surface mixture (black line) undergoes phase separation into two distinct phases: a protein-rich phase (red line) and a protein-poor phase (blue line) as a function of the intermediate bulk volume fraction, ϕ. Additionally, when the bulk protein volume fraction (ϕ) exceeds the saturation level, a 3D bulk condensate forms (blue region) that coexist with 2D surface condensates on the membrane. (c) The phase diagram illustrates that intermediate receptor area fractions (ϕ_R^0) and large bulk protein volume fractions (ϕ) promote surface phase separation on the membrane. The basic parameter values are: $\omega_{PR} = -1$, $\chi = 2.5$, $n = n_{PR} = n_R = 1$, $\omega = \omega_R = 0$, $\chi_{P\text{-}L} = 0$, and $\chi_{PR\text{-}R} = 0$.

₂₃₃ where ϕ_R and ϕ_{PR} are the area fractions of free and bound receptor on the membrane. By com-
₂₃₄ bining the conservation law in Eq. (16d) with the general binding relation (16a), we arrive at a *gen-*
₂₃₅ *eralized non-dilute binding isotherm*. Upon choosing a specific form of the free energy densities f_b,
₂₃₆ f_m and J in Eq. (4), Eqs. (16) can only be solved numerically due to the non-linear dependence of
₂₃₇ the activity coefficients on all the volume fractions of the components at non-dilute conditions.

₂₃₈   The non-dilute binding isotherm can be written in a simpler form for the special case of an
₂₃₉ unimolecular binding process ($\alpha = \beta = 1$). Defining a quantity commonly studied in adsorption
₂₄₀ isotherms *Ayawei et al. (2017)*, the binding fraction $\theta = \phi_{PR}/\phi_R^0$, we obtain:

$$\theta = \frac{\phi}{K_L + \phi},\tag{17}$$

₂₄₁ where $K_L = \exp(\Omega)$ is a *generalized Langmuir coefficient*. Here, Ω is the generalized binding affinity
₂₄₂ given in Eq. (16b), which describes the binding energy per molecule at non-dilute conditions.

₂₄₃   In the ideal case where the binding sites are homogeneous and receptors are dilute, the effects
₂₄₄ of intermolecular interactions in the activity coefficients vanish in Eq. (17). As a result, one obtains
₂₄₅ the classical Langmuir isotherm *Langmuir (1918)*, and K_L reduces to the Langmuir constant $K_L = \exp(\Delta\mu_0/k_B T)$, that is determined solely by the dilute binding affinity $\Delta\mu_0$.
₂₄₆

₂₄₇   In summary, starting from the general equilibrium conditions and incorporating mass conser-
₂₄₈ vation (16d) and non-ideal interactions via activity coefficients, we have derived a general binding
₂₄₉ isotherm applicable to non-dilute conditions (16a). This result not only generalizes the theory of
₂₅₀ surface binding to non-dilute conditions recovering the classical Langmuir equation in the dilute
₂₅₁ limit, it also describes phase transitions at the binding surface leading to a heterogeneous distri-
₂₅₂ bution of bound molecules.

The non-dilute binding isotherm and surface phase separation

₂₅₄ In this section, we investigate the intricate behaviors of non-dilute surface binding by varying bulk
₂₅₅ and surface compositions. Using Flory-Huggins free energy densities for the bulk-membrane sys-
₂₅₆ tem (Appendix 1), we assume unimolecular binding ($\alpha = \beta = 1$) and examine the thermodynamic
₂₅₇ equilibrium satisfying the conditions of phase coexistence (13b)–(13d) and the generalized binding
₂₅₈ isotherm (17), we obtain

$$\Omega = \Delta\mu_0 + B(\phi, \phi_R, \phi_{PR}) - \alpha\,A(\phi, \phi_R, \phi_{PR}) - \beta\,C(\phi, \phi_R, \phi_{PR})\tag{18a}$$

259 with

$$A = -n \ln(1 - \phi) + n\,\chi\,(-2\phi)\,, \tag{18b}$$

$$B = -n_{PR} \ln(1 - \phi_R - \phi_{PR}) - n_{PR}\,\chi_{P\text{-}L}\,\phi_R + n_{PR}\,\chi_{PR\text{-}L}\left(-\phi_R - 2\phi_{PR}\right) + n_{PR}\chi_{PR\text{-}R}\,\phi_R, \tag{18c}$$

$$C = -n_R \ln(1 - \phi_R - \phi_{PR}) + n_R\,\chi_{P\text{-}L}\left(-2\phi_R - \phi_{PR}\right) - n_R\,\chi_{PR\text{-}L}\,\phi_{PR} + n_R\,\chi_{PR\text{-}R}\,\phi_{PR}, \tag{18d}$$

$$\Delta\mu_0 = \left(n_{PR}\,\chi_{PR\text{-}L} + n_{PR}\omega_{PR} - n_{PR} + 1\right)$$
$$- \alpha\left(n\,\chi + n\,\omega - n + 1\right) - \beta\left(n_R\,\chi_{P\text{-}L} + n_R\,\omega_R - n_R + 1\right). \tag{18e}$$

To align with physical constraints and simplify computational implementation, we set the basic
parameter values as follows: $\omega_{PR} = -1$, $\chi = 2.5$, $n = n_{PR} = n_R = 1$, $\omega = \omega_R = 0$, $\chi_{P\text{-}L} = 0$, and $\chi_{PR\text{-}R} = 0$.
Specifically, ω_{PR}, the internal free energy coefficient, denotes the binding affinity of proteins to the
membrane receptors. Setting $\omega_{PR} = -1$ reflects a moderate attractive interaction, consistent with
observed protein-receptor binding in similar systems. The parameter $\chi = 2.5$ ensures sufficient
phase separation tendencies in the bulk, while $n = n_{PR} = n_R = 1$ assumes comparable molecular
sizes between receptor-bound proteins, unbound receptors, and lipid molecules, simplifying the
modeling framework. Finally, $\chi_{P\text{-}L}$ and $\chi_{PR\text{-}R}$ are set to zero to focus on dominant interactions and
exclude secondary effects that do not significantly impact the observed behavior.

We first consider the scenario where phase separation does not occur on the surface, as the
attractive interactions between bound proteins ($\chi_{PR\text{-}L} = 2$) are insufficient to induce surface phase
separation. Under these assumptions, our analysis reveals that the binding fraction θ increases
monotonically with the bulk protein volume fraction ϕ in the non-dilute case (Fig. 2a) This behavior
qualitatively resembles the dilute case, which corresponds to a scenario where the total recep-
tor concentration is very low ($\phi_R^0 \ll 1$). However, in the non-dilute case, the binding fraction θ is
significantly higher than in the dilute case as the total receptor concentration ϕ_R^0 increases. This
enhancement is a direct consequence of non-dilute binding. At very high bulk protein volume frac-
tions (ϕ), the bulk undergoes phase separation, further altering the system's behavior.

We now examine the case where the bound proteins at the surface can undergo surface phase
separation, forming a protein-rich and a protein-poor phase leading to a heterogeneous distribu-
tion of bound proteins PR at the surface (see Fig. 2b). A sufficiently large bulk protein concentration
is needed to ensure enough proteins are bound to receptors at the surface. Moreover, interactions
need to be strong enough (e.g., $\chi_{PR\text{-}L} = 3.2$), e.g., the bound proteins attract each other sufficiently
strongly.

The effects of both the bulk protein volume fraction ϕ and the total amount of receptors on
the surface ϕ_R^0 are summarized as a phase diagram in Fig. 2c for a fixed value of $\chi_{PR\text{-}L} = 3.2$. It
shows that surface phase separation occurs in an intermediate range of both bulk protein and
total receptor area fraction ϕ_R^0. A too-large value of bulk protein volume fraction triggers phase
separation in the bulk independent of the receptor and surface binding. Moreover, a too-large
amount of receptors on the surface leads to a homogeneous state for volume fraction fractions
below bulk phase separation since the mixing entropy on the surface dominates. Similarly, if there
are too few receptors on the surface, the enthalpic gains related to the interaction among bound
proteins are also outcompeted by the mixing entropy. This overall indicates a reentrant behavior
of surface phase separation when varying the total receptor area fraction ϕ_R^0.

Effects of binding affinity and interactions on surface phase separation

We also investigated how the binding affinity (ω_{PR}) and the interaction strength between membrane-
bound proteins ($\chi_{PR\text{-}L}$) affect surface phase separation. We distinguish two scenarios (Fig. 3a): The
three-dimensional bulk is homogeneous, or the bulk is phase-separated.

In the absence of bulk phase separation, as shown in Fig. 3b, we find that upon increasing the
binding affinity (ω_{PR}) and membrane-bound interaction strength ($\chi_{PR\text{-}L}$) can induce surface phase
separation on the membrane, shifting the system from a homogeneous state to a phase-separated

Figure 3. Surface behavior with varying binding affinity and membrane-bound interaction strength. (a) Schematic illustrating bulk-membrane coupled system with surface condensates formed in the membrane S without and with bulk phase separation, respectively. (b) Phase diagram without bulk phase separation, indicating that strong binding affinity and (ω_{PR}) interaction strength ($\chi_{PR\text{-}L}$) among membrane-bound complexes promote surface phase separation. (c) If protein interaction in the bulk is strong and thus bulk phase separation occurs. With bulk condensates partially/completely wet on top of the membrane, the surface phase separation occurs in a much larger parameter regime. This illustrates that bulk protein properties, like interaction strength, also regulate the surface phase behavior. The basic parameter values we set in this study are: $\phi_R^0 = 0.5$, (a) $\chi = 0$, $\phi = c_{sat} = 0.145$, (b) $\chi = 2.5$, $\phi = c_{sat} = 0.145$. Here, c_{sat} represents the bulk volume fraction at the saturation level.

one. This result underscores the importance of binding affinity and membrane-bound complex
interactions as a driving force for surface phase separation.

When bulk phase is phase-separated, as depicted in Fig. 3c, the parameter range of (ω_{PR}) and
($\chi_{PR\text{-}L}$) leading to surface separation broadens significantly. Under these conditions, surface phase
separation and partial wetting are observed even at lower binding affinities (ω_{PR}) compared to
conditions without bulk phase separation. This finding suggests that interaction strength between
proteins in the bulk, which gives rise to bulk phase separation, tunes surface phase separation.
However, bulk phase separation is not a prerequisite for surface phase separation. Surface phase
separation is dominated by the ability to recruit proteins from the bulk through binding that is in
turn controlled by the binding affinity (ω_{PR}) and the interaction strength between bound proteins
($\chi_{PR\text{-}L}$).

Mechanism in tight junction formation

Figure 4. Illustration of tight junction initiation and in vitro reconstitution system. (a) Schematic representation of tight junction initiation during cell-cell contact. Upon contact between Cell 1 and Cell 2, receptors (magenta) on the plasma membrane (PM) interact with scaffold proteins such as ZO1 (green) to form condensates that contribute to tight junction formation *Beutel et al.* (*2019*). (b) In vitro reconstitution system showing the binding dynamics of ZO1 proteins (green) with receptors (magenta) on a supported lipid bilayer. The His(10)-tagged receptors are anchored to DGS-NTA (Ni) lipid via His-Ni interaction and can bind and unbind with the ZO1 proteins in the bulk.

313   Tight junctions are important membrane-associated structures for epithelial and endothelial

barrier formation. The phase separation of scaffold proteins ZO1 has been reported to be essential for tight junction assembly *Beutel et al.* (*2019*). However, how binding interplays with the spatiotemporal phase separation of ZO1 at the cell-cell contact site on the plasma membrane is not well understood. At the cell-cell interface, tight junction receptors including the Claudin family, the CTX family (JAMs) and the MARVEL family (Occludin, Tricellulin, MARVELD3), form multiple *trans-* and *cis-* interactions to mediate cell adhesion (Fig. 4a). Genetic mutation in these *trans-* and *cis-* interactions has been shown to induce tight junction and epithelial barriers deficiency *Kostrewa et al.* (*2001*); *Zhao et al.* (*2018b*); *Heinemann and Schuetz* (*2019*). We hypothesise that receptor oligomerization at the cell-cell interface drives ZO1 surface phase separation and tight junction formation. To test this idea, we use our theoretical model developed above to understand the mechanism for surface phase separation involved in the formation of tight junctions. We consider ZO1 as the scaffold proteins in the bulk, Claudin2 C terminal (CLDN2-ICD, referred as 1-mer hereafter), which is one of the essential receptors for tight junction function, as the receptors anchored in the membrane. CDLN2-ICD were tagged with His tag and anchored to the membrane with DGS-NTA (Ni) lipid as a mimic of the membrane-associated receptors via His-Ni interaction (Fig. 4b). To mimic the receptor oligomerization happening at the cell-cell contact site, ySMF domain, which is known to form stable 14-mer *Collins et al.* (*2003*), was introduced to the CDLN2-ICD construct to get 14-meric receptors (referred as 14-mer hereafter). Scaffold proteins ZO1 were then added to the bulk. See Appendix 4 for methods and materials used in the *in vitro* experiments.

Now we use our theory for non-dilute binding to membrane-like surfaces and surface phase separation of bound proteins to model an in vitro reconstitution system for tight junction initiation. We will show that the theory will be instrumental to characterize binding affinities at non-dilute conditions and decipher the role of the oligomerization state of the receptor. As a crucial step in this analysis, we estimate the fundamental model parameter values for ZO1 protein binding to claudin receptors by integrating image analysis and experimental data (see Appendix 5 for details). These parameter values, which form the basis of subsequent modeling, are summarized in Table 1.

Receptor Oligomerization aids surface Phase Transitions

We analyzed how the oligomerization state of the receptor affects the surface phase transition. In this study, we used two kinds of receptor oligomerization states: 1-mer and 14-mer. By fitting our model (13) to the experimental data (see Appendix 6) for both receptor states (Fig. 5c and 5d and Fig. 6c and 6d), we determined the changes in receptor binding affinity ω_{PR} and self-interaction parameter $\chi_{PR\text{-}L}$ of membrane-bound ZO1 complexes. A key finding is that the binding affinity of the 14-meric receptor to ZO1 protein ($\omega_{PR} = 3.83$, Fig. 6) is much stronger compared to the monomeric receptor ($\omega_{PR} = -1.09$, Fig. 5). Furthermore, we found that the lower binding affinity of the 1-mer is dominantly caused by a reduced binding fraction, i.e., only a small fraction of 1-mer receptors can bind to ZO1 (see Appendix 7 for details); a mechanism previously reported in Ref. *Erlendsson et al.* (*2019*). Moreover, from the fits, we obtained that the bound ZO1-14-mer complexes strongly attract each other, i.e., they have a positive self-interaction parameter (Fig. 6). While the fitted self-interaction of the ZO1-1-mer ($\chi_{PR\text{-}L} = -0.4$) complex was lower compared to the 14-mer($\chi_{PR\text{-}L} = 1.09$), a wide range of self-interaction values provide similar good fits (blue shade in Appendix 6-fig. 1c), indicating that in the low-affinity binding regime of the 1-mer receptor self-interactions at the surface play a minor role (see Appendix 6-fig. 1c and its related discussion).

In summary, our results suggest that receptor oligomerization promotes surface phase separation through two synergistic mechanisms. First, increased binding affinity leads to a higher concentration of ZO1 on the membrane. Second, the self-interaction of the receptor-protein complex lowers the saturation concentration required for surface phase separation.

Using the parameters obtained from the fits, we determine the phase diagram for the monomeric (Fig. 5e) and 14-meric receptors (Fig. 6e). These phase diagrams show that there is no surface phase transition for monomeric receptors even for higher ZO1 concentrations, while there is a large region of surface phase separation for the 14-meric receptors.

Figure 5. Surface stays homogeneous with 1-mer receptor anchored on the membrane. (a) Fluorescence images of ZO1 and 1-mer at 1-mer receptor titrations, ranging from 19386 to 102689 μm^{-2}. (b) Fluorescence images of ZO1 and 1-mer showing the effect of ZO1 titration (0 to 800 nM) on 1-mer binding. Scalar bars in the bottom panels represent 2 μm. (c) The linear relationship between membrane-bound ZO1 density and receptor density on the membrane. This is because, in this study, the binding is saturated for all the receptor densities. In this figure, the dots with error bars represent the experimental data (quantified from figure (a)), and the solid line depicts the theoretical prediction. (d) Curve of membrane-bound ZO1 density as a function of ZO1 bulk density compared between experimental data (points) and theoretical prediction (line). The plateau is obtained from the binding fraction limit for 1-mer, $c = 0.014$. See Appendix 7 for details. The data is a fit to the membrane-bound ZO1 density observed in panel (b). (e) Phase diagram illustrating the homogeneous surface state across a physiological range of receptor and ZO1 densities.

Based on the agreements between experiments and our thermodynamic model, we conclude that a surface phase transition drives the formation of ZO1-rich surface condensates on membrane-like surfaces. The surface phase transition is controlled by the receptor's oligomerization state, suggesting cell adhesion-induced oligomerization could act as the key switch for ZO1 surface condensation at cell-cell contacts.

Spatial Pattern Formation of ZO1 Surface Condensates

When proteins bound to the surface undergo phase separation, multiple condensates form and exhibit a slow ripening kinetics. Theoretical results on surface binding, surface condensates growth and ripening can be obtained from numerically solving the dynamic equations (21) in a three-dimensional bulk volume V with surface S (Figure 7(a), see Appendix 2 and Appendix 3 for details). Both theoretical predictions and reconstitution experiments reveal an initial stage where surface condensates form and grow rapidly, followed by a prolonged ripening phase characterized by slow kinetics (see Appendix 8, Fig. 1). During this ripening stage, the average condensate areas predicted by the theory align well with experimental observations. Importantly, over a time course of approximately 15 minutes, the amount of bound proteins remains nearly constant as the surface condensates ripen, with only a slight reduction in the number of condensates. To further validate the model, we compared the condensate patterns from experiments and theory at time $t = 15$ min for different receptor titration levels ranging from 25 to 3800 PN/μm^2 and ZO1 titration levels from 4 to 100 nM (Fig. 7(b,c)). The patterns show quantitative agreement between theory and experiments, providing strong evidence that the theoretical framework effectively cap-

Figure 6. Surface condensates form with 14-mer Receptor on the membrane (a) Fluorescence images showing the distribution of 14-mer receptors and ZO1 across a range of receptor titrations, from 7 to 2744 μm^{-2}. Surface phase separation occurs at intermediate receptor densities. Too few or too many receptors result in a homogeneous membrane phase: a dilute phase at low receptor densities and a dense phase at high receptor densities. (b) Fluorescence images of 14-mer receptors and ZO1 as a function of increasing ZO1 bulk concentration, from 4 to 14000 nM, demonstrating the emergence of phase-separated domains. Since receptor density remains constant during this titration, the area of surface condensates does not change with increasing ZO1 bulk concentration. Once the ZO1 bulk concentration exceeds the saturation level, bulk condensates form in the presence of surface condensates (refer to Fig. 3a for illustration). Scalar bars in the bottom panels represent 2 μm. (c) Membrane-bound ZO1 density as a function of receptor density on the membrane, showing how receptor density influences surface phase behavior. In this figure, the dots with error bars represent the experimental data (quantified from figure (a)), and the solid line depicts the theoretical prediction. (d) Membrane-bound ZO1 density dependence on ZO1 bulk concentration, comparing experimental data with theoretical predictions. The data is a fit to the membrane-bound ZO1 density observed in figure (b), revealing a transition from a homogeneous surface phase to a two-phase coexisting state (protein-poor and protein-rich phases). (e) Phase diagram with respect to the receptor density on the membrane and Bulk ZO1 density, indicating that 14-mer receptor promotes surface phase separation.

tures the non-dilute surface binding mechanisms and the condensate formation process on the surface. Furthermore, we observed that the receptor density on the membrane significantly influences the spatial patterns of surface condensates. At lower receptor densities, condensates tend to be sparse and less connected, while higher receptor densities promote larger and more interconnected condensates, consistent with predictions from the theoretical model (Fig. 7(b,c)).

Conclusion

In this study, we developed a comprehensive theoretical framework to describe the binding of soluble scaffold proteins to lipid membranes via interactions with receptor proteins embedded in the membrane. The framework integrates a kinetic model based on non-equilibrium thermodynamics and equilibrium conditions derived from thermodynamic theory for non-dilute surface binding, leading to a generalized binding isotherm. This isotherm extends classical adsorption models by accounting for the non-dilute nature of surface-bound complexes and their interactions.

Using this framework, we explored the behaviors of non-dilute surface binding under varying bulk and surface compositions, binding affinities, and interaction strengths between membrane-bound complexes. The model also enabled us to simulate and analyze surface phase transitions in

Figure 7. Snapshots of ZO1 surface condensates from simulation (top) and experiments (bottom) when titrating the 14-mer receptor and ZO1 bulk concentration. (a) Schematic of the system in our theoretical model, with 3D bulk volume V and membrane surface S, where surface condensates formed via surface phase separation. (b) Snapshots of ZO1 from simulation (top row) and experimental observations (bottom) under varying receptor titration levels, from 25 to 3800 μm^{-2}, at 15 mins. (c) Snapshots of ZO1 from simulation (top row) and experimental observations under varying ZO1 concentration levels, from 4 to 100 nM, at 15 min. The experimental results match well with the theoretical predictions from perspectives of both condensate pattern and condensate size. Scale bars in the bottom panels represent 2 μm. we set the diffusivity coefficient value as $D_i = 0.01\,\mu m^2 s^{-1}$, $i = P, PR, R$. The gradient coefficients $\kappa = \kappa_{PR} = \kappa_R = 2 \times 10^{-5}\mu m^{-1}$.

an in vitro reconstitution system for tight junction initiation. A critical insight from this work is the role of receptor oligomerization in facilitating surface phase transitions, highlighting its importance in membrane-associated condensation phenomena.

Our findings demonstrate that receptor density significantly influences the spatial patterns of surface condensates. At lower receptor densities, condensates are sparse and less connected, whereas higher receptor densities promote larger and more interconnected structures. Moreover, the patterns observed in simulations show quantitative agreement with experimental results, providing strong evidence that the theoretical framework accurately captures the mechanisms of non-dilute surface binding and condensate formation processes.

In summary, this work bridges theoretical modeling and experimental validation, advancing our understanding of non-dilute surface binding and phase separation on biological membranes. This framework offers a versatile tool for further investigating multicomponent systems, membrane curvature effects, and the dynamic organization of cellular interfaces, with potential applications in both biological research and biomimetic design.

Acknowledgments

We thank I. LuValle-Burke for insightful discussions on the binding proteins. We thank Lars Hubatsch on the discussion related to the Tight Juction. We specially thank Frank Jülicher for the collaboration on the fundamental theory on prewetting and surface phase separation (Ref. *Zhao et al.* (*2021*, 2024); *Liese et al.* (*2023*)). X. Zhao acknowledges the "FoSE New Researchers Grant" of the University of Nottingham Ningbo China for financial support. D. Sun acknowledges the support from the PEPC facility at MPI-CBG for protein purification. D. Sun was supported by a seed grant of the DFG funded "Physics of Life" Excellent Cluster at TU-Dresden. C. Weber and A. Honigmann acknowledge the SPP 2191 "Molecular Mechanisms of Functional Phase Separation" of the German Science Foundation for financial support and for providing an excellent collaborative environment.

References

Ayawei N, Ebelegi AN, Wankasi D. Modelling and interpretation of adsorption isotherms. Journal of Chemistry. 2017; 2017(1):3039817. doi: 10.1155/2017/3039817.

Beutel O, Maraspini R, Pombo-García K, Martin-Lemaitre C, Honigmann A. Phase separation of zonula occludens proteins drives formation of tight junctions. Cell. 2019 10; 179:923–936.e11. doi: 10.1016/j.cell.2019.10.011.

Boija A, Klein IA, Sabari BR, Dall'Agnese A, Coffey EL, Zamudio AV, Li CH, Shrinivas K, Manteiga JC, Hannett NM, Abraham BJ, Afeyan LK, Guo YE, Rimel JK, Fant CB, Schuijers J, Lee TI, Taatjes DJ, Young RA. Transcription factors activate genes through the phase-separation capacity of their activation domains. Cell. 2018 12; 175. doi: 10.1016/j.cell.2018.10.042.

Brangwynne CP, Eckmann CR, Courson DS, Rybarska A, Hoege C, Gharakhani J, Jülicher F, Hyman AA. Germline P granules are liquid droplets that localize by controlled dissolution/condensation. Science. 2009 5; 324(5935):1729–1732. doi: 10.1126/science.1172046.

Brangwynne CP, Mitchison TJ, A HA. Active liquid-like behavior of nucleoli determines their size and shape in Xenopus laevis oocytes. Proceedings of the National Academy of Sciences. 2011 1; 108(11):4334–4339. doi: 10.1073/pnas.1017150108.

CASE LB, ZHANG X, DITLEV JA, , ROSEN MK. Stoichiometry controls activity of phase-separated clusters of actin signaling proteins. Science. 2019; 363(6431):1093–1097. doi: 10.1126/science.aau6313.

CJ D, R P. P-bodies and stress granules: possible roles in the control of translation and mRNA degradation. Cold Spring Harbor Perspectives in Biology. 2012 7; 4. doi: 10.1101/cshperspect.a012286.

Collins BM, Cubeddu L, Naidoo N, Harrop SJ, Kornfeld GD, Dawes IW, Curmi PMG, Mabbutt BC. Homomeric ring assemblies of eukaryotic Sm proteins have affinity for both RNA and DNA: crystal structure of an oligomeric complex of yeast SmF*. Journal of Biological Chemistry. 2003; 278(19):17291–17298. doi: 10.1074/jbc.M211826200.

Erlendsson S, Thorsen TS, Vauquelin G, Ammendrup-Johnsen I, Wirth V, Martinez KL, Teilum K, Gether U, Madsen KL. Mechanisms of PDZ domain scaffold assembly illuminated by use of supported cell membrane sheets. eLife. 2019 1; 8:e39180. doi: 10.7554/eLife.39180.

Fritsch AW, Diaz-Delgadillo AF, Adame-Arana O, Hoege C, Mittasch M, Kreysing M, Leaver M, Hyman AA, Jülicher F, Weber CA. Local thermodynamics govern formation and dissolution of Caenorhabditis elegans P granule condensates. Proceedings of the National Academy of Sciences. 2021 7; 118(37):e2102772118. doi: 10.1073/pnas.2102772118.

Gall JG. The centennial of the Cajal body. Nature Reviews Molecular Cell Biology. 2003 12; 4:975–980. doi: 10.1038/nrm1262.

GrandPre T, Pyo AGT, Wingreen NS. Membrane wetting by biomolecular condensates is facilitated by mobile tethers. bioRxiv. 2024; doi: 10.1101/2024.12.04.626804.

Heinemann U, Schuetz A. Structural features of tight-junction proteins. International Journal of Molecular Sciences. 2019; 20(23):6020. doi: 10.3390/ijms20236020.

Jawerth L, Fischer-Friedrich E, Saha S, Wang J, Franzmann T, Zhang X, Sachweh J, Ruer M, Ijavi M, Saha S, Mahamid J, Hyman AA, Jülicher F. Protein condensates as aging Maxwell fluids. Science. 2020; 370(6522):1317–1323. doi: 10.1126/science.aaw4951.

JR B, R P. Eukaryotic stress granules: the ins and outs of translation. Molecular Cell. 2009 12; 36(6):932–941. doi: 10.1016/j.molcel.2009.11.020.

Kostrewa D, Brockhaus M, D'Arcy A, Dale GE, Nelboeck P, Schmid G, Mueller F, Bazzoni G, Dejana E, Bartfai T, Winkler FK, Hennig M. X-ray structure of junctional adhesion molecule: structural basis for homophilic adhesion via a novel dimerization motif. The EMBO Journal. 2001; 20(16):4391–4398. doi: 10.1093/emboj/20.16.4391.

Langmuir I. The adsorption of gases on plane surfaces of glass, mica and platinum. Journal of the American Chemical Society. 1918 6; 40(9):1361–1403. doi: 10.1021/ja02242a004.

Li P, Banjade S, Cheng HC, Kim S, Chen B, Guo L, Llaguno M, Hollingsworth JV, King DS, Banani SF, Russo PS, Jiang QX, Nixon BT, Rosen MK. Phase transitions in the assembly of multivalent signalling proteins. Nature. 2012; 483(7389):336–340. doi: 10.1038/nature10879.

Li R, Li T, Lu G, Cao Z, Chen B, Wang Y, Du J, Li P. Programming cell-surface signaling by phase-separation-controlled compartmentalization. Nature Chemical Biology. 2022 11; 18(12):1351–1360. doi: 10.1038/s41589-022-01192-3.

Liese S, Zhao X, Weber CA, Jülicher F. Chemically active wetting. arXiv:231207239. 2023 12; doi: 10.48550/arXiv.2312.07239.

Lu Y, Wu T, Gutman O, Lu H, Zhou Q, Henis YI, Luo K. Phase separation of TAZ compartmentalizes the transcription machinery to promote gene expression. Nature Cell Biology. 2020 3; 22(4):453–464. doi: 10.1038/s41556-020-0485-0.

Mahen R, Venkitaraman AR. Pattern formation in centrosome assembly. Current Opinion in Cell Biology. 2012 2; 24(1):14–23. doi: 10.1016/j.ceb.2011.12.012.

Otani T, Nguyen TP, Tokuda S, Sugihara K, Sugawara T, Furuse K, Miura T, Ebnet K, Furuse M. Claudins and JAM-A coordinately regulate tight junction formation and epithelial polarity. Journal of Cell Biology. 2019 08; 218(10):3372–3396. doi: 10.1083/jcb.201812157.

Pombo-García K, Adame-Arana O, Martin-Lemaitre C, Jülicher F, Honigmann A. Membrane prewetting by condensates promotes tight-junction belt formation. Nature. 2024 8; 632:647–655. doi: 10.1038/s41586-024-07726-0.

Riback JA, Katanski CD, Kear-Scott JL, Pilipenko EV, Rojek AE, Sosnick TR, Drummond DA. Stress-triggered phase separation is an adaptive, evolutionarily tuned response. Cell. 2017 3; 168:1028–1040. doi: 10.1016/j.cell.2017.02.027.

Rouches M, Veatch SL, Machta BB. Surface densities prewet a near-critical membrane. Proceedings of the National Academy of Sciences. 2021; 118(40):e2103401118. doi: 10.1073/pnas.2103401118.

Snead WT, Jalihal AP, Gerbich TM, Fritsch AW, Higgs HN, Gladfelter AS. Membrane surfaces regulate assembly of ribonucleoprotein condensates. Nature Cell Biology. 2022; 24:461–470. doi: 10.1038/s41556-022-00882-3.

Su X, Ditlev JA, Hui E, Xing W, Banjade S, Okrut J, King DS, Taunton J, Rosen MK, Vale RD. Phase separation of signaling molecules promotes T cell receptor signal transduction. Science. 2016 4; 352(6285):595–599. doi: 10.1126/science.aad9964.

Van Kampen N. Nonlinear irreversible processes. Physica. 1973; 67(1):1–22.

Wang B, Zhang L, Dai T, Qin Z, Huasong Lu LZ, Zhou F. Liquid–liquid phase separation in human health and diseases. Signal Transduction and Targeted Therapy. 2021 8; 6(1):290. doi: 10.1038/s41392-021-00678-1.

Weber CA, Zwicker D, Jülicher F, Lee CF. Physics of active emulsions. Reports on Progress in Physics. 2019 4; 82(6):064601. doi: 10.1088/1361-6633/ab052b.

Wiegand T, Hyman AA. Drops and fibers — how biomolecular condensates and cytoskeletal filaments influence each other. Emerging Topics in Life Sciences. 2020 10; 4(3):247–261. doi: 10.1042/ETLS20190174.

Wilhelm J, Kühn S, Tarnawski M, Gotthard G, Tünnermann J, Tänzer T, Karpenko J, Mertes N, Xue L, Uhrig U, Reinstein J, Hiblot J, Johnsson K. Kinetic and structural characterization of the self-labeling protein tags HaloTag7, SNAP-tag, and CLIP-tag. Biochemistry. 2021 8; 60(33):2560–2575. doi: 10.1021/acs.biochem.1c00258.

Xiao Q, McAtee CK, Su X. Phase separation in immune signalling. Nature Reviews Immunology. 2022 4; 22(3):188–199. doi: 10.1038/s41577-021-00572-5.

Zeng M, Chen X, Guan D, Xu J, Wu H, Tong P, Zhang M. Reconstituted Postsynaptic Density as a Molecular Platform for Understanding Synapse Formation and Plasticity. Cell. 2018; 174:1172 – 1187.e16. doi: 10.1016/j.cell.2018.06.047.

Zeng M, Díaz-Alonso J, Ye F, Chen X, Xu J, Ji Z, Nicoll RA, Zhang M. Phase separation-mediated TARP/MAGUK complex condensation and AMPA receptor synaptic transmission. Neuron. 2019 11; 104:529–543.e6. doi: 10.1016/j.neuron.2019.08.001.

Zhao J, Yang X, Li J, Wang Q. Energy stable numerical schemes for a hydrodynamic model of Nematic liquid crystals. SIAM Journal on Scientific Computing. 2016; 38(5):A3264–A3290. doi: 10.1137/15M1024093.

Zhao J, Yang X, Gong Y, Zhao X, Yang X, Li J, Wang Q. A general strategy for numerical approximationss of non-equilibrium models-part I: thermodynamic systems. International Journal of Numerical Analysis & Modeling. 2018 8; 15(6):884–918. http://global-sci.org/intro/article_detail/ijnam/12613.html.

Zhao J, Krystofiak ES, Ballesteros A, Cui R, Van Itallie CM, Anderson JM, Fenollar-Ferrer C, Kachar B. Multiple claudin–claudin cis interfaces are required for tight junction strand formation and inherent flexibility. Communications Biology. 2018; 1. doi: 10.1038/s42003-018-0051-5.

Table 1. Model parameter value and their dimensionless values taken from the experimental data.
We note that the molecule sizes v_{PR}, v_R are values for 14-mer receptor. All other parameter values are shared in all the studies.

Parameter name	Symbol	Value [units]	Value
Characteristic scales			
Time scale	$t_0 = \Lambda^{-1}$	100 [s]	1
Length scale	l	10 [μm]	1
Receptor valency			
Number of receptors per ZO1 bound	β	v_{PR}/v_R	1
Molecular area/volume			
Molecular area of free lipid in the membrane	v_L	$1.53^2\pi[nm^2]$	$1.53^2\pi/l^2$
Molecular area of bound ZO1 in the membrane	v_{PR}	$(9.2/2)^2\pi[nm^2]$	$(9.2/2)^2\pi/l^2$
Molecular volume of solvent in the bulk	v	$\frac{4\pi}{3}(9.2/2)^3[nm^3]$	$\frac{4\pi}{3}(9.2/2)^3/l^3$
Molecular volume of scaffold in the bulk	v_P	$\frac{4\pi}{3}(9.2/2)^3[nm^3]$	$\frac{4\pi}{3}(9.2/2)^3/l^3$
Interaction parameter			
between free receptor and bound ZO1 in the membrane	$\chi_{PR\text{-}R}$	0 [$k_B T$]	0
between free receptor and free lipid	$\chi_{P\text{-}L}$	0 [$k_B T$]	0
between ZO1 and solvent in the bulk	χ	0 [$k_B T$]	0
Conformational coefficient			
Gradient coefficient of bound ZO1 in the membrane	κ_{PR}	2×10^{-5} [μm^{-1}]	2×10^{-4}
Gradient coefficient of free receptor in the membrane	κ_R	2×10^{-5} [μm^{-1}]	2×10^{-4}
Gradient coefficient of scaffolds in the bulk	κ	2×10^{-5} [μm^{-1}]	2×10^{-4}
Mobility coefficients			
Diffusion coefficient of scaffold in the bulk	$\Lambda_{P,0}k_B T$	0.01 [$\mu m^2 s^{-1}$]	0.01
Diffusion coefficient of bound scaffold in the membrane	$\Lambda_{PR,0}k_B T$	0.01 [$\mu m^2 s^{-1}$]	0.01
Diffusion coefficient of free receptor in the membrane	$\Lambda_{R,0}k_B T$	0.01 [$\mu m^2 s^{-1}$]	0.01

Zhao X, Wang Q. A second order fully-discrete linear energy stable scheme for a binary compressible viscous
fluid model. Journal of Computational Physics. 2019 10; 395:382–409. doi: 10.1016/j.jcp.2019.06.030.

Zhao X, Bartolucci G, Honigmann A, Jülicher F, Weber CA. Thermodynamics of wetting, prewetting and surface
phase transitions with surface binding. New Journal of Physics. 2021 12; 23:123003. doi: 10.1088/1367-
2630/ac320b.

Zhao X, Liese S, Honigmann A, Jülicher F, Weber CA. Theory of wetting dynamics with surface binding. New
Journal of Physics. 2024 9; 26:103025. doi: 10.1088/1367-2630/ad80bb.

Zheng Q, Chen Y, Chen D, Zhao H, Feng Y, Meng Q, Zhao Y, Zhang H. Calcium transients on the ER surface
trigger liquid-liquid phase separation of FIP200 to specify autophagosome initiation sites. Cell. 2022 10;
185(22):4082–4098.e22. doi: 10.1016/j.cell.2022.09.001.

Appendix 1

Free energies

To calculate equilibrium conditons (13), we define free energies for the bulk-membrane systems characterizing the energy of the polymer mixtures in the bulk and membrane, respectively.

We use Flory-Huggins free energies in the bulk and membrane, respectively:

$$f_b(\phi) = \frac{k_B T}{v}\left[\frac{\phi}{n}\ln\phi + (1-\phi)\ln(1-\phi) + \chi\phi(1-\phi) + \omega\phi \right], \tag{19a}$$

$$f_m(\phi_{PR}, \phi_R) = \frac{k_B T}{v_L}\left[\frac{\phi_{PR}}{n_{PR}}\ln\phi_{PR} + \frac{\phi_R}{n_R}\ln\phi_R + (1-\phi_{PR}-\phi_R)\ln(1-\phi_{PR}-\phi_R) \right. \tag{19b}$$

$$+ \chi_{PR\text{-}R}\phi_{PR}\phi_R + \chi_{PR\text{-}L}\phi_{PR}(1-\phi_{PR}-\phi_R) + \chi_{P\text{-}L}\phi_R(1-\phi_{PR}-\phi_R)$$

$$\left. + \omega_{PR}\phi_{PR} + \omega_R\phi_R \right].$$

We take the internal free energy of proteins in the bulk as ω, denote the internal free energies of receptor-protein complex and free receptor as ω_{PR} and ω_R. We note that v_L represents the area of an effective free lipid patch to avoid considering molecules of very different sizes. As a consequence, to obtain the actual Flory-Huggins parameters corresponding to the lipid-complex interaction, it need to be rescaled by the number of lipids in each lipid patch.

Furthermore, we consider the coupling free energy between the bulk and membrane mixtures. In this study, we focus on the surface phase separation, not the extended prewetting layer or wetting condensates on top of the membrane-bound layer. The coupling free energy only influences on the wetting and prewetting and thus for simplicity, we set coupling free energy as zero in this study. See *Zhao et al.* (*2024*) for more complex coupling free energies.

If the free lipid area fraction is zero ($\phi_l = 0$), our model reduces to the binary mixture in the model discussed recently *Zhao et al.* (*2021*). Without receptors in the membrane ($\phi_R + \phi_{PR} = 0$), our model yields two decoupled systems: a two-dimensional membrane and a three-dimensional binary mixture in the bulk. If $\phi_l > 0$ and $\phi_R > 0$, our model corresponds to a ternary mixture in the membrane coupled to a binary mixture in the bulk. This model accounts for one more component, i.e., the receptor-protein complex in comparison to the thermodynamic model developed in the paper *Rouches et al.* (*2021*). In addition, our model describes the binding/unbinding events of proteins to the free receptors (or tethers) anchored in the membrane.

We consider a finite system where the bulk has a volume $V = L_z L^2$, and L_z denotes the height orthogonal to the membrane surface which has a surface area of L^2. For simplicity, we set $L_z = L$ in this study.

Appendix 2

Non-dimensionalization of the kinetic model

To solve equations (6) with boundary conditions (7) numerically we rescale length $x \rightarrow x/\ell$, where $\ell = 10\,\mu m$ is the experimental field of view, and rescale time $t \rightarrow t \cdot \Lambda^{-1}$. The non-dimensional quantities are:

$$\tilde{J} = J\frac{v_L}{k_B T}\,, \quad \tilde{\kappa}_{PR} = \frac{1}{\ell^2}\kappa_{PR}\frac{v_{PR}}{k_B T} = \frac{v_{PR}}{\ell^2}\,, \quad \tilde{\kappa} = \frac{1}{\ell^2}\kappa\frac{v_P}{k_B T} = \frac{v_P^{2/3}}{\ell^2}\,, \quad \bar{\kappa} = \kappa\frac{v_L}{k_B T}\frac{1}{\ell} = \frac{v_L}{v_P^{1/3}\ell}\,, \quad (20)$$

where $\tilde{J}$ is the non-dimensional binding rate between membrane and bulk. $\tilde{\kappa}_s$, $\tilde{\kappa}$, and $\bar{\kappa}$ are the dimensional gradient coefficients in the 2D membrane, 3D bulk and bulk boundaries adjacent to the membrane.

We achieve the following non-dimensional equations:

$$\partial_t \phi = \mathcal{D}_P \nabla \cdot (\phi(1 - \phi)\nabla\tilde{\mu})\,, \qquad\qquad x \in V\,, \quad (21a)$$

$$\partial_t \phi_{PR} = \mathcal{D}_{PR} \nabla \cdot (\phi_{PR}(1 - \phi_{PR} - \phi_R)\nabla\tilde{\mu}_{PR}) - \left[\exp\left(\alpha\,\tilde{\mu} + \beta\,\tilde{\mu}_R\right) - \exp\left(\tilde{\mu}_{PR}\right)\right]\,, \quad x \in S\,, \quad (21b)$$

$$\partial_t \phi_R = \mathcal{D}_R \nabla \cdot (\phi_R(1 - \phi_R - \phi_{PR})\nabla\tilde{\mu}_R) + \left[\exp\left(\alpha\,\tilde{\mu} + \beta\,\tilde{\mu}_R\right) - \exp\left(\tilde{\mu}_{PR}\right)\right]\,, \qquad x \in S\,, \quad (21c)$$

with the non-dimensional boundary equations

$$\boldsymbol{n}\,\mathcal{D}_P \cdot \phi(1 - \phi)\nabla\tilde{\mu} = \alpha\,\frac{1}{\ell}\frac{v_P}{v_{PR}}\left[\exp\left(\alpha\,\tilde{\mu} + \beta\,\tilde{\mu}_R\right) - \exp\left(\tilde{\mu}_{PR}\right)\right]\,, \qquad x \in S\,, \quad (21d)$$

$$\boldsymbol{n} \cdot \bar{\kappa}\nabla\phi = -\frac{\partial\tilde{J}}{\partial\phi}\,, \qquad\qquad x \in S\,, \quad (21e)$$

$$-\boldsymbol{n} \cdot (\mathcal{D}_P\,\phi(1 - \phi)\nabla\tilde{\mu}) = 0\,, \qquad\qquad x \in \partial V\,, \quad (21f)$$

$$\boldsymbol{n} \cdot \nabla\phi = 0\,, \qquad\qquad x \in \partial V\,. \quad (21g)$$

In the equations above, chemical potentials and coupling free energy J are measured in units of $k_B T$, $\tilde{\mu} \equiv \mu/(k_B T)$ and $\tilde{J} \equiv v_{PR}J/(k_B T)$. We have also introduced the non-dimensional parameters:

$$\mathcal{D}_i = \frac{D_i}{\Lambda\ell^2}\,, \qquad\qquad (21h)$$

where $i = P, PR, R$. Here, $\mathcal{D}_i$ are inverse "Damköhler numbers", which compares the time-scale of a chemical reaction, i.e., binding, with the time-scale of diffusive transport. All the parameter values and their dimensionless values in the following study are given in Table 1.

 Appendix 3

Numerical methods for the kinetic model

In our numerical simulations, the dimensionless governing equations (21) are solved in a three-dimensional domain using a second-order central finite difference method for spatial discretization and the Crank-Nicolson method for time integration. To ensure unconditional energy stability, we apply the Energy Quadratization (EQ) *Zhao et al.* (*2018a*, 2016); *Zhao and Wang* (*2019*) method, which guarantees that the system's total energy dissipates while conserving mass throughout the simulation. The computational domain size is set to $L = 1$ in each spatial direction, with $N = 128$ grid points per direction, providing sufficient spatial resolution. The time step is set to $\Delta t = 10^{-3}$, offering both stability and accuracy during time integration. This setup allows for precise tracking of the system's kinetics, with a consistent energy dissipation mechanism throughout the evolution of the system.

Appendix 4

Methods and materials

Protein purification and labeling

1-mer, 14-mer, and ZO1-GFP constructs design The last C-terminal seven amino acids
from human CLDN2 protein was used as the 1-mer receptor for recruiting ZO1 protein to
the artificial membranes. To make the 1-mer receptor easier to purify, we fused a SNAP
tag on the N terminal to get a recombinant protein SNAP-CLDN2-C7 (referred to as 1-mer).
We fused ySMF, a table 14-meric domain, to get a recombinant protein ySMF-CLDN2-C7
(referred to as 14-mer). His tags were used at the N-terminal of both receptors for affin-
ity purification. The human ZO1 sequence was fused with an N-terminal HIS-MBP tag for
affinity purification and a C-terminal mGFP tag for visualization.

Receptor protein purification For 1-mer and 14-mer receptors, we used Escherichia
coli Rosetta cells to recombinantly express proteins. Proteins were expressed in the Es-
cherichia coli Rosetta cells in LB medium at 16 °C overnight and induced with 0.2 mM IPTG
when OD600 reached 0.6 - 0.8. All the following steps were carried out at 4 °C. Cell pellets
were collected, washed, and resuspended with a lysis buffer (20 mM Hepes, pH 7.4, 500
mM NaCl, 1 mM MgCl2, 1x protease inhibitor cocktail, 1x benzonase). Cell pellets were bro-
ken with LM20 Microfluidizer, 20,000 psi, 2 runs, and clarified by centrifugation at 17,000
g, at 4 °C, for 30 min. Proteins were then purified with metal ion affinity chromatography
(Ni-NTA) resin (IMAC, 5mL HiTrap Chelating, GE Healthcare) and followed by size exclusion
chromatography with Superdex 200 increase 10/300 GL column on AKTA pure FPLC system
(GE). Proteins were collected, aliquoted, and frozen in liquid nitrogen and stored at -80°C in
20mM Hepes, pH 7.4, 500 mM NaCl, 1 mM DTT, and 5% Glycerol.

ZO1-GFP protein purification we used insect cells to recombinantly express ZO1-GFP
proteins using the baculovirus expression system. SF9-ESF S. frugiperda cells were infected
with ZO1-GFP expression baculoviruses, and then cultured at 27 °C in ESF 921 insect cell
culture medium supplemented with 2% fetal bovine serum for 3 days. Cell pellets were
then collected, washed, and resuspended with a lysis buffer (20 mM Hepes, pH 7.4, 500
mM NaCl, 1 mM MgCl2, 1 x protease inhibitor cocktail, 1x benzonase). Cell pellets were
broken with LM20 Microfluidizer, 5,000 psi, 2 runs, and clarified by centrifugation at 17,000
g, at 4 °C, for 30 min. Proteins were then purified with metal ion affinity chromatography
(Ni-NTA) resin (IMAC, 5 mL HiTrap Chelating, GE Healthcare), and followed by amylose resin
(NEB). Finally, size exclusion chromatography was performed with the Superose 6 column
on the AKTA pure FPLC system (GE). Proteins were collected, aliquoted, and frozen in liquid
nitrogen and stored at -80 °C in 20 mM Hepes, pH 7.4, 500 mM NaCl, 1 mM DTT, and 5%
Glycerol. His-MBP tag was cleaved and removed with amylose resin before being used for
respective assays.

Receptor labeling with NHS-ester dye 1-mer and 14-mer receptors were labeled with
DyLight 650 NHS Ester (Thermo Scientific, Product No.62262) dye. Highly purified proteins
were prepared in 0.05 M sodium borate buffer at pH 8.5 (Thermo Scientific BupH Borate
Buffer Packs, Product No. 28384). DyLight NHS Ester dyes were dissolved in DMSO. When
labeling the protein, the dye and protein were mixed at a molar ratio of 1:1, and incubated at
RT for 1hr. Free dye was removed from the protein by desalting columns (Thermo Scientific
Zeba Spin Desalting Columns, Product No. 89882) with buffer containing 20 mM Hepes, pH
7.4, 500 mM NaCl, 1 mM DTT, 5% Glycerol. Fluorescence labeling efficiency was measured
with Nanodrop 2000 (ThermoFisher). The labeled and unlabeled proteins were mixed to get
a final stock of 5% labeled.

In vitro reconstitution of tight junction initiation

Liposome preparation Phospholipids including POPC (Avanti, Product No. 850457), certain amounts of DGS-NTA(Ni) (Avanti, Product No. 790404), 0.1% PEG5000PE (Avanti, Product No. 880230) and 0.1% Rhodamine-PE (Avanti, Product No. 810150) were mixed in chloroform in glass bottles, and dried under vacuum for 2 hrs or overnight. 500 uL buffer (20 mM Hepes, pH 7.4, 150 mM NaCl, and 10 mM MgCl2) was added to the dried lipid film to make a final lipid concentration of 1 mM and resuspended by shaking at a speed of 180 rpm, at 37 °C for 1 hr. The lipid mixture was then transferred to the Eppendorf tube and went through freeze-thaw for 15 runs until the lipid mixture became clear. The liposomes were further clarified by centrifugation at 17,000 g for 30 min and stored at -80°C for long-term storage or 4 °C for use within one week.

Supported lipid bilayer (SLBs) preparation Glass-bottomed 96-well plates (Greiner, Product No. 655891) were pre-cleaned with 2% Hellmanex II overnight and 6 M NaOH for 30 min at RT twice. Before adding liposomes, wells were equilibrated with buffers (20 mM Hepes, pH 7.4, 150 mM NaCl, and 10 mM MgCl2) for 5 min, and left a 60 uL buffer in the well. 20 uL 1 mM liposomes were added and incubated for 20 min. 20 uL 5 M NaCl was then added for another 20 min for the liposomes to further collapse on the glass bottom. Excess liposomes were intensively washed away by pipetting in and out buffers (20 mM Hepes, pH 7.4, 150 mM NaCl, and 10 mM MgCl2) 8 to 10 times. The quality of SLBs was checked by FRAP under a confocal microscope.

Surface phase separation assay SLBs with a certain amount of DGS-NTA(Ni) lipids were blocked with 1 mg/ml BSA for 30 min. 500 nM monomeric or 14-meric receptors were added and incubated for 30 min. Excess receptors were washed away by pipetting in and out buffer (20 mM Hepes, pH 7.4, 150 mM NaCl and 10 mM MgCl2) 8 to 10 times. The images of receptors on SLBs before adding ZO1 were taken with a confocal microscope. The dynamics of receptors on SLBs before adding ZO1 were measured by FRAP. Certain amounts of ZO1-GFP were added for 30 min to form ZO1 surface condensates. 3 images for each well were randomly taken under a confocal microscope. Receptor and ZO1 densities were converted from fluorescence intensity to density numbers with calibration curves obtained with FCS measurement.

Fluorescence correlation spectroscopy (FCS) mGFP and 5% DyLight650 labeled 14-meric receptors were anchored to SLBs via His tag and DGS-NTA(Ni) interaction and excited with 488 nm and 640 nm pulsed laser (Olympus 60x NA = 1.2 water objective). Fluorescence fluctuations were recorded with a time resolution of 500 ns for 10 s. Auto-correlation of the photon traces was performed in MATLAB using a multiple tau correlator. The resulting correlation curves were fitted according to the standard 2D diffusion model including one triplet component using MTALAB (Elson, 2011). The mean particle number N was obtained from the fitting. mGFP and ZO1-mGFP were excited with a 488 nm laser in the bulk. The resulting correlation curves were fitted according to the standard 3D diffusion model including one triplet component using MTALAB (Elson, 2011). The comparison of brightness between the two molecules was calculated by the counts per molecule.

Protein density calibration on the membrane with FCS FCS was performed on SLBs functionalized with different amounts of mGFP and 14-mer-DyLight 650. Mean particle numbers were obtained from FCS fitting. Confocal images were taken for the SLBs functionalized with mGFP and 14-mer-DyLight 650 for extracting the mean fluorescence intensity. The mean fluorescence intensity of mGFP was normalized to ZO1-mGFP by the brightness comparison obtained from FCS measurement in the bulk. Calibration curves were obtained by plotting mean particle numbers as a function of fluorescence mean intensity.

Microscopy Confocal imaging and FCS measurement were performed on a commercial confocal STED microscope (Abberior Instruments, Göttingen, Germany) with pulsed laser

excitation (490 nm, 560 nm, 640 nm, 40MHz) and 60x water or 100x oil objectives (Olympus).

Quantification and statistical analysis Images were analyzed with FIJI (https://fiji.sc/) or MATLAB (Mathworks). Statistical details for each experiment can be found in the figure legends and the corresponding methods. Data shown with error bars present the standard deviation (SD). All the experiments were performed more than three times and quantified independently.

Estimation of model parameter for ZO proteins binding to claudin receptors

We analyse the microscope images and compute the density of the components in the membrane and bulk. For this purpose, we define a mask to find the dense and dilute phases, respectively. Specifically, we use a Gaussian filter with a two-dimensional Gaussian smoothing kernel with standard deviation of 0.8 to smooth the raw receptor intensity data. Then, we set the area with $I_{\mathrm{receptor}} \geq 60$ for dense phase(red curve in Appendix 5-fig. 1a), while $I_{\mathrm{receptor}} \leq 10$ for dilute phase(white curve in Appendix 5-fig. 1a). The area in between is the interface area (see Appendix 5-fig. 1). Once we define the dense and dilute phases in the images, we can quantify the density in the dense and dilute phases correspondingly.

Appendix 5—figure 1. (a) Illustration of the mask for receptor intensity to determine the surface concentrations inside and outside of the surface condensates in the membrane. The blue area represents the dilute phase and the yellow area the dense phase. Mean absolute error (MAE) color maps with respect to interaction parameters $\chi_{PR\text{-}L}$ and internal free energy ω_{PR} for 1-mer receptor and 14-mer receptor are shown in (b) and (c), respectively. For the considered parameter range, MAE is minimized for 14-mer at $\chi_{PR\text{-}L} = 1.09$ and $\omega_{PR} = -1.37$, and for 1-mer at $\chi = -0.4$ and $\omega_{PR} = 1.1$. We note that though there is a minimum of the MSE corresponding to at $\chi_{PR\text{-}L} = -0.4$, the color map in figure (c) suggests that the $\chi_{PR\text{-}L}$ value for the 1-mer cannot be determined as there is a "valley" of similar MAE value along $\chi_{PR\text{-}L}$.

According to the experimental studies, membrane phase separation occurs for bulk protein concentrations of a few tens of nM. For concentrations in the order of 10 nM, there is no phase separation in the bulk since the bulk saturation concentration is about $10\,\mu M$. Thus, we simplify our model by adapting it in an approximate fashion to these experimental conditions by setting bulk interaction parameter $\chi = 0$ and use, for simplicity, equivalent solvent and solute molecule size, i.e., $n = 1$. Furthermore, we choose the coupling free energy equals to zero, implying that the profile along z, $\phi(z)$ is constant at thermodynamic equilibrium. Note that during the time evolution, the bulk volume fraction described by Eq. (6a) can be heterogeneous. At thermodynamic equilibrium, however, the bulk can be thought as a finite homogeneous reservoir for proteins.

We roughly estimated the size of 14-mer receptor and 1-mer receptor from the structure in Pymol based on their structures *Collins et al. (2003)*; *Wilhelm et al. (2021)* as: $d_{14\text{-mer Receptor}} = 9.2\,nm$, $d_{1\text{-mer Receptor}} = 3.06\,nm$,. The corresponding molecule diameter of the ZO1 protein is achieved from AFM, $d_{ZO1} = 9.2\,nm$, We choose the size of the lipid patch in the membrane as $d_{\mathrm{lipid}} = 3.06\,nm$. This is an effective free lipid patch to avoid considering molecules of very different sizes as we mentioned before. We note that the molecule area ratios for 14-mer receptor are $n_{PR} = n_R = 9$ with lipid patch as a reference component in the membrane, while the ones for 1-mer are $n_{PR} = n_R = 1$.

With the given molecule sizes, we determine that the ZO1 bulk volume fraction is exceedingly small (approximately 10^{-5}) when the ZO1 number density is $50\,nM$ in the bulk. From a numerical standpoint, such a low volume fraction requires a very fine mesh size in both time and space to accurately sample the curvature of the free energy density. This significant curvature originates from the entropic term, and mathematically, from the logarithmic dependence in the free energy. To circumvent the need for extremely fine numerical grids and thus conserve computational resources, we scale the bulk volume fraction ϕ^{raw} with a scaling factor such that the scaled volume fraction ϕ equals 0.1 when the ZO1 number density is $50\,nM$ in the bulk, i.e. $\phi = C_0 \phi^{\text{raw}}$, where $C_0 = 8146$.

 Appendix 6

Model fitting to experimental data of binding ZO1 proteins

Based on the previous discussions, our model has two remaining unknown parameters: the interaction parameter between the receptor-protein complex and free lipid, $\chi_{PR\text{-}L}$, and the internal free energy difference between the bound and unbound proteins, ω_{PR}. To determine these parameters, we fit our thermodynamic model (Eq. (13)) to experimental data.

We employ two sets of experimental data: one set varies the bulk ZO1 concentration, and the other varies the receptor amount in the membrane. The optimal values of the interaction parameter $\chi_{PR\text{-}L}$ and the internal free energy coefficient ω_{PR} are obtained by minimizing the mean absolute error (MAE), defined as:

$$MAE = \frac{1}{n} \sum_{i=1}^{n} |\phi_{PR}^{\text{data}} - \phi_{PR}^{\text{simu}}|, \tag{22}$$

where n represents the total number of data points in the experiments. Here, ϕ_{PR}^{data} denotes the area fraction of the receptor-ZO1 complex, and ϕ_{PR}^{simu} is the simulated area fraction solved from the equilibrium condition (Eq. (13)).

Strikingly, we found very good agreement between the model and the experimental measurements(see Fig. 5c and 5d and Fig. 6c and 6d). In particular, the model recapitulated the experimental ZO1 and receptor (14-mer) concentrations at which surface condensates appear as well as the concentrations in the condensed and dilute phases(see Fig. 6c and 6d). Consistent with the experimental data, the model predicted a jump in the difference in ZO1 concentrations between condensates and their surrounding dilute phase. The binding affinity obtained in the fitting is $\omega_{PR} = -1.09$ for 1-mer and $\omega_{PR} = 3.83$ for 14-mer; the interaction strength parameter value is $\chi_{PR\text{-}L} = -0.4$ for 1-mer and $\chi_{PR\text{-}L} = 1.09$ for 14-mer.

Binding fraction

In the experiments, we observed that maximal fraction of bound receptors is different for different oligomerization states of the receptor, i.e., 1-mer and 14-mer. Specifically, while all 14-mer receptors can be bound by proteins from the bulk, only a fraction of 1-mer receptors are available for binding. This phenomenon was already reported in Ref. ***Erlendsson et al.*** (***2019***) and tracked back to a molecular switch mechanism.

We quantify the binding fraction c as the ratio between total membrane-bound protein area fraction and total free receptor area fraction at saturated state, i.e. $c = \bar{\phi}_m/\phi_R^0$. Based on the experimental data, we obtain that $c = 1$ for the 14-mer receptors and $c = 0.014$ for 1-mer receptors. In our model, we include the reduced binding fraction mechanism in the following way:

We calculate the membrane-bound protein area fraction ϕ_{PR} from the thermodynamic conditions ((13)). Then, (1) If $\phi_{PR} \leq c\phi_R^0$, a finite binding fraction c has no effects. (2) If $\phi_{PR} > c\phi_R^0$, we use

$$\phi_{PR} = c\,\phi_R^0,$$
$$\phi_R = \phi_R^0 - c\,\phi_R^0,$$

where ϕ_R^0 is the average area fraction of the free receptors in the membrane initially.

Kinetic properties of the surface phase separation

Appendix 8—figure 1. (a) Time-resolved snapshots illustrating the emergence and growth of ZO1-rich condensates on a model membrane surface (top row: theoretical simulation; bottom row: experimental observation). Initially uniform at $t = 0$, the membrane surface progressively develops well-defined droplet patterns by 15 minutes. The system includes 100 nM ZO1 in the bulk and 800 μm^{-2} 14-mer Receptor in the membrane. The scale bar in the experimental images represents 2 μm. (b) Temporal evolution of the total membrane-bound ZO1 fraction, comparing the theoretical prediction (blue line) and experimental measurement (black line). Both approaches show an increase over time, and they closely agree on the final coverage levels. (c) Average droplet area as a function of time, demonstrating the consistency between theory (blue line) and experiment (black line) in capturing the size growth and stabilization of condensates. (d) Time-dependent count of individual droplets, showing that both the model (blue line) and the experimental data (black line) reproduce the overall trend in droplet formation and coalescence. we set the diffusivity coefficient value as $D_i = 0.01\ \mu m^2 s^{-1}$, $i = P, PR, R$. The gradient coefficients $\kappa = \kappa_{PR} = \kappa_R = 2 \times 10^{-5} \mu m^{-1}$.

We compare the simulated formation and evolution of ZO1-rich regions on the membrane (top row) to those observed experimentally (bottom row) (see Fig. 1). At the start, both simulation and experiment show a relatively uniform distribution of receptors and ZO1. Over time, small clusters begin to appear and grow more distinct, and by 15 minutes both systems feature well-defined, separated condensates.

The overall pattern of how these clusters develop is quite similar in both simulation and experiment. We observe that surface condensates rapidly form once the critical concentrations of receptor-ZO1 complex are exceeded. However, after this initial rapid phase, the system transitions to much slower coarsening kinetics (Appendix 8-fig. 1, movie S1). This slow-down in coarsening kinetics is consistent with the experimental data and can be explained theoretically by two possible mechanisms. First, the depletion of receptors in the dilute phase due to co-condensation of receptors and scaffold proteins likely reduces the ripening fluxes between surface condensates (Appendix 8-fig. 1a). This reduction in flux slows down the further growth and coalescence of condensates, effectively stalling the kinetic process. Alternatively, the slow-down may be attributed to an ageing process *Jawerth et al.* (*2020*) of the surface condensates and could be explained by the limited movement

of surface condensates on the membrane with the gain of mass *Snead et al.* (*2022*).

These results illustrate that the theoretical framework, incorporating appropriate diffusivity and interaction parameters, closely reproduces not only the qualitative spatial patterns observed experimentally but also key quantitative descriptors of ZO1 droplet dynamics, including membrane coverage(see Fig. 7b, Fig. 7c and Appendix 8-fig. 1a), droplet size(see Appendix 8-fig. 1c), and droplet number(see Appendix 8-fig. 1d). However, the timing of the initial binding and recruitment does not fully align with the experimental observations shown in Appendix 8-fig. 1(b). This discrepancy may also stem from differences between the model's assumed initial conditions and those in the experimental setup, as well as the limited spatiotemporal resolution of the experimental measurements. These factors lie beyond the scope of this manuscript. Nevertheless, the strong overall agreement indicates that our model provides a valuable representation of the key mechanisms behind ZO1 surface condensate formation.