Warming increases richness and shapes assemblages of eukaryotic parasitic plankton

Amruta Rajarajan*[1], Sławek Cerbin[2], Kingsly C. Beng[1, 3§, 4§, 5§], Michael T. Monaghan[1, 6], Justyna Wolinska[1, 6]

[1]Dept. of Evolutionary and Integrative Ecology, Leibniz Institute of Freshwater Ecology and Inland Fisheries (IGB), Müggelseedamm 301, 12587 Berlin, Germany

[2]Dept. of General Zoology, Faculty of Biology, Adam Mickiewicz University, Poznań, Poland

[3]Focus Group Marine Molecular Ecology, Helmholtz Institute for Functional Marine Biodiversity, University of Oldenburg (HIFMB), Oldenburg Germany

[4]Alfred Wegener Institute (AWI), Helmholtz Center for Polar and Marine Research, Bremerhaven, Germany

[5]Thünen Institute of Sea Fisheries, Bremerhaven, Germany

[6]Institut für Biologie, Freie Universität Berlin, Germany

*Corresponding author email: amruta.rajarajan@igb-berlin.de, amrutarajarajan@gmail.com

§Present address

Abstract

Background

Anthropogenic activities have led to a global rise in water temperatures, prompting increased

interest in how warming affects infectious disease ecology. While most studies have focused

on individual host-parasite systems, there is a gap in understanding the impact of warming

on multi-host, multi-parasite assemblages in natural ecosystems. To address this gap, we

investigated freshwater eukaryotic parasite communities in ten natural lakes near Konin,

Poland: five artificially heated and five non-heated "control" lakes. Since 1958, the heated

lakes have experienced a mean annual temperature increase of 2 °C due to hot water

discharge from two adjacent power plants. We collected seasonal environmental DNA

(eDNA) samples from surface waters over a two-year period and applied targeted

metabarcoding to compare the richness and distribution of eukaryotic parasites across lake

types, with a focus on protists and fungi.

Results

Using literature searches and sequence metadata from GenBank, we identified putative

parasites which included Alveolates, Stramenopiles, basal Fungi and Ichthyosporeans, as

well as their associated hosts. Heated lakes harboured distinct parasite assemblages, with

higher richness of chytrids and aphelids, suggesting thermal preferences among certain

freshwater microeukaryotic parasites. Other groups exhibited clear seasonal trends, with

richness of oomycetes peaking in spring and summer, and that of Cryptomycota in winter

and autumn. A general linear model revealed a marginally positive correlation between

chytrid parasite richness and richness of their green algal, diatom, and dinoflagellate hosts.

Post-hoc analyses indicated that heated lakes exhibited greater seasonal variation in chytrid

parasite richness and a stronger correlation between host and parasite richness than control

lakes.

Conclusion

These findings demonstrate that warming can induce strong shifts in the richness and

53 assemblages of freshwater microeukaryotic parasites. Using chytrids as a focal group, we

54 additionally demonstrate that warming may amplify seasonal variation in parasite richness

55 and strengthen host-parasite richness relationships.

56 Introduction

Warming in aquatic environments has led to an average increase in freshwater temperatures

of 0.34 °C per decade (O'Reilly et al., 2015). Elevated temperatures can drive deoxygenation

(Jane et al., 2021), changes in seasonality and reduced ice cover (Grant et al., 2021), shifts

in lake productivity dynamics (Gilarranz et al., 2022), and alterations in mixing regimes

(Woolway & Merchant, 2019). These changes have cascading effects on species

distributions and food webs (Bartley et al., 2019). The magnitude of warming-driven changes

on aquatic life is unprecedented, highlighting a critical need to understand their broader

ecological implications (IPCC 2023).

While the effects of warming on plankton have been widely studied due to their central role in

nutrient cycling (Bakhtiyar et al., 2020; Naselli-Flores & Padisák, 2023), the parasites of

plankton remain relatively understudied despite their significant ecological contributions. For

instance, chytrid fungi (Chytridiomycota) infect various phytoplankton, including green algae

(reviewed in Frenken et al., 2017), diatoms (reviewed in Danz & Quandt, 2023), and

dinoflagellates (Fernández-Valero et al., 2023, 2024). By producing motile zoospores,

chytrids create additional trophic links (mycoloops) that sustain zooplankton populations

(Abonyi et al., 2024; Kagami et al., 2014). Chytrid outbreaks can exert top-down control on

toxic cyanobacterial (Gleason et al., 2015) and dinoflagellate blooms (Lepelletier et al.,

2014). Other parasitic groups, such as Cryptomycota, hyperparasitize chytrids (James et al.,

2013; Powell & Letcher, 2019), while Aphelidiomycota (aphelids) (Jephcott et al., 2017) and

Oomycota (oomycetes) (Thines, 2018) infect various phytoplankton hosts. Perkinsea, a

group previously considered to exclusively contain intracellular parasites of green algae, was

recently discovered to be more diverse, with potential roles in ecosystem processes beyond

parasitism (Metz et al., 2023). Other highly diverse parasite groups include Apicomplexans

(Gad et al., 2023) and Cercozoans (Lefèvre et al., 2008), although their contribution to

aquatic ecosystem processes is less understood.

The influence of warming on infectious disease extends beyond parasite development to

affect host abundance, physiology, and immunity (King et al., 2023). For instance, a four-

month indoor mesocosm study found that 4 °C elevation in spring temperature (to 21 °C)

shortened the bloom duration of the diatom *Synedra* sp., resulting in an earlier end to the

epidemic of its chytrid parasite *Zygorhizidium planktonicum* (Frenken et al., 2016).

Conversely, an outdoor mesocosm study of amphipod host communities exposed to the

generalist trematode parasite *Maritrema novaezealandensis* under a 4 °C summer

temperature increase (to 25 °C) showed elevated infection prevalence, along with reduced

host species richness and abundance (Mouritsen et al., 2018). These findings demonstrate

how warming can impact disease dynamics in individual host-parasite systems within aquatic

environments.

Parasite species richness is integral to ecosystem health (Hudson et al., 2006) is primarily

influenced by the host distribution and richness of their hosts in distinct ways. Chytrid

parasite distributions are closely linked to the spatial co-occurrence of their green algal and

diatom hosts (Ilicic et al., 2024). More broadly, parasite richness also correlates positively

with host richness across biomes, according the host-diversity-begets-parasite-diversity

hypothesis (Kamiya et al., 2014; Krasnov & Poulin, 2015). For example, ponds with higher

amphibian host richness harbored higher richness of trematode parasites (Johnson et al.,

2016). Changes in host and parasite species richness can, in turn, affect disease dynamics

in complex ways. In one case, elevated amphibian host richness reduced infection

prevalence of the trematode *Ribeiroia ondatrae* (dilution effect) but amplified parasite

competition (Johnson et al., 2013).

Despite these insights, our understanding of how global warming affects parasite richness

remains incomplete (Sures et al., 2023). Warming-driven changes in species assemblages

can lead to 'ecological fitting' due to novel host-parasite encounters and the resulting

increased likelihood of host-switching (Wells & Clark, 2019). This could potentially increase

phenotypic and genotypic diversity of parasites (Agosta et al., 2010). Moreover,

anthropogenic influence can have unexpected impacts on the generally positive correlation

between host and parasite richness in natural ecosystems. For instance, islands that

experienced extensive fishing exhibited higher fish host richness, but lower helminth parasite

richness, compared to unfished islands (Wood et al., 2018). In contrast, warming-driven

habitat loss of host species reduced helminth parasite richness (Carlson et al., 2017). These

inconsistencies highlight the need for long-term studies of multi-host, multi-parasite systems

to better understand the role of warming in shaping disease ecology.

One promising approach to study the effects of warming on parasite assemblages involves *in*

situ sampling of natural lakes. The Konin lake complex in Poland offers an ideal opportunity.

Since 1958, five of these lakes have been artificially warmed by hot water discharge from two

nearby power plants, resulting in altered abiotic conditions and changes in plankton

community composition compared to the five nearby non-heated (hereafter "control") lakes

(Zdanowski et al., 2020). The heated lakes are on average 2 °C warmer (Beng et al., 2023),

mimicking the projected global freshwater temperature increases by 2100 (IPCC 2023).

Additionally, these lakes experience reduced winter ice cover (Socha & Zdanowski, 2001).

Warming has led to shifts in *Daphnia* species composition in these lakes, favoring the more

heat-tolerant *D. galeata* (Dziuba et al., 2020). Moreover, the heated lakes exhibit higher

richness and distinct community composition of various phytoplankton (green algae, protists,

fungi) and zooplankton (crustaceans, rotifers) taxa (Beng et al., 2023; Ejsmont-Karabin et al.,

2020). There is also an increased abundance of invasive species, including the tape grass

Vallisneria spiralis (Socha & Hutorowicz, 2009) and the mussels *Corbicula fluminea*

(Urbańska et al., 2018) and *Sinanodonta woodiana* (Urbańska et al., 2019). *Sinanodonta*

woodiana occurrence further favors the presence of its parasitic worm *Aspidogaster*

conchicola (Yuryshynets & Krasutska, 2009). Heated lakes also harbor introduced species of

fish (Kapusta & Bogacka-Kapusta, 2015). These changes in environmental conditions and

host community composition provide a valuable opportunity to investigate the effects of

warming on parasite diversity in natural settings. Although warming reduces *Caullerya*

mesnili prevalence in *Daphnia* hosts (Dziuba et al., 2023), comprehensive investigations into

differences in parasite assemblages between heated and control lakes are still lacking.

This study leverages this unique set of artificially heated and control lakes in Poland to

explore ca. 60 years of warming effects on eukaryotic parasitic plankton. Using eDNA

metabarcoding of seasonally collected surface water samples (winter, spring, summer,

autumn) over two years from five heated and five control lakes to address the following

questions: (i) Which parasites are present in these lakes? (ii) Do certain parasites exhibit a

preference for habitat types (heated or non-heated), leading to distinct assemblages in

heated versus control lakes? (iii) Does parasite richness differ between heated and control

lakes? and (iv) Does parasite richness correlate with host richness?

Material and methods

Study sites and eDNA sampling

Information on study sites and the timing and locations of eDNA sampling is adapted from

Beng et al., (2023). This study includes five thermally altered ("heated") lakes (Gosławskie,

Licheńskie, Mikorzyńskie, Pątnowskie and Ślesińskie) and five control lakes (Budzisławskie,

Gopło, Skulskie, Skulska Wieś and Suszewskie). Each lake was sampled once per season

throughout 2020 and 2021, with the exception that no summer samples were obtained in

2021 and all lakes were sampled twice in autumn 2020 (Fig. S1, Table S1). All lakes, except

for the control lakes Suszewskie and Budzisławskie, are connected by water channels,

potentially enabling dispersal, migration and colonization (Dziuba et al., 2017; Hillbricht-

Ilkowska & Zdanowski, 1988).

Water from the lakes Gosławskie, Licheńskie, Mikorzyńskie, Pątnowskie and Ślesińskie has

been used to cool lignite-combusting power plants situated in Konin (since 1958) and Pątnów

(since 1970), with the heated effluent subsequently discharged back into the lakes. This

discharge occurs year-round in all heated lakes except Ślesińskie, which receives thermal

effluent only from May to September. On average, heated lakes exhibit water temperatures 2

°C higher than control lakes. Beyond temperature, these lakes also show elevated

concentrations of total and soluble reactive phosphorus, particularly in the autumn (Beng et

al., 2023). While used for recreation and fish-farming, regular monitoring confirms that hot

water discharge does not cause chemical pollution. All study lakes are subject to some level

of anthropogenic disturbance, including agricultural runoff, aquaculture and urbanization.

To obtain eDNA samples, 2 L of surface water was collected at each sampling event using a

clean HDPE bottle. Water was filtered through a glass fiber mesh (Whatman GF/F, 25 mm

diameter, 0.7 µm pore size) using vacuum filtration at 200 mbar. Filters were first frozen at -

80 °C, freeze-dried at -45 °C for 8 hours and finally stored at -20 °C until DNA extraction.

171 Generation of sequence data

172 Molecular laboratory procedures and sequencing

Molecular laboratory procedures are reproduced from Beng et al., (2023). Filters containing

eDNA were pulverized by adding one stainless steel bead (5 mm diameter, Qiagen GmbH,

Hilden, Germany) and tissueLyser (Qiagen), and shaking three times for 90 seconds at 30

Hz in a mixer mill MM301 (Retsch GmbH, Haan, Germany). The tubes were briefly

centrifuged to collect the material into a pellet. DNA was extracted using the NucleoSpin

Plant II extraction kit (Sinhahery-Nagel, Düren, Germany), following the manufacturer's

protocol, and stored in TE buffer at -20 °C until further processing.

For DNA amplification, two primer sets were used: "EUK15" TAReuk454FWD1: 5'-

CCAGCASCYGCGGTAATTCC-3'; TAReukREV3: 5'-ACTTTCGTTCTTGATYRA-3'; both

primers from Stoeck et al., (2010), targeting a fragment of the 18S rRNA gene of eukaryotes,

and "CHY" ITS4ngsF: 5'-GCATATCAATAAGCGSAGGA-3' (Van den Wyngaert et al., 2022);

LF402R: 5'-TTCMCTTTNMRCAATTTCAC-3' (Thomé et al., 2024), targeting a fragment of

the 28S rRNA gene of fungi. For the "EUK15" primer set, 10 ng template DNA was used in

25-µL PCR reactions with the following conditions: an initial denaturation at 95 °C for 30

seconds, followed by 30 cycles (95 °C for 30 s, 45 °C at 30 s, 72 °C at 30 s), with a final

extension of 72 °C for 5 minutes. PCR products were normalized to 5 ng/µL and used as

templates for the second PCR, where Illumina Nextera indexes were added to 25-µL PCRs

under these conditions: 95 °C for 2 minutes, followed by 8 cycles (95 °C for 20 s, 52 °C for

30 s, 72 °C for 30 s), and a final extension at 72 °C for 3 minutes. For the "CHY" primer set,

10 ng template DNA was used in 25-µL PCRs with the following conditions: an initial

denaturation at 96 °C for 30 seconds, followed by 20 cycles (90 °C for 30 s, 50 °C for 30 s,

72 °C for 60 s), with a final extension of 72 °C for 3 minutes. 1 µL of PCR product was used

as template to repeat the enrichment PCR using the same conditions except that the final

extension at 72 °C was for 5 minutes. PCR products were purified using a magnetic bead

protocol (Agencourt AMPure XP, Beckman Coulter, Indianapolis, IN, USA), following the

manufacturer's instructions. DNA was quantified using a Quantus fluorometer and

QuantiFluor dsDNA system (Promega, Madison, WI, USA) and PCR products were

normalized to a concentration of 5 ng/µL for use in indexing. Indexing PCR was performed

with the following cycling conditions: initial denaturation at 95 °C for 2 min, followed by 8

cycles (95 °C for 20 s, 52 °C for 30 s, 72 °C for 30 s), with a final extension at 72 °C for 3

min. Barcoded samples were pooled in equimolar concentrations and paired-end sequenced

on the Illumina MiSeq (600 cycles, v3 chemistry) platform at the Berlin Center for Genomics

in Biodiversity Research (BeGenDiv). Each sequencing run included one no-template

negative control.

Bioinformatics

Each library (EUK15 and CHY) was sequenced in two runs: the first run included samples

from winter 2020, spring 2020, summer 2020 and autumn 2020, while the second run

included samples from winter 2021, spring 2021, autumn 2021 and an additional set of

samples taken in autumn 2020. Bioinformatic processing was performed as described in

Beng et al., (2023). The first sequencing run yielded 2.4 million raw reads for EUK15 and 1.9

million raw reads for CHY (Beng et al., 2023), while the second run yielded 643,742 raw

reads for EUK15 and 741,018 raw reads for CHY. Primer sequences were removed with

cutadapt (Martin, 2011) and reads were processed using the package DADA2 v 1.28

(Callahan et al., 2016) in R v 4.3.0 (R Core Team, 2023). Forward and reverse reads were

respectively: filtered with maximum expected errors set to 2.6 and 2, then truncated at 260

bp and 200 bp respectively before merging (see separate supplementary R scripts). Quality

filtering and *de novo* chimera removal retained 59% of EUK15 and 58.3% of CHY reads.

Reads from the two runs were combined into a "protist dataset" (EUK15) and a "fungal

dataset" (CHY). Amplicon Sequence Variants (ASVs) were identified for each dataset using

the *dada* function, and ASV-count tables were prepared. Taxonomic assignment of ASVs in

the protist dataset was conducted using the Protist Ribosomal database PR2 v.5.0.0 (Guillou

et al., 2013). For the fungal dataset, taxonomic assignment was performed using BLAST+

with a custom database (see below).

Despite the lower sequencing depth in the second runs, all rarefaction curves plateaued (Fig.

S2), confirming sufficient sequencing depth. Thus, all samples were retained for further

analyses without rarefaction. Contaminant ASVs, defined as those present in negative

controls, were removed using the phyloseq package in R (McMurdie & Holmes, 2013).

Specifically, 17 contaminant ASVs were removed from the first run of the protist dataset,

while 3 and 2 contaminant ASVs were removed from the first and second runs of the fungal

dataset, respectively, comprising <0.01% reads in each dataset. Bioinformatic processing

resulted in 3297 ASVs in the protist dataset and 4243 ASVs in the fungal dataset.

Classification of ASVs as parasites

ASVs were classified as parasites using a modified approach from Beng et al. (2021)

(detailed in Fig S3). Two custom parasite sequence databases were created for this purpose:

one containing sequences from the NCBI nucleotide database representing 10 higher taxa

discovered in the protist dataset (Fig. S4), and the other containing NCBI sequences of 9

taxa discovered in the fungal dataset (Fig. S5). Additionally, we incorporated (i) ribosomal

sequences from NCBI (18S, 28S, and ITS), (ii) sequences from parasite groups reported to

occur in the Konin lake complex (Cestoda, Nematoda, Monogenea, Digenea, Trematoda and

Acanthocephala), which infect fish (Pojmańska et al., 2012), mussels (Yuryshynets &

Krasutska, 2009), and snails (Cichy et al., 2020), and (iii) sequences of eukaryotic taxonomic

groups known to contain parasite species, identified through Web of Science literature

searches performed for each unique phylum listed in the Fungal Names database (F. Wang

et al., 2023), using the terms "[Phylum name] AND [parasit* OR pathogen*]" (Table S2).

Duplicate sequences were removed using seqkit (Shen et al., 2016). Each ASV was then

identified to species- or higher taxonomic level using its respective custom database and

BLAST+ on a high-performance computing cluster (Bennett et al., 2020). Only the top ten

hits with ≥90% sequence similarity and 100% query cover were processed further. For each

ASV, "host" entries within GenBank metadata of the top-bitscore BLAST hit (Beng et al.,

2021) were extracted using the rentrez package (Winter, 2017).

Species names from the protist and fungal datasets (for ASVs that were assigned to species

level) were used for literature searches on Web of Science with the search terms "[Species

name] AND [parasit* OR pathogen* OR infect*]". Relevant articles were checked for

evidence of parasitism and host association. BLAST hits of ASVs with no associated host

entries in GenBank or host information from literature searches were classified as non-

parasitic (Fig. S3).

Host entries obtained from GenBank were manually categorized into host groups (see Fig.

1). Only ASVs associated with live hosts were classified as parasites; those associated with

abiotic environmental samples (e.g., water, soil, mud sediment) or non-living substrates (e.g.,

"decaying log", "fallen branch") were excluded. ASVs representing symbionts (e.g., algal

endosymbionts of protists, lichen-associated ASVs, dinoflagellate symbionts of sponges) or

organisms within the pathobiome of diseased hosts were also excluded. This rigorous

classification ensured that only ASVs with confirmed parasitic lifestyles were included in the

final analysis, enabling an accurate assessment of parasite richness and distribution across

the studied lakes (Fig. S3).

Statistical analyses

Individual lakes of each type (heated, control) were considered replicates in all statistical analyses, but included independent sampling events over two consecutive years (Table S1). Sample sizes differed for the seasons in our analyses: winter (n = 10), spring (n = 10), summer (n = 5) and autumn (n = 15).

ASVs indicative of heated or control lakes

To assess whether certain parasite ASVs were indicative of either heated or control lakes, we used the *signassoc* function (9999 permutations) from the indicspecies package in R. This analysis was performed using presence-absence data for parasite ASVs (De Caceres & Legendre, 2009) (mode=0). Multiple testing correction was implemented using the Sidak method within the indicspecies package.

ASV richness differences in dominant parasite groups between heated and control lakes

Parasite groups were scrutinized to identify those associated with aquatic hosts (see Fig. 1). The four parasite groups with the highest ASV count were selected for comparisons across lake types and seasons. For example, in the protist dataset, 66 ASVs were identified as Gyrista (Stramenopiles phylum), but only 31 Gyrista ASVs belonged to Peronosporomycetes, after excluding terrestrial hosts. Consequently, only Peronosporomycetes (=Oomycetes, Subdivision: Gyrista), representing 7.9% of ASVs in the protist parasite dataset, were included in ASV richness analyses. Additionally, 216 ASVs in the protist dataset were identified as fungi but only fungi from the fungal dataset were considered for further analyses. In the fungal dataset, 297 and 86 ASVs belonged to Ascomycota and Basidiomycota, respectively, but 86.8% and 87.2% of these ASVs were associated with terrestrial hosts (such as metazoans, plants and humans; see Fig. 1B) and were therefore excluded. Chytrids (Phylum: Chytridiomycota), aphelids (Phylum: Aphelidiomycota) and Cryptomycota infecting aquatic hosts were analyzed, collectively representing 50.3% of ASVs in the fungal parasite dataset.

The ASV richness for each selected group was log-transformed and analyzed using a

Generalized Linear Mixed Model (GLMM). The most parsimonious model with the best fit

was used to examine parasite richness across lake types, seasons and years: parasite ASV

richness ~ lake type * season + (1 | lake) + (1 | sequencing run) + (1 | year), family = poisson,

using the *glmer* function of the lme4 package (Bates et al., 2024) in R. The significance of

fixed effects was assessed with a Type III ANOVA using the *Anova* function of the car

package (Fox et al., 2019). For chytrid parasite richness, a separate model was employed to

investigate its correlation with host richness (see below).

<u>Correlation of chytrid parasite richness and host richness</u>

A second GLMM was used to test whether differences in chytrid parasite richness correlated

with differences in host richness across heated and control lakes (Beng et al., 2023). Here,

"chytrid parasite richness" was defined as the count of chytrid parasite ASVs in the fungal

dataset, while "host richness" referred to the count of ASVs from the Subdivisions

Chlorophyta, Charophyta, Chromista, Gyrista, and Dinoflagellata in the protist dataset. These

host groups were selected because 98% of chytrid parasite ASVs were associated with one

or more of these groups (see Fig. 1B). 2% chytrid parasite ASVs were excluded from this

analysis because they associated with salamander hosts, and the richness of salamander

hosts was not surveyed (see Table S3). Both host and parasite richness values were log-

transformed. We then used the following GLMM structure: chytrid parasite richness ~ lake

type * season + host ASV richness + (1 | lake) + (1 | sequencing run) + (1 | year), family =

poisson. Fixed effects were evaluated with a Type III ANOVA. After observing significant

effects of host richness, lake type, season and lake type × season interaction on parasite

richness (see Results), we calculated confidence intervals and visualized a general linear

model for host and parasite richness, separately for each lake type (see Fig. 4). The model

was plotted using the "geom_smooth" argument (method="glm", se=TRUE) from the ggplot2

package (Wickham, 2016) in R.

After discovering the significant effect of lake type, host richness and the lake type × season

interaction on chytrid parasite richness (see Table 1), we computed partial R^2 values to

assess the effect sizes of host richness and season on parasite richness. This was done

separately for heated and control lakes. Random effects were excluded from this test

because mixed-effects structures resulted in a singular fit. Instead, GLMs were specified for

each lake type with the structure: parasite richness ~ host richness + season. Partial R^2

values for each fixed effect were computed using the rsq package in R (Zhang, 2024).

Results

Characterization of parasite assemblages and hosts in the protist and fungal datasets

In the protist dataset, 390 ASVs (12% of the dataset) were classified as parasites. These

included: 43 ASVs confirmed as parasitic based on both literature evidence and GenBank

host entries, 55 ASVs classified as parasites using literature evidence alone, and 292 ASVs

classified based solely on GenBank host entries (Table S2). A total of 151 parasite ASVs

(39%) were identified to the species level, representing 54 species from 23 genera (Table

S3), which were further grouped into 14 higher taxa (Fig. 1A). The highest ASV richness was

observed among Fungi, Gyrista, and Ichthyosporea (Fig 1A). Most parasite ASVs (66%)

were associated with hosts such as green algal, diatom, or terrestrial-plant hosts (Fig. 1A,

Table S3).

In the fungal dataset, 683 ASVs (16% of the dataset) were identified as parasites: 198 ASVs

were classified as parasitic using both literature evidence and GenBank host entries, 298

ASVs based on GenBank host entries alone, and 187 ASVs using literature searches of

species names. A total of 496 ASVs (73% of parasite ASVs) showed >90% sequence

similarity and 100% query cover with known species in GenBank (Table S4). Based on these

criteria, the fungal dataset contained 268 putative species across 218 genera, classified into

15 higher taxa. The highest ASV richness was observed among Ascomycota,

Chytridiomycota and Cryptomycota (Fig. 1B). Similar to the protist dataset, a notable

proportion of parasite ASVs (45%) in the fungal dataset were linked to hosts such as green

algae, diatoms, or terrestrial plants (Fig. 1B, Table S4).

Figure 1. Parasite ASV count in the (A) protist and (B) fungal datasets. Colors represent broad host

taxa to which identified hosts of the parasite ASVs belong (see Methods and Fig. S3 for details). Note

that Fungal ASVs from the protist dataset were not considered for statistical analyses.

ASVs indicative of lake type

A total of 39 parasite ASVs were indicators of lake type: 5 ASVs were specific to control

lakes, and 34 ASVs were specific to heated lakes (Fig. 2). Among these were 19 chytrids, 8

aphelids, 4 Cryptomycota, 7 protists from diverse lineages, and 1 metazoan (Table S5).

Control lakes indicators included chytrids ASVs (263 and 34) from unclassified members of

the Order Rhizophydiales. In contrast, heated-lake indicators comprised various taxa,

including chytrid species *Dinomyces arenysensis*, *Dangeardia mamillata*, *Rhizophydium*

chlorogonii, *Quaeritorhiza haematococci* and an unknown Rhizophydiales parasite (ASV

210). Other heated-lake indicators were *Elliptio complanata* (Unionidae mussel), *Ustilago*

esculenta (Basidiomycete fungus) along with unclassified ASVs from Dinophyceae (ASV

110) and Peronosporales fungi (ASV 286). Additionally, *Aphelidium desmodesmi* and four

unclassified ASVs from the Family Aphelidiaceae (ASVs 60, 220, 194 and 101) showed a

preference for heated lakes.

Figure 2. Relative abundances of ASVs indicative of control (left panel) and heated (right panel) lakes. Each square represents the relative abundance of

indicator ASVs (rows) in individual lakes (columns), grouped by the season of sampling indicated on the x-axis. Sampling occurred over two consecutive years

(See Methods). Of the 39 indicator ASVs identified (Table S5), 29 are presented here, as they were present in more than 20% of samples. ASV numbers are

specified in brackets. For ASVs in the protist dataset (marked with an asterisk), the lowest taxonomic classification assigned using the PR2 database is reported,

with values in the blue "%" column indicating sequence similarity to the BLAST hit used to obtain host information (host entry from GenBank metadata). For ASVs

from the fungal dataset, the lowest known classification from the BLAST hit is reported, with the blue "%" values indicating sequence similarity between the ASV

and the BLAST hit. The classification of *E. complanata* as a parasite was based on a literature search (Table S5).

Parasite ASV richness

Among the four parasite groups with the highest ASV richness - chytrids, aphelids, Peronosporomycetes (Oomycetes), and Cryptomycota - chytrids and aphelids had significantly higher ASV richness in heated lakes (Fig. 3, Table 1), with a significant interaction between lake type and season. Chytrid ASV richness peaked in heated lakes during autumn and summer (Fig. 3A). Aphelid richness was higher in heated lakes in all seasons except spring (Fig. 3B). Peronosporomycete ASV richness varied seasonally, with higher values in spring and a peak in summer (Fig. 3C). Both Peronosporomycetes and Cryptomycota exhibited lower ASV richness in 2021 compared to 2020 (Fig. 3C, D, Table 1).

383 **Table 1.** Results of a type III ANOVA of GLMM investigating ASV richness for the four most ASV-

384 diverse parasite groups across lake type and season. Lake and sequencing run were specified as

385 random effects. The chytrid parasite richness model uniquely incorporated host richness (Chlorophyta,

386 Charophyta, Chromista, Gyrista, and Dinoflagellata) as a predictor. Significant results (*$p<0.05$) are

387 marked with an asterisk.

Parasite group	Fixed effect	χ^2	*p*-value
Chytrids	Lake type	50.1	<0.001*
	Season	4.2	0.241
	Host richness	3.4	0.066
	Lake type × season	23.4	<0.001*
Aphelids	Lake type	12.1	<0.001*
	Season	6.3	0.097
	Lake type × season	19.9	<0.001*
Peronosporomycetes	Lake type	0.00	1
(=Oomycetes)	Season	6.4	0.094
	Lake type × season	1.6	0.662
Cryptomycota	Lake type	1.5	0.216
	Season	8.4	0.038*
	Lake type × season	1.3	0.073

390 **Figure 3.** ASV richness of the four most ASV-rich parasite groups: (A) chytrids, (B) aphelids, (C) Peronosporomycetes (=Oomycetes) and (D) Cryptomycota.

391 Colors indicate lake type and symbols indicate the season of sample collection. Filled symbols denote individual data points, whereas larger open symbols

392 indicate seasonal mean values. Note that the y-axis scales differ across panels to optimize visualization.

Correlation between host and parasite richness

Chytrid parasite richness showed a marginally significant positive correlation with host richness (Table 1, Fig. 4). Heated lakes harbored particularly high host richness during summer (Fig. 4), which aligned with the observed elevated chytrid richness for that season (Fig. 3A). However, when examining the correlation between parasite and host richness separately within each lake type, a deviation from the linear model was found: chytrid richness was exceptionally high in heated lakes, despite comparable host richness between heated and control lakes (Fig. 4). This discrepancy aligned with the significant interaction between lake type and season.

Post-hoc tests conducted separately for each lake type showed that in heated lakes, both host richness (partial $R^2 = 0.72$, $p < 0.05$) and season (partial $R^2 = 0.58$, $p < 0.05$) strongly correlated with variation in parasite richness. In control lakes, the correlation between parasite richness and host richness (partial $R^2 = 0.32$, $p < 0.05$) and season (partial $R^2 = 0.11$, $p < 0.05$) was less pronounced (Table S6). These results indicated that while host richness and season significantly impacted chytrid parasite richness in both lake types (Table S6), heated lakes exhibited stronger seasonal variation and a higher correlation with host richness.

Figure 4. Correlation between chytrid parasite ASV richness (from the fungal dataset) and host ASV

richness across Chlorophyta, Charophyta, Chromista, Gyrista, and Dinoflagellata (from the protist

dataset). Shaded areas represent 95% confidence intervals. Colors differentiate between lake types, and

symbols indicate the season of sample collection.

Discussion

Heated lakes received hot water discharge since 1958 and are on average 2 °C warmer,

allowing us to investigate the long-term effects of warming on parasite assemblages. We

identified parasites from seasonally sampled eDNA to assess differences in parasite species

richness and assemblages between five artificially heated and five non-heated control lakes

over two years, using parasite-specific sequence databases, their metadata, and literature. Our

rigorous approach of classifying ASVs as parasites identified a diverse array of confirmed

aquatic parasites, spanning major eukaryotic taxa Alveolates (Apicomplexa, Ciliophora,

Dinoflagellata), Stramenopiles (Bigyra, Oomycetes), basal fungi (Chytrids, Rozellomycota) and

Ichthyosporeans, highlighting the undocumented parasite diversity in natural lakes.

We found that certain parasite species exhibit a strong preference for either heated or control

habitats, indicating that long-term warming can shape assemblages of parasite species. Chytrid

and aphelid parasite richness was notably higher in heated lakes, supporting the idea that

warming can enhance genotypic richness of parasites (Agosta et al., 2010). However, this trend

was not observed for Peronosporomycetes (=Oomycetes) or Cryptomycota, which instead

showed similar seasonal patterns of richness in all lakes (Peronosporomycete richness peaked

in spring and summer, Cryptomycota richness peaked in autumn and winter). As predicted by

the host-diversity-begets-parasite-diversity hypothesis (Kamiya et al., 2014; Krasnov & Poulin,

2015), chytrid parasite richness showed a positive correlation with host richness, but warming

altered the strength of the correlation. Post-hoc analyses revealed that season and host

richness correlated more strongly with parasite richness in heated lakes than control lakes,

suggesting that warming may both amplify seasonal variation and increase the strength of

correlation between host and parasite richness. Overall, these findings support the idea that

thermal conditions strongly influence the composition and dynamics of parasite communities,

possibly through a combination of direct effects on parasites and indirect effects via hosts.

The Konin lake complex harbors ecologically relevant parasites, though a

majority are unidentified

Our study identified 269 parasites that were assigned to species level in the protist dataset, or

were >98% similar to a known parasite in GenBank. However, 86% of parasite ASVs in the

protist dataset and 53% parasite ASVs in the fungal dataset could not be identified to species

level (see Tables S3 & S4). This highlights the acute shortage of taxonomic information and

reference sequences for freshwater parasites. Despite limited research, microeukaryotic

parasites are known to impact ecosystem function. For example, zoospores of the chytrid

Zygorhizydium affluens released from parasitized diatom blooms, provide a nutritious food

source for *Daphnia*, thereby establishing a trophic link between producers and primary

consumers (Kagami et al., 2007). In addition to *Z. affluens* (ASVs 31 and 717 in the current

study)*,* other parasite ASVs were >95% similar to known but as yet uncharacterized chytrid

parasites, such as the chitin-degrading *Rhizoclosmatium pessaminum* (ASV 1901) (Powell et

al., 2019), *Endochytrium ramosum* (ASVs 1418 and 2290), and *Staurastromyces oculus* (ASVs

1236, 1328 and 3041). These chytrids parasitize the diatom *Asterionella* (Rad-Menéndez et al.,

2018), and green algae *Chara*, *Cladophora* (Simmons et al., 2020), and *Straurastrum* (Van den

Wyngaert et al., 2017, see Table S43). Other ecologically relevant parasites detected in our

study included two ASVs (181 and 1195; see Table S3) that showed >98% sequence similarity

with *Psorospermium haeckeli* (Class: Ichythyophonida), a parasite of the crayfish *Ascatus*

ascatus (Rug & Vogt, 1994)*,* a host species that is impacted by the devastating Europe-wide

crayfish plague (Wiśniewski et al., 2020)*.* We also identified four ASVs (1156, 2064, 1231, and

1305, see Table S3) of the Apicomplexan parasite *Nematopsis temporariae*, which infects a

range of amphibian hosts (Chambouvet et al., 2016).

ASV 286 (Family: Peronosporales) showed a strong preference for heated lakes (see Fig. 2)

and exhibited 96.4% sequence similarity with *Pythium insidiosum*, a generalist oomycete

parasite known to cause devastating root rot in commercial crops (Golińska & Świecimska,

2020), as well as disease in mammals (Mendoza, 2009) including humans (Mendoza et al.,

1993). *Pythium* releases motile zoospores from infected plant roots or skin lesions (Mendoza et

al., 1993). The close similarity of ASV 286 to a clinical sample from a patient with pythiosis (see

Krajaejun et al., (2014) and GenBank BBXB02000196) suggests that heated lakes might

function as a reservoir, providing suitable habitats or alternative hosts for this parasite.

We additionally found three parasite ASVs that showed 90-93% sequence similarity to

Batrachochytrium salamandrivorans (*Bsal*), of which ASV 764 was predominantly detected in

winter within heated lakes Gosławskie, Licheńskie, Mikorzyńskie, and Pątnowskie (see Table

S5, Fig. S6). *Bsal* is responsible for global chytridiomycosis outbreaks in amphibians (Balàž et

al., 2017; Laking et al., 2017; Yap et al., 2015). *Bsal* produces motile, water-borne transmission

stages (zoospores) that are active within a thermal range of 10-15 °C but capable of surviving at

temperatures as low as 5 °C (Martel et al., 2013). Consequently, epidemics are generally

observed during colder months, such as late autumn, winter or spring (Schulz et al., 2020).

Geospatial models predict a reduction in climatically suitable habitats for *Bsal* under global

warming scenarios (Grisnik et al., 2023). However, recent chytridiomycosis outbreaks at water

temperatures as high as 26.4 °C (Laking et al., 2017) suggest a broader thermal niche than

previously anticipated. Although *Bsal* infections have so far not been detected in Poland, its

sister species *B. dendrobatidis* is widespread (Palomar et al., 2021). The detection of *Bsal* in

the heated Konin lakes during winter suggests that warm-winter conditions may provide a

favourable environment for this parasite. These findings underscore the need for additional

research, particularly regarding host demographics (Bielby et al., 2021), to assess the potential

presence and implications of *Bsal* in the heated environments of the Konin lakes.

Warming-induced shifts in parasite distribution and elevation in richness are

mirrored by similar shifts in host taxa

A total of 39 parasite ASVs exhibited a strong preference for specific lake types, indicating

distinct parasite assemblages in heated versus non-heated habitats. Of these, 34 were more

prevalent in heated lakes, including *Aphelidium desmodesmi* and four other ASVs belonging to

Aphelidiaceae. Although the thermal preference of aphelid parasites remains underexplored, *A.*

desmodesmi infections in the green alga *Scenedesmus* were previously shown to increase

under elevated summer temperatures (22-25 °C) and phosphorus-limiting conditions in a

mesocosm study (Alam et al., 2024). Consistent with this, green algal and diatom hosts of *A.*

desmodesmi, including *Desmodesmus armatus* (=*Scenedesmus armatus*, ASV 27), have

shown a preference for, and higher relative abundance in heated lakes (Beng et al., 2023). *D.*

armatus typically shows optimum growth at high temperatures (~27 °C) (S. Wang et al., 2019),

matching the summer water temperatures of the heated Konin lakes. This suggests that

warming may increase aphelid parasite richness both by directly facilitating epidemics at

elevated temperatures and indirectly by promoting host species growth. Similarly, the parasitic

dinoflagellate *Dissodinium pseudolunula* (ASV 110), which infects copepod eggs, also showed

a preference for heated lakes, suggesting that warming may favor this parasite as well.

Conversely, two unidentified chytrid ASVs (34 and 263; Order: Rhizophydiales) showed a

preference for non-heated lakes and were associated with the diatom host *Ulnaria* (=*Synedra*),

which also showed a preference for control lakes (Beng et al., 2023). The thermal preferences

of other indicator parasites, have not been extensively investigated in semi-natural or laboratory

settings. Overall, these findings suggest that long-term warming has led to the development of

distinct parasite assemblages, likely influenced by host distribution patterns, despite the

potential flow of water between heated and non-heated lakes (Zdanowski et al., 2020).

The richness of chytrid parasites in our study (which was elevated in heated lakes) showed a

general positive correlation with host richness, despite known variation in the extent of their

host-specificity. The host-diversity-begets-parasite-diversity hypothesis is proposed to be driven

by host-specificity of parasites (Krasnov & Poulin, 2015). While some chytrid parasites are

highly host-specific (Van den Wyngaert et al., 2018), other evidence suggests that the

specificity might be flexible, particularly during the initial "attraction" phase when motile

zoospores find and attach to potential host cells. The chemical cues that attract motile

zoospores, such as dimethylsulfoniopropionate (DMSP), are derived from algal cells and are

non-specific, suggesting that the attraction might not be limited to particular algal species

(Frenken et al., 2017; Vallet, 2024). Additionally, genotype-specificity of some chytrids may shift

across thermal regimes, as seen in *Zygorhizidium planktonicum* infecting the diatom *A. formosa*

(Gsell et al., 2013). Nevertheless, the positive correlation between host and parasite richness in

our study suggests some degree of specificity between chytrid parasites and their compatible

host species (or genotypes). Increased host richness may also support a greater richness of

generalist parasites. Chytrids that exhibit a 'generalist' versus 'specialist' infection strategy often

occur at different times of the year (Kagami et al., 2021; Van den Wyngaert et al., 2022). For

example, the seasonal abundance of chytrid parasites often corresponds with fluctuations in

their specific phytoplankton host taxa (e.g. diatoms, cyanobacteria) (Rasconi et al., 2012),

suggesting that parasites may "track" the most abundant host species. During spring diatom

blooms, chytrids tend to specialize on the dominant diatom species, while an even host

community towards the end of the bloom supports the persistence of generalist chytrids

(Sassenhagen et al., 2023). Thus, for chytrid parasites, increased host richness may support

higher parasite richness regardless of whether the parasites are specialist or generalist,

potentially at different times of the year, contributing to their overall increased richness in heated

lakes.

Warming alters seasonal richness patterns of parasitic chytrids, but not

Peronosporomycetes (=Oomycetes) or Cryptomycota

The richness of chytrid parasites in heated lakes exceeded expectations based on host richness

alone during autumn. One potential explanation for this finding is the significantly higher soluble

phosphorus content observed in heated lakes during this season (Beng et al., 2023). Phosphate

levels can affect both hosts and parasites. Elevated phosphorus is linked to increased infection

prevalence of some chytrids, such as *Rhizosiphon crassum* infecting its cyanobacterial host

Anabaena flosaquae (Rasconi et al., 2012). Conversely, low phosphorus levels can reduce the

growth rate of diatom hosts such as *A. formosa* without impacting the growth rate of their

Rhizophydium parasites, leading to epidemics at lower host densities (Bruning, 1991). Heated

lakes harbor cyanobacterial blooms in the autumn, which could grow to high densities due to

elevated phosphorus, and serve as alternative hosts – albeit none of the chytrid parasites in our

study associated with cyanobacterial hosts (see Fig. 1B). Thus, the increase in phosphorus

levels in heated lakes during autumn (Beng et al., 2023) could underlie the unexpectedly high

chytrid parasite richness in autumn.

Warming is hypothesized to induce shifts in the host range of parasites (Wells & Clark, 2019), or

favor generalist parasites due to their ability to infect multiple hosts (Cizauskas et al., 2017).

Four indicator parasites - the mussel *Elliptio complanata,* fungus *Ustilago esculenta,* and

chytrids *Dinomyces arenysensis,* and *Dangeardia mamillata*, all known host generalists – were

more prevalent in heated lakes (see Fig. 2). *E. complanata* (ASV 462) is an ectoparasite

(Gendron et al., 2019) that can attach to at least 40 fish host species spanning 14 families

(Lellis et al., 2013). *U. esculenta* belongs to a hybridizing species complex of multicellular fungi

that parasitize multiple hosts within the *Poaceae* grass family (reviewed in McTaggart et al.,

2012)*.* The chytrid *Di. arenysensis* infects at least four genera of dinoflagellates (Lepelletier et

al., 2014), whereas *Da. mamillata* infects green algae from at least three genera (Van den

Wyngaert et al., 2018). Other indicator taxa, including the aphelid *Aphelidium desmodesmi* and

the chytrids *Quaeritorhiza haematococci* and *Rhizophydium chlorogonii* (=*Aquamyces*

chlorogonii), were also more prevalent in heated lakes, but their host ranges remain

unknown.(Beng et al., 2023)(Hutorowicz, 2006)(Dunn et al., 2008)Further research on host

ranges is needed to test (Agosta et al., 2010; Wells & Clark, 2019) the potential for warming to

favour generalist parasites (Cizauskas et al., 2017).

In contrast with chytrids, warming did not alter seasonal richness patterns of oomycetes or

Cryptomycota. Oomycete richness peaked during spring and summer in the current study,

potentially aligning with the elevated abundance of airborne (Lang-Yona et al., 2018) and

freshwater (Hallett & Dick, 1981) oomycetes documented in spring and summer. However,

there are no studies investigating seasonal variation in oomycete parasite species richness,

especially among freshwater taxa. Cryptomycota are known to hyperparasitize chytrids

(reviewed in Bermúdez-Cova et al., 2023), and are thus proposed to impact phytoplankton

communities (Gleason et al., 2017). In the current study, Cryptomycota richness peaked in

autumn and winter. This could potentially be linked with the peak in parasitic chytrid richness in

autumn; however, further investigation at greater temporal resolution is required to confirm this.

Furthermore, not all Cryptomycota hyperparasitize chytrids: some are also reported to infect

amoebae (Seto et al., 2023), highlighting how research on the host ranges or thermal

preferences of these enigmatic groups (Cryptomycota and oomycetes) is currently insufficient to

model expectations of their potential responses to warming.

Limitations of eDNA-based monitoring for disease ecology

eDNA metabarcoding is a powerful tool for biodiversity-monitoring, but has well-documented

limitations (Garlapati et al., 2019). Two key limitations are the lack of quantitative estimates of

taxon abundance and the inability to distinguish between the presence of live organisms and

that of dormant life stages such as spores. Overcoming these issues by complementing eDNA

metabarcoding with quantitative and organismal approaches could improve our insight into the

effects of warming on disease.

One important aspect of disease ecology is the abundance of hosts, which cannot be estimated

from a metabarcoding approach alone, and which significantly impacts parasitism in the

environment. In our study, heated lakes exhibited stronger seasonal variation in parasite

richness than control lakes, which may be due to the unexpectedly high chytrid parasite

richness in autumn. This suggests that seasonal ecosystem-wide processes driven by warming

may contribute to the peak in parasite richness in autumn. Moreover, the higher parasite

richness and distinct species assemblages in heated lakes, combined with previously reported

elevation in host richness in these environments (Beng et al., 2023), raises the possibility that

warming may foster novel host-parasite encounters. For instance, heated lakes harbor the

invasive tape grass *Vallisneria spiralis*, which peaks in biomass during autumn (Hutorowicz,

2006). *Vallisneria* macrophytes can increase the abundance of their epibiont taxa that include

green algae, diatoms, and dinoflagellates (Dunn et al., 2008), all of which serve as

phytoplankton hosts for a broad range of chytrid parasites (see Fig. 1B). We speculate that this

may promote abundance of hosts and generate 'hotspots' for novel encounters between

potential host species and motile zoospores of generalist parasites. This conceivably increases

the likelihood of host shifts, and potentially contributes to the high chytrid parasite richness in

heated lakes observed in autumn. Such hypothesized outcomes of warming (Agosta et al.,

2010) require quantifying host and parasite abundance in addition to richness to gain a

complete understanding of the effects of warming on disease.

Another limitation of eDNA is that detected taxa may not be biologically active, or may originate

from dormant dispersal stages. This aligns with our observation that 27% of parasite ASVs from

the protist dataset and 41% from the fungal dataset were unexpectedly associated with

terrestrial hosts, including plants, animals and humans (See Fig. 1). As one example, the fungal

ASV 259 shared 100% sequence similarity with *Botrytis cineria* (Phylum: Ascomycota, GenBank

AY544651), a pathogen that disperses through airborne spores and causes grey mold in

terrestrial plant hosts (Williamson et al., 2007). *B. cineria* has also been shown to cause severe

developmental abnormalities and fitness loss in aquatic organisms like the zebrafish *Danio rerio*

(Shi et al., 2023, see Table S4). This illustrates how some fungal parasites, although typically

associated with terrestrial hosts, may impact aquatic species. Despite such a possibility, we

cannot rule out that this taxon was detected from dormant spores. eDNA-based detection must

therefore be complemented with organismal approaches such as microscopic counts to confirm

the presence and abundance of live parasites *in situ*.

Conclusion

Our study revealed a vast diversity of eukaryotic parasitic plankton species, with warming

shown to increase parasite richness for some groups such as chytrids and aphelids. Parasites

exhibited distinct habitat preferences, favouring either heated or non-heated lakes, resulting in

unique parasite species assemblages. These preferences might be linked to host species

distributions. For chytrid parasites, increased richness in heated lakes can be partially attributed

to greater host richness in these environments. For chytrid parasites, warming amplified

seasonal variation in richness and increased the strength of correlation between host and

parasite richness. Habitat preferences of parasites suggest that warming can shape species

assemblages by impacting both abiotic factors (such as thermal conditions in heated lakes) and

biotic factors (such as host richness), demonstrating how parasite richness and assemblages

may shift under global change.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

Raw sequence data for the first sequencing run are available in the European Nucleotide

Database (ENA) under BioProject ID PRJEB54709 and those of the second run are available at

NCBI Sequence Read Archive (SRA) under BioProject ID PRJNA1201567. Reproducible R

code for bioinformatic processing is provided in supplementary R scripts.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was funded by a joint Beethoven Life-1 grant from the German Science Foundation

(WO 1587/9-1 to JW) and the National Science Centre, Poland (2018/31/F/NZ8/01986 to SC).

Authors' contributions

JW and SC secured funding and designed the project with MTM. JW and SC collected eDNA

samples. AR and KCB performed bioinformatic processing of samples. AR performed ecological

analyses and wrote the manuscript with input from all co-authors.

Acknowledgement

We thank Elisabeth Funke for conducting metabarcoding library preparation, and Sarah

Sparmann and Susan Mbedi at the Berlin Center for Genomics in Biodiversity Research

(BeGenDiv: https://begendiv.de/) for sequencing support. We further thank the Disease

Evolutionary Ecology, and Molecular Ecology & Genomics groups at IGB Berlin for constructive

feedback, as well as Nedim Tüzün for stimulating discussions. We are also grateful to two

anonymous reviewers whose constructive feedback significantly improved our manuscript.

References

Abonyi, A., Fornberg, J., Rasconi, S., Ptacnik, R., Kainz, M. J., & Lafferty, K. D. (2024). The chytrid insurance hypothesis: integrating parasitic chytrids into a biodiversity–ecosystem functioning framework for phytoplankton–zooplankton population dynamics. *Oecologia*, *204*(2), 279–288. https://doi.org/10.1007/s00442-024-05519-w

Agosta, S. J., Janz, N., & Brooks, D. R. (2010). How specialists can be generalists: Resolving the and "parasite paradox" and implications for emerging infectious disease. *Zoologia*, *27*(2), 151–162. https://doi.org/10.1590/S1984-46702010000200001

Alam, M. M., Hodaei, M., Hartnett, E., Gincley, B., Khan, F., Kim, G. Y., Pinto, A. J., & Bradley, I. M. (2024). Community structure and function during periods of high performance and system upset in a full-scale mixed microalgal wastewater resource recovery facility. *Water Research*, *259*. https://doi.org/10.1016/j.watres.2024.121819

Bakhtiyar, Y., Arafat, M. Y., Andrabi, S., & Tak, H. I. (2020). Zooplankton: The Significant Ecosystem Service Provider in Aquatic Environment. In *Bioremediation and Biotechnology, Vol 3* (pp. 227–244). Springer International Publishing. https://doi.org/10.1007/978-3-030-46075-4_10

Balàž, V., Gortázar Schmidt, C., Murray, K., Carnesecchi, E., Garcia, A., Gervelmeyer, A., Martino, L., Munoz Guajardo, I., Verdonck, F., Zancanaro, G., & Fabris, C. (2017). Scientific and technical assistance concerning the survival, establishment and spread of *Batrachochytrium salamandrivorans (Bsal)* in the EU. *EFSA Journal*, *15*(2). https://doi.org/10.2903/j.efsa.2017.4739

Bartley, T. J., McCann, K. S., Bieg, C., Cazelles, K., Granados, M., Guzzo, M. M., MacDougall, A. S., Tunney, T. D., & McMeans, B. C. (2019). Food web rewiring in a changing world. In *Nature Ecology and Evolution* (Vol. 3, Issue 3, pp. 345–354). Nature Publishing Group. https://doi.org/10.1038/s41559-018-0772-3

Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R., Singmann, H., Scheipl, F., Grothendieck, G., Green, P., Fox, J., Bauer, A., Krivitsky, P., Tanaka, E., & Jagan, M. (2024). Linear Mixed-Effects Models using "Eigen" and S4 (package lme4). *CRAN*.

Beng, K. C., Cerbin, S., Monaghan, M. T., & Wolinska, J. (2023). Long-term effects of surface-water temperature increase on plankton communities in artificially heated lakes: insights from eDNA metabarcoding. *Authorea Preprints*. https://doi.org/10.22541/au.168639128.85081228/v1

Beng, K. C., Wolinska, J., Funke, E., Van Den Wyngaert, S., Gsell, A. S., & Monaghan, M. T. (2021). Temporal dynamics of freshwater planktonic parasites inferred using a DNA metabarcoding time-series. *Parasitology*, *148*(13), 1602–1611. https://doi.org/10.1017/S0031182021001293

Bennett, L., Melchers, B., & Proppe, B. (2020). *Curta: A General-purpose High-Performance Computer at ZEDAT, Freie Universität Berlin*. https://refubium.fu-berlin.de/handle/fub188/26993

Bermúdez-Cova, M. A., Haelewaters, D., Bekker, C. de, Piepenbring, M., Schoutteten, N., & Quandt, C. A. (2023). *Hyperparasitic fungi—definitions, diversity, ecology, and research*. https://doi.org/10.22541/au.168787020.07281183/v2

Bielby, J., Price, S. J., Monsalve-CarcaÑo, C., & Bosch, J. (2021). Host contribution to parasite
persistence is consistent between parasites and over time, but varies spatially. *Ecological
Applications*, *31*(3). https://doi.org/10.1002/eap.2256

Bruning, K. (1991). Effects of phosphorus limitation on the epidemiology of a chytrid
phytoplankton parasite. *Freshwater Biology*, *25*(3), 409–417.
https://doi.org/10.1111/j.1365-2427.1991.tb01385.x

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P.
(2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature
Methods*, *13*(7), 581–583. https://doi.org/10.1038/nmeth.3869

Calvin, K., Dasgupta, D., Krinner, G., Mukherji, A., Thorne, P. W., Trisos, C., Romero, J.,
Aldunce, P., Barrett, K., Blanco, G., Cheung, W. W. L., Connors, S., Denton, F., Diongue-
Niang, A., Dodman, D., Garschagen, M., Geden, O., Hayward, B., Jones, C., … Ha, M.
(2023). *IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working
Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on
Climate Change [Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva,
Switzerland.* (P. Arias, M. Bustamante, I. Elgizouli, G. Flato, M. Howden, C. Méndez-
Vallejo, J. J. Pereira, R. Pichs-Madruga, S. K. Rose, Y. Saheb, R. Sánchez Rodríguez, D.
Ürge-Vorsatz, C. Xiao, N. Yassaa, J. Romero, J. Kim, E. F. Haites, Y. Jung, R. Stavins, …
C. Péan, Eds.). https://doi.org/10.59327/IPCC/AR6-9789291691647

Carlson, C. J., Burgio, K. R., Dougherty, E. R., Phillips, A. J., Bueno, V. M., Clements, C. F.,
Castaldo, G., Dallas, T. A., Cizauskas, C. A., Cumming, G. S., Doña, J., Harris, N. C.,
Jovani, R., Mironov, S., Muellerklein, O. C., Proctor, H. C., & Getz, W. M. (2017). Parasite
biodiversity faces extinction and redistribution in a changing climate. *Science Advances*, *3*.
https://www.science.org

Chambouvet, A., Valigurová, A., Pinheiro, L. M., Richards, T. A., & Jirků, M. (2016). *Nematopsis
temporariae* (Gregarinasina, Apicomplexa, Alveolata) is an intracellular infectious agent of
tadpole livers. *Environmental Microbiology Reports*, *8*(5), 675–679.
https://doi.org/10.1111/1758-2229.12421

Cichy, A., Marszewska, A., Ciapka, P., & Żbikowska, E. (2020). Different thermal conditions of
lakes affect host–parasite systems: A case study of *Viviparus contectus* (Millet, 1813) and
digenean trematodes. *Freshwater Biology*, *65*(3), 417–425.
https://doi.org/10.1111/fwb.13434

Cizauskas, C. A., Carlson, C. J., Burgio, K. R., Clements, C. F., Dougherty, E. R., Harris, N. C.,
& Phillips, A. J. (2017). Parasite vulnerability to climate change: An evidence-based
functional trait approach. *Royal Society Open Science*, *4*(1).
https://doi.org/10.1098/rsos.160535

Danz, A., & Quandt, C. A. (2023). A review of the taxonomic diversity, host–parasite
interactions, and experimental research on chytrids that parasitize diatoms. In *Frontiers in
Microbiology* (Vol. 14). Frontiers Media SA. https://doi.org/10.3389/fmicb.2023.1281648

De Caceres, M., & Legendre, P. (2009). Associations between species and groups of sites:
indices and statistical inference. *Ecology*, *90*(12), 3566–3574.

Dunn, A. E., Dobberfuhl, D. R., & Casamatta, D. A. (2008). A survey of algal epiphytes from
Vallisneria americana Michx. (Hydrocharitaceae) in the lower St. Johns River, Florida.
Southeastern Naturalist, *7*(2), 229–244. https://doi.org/10.1656/1528-
7092(2008)7[229:ASOAEF]2.0.CO;2

Dziuba, M. K., Cerbin, S., & Wejnerowski, L. (2017). Is bigger better? A possibility for adaptation of *Daphnia* to filamentous cyanobacteria in the face of global warming. *Hydrobiologia*, *798*(1), 105–118. https://doi.org/10.1007/s10750-015-2438-3

Dziuba, M. K., Herdegen-Radwan, M., Pluta, E., Wejnerowski, Ł., Szczuciński, W., & Cerbin, S. (2020). Temperature increase altered *Daphnia* community structure in artificially heated lakes: a potential scenario for a warmer future. *Scientific Reports*, *10*(1). https://doi.org/10.1038/s41598-020-70294-6

Dziuba, M. K., Manzi, F., Cerbin, S., & Wolinska, J. (2023). Can climate warming save *Daphnia* from parasites? Reduced parasite prevalence in *Daphnia* populations from artificially heated lakes. *Limnology and Oceanography*, *68*(1), 181–191. https://doi.org/10.1002/lno.12257

Ejsmont-Karabin, J., Hutorowicz, A., Kapusta, A., Stawecki, K., Tunowski, J., & Zdanowski, B. (2020). Rotifers in heated Konin lakes-A review of long-term observations. *Water (Switzerland)*, *12*(6). https://doi.org/10.3390/w12061660

Fernández-Valero, A. D., Karpov, S. A., Sampedro, N., Gordi, J., Timoneda, N., Garcés, E., & Reñé, A. (2024). Newly identified diversity of Dinomycetaceae (Rhizophydiales, Chytridiomycota), a family of fungal parasites of marine dinoflagellates. *European Journal of Protistology*, *93*. https://doi.org/10.1016/j.ejop.2024.126053

Fernández-Valero, A. D., Reñé, A., Timoneda, N., Pou-Solà, N., Gordi, J., Sampedro, N., & Garcés, E. (2023). The succession of epiphytic microalgae conditions fungal community composition: how chytrids respond to blooms of dinoflagellates. *ISME Communications*, *3*(1). https://doi.org/10.1038/s43705-023-00304-x

Fox, J., Weisberg, S., Price, B., Adler, D., Bates, D., Baud-Bovy, G., Bolker, B., Ellison, S., Firth, D., Friendly, M., Gorjanc, G., Graves, S., Heiberger, R., Krivitsky, P., Laboissierre, R., Maechler, M., Monette, G., Murdoch, D., Nilsson, H., … Zeilais, A. (2019). car: Companion to Applied Regression. *CRAN*.

Frenken, T., Alacid, E., Berger, S. A., Bourne, E. C., Gerphagnon, M., Grossart, H. P., Gsell, A. S., Ibelings, B. W., Kagami, M., Küpper, F. C., Letcher, P. M., Loyau, A., Miki, T., Nejstgaard, J. C., Rasconi, S., Reñé, A., Rohrlack, T., Rojas-Jimenez, K., Schmeller, D. S., … Agha, R. (2017). Integrating chytrid fungal parasites into plankton ecology: research gaps and needs. In *Environmental Microbiology* (Vol. 19, Issue 10, pp. 3802–3822). Blackwell Publishing Ltd. https://doi.org/10.1111/1462-2920.13827

Frenken, T., Velthuis, M., de Senerpont Domis, L. N., Stephan, S., Aben, R., Kosten, S., van Donk, E., & Van de Waal, D. B. (2016). Warming accelerates termination of a phytoplankton spring bloom by fungal parasites. *Global Change Biology*, *22*(1), 299–309. https://doi.org/10.1111/gcb.13095

Gad, M., Fawzy, M. E., Al-Herrawy, A. Z., Abdo, S. M., Nabet, N., & Hu, A. (2023). PacBio next-generation sequencing uncovers Apicomplexa diversity in different habitats. *Scientific Reports*, *13*(1). https://doi.org/10.1038/s41598-023-40895-y

Garlapati, D., Charankumar, B., Ramu, K., Madeswaran, P., & Ramana Murthy, M. V. (2019). A review on the applications and recent advances in environmental DNA (eDNA) metagenomics. *Reviews in Environmental Science and Biotechnology*, *18*(3), 389–411. https://doi.org/10.1007/s11157-019-09501-4

Gendron, A. D., Sanchez, D., Douville, M., & Houde, M. (2019). Stress-related gene transcription in fish exposed to parasitic larvae of two freshwater mussels with divergent

790   infection strategies. *Diseases of Aquatic Organisms*, *132*(3), 191–202.
791   https://doi.org/10.3354/dao03319

792 Gilarranz, L. J., Narwani, A., Odermatt, D., Siber, R., Dakos D Edited, V., & Hastings, A. (2022).
793   *Regime shifts, trends, and variability of lake productivity at a global scale*.
794   https://doi.org/10.1073/pnas

795 Gleason, F. H., Jephcott, T. G., Küpper, F. C., Gerphagnon, M., Sime-Ngando, T., Karpov, S.
796   A., Guillou, L., & van Ogtrop, F. F. (2015). Potential roles for recently discovered chytrid
797   parasites in the dynamics of harmful algal blooms. In *Fungal Biology Reviews* (Vol. 29,
798   Issue 1, pp. 20–33). Elsevier Ltd. https://doi.org/10.1016/j.fbr.2015.03.002

799 Gleason, F. H., Scholz, B., Jephcott, T. G., van Ogtrop, F. F., Henderson, L., Lilje, O.,
800   Kittelmann, S., & Macarthur, D. J. (2017). Key Ecological Roles for Zoosporic True Fungi in
801   Aquatic Habitats. *Microbiology Spectrum*, *5*(2). https://doi.org/10.1128/microbiolspec.funk-
802   0038-2016

803 Golińska, P., & Świecimska, M. (2020). *Pythium*: Diseases and Their Management. In M. Rai, K.
804   A. Abd-Elsalam, & A. P. Ingle (Eds.), *Pythium*. CRC Press.
805   https://doi.org/10.1201/9780429296406

806 Grant, L., Vanderkelen, I., Gudmundsson, L., Tan, Z., Perroud, M., Stepanenko, V. M.,
807   Debolskiy, A. V., Droppers, B., Janssen, A. B. G., Woolway, R. I., Choulga, M., Balsamo,
808   G., Kirillin, G., Schewe, J., Zhao, F., del Valle, I. V., Golub, M., Pierson, D., Marcé, R., …
809   Thiery, W. (2021). Attribution of global lake systems change to anthropogenic forcing.
810   *Nature Geoscience*, *14*(11), 849–854. https://doi.org/10.1038/s41561-021-00833-x

811 Grisnik, M., Gray, M. J., Piovia-Scott, J., Carter, E. D., & Sutton, W. B. (2023). Incorporating
812   caudate species susceptibilities and climate change into models of *Batrachochytrium*
813   *salamandrivorans* risk in the United States of America. *Biological Conservation*, *284*.
814   https://doi.org/10.1016/j.biocon.2023.110181

815 Gsell, A. S., de Senerpont Domis, L. N., van Donk, E., & Ibelings, B. W. (2013). Temperature
816   Alters Host Genotype-Specific Susceptibility to Chytrid Infection. *PLoS ONE*, *8*(8).
817   https://doi.org/10.1371/journal.pone.0071737

818 Guillou, L., Bachar, D., Audic, S., Bass, D., Berney, C., Bittner, L., Boutte, C., Burgaud, G., De
819   Vargas, C., Decelle, J., Del Campo, J., Dolan, J. R., Dunthorn, M., Edvardsen, B.,
820   Holzmann, M., Kooistra, W. H. C. F., Lara, E., Le Bescot, N., Logares, R., … Christen, R.
821   (2013). The Protist Ribosomal Reference database (PR2): A catalog of unicellular
822   eukaryote Small Sub-Unit rRNA sequences with curated taxonomy. *Nucleic Acids*
823   *Research*, *41*(D1). https://doi.org/10.1093/nar/gks1160

824 Hallett, I. C., & Dick, M. W. (1981). Seasonal and Diurnal Fluctuations of Oomycete Propagule
825   Numbers in the Free Water of a. In *Source: Journal of Ecology* (Vol. 69, Issue 2).
826   https://www.jstor.org/stable/2259691

827 Hillbricht-Ilkowska, A., & Zdanowski, B. (1988). Changes in lake ecosystems connected with the
828   power-generating industry (the outline of problem); the Konin Lakes (Poland) as the study
829   sites. *Ekol. Pol*, *36*, 5–21.

830 Hudson, P. J., Dobson, A. P., & Lafferty, K. D. (2006). Is a healthy ecosystem one that is rich in
831   parasites? *Trends in Ecology and Evolution*, *21*(7), 381–385.
832   https://doi.org/10.1016/j.tree.2006.04.007

833 Hutorowicz, A. (2006). *Vallisneria spiralis L.* (Hydrocharitaceae) in Lakes in the Vicinity of Konin
834   (Kujawy Lakeland). *Biodiv. Res. Conserv.*, *2*(1), 154–158. www.brc.amu.edu.pl

Ilicic, D., Woodhouse, J., Karsten, U., Schimani, K., Zimmermann, J., & Grossart, H. P. (2024).
Chytrid fungi infecting Arctic microphytobenthic communities under varying salinity
conditions. *Scientific Reports*, *14*(1), 25821. https://doi.org/10.1038/s41598-024-77202-2

James, T. Y., Pelin, A., Bonen, L., Ahrendt, S., Sain, D., Corradi, N., & Stajich, J. E. (2013).
Shared signatures of parasitism and phylogenomics unite Cryptomycota and Microsporidia.
Current Biology, *23*(16), 1548–1553. https://doi.org/10.1016/j.cub.2013.06.057

Jane, S. F., Hansen, G. J. A., Kraemer, B. M., Leavitt, P. R., Mincer, J. L., North, R. L., Pilla, R.
M., Stetler, J. T., Williamson, C. E., Woolway, R. I., Arvola, L., Chandra, S., DeGasperi, C.
L., Diemer, L., Dunalska, J., Erina, O., Flaim, G., Grossart, H. P., Hambright, K. D., …
Rose, K. C. (2021). Widespread deoxygenation of temperate lakes. *Nature*, *594*(7861),
66–70. https://doi.org/10.1038/s41586-021-03550-y

Jephcott, T. G., van Ogtrop, F. F., Gleason, F. H., Macarthur, D. J., & Scholz, B. (2017). The
ecology of chytrid and aphelid parasites of phytoplankton. In J. Dighton & J. F. White
(Eds.), *The Fungal Community* (pp. 239–255). CRC Press.
https://doi.org/10.1201/9781315119496

Johnson, P. T. J., Preston, D. L., Hoverman, J. T., & LaFonte, B. E. (2013). Host and parasite
diversity jointly control disease risk in complex communities. *Proceedings of the National
Academy of Sciences of the United States of America*, *110*(42), 16916–16921.
https://doi.org/10.1073/pnas.1310557110

Johnson, P. T. J., Wood, C. L., Joseph, M. B., Preston, D. L., Haas, S. E., & Springer, Y. P.
(2016). Habitat heterogeneity drives the host-diversity-begets-parasite-diversity
relationship: evidence from experimental and field studies. *Ecology Letters*, *19*(7), 752–
761. https://doi.org/10.1111/ele.12609

Kagami, M., Miki, T., & Takimoto, G. (2014). Mycoloop: Chytrids in aquatic food webs. *Frontiers
in Microbiology*, *5*(APR). https://doi.org/10.3389/fmicb.2014.00166

Kagami, M., Seto, K., Nozaki, D., Nakamura, T., Wakana, H., & Wurzbacher, C. (2021). Single
dominant diatom can host diverse parasitic fungi with different degree of host specificity.
Limnology and Oceanography, *66*(3), 667–677. https://doi.org/10.1002/lno.11631

Kagami, M., von Elert, E., Ibelings, B. W., de Bruin, A., & Van Donk, E. (2007). The parasitic
chytrid *Zygorhizidium* facilitates the growth of the Cladoceran zooplankter, *Daphnia* in
cultures of the inedible alga *Asterionella*. *Proceedings of the Royal Society B: Biological
Sciences*, *274*(1617), 1561–1566. https://doi.org/10.1098/rspb.2007.0425

Kamiya, T., O'Dwyer, K., Nakagawa, S., & Poulin, R. (2014). Host diversity drives parasite
diversity: Meta-analytical insights into patterns and causal mechanisms. *Ecography*, *37*(7),
689–697. https://doi.org/10.1111/j.1600-0587.2013.00571.x

Kapusta, A., & Bogacka-Kapusta, E. (2015). Non-native fish species in heated lakes: Origins
and present status. *Archives of Polish Fisheries*, *23*(2), 121–129.
https://doi.org/10.1515/aopf-2015-0014

King, K. C., Hall, M. D., & Wolinska, J. (2023). Infectious disease ecology and evolution in a
changing world. In *Philosophical Transactions of the Royal Society B: Biological Sciences*
(Vol. 378, Issue 1873). Royal Society Publishing. https://doi.org/10.1098/rstb.2022.0002

Krajaejun, T., Lerksuthirat, T., Garg, G., Lowhnoo, T., Yingyong, W., Khositnithikul, R.,
Tangphatsornruang, S., Suriyaphol, P., Ranganathan, S., & Sullivan, T. D. (2014).
Transcriptome analysis reveals pathogenicity and evolutionary history of the pathogenic

oomycete *Pythium insidiosum*. *Fungal Biology*, *118*(7), 640–653.
https://doi.org/10.1016/j.funbio.2014.01.009

Krasnov, B. R., & Poulin, R. (2015). Relationships between parasite diversity and host diversity.
In S. Morand, B. R. Krasnov, & D. T. J. Littlewood (Eds.), *Parasite Diversity and*
Diversification (pp. 27–38). Cambridge University Press.
https://doi.org/10.1017/CBO9781139794749

Laking, A. E., Ngo, H. N., Pasmans, F., Martel, A., & Nguyen, T. T. (2017). *Batrachochytrium*
salamandrivorans is the predominant chytrid fungus in Vietnamese salamanders. *Scientific*
Reports, *7*(1), 44443. https://doi.org/10.1038/srep44443

Lang-Yona, N., Pickersgill, D. A., Maurus, I., Teschner, D., Wehking, J., Thines, E., Poschl, U.,
Després, V. R., & Frohlich-Nowoisky, J. (2018). Species Richness, rRNA Gene
Abundance, and Seasonal Dynamics of Airborne Plant-Pathogenic Oomycetes. *Frontiers*
in Microbiology, *9*(NOV). https://doi.org/10.3389/fmicb.2018.02673

Lefèvre, E., Roussel, B., Amblard, C., & Sime-Ngando, T. (2008). The molecular diversity of
freshwater picoeukaryotes reveals high occurrence of putative parasitoids in the plankton.
PLoS ONE, *3*(6). https://doi.org/10.1371/journal.pone.0002324

Lellis, W. A., White, B. S. J., Cole, J. C., Johnson, C. S., Devers, J. L., Gray, E. V. S., &
Galbraith, H. S. (2013). Newly documented host fishes for the eastern elliptio mussel
Elliptio complanata. *Journal of Fish and Wildlife Management*, *4*(1), 75–85.
https://doi.org/10.3996/102012-JFWM-094

Lepelletier, F., Karpov, S. A., Alacid, E., Le Panse, S., Bigeard, E., Garcés, E., Jeanthon, C., &
Guillou, L. (2014). *Dinomyces arenysensis gen. et sp. nov.* (Rhizophydiales,
Dinomycetaceae fam. nov.), a Chytrid Infecting Marine Dinoflagellates. *Protist*, *165*(2),
230–244. https://doi.org/10.1016/j.protis.2014.02.004

Martel, A., Spitzen-Van Der Sluijs, A., Blooi, M., Bert, W., Ducatelle, R., Fisher, M. C., Woeltjes,
A., Bosman, W., Chiers, K., Bossuyt, F., & Pasmans, F. (2013). *Batrachochytrium*
salamandrivorans sp. nov. causes lethal chytridiomycosis in amphibians. *Proceedings of*
the National Academy of Sciences of the United States of America, *110*(38), 15325–15329.
https://doi.org/10.1073/pnas.1307356110

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet Journal*, *17*(1), 10–12. http://www-huber.embl.de/users/an-

McMurdie, P. J., & Holmes, S. (2013). Phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. *PLoS ONE, 8*(4).
https://doi.org/10.1371/journal.pone.0061217

McTaggart, A. R., Shivas, R. G., Geering, A. D. W., Vánky, K., & Scharaschkin, T. (2012). A
review of the *Ustilago-Sporisorium-Macalpinomyces* complex. *Persoonia: Molecular*
Phylogeny and Evolution of Fungi, *29*, 55–62. https://doi.org/10.3767/003158512X660283

Mendoza, L. (2009). *Pythium insidiosum* and mammalian hosts. In G. W. Beakes & S. Sekimoto
(Eds.), *Oomycete genetics and genomics: diversity, interactions, and research tools*.

Mendoza, L., Hernandez, F., & Ajello3, L. (1993). Life Cycle of the Human and Animal
Oomycete Pathogen *Pythium insidiosum*. In *JOURNAL OF CLINCAL MICROBIOLOGY*
(Vol. 31, Issue 11). https://journals.asm.org/journal/jcm

Metz, S., Itoïz, S., Obiol, A., Derelle, E., Massana, R., Berney, C., de Vargas, C., Soudant, P.,
Monier, A., & Chambouvet, A. (2023). Global perspective of environmental distribution and

diversity of Perkinsea (Alveolata) explored by a meta-analysis of eDNA surveys. *Scientific*
Reports, *13*(1), 1–13. https://doi.org/10.1038/s41598-023-47378-0

Mouritsen, K. N., Sørensen, M. M., Poulin, R., & Fredensborg, B. L. (2018). Coastal ecosystems
on a tipping point: Global warming and parasitism combine to alter community structure
and function. *Global Change Biology*, *24*(9), 4340–4356. https://doi.org/10.1111/gcb.14312

Naselli-Flores, L., & Padisák, J. (2023). Ecosystem services provided by marine and freshwater
phytoplankton. *Hydrobiologia*, *850*(12–13), 2691–2706. https://doi.org/10.1007/s10750-
022-04795-y

O'Reilly, C. M., Sharma, S., Gray, D. K., Hampton, S. E., Read, J. S., Rowley, R. J., Schneider,
P., Lenters, J. D., McIntyre, P. B., Kraemer, B. M., Weyhenmeyer, G. A., Straile, D., Dong,
B., Adrian, R., Allan, M. G., Anneville, O., Arvola, L., Austin, J., Bailey, J. L., … Zhang, G.
(2015). Rapid and highly variable warming of lake surface waters around the globe.
Geophysical Research Letters, *42*(24), 10773–10781.
https://doi.org/10.1002/2015GL066235

Palomar, G., Jakobik, J., Bosch, J., Kolenda, K., Kaczmarski, M., Josko, P., Roces-Diaz, J. V.,
Stachyra, P., Thumsova, B., Zielinski, P., & Pabijan, M. (2021). Emerging infectious
diseases of amphibians in Poland: distribution and environmental drivers. *Diseases of*
Aquatic Organisms, *147*, 1–12.

Pojmańska, T., Salamatin, R., Sulgostowska, T., Cielecka, D., Okulewicz, A., Niewiadomska, K.,
& Grytner-Zięcina, B. (2012). The Polish collection of parasitic helminths. *Annals of*
Parasitology, *58*(2), 75–86. http://www.helminths.eu.http://www.helminths.eu

Powell, M. J., & Letcher, P. M. (2019). Ultrastructure of early stages of *Rozella allomycis*
(Cryptomycota) infection of its host, *Allomyces macrogynus* (Blastocladiomycota). *Fungal*
Biology, *123*(2), 109–116. https://doi.org/10.1016/j.funbio.2018.11.009

Powell, M. J., Letcher, P. M., Davis, W. J., Lefèvre, E., Brooks, M., & Longcore, J. E. (2019).
Taxonomic summary of *Rhizoclosmatium* and description of four new *Rhizoclosmatium*
species (Chytriomycetaceae, Chytridiales). *Phytologia*, *101*(2), 139–163.
www.phytologia.org

Rad-Menéndez, C., Gerphagnon, M., Garvetto, A., Arce, P., Badis, Y., Sime-Ngando, T., &
Gachon, C. M. M. (2018). Rediscovering *Zygorhizidium affluens* Canter: Molecular
taxonomy, infectious cycle, and cryopreservation of a chytrid infecting the bloom-forming
diatom *Asterionella formosa*. *Applied and Environmental Microbiology*, *84*(23).
https://doi.org/10.1128/AEM.01826-18

Rasconi, S., Niquil, N., & Sime-Ngando, T. (2012). Phytoplankton chytridiomycosis: Community
structure and infectivity of fungal parasites in aquatic ecosystems. *Environmental*
Microbiology, *14*(8), 2151–2170. https://doi.org/10.1111/j.1462-2920.2011.02690.x

R Core Team. (2023). *R: A Language and Environment for Statistical Computing*.
https://www.R-project.org/

Rug, M., & Vogt, G. (1994). Developmental stages of the crayfish parasite *Psorospermium*
haeckeli in thoracic arteries of *Ascatus ascatus*. *Journal of Invertebrate Pathology*, *64*,
153–155.

Sassenhagen, I., Langenheder, S., & Lindström, E. S. (2023). Infection strategies of different
chytrids in a diatom spring bloom. *Freshwater Biology*, *68*(6), 972–986.
https://doi.org/10.1111/fwb.14079

Schulz, V.⬚;, Schulz, A.⬚;, Klamke, M.⬚;, Preissler, K.⬚;, Sabino-Pinto, J.⬚;, Müsken, M.⬚;,
Schlüpmann, M.⬚;, Heldt, L.⬚;, Kamprad, F.⬚;, Enss, J.⬚;, Schweinsberg, M.⬚;, Virgo,
J.⬚;, Rau, H.⬚;, Veith, M.⬚;, Lötters, S.⬚;, Wagner, N.⬚;, Steinfartz, S.⬚;, & Vences, M.
(2020). Batrachochytrium salamandrivorans *in the Ruhr district, Germany: History,*
distribution, decline dynamics and disease symptoms of the salamander plague.
http://www.salamandra-journal.com

Seto, K., Simmons, D. R., Quandt, C. A., Frenken, T., Dirks, A. C., Clemons, R. A., McKindles,
K. M., McKay, R. M. L., & James, T. Y. (2023). A combined microscopy and single-cell
sequencing approach reveals the ecology, morphology, and phylogeny of uncultured
lineages of zoosporic fungi. *MBio*, *14*(4), e0131323. https://doi.org/10.1128/mbio.01313-23

Shen, W., Le, S., Li, Y., & Hu, F. (2016). SeqKit: A cross-platform and ultrafast toolkit for
FASTA/Q file manipulation. *PLoS ONE*, *11*(10).
https://doi.org/10.1371/journal.pone.0163962

Shi, S., Wang, J., Liu, C., & Zheng, L. (2023). Developmental toxicity and inflammatory
response induced by Botrytis cinerea in zebrafish (*Danio rerio*) larvae. *Comparative*
Biochemistry and Physiology Part - C: Toxicology and Pharmacology, *267*.
https://doi.org/10.1016/j.cbpc.2023.109575

Simmons, D. R., Bonds, A. E., Castillo, B. T., Clemons, R. A., Glasco, A. D., Myers, J. M.,
Thapa, N., Letcher, P. M., Powell, M. J., Longcore, J. E., & James, T. Y. (2020). The
Collection of Zoosporic Eufungi at the University of Michigan (CZEUM): Introducing a new
repository of barcoded Chytridiomyceta and Blastocladiomycota cultures. *IMA Fungus*,
11(1). https://doi.org/10.1186/s43008-020-00041-z

Socha, D., & Hutorowicz, A. (2009). Changes in the quantitative relations of the phytoplankton
in heated lakes. *Archives of Polish Fisheries*, *17*(4), 239–251.
https://doi.org/10.2478/v10086-009-0017-9

Socha, D., & Zdanowski, B. (2001). Aquatic ecosystems in the vicinity of Konin. *Biblioteka*
Monitoringu OErodowisk, 75.

Stoeck, T., Bass, D., Nebel, M., Christen, R., Jones, M. D. M., Breiner, H. W., & Richards, T. A.
(2010). Multiple marker parallel tag environmental DNA sequencing reveals a highly
complex eukaryotic community in marine anoxic water. *Molecular Ecology*, *19*(SUPPL. 1),
21–31. https://doi.org/10.1111/j.1365-294X.2009.04480.x

Sures, B., Nachev, M., Schwelm, J., Grabner, D., & Selbach, C. (2023). Environmental
parasitology: stressor effects on aquatic parasites. *Trends in Parasitology*, *39*(6), 461–474.
https://doi.org/10.1016/j.pt.2023.03.005

Thines, M. (2018). Oomycetes. In *Current Biology* (Vol. 28, Issue 15, pp. R812–R813). Cell
Press. https://doi.org/10.1016/j.cub.2018.05.062

Thomé, P. C., Wolinska, J., Van Den Wyngaert, S., Reñé, A., Ilicic, D., Agha, R., Grossart, H.
P., Garcés, E., Monaghan, M. T., & Strassert, J. F. H. (2024). Phylogenomics including
new sequence data of phytoplankton-infecting chytrids reveals multiple independent
lifestyle transitions across the phylum. *Molecular Phylogenetics and Evolution*, *197*.
https://doi.org/10.1016/j.ympev.2024.108103

Urbańska, M., Andrzejewski, W., Riccardi, N., Gierszal, H., & Golski, J. (2018). The Invasive
Asian Clam *Corbicula fluminea* in Polish Rivers: The Importance of Thermal Discharge
from Power Plants for Its Spread. *Polish Journal of Ecology*, *66*(1), 70–75.
https://doi.org/10.3161/15052249PJE2018.66.1.008

Urbańska, M., Kirschenstein, M., Obolewski, K., & Ożgo, M. (2019). Silent invasion:
Sinanodonta woodiana successfully reproduces and possibly endangers native mussels in
the north of its invasive range in Europe. *International Review of Hydrobiology*, *104*(5–6),
127–136. https://doi.org/10.1002/iroh.201801971

Vallet, M. (2024). Chemical ecology of plankton parasitism in algae. In *Botanica Marina* (Vol. 67,
Issue 3, pp. 211–229). Walter de Gruyter GmbH. https://doi.org/10.1515/bot-2023-0062

Van den Wyngaert, S., Ganzert, L., Seto, K., Rojas-Jimenez, K., Agha, R., Berger, S. A.,
Woodhouse, J., Padisak, J., Wurzbacher, C., Kagami, M., & Grossart, H. P. (2022).
Seasonality of parasitic and saprotrophic zoosporic fungi: linking sequence data to
ecological traits. *ISME Journal*, *16*(9), 2242–2254. https://doi.org/10.1038/s41396-022-
01267-y

Van den Wyngaert, S., Rojas-Jimenez, K., Seto, K., Kagami, M., & Grossart, H. P. (2018).
Diversity and Hidden Host Specificity of Chytrids Infecting Colonial Volvocacean Algae.
Journal of Eukaryotic Microbiology, *65*(6), 870–881. https://doi.org/10.1111/jeu.12632

Van den Wyngaert, S., Seto, K., Rojas-Jimenez, K., Kagami, M., & Grossart, H. P. (2017). A
New Parasitic Chytrid, *Staurastromyces oculus* (Rhizophydiales, Staurastromycetaceae
fam. nov.), Infecting the Freshwater Desmid *Staurastrum sp. Protist*, *168*(4), 392–407.
https://doi.org/10.1016/j.protis.2017.05.001

Wang, F., Wang, K., Cai, L., Zhao, M., Kirk, P. M., Fan, G., Sun, Q., Li, B., Wang, S., Yu, Z.,
Han, D., Ma, J., Wu, L., & Yao, Y. (2023). Fungal names: a comprehensive nomenclatural
repository and knowledge base for fungal taxonomy. *Nucleic Acids Research*, *51*(D1),
D708–D716. https://doi.org/10.1093/nar/gkac926

Wang, S., Cao, M., Wang, B., Deng, R., Gao, Y., & Liu, P. (2019). Optimization of growth
requirements and scale-up cultivation of freshwater algae *Desmodesmus armatus* using
response surface methodology. *Aquaculture Research*, *50*(11), 3313–3325.
https://doi.org/10.1111/are.14290

Wells, K., & Clark, N. J. (2019). Host Specificity in Variable Environments. In *Trends in
Parasitology* (Vol. 35, Issue 6, pp. 452–465). Elsevier Ltd.
https://doi.org/10.1016/j.pt.2019.04.001

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*.

Williamson, B., Tudzynski, B., Tudzynski, P., & Van Kan, J. A. L. (2007). *Botrytis cinerea*: The
cause of grey mould disease. In *Molecular Plant Pathology* (Vol. 8, Issue 5, pp. 561–580).
https://doi.org/10.1111/j.1364-3703.2007.00417.x

Winter, D. J. (2017). An R package for the NCBI eUtils API. *PeerJ Preprints*.

Wiśniewski, K., Szarmach, D., & Poznańska-Kakareko, M. (2020). The role of abiotic and biotic
factors in interspecific competition of Polish crayfish-Comprehensive literature review. In
Oceanological and Hydrobiological Studies (Vol. 49, Issue 4, pp. 428–441). De Gruyter
Open Ltd. https://doi.org/10.1515/ohs-2020-0038

Wood, C. L., Zgliczynski, B. J., Haupt, A. J., Guerra, A. S., Micheli, F., & Sandin, S. A. (2018).
Human impacts decouple a fundamental ecological relationship—The positive association
between host diversity and parasite diversity. *Global Change Biology*, *24*(8), 3666–3679.
https://doi.org/10.1111/gcb.14159

Woolway, R. I., & Merchant, C. J. (2019). Worldwide alteration of lake mixing regimes in
response to climate change. *Nature Geoscience*, *12*(4), 271–276.
https://doi.org/10.1038/s41561-019-0322-x

Yap, T. A., Koo, M. S., Ambrose, R. F., Wake, D. B., & Vredenburg, V. T. (2015). Averting a
North American biodiversity crisis. *Science, 349*(6247), 481–482.
https://doi.org/10.1126/science.aab1052

Yuryshynets, V., & Krasutska, N. (2009). Records of the parasitic worm *Aspidogaster*
conchicola (Baer 1827) in the Chinese pond mussel *Sinanodonta woodiana* (Lea 1834) in
Poland and Ukraine. *Aquatic Invasions*, *4*(3), 491–494.
https://doi.org/10.3391/ai.2009.4.3.9

Zdanowski, B., Napiórkowska-Krzebietke, A., Stawecki, K., Świątecki, A., Babko, R., Bogacka-
Kapusta, E., Czarnecki, B., & Kapusta, A. (2020). *Polish River Basins and Lakes-Part I* (E.
Korzeniewska & M. Harnisz, Eds.). Springer. http://www.springer.com/series/698

Zhang, D. (2024). *Package "rsq". R package version 4.2-0.*

1068