Warming increases richness and shapes assemblages of eukaryotic parasitic plankton

Amruta Rajarajan*[1], Sławek Cerbin[2], Kingsly C. Beng[1, 3§, 4§, 5§], Michael T. Monaghan[1, 6], Justyna Wolinska[1, 6]

[1]Dept. of Evolutionary and Integrative Ecology, Leibniz Institute of Freshwater Ecology and Inland Fisheries (IGB), Müggelseedamm 301, 12587 Berlin, Germany

[2]Dept. of General Zoology, Faculty of Biology, Adam Mickiewicz University, 61-614 Poznań, Poland

[3]Focus Group Marine Molecular Ecology, Helmholtz Institute for Functional Marine Biodiversity, University of Oldenburg (HIFMB), 26129 Oldenburg, Germany

[4]Alfred Wegener Institute (AWI), Helmholtz Center for Polar and Marine Research, 27570 Bremerhaven, Germany

[5]Thünen Institute of Sea Fisheries, 38116 Braunschweig, Bremerhaven, Germany

[6]Institut für Biologie, Freie Universität Berlin, 14195 Berlin, Germany

*Corresponding author email: amruta.rajarajan@igb-berlin.de, amrutarajarajan@gmail.com

§Present address

Abstract

Anthropogenic activities have led to a global rise in water temperatures, leading to an increased interest in how warming affects infectious disease ecology. While most studies have focused on individual host-parasite systems, there is a gap in understanding the impact of warming on multi-host, multi-parasite assemblages in natural ecosystems. To address this gap, we examined freshwater eukaryotic parasite communities in ten natural lakes near Konin, Poland: five artificially heated and five nearby non-heated "control" lakes. Since 1958, the heated lakes have experienced a mean annual temperature increase of 2°C due to hot water discharge from two adjacent power plants. We collected seasonal environmental DNA (eDNA) samples from surface waters over a two-year period and applied targeted metabarcoding to compare the richness and distribution of eukaryotic parasites across lake types, with a focus on protists and fungi. Using literature searches and sequence metadata from GenBank, we identified putative parasites which included Alveolates, Stramenopiles, basal Fungi and Ichthyosporeans, as well as their associated hosts. Heated lakes exhibited distinct parasite assemblages, with higher richness of parasitic chytrids and aphelids particularly during summer, suggesting thermal preferences among some freshwater parasite species. A general linear model suggests that increased chytrid richness in heated lakes may be partly due to higher richness of their green algal, diatom- and dinoflagellate hosts. These findings highlight how warming can drive shifts in parasite richness and assemblages, potentially altering host-parasite dynamics in freshwater ecosystems.

Introduction

Warming in aquatic environments has led to an average increase in freshwater temperatures of 0.34°C per decade. Elevated temperatures lead to deoxygenation (Jane et al., 2021), changes in seasonality and reduced ice cover (Grant et al., 2021), shifts in lake productivity dynamics (Gilarranz et al., 2022), and alterations in mixing regimes (Woolway & Merchant, 2019). These changes have cascading effects on species distributions and food webs (Bartley et al., 2019). The magnitude of warming-driven changes on aquatic life is unprecedented, highlighting a critical need to understand their broader ecological implications (IPCC 2023).

While the effects of warming on plankton have been widely studied due to their central role in nutrient cycling (Bakhtiyar et al., 2020; Naselli-Flores & Padisák, 2023), the parasites of plankton remain understudied despite their significant ecological contributions. Chytrid fungi (Chytridiomycota) infect various phytoplankton, including green algae (reviewed in Frenken et al., 2017), diatoms (reviewed in Danz & Quandt, 2023), and dinoflagellates (Fernández-Valero et al., 2023, 2024). By producing motile zoospores, chytrids create additional trophic links (mycoloops) that sustain zooplankton populations (Abonyi et al., 2024; Kagami et al., 2014). Chytrid outbreaks can exert top-down control on toxic cyanobacterial (Gleason et al., 2015) and dinoflagellate blooms (Lepelletier et al., 2014). Other parasitic groups, such as Cryptomycota, hyperparasitize chytrids (James et al., 2013; Powell & Letcher, 2019), while Aphelidiomycota (aphelids) (Jephcott et al., 2017) and Oomycota (oomycetes) (Thines, 2018) infect various phytoplankton hosts. Perkinsea, a group previously considered to exclusively contain intracellular parasites of green algae, was recently discovered to be more diverse, with potential roles in various ecosystem processes beyond parasitism (Metz et al., 2023).

The influence of warming on infectious disease extends beyond parasite development to affect host abundance, physiology, and immunity (King et al., 2023). For instance, a four-month indoor mesocosm study found that 4°C elevation in spring temperature (to 21°C)

shortened the bloom duration of the diatom *Synedra* sp., resulting in an earlier end to the

epidemic of its chytrid parasite *Zygorhizidium planktonicum* (Frenken et al., 2016).

Conversely, an outdoor mesocosm study of amphipod host communities exposed to the

generalist trematode parasite *Maritrema novaezealandensis* under a 4°C summer temperature

increase (to 25°C) showed elevated infection prevalence, along with reduced host species

richness and abundance (Mouritsen et al., 2018). These findings demonstrate how warming

can impact disease dynamics in individual host-parasite systems within aquatic environments.

Parasite richness, integral to ecosystem health (Hudson et al., 2006) is influenced by host

distribution and richness in distinct ways. Using theoretical models, Braga et al., (2018)

demonstrated that habitat heterogeneity fosters host heterogeneity, which in turn increases

genotypic richness of parasites. Host richness is positively correlated with parasite richness,

according the host-diversity-begets-parasite-diversity hypothesis (Kamiya et al., 2014;

Krasnov & Poulin, 2015). For example, ponds with higher amphibian host richness harbored

higher richness of trematode parasites (Johnson et al., 2016). Changes in host and parasite

species richness can, in turn, affect disease dynamics in complex ways. In one case, High

amphibian host richness reduced infection prevalence of the trematode *Ribeiroia ondatrae*

(dilution effect) but amplified parasite competition (Johnson et al., 2013).

Despite these insights, our understanding of how global warming affects parasite richness and

the potential consequences for disease dynamics remains incomplete (Sures et al., 2023).

Warming-driven changes in species assemblages can lead to 'ecological fitting' due to novel

host-parasite encounters and the resulting increased likelihood of host-switching (Wells &

Clark, 2019). This could potentially increase phenotypic and genotypic diversity of parasites

(Agosta et al., 2010). Moreover, anthropogenic influence can have unexpected impacts on the

generally positive correlation between host and parasite richness in natural ecosystems. For

instance, islands that experienced extensive fishing exhibited higher fish host richness, but

lower helminth parasite richness, compared to unfished islands (Wood et al., 2018). In

contrast, warming-driven habitat loss of host species reduced helminth parasite richness

(Carlson et al., 2017). Long-term empirical studies in multi-host, multi-parasite systems are

essential for understanding warming effects on disease ecology.

One promising approach to study the effects of warming on parasite assemblages involves *in*

situ sampling of natural lakes. The Konin lake complex in Poland offers an ideal opportunity.

Since 1958, five of these lakes have been artificially warmed by hot water discharge from two

nearby power plants, resulting in altered abiotic conditions and changes in plankton

community composition compared to the five nearby non-heated (hereafter "control") lakes

(Zdanowski et al., 2020). The heated lakes are on average 2°C warmer (Beng et al., 2023),

mimicking the projected global freshwater temperature increases by 2100 (IPCC 2023).

Additionally, these lakes experience reduced winter ice cover (Socha & Zdanowski, 2001).

Warming has led to shifts in *Daphnia* species composition in these lakes, favoring the more

heat-tolerant *D. galeata* (Dziuba et al., 2020). Moreover, the heated lakes exhibit higher

richness and distinct community composition of various phytoplankton (green algae, protists,

fungi) and zooplankton (crustaceans, rotifers) taxa (Beng et al., 2023; Ejsmont-Karabin et al.,

2020). There is also an increased abundance of invasive species, including the tape grass

Vallisneria spiralis (Socha & Hutorowicz, 2009) and the mussels *Corbicula fluminea*

(Urbańska et al., 2018) and *Sinanodonta woodiana* (Urbańska et al., 2019). *Sinanodonta*

woodiana occurrence further favors the presence of its parasitic worm *Aspidogaster*

conchicola (Yuryshynets & Krasutska, 2009). Heated lakes also harbor introduced species of

fish (Kapusta & Bogacka-Kapusta, 2015). These changes in environmental conditions and

host community composition provide a valuable opportunity to investigate the effects of

warming on parasite diversity in natural settings. Although warming reduces *Caullerya*

mesnili prevalence in *Daphnia* hosts (Dziuba et al., 2023), comprehensive investigations into

differences in parasite assemblages between heated and control lakes are still lacking.

This study leverages this unique set of artificially heated and control lakes in Poland to

explore ca. 60 years of warming effects on eukaryotic parasitic plankton. Using eDNA

metabarcoding of seasonally collected surface water samples (winter, spring, summer,

autumn) from five heated and five control lakes, and addressed the questions: (i) Which

parasites are present in the study sites? (ii) Do certain parasites exhibit a preference for habitat

types (heated or non-heated), leading to distinct assemblages in heated versus control lakes?

(iii) Does parasite richness differ between heated and control lakes? and (iv) Is parasite

richness correlated with host richness?

Material and methods

Generating sequence data

<u>eDNA sampling, molecular laboratory procedures and sequencing</u>

Detailed description of eDNA sampling dates, locations, and molecular laboratory procedures

are provided in Beng et al., (2023). Surface water samples were collected from five heated

lakes (Gosławskie, Licheńskie, Mikorzyńskie, Pątnowskie and Ślesińskie) and five control

lakes (Budzisławskie, Gopło, Skulskie, Skulska Wieś and Suszewskie) once per season

throughout 2020 and 2021 (except that no summer samples were obtained in 2021, while all

lakes were sampled twice in autumn 2020). For DNA amplification, two primer sets were

used: "EUK15" TAReuk454FWD1: 5'-CCAGCASCYGCGGTAATTCC-3'; TAReukREV3:

5'-ACTTTCGTTCTTGATYRA-3'; both primers from Stoeck et al., (2010), targeting a

fragment of the 18S rRNA gene of eukaryotes, and "CHY" ITS4ngsF: 5'-

GCATATCAATAAGCGSAGGA-3' (Van den Wyngaert et al., 2022); LF402R: 5'-

TTCMCTTTNMRCAATTTCAC-3' (Thomé et al., 2024), targeting a fragment of the 28S

rRNA gene of fungi. For the "EUK15" primer set, 10 ng template DNA was used in 25-µL

PCR reactions with the following conditions: an initial denaturation at 95°C for 30 seconds,

followed by 30 cycles (95°C for 30 s, 45°C at 30 s, 72°C at 30 s), with a final extension of

72°C for 5 minutes. The PCR products were normalized to 5 ng/µL and used as templates for

the second PCR, where Illumina Nextera indexes were added to 25-µL PCRs under these

conditions: 95°C for 2 minutes, followed by 8 cycles (95°C for 20 s, 52°C for 30 s, 72°C for

30 s), and a final extension at 72°C for 3 minutes. PCR of the "CHY" primer set followed the

protocol by Velthuis et. al. (unpubl). Barcoded samples were pooled in equimolar

concentrations and paired-end sequenced on the Illumina MiSeq v3 (600 cycles) platform.

Each sequencing run included one no-template negative control.

Bioinformatics

Each library (EUK15 and CHY) was sequenced in two runs: the first run included samples

from January 2020 (winter 2020), May 2020 (spring 2020), July 2020 (summer 2020) and

October 2020 (autumn 2020), while the second run included samples from November 2020

(autumn 2020), December 2020 (winter 2021), April 2021 (spring 2021) and October 2021

(autumn 2021). Bioinformatic processing followed the protocol outlined in Beng et al.,

(2023). The first sequencing run yielded 2.4 million raw reads for EUK15 and 1.9 million raw

reads for CHY (Beng et al., 2023), while the second run yielded 643,742 raw reads for

EUK15 and 741,018 raw reads for CHY. Primer sequences were removed with cutadapt

(Martin, 2011) and reads were processed using the package DADA2 v 1.28 (Callahan et al.,

2016) in R v 4.3.0 (R Core Team, 2023) (see supp R script). Quality filtering and *de novo*

chimera removal retained 59% of EUK15 and 58.3% of CHY reads. Reads from the two runs

were combined into a "protist dataset" (EUK15) and a "fungal dataset" (CHY). Amplicon

Sequence Variants (ASVs) were identified for each dataset using the *dada* function, and

ASV-count tables were prepared. Taxonomic assignments of ASVs in the protist dataset were

conducted using the Protist Ribosomal database PR2 v.5.0.0 (Guillou et al., 2013). For the

fungal dataset, taxonomic assignment was performed using BLAST+ with a custom database

(see below).

Despite the relatively lower sequencing depth in the second runs, all rarefaction curves

plateaued (Fig. S1), confirming sufficient sequencing depth. Thus, all samples were retained

for further analyses. Contaminant ASVs, defined as those present in negative controls, were

removed using the phyloseq package in R (McMurdie & Holmes, 2013). Specifically, 17

contaminant ASVs were removed from the first run of the protist dataset, while 3 and 2

contaminant ASVs were removed from the first and second runs of the fungal dataset,

respectively, comprising <0.01% reads in each dataset. In total, bioinformatic processing

resulted in 3297 ASVs in the protist dataset and 4243 ASVs in the fungal dataset.

Classification of ASVs as parasites

ASVs were classified as parasites using a modified approach from Beng et al. (2021) (detailed

in Fig S2). Two custom parasite sequence databases were created for this purpose: one

containing sequences from the NCBI BLAST nucleotide database representing 10 higher taxa

discovered in the protist dataset (Fig. S3), and the other containing NCBI sequences of 9 taxa

discovered in the fungal dataset (Fig. S4). Additionally, we incorporated (i) ribosomal

sequences from NCBI (18S, 28S, and ITS), (ii) sequences from parasite groups reported to

occur in the Konin lake complex (Cestoda, Nematoda, Monogenea, Digenea, Trematoda and

Acanthocephala), which infect fish (Pojmańska et al., 2012), mussels (Yuryshynets &

Krasutska, 2009), and snails (Cichy et al., 2020), and (iii) sequences of eukaryotic taxonomic

groups known to contain parasite species, identified through Web of Science literature

searches performed for each unique phylum listed in the Fungal Names database (F. Wang et

al., 2023), using the terms "[Phylum name] AND [parasit* OR pathogen*]" (Table S1).

Duplicate sequences were removed using seqkit (Shen et al., 2016). Each ASV was then

identified to species- or higher taxonomic level using its respective custom database and

BLAST+ on a high-performance computing cluster (Bennett et al., 2020). Only the top ten

hits with ≥90% sequence similarity and 100% query cover were processed further. For each

ASV, "host" entries within GenBank metadata of the top-bitscore BLAST hit (Beng et al.,

2021) were extracted using the rentrez package (Winter, 2017).

Species names from the protist and fungal datasets (for ASVs that were assigned to species

level) were used for literature searches on Web of Science with the search terms "[Species

name] AND [parasit* OR pathogen* OR infect*]". Relevant articles were checked for

evidence of parasitism and host association. BLAST hits of ASVs with no associated host

entries in GenBank or host information from literature searches were classified as non-

parasitic (Fig. S2).

Host entries obtained from GenBank were manually categorized into host groups (see Fig. 1).

Only ASVs associated with living hosts were classified as parasites. ASVs associated with

environmental samples (e.g., water, soil, mud sediment) or non-living substrates (e.g.,

"decaying log", "fallen branch") were excluded. ASVs representing symbionts (e.g., algal

endosymbionts of protists, lichen-associated ASVs, dinoflagellate symbionts of sponges) or

organisms within the pathobiome of diseased hosts were also excluded. This rigorous

classification ensured that only ASVs with confirmed parasitic lifestyles were included in the

final analysis, enabling an accurate assessment of parasite richness and distribution across the

studied lakes (Fig. S2).

Statistical analyses

<u>ASVs indicative of heated or control lakes</u>

To assess whether certain parasite ASVs were indicative of either heated or control lakes, we

used the *signassoc* function (9999 permutations) from the indicspecies package in R. This

analysis was performed using presence-absence data for parasite ASVs (mode=0). Multiple

testing correction was implemented using the Sidak method within the indicspecies package

(De Caceres & Legendre, 2009).

ASV richness differences in dominant parasite groups between heated and control lakes

Parasite groups were scrutinized to identify those associated with aquatic hosts (see Fig. 1).

The four parasite groups with the highest ASV count were selected for comparisons across

lake types, seasons and years. For example, in the protist dataset, 66 ASVs were identified as

Gyrista (Stramenopiles phylum), but only 31 Gyrista ASVs belonged to Peronosporomycetes,

after excluding terrestrial hosts. Consequently, only Peronosporomycetes (=Oomycetes,

Subdivision: Gyrista), representing 7.9% of ASVs in the protist parasite dataset, were

included in ASV richness analyses. Additionally, 216 ASVs in the protist dataset were

identified as fungi but only fungi from the fungal dataset were considered for further analyses.

In the fungal dataset, 297 and 86 ASVs belonged to Ascomycota and Basidiomycota,

respectively, but 86.8% and 87.2% of these ASVs were associated with terrestrial hosts (such

as metazoans, plants and humans; see Fig. 1B) and were therefore excluded. Chytrids

(Phylum: Chytridiomycota), aphelids (Phylum: Aphelidiomycota) and Cryptomycota

infecting aquatic hosts were analyzed, collectively representing 50.3% of ASVs in the fungal

parasite dataset.

The ASV richness for each selected group was log-transformed and analyzed using

Generalized Linear Mixed Model (GLMM). The most parsimonious model with the best fit

was used to examine parasite richness across lake types, seasons and years: parasite ASV

richness ~ lake type * season + year + (1 | lake) + (1 | sampling event), family = poisson,

using the *glmer* function of the lme4 package (Bates et al., 2024) in R. The significance of

fixed effects was assessed with a Type III ANOVA using the *Anova* function of the car

package (Fox et al., 2019). For chytrid parasite richness, a separate model was employed to

investigate its correlation with host richness (see below).

Correlation of chytrid parasite richness and host richness

A GLMM was used to test whether differences in chytrid parasite richness correlated with

differences in host richness across heated and control lakes (Beng et al., 2023). Here, "chytrid

parasite richness" was defined as the count of chytrid parasite ASVs in the fungal dataset,

while "host richness" referred to the count of ASVs from the Subdivisions Chlorophyta,

Charophyta, Chromista, Gyrista, and Dinoflagellata in the protist dataset, as 88% of chytrid

parasite ASVs were associated with one or more of these groups (see Fig. 1B). Both host and

parasite richness values were log-transformed. We then used the following GLMM structure:

chytrid parasite richness ~ lake type * season + year + host ASV richness + (1 | lake) + (1 |

sampling event), family = poisson. Fixed effects were evaluated with a Type III ANOVA.

After observing significant effects of host richness, lake type, season and lake type × season

interaction on parasite richness (see Results), we calculated confidence intervals and

visualized a general linear model for host and parasite richness, separately for each lake type

(see Fig. 4). The model was plotted using the "geom_smooth" argument (method="glm",

se=TRUE) from the ggplot2 package (Wickham, 2016) in R.

Results

Characterization of parasite assemblages and hosts in the protist and fungal datasets

In the protist dataset, 390 ASVs (12% of the dataset) were classified as parasites. These

included: 43 ASVs confirmed as parasitic based on both literature evidence and GenBank

host entries, 55 ASVs classified as parasites using literature evidence alone, and 292 ASVs

classified based solely on GenBank host entries (Table S2). A total of 151 parasite ASVs

(39%) were identified to the species level, representing 54 species from 23 genera (Table S2),

which were further grouped into 14 higher taxa (Fig. 1A). The highest ASV richness was

observed among Fungi, Gyrista, and Ichthyosporea (Fig 1A). Most parasite ASVs (66%) were

associated with hosts such as green algal, diatom, or terrestrial-plant hosts (Fig. 1A, Table

S2).

In the fungal dataset, 683 ASVs (16% of the dataset) were identified as parasites: 198 ASVs

were classified as parasitic using both literature evidence and GenBank host entries, 298

ASVs based on GenBank host entries alone, and 187 ASVs using literature searches of

species names. A total of 496 ASVs (73% of parasite ASVs) showed >90% sequence

similarity and 100% query cover with known species in GenBank (Table S3). Based on these

criteria, the fungal dataset contained 268 putative species across 218 genera, classified into 15

higher taxa. The highest ASV richness was observed among Ascomycota, Chytridiomycota

and Cryptomycota (Fig. 1B). Similar to the protist dataset, a notable proportion of parasite

ASVs (45%) in the fungal dataset were linked to hosts such as green algae, diatoms, or

terrestrial plants (Fig. 1B, Table S3).

Figure 1. Parasite ASV count in the (A) protist and (B) fungal datasets. Colors represent broad host

taxa to which identified hosts of the parasite ASVs belong (see Methods and Fig. S2 for details). Note

that Fungal ASVs from the protist dataset were not considered for statistical analyses.

ASVs indicative of lake type

A total of 39 parasite ASVs were indicators of lake type: 5 ASVs were specific to control lakes, and 34 ASVs were specific to heated lakes (Fig. 2). Among these were 19 chytrids, 8 aphelids, 4 Cryptomycota, 7 protists from diverse lineages, and 1 metazoan (Table S4). Control lakes indicators included chytrids ASVs (263 and 34) from unclassified members of the Order Rhizophydiales. Heated-lake indicators comprised various taxa, including chytrid species *Dinomyces arenysensis*, *Dangeardia mamillata*, *Rhizophydium chlorogonii*, *Quaeritorhiza haematococci* and an unknown Rhizophydiales parasite (ASV 210). Other heated-lake indicators were *Elliptio complanata* (Unionidae mussel), *Ustilago esculenta* (Basidiomycete fungus) along with unclassified ASVs from Dinophyceae (ASV 110) and Peronosporales fungi (ASV 286). Additionally, *Aphelidium desmodesmi* and four unclassified ASVs from the Family Aphelidiaceae (ASVs 60, 220, 194 and 101) showed a preference for heated lakes.

Figure 2. Relative abundances of ASVs indicative of control (left panel) and heated (right panel)

lakes. Of the 39 indicator ASVs identified (Table S4), 29 are presented here, as they were present in

more than 20% of samples. ASV numbers are specified in brackets. For ASVs in the protist dataset

(marked with an asterisk), the lowest taxonomic classification assigned using the PR2 database is

reported, with values in the blue "%" column indicating sequence similarity to the BLAST hit used to

obtain host information (host entry from GenBank metadata). For ASVs from the fungal dataset, the

lowest known classification from the BLAST hit is reported, with the blue "%" values indicating

sequence similarity between the ASV and the BLAST hit. The classification of *E. complanata* as a

parasite was based on a literature search (Table S2).

Parasite ASV richness

Among the four parasite groups with the highest richness - chytrids, aphelids, Peronosporomycetes (Oomycetes), and Cryptomycota - chytrids and aphelids had significantly higher ASV richness in heated lakes (Fig. 3, Table 1), with a significant interaction between lake type and season. Chytrid ASV richness peaked during summer and autumn (Fig. 3A), while aphelid richness was lowest in spring (Fig. 3B). Peronosporomycete ASV richness varied seasonally, with higher values in spring and a peak in summer (Fig. 3C). Both Peronosporomycetes and Cryptomycota exhibited lower ASV richness in 2021 compared to 2020 (Fig. 3C, D, Table 1).

317 **Table 1.** Results of a type III ANOVA of GLMM investigating ASV richness for the four most ASV-

318 diverse parasite groups across lake type, season and year. Lake and sampling event were specified as

319 random effects. The chytrid parasite richness model uniquely incorporated host richness (Chlorophyta,

320 Charophyta, Chromista, Gyrista, and Dinoflagellata) as a predictor. Significant results (*$p<0.05$) are

321 marked with an asterisk.

Parasite group	Fixed effect	χ^2	*p*-value
Chytrids	Lake type	53.6	<0.001*
	Season	5.2	0.155
	Year	2.3	0.133
	Host richness	7.5	0.006*
	Lake type × season	23.2	<0.001*
Aphelids	Lake type	12.37	<0.001*
	Season	7.04	0.071
	Year	2.78	0.095
	Lake type × season	19.89	<0.001*
Peronosporomycetes	Lake type	0.00	1
(=Oomycetes)	Season	8.29	0.04*
	Year	20.05	<0.001*
	Lake type × season	1.33	0.722
Cryptomycota	Lake type	1.64	0.199
	Season	4.47	0.215
	Year	7.43	0.006*
	Lake type × season	1.22	0.748

Figure 3. ASV richness of the four most ASV-rich parasite groups: (A) chytrids, (B) aphelids, (C)

Peronosporomycetes (=Oomycetes) and (D) Cryptomycota. Colors indicate lake type and symbols

indicate the season of sample collection. Filled symbols denote individual data points, whereas larger

open symbols indicate seasonal mean values. Note that the y-axis scales differ across panels to

optimize visualization.

Correlation between host and parasite richness

Chytrid parasite richness was positively correlated with host richness (Table 1, Fig. 4). Heated

lakes harbored particularly high host richness during summer (Fig. 4), which aligned with the

significantly elevated chytrid richness observed in summer (Fig. 3A). When examining the

correlation between parasite and host richness separately within each lake type, a deviation from

the linear model was found in during autumn: chytrid parasite richness was exceptionally high in

heated lakes, despite comparable host richness between heated and control lakes. This

discrepancy aligned with the significant interaction between lake type and season.

Figure 4. Correlation between chytrid parasite ASV richness (from the fungal dataset) and host ASV

richness across Chlorophyta, Charophyta, Chromista, Gyrista, and Dinoflagellata (from the protist

dataset). Shaded areas represent 95% confidence intervals. Colors differentiate between lake types, and

symbols indicate the season of sample collection.

Discussion

Heated lakes received hot water discharge since 1958 and are on average 2°C warmer, allowing us to investigate the long-term effects of warming on parasite assemblages. We identified parasites from seasonally sampled eDNA to assess differences in parasite species richness and assemblages between five artificially heated and five non-heated control lakes over two years, using parasite-specific sequence databases, their metadata and literature. Our rigorous approach of classifying ASVs as parasites identified a diverse array of confirmed aquatic parasites, spanning major eukaryotic taxa Alveolates (Apicomplexa, Ciliophora, Dinoflagellata), Stramenopiles (Bigyra, Oomycetes), basal fungi (Chytrids, Rozellomycota) and Ichthyosporeans, highlighting the undocumented parasite diversity in natural lakes. We found that certain parasite species exhibit a strong preference for either heated or control habitats, indicating that long-term warming can shape assemblages of parasite species. Chytrid and aphelid parasite richness was notably higher in heated lakes, supporting the idea that warming can enhance genotypic richness of parasites (Agosta et al., 2010). However, this trend was not observed for Peronosporomycetes (=Oomycetes) or Cryptomycota, which showed no difference in parasite richness between heated and control lakes. As predicted by the host-diversity-begets-parasite-diversity hypothesis (Kamiya et al., 2014; Krasnov & Poulin, 2015), chytrid parasite richness showed positively correlated with host richness, suggesting that increased chytrid parasite richness in heated lakes may result directly from favourable abiotic conditions. Overall, these findings support the idea that thermal conditions strongly influence the composition and dynamics of parasite communities, possibly through a combination of direct effects on parasites and indirect effects via hosts.

Some parasite ASVs were >95% similar to known chytrid parasites, such as *Zygorhizidium affluens* (ASVs 31 and 717), the chitin-degrading *Rhizoclosmatium pessaminum* (ASV 1901) (Powell et al., 2019), *Endochytrium ramosum* (ASVs 1418 and 2290), and *Staurastromyces*

oculus (ASVs 1236, 1328 and 3041). These chytrids parasitize diatoms *Asterionella* (Rad-

Menéndez et al., 2018), and green algae, including *Chara*, *Cladophora* (Simmons et al., 2020),

and *Straurastrum* (Van den Wyngaert et al., 2017, see Table S3). Additionally, two ASVs (181

and 1195; see Table S2) showed >98% sequence similarity with *Psorospermium haeckeli* (Class:

Ichythyophonida), a parasite of the crayfish *Ascatus ascatus* (Rug & Vogt, 1994)*, a host species

that is impacted by the devastating Europe-wide crayfish plague (Wiśniewski et al., 2020). We

also identified four ASVs (1156, 2064, 1231, and 1305, see Table S2) of the Apicomplexan

parasite *Nematopsis temporariae*, which infects a variety of amphibian hosts (Chambouvet et al.,

2016). Microeukaryotic parasites can impact ecosystem function. For example, zoospores of *Z.*

affluens, released from parasitized diatom blooms, provide a nutritious food source for *Daphnia*,

thereby establishing a trophic link between producers and primary consumers (Kagami et al.,

2007).

Three parasite ASVs showed 90-93% sequence similarity to *Batrachochytrium*

salamandrivorans (*Bsal*), a pathogen responsible for global chytridiomycosis outbreaks in

amphibians (Balàž et al., 2017; Laking et al., 2017; Yap et al., 2015). *Bsal* produces motile,

water-borne transmission stages (zoospores) that are active within a thermal range of 10-15°C

but capable of surviving at temperatures as low as 5° (Martel et al., 2013). Consequently, *Bsal*

epidemics are generally observed during colder months, such as late autumn, winter or spring

(Schulz et al., 2020). Geospatial models predict a reduction in climatically suitable habitats for

Bsal under global warming scenarios (Grisnik et al., 2023). However, recent chytridiomycosis

outbreaks at water temperatures as high as 26.4°C (Laking et al., 2017) suggest a broader

thermal niche than previously anticipated. Although *Bsal* infections have so far not been detected

in Poland, its sister species *B. dendrobatidis* is widespread (Palomar et al., 2021). Notably, ASV

764 - closely related to *Bsal* - was predominantly detected in winter within heated lakes

Gosławskie, Licheńskie, Mikorzyńskie, and Pątnowskie (see Table S4). This suggests that warm-

winter conditions may provide a favourable environment for this parasite. These findings

underscore the need for additional research, particularly regarding host demographics (Bielby et

al., 2021), to assess the potential presence and implications of *Bsal* in the heated environments of

the Konin lakes.

A total of 39 parasite ASVs exhibited a strong preference for specific lake types, indicating

distinct parasite assemblages in heated versus non-heated habitats. Of these, 34 were more

prevalent in heated lakes, including *Aphelidium desmodesmi* and four other ASVs belonging to

Aphelidiaceae. Although the thermal preference of aphelid parasites remains underexplored, *A.*

desmodesmi infections in the green alga *Scenedesmus* were previously shown to increase under

elevated summer temperatures (22-25°C) and phosphorus-limiting conditions in a mesocosm

study (Alam et al., 2024). Consistent with this, green algal and diatom hosts of *A. desmodesmi*,

including *Desmodesmus armatus* (=*Scenedesmus armatus*, ASV 27), have shown a preference

for heated lakes (Beng et al., 2023). *D. armatus* typically shows optimum growth at high

temperatures (~27°C) (S. Wang et al., 2019), matching the summer water temperatures of the

heated Konin lakes. This suggests that warming may increase aphelid parasite richness both by

directly facilitating epidemics at elevated temperatures and indirectly by promoting host species

growth. The thermal preferences of other indicator parasites, however, remains unknown.

Additionally, one ASV (1558) that was more prevalent in heated lakes showed 92% similarity to

a Cryptomycota strain that infects rotifer hosts (Seto et al., 2023, see Table S4). Similarly, the

parasitic dinoflagellate *Dissodinium pseudolunula* (ASV 110), which infects copepod eggs, also

showed a preference for heated lakes, suggesting that warming may favor this parasite as well.

Conversely, two unidentified chytrid ASVs (34 and 263; Order: Rhizophydiales) showed a

preference for non-heated lakes and were associated with the diatom host *Ulnaria* (=*Synedra*),

which also showed a preference for control lakes (Beng et al., 2023). These findings suggest that

long-term warming has led to the development of distinct parasite assemblages, likely influenced

by host distribution patterns, despite the potential flow of water between heated and non-heated

lakes (Zdanowski et al., 2020).

Additionally, 27% of the ASVs from the protist dataset and 41% from the fungal dataset detected

in the Konin lakes were unexpectedly associated with terrestrial hosts, including plants, animals

and humans (See Fig. 1). For example, ASV 286 (Family: Peronosporales) showed a strong

preference for heated lakes (see Fig. 2) and exhibited 96.4% sequence similarity with *Pythium*

insidiosum, a generalist oomycete parasite known to cause devastating root rot in commercial

crops (Golińska & Świecimska, 2020), as well as disease in mammals (Mendoza, 2009) and

humans (Mendoza et al., 1993). *Pythium* releases motile zoospores from infected plant roots or

skin lesions (Mendoza et al., 1993). The close similarity of ASV 286 to a clinical sample from a

patient with pythiosis (GenBank BBXB02000196) suggests that heated lakes might function as a

reservoir, providing suitable habitats or alternative hosts for this parasite (Krajaejun et al., 2014).

Another fungal ASV (259) did not show a preference for lake type but shared 100% sequence

similarity with *Botrytis cineria* (Phylum: Ascomycota, GenBank AY544651), a well-known

airborne plant pathogen that causes grey mold in terrestrial plants (Williamson et al., 2007).

However, *B. cineria* has also been shown to cause severe developmental abnormalities and

fitness loss in aquatic organisms like zebrafish *Danio rerio* (Shi et al., 2023, also see Table S3).

This suggests that some fungal parasites, although typically associated with terrestrial plants,

may disperse to aquatic habitats and potentially impact aquatic species.

The richness of chytrid parasites in our study showed a positive correlation with host richness,

despite known variation in the extent of their host-specificity. The host-diversity-begets-parasite-

diversity hypothesis is proposed to be driven by host-specificity of parasites (Krasnov & Poulin,

2015). While some chytrid parasites are highly host-specific (Van den Wyngaert et al., 2018),

other evidence suggests that the specificity might be flexible, particularly during the initial

"attraction" phase when motile zoospores find and attach to potential host cells. The chemical

cues that attract motile zoospores, such as dimethylsulfoniopropionate (DMSP), are derived from

algal cells and are non-specific, suggesting that the attraction might not be limited to particular

algal species (Frenken et al., 2017; Vallet, 2024). Additionally, genotype specificity of some

chytrids may shift across thermal regimes, as seen in *Zygorhizidium planktonicum* infecting the

diatom *A. formosa* (Gsell et al., 2013). Nevertheless, the positive correlation between host and

parasite richness in our study suggests some degree of specificity between chytrid parasites and

their compatible host species (or genotypes). Increased host richness may also support a greater

richness of generalist parasites. Chytrids that exhibit a 'generalist' versus 'specialist' infection

strategy often occur at different times of the year (Kagami et al., 2021; Van den Wyngaert et al.,

2022). For example, the seasonal abundance of chytrid parasites often corresponds with

fluctuations in their specific phytoplankton host taxa (e.g. diatoms, cyanobacteria) (Rasconi et

al., 2012), suggesting that parasites may "track" the most abundant host species. During spring

diatom blooms, chytrids tend to specialize on the dominant diatom species, while an even host

community towards the end of the bloom supports the persistence of generalist chytrids

(Sassenhagen et al., 2023). Thus, for chytrid parasites, increased host richness may support

higher parasite richness regardless of whether the parasites are specialist or generalist, potentially

at different times of the year, contributing to their overall increased richness in heated lakes.

A deviation from the expected host-diversity-begets-parasite-diversity trend occurred in heated

lakes during autumn. The richness of chytrid parasites in heated lakes exceeded expectations

based on host richness alone during autumn. One potential explanation for this finding is the

significantly higher soluble phosphorus content observed in heated lakes during this season

(Beng et al., 2023). Phosphate levels can affect both hosts and parasites. Elevated phosphorus is

linked to increased infection prevalence of some chytrids, such as *Rhizosiphon crassum* infecting

its cyanobacterial host *Anabaena flosaquae* (Rasconi et al., 2012). Conversely, low phosphorus

levels can reduce the growth rate of diatom hosts such as *A. formosa* without impacting the

growth rate of *Rhizophydium* parasites, leading to epidemics at lower host densities (Bruning,

1991). Heated lakes harbor cyanobacterial blooms in the autumn, which could grow to high

densities due to elevated phosphorus and serve as alternative hosts – albeit none of the chytrid

parasites in our study associated with cyanobacterial hosts (see Fig. 1B). Thus, the increase in

phosphorus levels in heated lakes during autumn (Beng et al., 2023) could underlie the

unexpectedly high chytrid parasite richness in autumn.

The unexpectedly high chytrid richness in autumn may also be related to other ecosystem-wide

processes driven by warming. Warming is hypothesized to induce shifts in the host range of

parasites (Wells & Clark, 2019), or favor generalist parasites due to their ability to infect

multiple hosts (Cizauskas et al., 2017). Our study offers insights into how warming-driven

changes in aquatic ecosystems may impact parasite presence and richness. First, four indicator

parasites - the mussel *Elliptio complanata,* fungus *Ustilago esculenta,* and chytrids *Dinomyces*

arenysensis, and *Dangeardia mamillata*, all known host generalists – were more prevalent in

heated lakes (see Fig. 2). *E. complanata* (ASV 462) is an ectoparasite (Gendron et al., 2019) that

can attach to at least 40 fish host species spanning 14 families (Lellis et al., 2013). *U. esculenta*

belongs to a hybridizing species complex of multicellular fungi that parasitize multiple hosts

within the *Poaceae* grass family (reviewed in McTaggart et al., 2012*).* The chytrid *Di.*

arenysensis infects at least four genera of dinoflagellates (Lepelletier et al., 2014), whereas *Da.*

mamillata infects green algae from at least three genera (Van den Wyngaert et al., 2018). Other

indicator taxa, including the aphelid *Aphelidium desmodesmi* and the chytrids *Quaeritorhiza*

haematococci and *Rhizophydium chlorogonii* (=*Aquamyces chlorogonii*), were also more

prevalent in heated lakes, but their host ranges remain unknown. Second, the higher parasite

richness and distinct species assemblages in heated lakes, combined with previously reported

increases in host richness in these environments (Beng et al., 2023), raises the possibility that

warming may foster novel host-parasite encounters. For instance, heated lakes harbor the

invasive tape grass *Vallisneria spiralis*, which peaks in biomass during autumn (Hutorowicz,

2006). *Vallisneria* macrophytes can sustain rich epibiont communities that include green algae,

diatoms, and dinoflagellates (Dunn et al., 2008). We speculate that this may generate 'hotspots'

for novel encounters between potential host species and motile zoospores of generalist parasites.

This may increase the likelihood of host shifts and could contribute to the unexpectedly high

chytrid parasite richness in heated lakes observed in autumn. It is also possible for a deviation

from the host-diversity-begets-parasite-diversity trend to be linked to parasites with large host

ranges. For example, the diversity of some microsporidian parasites shows no association with

that of their amphipod hosts, likely due to their ability to switch hosts (Prati et al., 2022). Further

research on host ranges and disease dynamics is needed to test these hypothesized consequences

of long-term warming (Agosta et al., 2010; Wells & Clark, 2019), particularly the potential for

warming to favour generalist parasites.

Conclusion

Overall, our study revealed a vast diversity of eukaryotic parasitic plankton species, with

warming shown to increase parasite richness for some groups such as chytrids and aphelids.

Parasites exhibited distinct habitat preferences, favouring either heated or non-heated lakes,

resulting in unique parasite species assemblages. These preferences might be linked to host

species distributions. For chytrid parasites, increased richness in heated lakes can be partially

attributed to greater host richness in these environments. The observed habitat preferences

suggest that warming can shape species assemblages by impacting both abiotic factors (such as

thermal conditions in heated lakes) and biotic factors (such as host richness), potentially

impacting ecological interactions within freshwater ecosystems.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

Raw sequence data for the first sequencing run are available in the European Nucleotide Database (ENA) under BioProject ID PRJEB54709 and those of the second run are available at NCBI Sequence Read Archive (SRA) under BioProject ID PRJNA1201567. Reproducible R code for bioinformatic processing is provided in supplementary R scripts.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was funded by a joint Beethoven Life-1 grant from the German Science Foundation (WO 1587/9-1 to JW) and the National Science Centre, Poland (2018/31/F/NZ8/01986 to SC).

Authors' contributions

JW and SC secured funding and designed the project with MTM. JW and SC collected eDNA samples. AR and KCB performed bioinformatic processing of samples. AR performed ecological analyses and wrote the manuscript with input from all co-authors.

Acknowledgement

We thank Elisabeth Funke for conducting metabarcoding library preparation, and Sarah Sparmann and Susan Mbedi at the Berlin Center for Genomics in Biodiversity Research (BeGenDiv: https://begendiv.de/) for sequencing support. We further thank the Disease Evolutionary Ecology, and Molecular Ecology & Genomics groups at IGB Berlin for constructive feedback, as well as Nedim Tüzün for stimulating discussions.

References

Abonyi, A., Fornberg, J., Rasconi, S., Ptacnik, R., Kainz, M. J., & Lafferty, K. D. (2024). The chytrid insurance hypothesis: integrating parasitic chytrids into a biodiversity–ecosystem functioning framework for phytoplankton–zooplankton population dynamics. *Oecologia*, *204*(2), 279–288. https://doi.org/10.1007/s00442-024-05519-w

Agosta, S. J., Janz, N., & Brooks, D. R. (2010). How specialists can be generalists: Resolving the and "parasite paradox" and implications for emerging infectious disease. *Zoologia*, *27*(2), 151–162. https://doi.org/10.1590/S1984-46702010000200001

Alam, M. M., Hodaei, M., Hartnett, E., Gincley, B., Khan, F., Kim, G. Y., Pinto, A. J., & Bradley, I. M. (2024). Community structure and function during periods of high performance and system upset in a full-scale mixed microalgal wastewater resource recovery facility. *Water Research*, *259*. https://doi.org/10.1016/j.watres.2024.121819

Bakhtiyar, Y., Arafat, M. Y., Andrabi, S., & Tak, H. I. (2020). Zooplankton: The Significant Ecosystem Service Provider in Aquatic Environment. In *Bioremediation and Biotechnology, Vol 3* (pp. 227–244). Springer International Publishing. https://doi.org/10.1007/978-3-030-46075-4_10

Balàž, V., Gortázar Schmidt, C., Murray, K., Carnesecchi, E., Garcia, A., Gervelmeyer, A., Martino, L., Munoz Guajardo, I., Verdonck, F., Zancanaro, G., & Fabris, C. (2017). Scientific and technical assistance concerning the survival, establishment and spread of *Batrachochytrium salamandrivorans (Bsal)* in the EU. *EFSA Journal*, *15*(2). https://doi.org/10.2903/j.efsa.2017.4739

Bartley, T. J., McCann, K. S., Bieg, C., Cazelles, K., Granados, M., Guzzo, M. M., MacDougall, A. S., Tunney, T. D., & McMeans, B. C. (2019). Food web rewiring in a changing world. In *Nature Ecology and Evolution* (Vol. 3, Issue 3, pp. 345–354). Nature Publishing Group. https://doi.org/10.1038/s41559-018-0772-3

Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R., Singmann, H., Scheipl, F., Grothendieck, G., Green, P., Fox, J., Bauer, A., Krivitsky, P., Tanaka, E., & Jagan, M. (2024). Linear Mixed-Effects Models using "Eigen" and S4 (package lme4). *CRAN*.

Beng, K. C., Cerbin, S., Monaghan, M. T., & Wolinska, J. (2023). Long-term effects of surface-water temperature increase on plankton communities in artificially heated lakes: insights from eDNA metabarcoding. *Authorea Preprints*. https://doi.org/10.22541/au.168639128.85081228/v1

Beng, K. C., Wolinska, J., Funke, E., Van Den Wyngaert, S., Gsell, A. S., & Monaghan, M. T. (2021). Temporal dynamics of freshwater planktonic parasites inferred using a DNA metabarcoding time-series. *Parasitology*, *148*(13), 1602–1611. https://doi.org/10.1017/S0031182021001293

Bennett, L., Melchers, B., & Proppe, B. (2020). *Curta: A General-purpose High-Performance Computer at ZEDAT, Freie Universität Berlin*. https://refubium.fu-berlin.de/handle/fub188/26993

Benny, G. L., Ho, H. M., Lazarus, K. L., & Smith, M. E. (2016). Five new species of the obligate mycoparasite *Syncephalis* (Zoopagales, Zoopagomycotina) from soil. *Mycologia*, *108*(6), 1114–1129. https://doi.org/10.3852/15-073

Bielby, J., Price, S. J., Monsalve-CarcaÑo, C., & Bosch, J. (2021). Host contribution to parasite persistence is consistent between parasites and over time, but varies spatially. *Ecological Applications*, *31*(3). https://doi.org/10.1002/eap.2256

Braga, M. P., Araujo, S. B. L., Agosta, S., Brooks, D., Hoberg, E., Nylin, S., Janz, N., & Boeger, W. A. (2018). Host use dynamics in a heterogeneous fitness landscape generates oscillations in host range and diversification. *Evolution*, *72*(9), 1773–1783. https://doi.org/10.1111/evo.13557

Bruning, K. (1991). Effects of phosphorus limitation on the epidemiology of a chytrid phytoplankton parasite. *Freshwater Biology*, *25*(3), 409–417. https://doi.org/10.1111/j.1365-2427.1991.tb01385.x

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*, *13*(7), 581–583. https://doi.org/10.1038/nmeth.3869

Calvin, K., Dasgupta, D., Krinner, G., Mukherji, A., Thorne, P. W., Trisos, C., Romero, J., Aldunce, P., Barrett, K., Blanco, G., Cheung, W. W. L., Connors, S., Denton, F., Diongue-Niang, A., Dodman, D., Garschagen, M., Geden, O., Hayward, B., Jones, C., … Ha, M. (2023). *IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland.* (P. Arias, M. Bustamante, I. Elgizouli, G. Flato, M. Howden, C. Méndez-Vallejo, J. J. Pereira, R. Pichs-Madruga, S. K. Rose, Y. Saheb, R. Sánchez Rodríguez, D. Ürge-Vorsatz, C. Xiao, N. Yassaa, J. Romero, J. Kim, E. F. Haites, Y. Jung, R. Stavins, … C. Péan, Eds.). https://doi.org/10.59327/IPCC/AR6-9789291691647

Carlson, C. J., Burgio, K. R., Dougherty, E. R., Phillips, A. J., Bueno, V. M., Clements, C. F., Castaldo, G., Dallas, T. A., Cizauskas, C. A., Cumming, G. S., Doña, J., Harris, N. C., Jovani, R., Mironov, S., Muellerklein, O. C., Proctor, H. C., & Getz, W. M. (2017). Parasite biodiversity faces extinction and redistribution in a changing climate. *Science Advances*, *3*. https://www.science.org

Chambouvet, A., Valigurová, A., Pinheiro, L. M., Richards, T. A., & Jirků, M. (2016). *Nematopsis temporariae* (Gregarinasina, Apicomplexa, Alveolata) is an intracellular infectious agent of tadpole livers. *Environmental Microbiology Reports*, *8*(5), 675–679. https://doi.org/10.1111/1758-2229.12421

Cichy, A., Marszewska, A., Ciapka, P., & Żbikowska, E. (2020). Different thermal conditions of lakes affect host–parasite systems: A case study of *Viviparus contectus* (Millet, 1813) and digenean trematodes. *Freshwater Biology*, *65*(3), 417–425. https://doi.org/10.1111/fwb.13434

Cizauskas, C. A., Carlson, C. J., Burgio, K. R., Clements, C. F., Dougherty, E. R., Harris, N. C., & Phillips, A. J. (2017). Parasite vulnerability to climate change: An evidence-based

functional trait approach. *Royal Society Open Science*, *4*(1).
https://doi.org/10.1098/rsos.160535

Danz, A., & Quandt, C. A. (2023). A review of the taxonomic diversity, host–parasite
interactions, and experimental research on chytrids that parasitize diatoms. In *Frontiers in Microbiology* (Vol. 14). Frontiers Media SA. https://doi.org/10.3389/fmicb.2023.1281648

De Caceres, M., & Legendre, P. (2009). Associations between species and groups of sites:
indices and statistical inference. *Ecology*, *90*(12), 3566–3574.

Dunn, A. E., Dobberfuhl, D. R., & Casamatta, D. A. (2008). A survey of algal epiphytes from
Vallisneria americana Michx. (Hydrocharitaceae) in the lower St. Johns River, Florida.
Southeastern Naturalist, *7*(2), 229–244. https://doi.org/10.1656/1528-
7092(2008)7[229:ASOAEF]2.0.CO;2

Dziuba, M. K., Herdegen-Radwan, M., Pluta, E., Wejnerowski, Ł., Szczuciński, W., & Cerbin,
S. (2020). Temperature increase altered *Daphnia* community structure in artificially heated
lakes: a potential scenario for a warmer future. *Scientific Reports*, *10*(1).
https://doi.org/10.1038/s41598-020-70294-6

Dziuba, M. K., Manzi, F., Cerbin, S., & Wolinska, J. (2023). Can climate warming save *Daphnia*
from parasites? Reduced parasite prevalence in *Daphnia* populations from artificially
heated lakes. *Limnology and Oceanography*, *68*(1), 181–191.
https://doi.org/10.1002/lno.12257

Ejsmont-Karabin, J., Hutorowicz, A., Kapusta, A., Stawecki, K., Tunowski, J., & Zdanowski, B.
(2020). Rotifers in heated Konin lakes-A review of long-term observations. *Water
(Switzerland)*, *12*(6). https://doi.org/10.3390/w12061660

Fernández-Valero, A. D., Karpov, S. A., Sampedro, N., Gordi, J., Timoneda, N., Garcés, E., &
Reñé, A. (2024). Newly identified diversity of Dinomycetaceae (Rhizophydiales,
Chytridiomycota), a family of fungal parasites of marine dinoflagellates. *European Journal
of Protistology*, *93*. https://doi.org/10.1016/j.ejop.2024.126053

Fernández-Valero, A. D., Reñé, A., Timoneda, N., Pou-Solà, N., Gordi, J., Sampedro, N., &
Garcés, E. (2023). The succession of epiphytic microalgae conditions fungal community
composition: how chytrids respond to blooms of dinoflagellates. *ISME Communications*,
3(1). https://doi.org/10.1038/s43705-023-00304-x

Fox, J., Weisberg, S., Price, B., Adler, D., Bates, D., Baud-Bovy, G., Bolker, B., Ellison, S.,
Firth, D., Friendly, M., Gorjanc, G., Graves, S., Heiberger, R., Krivitsky, P., Laboisierre,
R., Maechler, M., Monette, G., Murdoch, D., Nilsson, H., … Zeilais, A. (2019). car:
Companion to Applied Regression. *CRAN*.

Frenken, T., Alacid, E., Berger, S. A., Bourne, E. C., Gerphagnon, M., Grossart, H. P., Gsell, A.
S., Ibelings, B. W., Kagami, M., Küpper, F. C., Letcher, P. M., Loyau, A., Miki, T.,
Nejstgaard, J. C., Rasconi, S., Reñé, A., Rohrlack, T., Rojas-Jimenez, K., Schmeller, D. S.,
… Agha, R. (2017). Integrating chytrid fungal parasites into plankton ecology: research
gaps and needs. In *Environmental Microbiology* (Vol. 19, Issue 10, pp. 3802–3822).
Blackwell Publishing Ltd. https://doi.org/10.1111/1462-2920.13827

Frenken, T., Velthuis, M., de Senerpont Domis, L. N., Stephan, S., Aben, R., Kosten, S., van Donk, E., & Van de Waal, D. B. (2016). Warming accelerates termination of a phytoplankton spring bloom by fungal parasites. *Global Change Biology*, *22*(1), 299–309. https://doi.org/10.1111/gcb.13095

Gendron, A. D., Sanchez, D., Douville, M., & Houde, M. (2019). Stress-related gene transcription in fish exposed to parasitic larvae of two freshwater mussels with divergent infection strategies. *Diseases of Aquatic Organisms*, *132*(3), 191–202. https://doi.org/10.3354/dao03319

Gilarranz, L. J., Narwani, A., Odermatt, D., Siber, R., Dakos D Edited, V., & Hastings, A. (2022). *Regime shifts, trends, and variability of lake productivity at a global scale*. https://doi.org/10.1073/pnas

Gleason, F. H., Jephcott, T. G., Küpper, F. C., Gerphagnon, M., Sime-Ngando, T., Karpov, S. A., Guillou, L., & van Ogtrop, F. F. (2015). Potential roles for recently discovered chytrid parasites in the dynamics of harmful algal blooms. In *Fungal Biology Reviews* (Vol. 29, Issue 1, pp. 20–33). Elsevier Ltd. https://doi.org/10.1016/j.fbr.2015.03.002

Golińska, P., & Świecimska, M. (2020). *Pythium*: Diseases and Their Management. In M. Rai, K. A. Abd-Elsalam, & A. P. Ingle (Eds.), *Pythium*. CRC Press. https://doi.org/10.1201/9780429296406

Grant, L., Vanderkelen, I., Gudmundsson, L., Tan, Z., Perroud, M., Stepanenko, V. M., Debolskiy, A. V., Droppers, B., Janssen, A. B. G., Woolway, R. I., Choulga, M., Balsamo, G., Kirillin, G., Schewe, J., Zhao, F., del Valle, I. V., Golub, M., Pierson, D., Marcé, R., … Thiery, W. (2021). Attribution of global lake systems change to anthropogenic forcing. *Nature Geoscience*, *14*(11), 849–854. https://doi.org/10.1038/s41561-021-00833-x

Grisnik, M., Gray, M. J., Piovia-Scott, J., Carter, E. D., & Sutton, W. B. (2023). Incorporating caudate species susceptibilities and climate change into models of *Batrachochytrium salamandrivorans* risk in the United States of America. *Biological Conservation*, *284*. https://doi.org/10.1016/j.biocon.2023.110181

Gsell, A. S., de Senerpont Domis, L. N., van Donk, E., & Ibelings, B. W. (2013). Temperature Alters Host Genotype-Specific Susceptibility to Chytrid Infection. *PLoS ONE*, *8*(8). https://doi.org/10.1371/journal.pone.0071737

Guillou, L., Bachar, D., Audic, S., Bass, D., Berney, C., Bittner, L., Boutte, C., Burgaud, G., De Vargas, C., Decelle, J., Del Campo, J., Dolan, J. R., Dunthorn, M., Edvardsen, B., Holzmann, M., Kooistra, W. H. C. F., Lara, E., Le Bescot, N., Logares, R., … Christen, R. (2013). The Protist Ribosomal Reference database (PR2): A catalog of unicellular eukaryote Small Sub-Unit rRNA sequences with curated taxonomy. *Nucleic Acids Research*, *41*(D1). https://doi.org/10.1093/nar/gks1160

Hudson, P. J., Dobson, A. P., & Lafferty, K. D. (2006). Is a healthy ecosystem one that is rich in parasites? *Trends in Ecology and Evolution*, *21*(7), 381–385. https://doi.org/10.1016/j.tree.2006.04.007

Hutorowicz, A. (2006). *Vallisneria spiralis L.* (Hydrocharitaceae) in Lakes in the Vicinity of Konin (Kujawy Lakeland). *Biodiv. Res. Conserv.*, *2*(1), 154–158. www.brc.amu.edu.pl

James, T. Y., Pelin, A., Bonen, L., Ahrendt, S., Sain, D., Corradi, N., & Stajich, J. E. (2013). Shared signatures of parasitism and phylogenomics unite Cryptomycota and Microsporidia. *Current Biology*, *23*(16), 1548–1553. https://doi.org/10.1016/j.cub.2013.06.057

Jane, S. F., Hansen, G. J. A., Kraemer, B. M., Leavitt, P. R., Mincer, J. L., North, R. L., Pilla, R. M., Stetler, J. T., Williamson, C. E., Woolway, R. I., Arvola, L., Chandra, S., DeGasperi, C. L., Diemer, L., Dunalska, J., Erina, O., Flaim, G., Grossart, H. P., Hambright, K. D., … Rose, K. C. (2021). Widespread deoxygenation of temperate lakes. *Nature*, *594*(7861), 66–70. https://doi.org/10.1038/s41586-021-03550-y

Jephcott, T. G., van Ogtrop, F. F., Gleason, F. H., Macarthur, D. J., & Scholz, B. (2017). The ecology of chytrid and aphelid parasites of phytoplankton. In J. Dighton & J. F. White (Eds.), *The Fungal Community* (pp. 239–255). CRC Press. https://doi.org/10.1201/9781315119496

Johnson, P. T. J., Preston, D. L., Hoverman, J. T., & LaFonte, B. E. (2013). Host and parasite diversity jointly control disease risk in complex communities. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(42), 16916–16921. https://doi.org/10.1073/pnas.1310557110

Johnson, P. T. J., Wood, C. L., Joseph, M. B., Preston, D. L., Haas, S. E., & Springer, Y. P. (2016). Habitat heterogeneity drives the host-diversity-begets-parasite-diversity relationship: evidence from experimental and field studies. *Ecology Letters*, *19*(7), 752–761. https://doi.org/10.1111/ele.12609

Kagami, M., Miki, T., & Takimoto, G. (2014). Mycoloop: Chytrids in aquatic food webs. *Frontiers in Microbiology*, *5*(APR). https://doi.org/10.3389/fmicb.2014.00166

Kagami, M., Seto, K., Nozaki, D., Nakamura, T., Wakana, H., & Wurzbacher, C. (2021). Single dominant diatom can host diverse parasitic fungi with different degree of host specificity. *Limnology and Oceanography*, *66*(3), 667–677. https://doi.org/10.1002/lno.11631

Kagami, M., von Elert, E., Ibelings, B. W., de Bruin, A., & Van Donk, E. (2007). The parasitic chytrid *Zygorhizidium* facilitates the growth of the Cladoceran zooplankter, *Daphnia* in cultures of the inedible alga *Asterionella*. *Proceedings of the Royal Society B: Biological Sciences*, *274*(1617), 1561–1566. https://doi.org/10.1098/rspb.2007.0425

Kamiya, T., O'Dwyer, K., Nakagawa, S., & Poulin, R. (2014). Host diversity drives parasite diversity: Meta-analytical insights into patterns and causal mechanisms. *Ecography*, *37*(7), 689–697. https://doi.org/10.1111/j.1600-0587.2013.00571.x

Kapusta, A., & Bogacka-Kapusta, E. (2015). Non-native fish species in heated lakes: Origins and present status. *Archives of Polish Fisheries*, *23*(2), 121–129. https://doi.org/10.1515/aopf-2015-0014

King, K. C., Hall, M. D., & Wolinska, J. (2023). Infectious disease ecology and evolution in a changing world. In *Philosophical Transactions of the Royal Society B: Biological Sciences* (Vol. 378, Issue 1873). Royal Society Publishing. https://doi.org/10.1098/rstb.2022.0002

Krajaejun, T., Lerksuthirat, T., Garg, G., Lowhnoo, T., Yingyong, W., Khositnithikul, R., Tangphatsornruang, S., Suriyaphol, P., Ranganathan, S., & Sullivan, T. D. (2014). Transcriptome analysis reveals pathogenicity and evolutionary history of the pathogenic

oomycete *Pythium insidiosum. Fungal Biology*, *118*(7), 640–653.
https://doi.org/10.1016/j.funbio.2014.01.009

Krasnov, B. R., & Poulin, R. (2015). Relationships between parasite diversity and host diversity.
In S. Morand, B. R. Krasnov, & D. T. J. Littlewood (Eds.), *Parasite Diversity and
Diversification* (pp. 27–38). Cambridge University Press.
https://doi.org/10.1017/CBO9781139794749

Laking, A. E., Ngo, H. N., Pasmans, F., Martel, A., & Nguyen, T. T. (2017). *Batrachochytrium
salamandrivorans* is the predominant chytrid fungus in Vietnamese salamanders. *Scientific
Reports*, *7*(1), 44443. https://doi.org/10.1038/srep44443

Lellis, W. A., White, B. S. J., Cole, J. C., Johnson, C. S., Devers, J. L., Gray, E. V. S., &
Galbraith, H. S. (2013). Newly documented host fishes for the eastern elliptio mussel
Elliptio complanata. Journal of Fish and Wildlife Management, *4*(1), 75–85.
https://doi.org/10.3996/102012-JFWM-094

Lepelletier, F., Karpov, S. A., Alacid, E., Le Panse, S., Bigeard, E., Garcés, E., Jeanthon, C., &
Guillou, L. (2014). *Dinomyces arenysensis gen. et sp. nov.* (Rhizophydiales,
Dinomycetaceae fam. nov.), a Chytrid Infecting Marine Dinoflagellates. *Protist*, *165*(2),
230–244. https://doi.org/10.1016/j.protis.2014.02.004

Martel, A., Spitzen-Van Der Sluijs, A., Blooi, M., Bert, W., Ducatelle, R., Fisher, M. C.,
Woeltjes, A., Bosman, W., Chiers, K., Bossuyt, F., & Pasmans, F. (2013).
Batrachochytrium salamandrivorans sp. nov. causes lethal chytridiomycosis in amphibians.
Proceedings of the National Academy of Sciences of the United States of America, *110*(38),
15325–15329. https://doi.org/10.1073/pnas.1307356110

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet Journal, *17*(1), 10–12. http://www-huber.embl.de/users/an-

McMurdie, P. J., & Holmes, S. (2013). Phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. *PLoS ONE*, *8*(4).
https://doi.org/10.1371/journal.pone.0061217

McTaggart, A. R., Shivas, R. G., Geering, A. D. W., Vánky, K., & Scharaschkin, T. (2012). A
review of the *Ustilago-Sporisorium-Macalpinomyces* complex. *Persoonia: Molecular
Phylogeny and Evolution of Fungi*, *29*, 55–62. https://doi.org/10.3767/003158512X660283

Mendoza, L. (2009). *Pythium insidiosum* and mammalian hosts. In G. W. Beakes & S. Sekimoto
(Eds.), *Oomycete genetics and genomics: diversity, interactions, and research tools*.

Mendoza, L., Hernandez, F., & Ajello3, L. (1993). Life Cycle of the Human and Animal
Oomycete Pathogen *Pythium insidiosum*. In *JOURNAL OF CLINCAL MICROBIOLOGY*
(Vol. 31, Issue 11). https://journals.asm.org/journal/jcm

Metz, S., Itoïz, S., Obiol, A., Derelle, E., Massana, R., Berney, C., de Vargas, C., Soudant, P.,
Monier, A., & Chambouvet, A. (2023). Global perspective of environmental distribution
and diversity of Perkinsea (Alveolata) explored by a meta-analysis of eDNA surveys.
Scientific Reports, *13*(1), 1–13. https://doi.org/10.1038/s41598-023-47378-0

Mouritsen, K. N., Sørensen, M. M., Poulin, R., & Fredensborg, B. L. (2018). Coastal ecosystems
on a tipping point: Global warming and parasitism combine to alter community structure
and function. *Global Change Biology*, *24*(9), 4340–4356. https://doi.org/10.1111/gcb.14312

Naselli-Flores, L., & Padisák, J. (2023). Ecosystem services provided by marine and freshwater
phytoplankton. *Hydrobiologia*, *850*(12–13), 2691–2706. https://doi.org/10.1007/s10750-
022-04795-y

Palomar, G., Jakobik, J., Bosch, J., Kolenda, K., Kaczmarski, M., Josko, P., Roces-Diaz, J. V.,
Stachyra, P., Thumsova, B., Zielinski, P., & Pabijan, M. (2021). Emerging infectious
diseases of amphibians in Poland: distribution and environmental drivers. *Diseases of*
Aquatic Organisms, *147*, 1–12.

Pojmańska, T., Salamatin, R., Sulgostowska, T., Cielecka, D., Okulewicz, A., Niewiadomska,
K., & Grytner-Zięcina, B. (2012). The Polish collection of parasitic helminths. *Annals of*
Parasitology, *58*(2), 75–86. http://www.helminths.eu.http://www.helminths.eu

Powell, M. J., & Letcher, P. M. (2019). Ultrastructure of early stages of *Rozella allomycis*
(Cryptomycota) infection of its host, *Allomyces macrogynus* (Blastocladiomycota). *Fungal*
Biology, *123*(2), 109–116. https://doi.org/10.1016/j.funbio.2018.11.009

Powell, M. J., Letcher, P. M., Davis, W. J., Lefèvre, E., Brooks, M., & Longcore, J. E. (2019).
Taxonomic summary of *Rhizoclosmatium* and description of four new *Rhizoclosmatium*
species (Chytriomycetaceae, Chytridiales). *Phytologia*, *101*(2), 139–163.
www.phytologia.org

Prati, S., Grabner, D. S., Pfeifer, S. M., Lorenz, A. W., & Sures, B. (2022). Generalist parasites
persist in degraded environments: A lesson learned from microsporidian diversity in
amphipods. *Parasitology*, *149*(7), 973–982. https://doi.org/10.1017/S0031182022000452

Rad-Menéndez, C., Gerphagnon, M., Garvetto, A., Arce, P., Badis, Y., Sime-Ngando, T., &
Gachon, C. M. M. (2018). Rediscovering *Zygorhizidium affluens* Canter: Molecular
taxonomy, infectious cycle, and cryopreservation of a chytrid infecting the bloom-forming
diatom *Asterionella formosa*. *Applied and Environmental Microbiology*, *84*(23).
https://doi.org/10.1128/AEM.01826-18

Rasconi, S., Niquil, N., & Sime-Ngando, T. (2012). Phytoplankton chytridiomycosis:
Community structure and infectivity of fungal parasites in aquatic ecosystems.
Environmental Microbiology, *14*(8), 2151–2170. https://doi.org/10.1111/j.1462-
2920.2011.02690.x

R Core Team. (2023). *R: A Language and Environment for Statistical Computing*.
https://www.R-project.org/

Rug, M., & Vogt, G. (1994). Developmental stages of the crayfish parasite *Psorospermium*
haeckeli in thoracic arteries of *Ascatus ascatus*. *Journal of Invertebrate Pathology*, *64*, 153–
155.

Sassenhagen, I., Langenheder, S., & Lindström, E. S. (2023). Infection strategies of different
chytrids in a diatom spring bloom. *Freshwater Biology*, *68*(6), 972–986.
https://doi.org/10.1111/fwb.14079

Schulz, V. ;, Schulz, A. ;, Klamke, M. ;, Preissler, K. ;, Sabino-Pinto, J. ;, Müsken, M. ;,
Schlüpmann, M. ;, Heldt, L. ;, Kamprad, F. ;, Enss, J. ;, Schweinsberg, M. ;, Virgo, J. ;, Rau,
H. ;, Veith, M. ;, Lötters, S. ;, Wagner, N. ;, Steinfartz, S. ;, & Vences, M. (2020).
Batrachochytrium salamandrivorans *in the Ruhr district, Germany: History, distribution,*
decline dynamics and disease symptoms of the salamander plague. http://www.salamandra-
journal.com

Seto, K., Simmons, D. R., Quandt, C. A., Frenken, T., Dirks, A. C., Clemons, R. A., McKindles,
K. M., McKay, R. M. L., & James, T. Y. (2023). A combined microscopy and single-cell
sequencing approach reveals the ecology, morphology, and phylogeny of uncultured
lineages of zoosporic fungi. *MBio*, *14*(4), e0131323. https://doi.org/10.1128/mbio.01313-23

Shen, W., Le, S., Li, Y., & Hu, F. (2016). SeqKit: A cross-platform and ultrafast toolkit for
FASTA/Q file manipulation. *PLoS ONE*, *11*(10).
https://doi.org/10.1371/journal.pone.0163962

Shi, S., Wang, J., Liu, C., & Zheng, L. (2023). Developmental toxicity and inflammatory
response induced by Botrytis cinerea in zebrafish (*Danio rerio*) larvae. *Comparative*
Biochemistry and Physiology Part - C: Toxicology and Pharmacology, *267*.
https://doi.org/10.1016/j.cbpc.2023.109575

Simmons, D. R., Bonds, A. E., Castillo, B. T., Clemons, R. A., Glasco, A. D., Myers, J. M.,
Thapa, N., Letcher, P. M., Powell, M. J., Longcore, J. E., & James, T. Y. (2020). The
Collection of Zoosporic Eufungi at the University of Michigan (CZEUM): Introducing a
new repository of barcoded Chytridiomyceta and Blastocladiomycota cultures. *IMA*
Fungus, *11*(1). https://doi.org/10.1186/s43008-020-00041-z

Socha, D., & Hutorowicz, A. (2009). Changes in the quantitative relations of the phytoplankton
in heated lakes. *Archives of Polish Fisheries*, *17*(4), 239–251.
https://doi.org/10.2478/v10086-009-0017-9

Socha, D., & Zdanowski, B. (2001). Aquatic ecosystems in the vicinity of Konin. *Biblioteka*
Monitoringu OErodowisk, 75.

Stoeck, T., Bass, D., Nebel, M., Christen, R., Jones, M. D. M., Breiner, H. W., & Richards, T. A.
(2010). Multiple marker parallel tag environmental DNA sequencing reveals a highly
complex eukaryotic community in marine anoxic water. *Molecular Ecology*, *19*(SUPPL. 1),
21–31. https://doi.org/10.1111/j.1365-294X.2009.04480.x

Sures, B., Nachev, M., Schwelm, J., Grabner, D., & Selbach, C. (2023). Environmental
parasitology: stressor effects on aquatic parasites. *Trends in Parasitology*, *39*(6), 461–474.
https://doi.org/10.1016/j.pt.2023.03.005

Thines, M. (2018). Oomycetes. In *Current Biology* (Vol. 28, Issue 15, pp. R812–R813). Cell
Press. https://doi.org/10.1016/j.cub.2018.05.062

Thomé, P. C., Wolinska, J., Van Den Wyngaert, S., Reñé, A., Ilicic, D., Agha, R., Grossart, H.
P., Garcés, E., Monaghan, M. T., & Strassert, J. F. H. (2024). Phylogenomics including new
sequence data of phytoplankton-infecting chytrids reveals multiple independent lifestyle
transitions across the phylum. *Molecular Phylogenetics and Evolution*, *197*.
https://doi.org/10.1016/j.ympev.2024.108103

Urbańska, M., Andrzejewski, W., Riccardi, N., Gierszal, H., & Golski, J. (2018). The Invasive
Asian Clam *Corbicula fluminea* in Polish Rivers: The Importance of Thermal Discharge
from Power Plants for Its Spread. *Polish Journal of Ecology*, *66*(1), 70–75.
https://doi.org/10.3161/15052249PJE2018.66.1.008

Urbańska, M., Kirschenstein, M., Obolewski, K., & Ożgo, M. (2019). Silent invasion:
Sinanodonta woodiana successfully reproduces and possibly endangers native mussels in
the north of its invasive range in Europe. *International Review of Hydrobiology*, *104*(5–6),
127–136. https://doi.org/10.1002/iroh.201801971

Vallet, M. (2024). Chemical ecology of plankton parasitism in algae. In *Botanica Marina* (Vol.
67, Issue 3, pp. 211–229). Walter de Gruyter GmbH. https://doi.org/10.1515/bot-2023-0062

Van den Wyngaert, S., Ganzert, L., Seto, K., Rojas-Jimenez, K., Agha, R., Berger, S. A.,
Woodhouse, J., Padisak, J., Wurzbacher, C., Kagami, M., & Grossart, H. P. (2022).
Seasonality of parasitic and saprotrophic zoosporic fungi: linking sequence data to
ecological traits. *ISME Journal*, *16*(9), 2242–2254. https://doi.org/10.1038/s41396-022-
01267-y

Van den Wyngaert, S., Rojas-Jimenez, K., Seto, K., Kagami, M., & Grossart, H. P. (2018).
Diversity and Hidden Host Specificity of Chytrids Infecting Colonial Volvocacean Algae.
Journal of Eukaryotic Microbiology, *65*(6), 870–881. https://doi.org/10.1111/jeu.12632

Van den Wyngaert, S., Seto, K., Rojas-Jimenez, K., Kagami, M., & Grossart, H. P. (2017). A
New Parasitic Chytrid, *Staurastromyces oculus* (Rhizophydiales, Staurastromycetaceae
fam. nov.), Infecting the Freshwater Desmid *Staurastrum sp. Protist*, *168*(4), 392–407.
https://doi.org/10.1016/j.protis.2017.05.001

Wang, F., Wang, K., Cai, L., Zhao, M., Kirk, P. M., Fan, G., Sun, Q., Li, B., Wang, S., Yu, Z.,
Han, D., Ma, J., Wu, L., & Yao, Y. (2023). Fungal names: a comprehensive nomenclatural
repository and knowledge base for fungal taxonomy. *Nucleic Acids Research*, *51*(D1),
D708–D716. https://doi.org/10.1093/nar/gkac926

Wang, S., Cao, M., Wang, B., Deng, R., Gao, Y., & Liu, P. (2019). Optimization of growth
requirements and scale-up cultivation of freshwater algae *Desmodesmus armatus* using
response surface methodology. *Aquaculture Research*, *50*(11), 3313–3325.
https://doi.org/10.1111/are.14290

Wells, K., & Clark, N. J. (2019). Host Specificity in Variable Environments. In *Trends in
Parasitology* (Vol. 35, Issue 6, pp. 452–465). Elsevier Ltd.
https://doi.org/10.1016/j.pt.2019.04.001

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*.

Williamson, B., Tudzynski, B., Tudzynski, P., & Van Kan, J. A. L. (2007). *Botrytis cinerea*: The
cause of grey mould disease. In *Molecular Plant Pathology* (Vol. 8, Issue 5, pp. 561–580).
https://doi.org/10.1111/j.1364-3703.2007.00417.x

Winter, D. J. (2017). An R package for the NCBI eUtils API. *PeerJ Preprints*.

Wiśniewski, K., Szarmach, D., & Poznańska-Kakareko, M. (2020). The role of abiotic and biotic
factors in interspecific competition of Polish crayfish-Comprehensive literature review. In

Oceanological and Hydrobiological Studies (Vol. 49, Issue 4, pp. 428–441). De Gruyter Open Ltd. https://doi.org/10.1515/ohs-2020-0038

Wood, C. L., Zgliczynski, B. J., Haupt, A. J., Guerra, A. S., Micheli, F., & Sandin, S. A. (2018). Human impacts decouple a fundamental ecological relationship—The positive association between host diversity and parasite diversity. *Global Change Biology*, *24*(8), 3666–3679. https://doi.org/10.1111/gcb.14159

Woolway, R. I., & Merchant, C. J. (2019). Worldwide alteration of lake mixing regimes in response to climate change. *Nature Geoscience*, *12*(4), 271–276. https://doi.org/10.1038/s41561-019-0322-x

Yamato, S., Sasaki, N., Komatsu, K., Mashimo, K., & Motobayashi, T. (2024). Temporal changes in colonization levels of *Ustilago esculenta* in different parts of *Zizania latifolia* plants competent and incompetent for stem gall formation. *Journal of General Plant Pathology*, *90*(1), 14–22. https://doi.org/10.1007/s10327-023-01152-2

Yap, T. A., Koo, M. S., Ambrose, R. F., Wake, D. B., & Vredenburg, V. T. (2015). Averting a North American biodiversity crisis. *Science*, *349*(6247), 481–482. https://doi.org/10.1126/science.aab1052

Yuryshynets, V., & Krasutska, N. (2009). Records of the parasitic worm *Aspidogaster conchicola* (Baer 1827) in the Chinese pond mussel *Sinanodonta woodiana* (Lea 1834) in Poland and Ukraine. *Aquatic Invasions*, *4*(3), 491–494. https://doi.org/10.3391/ai.2009.4.3.9

Zdanowski, B., Napiórkowska-Krzebietke, A., Stawecki, K., Świątecki, A., Babko, R., Bogacka-Kapusta, E., Czarnecki, B., & Kapusta, A. (2020). *Polish River Basins and Lakes-Part I* (E. Korzeniewska & M. Harnisz, Eds.). Springer. http://www.springer.com/series/698