Molecular mechanisms underlying p62-dependent secretion of the Alzheimer-associated ubiquitin variant, UBB[+1]

Ajay R. Wagh[1] and Michael H. Glickman[1]

[1] The Faculty of Biology, Technion Israel Institute of Technology, Haifa, 3200003, Israel

Corresponding author: The Faculty of Biology #438, Technion Israel Institute of Technology, Haifa, 3200003, Israel

Email: glickman@technion.ac.il

Preprint: BioRxiv preprint **doi:** https://doi.org/10.1101/2024.12.31.630908

Classification: Biological Sciences / Biochemistry

Keywords: Alzheimer's disease; autophagy; autophagosome; lysosome; neurodegeneration; p62 bodies; proteasome; protein aggregates; secretory autophagy; SNARE; SQSTM1

2. Abstract

UBB[+1], a ubiquitin variant protein resulting from a frameshift in the ubiquitin-B gene, is a pathological hallmark of Alzheimer disease (AD). At the cellular level, UBB[+1] disrupts the ubiquitin-proteasome system while inducing autophagy. Notably, UBB[+1] itself is secreted via autophagosome-like vesicles. Here, we demonstrate that UBB[+1] can be removed from the cell by degradative and secretory autophagy. SQSTM1/p62 functions as a pivotal ubiquitin receptor for UBB[+1], recognizing its ubiquitin domain and facilitating loading into autophagosomes. Oligomerization of SQSTM1/p62 was critical to isolate UBB[+1] in bodies preventing its aggregation. Intriguingly, both gain- and loss-of-function SQSTM1/p62 suppressed UBB[+1] secretion, causing intracellular retention: SQSTM1/p62 knockout led to UBB[+1] accumulation in insoluble aggregates, while its overexpression promoted the formation of p62-UBB[+1] bodies. We further identified distinct roles for SNARE-mediated membrane fusion in secretory autophagy of UBB[+1]. Specifically, the R-SNARE SEC22B and the Q-SNAREs Syntaxin-4 (STX4) and SNAP23 participated in UBB[+1] exocytosis. Disruption of SEC22B impaired the fusion of UBB[+1]-containing autophagosomes with the plasma membrane, reducing UBB[+1] secretion without affecting its intracellular turnover. Inhibition of lysosomes partially stabilized UBB[+1] indicating that degradation and secretion are complementary processes that determine the fate of UBB[+1]. This study elucidates the dual roles of autophagy in managing neurotoxic proteins, highlighting SQSTM1/p62 as a key mediator of UBB[+1] trafficking and secretion. Although ubiquitin typically acts as a degradation signal, our findings reveal a rare instance of a ubiquitin-related protein driving secretory autophagy. These findings advance our understanding of cellular

mechanisms underlying the clearance of misfolded proteins in neurodegenerative

diseases.

3. Significance

To maintain health, cells must remove toxic proteins, typically by ubiquitin-dependent

degradation. Neurons are particularly sensitive since they are not dividing and cannot

replace damaged cells. This study suggests that autophagy, a pathway for degrading

proteins, can also secrete harmful proteins, highlighting a new pathway for tackling

neurodegenerative diseases. Specifically, UBB[+1], a variant of the ubiquitin protein linked

to Alzheimer's disease, is recognized by a central autophagy component, p62,

packaged into vesicles and secreted from cells. When p62 is absent, UBB[+1]

accumulates inside cells, which can be harmful. This work identifies UBB[+1] as a novel

cargo for secretory autophagy, extending the understanding of how cells may handle

proteotoxic stress beyond classical degradation pathways.

4. Main Text

Introduction:

Autophagy is traditionally viewed as a degradative process (1), but growing evidence

has revealed its role in vesicle production (2), the secretion of proteins such as pro-

inflammatory cytokines (3), or lysozyme release (4). These processes, collectively

termed autophagy-associated secretion, or secretory autophagy, define an autophagy

pathway that plays an active role in maintaining cellular homeostasis in both healthy

and disease states (2-4). Unlike conventional degradative autophagy, where cytosolic

cargoes are broken down, secretory autophagy involves the transport of these cargoes

within autophagosomes to the extracellular environment through exocytosis (5). This

secretory autophagy machinery has been implicated in the exocytosis of various

elements, including cytokines (such as IL-1β, IL-6, IL-18, and TNF-α), aggregated

proteins, organelles, and microbial components (5, 6). An important role of degradative

autophagy, by contrast, is the removal of ubiquitinated cargo such as ubiquitinated

proteins, ubiquitinated aggregates (aggrephagy), and ubiquitinated organelles (e.g.,

mitophagy) (7, 8). While many studies have explored autophagy-associated protein

secretion/release, few have focused on the recruitment of cargo into secretory

autophagosomes.

Several dedicated ubiquitin receptors facilitate the loading of cargo into

autophagosomes, serving as bridges between ubiquitinated cargo and the isolation

membrane (9). These receptors possess specific domains: a ubiquitin recognition

element on one side and an LC3-interacting region (LIR) on the other. One classical

autophagy receptor is p62/sequestosome 1 (SQSTM1), which plays a crucial role in

clearing ubiquitinated proteins (10, 11). SQSTM1/p62 abnormal expression or

SQSTM1gene mutation is tightly associated with various diseases including AD.

SQSTM1/p62 has a multidomain structure tailored for this function: its N-terminal PB1

domain mediates self-oligomerization, a central LIR motif targets it to the phagophore,

and a C-terminal ubiquitin-associated (UBA) domain facilitates ubiquitin binding (9, 12,

13). Through its UBA domain, SQSTM1/p62 noncovalently binds monoubiquitin and

Lys48- or Lys63-linked polyubiquitin chains (14-16). Notably, proteins with UBA

domains generally exhibit a stronger preference for binding polyubiquitin chains over

monoubiquitin (17, 18).

Misfolded protein aggregation in both intracellular and extracellular environments is a

key factor in the development of Alzheimer disease (AD). Neurons rely on autophagy for

the removal of defective proteins to maintain neuronal health (19-22). The primary

pathological hallmarks of AD are the formation of amyloid plaques, which are primarily

composed of aggregated amyloid-β (Aβ) peptides, and intracellular neurofibrillary

tangles containing hyper-phosphorylated tau (23-25). UBB[+1] is a non-hereditary

mutation of ubiquitin that results from a transcription error leading to a dysfunctional

ubiquitin. UBB[+1] is associated with AD by accumulating in both extracellular plaques

and intracellular tangles of patients (26). UBB[+1] arises from transcriptional frameshift

errors in the ubiquitin B gene and accumulates robustly in neurofibrillary tangles and

amyloid plaques of advanced Alzheimer's disease patients (26-28). A transgenic mouse

model further confirmed UBB[+1] deposition in hippocampal and brainstem regions

associated with AD-like phenotypes, including memory deficits and proteostasis

dysfunction (29, 30). Human post-mortem studies have also reported UBB[+1]

accumulation as early as Braak stages III–IV (25), highlighting its progressive buildup.

Although the precise ratio of UBB[+1] to wild-type ubiquitin in vivo is difficult to measure,

its persistent presence across disease stages points to its biological significance.

This dual intra- and extra-cellular localization raises questions about the pathway by

which UBB[+1] exits the cell. Indeed, in our previous study, we investigated this pathway

and demonstrated that autophagy components play a role in the unconventional

secretion of UBB[+1] (31). Unconventional secretion involves the packaging of cytosolic

proteins into vesicles which ultimately fuse to the plasma membrane to release their

contents into the extracellular space (32, 33). This process is mediated by SNARE

proteins, which catalyze membrane fusion. Among the 38 known mammalian SNARE

proteins, each is selectively localized to specific cellular membranes, thereby defining

distinct trafficking pathways (34). SNARE proteins are classified based on the central

amino acid residues in their SNARE motifs: Q-SNAREs for glutamine and R-SNAREs

for arginine. A functional SNARE complex required for membrane fusion consists of a

parallel four-helix bundle formed by one Qa-, Qb-, Qc-, and R-SNARE. Specific sets of

these proteins catalyze distinct trafficking pathways.

In this study, we show that autophagy-mediated secretion of UBB^{+1} is dependent on

SQSTM1/p62. Knockout (KO) of p62 in HeLa cells significantly disrupts UBB^{+1}

recruitment to SEC22B-containing secretory autophagosome-like vesicles, resulting in

the formation of intracellular aggregates and a marked inhibition of its secretion. We

further elucidate the participation of the SNARE proteins, SEC22B, STX4 and SNAP23

in UBB^{+1} exocytosis. These findings provide novel perspectives on the cellular

mechanisms involved in the removal of misfolded proteins that can cause significant

damage to the cell and lead to neurodegenerative diseases.

Results:

Secretion of UBB^{+1} shares properties with secretory autophagy

In secretory autophagy, SEC22B is one of the factors responsible for mediating fusion

with the plasma membrane and plays an essential role in the clearance of an

autophagic substrate (35, 36). We generated *sec22b* KO HeLa cells (**Fig. S1A)** and

confirmed that loss of SEC22B impairs autophagic flux, resulting in autophagosome

accumulation. Expression of UBB^{+1} induces LC3B-II, which can be visualized as puncta

(**Fig. 1A and B and Fig. S1B**). In *sec22b* KO cells, UBB^{+1} secretion is suppressed, and

colocalization with LC3B is more readily observed, consistent with impaired autophagic

flux and blocked vesicle clearance (**Fig. 1B and Fig. S1C**).

To confirm whether these autophagosomes are involved in UBB^{+1} secretion, we utilized

sec22b KO cells overexpressing UBB^{+1} and evaluated its release to the growth media

(**Fig. S1D**). Treatment of UBB^{+1}-expressing *sec22b* KO cells with cycloheximide (CHX),

a protein synthesis inhibitor, showed that its secretion was significantly reduced

compared to wild-type cells (**Fig. 1C**). To rule out the possibility that secretion was

caused by cytotoxicity or apoptosis under CHX treatment or other experimental

conditions, we assessed cleaved caspase-3 levels by Western blot. No cleaved

caspase-3 was detected in WT or *sec22b* KO cells expressing UBB^{+1} with or without

CHX or CHX + Bafilomycin A1. In contrast, a strong cleaved caspase-3 band was

observed in cells treated with Venetoclax (ABT-199), a known inducer of intrinsic

apoptosis (37), confirming the effectiveness of the assay (**Fig. S1G**). These results

demonstrate that UBB^{+1} secretion under these conditions is not due to apoptotic cell

death.

At the plasma membrane, secretory autophagosomes containing SEC22B, an R-

SNARE, fuse with Q-SNAREs such as STX3 and STX4 (38, 39). As shown in **Figure**

1D, knockdown of STX4 by dicer-substrate small interfering RNA (DsiRNA) reduced

UBB^{+1} secretion. Conversely, knockdown of STX3 had no effect on UBB^{+1} secretion

(**Fig. 1D**). To further clarify the role of SNARE machinery, we performed knockdown

experiments targeting additional Q-SNARE proteins SNAP23 and SNAP29. Knockdown

of SNAP23 significantly impaired secretion of UBB^{+1}, similarly to STX4 knockdown,

while SNAP29 knockdown had no significant effect compared to cells treated with non-

targeting (NT) siRNA (**Fig. 1D**). These findings indicate that SNAP23, together with

STX4, are Q-SNAREs participating in fusion of UBB[+1]-containing secretory

autophagosomes with the plasma membrane. Alternatively, in degradative autophagy,

STX17 facilitates autophagosome fusion with the lysosome (40-42). Knockdown of

STX17 had no significant effect on UBB[+1] secretion (**Fig. S1E**), indicating either that

STX17 is redundant, or that degradative autophagy is not implicated in the secretion of

UBB[+1]. Similarly, we did not find evidence for exosome-mediated secretion of UBB[+1], as

knockdown of TSG101, a key regulator of exosome biogenesis, did not reduce UBB[+1]

secretion (**Fig. S1F**). Consistent with this, we have previously shown that ALIX knockout

HeLa cells also secrete UBB[+1] efficiently, further supporting that exosome pathways are

not involved in UBB[+1] secretion (31).

To investigate the relationship between secretion and intracellular turnover dynamics,

we expressed UBB[+1] in HeLa cells and treated them with CHX. Intracellular UBB[+1]

levels decreased continuously over 4 to 8 h of CHX treatment, indicating a biological

half-life of roughly eight hours (**Fig. 1E**). Nevertheless, the intracellular turnover of

UBB[+1] persisted despite the absence of SEC22B, while its secretion was significantly

impaired under the same conditions (**Fig. 1C**). Taken together, these results highlight

the roles of SEC22B and STX4/SNAP23 in the secretory autophagy machinery for

UBB[+1] secretion, while an alternative mechanism is likely operating to clear the

intracellular buildup of UBB[+1].

Secretion and degradation of UBB[+1] both account for its intracellular turnover

dynamics

To determine if intracellular degradation contributes to UBB^{+1} turnover, its cellular half-

life was evaluated upon addition of lysosomal (Bafilomycin) or proteasomal (Velcade)

inhibitors and CHX treatment. As shown in **Figure 2A**, intracellular UBB^{+1} levels

significantly decreased during CHX treatment, irrespective of inhibitor treatment. CHX

and Velcade efficiency were confirmed as shown in **Figure S2A and S2B**. These

turnover dynamics indicate that intracellular degradation alone cannot fully account for

the clearance of UBB^{+1} from the cells. In contrast, the effect of intracellular proteolysis

inhibitors on extracellular UBB^{+1} levels was minimal (**Fig. 2B**), suggesting that secretion

serves as a compensatory mechanism when degradation pathways are impaired. In

sec22b-deficient cells, where UBB^{+1} secretion is significantly impaired but not

completely abolished, treatment with Bafilomycin further enhanced extracellular UBB^{+1}

levels (**Fig. 2C and 2D**). This suggests that when lysosomal degradation is blocked,

UBB^{+1} may be redirected toward alternative vesicular pathways for secretion, even in

the absence of SEC22B. In parallel, the intracellular pool of UBB^{+1} was partially

stabilized (**Fig. 2E and 2F**). These data support our earlier observation (31) that UBB^{+1}

degradation and secretion are complementary processes that together determine the

fate of UBB^{+1}. We conclude that UBB^{+1}-containing autophagosome-like vesicles can be

targeted either to the lysosome or to the plasma membrane, effectively distributing their

cargo between these two destinations. This observation prompted further investigation

into the mechanisms of how UBB^{+1} is packaged into these vesicles.

SQSTM1/p62 is required for efficient clearance of UBB^{+1} aggregates and its

secretion

The uptake of ubiquitinated cargo into autophagosomes generally relies on specialized

ubiquitin receptors. To test whether any of the known autophagy-related ubiquitin

receptors are responsible for directing UBB^{+1} to secretion, we obtained individual and

combinatorial knockouts of TAX1BP1, NBR1, OPTN, NDP52, and SQSTM1/p62 (**Fig.**

S3A-D). A *penta* KO (5 KO) of all five receptors (*tax1bp1/optn/ndp52/nbr1/p62)*

abolished secretion of UBB^{+1} (**Fig. 3A**). Since TAX1BP1 contains a UBZ (Ubiquitin-

Binding Zinc Finger) domain with specificity for monoubiquitin, we considered it a

potential candidate for association with the "single ubiquitin domain protein", UBB^{+1}.

However, an individual *tax1bp1* KO showed no significant effect on UBB^{+1} secretion

(**Fig. S3E**). Even a *triple* KO (TKO) of a subset of these receptors (*optn/ndp52/tax1bp1)*

showed no significant effect on UBB^{+1} secretion (**Fig. 3B and 3C**). Surprisingly, the

individual *p62* KO exhibited a complete loss of UBB^{+1} secretion (**Fig. 3D and 3E**),

indicating that SQSTM1/p62 is essential and sufficient for UBB^{+1} secretion. In addition

to its known role in aggrephagy (43-45), this study reveals that SQSTM1/p62 also plays

a role in secretory autophagy.

Having uncovered an exclusive role for SQSTM1/p62 in UBB^{+1} secretion we turned our

attention to the intracellular fate of UBB^{+1}. How is its accumulation managed in the

absence of secretion? Amorphous UBB^{+1} clusters were visualized in a KO of

SQSTM1/p62 by immunofluorescence (**Fig. 3F**). Both *p62* KO and *penta* KO (**Fig. S3F**)

cells accumulated UBB^{+1}-positive material, showing a similar spread-out distribution,

suggesting that loss of SQSTM1/p62 alone is sufficient to disrupt UBB^{+1}. Does the

involvement of SQSTM1/p62 in elimination of these UBB^{+1} puncta imply that intracellular

UBB^{+1} is prone to aggregation? Indeed, the UBB^{+1} protein accumulated in insoluble

fractions of *p62* KO cell extracts (**Fig. 3G**). To conclude, *SQSTM1/p62* deficient cells

displayed a significant block in UBB^{+1} aggregate clearance, highlighting SQSTM1/p62's

role as a specific receptor for UBB^{+1}, potentially for both degradative and secretory

autophagy.

Functional association between SQSTM1/p62 and the ubiquitin domain of UBB^{+1}

Since SQSTM1/p62 interferes with UBB^{+1} aggregate formation on the one hand and is

required for its secretion on the other hand, we explored the potential for direct

interactions between these two proteins. As shown in **Figure 4A**, immunofluorescence

of UBB^{+1}-expressing HeLa cells revealed co-localization with endogenous SQSTM1/p62

in the form of intracellular puncta. However, a significant portion of UBB^{+1} was

delocalized throughout the cell. This could suggest that the levels of endogenous

SQSTM1/p62 were insufficient to associate with the heterologously expressed UBB^{+1}.

To address this, we repeated the experiment in cells overexpressing SQSTM1/p62.

Immunofluorescence analysis of co-expressing cells showed prominent perinuclear

puncta containing both proteins.

Previous studies have reported the formation of membrane-less inclusion bodies

composed of SQSTM1/p62 and ubiquitinated substrates upon ectopic expression of

SQSTM1/p62, termed 'p62-bodies' or 'sequestosomes' (45, 46). We similarly observed

the formation of such p62-bodies when SQSTM1/p62 was overexpressed in the current

study (**Fig. S4A**). However, in UBB^{+1}-expressing cells, these bodies were organized in

an ordered, perinuclear pattern (**Fig. 4B and S4B**) strengthening our conclusion that the

two proteins are co-localized. The presence of a potential (ubiquitinated) substrate,

such as UBB^{+1}, appears to influence the intracellular localization and organization of

these p62-bodies.

SQSTM1/p62 is a critical ubiquitin-binding protein that facilitates the shuttling of

ubiquitinated cargo towards autophagosomes. To elucidate the molecular mechanisms

of recognition, we mutated key recognition elements within the ubiquitin domain of

UBB^{+1}. Previous studies have identified two hydrophobic surface residues on ubiquitin,

Phe4 (F4) and Ile44 (I44), that are recognized directly by the overwhelming majority of

Ub receptors (47). To investigate the role of these hydrophobic patches in UBB^{+1}

recognition, we generated a double substitution F4A and I44A in UBB^{+1} and co-

expressed UBB$^{+1\ F4A,\ I44A}$ with SQSTM1/p62 in HeLa cells. Immunofluorescence showed

a complete loss in co-localization between SQSTM1/p62 and UBB$^{+1\ F4A,\ I44A}$ and

disruption of the characteristic perinuclear pattern of UBB^{+1}-containing *p62-bodies* (**Fig.**

4C). A significant reduction in secretion of the hydrophobic patch mutant compared to

WT UBB^{+1} (**Fig. 4D**) further supported the importance of the ubiquitin domain for

recognition of UBB^{+1} by SQSTM1/p62 and for its secretion. Additionally, the

hydrophobic patch mutant led to undetectable ubiquitination on UBB$^{+1\ F4A,\ I44A}$ (**Fig. 4D**).

Therefore, our next question was whether the binding of SQSTM1/p62 is directly to the

surface of UBB^{+1} or to its ubiquitinated form.

Co-immunoprecipitation (co-IP) was employed to investigate whether SQSTM1/p62

preferentially binds to the ubiquitinated or non-ubiquitinated forms of UBB^{+1}. First,

immunoprecipitation (IP) of MYC-tagged UBB^{+1} co-precipitated HA-p62 (**Fig. 4E**),

indicating a specific association between the two proteins. Next, precipitation of

endogenous SQSTM1/p62 co-immunoprecipitated UBB^{+1} (**Fig. 4F**), further supporting a

physical association. Consistent with a requirement for an intact ubiquitin surface, the

hydrophobic-patch mutant MYC-UBB^{+1} $^{F4A, I44A}$ failed to co-precipitate endogenous

SQSTM1/p62 and showed negligible colocalization by immunofluorescence (**Fig. 4F**).

To test whether additional ubiquitin chains are required for this association, we blocked

all cellular ubiquitination by inhibiting the ubiquitin-activating enzyme (E1). This

treatment depleted polyubiquitin conjugates and eliminated ubiquitination of UBB^{+1} (**Fig.**

S4C and S4D). The remaining unmodified UBB^{+1} nevertheless interacted with

SQSTM1/p62 in co-IP assays and colocalized in cells (**Fig. 4G and 4H**). Across all co-

IP's, we noticed that the ratio between ubiquitinated and unmodified UBB^{+1} in the co-IP

eluate was unaltered relative to the ratio in whole cell extract (**Fig. 4F**) indicating that

SQSTM1/p62 does not have a marked preference for one form or the other**.**

Collectively, these findings identify SQSTM1/p62 as an interacting partner of UBB^{+1}

independent of its ubiquitination state. Through this association, SQSTM1/p62 plays a

crucial role in regulating intracellular levels of UBB^{+1} by alleviating its aggregation and

promoting its secretion.

Mapping SQSTM1/p62 domains involved in UBB^{+1} binding and secretion

As a scaffold protein, SQSTM1/p62 has multiple functional domains, through which it

associates with its interacting partners. Beyond a C-terminal UBA domain that binds

ubiquitinated cargo (15, 48-50), SQSTM1/p62 contains an N-terminal PB1 domain

which drives self-oligomerization, and an LC3-interacting region (LIR) motif mediating

the interaction with ATG8 family proteins (51-53). To better understand the role of these

domains in regulating UBB^{+1} secretion, we constructed SQSTM1/p62 lacking its PB1

domain (p62ΔPB1), a version with a non-functional LIR motif (p62 $^{W340A, L343A}$), and a

UBA domain mutant (p62 ^{M404V}), which disrupts hydrophobic interactions essential for

ubiquitin binding. All constructs were HA-tagged and expressed in *p62* KO HeLa cells

co-expressing UBB^{+1} (**Fig. 5A**).

In contrast to full-length (FL) SQSTM1/p62, overexpression of truncated SQSTM1/p62

(p62ΔPB1) lacking the self-oligomerization property did not form *p62-bodies* with UBB^{+1}

(**Fig. 5B**), emphasizing the essential role of the PB1 domain in body formation. By

contrast, the construct lacking a functional LIR motif (p62 $^{W340A,\ L343A}$) displayed the

characteristic perinuclear *p62-bodies* (**Fig. 5B**), supporting a model whereby *p62-bodies*

form upstream to autophagosomes. Strikingly, the M404V mutation in the UBA domain

abolished p62's ability to co-localize with UBB^{+1}, indicating that UBA-dependent binding

to ubiquitinated UBB^{+1} is critical for cargo engagement (**Fig. 5B**).

In *p62* KO cells, expression of UBB^{+1} alone resulted in a complete loss of secretion

compared to WT cells. Overexpression of HA-p62FL in the KO background restored

secretion, demonstrating that full-length p62 is sufficient to rescue this defect (**Fig. 5G**).

In contrast, none of the mutant constructs (p62ΔPB1, p62W340A/L343A, or

p62M404V), expressed at comparable levels (**Fig. 5B**), were able to rescue secretion

(**Fig. 5D–F**). These findings underscore the requirement for all three functional modules

(PB1, LIR, and UBA) in facilitating efficient UBB^{+1} sequestration and routing into the

secretory autophagy pathway. To exclude the possibility that these results reflect

artifacts of SQSTM1/p62 overexpression, we performed dose-dependent titration

experiments in both WT and *p62* KO cells (**Fig. S5A–C**). Increasing plasmid

concentrations of HA-p62FL (0 ng, 100 ng, 250 ng, 500 ng, 1 µg) demonstrated that at

moderate expression levels, HA-p62FL reproducibly enhanced UBB^{+1} secretion,

whereas very high overexpression led to a dominant-negative effect through the

formation of large p62-bodies (**Fig. S5B**). Expression levels of the different HA-p62

constructs were validated by both immunofluorescence and western blotting, confirming

comparability across constructs (**Fig. 5B**). Together, these results establish HA-p62FL

as a robust rescue reference and demonstrate that disruption of the PB1, LIR, or UBA

domains abolishes the ability of SQSTM1/p62 to restore UBB^{+1} secretion.

.

Without a functional LIR motif, SQSTM1/p62 is not expected to shuttle ubiquitinated

cargo to LC3-containing phagophores, explaining the inability of this mutant to support

UBB^{+1} secretion (**Fig. 5E**). However, why would a SQSTM1/p62 construct lacking a self-

oligomerization (PB1) domain be unable to promote the secretion of UBB^{+1} (**Fig. 5D**)?

In light of the inability of p62ΔPB1 to co-localize with UBB^{+1} (**Fig. 5C**), we proposed that

the PB1 may potentially participate in UBB^{+1} binding. To further define the binding

requirements, we next evaluated the role of the UBA domain using an HA-p62^{M404V}

mutant known to abolish binding to polyubiquitin (54-56). Co-immunoprecipitation and

immunofluorescence assays showed that HA-p62^{M404V} failed to interact with or form p62

bodies with UBB^{+1} (**Fig. 5C and 5J**). Moreover, when expressed in p62 KO cells, this

mutant was unable to rescue UBB^{+1} secretion (**Fig. 5F**), in contrast to HA-p62 FL (**Fig.**

5G). Importantly, Western blot analysis demonstrated that expression levels of HA-p62

FL, HA-p62ΔPB1, HA-p62 $^{W340A, L343A}$, and HA-p62 ^{M404V} were all comparable and within

moderate ranges, suggesting that the failure to rescue secretion was specifically due to

impaired function rather than artifacts from excessive overexpression (**Fig. 5B**). These

results strongly support the notion that the intact UBA domain and specifically residue

M404 is essential for p62-mediated UBB[+1] recognition, aggregation, and secretion.

It has been reported that the deletion of the PB1 domain or oligomerization-inhibiting

mutations resulted in reduced interaction with both LC3B and ubiquitin in pull-down

assays, suggesting that oligomerization may increase the interaction with these binding

partners (8, 45). Therefore, we tested whether p62ΔPB1 could bind to UBB[+1]. A

pulldown assay of HA-p62ΔPB1 showed no evidence of interaction with UBB[+1] (**Fig.**

5G). Apparently, due to the loss of oligomerization of SQSTM1/p62, p62ΔPB1 was

insufficient to trap UBB[+1] despite containing an intact UBA domain. The inherent

property of SQSTM1/p62 to oligomerize and form tandem interactions with multiple Ub

domains may underline the body formation we witnessed with UBB[+1]. Lacking the ability

to self-oligomerize, p62ΔPB1 can only bind a single unit at a time, which for a single

ubiquitin domain protein such as UBB[+1] is insufficient, apparently, for tight binding. For

this reason, we propose that p62ΔPB1 was unable to sequester UBB[+1] or package it

into autophagosomes.

To gain additional structural insight into the interaction between UBB[+1] and the UBA

domain of SQSTM1/p62, we used AlphaFold3 to model a complex between the two

proteins based on their primary sequences (**Fig. S5D and S5E**). The predicted structure

revealed that residue M404 in the p62 UBA domain is positioned near to a conserved

hydrophobic patch on UBB[+1] comprising residues Isoleucine 44, Leucine 8, and Valine

70. When this methionine was mutated to valine (V404), the predicted distance between

V404 and these hydrophobic residues increased, suggesting a loss of stabilizing

hydrophobic contacts. These structural observations are consistent with our biochemical

and imaging data, which show that the M404V mutation disrupts UBB^{+1} binding, impairs

p62-body formation, and blocks UBB^{+1} secretion (**Fig. 5C and 5F**). This supports a

critical role for M404 in mediating UBB^{+1} recognition by p62 through hydrophobic

interaction.

Discussion:

Accumulation of UBB^{+1} is toxic to cells, particularly neurons, where it disrupts cellular

homeostasis. Recently, UBB^{+1} has been identified as a key factor in inducing AD-like

pathology (25). Cells appear to have evolved multiple mechanisms to counteract

UBB^{+1}-associated stress by reducing its intracellular levels. UBB^{+1} can be partially

degraded by the proteasome (57, 58), and findings from this study suggest that

lysosomes also contribute to its intracellular turnover. Cells can also actively secrete

UBB^{+1} via an unconventional autophagosome-like vesicle-mediated pathway (31).

Overexpression of UBB^{+1} in cultured cells enables detailed mechanistic analysis of its

trafficking and clearance, a widely used approach for studying aggregation-prone

proteins implicated in neurodegeneration (31, 59, 60). Insights gained from such models

have consistently mirrored findings in neurons and patient-derived tissues, thereby

reinforcing the physiological relevance of our observations. In this study, we identified a

novel function of secretory autophagy in mitigating UBB^{+1} toxicity. Our findings show

that UBB^{+1} is actively regulated by an autophagy-mediated vesicle release mechanism,

with the autophagy receptor SQSTM1/p62 playing a central role. SQSTM1/p62

promotes the sequestration of UBB^{+1} into specialized *p62-bodies*, a process that occurs

independently of UBB^{+1}'s ubiquitination status (**Fig. 3 and 4**). The N-terminal PB1

domain of SQSTM1/p62 plays a critical role in its oligomerization, allowing the formation

of higher-order structures that are essential for the efficient sequestration of

ubiquitinated cargo (45, 61). This oligomerization facilitates the clustering of

ubiquitinated proteins into *p62-bodies*, which act as scaffolds for selective autophagy.

Likewise, our results revealed that the PB1 domain is essential for the sequestration of

UBB[+1] by SQSTM1/p62, and in its absence, *p62-body* formation was disrupted (**Fig. 5**).

The measured affinity of the UBA domain of SQSTM1/p62 for monoubiquitin is relatively

weak (K_d = *540 µM*) (15). The PB1 domain facilitates higher ordered oligomers of

SQSTM1/p62, which interact with multiple ubiquitin molecules (45), which in our case

explains how SQSTM1/p62 can bind the non-ubiquitinated form of UBB[+1].

A limitation of these assays is that they rely on overexpression of p62 constructs. While

our titration experiments confirmed that moderate expression levels reproducibly restore

UBB[+1] secretion, we also observed that very high overexpression can reduce secretion

due to dominant-negative effects caused by excessive *p62-body* formation. These

findings indicate that dosage is an important variable when interpreting rescue

experiments and should be considered in future studies using more physiological

expression systems.

The ensuing *p62-bodies* help isolate UBB[+1] from the cytosol, preventing its aggregation

and facilitating its recruitment into autophagosome-like vesicles for secretion (**Fig. 6)**.

SQSTM1/p62 is shared between degradative and secretory autophagy. Recent studies

have begun to uncover roles for SQSTM1/p62 in the secretion of other

neurodegenerative disease–related proteins. For example, a recent report (62)

demonstrated that SQSTM1/p62 facilitates the unconventional secretion of alpha-

synuclein from neurons using mechanisms that overlap with those identified here for

UBB[+1]. In that study, SQSTM1/p62 was shown to interact with intracellular alpha-

synuclein and direct it to autophagosome-derived secretory vesicles, independent of

canonical lysosomal degradation. This emerging role of SQSTM1/p62 as a gatekeeper

not only for degradation but also for secretion of misfolded or aggregation-prone

proteins underscores a broader neuroprotective strategy employed by cells. Our

findings extend this concept to UBB[+1], an aberrant ubiquitin variant associated with AD,

highlighting that SQSTM1/p62-driven secretory autophagy may be generalizable for

mitigating proteotoxic stress across multiple neurodegenerative contexts. These

parallels suggest that similar vesicular routes may govern the disposal of different

pathological proteins, offering potential targets for modulating disease progression.

The mere presence of UBB[+1] increases autophagic vesicle formation (31, 63).

Supporting this observation, our results show that intracellular UBB[+1] accumulation in

the absence of SQSTM1/p62 strongly induces LC3B-II conversion, indicating that

aggregated UBB[+1] itself is sufficient to trigger autophagic responses independently of

SQSTM1/p62 (**Fig. S3G**). Future studies should further explore whether such UBB[+1]-

induced autophagy is predominantly degradative or can also stimulate SEC22B-

mediated secretory pathways under different cellular conditions, such as nutrient

deprivation. Once packaged into autophagic vesicles aided by SQSTM1/p62, a subset

of UBB[+1]-containing autophagosomes bypass lysosomal degradation.

Depletion of SEC22B, an R-SNARE involved in unconventional protein secretion [3, 31],

also reduced UBB[+1] secretion. In the absence of SEC22B, UBB[+1] colocalized with LC3-

containing autophagosomes (**Fig. 1**). Although SEC22B plays a significant role in UBB[+1]

secretion, our findings suggest that additional compensatory secretory pathways may

exist, particularly under conditions where intracellular degradation is impaired. This may

either involve other SNAREs or non-canonical vesicular trafficking pathways

independent of classical secretory autophagy.

Knockdown of syntaxin 17 (STX17), a Qa-SNARE essential for autophagosome-

lysosome fusion [35], did not impair UBB^{+1} secretion (**Fig. S1C**). Instead, we identified

syntaxin 4 (STX4) as the critical Qa-SNARE mediating the fusion of UBB^{+1}-containing

vesicles with the plasma membrane. Knockdown of STX4 significantly reduced UBB^{+1}

secretion, whereas knockdown of syntaxin 3 (STX3) had no discernible effect. In

addition, we explored the involvement of SNAP23 and SNAP29, known regulators of

vesicle-plasma membrane fusion. Knockdown experiments revealed that depletion of

SNAP23 significantly impaired UBB^{+1} secretion, like the effect observed with STX4

knockdown, whereas SNAP29 depletion showed no significant effect (**Fig. 1D**). While

some autophagosomes target UBB^{+1} for lysosomal degradation, others are secreted

through an unconventional autophagy-related pathway mediated by a SNARE complex

including SEC22B, STX4, and SNAP23, facilitating the fusion of UBB^{+1}-loaded vesicles

with the plasma membrane (**Fig. 6**).

This work identifies UBB^1 as a novel cargo for secretory autophagy specifically targeted

through its ubiquitin domain. UBB^{+1} contains an internal, non-cleavable ubiquitin domain

that is integral to its structure. In this study, we provide initial evidence suggesting that

this ubiquitin domain participates in the secretion of UBB^{+1} and may contribute to its

recognition for secretory autophagy. Specifically, we demonstrate that UBB^{+1}

colocalizes with the autophagy receptor SQSTM1/p62 and that SQSTM1/p62 physically

associates with UBB^{+1} through its ubiquitin domain. Furthermore, disruption of the

hydrophobic patch within this ubiquitin domain significantly impairs SQSTM1/p62

association and inhibits UBB^{+1} secretion. Together, these findings support the idea that

the internal ubiquitin domain of UBB^{+1} plays an important role in its selective recruitment

into the secretory autophagy pathway.

In summary, our study exposes a dual role of autophagy in both intracellular protein

quality control and a distinct pathway for UBB^{+1} secretion mediated by SQSTM1/p62

and SEC22B. Secretion appears to represent an active strategy for alleviating UBB^{+1}-

induced cellular stress. While ubiquitin is traditionally viewed as a degradation signal,

this work highlights a unique example of a ubiquitin-related protein driving secretory

autophagy. This process represents a specialized form of secretory autophagy that

requires a ubiquitin receptor, yet bypasses lysosomal degradation, utilizing the

SEC22B-STX4/SNAP23 SNARE complexes for vesicle-plasma membrane fusion.

These findings provide new insights into how cells manage UBB^{+1} toxicity and highlight

the broader implications of secretory autophagy in proteostasis, intercellular

communication, and neurodegenerative disease progression. Future studies could

focus on identifying additional factors regulating this pathway and exploring the

extracellular ramifications of UBB^{+1} in health and disease.

Secretion raises intriguing questions about the fate of UBB^{+1} in the extracellular

environment. While this mechanism may initially benefit the cell by alleviating

intracellular stress, the extracellular propagation of UBB^{+1} could have pathological

consequences. The potential involvement of UBB^{+1} in extracellular amyloid plaque

formation or spreading of toxic effects warrants further investigation.

Material and methods:

Chemicals and Antibodies:

DL-dithiothreitol (DTT, D0632), iodoacetamide (I1149), IGEPAL CA-630 (NP40, I3021),

bovine serum albumin (BSA, A7906), sodium dodecyl sulfate (SDS, L3771), Tween-20

(P9416), ethylenediaminetetraacetic acid disodium salt dehydrate (EDTA, E4884),

bafilomycin A1 (B1793) and Velcade (504314) were purchased from Sigma-Aldrich.

TAK-243 (MLN7243) (S8341), Venetoclax (ABT-199; HY-15531) was purchased from

Selleckchem. Complete protease inhibitor cocktail (04693116001) was purchased from

Roche Diagnostic.

Antibodies and their manufacturers were: rabbit anti-UBB^{+1} (WB 1:1000, IF 1:250

custom made by Sigma for our lab) (25, 31). Rabbit anti-GAPDH (WB 1:10,000,

G9545), mouse anti-HA tag (WB 1:2000, IF 1:250, H3663) were from Sigma-Aldrich.

Mouse anti-CLUSTERIN (WB 1:1000; 166907), mouse anti-UBIQUITIN (WB 1:1000,

8017), mouse anti-ATG5 (WB 1:1000, 133158), mouse anti-SEC22B (WB 1:1000,

101267), mouse anti-ACTIN (WB 1:1000, 47778) and mouse anti-TSG101 (WB 1:1000,

7964), mouse anti-SNAP23 (WB 1:1000, 166244), mouse anti-SNAP29 (WB 1:1000,

390602) were from Santa Cruz Biotechnology. Rabbit anti-LC3 (WB: 1:2000, 51520)

was from Abcam. Mouse anti-SQSTM1/p62 (WB 1:1000, IF, 1:250 88588), mouse anti-

STX4 (WB 1:1000, E6W7B), mouse anti-STX17 (WB 1:1000, D3D7H), rabbit anti-

cleaved caspase 3 (WB 1:1000, 9661) were from Cell Signaling Technology. Rabbit

anti-STX3 (WB 1:1000, Synaptic systems, 110032). Alexa Fluor 488- and 594-

conjugated secondary antibodies were from Thermo Fisher (A21206 and A11032

respectively). Peroxidase-conjugated anti-mouse (WB 1:10,1000, 115-005-003 and anti-

rabbit secondary antibodies (WB 1:10,000; 111-005-144) were from Jackson lab.

Cell lines, plasmid, and transfection

HeLa cells (ATCC RR-B51S) were routinely grown at 37 ℃ in Dulbecco's modified

essential media (DMEM, Sigma-Aldrich, D6429), supplemented with 10% FBS (Sigma,

F7524), 100 U/ml Penicillin/Streptomycin, 2 mM L-Glutamine (L-Gln). Triple

(optn/*ndp52*/*tax1bp1)* and penta (*tax1bp1*/*optn*/*ndp52*/*nbr1*/*p62)* knockout (KO) HeLa

cell lines were kindly provided by Prof. Richard Youle (NINDS, NIH, USA). The

individual knockouts of *taxbp1*, *sqstm1/p62* and *sec22b* in HeLa cell lines were

generated by CRISPR-Cas9 approach. We used the following targeting sequence to

design guide RNAs:

SEC22B: 5' – CACCGGAGTGCAAATCTTCTAGGT – 3'.

TAX1BP1: 5' – CACCGTAGTGGTGACCACAAAAGC – 3',

SQSTM1/p62: 5' – CACCGCCGAATCTACATTAAAGAGAA – 3' and

The guide RNA oligonucleotides were annealed and ligated to BpiI digested pSp-Cas9-

BB-2A-Puro (PX459) V 2.0 vector (a gift from Feng Zhang; Addgene Plasmid #62988).

Cells were transfected with respective plasmids using Lipofectamine-3000 (Thermo

Fischer, L3000001) according to the manufacturer's instructions followed by selection

with Puromycin (2-3 μg/ml). The individual clones obtained after selection were

expanded to screen them for the expression by immunoblotting using specific

antibodies.

Plasmids encoding MYC-tagged UBB^{+1} was described previously (31). SQSTM1/p62

with C-terminal HA tag (HA-SQSTM1/p62, 452 residues) was obtained from addgene

(28027, deposited by Qing Zhong). The coding sequence of the SQSTM1/p62 without

PB1 (HA-SQSTM1/p62ΔPB1, 330 residues) domain was subcloned from the full-length

HA-tagged SQSTM1/p62 plasmid.

PCR-based site-directed mutagenesis (SDM) was used to generate specific point

mutations in the full-length HA-tagged SQSTM1/p62 plasmid. To disrupt the LC3-

interacting region (LIR), tryptophan at position 340 and leucine at position 343 were

substituted with alanine, generating the HA-SQSTM1/p62 W340A/L343A mutant. The

primer pair used for this mutation was:

Forward: 5′-GGAGATGATGACGCGACCCATGCGTCTTCAAAAG-3′ and

Reverse: 5′-CTTTTGAAGACGCATGGGTCGCGTCATCATCTCC-3′.

Additionally, to disrupt the ubiquitin-binding function of the UBA domain, methionine at

position 404 was substituted with valine (M404V) in the same plasmid using the

following primer pair:

Forward: 5′-CAGATGCTGTCCGTGGGCTTCTCTG-3′ and

Reverse: 5′-CAGAGAAGCCCACGGACAGCATCTG-3′.

All site-directed mutagenesis reactions were performed using high-fidelity polymerase

and verified by DNA sequencing.

Protein secretion experiments.

To measure UBB^{+1} secretion, cells (1×10^5) were seeded and cultured to ~70–80% confluency. Cells were transfected with MYC-UBB^{+1} along with the indicated control or effector plasmids. After 24 h, the culture medium was replaced with 1.5 ml of fresh complete DMEM. Cells were incubated for an additional 16 h, and the conditioned media were then collected. Media were sequentially centrifuged, first at $1,000 \times g$ for 5 min to remove detached cells, and then at $10,000 \times g$ for 30 min to eliminate debris. Equal volumes of cleared media and corresponding whole-cell lysates (WCL) were resolved by SDS-PAGE and analyzed by immunoblotting. Secretion efficiency was assessed by normalizing the UBB^{+1} signal in the media to its expression level in the WCL.

Immunoblotting and immunoprecipitation

For immunoblotting and co-immunoprecipitation, cells harvested 24 h or 48 h post transfection were lysed in NP40 lysis buffer (50 mM Tris-HCl pH 7.6, 150 mM NaCl, 5 mM MgCl2, 1 mM EDTA, 1 mM DTT, 1% NP-40, 5% glycerol) supplemented with protease and deubiquitinase inhibitors (20 mM Iodoacetamide). Protein concentration was measured using Bradford reagent (Thermo Scientific). The insoluble fraction consisting of the remaining pellet was washed once with NP-40 lysis buffer, spun at $17,000 \times g$ for 10 min at 4°C, then lysed in 1 X SDS sample buffer and heated to 95°C for 10 min. Immunoblotting was performed according to a previously described procedure (31, 64).

For co-immunoprecipitation of overexpressed MYC-UBB^{+1} and HA-p62 variants, Cells

were co-transfected with the indicated plasmids for 24 h. Subsequently, they were

washed three times with ice-cold PBS (Sigma, D8537) and lysed in 500 µl of ice-cold

NP40 lysis buffer. Cell lysates were incubated for 2 h with anti-MYC magnetic beads

(Bimake, B26301) and anti-HA magnetic beads (MedChemExpress, HY-K0201).

Precipitated complexes were washed with lysis buffer and eluted with MYC and HA

peptides (GenScript, RP11731 and RP11735S), respectively at a concentration of 400

µg/ml.

For co-immunoprecipitation of endogenous SQSTM1/p62, the cell lysates were

incubated for 2–4 h with specific antibody followed by 1–2 h with protein-A/G plus

coupled Sepharose beads (GE Healthcare, 17–0885-01). For control experiments,

lysates were also incubated with protein A/G beads alone (without antibody) to assess

non-specific binding to the beads. To resolve protein complexes for denaturing

immunoprecipitation, cell pellets were lysed in 50 µl of ice-cold NP40 lysis buffer

supplemented with 1% SDS. Immunocomplexes were washed with NP-40 lysis buffer

and elution was performed by boiling for 5 min in 2x SDS-PAGE loading buffer. Equal

amounts of proteins were fractionated in a SurePAGE^{TM}, Bis-Tris 4–20% gradient gels

(GenScript, M00655). After transferring to nitrocellulose membranes (Millipore), the

blots were blocked in Tris-buffered saline containing 5% non-fat milk and 0.1% Tween-

20 and incubated with primary antibodies either for 1 h at room temperature or overnight

at 4°C followed by incubation for 1 h with the appropriate horseradish peroxidase-

conjugated secondary antibodies. Western blot images were visualized by enhanced

chemiluminescence (ECL).

Immunofluorescence and confocal microscopy

HeLa cells were grown to semi-confluency on glass cover slips and transfected with the indicated plasmids using the Lipofectamine 3000 kit as recommended by the manufacturers. After treatment for the indicated time the cells were fixed in 4% paraformaldehyde (Merck, 100496) and permeabilized with 0.1% Triton X-100 in phosphate buffer saline (PBS) for 5 min RT. Blocking in 5% bovine serum albumin (BSA) (Sigma, A7906) in PBS for 30 min at RT was performed before antibody labeling. The cells were labeled in 5% BSA-PBS using rabbit anti-UBB[+1], mouse anti-HA, mouse anti-SQSTM1/p62, mouse anti-MYC and rabbit anti-LC3 followed by the appropriate Alexa Fluor 488 or 594 conjugated secondary antibodies. The nuclei were stained with 2.5 µg/ml DAPI (Sigma, D9542) in PBS for 10 min. The coverslips were again washed with PBS thrice and finally mounted with Fluoromount G (Thermo) on glass slides and stored at −20 °C until image acquisition. Images were acquired using Zen blue software on an LSM 880 confocal microscope system (ZEISS) with a Plan Apochromat X63 1.4 numerical aperture objective and oil immersion. The colocalization analysis was performed by using the Profile analysis tool in the ZEN lite (V 3.0) software (Zeiss) by fluorescence intensity line measurement approach.

Gene silencing by RNA interference

All gene specific DsiRNA oligonucleotides were purchased from Integrated DNA Technologies. DsiRNA was transfected with lipofectamine-3000 transfection reagent (Thermo Fischer, L3000001) at a final concentration of 25 nM. Following are DsiRNA constructs used in this study (**Table 1**),

Gene	Accession number	DsiRNA (IDT)
Human Stx3	NM_004177.5	hs.Ri.STX3.13.1,
Human Stx4	NM_001272096	hs.Ri.STX4.13.2
Human SNAP23	NM_003825	hs.Ri.SNAP23.13
Human SNAP29	NM_004782	hs.Ri.SNAP29.13
Human Stx17	NM_017919.3	hs.Ri.STX17.13.2
Human TSG101	NM_000011.10	hs.Ri.TSG101.13.2

Statistical analysis

All statistical analyses were performed using GraphPad Prism (version 5.0). Data are

presented as mean ± standard deviation (SD) as indicated in figure legends. For

comparisons between two groups, an unpaired two-tailed Student's t-test was used. For

multiple group comparisons, such as the quantification of p62 condensate volume and

number per cell, statistical significance was assessed using one-way ANOVA followed

by Tukey's multiple comparisons test. A p-value > 0.05 was considered not significant

(ns). Statistical significance is indicated as follows: *p < 0.05, **p < 0.01, and ***p <

0.001.

AlphaFold-based modeling of UBB^{+1}–p62 UBA interaction:

Protein sequences of full length UBB^{+1} and the UBA domain of human SQSTM1/p62

(residues 387–436) were submitted to the AlphaFold3-based ColabFold pipeline (65) for

structural prediction of a protein-protein complex. The top-ranked predicted model

(based on the predicted alignment error and pLDDT scores) was selected for

visualization. Structural images were rendered using PyMOL (v2.5.4) Color coding was

applied as follows: UBB^{+1} in green, its C-terminal tail in red, the p62 UBA domain in

magenta, Met404 in orange, and the hydrophobic patch residues of UBB^{+1} (Ile44, Val70,

and Leu8) in blue. Representative structural snapshots were rendered and included as

Supplementary Fig. S5D.

5. Acknowledgments and funding sources

Acknowledgments: We sincerely thank Prof. Richard Youle and his team for kindly

providing us with the penta knockout (5KO) and triple knockout (TKO) HeLa cell lines.

We thank Dr. Nitsan Dahan and all members of the Microscopy and sorting units of the

LS&E institute at the Technion. The authors thank Dr. Inbal Maniv, Noa Reis, and all the

Glickman lab members for their intellectual input and critical feedback on the

manuscript.

Funding: This work was partially funded by ISF BRG grant 2640/23, and by the

European Union ERC grant (UbWan, 101142726) to MHG. Views and opinions

expressed are however those of the author(s) only and do not necessarily reflect those

of the European Union or the European Research Council Executive Agency. Neither

the European Union nor the granting authority can be held responsible for them. We

select open access license.

Abbreviations: AD, Alzheimer disease; CHX, cycloheximide; DsiRNAs: dicer-substrate

small interfering RNA; KO, knockout; 5 KO, penta knockout; TKO, triple knockout LC3,

microtubule associated protein 1 light chain 3; SNARE, SNAP soluble NSF attachment

protein receptor; SQSTM1/p62, sequestosome 1; STX, syntaxin; TAX1BP1, Tax1

binding protein 1; WB, Western blot; WT, wild type

645
646   **Disclosure of competing interests:** The authors declare no potential conflicts of

647   interest.

648   **Data Sharing:** All relevant data is included with the submission.

649   **Author contributions:** ARW conducted all experiments. MHG obtained funding and

650   supervised. Both authors wrote the manuscript.

6. References:

1. J. Kaur, J. Debnath, Autophagy at the crossroads of catabolism and anabolism. *Nat Rev Mol Cell Biol* **16**, 461-472 (2015).
2. H. Guo *et al.*, Atg5 Disassociates the V(1)V(0)-ATPase to Promote Exosome Production and Tumor Metastasis Independent of Canonical Macroautophagy. *Dev Cell* **43**, 716-730 e717 (2017).
3. M. Zhang, S. J. Kenny, L. Ge, K. Xu, R. Schekman, Translocation of interleukin-1beta into a vesicle intermediate in autophagy-mediated secretion. *Elife* **4** (2015).
4. S. Bel *et al.*, Paneth cells secrete lysozyme via secretory autophagy during bacterial infection of the intestine. *Science* **357**, 1047-1052 (2017).
5. M. Ponpuak *et al.*, Secretory autophagy. *Curr Opin Cell Biol* **35**, 106-116 (2015).
6. S. Jiang, N. Dupont, E. F. Castillo, V. Deretic, Secretory versus degradative autophagy: unconventional secretion of inflammatory mediators. *J Innate Immun* **5**, 471-479 (2013).
7. Z. Xie, D. J. Klionsky, Autophagosome formation: core machinery and adaptations. *Nat Cell Biol* **9**, 1102-1109 (2007).
8. V. Kirkin, D. G. McEwan, I. Novak, I. Dikic, A role for ubiquitin in selective autophagy. *Mol Cell* **34**, 259-269 (2009).
9. T. Lamark, V. Kirkin, I. Dikic, T. Johansen, NBR1 and p62 as cargo receptors for selective autophagy of ubiquitinated targets. *Cell Cycle* **8**, 1986-1990 (2009).
10. S. Pankiv *et al.*, p62/SQSTM1 binds directly to Atg8/LC3 to facilitate degradation of ubiquitinated protein aggregates by autophagy. *J Biol Chem* **282**, 24131-24145 (2007).
11. M. Komatsu *et al.*, Homeostatic levels of p62 control cytoplasmic inclusion body formation in autophagy-deficient mice. *Cell* **131**, 1149-1163 (2007).
12. K. Hofmann, P. Bucher, The UBA domain: a sequence motif present in multiple enzyme classes of the ubiquitination pathway. *Trends Biochem Sci* **21**, 172-173 (1996).
13. J. Schultz, F. Milpetz, P. Bork, C. P. Ponting, SMART, a simple modular architecture research tool: identification of signaling domains. *Proc Natl Acad Sci U S A* **95**, 5857-5864 (1998).
14. M. L. Seibenhener *et al.*, Sequestosome 1/p62 is a polyubiquitin chain binding protein involved in ubiquitin proteasome degradation. *Mol Cell Biol* **24**, 8055-8068 (2004).
15. J. Long *et al.*, Ubiquitin recognition by the ubiquitin-associated domain of p62 involves a novel conformational switch. *J Biol Chem* **283**, 5427-5440 (2008).
16. H. B. Yu *et al.*, Sequestosome-1/p62 is the key intracellular target of innate defense regulator peptide. *J Biol Chem* **284**, 36007-36011 (2009).
17. J. H. Son, J. H. Shim, K. H. Kim, J. Y. Ha, J. Y. Han, Neuronal autophagy and neurodegenerative diseases. *Exp Mol Med* **44**, 89-98 (2012).
18. C. R. Wilkinson *et al.*, Proteins containing the UBA domain are able to bind to multi-ubiquitin chains. *Nat Cell Biol* **3**, 939-943 (2001).
19. A. K. H. Stavoe, E. L. F. Holzbaur, Autophagy in Neurons. *Annu Rev Cell Dev Biol* **35**, 477-500 (2019).
20. A. Kulkarni, J. Chen, S. Maday, Neuronal autophagy and intercellular regulation of homeostasis in the brain. *Curr Opin Neurobiol* **51**, 29-36 (2018).
21. Y. Liang, S. Sigrist, Autophagy and proteostasis in the control of synapse aging and disease. *Curr Opin Neurobiol* **48**, 113-121 (2018).
22. D. Azarnia Tehran, M. Kuijpers, V. Haucke, Presynaptic endocytic factors in autophagy and neurodegeneration. *Curr Opin Neurobiol* **48**, 153-159 (2018).

23.   P. Cras *et al.*, Senile plaque neurites in Alzheimer disease accumulate amyloid precursor protein.
Proc Natl Acad Sci U S A **88**, 7552-7556 (1991).

24.   G. K. Gouras, T. T. Olsson, O. Hansson, beta-Amyloid peptides and amyloid plaques in
Alzheimer's disease. *Neurotherapeutics* **12**, 3-11 (2015).

25.   I. Maniv *et al.*, Altered ubiquitin signaling induces Alzheimer's disease-like hallmarks in a three-
dimensional human neural cell culture model. *Nat Commun* **14**, 5922 (2023).

26.   F. W. van Leeuwen *et al.*, Frameshift mutants of beta amyloid precursor protein and ubiquitin-B
in Alzheimer's and Down patients. *Science* **279**, 242-247 (1998).

27.   K. Banasiak, N. A. Szulc, W. Pokrzywa, The Dose-Dependent Pleiotropic Effects of the UBB(+1)
Ubiquitin Mutant. *Front Mol Biosci* **8**, 650730 (2021).

28.   S. Oddo, The ubiquitin-proteasome system in Alzheimer's disease. *J Cell Mol Med* **12**, 363-373
(2008).

29.   R. J. Gentier *et al.*, Localization of mutant ubiquitin in the brain of a transgenic mouse line with
proteasomal inhibition and its validation at specific sites in Alzheimer's disease. *Front Neuroanat*
9, 26 (2015).

30.   M. Irmler *et al.*, Long-term proteasomal inhibition in transgenic mice by UBB(+1) expression
results in dysfunction of central respiration control reminiscent of brainstem neuropathology in
Alzheimer patients. *Acta Neuropathol* **124**, 187-197 (2012).

31.   A. R. Wagh, P. Sulakshane, M. H. Glickman, Alzheimer's disease-associated mutant ubiquitin
(UBB(+1)) is secreted through an autophagosome-like vesicle-mediated unconventional
pathway. *Biochim Biophys Acta Gene Regul Mech* **1866**, 194936 (2023).

32.   W. Nickel, Pathways of unconventional protein secretion. *Curr Opin Biotechnol* **21**, 621-626
(2010).

33.   W. Nickel, C. Rabouille, Mechanisms of regulated unconventional protein secretion. *Nat Rev Mol*
Cell Biol **10**, 148-155 (2009).

34.   W. Hong, SNAREs and traffic. *Biochim Biophys Acta* **1744**, 120-144 (2005).

35.   T. Kimura *et al.*, Dedicated SNAREs and specialized TRIM cargo receptors mediate secretory
autophagy. *EMBO J* **36**, 42-60 (2017).

36.   M. Renna *et al.*, Autophagic substrate clearance requires activity of the syntaxin-5 SNARE
complex. *J Cell Sci* **124**, 469-482 (2011).

37.   A. J. Souers *et al.*, ABT-199, a potent and selective BCL-2 inhibitor, achieves antitumor activity
while sparing platelets. *Nat Med* **19**, 202-208 (2013).

38.   I. Cebrian *et al.*, Sec22b regulates phagosomal maturation and antigen crosspresentation by
dendritic cells. *Cell* **147**, 1355-1368 (2011).

39.   K. Arasaki, C. R. Roy, Legionella pneumophila promotes functional interactions between plasma
membrane syntaxins and Sec22b. *Traffic* **11**, 587-600 (2010).

40.   E. Itakura, C. Kishi-Itakura, N. Mizushima, The hairpin-type tail-anchored SNARE syntaxin 17
targets to autophagosomes for fusion with endosomes/lysosomes. *Cell* **151**, 1256-1269 (2012).

41.   S. Takats *et al.*, Autophagosomal Syntaxin17-dependent lysosomal degradation maintains
neuronal function in Drosophila. *J Cell Biol* **201**, 531-539 (2013).

42.   I. Koyama-Honda, N. Mizushima, Transient visit of STX17 (syntaxin 17) to autophagosomes.
Autophagy **18**, 1213-1215 (2022).

43.   R. Trapannone, J. Romanov, S. Martens, p62 and NBR1 functions are dispensable for aggrephagy
in mouse ESCs and ESC-derived neurons. *Life Sci Alliance* **6** (2023).

44.   G. Zaffagnini *et al.*, p62 filaments capture and present ubiquitinated cargos for autophagy.
EMBO J **37** (2018).

45.   B. Wurzer *et al.*, Oligomerization of p62 allows for selection of ubiquitinated cargo and isolation
membrane during selective autophagy. *Elife* **4**, e08941 (2015).

46. S. Berkamp, S. Mostafavi, C. Sachse, Structure and function of p62/SQSTM1 in the emerging
framework of phase separation. *FEBS J* **288**, 6927-6941 (2021).

47. R. K. Singh, Y. Kazansky, D. Wathieu, D. Fushman, Hydrophobic Patch of Ubiquitin is Important
for its Optimal Activation by Ubiquitin Activating Enzyme E1. *Anal Chem* **89**, 7852-7860 (2017).

48. M. Cabe, D. J. Rademacher, A. B. Karlsson, S. Cherukuri, J. C. Bakowska, PB1 and UBA domains of
p62 are essential for aggresome-like induced structure formation. *Biochem Biophys Res*
Commun **503**, 2306-2311 (2018).

49. T. Johansen, T. Lamark, Selective autophagy mediated by autophagic adapter proteins.
Autophagy **7**, 279-296 (2011).

50. R. K. Vadlamudi, I. Joung, J. L. Strominger, J. Shin, p62, a phosphotyrosine-independent ligand of
the SH2 domain of p56lck, belongs to a new class of ubiquitin-binding proteins. *J Biol Chem* **271**,
20235-20237 (1996).

51. X. Wang, E. J. Terpstra, Ubiquitin receptors and protein quality control. *J Mol Cell Cardiol* **55**, 73-
84 (2013).

52. K. Zientara-Rytter, S. Subramani, The Roles of Ubiquitin-Binding Protein Shuttles in the
Degradative Fate of Ubiquitinated Proteins in the Ubiquitin-Proteasome System and Autophagy.
Cells **8** (2019).

53. Y. Ichimura *et al.*, Structural basis for sorting mechanism of p62 in selective autophagy. *J Biol*
Chem **283**, 22847-22857 (2008).

54. R. Layfield *et al.*, Structural and functional studies of mutations affecting the UBA domain of
SQSTM1 (p62) which cause Paget's disease of bone. *Biochem Soc Trans* **32**, 728-730 (2004).

55. J. R. Cavey *et al.*, Loss of ubiquitin-binding associated with Paget's disease of bone p62
(SQSTM1) mutations. *J Bone Miner Res* **20**, 619-624 (2005).

56. E. R. Barrow *et al.*, Discovery of SQSTM1/p62-dependent P-bodies that regulate the NLRP3
inflammasome. *Cell Rep* **43**, 113935 (2024).

57. D. Krutauz *et al.*, Extended ubiquitin species are protein-based DUB inhibitors. *Nat Chem Biol* **10**,
664-670 (2014).

58. P. van Tijn *et al.*, Dose-dependent inhibition of proteasome activity by a mutant ubiquitin
associated with neurodegenerative disease. *J Cell Sci* **120**, 1615-1623 (2007).

59. K. Lindsten *et al.*, Mutant ubiquitin found in neurodegenerative disorders is a ubiquitin fusion
degradation substrate that blocks proteasomal degradation. *J Cell Biol* **157**, 417-427 (2002).

60. B. M. Verheijen, R. J. G. Gentier, D. Hermes, F. W. van Leeuwen, D. A. Hopkins, Selective
Transgenic Expression of Mutant Ubiquitin in Purkinje Cell Stripes in the Cerebellum. *Cerebellum*
16, 746-750 (2017).

61. G. Bjorkoy *et al.*, p62/SQSTM1 forms protein aggregates degraded by autophagy and has a
protective effect on huntingtin-induced cell death. *J Cell Biol* **171**, 603-614 (2005).

62. Y. Nakamura, T. Sawai, K. Kakiuchi, S. Arawaka, Neuronal activity promotes secretory autophagy
for the extracellular release of alpha-synuclein. *J Biol Chem* **300**, 107419 (2024).

63. X. Chen, A. J. Munoz-Arellano, D. Petranovic, UBB(+1) reduces amyloid-beta cytotoxicity by
activation of autophagy in yeast. *Aging (Albany NY)* **13**, 23953-23980 (2021).

64. I. Sahu *et al.*, The 20S as a stand-alone proteasome in cells can degrade the ubiquitin tag. *Nat*
Commun **12**, 6173 (2021).

65. M. Mirdita *et al.*, ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679-682
(2022).

7. Figure legends:

Figure 1. Q- and R-SNAREs involved in UBB^{+1} secretion.

(**A**) Wild-type (WT) and *sec22b* knockout (KO) HeLa cells were transiently transfected with a MYC-UBB^{+1} expressing plasmid or treated with transfection reagent alone (mock control). After 24 h, cells were incubated with fresh DMEM for an additional 16 h. Conditioned media and whole-cell lysates (WCL) were collected, resolved by 15% SDS-PAGE, and immunoblotted with specific antibodies. UBB^{+1} and LC3 were detected to assess secretion and autophagy induction, respectively, GAPDH was used as a loading control for WCL. Secreted UBB^{+1} in the media was detected using anti-UBB^{+1} antibody. Equal volumes of media were collected from cultures of comparable cell numbers across all conditions to ensure consistency in secretion analysis. Band intensities of LC3-II were quantified relative to GAPDH and are indicated below the blots.

(**B**) Intracellular distribution of UBB^{+1} (Green) and LC3 (Red) was visualized by confocal immunofluorescence microscopy using specific antibodies. Nuclei were counterstained with DAPI (blue). In wild-type HeLa cells expressing MYC-UBB^{+1}, LC3-positive autophagic vesicles primarily accumulated in the juxtanuclear region with limited colocalization with UBB^{+1} puncta. In contrast, *sec22b* KO cells displayed prominent colocalization of UBB^{+1} with LC3-positive autophagosomes, consistent with impaired autophagosome maturation and reduced vesicle clearance. Scale bar: 5 µm. Quantification of UBB^{+1}–LC3 colocalized puncta was performed by analyzing 100 randomly selected cells per condition from three independent experiments. Bar graphs

show mean ± SD. Statistical significance was determined using an unpaired two-tailed

Student's t-test; *, **, and *** denote p < 0.05, 0.01, and 0.001, respectively; ns = non-

significant.

(**C**) WT and *sec22b* KO HeLa cells expressing MYC-UBB[+1] were incubated with 100

µg/ml cycloheximide (CHX) for the indicated time periods. Conditioned media were

collected in equal volumes from comparable cell numbers across all experimental

conditions and immunoblotted for UBB[+1] (top). Densitometric analysis of UBB[+1]

secretion (bottom) was performed by quantifying the UBB[+1] band intensity. For each

time point, UBB[+1] levels in *sec22b* KO cells were compared to WT cells. Statistical

comparisons between WT and *sec22b* KO are indicated in the graph: comparisons at 0

h are shown by round dotted lines, at 4 h by solid lines, and at 8 h by square dotted

lines. Data represents the mean ± SD from three independent experiments. Statistical

significance was determined using an unpaired two-tailed t-test. *, **, and *** denote p <

0.05, 0.01, and 0.001, respectively; ns = non-significant. In the bar graph, *sec22b* KO

data are shown in [gray], and WT data are shown in [black].

(**D**) HeLa cells expressing MYC-UBB[+1] were transfected with 25 nM DsiRNAs targeting

STX3, STX4, SNAP23, or SNAP29, or with a non-targeting (NT) control (see Table 1 for

sequences). Forty-eight hours after transfection, cells were incubated in fresh media for

16 h. Conditioned media and corresponding whole-cell lysates (WCLs) were collected

and analyzed by SDS-PAGE followed by immunoblotting. Seven blots show: STX3,

STX4, SNAP23, and SNAP29 knockdown efficiency, GAPDH as the WCL loading

control, UBB[+1] levels in WCLs, and secreted UBB[+1] in the media. Conditioned media

were collected in equal volumes from comparable cell numbers across all experimental

conditions and immunoblotted for UBB^{+1}. STX4 and SNAP23 knockdown significantly

reduced UBB^{+1} secretion, whereas STX3 and SNAP29 knockdown had no notable

effect. Data are representative of three independent experiments; quantification and

statistical analysis were performed using one-way ANOVA. *, **, and *** denote p <

0.05, 0.01, and 0.001, respectively; ns = non-significant.

(**E**) WT and *sec22b* KO HeLa cells expressing MYC-UBB^{+1} were incubated with 100

µg/ml cycloheximide (CHX) for the indicated periods. The cells were then harvested,

and WCL immunoblotted for UBB^{+1} (top). Densitometric analysis of the UBB^{+1} relative to

untreated cells (normalized to GAPDH content) was averaged from three independent

experiments (bottom). The error bars represent the means ± SD. Statistical significance

was determined using an unpaired two-tailed t-test. For all statistical tests *, **,

***, *p* < 0.05, 0.01, and 0.001, respectively, ns = non-significant.

Figure 2. Turnover and Secretion Dynamics of UBB^{+1} Under Cycloheximide

Treatment".

(**A and B**) WT HeLa cells expressing MYC-UBB^{+1} were treated with 100 µg/mL

cycloheximide (CHX) alone or in combination with Bafilomycin A1 (100 nM) or Velcade

(100 nM) for the indicated time points (0, 4, and 8 h). Whole cell lysates (WCL) **(A)** and

conditioned media **(B)** were collected and immunoblotted for UBB^{+1}. Equal volumes of

conditioned media were harvested from comparable cell numbers across all conditions.

Representative UBB^{+1} blots are shown. Densitometric quantification of UBB^{+1} levels

from three independent experiments was performed.

For WCL (A), UBB^{+1} band intensities were normalized to GAPDH and expressed as

percentage degradation relative to the 0 h time point (set as 100%).

For conditioned media (B), UBB^{+1} band intensities were quantified and plotted as

secreted levels. Statistical comparisons were performed between CHX-alone and each

of the inhibitor-treated conditions (CHX + Bafilomycin A1 or CHX + Velcade) at each

time point. Comparisons at 0 h are indicated by round dotted lines, at 4 h by solid lines,

and at 8 h by square dotted lines in the graph. Bar graphs represent mean ± SD from

three biologically independent experiments. Statistical significance was determined

using one-way ANOVA followed by Tukey's multiple comparisons test. For all statistical

tests: *, **, and ***, indicate $p < 0.05$, 0.01, and 0.001, respectively; ns = non-significant.

(C–F) Conditioned media **(C, D)** and whole-cell lysates (WCL; **E, F**) were collected from

wild-type (WT) and *sec22b* KO HeLa cells expressing MYC-UBB^{+1} after treatment with

100 µg/mL cycloheximide (CHX) alone or in combination with Bafilomycin A1 (100 nM)

for the indicated time points (0 h and 8 h). Equal volumes of conditioned media were

harvested from comparable cell numbers across all conditions and immunoblotted for

UBB^{+1}. For WCL (**E, F**), GAPDH was used as a loading control. UBB^{+1} band intensities

were normalized to GAPDH and expressed as a percentage of the 0 h time point (set to

100%). Representative immunoblots and densitometric quantification from three

independent biological replicates are shown. For conditioned media (**C, D**), statistical

comparisons were made between WT and *sec22b* KO cells for each of the following

conditions: CHX 0 h (round dotted lines), CHX 8 h (solid lines), CHX + Bafilomycin A1

(8 h) (square dotted lines). Bar graphs represent, mean ± SD from three independent

experiments. Statistical significance was assessed using an unpaired two-tailed t-test to

compare WT and *sec22b* KO samples at each respective time point. Asterisks denote

significance thresholds: *, **, and *** correspond to $p < 0.05$, $p < 0.01$, and $p < 0.001$,

respectively; ns = not significant. In the graphs, WT and *sec22b* KO data are shown in

black and gray, respectively.

Figure 3. SQSTM1/p62 knockout hinders the efficient clearance of UBB[+1]

(**A-E**) Wild type (WT), *TAX1BP1/OPTN/NDP52/NBR1/p62* penta knockout (5KO),

OPTN/NDP52/TAX1BP1 triple knockout (TKO), and *p62* KO HeLa cells were transiently

transfected with or without MYC-UBB[+1]. After 24 h, cells were incubated with fresh

DMEM for 16 h. Conditioned media and whole cell lysates (WCL) were resolved by 15%

SDS-PAGE and immunoblotted for UBB[+1]. GAPDH was used as the loading control for

WCL. Conditioned media were collected in equal volumes from comparable cell

numbers across all experimental conditions and immunoblotted for UBB[+1].

(**F**) Representative immunofluorescence staining of WT and *p62* KO HeLa cells

expressing MYC-UBB[+1]. Cells were fixed, permeabilized, and immunostained with a

monoclonal antibody against UBB[+1] (green), and nuclei were stained with DAPI (blue).

Scale bars: 5 μm. Statistical significance was determined by unpaired two-tailed t-test,

with data showing the mean ± SD from approximately 100 cells.

(**G**) Representative immunoblot of UBB[+1] in insoluble fractions harvested from WT and

p62 KO HeLa cells transfected with MYC-UBB[+1] plasmid. Cells were lysed with NP40-

lysis buffer, clarified at 15,000 x g and the pellet resuspended in 1% SDS. Ponceau

staining was used for loading control. Data represents the mean ± SD from at least

three independent experiments. For all statistical tests, *, **, ***, p < 0.05, 0.01, and

0.001, respectively.

Figure 4. Evidence of direct interaction between SQSTM1/p62 and UBB^{+1}

(A) Immunofluorescence staining of endogenous SQSTM1/p62 in HeLa cells

expressing MYC-UBB^{+1}. Cells were grown in complete media, fixed, permeabilized, and

stained with monoclonal antibodies for UBB^{+1} (green) and SQSTM1/p62 (red). Merged

images show colocalization, with nuclei stained using DAPI (blue). Scale bar: 2 μm.

(B) Immunofluorescence staining of overexpressed HA-p62 in HeLa cells co-expressing

MYC-UBB^{+1}. Cells were fixed, permeabilized, and stained with monoclonal antibodies

for UBB^{+1} (green) and HA (red). Merged images are shown, with nuclei counterstained

using DAPI (blue). Scale bar: 2 μm.

(C) Immunofluorescence staining of HeLa cells expressing HA-p62 and MYC-UBB$^{+1\ F4A,}$

^{I44A}. Superscript denotes point mutations (F4A and I44A) introduced into the UBB^{+1}

protein. UBB$^{+1\ F4A,\ I44A}$ showed no clear colocalization with HA-p62 condensates. Scale

bar: 2 μm.

Quantification of condensate volume per cell and condensate number per cell was

performed using Imaris-based image analysis. Data represents a minimum of 100 cells

per condition, pooled from three independent experiments. In both the condensate

volume and condensate number graphs, statistical comparisons were made between

the UBB^{+1}-only condition and each of the following: UBB^{+1} + HA-p62 and UBB$^{+1\ F4A,\ I44A}$

+ HA-p62, as indicated by solid lines. Bar graphs represent mean ± SD. Statistical

significance was determined using one-way ANOVA followed by Tukey's multiple

comparisons test. For all statistical tests: *, **, and ***, indicate $p < 0.05$, 0.01, and

0.001, respectively.

(D) Immunoblot analysis of conditioned media and whole cell lysates (WCL) from HeLa

cells expressing MYC-UBB^{+1} or the hydrophobic patch mutant MYC-UBB^{+1} $^{F4A, I44A}$.

Cells were transfected for 24 h and then incubated in fresh DMEM for 16 h. Conditioned

media and WCL were resolved by 15% SDS-PAGE and immunoblotted using specific

antibodies against UBB^{+1}. GAPDH was used as a loading control for WCL. Conditioned

media were collected in equal volumes from comparable cell numbers across all

experimental conditions to ensure consistent comparison of secreted protein levels.

(E) Co-immunoprecipitation (co-IP) of HA-tagged p62 with MYC-UBB^{+1} in HeLa cells.

Cells were co-transfected with HA-p62 and MYC-UBB^{+1}, and lysates were subjected to

immunoprecipitation under non-denaturing conditions using MYC- or FLAG-conjugated

magnetic beads. Bound proteins were eluted and analysed by immunoblotting (IB) with

anti-HA and anti-UBB^{+1} antibodies to assess interaction between p62 and UBB^{+1}.

(F) Co-immunoprecipitation of endogenous SQSTM1/p62 with MYC-UBB^{+1} in HeLa

cells. Lysates from cells expressing MYC-UBB^{+1} were immunoprecipitated using an

anti-p62 antibody under non-denaturing conditions. Immunoprecipitates were resolved

by SDS-PAGE and probed with anti-p62 and anti-UBB^{+1} antibodies. The anti-UBB^{+1}

antibody detected both ubiquitinated and non-ubiquitinated forms of UBB^{+1}, while the

immunoglobulin heavy chain is visible as a ~50 kDa band (indicated by an asterisk).

Negative controls included: (1) a beads-only (no antibody) IP control to evaluate non-

specific binding, and (2) IP from untransfected cells lacking MYC-UBB^{+1} expression to

assess background signal.

(G) Co-immunoprecipitation of endogenous SQSTM1/p62 with MYC-UBB^{+1} following

ubiquitination inhibition by TAK243. HeLa cells were transfected with MYC-UBB^{+1} and

treated with TAK243 (1 µM, 3 h) to inhibit ubiquitin-activating enzyme (E1) activity.

Lysates were immunoprecipitated using an anti-p62 antibody and analysed by

immunoblotting with anti-p62 and anti-UBB^{+1} antibodies. Both ubiquitinated and non-

ubiquitinated forms of UBB^{+1} were detected in the input; however, only the non-

ubiquitinated form co-precipitated with p62 following TAK243 treatment. The antibody

heavy chain (~50 kDa) is marked with an asterisk. Negative controls included beads-

only IP and IP from untransfected cells to rule out non-specific interactions.

(H) Immunofluorescence staining of HeLa cells co-expressing HA-p62 and MYC-UBB^{+1}

after 3 h of TAK243 treatment (same as in panel G). Cells were fixed, permeabilized,

and stained with monoclonal antibodies for UBB^{+1} (green) and HA (red). Nuclei were

counterstained using DAPI (blue). Merged images and zoomed insets show the

intracellular distribution and colocalization of p62 and UBB^{+1} following ubiquitination

blockade. Scale bar: 2 µm.

Quantification of p62 condensate volume and condensate number per cell was

performed from the same experimental set shown in panel (H), using Imaris image

analysis software. Data was obtained from a minimum of 100 cells per condition, pooled

from three independent experiments. Statistical comparisons were made between the

UBB^{+1} + HA-p62 WT and UBB^{+1} + HA-p62 WT + TAK243 conditions. In the graphs,

comparisons are indicated by square-dotted lines. Bar graphs represent mean ± SD.

Statistical significance was assessed using one-way ANOVA followed by Tukey's

multiple comparisons test. For all statistical tests: *, **, and ***, indicate $p < 0.05$, 0.01,

and 0.001, respectively.

Figure 5. SQSTM1/p62 domains required for functional association with UBB[+1].

(A) Schematic representation of functional domains in full-length SQSTM1/p62 (FL), the

ΔPB1 deletion mutant (lacking the N-terminal PB1 oligomerization domain; p62ΔPB1),

the LC3-interacting region (LIR) mutant (W340A, L343A; denoted p62 [W340A, L343A]), and

the UBA domain point mutant (p62 [M404V]).

(B) HeLa cells were transiently transfected with plasmids encoding C-terminally HA-

tagged SQSTM1/p62 constructs: FL, ΔPB1, WA/LA, and MV. After 24 h, whole-cell

lysates were prepared and subjected to SDS-PAGE followed by immunoblotting using

an anti-HA antibody to assess expression levels. All constructs were expressed at

comparable levels, confirming that functional differences observed in secretion and

localization assays (**Fig. 5**) are not due to unequal expression. The predicted molecular

weight of full-length p62 is 47 kDa; however, it migrates above 60 kDa due to post-

translational modifications. The truncated ΔPB1 construct has a calculated MW of 35.5

kDa, also appearing as a modified higher MW form.

(C) Immunofluorescence staining of HeLa cells co-expressing MYC-UBB[+1] with either

full-length HA-p62 (FL), HA-p62ΔPB1 (lacking the N-terminal PB1 oligomerization

domain), HA-p62 [W340A, L343A] (LC3-interacting region mutant; denoted WA/LA), or HA-

p62 ^{M404V} (UBA domain mutant defective in ubiquitin binding). Cells were fixed 24 h

post-transfection, permeabilized, and stained with monoclonal antibodies specific for HA

(red) and UBB^{+1} (green); nuclei were counterstained with DAPI (blue). The ΔPB1

mutant failed to form *p62-bodies* due to impaired oligomerization, while the M404V

mutant showed markedly reduced colocalization and condensate formation with UBB^{+1}

due to loss of UBA-mediated binding. In contrast, both FL and WA/LA constructs

retained the ability to form p62-positive condensates. Scale bar: 2 µm.

Right: Quantification of p62-positive condensates per cell under each condition. Data

represent mean ± SD from >100 cells per group pooled from three biologically

independent experiments. Statistical significance was determined using one-way

ANOVA followed by Tukey's multiple comparisons test. For all statistical tests: *p < 0.05,

**p < 0.01, ***p < 0.001.

(D–G) Immunoblot analysis of conditioned media and whole-cell lysates (WCL) from WT

and *p62* KO HeLa cells co-transfected with MYC-UBB^{+1} and either HA-p62ΔPB1 **(D)**,

HA-p62 *W340A, L343A* **(E)**, HA-p62 *M404V* **(F)**, or full-length HA-p62 (FL) **(G)**. After 24

h of transfection, cells were incubated with fresh DMEM for an additional 16 h.

Conditioned media and WCL were collected, resolved by 10% or 15% SDS-PAGE, and

immunoblotted with the indicated antibodies. GAPDH was used as a loading control for

WCL. Equal volumes of conditioned media were harvested from comparable cell

numbers across all experimental conditions. Representative immunoblots showing

UBB^{+1} levels in the media are displayed.

(H) Densitometric quantification of UBB^{+1} levels in the conditioned media across p62 KO

cells co-expressing MYC-UBB^{+1} with either full-length HA-p62 or the indicated p62

mutant constructs (ΔPB1, W340A/L343A, or M404V). Quantification is based on three

biologically independent experiments. Statistical comparisons were performed between

the full-length p62 condition and each of the mutant constructs. Bar graphs represent

mean ± SD. Statistical significance was assessed using one-way ANOVA followed by

Tukey's multiple comparisons test. For all statistical tests: *$p < 0.05$, **$p < 0.01$, ***$p <$

0.001; ns = non-significant.

(I) Co-immunoprecipitation (Co-IP) of HA-tagged SQSTM1/p62 constructs with MYC-

UBB^{+1} from HeLa cell lysates. Cells were co-transfected with MYC-UBB^{+1} and either

full-length HA-p62 FL or HA-p62ΔPB1. After 24 h, cells were lysed under non-

denaturing conditions, and immunoprecipitation was performed using anti-HA magnetic

beads. Elution was carried out using HA peptide. Immunoprecipitates were analyzed by

SDS-PAGE and immunoblotting (IB) with anti-HA and anti-UBB^{+1} antibodies.

(J) Co-immunoprecipitation of MYC-UBB^{+1} or MYC-UBB^{+1} mutants with HA-p62 or HA-

p62M404V from HeLa cell lysates. Lanes 1–4 show 2% input: lane 1, MYC-UB and HA-

p62 (positive control); lane 2, MYC-UBB^{+1} $^{F4A, I44A}$ mutant and HA-p62 (negative control);

lane 3, MYC-UBB^{+1} and HA-p62; lane 4, MYC-UBB^{+1} and HA-p62 ^{M404V}. Lanes 5–8

display HA immunoprecipitations using anti-HA magnetic beads: lanes 5–7 with HA-

p62; lane 8 with HA-p62 ^{M404V}. Lanes 9–10 serve as additional negative controls in

which lysates were incubated with anti-FLAG magnetic beads: lane 9, HA-p62 + MYC-

UBB^{+1}; lane 10, HA-p62M404V + MYC-UBB^{+1}. Immunoprecipitates were probed using

anti-MYC and anti-HA antibodies. The results confirm a specific interaction between

wild type p62 and UBB^{+1}, which is abolished when either the hydrophobic patch of

UBB^{+1} (F4A, I44A) or the Met404 residue in the UBA domain of p62 is mutated.

Figure 6. SQSTM1/p62 and the Cellular fates of UBB^{+1}.

SQSTM1/p62 critically affects both the intra- and extra-cellular fates of UBB^{+1}: (1)

Autophagic clearance facilitated by SQSTM1/p62 and LC3, leading to either lysosomal

degradation or extracellular expulsion via autophagosome-plasma membrane fusion

mediated by SEC22B and STX4/SNAP23. (2) Sequestration into p62-induced round

bodies when SQSTM1/p62 is elevated relative to levels of LC3. (3) Aggregation and

intracellular retention in the absence – or insufficient levels – of SQSTM1/p62.

Gene	Accession number	DsiRNA (IDT)
Human Stx3	NM_004177.5	hs.Ri.STX3.13.1,
Human Stx4	NM_001272096	hs.Ri.STX4.13.2
Human SNAP23	NM_003825	hs.Ri.SNAP23.13
Human SNAP29	NM_004782	hs.Ri.SNAP29.13
Human Stx17	NM_017919.3	hs.Ri.STX17.13.2
Human TSG101	NM_000011.10	hs.Ri.TSG101.13.2

DsiRNA constructs used in this study (**Table 1**),

Figure 1

Figure 2

Figure 3

Figure 4

A — Only myc-UBB⁺¹

UBB⁺¹ SQSTM1/p62 DAPI

Merge / Inset

B — myc-UBB⁺¹ + HA-p62

Merge / Inset

C — myc-UBB⁺¹F4A, I44A + HA-p62

UBB⁺¹ HA DAPI

Merge / Inset

Mean Volume of p62 Condensates (µm³)
Only UBB⁺¹ ; UBB⁺¹ + HA-p62 ; UBB⁺¹ F4A, I44A + HA-p62
*** ***

p62 condensates per cell
Only UBB⁺¹ ; UBB⁺¹ + HA-p62 ; UBB⁺¹ F4A, I44A + HA-p62
*** ***

D

WCL
IB: UBB⁺¹ — UBB⁺¹ / UBB⁺¹ F4A, I44A
Ub-UBB⁺¹
15
UBB⁺¹
10
35 GAPDH

Media
IB: UBB⁺¹ — UBB⁺¹ / UBB⁺¹ F4A, I44A
15
UBB⁺¹
10

E

Co-IP
2% input / FLAG / MYC

IP: MYC IB: UBB⁺¹
20
Ub-UBB⁺¹
UBB⁺¹
10

IP: MYC IB: HA
60
HA-p62FL

F

2% input / No antibody (beads only) / UBB⁺¹ Co-IP
IP: SQSTM1/p62 IB: SQSTM1/p62
60 p62 *

IP: SQSTM1/p62 IB: UBB⁺¹
20 Ub-UBB⁺¹
10 UBB⁺¹

p62 IP (no UBB⁺¹) / 2% input / UBB⁺¹ F4A, I44A Co-IP
IB: SQSTM1/p62
60 p62 *

IB: UBB⁺¹
15 UBB⁺¹

G

TAK243 (E1 inhibitor) − / +
myc-UBB⁺¹ + / +
2% input ; Co-IP
p62 IP (no UBB⁺¹) ; No antibody (beads only)

IP: SQSTM1/p62 IB: SQSTM1/p62
p62 *
60

IP: SQSTM1/p62 IB: UBB⁺¹
20 Ub-UBB⁺¹
10 UBB⁺¹

H — myc-UBB⁺¹ + HA-p62 + TAK243

UBB⁺¹ HA DAPI

Merge / Inset

Mean Volume of p62 Condensates (µm³)
UBB⁺¹ + HA-p62 ; UBB⁺¹ + HA-p62 + TAK243
***

p62 condensates per cell
UBB⁺¹ + HA-p62 ; UBB⁺¹ + HA-p62 + TAK243
***

Figure 5

Figure 6