TITLE

Brain activation for language and its relationship to cognitive and linguistic measures

AUTHORS AND AFFILIATIONS:

Irene Balboni*[a,d], Alessandra Rampinini* [a], Olga Kepinska[b,c], Raphael Berthele #[d], Narly Golestani
#[a,b,c]

a. Department of Psychology, Faculty of Psychology and Education Science, University of
Geneva, Geneva, Switzerland
b. Vienna Cognitive Science Hub, University of Vienna, Vienna, Austria.
c. Department of Behavioural and Cognitive Biology, Faculty of Life Sciences, University
of Vienna, Vienna, Austria.
d. Institute of Multilingualism, University of Fribourg, Fribourg, Switzerland.

* Joint first author

Joint last author

Corresponding author:

Irene Balboni, irene.balboni@unifr.ch and irene.balboni@unige.ch
16

TITLE: Brain activation for language and its relationship to cognitive and linguistic measures

ABSTRACT

Language learning and use relies on both domain-specific and domain-general cognitive and sensory-motor functions. Evolutionary and developmental perspectives, as well as the use of language in interaction, suggest an overlap between language and other skills. Building on our previous behavioural findings, which outlined a consistent behavioural association between language and domain-general skills, we investigated brain-behavioural associations using story listening during fMRI and behavioural measures of language, reading, multilingual experience, cognition, musicality, arithmetic, and motor skills. Participants varied in multilingual language experience and reading aptitudes, including both typical (TRs) and dyslexic readers (DRs). Using multivariate Partial Least Squares correlation, we identified a main component linking cognitive, linguistic, and phonological measures to brain areas underlying lexico/semantics, combinatorial processing, and amodal semantic processing. A second analysis excluding DRs showed closer associations between cognitive/linguistic, literacy, phonological and memory processes within the same brain network as in the full sample. Here, we also isolated additional, complementary components, including one involving speed, automatization and lexical access, linked to auditory and motor brain areas. This suggests greater coherence and more integrated, 'expert' processing in TRs. This work is a first step in exploring complex relationships between language and non-linguistic functions that are important to it.

KEYWORDS

Language, Cognition, Brain-Behaviour Relationships, Dyslexia, Bilingualism

TITLE: Brain activation for language and its relationship to cognitive and linguistic measures

1. INTRODUCTION

Language is a key human ability and is represented by a complex system of spoken, signed or written symbols allowing us to communicate with others (Crystal and Robins 2024 Jul 25). It enables abstraction and is believed to support complex thinking and cognition (Carruthers 2002; Majid et al. 2004). Language is mostly learned and used during face-to-face interactions, and with the support of paralinguistic and non-verbal cues (Levinson and Holler 2014). Being a successful language user entails a remarkable number of processes and subroutines, and yet neurotypical individuals successfully learn and use a first (L1) and often a second (L2) language. The most established models of the neurobiology of language perception and production highlight the hierarchical nature of language, involving phonological, lexico-semantic and morphosyntactic information, subserved neurally by a left-lateralised perisylvian network including primarily dorsal motor/premotor areas and inferior frontal areas, and superior, middle and inferior temporal cortices (Hickok and Poeppel 2004; Hickok and Poeppel 2007; Rauschecker and Scott 2009; Rauschecker 2012; Rauschecker and Scott 2016). Some authors argue for a broader inclusion of areas in the so-called 'language network', like the inferior parietal cortex, underlying knowledge representation, as well as the anterior cingulate and dorsolateral prefrontal cortices for control functions (Price 2012; Hagoort 2014; Hagoort 2017; Hagoort and Özyürek 2024), while others take a conservative approach and consider only areas strictly involved in the access to meaning and sentence construction, including only left-lateralized lateral frontal (IFG, middle frontal) and superior temporal areas (Fedorenko et al. 2024).

Language is a uniquely human and relatively recent evolutionary skill, especially in the case of literacy. To understand how our brains evolved to accommodate such a new and complex skill,

different theories and findings suggest that specific aspects of language had to rely on pre-existing mechanisms that predated the emergence of language and that could be used and specialised for language because of their computational properties (Asano and Boeckx 2015; Asano et al. 2022). This has been described as neural reuse or 'exaptation' (Fitch 2011; Fitch 2012; Janacsek et al. 2022), and may explain why language has often been associated with other domain-general and domain-specific abilities and why the question of language specificity is difficult to resolve.

Different implementations could underlie this overlap between language-specific and other domain-general or domain-specific functions. One possibility is that different abilities might leverage the computational mechanisms of a specific brain area. The computational mechanisms themselves may be shared across domains, while domain-specificity may be shaped by the networks in which it is embedded (Asano and Boeckx 2015; Asano et al. 2022). This has been proposed to be the case for the IFG, specifically BA44/45, which is known to be involved in syntax processing in language, but is thought to have evolved for this due to its role in hierarchical motor planning (Asano and Boeckx 2015; Asano et al. 2022). As such, this region is thought to commonly underlie hierarchical processing for music, action, and language (Fitch 2011; Fitch and Martins 2014; Asano et al. 2021; Asano et al. 2022). A second possible mechanism that has been proposed to explain the neural underpinnings of language in brain regions originally involved in a different function is that of a complete replacement of old functions with new ones. A widely accepted example of this 'neuronal recycling' comes from research on developmental brain adaptations to reading, a late-emerging skill both in evolutionary and developmental terms. Initially involved in face and object perception in preliterate children, this area is repurposed for reading after acquisition. As a result, in adults, this area responds reliably to words as an integral part of the language system, while its sensitivity to faces and objects decreases (Cohen and Dehaene 2004;

Dehaene 2005; Dehaene et al. 2015; Dehaene-Lambertz et al. 2018). These different explanations

for shared mechanisms between language and other domain-general (e.g., cognition, perception,

motor skills) and domain-specific skills (e.g., musicality) processing can explain observed neural

overlap across language and other domains.

Beyond brain regions that may have originally served different purposes in ontogeny or phylogeny,

naturalistic use of language – especially under more demanding conditions – involves interaction

with other cognitive functions (executive control, memory) and modulation from bottom-up

sensory areas. This has been suggested by 'declarative-procedural' memory model of language,

one of the most prominent language models (Ullman 2004; Ullman 2016). The model posits that

the native language acquisition of lexicon and grammar depends on general-cognitive mechanisms

of declarative and procedural memory, respectively, with their associated neural networks playing

a crucial role in both language functions and general memory abilities. Moreover, given that

meaning interpretation is highly context-dependent (Hagoort and Van Berkum 2007) and the

processing of the speech input is influenced by top-down information (McClelland et al. 2006;

Davis and Johnsrude 2007; Rutten et al. 2019), it is more plausible that language processing

engages broader cognitive mechanisms. It is also important to consider that the use of language in

successful communication (i.e., in interaction with others) goes beyond the single word processing

paradigm often used in neuroimaging studies, and it is likely to rely on additional cognitive

processes, such as attention or theory of mind in order to deal with the demands like turn-taking

and tracking the narrative of the conversation (Hagoort 2014). Lastly, given the multimodal nature

of language, which requires the integration of oral, written and also paralinguistic information,

(Levinson and Holler 2014; Cohn and Schilperoord 2024; Hagoort and Özyürek 2024), effective

processing is likely to depend on the coordination of wide multimodal cortical areas and binding

of top-down and bottom-up information, likely to be enabled by subcortical structures, such as the

basal ganglia, thalamus and cerebellum (Asano et al. 2022; Janacsek et al. 2022; Turker et al. 2023).

Taken together, this literature suggests that the neural substrate supporting language may extend

beyond traditional models that include only perisylvian cortical regions and has led some

researchers to view language as "widely distributed throughout the brain" (Aliko et al. 2023;

Drijvers et al. 2025).

At the behavioural level, and in particular in the study of language aptitude (i.e., the potential for

language learning), various models and theories already consider skills beyond language-specific

abilities necessary to make a good language learner. Language aptitude has been linked to musical

abilities (Nardo and Reiterer 2009; Fonseca-Mora et al. 2011; Turker et al. 2017; Turker and

Reiterer 2021; Coumel et al. 2023), as well as general cognition (Doughty et al. 2010) with

particular attention to the role of memory (Wen 2012; Singleton 2017; Feng et al. 2021). Also, in

behavioural studies L1 processing language-specific abilities have been consistently found to

interact with general cognition during development (for a review: Kidd & Donnelly, 2020) and to

be supported by cognition in healthy older adults to ensure language comprehension despite

sensory and cognitive decline (Wingfield and Grossman 2006), particularly during the perception

of speech in noise (Wong et al. 2010). Moreover, language is associated with early motor abilities

(Alcock and Krawczyk 2010; Libertus and Violi 2016; Gonzalez et al. 2019), as well as music (for

an overview: Nayak et al., 2022) and numerical skills (Simmons and Singleton 2008; Purpura and

Lonigan 2013). It is worth noting that while these skills are positively linked to language abilities

in individuals with typical development, they also show deficits in dyslexia, a neurodevelopmental

disorder that leads to poor literacy outcomes (Lyon et al. 2003). Dyslexia has been associated with

impairments in motor control (Marchand-Krynski et al. 2017; Decarli et al. 2024) and musicality

(Overy et al. 2003; Christiner et al. 2022), with musical training being explored as a potential

remediation therapy (Overy 2003; Rolka and Silverman 2015). Additionally, difficulties in

numerical skills are common, as dyslexia and dyscalculia frequently co-occur and share genetic

influences (van Bergen et al. 2023)

In order to learn about the componential structure of language and cognition and of the relationships

therein, individual differences can be leveraged to explore which abilities stably associate and

dissociate in light of the variability (Bates et al. 1991; Kidd et al. 2018). Several behavioural studies

having assessed cognitive and linguistic performance used this approach to explore the complex

interplay between these domains, and showed in adults (Hintz, McQueen, et al. 2024; Hintz,

Voeten, et al. 2024) and in children (Berthele and Udry 2021; Udry and Berthele 2021) that

language-specific and domain-general cognition cannot be clearly separated. In light of these works

and of the above-described debate, in previous work we collected data from a relatively large

participant sample (n = 152) on behavioural measures of language, cognition, musicality,

arithmetic and motor skills, as well as on measures of language and reading experience to explore

patterns of dissociation and association between these domains (Rampinini et al. 2024). In order to

maximise inter-individual variability, we recruited people with a wide range of multilingual

language experience, and we also included a subgroup of adults who had previously been

diagnosed with dyslexia. Using exploratory graph analysis, we analysed the network of

relationships between individual variables while identifying underlying clusters. We confirmed

that language and cognition are stably associated, and that none of the identified clusters were

exclusively composed of language-specific or of domain-general cognitive skills. On the other

hand, musicality, a domain-specific skill often associated with language, was isolated from

language and cognition scores. The same pattern of isolation was present for multilingualism

measures and literacy, indicating that these measures are possibly more linked to experience and

education and that they are dissociated from performance in language/cognition.

Less work has explored the key dimensions supporting language processing beyond language-

specific skills and their corresponding neural correlates in a multivariate manner. Although recent

work has started to highlight the importance of multimodal and multivariate analyses of language

learning profiles, this approach remains relatively underexplored (Feng et al. 2021). In the present

work, we build upon our previous results at the behavioural level to extend our study to brain

activation for native language speech processing (story listening). L1 speech processing is an

informative and stable neural marker (Mahowald and Fedorenko 2016) of how language is

represented in the brain and of individual differences in language processing and experience.

Neural responses to L1 have been linked to outcomes in L2 learning (Díaz et al. 2022; Fuhrmeister

et al. 2023), multilingual experience (Jouravlev et al. 2020), and print processing that underlies

successful language reading (i.e. print-speech convergence; Rueckl et al. 2015). Furthermore, the

use of data collected during L1 as opposed to L2 speech listening allows individual variation to be

sampled while maintaining a degree of homogeneity in terms of age of acquisition and baseline

level of exposure. We included whole-brain activation, including subcortical areas, to assess brain-

behaviour associations in a fully data-driven manner. This allowed us to avoid limiting the analysis

to traditional language areas and implicitly favouring one 'language model' over another. To

uncover this relationship, we used Partial Least Squares Correlation (PLS), a method that identifies

components explaining common variance between two types of data.

2. METHODS

2.1.   Stimuli and procedure

2.1.1. Participants

152 participants took part in the study. Participants were healthy, mostly speaking French as a first

language and having knowledge of two or more languages. None of the participants had official

qualifications for language translation or interpretation, nor did they have professional musical

training. Participants who spoke more than 6 languages were included in the study even if French

was not their native language, if they reported advanced French proficiency. Figure 1 depicts the

multilingual background of the participants in the sample included in the analysis, TRs had on

average higher multilingualism scores than DRs (as shown also in the Supplementary results).

Details of the participants' language and demographic backgrounds are available in the

supplementary materials (Table 1). The final sample only included participants who did not have

missing data on any of the measures analysed. It consisted of 134 participants, aged between 18.1

and 47.2 years ($M = 24.0$, $SD = 4.9$, 92 females). A subset of these participants ($n = 25$) was

diagnosed with dyslexia before taking part in the study. A second analysis was run only on 109

typically reading participants (TRs), again selected based on having no missing data. In this

subsample, participants were between 18.2 and 47.2 years of age ($M = 24.1$, $SD = 5.07$, 74

females).

Figure 1. Language background of the participants in the sample. Each bar represents the language

experience of one participant; the height of the stacked bars within each bar represents the self-

reported speaking proficiency for individual languages (the taller the bar, the higher the speaking

proficiency for that language). Before plotting, the data was sorted based on cumulative language

experience values; as a result, the left-hand side of the figure displays hyperpolyglots' data while

the right-hand side shows data from monolinguals. Dyslexic readers are marked in dark pink.

2.1.2. Procedure

Each participant took part in 4 different sessions (spread over several testing days): session 1 to fill

in online questionnaires, session 2 to perform a portion of the tasks online while supervised by one

trained member of the team, session 3 to complete in-person tasks that required closer supervision

or standardised hardware and session 4 for neuroimaging and genetic data collection (the latter not

included in this study). Questionnaires (session 1) were delivered using Qualtrics XM©, online

tasks (sessions 2 and 3) were delivered using Gorilla, and the fMRI task (session 4) was presented

using Psychtoolbox3 (Matlab r2021b). An overview of all the tasks and behavioural measures is

presented in Table 1. Detailed description of the hardware, procedure, and extended descriptions

of the behavioural tasks and their references can be found in our previous work (Rampinini et al.

2024). All eligible participants provided signed informed consent for all subsequent experimental

procedures and data reuse. The study received ethical approval from the Geneva Cantonal Ethical

Commission (Protocol N. 2021-01004).

2.1.3. fMRI data collection

All participants took part in an MRI session where we collected brain structural (MPRAGE),

MP2RAGE, DWI) data, and functional imaging data during resting-state, and during music,

reading and language localizers, the latter both in the participants' L1 and L2. In this work, we only

analysed fMRI data from the language localiser in the L1, and used the T1 MPRAGE sequences

for co-registration. Images were acquired using a Siemens 3T Magnetom-Prisma scanner at

Campus Biotech (Geneva), equipped with a 64-channel head coil. T1-weighted MPRAGE

structural images were collected using 208 sagittal slices (TRs = 2,300 ms, TE = 3,26 ms, voxel

size: $1\times1\times1$ mm;). BOLD activation was measured using a whole-brain EPI sequence (acceleration

factor -1, FOV=224 mm, voxel size: $2\times2\times2$ mm, TRs = 2000,0 ms, TE=32,00 ms flip angle 75°,

interleaved acquisition, 72 slices).

During the fMRI task, participants passively listened to intact and degraded speech passages of the

story 'Alice in Wonderland' while fixating a black fixation cross on a white screen. The stimuli

and task are openly available (http://web.mit.edu/evlab/aliceloc/index.html), and the procedure for

the degradation of low-level auditory features is described in Malik-Moraleda and colleagues

(2022). We used the localiser to present intact and degraded passages for both the L1 and the L2

to each participant. In total, 3 runs were acquired, each including 5 fixations blocks of 12 seconds

and 16 blocks of 18 seconds each, 4 per each of the intact and degraded conditions. No response

was required, and participants took short, self-timed breaks between blocks. Of the 134 participants

included in the final analysis, a total of 127 listened to passages in French for the L1, while the

remaining 7 chose another language (this always either being their first language or a language that

felt native to them): 3 participants chose English, 1 chose Spanish, 1 chose German and 1 chose

Portuguese. More information on the individual profiles of each participant can be found in the

supplementary material (Table 1).

2.2. Analysis

2.2.1. Behavioural data

Behavioural data for the questionnaires and the tasks completed on the Gorilla online platform

were analysed and scored using in-house Python scripts. Tasks requiring manual scoring, such as

spelling or foreign speech sound imitation, were scored manually by native speakers. All measures,

with their corresponding tasks and scores, are outlined in Table 1. A more detailed description of

the behavioural tasks, together with descriptive results and reliability indices for each task, can be

found in Rampinini and colleagues (2024). The supplementary materials (Figure 1) include plots

with scores for all participants in all of the included behavioural measures, divided by reading

group.

Variable name	Task Name	Brief task description	Selected score	Domain
Reading history	AHRQ	Reading experience, enjoyment and familiarity with reading difficulties	AHRQ index	Language experience

TITLE: Brain activation for language and its relationship to cognitive and linguistic measures

			Accuracy	Domain-specific skills
Musical abilities	AMMA	Sensitivity to tonal and rhythm change	Accuracy	Domain-specific skills
Alerting network (ANT)	ANT-I	Ability to use an auditory alerting cue to speed the reaction to a target	RT no cue trials – RT cued trials	General cognition
Orienting network (ANT)	ANT-I	Ability to use an orienting cue to orient attention to the target location	RT incorrectly cued trials - RT correctly cued trials	General cognition
inhibition (ANT)	ANT-I	Ability to correctly report target direction irrespective of the surrounding flankers (Flanker task)	RT incongruent trials - RT congruent trials	General cognition
Grammar analysis	ArtGram	Ability to infer morphosyntactic characteristics of an artificial language and translate sentences after the presentation of some examples and dictionary	Accuracy	Language skills
Artificial grammar learning	Brocanto	Grammaticality judgment of sentences in an artificial grammar after repeated presentation of grammatical examples	Accuracy	Language skills
Visual memory span	Corsi blocks	Reproducing a visuo-spatial sequence	Span (forward and backward)	General cognition
CVLT short term	California Verbal Learning Test	Ability to recall a 16-item list immediately after presentation (5 trials) and after presenting an interference list.	Accuracy	General cognition
CVLT long term	California Verbal	Recognition, 20 minutes after encoding, of items memorized. Presentation of both	Accuracy	General cognition

TITLE: Brain activation for language and its relationship to cognitive and linguistic measures

	Learning Test	targets and semantic and phonological distractors		
Visual declarative memory	DecLearn	Ability to recognize both real and invented objects 10 minutes after incidental encoding	Accuracy	General cognition
Verbal memory span	Digit span	Short-term and working memory span for digits	Span (forward and backward)	General cognition
Foreign speech sounds imitation	Farsi uvular production	Imitation abilities for uvular sound production	Percentage of correct responses	
Finger tapping speed	Finger tapping	Tapping speed with dominant and non-dominant hands	Average tapping speed	Motor skills (speed)
Foreign speech sound discrimination	Hindi dental retroflex contrast	Adaptive task measuring ability to discriminate a dental-retroflex plosive consonant contrast	Accuracy weighted by difficulty	Language skills
Multilingualism (speaking)	LEAP-Q	Self-report of language experience: number of languages known, in what modality, manner and age of acquisition, proficiency and current exposure	Entropy scores for speaking	Language experience
Multilingualism (comprehension)	LEAP-Q	Self-report of language experience: number of languages known, in what modality, manner and age of acquisition, proficiency and current exposure	Entropy scores for comprehension	Language experience
Multilingual exposure	LEAP-Q	Self-report of language experience: number of languages known, in what modality, manner and age of acquisition, proficiency and current exposure	Entropy scores for exposure	Language experience

TITLE: Brain activation for language and its relationship to cognitive and linguistic measures

Motivation (attitude)	MFQ	Self-report for motivation for language learning	Total score	Language motivation
Motivation (openness)	MFQ	Self-report for motivation for language learning	Total score	Language motivation
Rote learning	MLAT 5	Vocabulary learning in an unknown language	Accuracy	Language skills
Musical training	MUSEBAQ	Questionnaire on musical experience and type and amount of musical training	Musical training index	Music experience
Phoneme suppression	Phoneme suppression	Repetition of nonword removing the first sound	Accuracy	Phonological awareness
Purdue pegboard dominant	Purdue pegboard test	Test for fine motor dexterity	Average number of completed trials, dominant hand only	Motor skills (dexterity)
RAN (RT)	Rapid automatized naming	Rapid naming of objects, colours, digits	Average reaction time (all categories)	Literacy mediator
Fluid intelligence	Raven's APM	Reasoning abilities, ability to complete an incomplete pattern	Accuracy	General cognition
Arithmetic	Revised tempo test	Speeded resolution of sums and subtractions	Maximum number of completed trials	Domain-specific skills
Word spelling	ECLA16	Spelling of words after auditory presentation	Accuracy	Literacy

Pseudoword spelling	ECLA16	Spelling of pseudowords after auditory presentation	Accuracy	Literacy
Spoonerism	Spoonerisms	Ability to transpose the initial sounds of two pseudowords	Accuracy	Phonological awareness
Procedural memory	Serial reaction time	Ability to tap following the pattern visually present on screen, implicit sequence presented across 5 blocks	Procedural learning gain	General cognition
Reading text (Alouette)	Alouette	Ability to correctly read a non-sensical text passage within 3 minutes.	Accuracy	Literacy
Reading text (Pollueur)	Pollueur	Ability to correct a text passage within 3 minutes.	Accuracy	Literacy
Word reading (RT)	ECLA16, EVALEC	Reading of regular and irregular words	Average reaction time per block	Literacy
Pseudoword reading (RT)	ECLA16, EVALEC	Reading of pseudowords	Average reaction time per block	Literacy
Nonword repetition	Nonword repetition	Repetition of nonwords with increasing span length	Accuracy	Phonological awareness

Table 1. Overview of the tasks, scores and resulting variables obtained from behavioural testing.

All the scores, including reaction times and other metrics used in this analysis, were transformed

such that higher values consistently indicate better performance.

2.2.2. Preprocessing fMRI data

Functional images were preprocessed using FSL 6.0. Preprocessing was carried out using FEAT

(FMRI Expert Analysis Tool, version 6.0) with a standard preprocessing pipeline that included the

following steps: removal of non-brain tissue using optiBET (Lutkenhoff et al. 2014), B0 unwarping

using fieldmap images and nonlinear registration to structural T1 images (BBR, FNIRT), and

spatial smoothing using a Gaussian kernel of 5 mm FWHM. Data were motion-corrected using

ICA-AROMA (Pruim et al. 2015) and no participant had an absolute displacement higher than 5

mm (which would have resulted in their exclusion from the sample). We then removed nuisance

regressors (white matter, WM, and cerebral-spinal fluid, CSF), performed high-pass filtering at 90

s and registered the preprocessed data to the MNI152 standard space (Grabner et al. 2006). The

BOLD response was modelled using a double-gamma haemodynamic response function (HRF).

Activation for intact and degraded L1 speech listening was modelled, and the intact *vs.* degraded

speech contrast was generated.

2.2.3. Extraction of individual level activity with parcellation

Once the intact *vs.* degraded brain activation contrast was computed for each participant, we

defined different brain parcels in order to maintain a whole-brain approach while reducing the

numbers of brain variables for the PLS analysis (i.e. as compared to a voxel-based approach). We

included all cortical and subcortical regions. For the cortical regions, a total of 397 parcels, or

regions of interest (ROIs) were identified. The cortex was segmented using the Glasser multimodal

parcellation (Glasser et al. 2016). The parcellation was adapted to volume space, registering it to

the same MNI152 template as the functional data (Horn 2016 Jun). The volume was then split by

hemisphere to obtain 180 parcels per hemisphere. In the results section, cortical ROI labels

correspond to each area's name in the original parcellation publication (Glasser et al. 2016). ROIs

for the hippocampus were excluded from the cortical parcellation because hippocampal areas were

defined using a subcortical atlas, as follows: subcortical regions comprising bilateral hippocampus,

thalamus, amygdala, pallidum, putamen, and caudate nucleus were identified using the Harvard-

Oxford subcortical atlas (Makris et al. 2006). Cerebellar regions were identified using a

probabilistic cerebellar atlas (Diedrichsen et al., 2009; full list of cerebellar ROIs in the

supplementary material). Both subcortical and cerebellar areas were extracted using a threshold of

50%. The full list of ROIs is presented in the supplementary materials. For each of the ROIs

obtained using the above parcellations, average contrast values for the difference in brain activation

for intact *vs.* degraded speech were extracted. The resulting brain dataset extracted for each

participant contained the average contrast values for each of the 397 ROIs.

2.2.4. PLS

We used PLS to evaluate the multivariate relationships between two data modalities, brain and

behavioural, aiming to isolate components that maximise shared variance between the two sets of

data. PLS is conceptually similar to principal component analysis, which finds patterns within a

single dataset, with the different that PLS extracts components that best explain the correlation

between two sets of variables. PLS works by computing linear combinations of variables from both

datasets to uncover hidden relationships. This makes it a powerful tool for studying complex

interactions between different types of data.

We implemented it by first z-scoring both datasets and correcting them for age, sex and handedness

through residual calculation (McIntosh and Lobaugh 2004; Krishnan et al. 2010; Krishnan et al.

2011). After data preprocessing and preparation, the analysis was performed using the myPLS

toolbox (https://github.com/danizoeller/myPLS). A correlation matrix between behavioural and

brain data was generated, and then the main components which explained the relationship between

the two modalities were extracted through singular value decomposition.

For each component, behavioural and brain weights were calculated: these explain how strongly

each single variable contributed to the multivariate correlation between the data modalities in that

component. Behavioural weights, or saliences, represent how strongly each variable contributes to

the correlation between brain and behaviour in each component. Brain weights, or saliences,

represent how strongly the intact versus degraded speech listening average contrast values within

each ROI contributes to the correlation between brain and behaviour in each component.

The significance of each component was assessed using 1000 permutations, and the stability of the

weights of each variable per component was evaluated using a bootstrapping procedure (500

samples). The significance of each variable was determined using a Z score $>=|3|$ threshold, which

corresponds to a two-tailed probability of 0.0013. Previous work suggested that a Z score $>=|2|$

(Krishnan et al. 2011) or $>=|2.57|$ (equivalent to a two-tailed probability of 0.01; McIntosh &

Lobaugh, 2004) could be used, but in the present analysis, given the high number of variables used

in both datasets, we chose a stricter threshold to consider the results robust/significant. The results

report weight means and standard deviation after bootstrapping for each variable/ROI. More details

on permutation, bootstrapping and thresholding procedures are available in the Supplementary

Materials (Supplementary methods section).

As mentioned in the Participant section, we initially ran the analysis on the full sample including

TRs and DRs ($n =134$), and then on a subsample including TR participants only ($n=109$). We could

not perform a separate analysis on the DRs alone due to their smaller sample size ($n =25$), which,

given the number of variables included in the analysis, would likely have resulted in overfitting

and poor generalization (Mihalik et al. 2022). To ensure that any potential differences between the

two analyses were due to sample characteristics and not to differences in sample size, we ran an

additional PLS analysis on a third subsample, hereafter referred to as the verification subsample.

This had the same sample size as the subsample of TRs only (n=109), but included the same

proportion of TRs and DRs as the full sample.

3. RESULTS

3.1.   Partial least square results

Overall, the analysis of the whole sample resulted in one significant component whereas the

analysis including only TR participants resulted in 3 significant components. The analysis from

this verification subsample only involving TR participants showed a substantial overlap with the

results in the whole sample, confirming that differences between the latter and the subsample were

likely due to differences driven by sample characteristics and not sample size reduction. All the

results for the verification subsample are available in the supplementary material.

3.1.1. Brain and behavioural relationships in the whole sample

We applied a PLS analysis to the 36 behavioural measures and to the average activation (contrast

values for intact *vs.* degraded L1 speech) from the 397 cortical and subcortical areas during L1

story listening. One component was significant after permutation testing ($p = .006$, $r = .34$). Figure

2 shows the contribution of each variable from both modalities to the brain-behavioural correlation

component (more details are available in the Results section of the Supplementary Material). The

component explained 36% of the brain-behavioural covariance.

At the behavioural level, the component was mostly positively associated with half of the

behavioural tasks (Figure 2A). This component identified an association between measures of

general cognition (memory in particular), language skills (with a prominent role of grammar) and

to a lesser extent of phonological skills. More specifically, a positive association was found

between the brain measures and measures of language and cognition that rely heavily on memory

processes (such as nonword repetition, declarative and verbal memory, and rote learning), pattern

recognition and learning (morphosyntactic patterns for a grammar analysis task, and visual patterns

for fluid intelligence scores), or a combination of both (as seen in artificial grammar learning).

Artificial grammar learning skills were, both in the full sample and in the typical readers only (see

next analysis) the strongest contributors to the component. Another set of measures pertain to the

auditory domain and to auditory working memory: musicality and phonological awareness

(including spoonerisms, nonword repetition, and foreign speech sound imitation), and to a literacy

measure involving sound-symbol correspondences (i.e. word spelling). Multilingualism scores,

with a stronger influence of multilingual comprehension than of speaking, also contribute

positively to this component. Moreover, there was a positive contribution of alerting skills, i.e. a

measure of response readiness and of sustained attention to incoming stimuli (Ishigami et al.,

2013), and of motor dexterity (Purdue pegboard test). Last, Multilingual exposure, differently from

multilingual comprehension and speaking abilities, contributed negatively to the component, with

a relatively low weight compared to positively associated variables.

All cortical and subcortical brain regions showing a significant contribution to the

component, and their weights, are shown in Figure 1B (results for cerebellar ROIs and uncorrected

results can be found in the Supplementary Materials). Overall, better performance on the above

tasks correlated with stronger activation of bilateral anterior temporal cortical areas (e.g. STGa,

STSda, STSva, TGd, TGv, TE1), posterior cingulate areas (e.g. RSC, v23abm, POS1), inferior

parietal regions (PGi, PGs, PFm), and temporo-parietal-junction (TPJ; TPOJ1, TPOJ2, TPOJ3).

The latter regions – inferior parietal cortex and adjacent temporo-parietal junction – showed

somewhat left *versus* right hemisphere lateralization, respectively. Other areas that displayed a

360 significant positive activation in relation to behavioural measures comprised predominantly left-

361 lateralised dorsolateral prefrontal (8v, 8c) and inferior frontal (IFJp, 45) regions. Better scores also

362 positively correlated with activity in medial temporal regions such as the entorhinal cortex (EC)

363 and perirhinal cortex (PeEC) in the left hemisphere, presubiculum (PreS) in the right hemisphere

364 and para-hippocampal (PHA3) areas bilaterally. Subcortically, Figure 2c shows a positive

365 correlation with bilateral amygdala and right hippocampus. None of the cerebellar regions were

366 significantly associated with the component.

368 Figure 2. Results of the behavioural, cortical and subcortical saliences/weights for the significant correlational PLS component in

369 the whole sample. Blue represents significant positive weights, red represents significant negative weights, and grey represent non-

370 significant ones. (A) Contribution of behavioural variables to the component (for all scores, higher values represent better

371 performance). (B) Contribution of cortical ROIs to the component. (C) Contribution of subcortical areas to the component. For each

372 component, mean weights, SD and brain uncorrected results can be found in the supplementary material, in the Supplementary

373 results section.

3.1.2.   Brain-Behavioural relationship in typical readers only

We applied a second PLS analysis to the same behavioural and brain measures presented in the

above section, but this time, we included only participants without a dyslexia diagnosis *(n = 109)*.

After permutation testing, we identified 3 significant components.

Component 1

The first component accounted for 44% of the variance *(p = .001, r = .45)*. Behaviourally, this

component resembles the one identified in the whole sample, but it is associated with a higher

number of memory measures and, differently from the whole-sample analysis, includes a larger

number of literacy measures (brain and behavioural saliences are shown in Figure 3).

A positive association was found with memory measures (such as Nonword repetition,

Visual declarative and Verbal memory span, and Rote learning), pattern recognition and learning

(Grammar analysis, Fluid intelligence, Artificial grammar learning), auditory measures including

working memory: musicality and phonological awareness (including Spoonerisms, Nonword

repetition, and Foreign speech sound imitation), and to a literacy measure involving sound-symbol

correspondences (i.e. Word spelling). Positive associations were present also for multilingualism

scores for comprehension and speaking, alerting skills and motor dexterity (Purdue pegboard test).

Moreover, focusing on the differences in the present TRs analysis compared to that in the full

sample, here component 1 was associated with additional measures for memory (CVLT long term,

Visual memory span), arithmetic, literacy and literacy mediators (RAN, both Reading text

measures, Reading history). Also, even though this component includes measures of phonological

abilities in both the TRs and the whole-sample analyses, in this analysis including only the TRs

group, the phonological measures play a more prominent role (i.e. higher weights). This suggests

a stronger influence of phonology, which is particularly interesting when considering that this

component also encompasses literacy measures. Finally, in the whole sample, multilingualism

exposure contributes negatively to the component.

At the brain level, the ROIs contributing to the first component were similar to those

contributing to the single component found in the whole sample, with more bilateral activation

especially in temporal areas. Higher behavioural scores were associated with more activity in

middle-to-superior posterior (STSdp, STSdp) and anterior (STSda, STSva, TE1a, STSva, STSda,

STGa, TGd, TGV) temporal areas, in frontal areas (IFJa and 45 on the left, 45 and 47l on the right),

in posterior inferior parietal regions (e.g. PGi, and surrounding clusters, again predominantly left-

lateralised) and in temporoparietal regions (e.g. TPOJ2), the latter again somewhat right lateralised

in extent. There was also a positive correlation with medial posterior cingulate (RSC, POS1, 23ab,

31pv and surrounding areas), and bilateral presubiculum (PreS) and parahippocampal (PHA1,

PHA2, PHA3) areas. Subcortically, areas that contribute positively to the relationship are the right

hippocampus and left amygdala, and the cerebellum on the right crus II. All results for cerebellar

ROIs and uncorrected results can be found in the Supplementary Materials.

TITLE: Brain activation for language and its relationship to cognitive and linguistic measures

Figure 3. Results of the behavioural, cortical and subcortical saliences/weights for the first significant correlational PLS component in the typical readers. Blue represents significant positive weights, red represents significant negative weights, and grey represents non-significant ones. (A) Contribution of behavioural variables to the component (for all scores, higher values represent better performance). (B) Contribution of cortical ROIs to the component. (C) Contribution of subcortical areas to the component. For each component, mean weights, SD and brain uncorrected results can be found in the supplementary material, in the Supplementary results section.

Component 2

The second component (Figure 4) accounted for 19% of the variance ($p = .012$, $r = .35$). This component was mostly negatively associated with tasks mainly related to motor and processing speed and lexical access compared to Component 1. Negative associations were present for motor speed, with Finger tapping speed (and not for dexterity, as in Component 1) and lexical access, reading and memory (RAN, CVLT short term, word and pseudoword reading, non-sensical text reading) and phonological perception. Two tasks measuring Visual declarative memory and

phonological skills (Spoonerisms) were also positively associated with the component, with overall

smaller weights compared to the negatively correlated measures.

At the brain level, mirroring the behavioural results, lower-level auditory and sensorimotor

areas were associated with this second component. Poorer performance on the tasks negatively

contributing to the component was associated with more activation in bilateral early auditory (A1,

MBelt, PBelt, LBelt, A4, A5) and other auditory temporal association areas (TA2). Moreover, left

primary motor (4) and bilateral sensorimotor cortices (areas 3a, 3b on the left, and areas 1, 2 on the

right) were involved. Smaller positive contributions came from bilateral opercular/insular regions

(e.g. OP4, 43, FOP1 bilaterally, OP1, RI, PFcm on the left). None of the subcortical or cerebellar

ROIs contributed significantly to this second component. All results for cerebellar ROI and

uncorrected results can be found in the Supplementary Materials.

Figure 4. Results of the behavioural and cortical and subcortical saliences/weight for the second significant correlational component

in the typical readers. Blue represents positive weights, red represents negative weights, and grey represent non-significant ones.

(A) Behavioural correlation for each individual variable to the component (for all scores higher values represent better performance).

(B) Cortical ROI correlations with the component. (C) Subcortical areas correlations with the component. For each component,

mean weights, SD and brain uncorrected results can be found in the supplementary material, in the Supplementary results section.

Component 3

The third component accounted for 8% of the variance ($p = .026$, $r = .67$). This component showed

the most complex pattern of results in the analysis. Figure 5A shows that it was behaviourally

negatively associated with measures of auditory and phonological skill (e.g. Nonword repetition,

Foreign speech sound discrimination, Phoneme suppression) and Verbal memory span score, and

positively associated with a mix of cognitive, linguistic, and multilingual experience measures,

some of which are negatively correlated with the previous components (e.g. Text reading, RAN,

Orienting).

In terms of relationships with brain activation, we observed a clearly lateralised pattern,

mostly in higher-level language regions (Figure 5B). In the left hemisphere, regions contributing

positively to the component included early auditory (MBelt, Pbelt) and associative auditory areas

(STSda, STSva, STSdp, STSvp, STGa, A4, A5), as well as some clusters in opercular regions

(45,43, OP4). Negative relationships were observed in right-lateralised areas such as the inferior

frontal (IFJp, IFJa, IFSp, IFSa, 44, 45) and dorsolateral prefrontal cortices (55b, 8c, P9-46v,

p47r), in the temporoparietal junction (e.g. TPOJ1, DTV), in the posterior STS and in other

medial clusters (SFL, SCEF). In the right hemisphere, positive associations were only found with

a left parahippocampal area (PHA2) and with one region in the posterior cingulate cortex

(v23ab). None of the subcortical regions were significantly associated with this component

(Figure 5c), and only the left cerebellar area VIIb negatively contributed, though marginally, to

the component (Supplementary material).

Figure 5. Results of the behavioural and cortical and subcortical saliences/weight for the second significant correlational component

in the typical readers. Blue represents positive weights, red represents negative weights, and grey represents non-significant ones.

(A) Behavioural correlation for each individual variable to the component (for all scores higher values represent better performance).

(B) Cortical ROI correlations with the component. (C) Subcortical areas correlations with the component. For each component,

mean weights, SD and brain uncorrected results can be found in the supplementary material, in the Supplementary results section.

4. Discussion

We investigated brain-behaviour associations by examining, in a data-driven and multivariate

manner, behavioural measures of linguistic, domain-general and domain-specific skills and brain

activation during story listening in the native language. We performed two PLS analyses, one on

the full sample including TRs and dyslexic DRs readers, and the other only in typical readers. The

former analysis allowed us to explore brain-behaviour relationships across a wider range of

performance – one which included a (language/reading) learning disorder, and the latter allowed

us to determine which relationships are more likely to be observed in typical readers. Results that

are consistent across the two analyses are likely to reflect more stable and potentially more

generalisable results. Results that differ between the two analyses are likely to reflect differences

arising from the additional inclusion of individuals with a previous history of reading disorder,

and/or differences arising from the inclusion of individuals with lower multilingual language

experience, given that DRs on average has a lower multilingual language experience and exposure

than the TRs.

In the next sections, we will discuss the brain-behaviour multivariate components arising from

the two analyses, in light of how brain activation during speech processing is related to positive or

negative relationships with subsets of tasks.

4.1.   Overview of the components

4.1.1. Component 1

Behaviourally, Component 1 (accounting for 36% of the brain-behavioural covariance in the full

sample, and 44% of the variance in the TRs) in both samples showed an association between

higher-level cognitive and linguistic skills, particularly for memory and grammar, and

phonological awareness. These were associated with neural activation in higher-level areas within

the language network known to be involved in combinatorial processes, in amodal semantic hubs,

and in areas involved in phonological processing. In the TRs sample, behaviourally this component

also included motor dexterity, and literacy measures, for text reading and spelling, with a heavier

involvement of memory and phonological skills. This suggests that while language and cognitive

measures are strongly and consistently associated across samples, literacy also becomes part of this

association when only typically reading participants are included.

Overall, there are substantial similarities between the brain-behavioural association

observed in the two analyses. In more detail, two of the tasks that positively contributed to this

component reflect grammatical rule learning and knowledge (i.e. Artificial grammar learning and

Grammar analysis) and pattern recognition (i.e. Fluid intelligence) skills, which reflect complex

pattern analysis, arguably important components of cognition and intelligence. In fact, Artificial

grammar learning was the strongest behavioural contributor to the brain-behavioural relationship.

Neurally, activity in the bilateral dorsal and ventral anterior temporal lobes (ATL) was one of the

strongest contributors to the brain-behaviour relationship of this component, again in both analyses.

The ATL has been shown to be sensitive to abstract syntactic structure (Fedorenko et al. 2010;

Pallier et al. 2011), and to be involved in lexico-semantic and compositional semantic processes

(Hickok and Poeppel 2007). The left IFG (area 45) also contributed to this component; this area is

known to support morphosyntactic processes (Hickok and Poeppel 2007) and is proposed to be

involved in the semantic combinatorics (Friederici 2006; Schell et al. 2017).

There were also positive behavioural contributions of tasks that support or are related to

lexico-semantic processing, such as memory and vocabulary learning tasks (e.g. Verbal memory,

Rote learning and Visual declarative memory scores). Both brain imaging research and

neuropsychological work on patients with stroke or neurodegeneration support the role of the ATL

as an amodal, domain-general hub for semantic and conceptual knowledge and memory (Lambon

Ralph et al. 2010; Lambon Ralph et al. 2016; Gainotti 2017). In both analyses, we found

contributions of other brain areas that are also known to be amodal semantic hubs, where unimodal

information converges: the bilateral IPL (PGi, PGs), bilateral posterior cingulate/retrosplenial (e.g.

RSC, POS1, v23ab), medial prefrontal (9m, left-lateralised in the whole sample and bilateral for

TRs), bilateral parahippocampus, and subiculum areas in the right hemisphere and entorhinal

cortex in the left hemisphere (Fernandino et al., 2016).

Another set of tasks positively correlated with this component pertained to phonological/auditory

processing and working memory (e.g. Foreign speech imitation, Spoonerisms, Phoneme

suppression, Musical abilities). These tasks contributed in both analyses, but more so in the TRs

analysis. These skills may partly drive the correlation with the activation in the posterior STS,

known to be involved in speech and phonological processing (Hickok and Poeppel 2007; Hein and

Knight 2008). Also, the relationship with phonological and language processing (imitation and

multilingualism) may be driven by the activation of the IPL, a central hub in the dorsal phonological

route of reading (Kuhl et al. 2020; Turker and Hartwigsen 2021). The IPL is also associated with

sound-based phonemic representations (Coslett and Schwartz 2018), perception of ambiguous

speech (Kilian-Hütten et al. 2011) and works in combination with IFG areas to support

phonological memory processes (Kellmeyer et al. 2013).

The prominent role of grammatical processing and of verbal declarative memory in this

first component inevitably recalls Ullman's declarative/procedural model (Ullman 2004; Ullman

2016). While the model predicts that in L1, lexicon depends on the declarative memory system and

grammar on the procedural one, it also posits that for L2, especially if explicitly learned and at low

level of proficiency, both of these language components would depend on explicit and effortful

retrieval from the declarative memory. The tasks associated with this component tapped into

memory for lexical items and memory for verbal items, as well as into the learning and analysis of

artificial languages, which are expected to assess L2 grammar learning processes. In line with this,

our results show that procedural memory (measured through a serial reaction time task) did not

contribute to the component. However, the declarative-based tasks that do contribute to the

component correlate positively with brain activation in areas supporting declarative processes, in

combination with a set of semantic and amodal conceptual hubs. This suggests that the learning

and use of novel (i.e. L2-like) grammatical rules and verbal memory may rely on declarative skills

and underlying medial temporal structures, as suggested by the declarative/procedural models, but

that they are additionally supported by combinatorial and amodal semantic systems.

4.1.2. Component 2

Component 2 (accounting for 19% of the brain-behavioural covariance), obtained in the analysis

with TRs only, represents a complementary, lower-level component compared to Component 1.

Behaviourally, there are negative contributions from tasks reflecting automatization, processing

and motor and processing speed and lexical access (Penolazzi et al. 2007). What appears to

commonly underlie most of the tasks showing negative weights is that most require fast processing

and quick reactivity (RAN, Orienting and Finger tapping speed). In terms of the brain, lower-level

auditory and sensorimotor cortices contribute positively to this component. The negative

relationship between the brain and most of the behavioural measures of this component could

reflect optimisation or efficiency, with lower behavioural scores being associated with increased

brain activation. There were also, however, positive contributions of Spoonerisms, measuring

phonological awareness, and Visual declarative memory to this component.

In more detail, bilateral activation of auditory and sensorimotor areas was associated with

the above tasks, with a left-lateralised bias particularly in superior temporal regions and in the

frontal opercular regions. Lower behavioural scores were associated with greater activation in

auditory areas including the planum temporale, which are important for both phonological and

orthographic processing in good readers (McNorgan et al. 2014). Moreover, greater activation was

present in the TPJ, known to be important for speech-related auditory-motor association, this being

a fundamental step in lexical access (Davis 2016 Jan 1) and being required in several of the tasks

that are negatively associated with this component.

Most of these tasks also require speech production, which can explain the involvement of

frontal opercular and motor areas. Bilaterally, the posterior frontal opercular/insular area 43 is

associated with movements such as swallowing, area FOP1 is associated with imagined

movements, and OP4 integrates sensorimotor information into actions (Baker et al. 2018).

There was also activation in the left motor (area 4) and sensorimotor areas (areas 3a and

3b). Areas 4 and 3b are thought to be involved in the control of hand/finger movements and in the

processing of finger-specific information, respectively (Baker et al. 2018). The involvement of

motor cortices is consistent with the correlation with our motor speed task. Moreover, left area 4

has been shown to be active during passage and alphabet writing tasks (Beeson et al. 2003).

4.1.3. Component 3

Component 3 (accounting for 8% of the brain-behavioural covariance), again identified only in

TRs, showed a more complex pattern of associations. Behaviourally, it was positively associated

with cognitive and multilingualism scores, and negatively with auditory/phonological abilities.

These performances were associated with more activation in left superior temporal and some

inferior frontal/opercular areas, and with less activation in right hemispheric IPL/TPj and

IFG/DLPFC areas.

In particular, better memory, inhibition and multilingualism scores were associated with lower

activity in the right inferior frontal sulcus regions (IFS; IFSa, IFSp, IFJa, IFJa), right middle frontal

gyrus (area 8, p9-46v), right IFG (44, 45, p47r), and IPL/TPJ regions (e.g., TPOJ1, STV, PFm,

PSL). The right IFS has been linked to music, reasoning, attention (Ruland et al. 2022) and to rule

implementation (Demanet et al. 2016). The right middle frontal gyrus has been shown to be

involved in attention re-orienting (Japee et al. 2015), and its connectivity to be negatively correlated

with numeracy (Koyama et al. 2017), which is also negatively associated in this component.

Moreover, the right IPL/TPJ region supports executive functions (including working memory and

inhibition) (Rothmayr et al. 2011; Igelström and Graziano 2017) and together with right IFG areas

it is part of the ventral attention network, underlying bottom-up attentional processes (Corbetta and

Shulman 2002; Igelström and Graziano 2017). At the same time, in this component, better scores

on these tasks are also associated with poorer phonological awareness/musical abilities, and with

greater engagement of left lateralized, early auditory (A5, A4, Pbelt, Mbelt), auditory association

(STSda, STSvp, STSdp) and opercular areas (43, 45, OP4).

The relationships within this component suggest that increased multilingualism is associated

with better short- and long-term memory for words and with better attention, but with poorer

phonological awareness and poorer performance on tasks requiring working and short-term

memory for sublexical (i.e. phoneme-level) information. This may be partly explained by the nature

of our sample. Some of the most multilingual speakers (poly- and hyperpolyglots) in this study

were highly proficient but non-native French speakers. Although the hyperpolyglots did not differ

from the other participants in their behavioural scores on these tasks (as shown in the

supplementary results, Table 10), the fact that they were performing tasks in a non-native language

may have led some of them to rely more on lower-level auditory (e.g. acoustic) processes and on

greater engagement of lower-level auditory regions to perform tasks requiring subtle phonological

manipulations. Moreover, within this third component, higher multilingual experience is associated

with clear brain functional lateralisation, suggesting greater engagement of a left-lateralised higher-

level auditory (i.e. belt and parabelt regions including STS subregions and inferior frontal (i.e. BA

45) language network, and lower (maybe optimised) engagement of right-lateralised attention and

executive areas. It should be noted, however, that this third (and last) component explains a smaller

percentage of the variance (8%) compared to the other two (63% in total), and that therefore these

results and interpretations may not be representative of broader and generalisable trends

615 **4.2.   Language and its components**

4.2.1. Language and cognition

Our data-driven analysis did not explicitly confirm or refute any of the traditional language models

presented in this paper, nor was it our goal to compare language models. We aimed at more broadly

exploring the relationship between performance on linguistic and non-linguistic tasks thought to

be involved in aspects of language, and brain activation during a story listening task. Our results

identified components that span most of the major areas considered essential for language

processing, as well as additional regions.

In both samples, we have been able to show that, at the behavioural level, cognition and language

present a stable association, suggesting a consistent overlap. This is in line with previous

behavioural analysis, with different statistical methods, in this sample (Rampinini et al. 2024), and

also in children tested with tasks tapping into similar domains (Udry and Berthele 2021). In both

of our analyses, measures including memory and grammar learning abilities were strong drivers of

the brain-behavioural association in the first, most important component. Previous work has

successfully shown dissociations between language-specific and cognitive processes at both the

brain and behavioural levels using finer-grained, targeted analyses, and areas responsive to

language have previously been shown to be less responsive to domain-general cognitive tasks

(Fedorenko and Varley 2016; Fedorenko et al. 2024). However, language does not occur in a

vacuum. Our analysis involved exploration of the relationship between behavioural scores and

language-specific activation rather than with task-free neural data (e.g. resting-state functional

connectivity or brain structural metrics). As such, our results strengthen the case for the connection

between language and cognition, since testing for associations with language-specific activation

during story listening could have biased results in favour of language-specific tasks and/or of brain

activation underlying language subcomponents (i.e., phonology vs. lexicon/semantics), while

minimizing the role of cognitive measures. However, a consistent association between language

and cognition was observed in the brain-behaviour components, despite the use of brain activation

for story listening in the analysis which could have favoured language-specific outcome.

As discussed in the introduction, the overlap between language-specific processing and other

domain-general and domain-specific abilities is predicted in light of evolutionary and

developmental mechanisms and the nature of language in interaction. Language is mostly used in

interaction with other individuals (Levinson and Holler 2014), and the use of language in real-life

contexts requires us to engage in other cognitive processes beyond simply retrieving semantic and

syntactic information, in order to keep track of the narrative, turn-taking and conversation (Hasson

et al. 2018). Moreover, language and cognitive processes mutually influence each other during

language learning (Kidd et al. 2018; Kidd and Donnelly 2020; Schwering and MacDonald 2020;

Donnelly and Kidd 2021). The principles of Hebbian plasticity (i.e., "what fires together, wires

together") suggest that, at the neural level, interactions between the brain functional underpinnings

of different processes can establish enduring functional connections between them. It is likely that

the interaction between these skills during development, language learning and language use results

in long-lasting connections between these domains throughout life, both at behavioural and neural

levels (Munakata and Pfaffly 2004; Caporale and Dan 2008). Additional support for this idea

comes from clinical work, where neurodegeneration has been observed to spread along established

brain networks (Bak and Chandran 2012; Zhou et al. 2012; Drzezga 2018). One example of this is

found in Motor Neuron Disease, in which the linguistic skills that are more impaired are those with

the closest brain functional links to the motor system, with earlier impairment of verb and action

processing than object naming (Bak and Chandran 2012). Taken together, these findings support

the view that language and cognition are deeply intertwined at both behavioural and neural levels,

shaped by shared developmental trajectories, overlapping neural circuitry, and continuous

interaction throughout life.

4.2.2. Complementary patterns for language and cognition

Examining the components obtained in the TRs reveals a pattern of complementarity, particularly

between the components explaining most of the variance, Components 1 and 2.

Component 1 was predominantly linked to higher-order cognitive and linguistic skills, as well as

domain-specific skills, and literacy measures such as spelling and text reading. It is known that

higher instructional time with opaque orthographies is associated with better performance on

spelling (Zaretsky et al. 2009). Higher instructional time with opaque orthographies is also related

to text reading fluency, which requires additional skills beyond the necessary automatized ability

to read single words, such as sequential processing of words (van Viersen et al. 2024). This could

potentially indicate that this component is linked to higher-level, well-established skills. Even in

the motor domain this component was more strongly associated with more complex motor abilities,

like dexterity and fine motor coordination as measured by the Purdue Pegboard Test, rather than

simple motor speed, assessed through speeded tapping. This component was reflected in the

cortical activation of higher-order language and cognitive areas, where increased activation

correlates with better behavioural performance. This suggests an active role of these regions during

language processing, particularly in individuals with automatised, proficient reading skills and

more multilingual experience.

Conversely, the second component captured a subset of lower-level abilities, including motor

speed, rapid lexical access, and reading single items and nonsensical text. This component was

mostly associated with activity in sensory and motor cortices. Higher activation in these regions

correlated with poorer behavioural performance, suggesting that people with lower abilities rely

more heavily on lower-level cortical regions.

This component appears to show a complementary pattern in terms of the behavioural tasks

involved, with Component 1 involving higher-level, well-established skills and Component 2

involving lower-level, speed-related skills. Furthermore, at the neural level, Component 1 is

primarily associated with higher-order association cortices, which are predominantly transmodal,

whereas Component 2 is primarily associated with lower-order sensorimotor cortices, which are

typically considered to be unimodal (Sydnor et al. 2021). The complementarity between the 2

components is likely to reflect an interplay between the two key aspects that characterise a typical

skilled reader. Individuals with stronger linguistic, cognitive, musical and higher-level reading

skills and with greater multilingual experience appear to use higher-level language and cognitive

brain areas more actively, while relying less on lower-level sensorimotor regions.

4.2.3. Literacy and other language functions

The association between language/cognitive scores and literacy may be unexpected given the previous work by our group (Rampinini et al. 2024) in which both in the full sample and in TRs only, literacy measures tended to consistently cluster separately from other language/cognitive scores. However, we need to consider that the analysis methods were different across studies, and that here, we explored the relationship between different data types, i.e. brain and behaviour, and not just between variables within one modality. We must consider that "cognitive processes and neural processes are not the same thing" (Valian, 2016, p. 570), and while literacy tasks can be dissociated from other language/cognitive skills at the behavioural level, they could rely on mostly shared underlying neural substrates.

In the present study, the differences between the analysis in the full sample (i.e. including the DRs) and the analysis including only the TRs showed that in the latter, these cognitive and linguistic processes are also associated with measures of literacy, particularly for reading fluency and history, while also showing a stronger association with phonological awareness (accompanied by more STG activation). This may suggest that in typical readers, literacy skills may rely on, or be more strongly linked to, more general cognitive/linguistic abilities and to their associated high-level neural mechanisms, reflecting a more specialised and expert, reading-ready cognitive architecture. On the other hand, in dyslexic readers, these literacy tasks engage different brain areas (i.e. not those language / cognitive ones associated with Component 1), perhaps as compensatory or more variable (e.g. noisy; Hancock et al., 2017) neural activations. Moreover, it suggests that when TRs only are considered, spoken language and literacy skills (along with phonological mediators) are associated to overlapping neural substrates. This finding is in line with the print-speech convergence hypothesis, which postulates that successful reading is supported by strong overlap

between the neural substrates underlying the processing of written and of spoken language

(Shankweiler et al. 2008; Rueckl et al. 2015; Preston et al. 2016).

4.2.4. Multilingualism

In this study, we characterised multilingualism using 3 separate entropy scores that

continuously captured participants' language experience in terms of proficiency, comprehension

abilities and current exposure. Importantly, multilingualism was not treated as an independent

variable or used to categorise participants into binary groups such as 'monolinguals' and 'bilinguals'.

While this approach does not allow us to draw strong conclusions regarding the differences in the

components based on language background, we consider that our approach offers several

advantages: (1) it provides more nuanced and ecologically valid measures of multilingualism

compared to strict dichotomies (e.g., mono- *versus* bilingual), and (2) it allows for the exploration

of brain-behavioural associations across the full spectrum of different aspects of the multilingual

experience (i.e. proficiency, comprehension abilities and exposure).

In terms of the role of the multilingualism variables in the obtained components,

multilingualism exposure contributed negatively to the first component in the whole sample only.

This negative contribution may be due to the differences between the three multilingualism

variables being measured. Multilingualism *exposure* assessed the number of languages currently

present in the participants' environment, whereas the other two scores — multilingualism *speaking*

and multilingualism *comprehension* — better capture language knowledge. The participants

showed considerable variation in these latter 2 measures, with some polyglots speaking or

understanding more than 10 languages. As a result, the current exposure measure may be less

informative than our two measures of multilingual knowledge, as polyglots might not have been

exposed daily to all the languages they know. In contrast, less multilingual participants (amongst

which we can find DRs) were likely exposed to their known languages daily due to the multilingual

nature of the region where data collection took place. Thus, while multilingual speaking and

multilingual comprehension scores are strongly positively associated, multilingual exposure

represents an anomaly. We may have very multilingual speakers with relatively low current

exposure, making this measure less informative. This could explain why only language knowledge,

represented by measures assessing cumulative experience with speaking and understanding one's

languages, were positively related to linguistic and cognitive skills. Conversely, current exposure

to language, which provides a snapshot of the language environment at the time of assessment, did

not show the same relationship. Once DRs were removed, multilingual exposure remained

negatively associated with the component but not significantly, possibly indicating the reduced

variability between these three measures given that DRs are less multilingual.

It is important to note that the participants in this study were, on average, highly multilingual,

with very few monolinguals and bilinguals. Future works is necessary to confirm that our results

are reproduced in the lower levels of the multilingual distribution.

756   **5. CONCLUSIONS**

In this paper, brain-behaviour associations for language-relevant skills and speech processing were

investigated using PLS correlation analysis. Participants with different levels of multilingual

experience and reading ability were included, and the analyses were performed first on the whole

sample and then on the subset of TRs only, to explore which associations were stable and which

were modulated by the presence or absence of DRs.

Overall, in both samples, a main component including cognitive, linguistic and

phonological measures was found to be associated with cortical areas involved in lexico/semantics,

combinatorial processing and amodal conceptual-semantic areas. Analyses excluding DRs suggest

that literacy and phonological processes may be more closely associated with language and

cognition in TRs than when DRs are included in the sample, pointing to more shared neural

resources between these processes in TRs. Furthermore, results from TRs only indicate that

additional components can be identified in populations without reading deficits, in particular, a

complementary component involving foreign speech processing and motor speed, automatization

and lexical access related to auditory and motor brain areas.

This work is a first step in exploring complex relationships underlying language use and

learning and domain-general and domain-specific non-linguistic skills, in a multivariate way. We

believe that language, as a multifaceted and multicomponent ability, should be studied

comprehensively, that it should include broad behavioural characterisation of speakers and that it

should ideally explore the relationship between different data modalities (e.g. brain function and

structure, functional and structural connectivity, genetics, etc.). We believe that this approach can

advance our understanding of language, of how individual differences arise and of how different

abilities interact, and hope that future work will continue to unveil new aspects of these complex

relationships.

6. ACKNOWLEDGEMENTS

We are very grateful for the statistical and methodological support for the PLS analysis offered by

Erik Ringen, Farnaz Delavari, and Maria Giulia Preti. We would like to thank Michael Erard,

Richard Simcott, Alexandre Teiga, Susan Fitzgerald, Lane Greene, Hugues Pluvinage and Swiss

in this study. We also thank Neiloufar Family and Sevil Maghsadhagh for rating our foreign speech

sound imitation task.

7. AUTHORS' CONTRIBUTIONS

writing—review & editing, I.B., A.R., O. K, R. B., N.G, funding acquisition, R.B. and N.G.

8. FUNDING

This work was supported by the Swiss National Science Foundation [Grant #100014_182381], and

by the NCCR Evolving Language, Swiss National Science Foundation [Agreement

#51NF40_180888].

9. SUPPLEMENTARY MATERIALS

Supplementary materials are openly available at the project's Open Science Framework page at

https://osf.io/4zcn2/?view_only=f1f5f11ba91c4e168e5321a198b4dc7a

10. REFERENCES

Alcock KJ, Krawczyk K. 2010. Individual differences in language development: relationship with

motor skill at 21 months. Dev Sci. 13(5):677–691. doi:10.1111/J.1467-7687.2009.00924.X.

[accessed 2024 Aug 23]. https://onlinelibrary.wiley.com/doi/full/10.1111/j.1467-

7687.2009.00924.x.

Aliko S, Wang B, Small SL, Skipper JI. 2023. The entire brain, more or less, is at work: 'Language

regions' are artefacts of averaging. bioRxiv.:2023.09.01.555886. doi:10.1101/2023.09.01.555886.

[accessed 2025 Mar 5]. https://www.biorxiv.org/content/10.1101/2023.09.01.555886v2.

Asano R, Boeckx C. 2015. Syntax in language and music: What is the right level of comparison?

Front Physiol. 6:140185. doi:10.3389/FPSYG.2015.00942/BIBTEX. [accessed 2025 Mar 6].

www.frontiersin.org.

Asano R, Boeckx C, Fujita K. 2022. Moving beyond domain-specific versus domain-general

options in cognitive neuroscience. Cortex. 154:259–268. doi:10.1016/J.CORTEX.2022.05.004.

Asano R, Boeckx C, Seifert U. 2021. Hierarchical control as a shared neurocognitive mechanism

for language and music. Cognition. 216:104847. doi:10.1016/J.COGNITION.2021.104847.

Bak TH, Chandran S. 2012. What wires together dies together: Verbs, actions and

neurodegeneration   in   motor   neuron   disease.   Cortex.   48(7):936–944.

doi:10.1016/J.CORTEX.2011.07.008.

Baker CM, Burks JD, Briggs RG, Sheets JR, Conner AK, Glenn CA, Sali G, McCoy TM, Battiste

JD, O'Donoghue DL, et al. 2018. A Connectomic Atlas of the Human Cerebrum—Chapter 3: The

Motor,   Premotor,   and   Sensory   Cortices.   Operative   Neurosurgery.   15(Suppl   1):S75.

doi:10.1093/ONS/OPY256. [accessed 2024 Aug 20]. /pmc/articles/PMC6887944/.

Bates E, Bretherton I, Sebestyen Snyder L. 1991. From First Words to Grammar: Individual

Differences and Dissociable Mechanisms - Elizabeth Bates, Inge Bretherton, Lynn Sebestyen

Snyder - Google Books. Cambridge University Press. [accessed 2024 Aug 25].

https://books.google.ch/books?hl=en&lr=&id=DBIQfHjSotEC&oi=fnd&pg=PR9&ots=SuhYruo

WUP&sig=b2oRCF9J-WWQ-L6_UuvABdqeBRU&redir_esc=y#v=onepage&q&f=false.

Beeson PM, Rapcsak SZ, Plante E, Chargualaf J, Chung A, Johnson SC, Trouard TP. 2003. The

neural substrates of writing: A functional magnetic resonance imaging study. Aphasiology. 17(6–

7):647–665.   doi:10.1080/02687030344000067.   [accessed   2024   Sep   1].

https://www.tandfonline.com/action/journalInformation?journalCode=paph20.

van Bergen E, Eveline L de Z, Sara A. Hart, Dorret I. Boomsma, Eco J. C. de Geus, Kees-Jan Kan.

2023.   Comorbidity   and   causality   among   ADHD,   dyslexia,   and   dyscalculia.

https://doi.org/10.31234/osf.io/epzgy.

Berthele R, Udry I. 2021. Individual differences in early instructed language learning: The role of

language aptitude, cognition, and motivation. Language science press.

Bickel B, Giraud AL, Zuberbühler K, van Schaik CP. 2024. Language follows a distinct mode of

extra-genomic evolution. Phys Life Rev. 50:211–225. doi:10.1016/J.PLREV.2024.08.003.

Caporale N, Dan Y. 2008. Spike timing-dependent plasticity: A Hebbian learning rule. Annu Rev

Neurosci.   31(Volume   31,   2008):25–46.

doi:10.1146/ANNUREV.NEURO.31.060407.125639/CITE/REFWORKS. [accessed 2024 Aug

29]. https://www.annualreviews.org/content/journals/10.1146/annurev.neuro.31.060407.125639.

Carruthers P. 2002. The cognitive functions of language. Behavioral and Brain Sciences.

25(6):657–674.   doi:10.1017/S0140525X02000122.   [accessed   2024   Aug   11].

https://www.cambridge.org/core/journals/behavioral-and-brain-sciences/article/abs/cognitive-

functions-of-language/0CDF19278D6DB1919B2DEE655893AB46.

Christiner M, Serrallach BL, Benner J, Bernhofs V, Schneider P, Renner J, Sommer-Lolei S, Groß

C. 2022. Examining Individual Differences in Singing, Musical and Tone Language Ability in

Adolescents and Young Adults with Dyslexia. Brain Sci. 12(6):744.

doi:10.3390/brainsci12060744. https://www.mdpi.com/2076-3425/12/6/744.

Cohen L, Dehaene S. 2004. Specialization within the ventral stream: the case for the visual word

form area. Neuroimage. 22(1):466–476. doi:10.1016/J.NEUROIMAGE.2003.12.049.

Cohn N, Schilperoord J. 2024. A Multimodal Language Faculty : A Cognitive Framework for

Human Communication. A Multimodal Language Faculty: A Cognitive Framework for Human

Communication.:1–352.

Corbetta M, Shulman GL. 2002. Control of goal-directed and stimulus-driven attention in the brain.

Nature Reviews Neuroscience 2002 3:3. 3(3):201–215. doi:10.1038/nrn755. [accessed 2024 Sep

2]. https://www.nature.com/articles/nrn755.

Coslett HB, Schwartz MF. 2018. The parietal lobe and language. Handb Clin Neurol. 151:365–

375. doi:10.1016/B978-0-444-63622-5.00018-8.

Coumel M, Groß C, Sommer-Lolei S, Christiner M. 2023. The Contribution of Music Abilities and

Phonetic Aptitude to L2 Accent Faking Ability. Languages. 8(1):68.

doi:10.3390/languages8010068.

Crystal D, Robins RH. 2024 Jul 25. Language | Definition, Types, Characteristics, Development,

& Facts | Britannica. Encyclopaedia Britannica. [accessed 2024 Aug 5].

https://www.britannica.com/topic/language.

Davis MH. 2016 Jan 1. The Neurobiology of Lexical Access. Neurobiology of Language.:541–

555. doi:10.1016/B978-0-12-407794-2.00044-4.

Davis MH, Johnsrude IS. 2007. Hearing speech sounds: Top-down influences on the interface

between   audition   and   speech   perception.   Hear   Res.   229(1–2):132–147.

doi:10.1016/J.HEARES.2007.01.014.

Decarli G, Franchin L, Vitali F. 2024. Motor skills and capacities in developmental dyslexia: A

systematic review and meta-analysis. Acta Psychol (Amst). 246:104269.

Dehaene S. 2005. Evolution of Human Cortical Circuits for Reading and Arithmetic: The Neuronal

Recycling   Hypothesis.   From   Monkey   Brain   to   Human   Brain.:133–158.

doi:10.7551/MITPRESS/3136.003.0012.   [accessed   2024   Aug   25].

https://direct.mit.edu/books/edited-volume/2477/chapter/66073/Evolution-of-Human-Cortical-

Circuits-for-Reading.

Dehaene S, Cohen L, Morais J, Kolinsky R. 2015. Illiterate to literate: behavioural and cerebral

changes induced by reading acquisition. Nature Reviews Neuroscience 2015 16:4. 16(4):234–244.

doi:10.1038/nrn3924. [accessed 2024 Aug 25]. https://www.nature.com/articles/nrn3924.

Dehaene-Lambertz G, Monzalvo K, Dehaene S. 2018. The emergence of the visual word form:

Longitudinal evolution of category-specific ventral visual areas during reading acquisition. PLoS

Biol.   16(3):e2004103.   doi:10.1371/JOURNAL.PBIO.2004103.   [accessed   2024   Jul   14].

https://journals.plos.org/plosbiology/article?id=10.1371/journal.pbio.2004103.

Demanet J, Liefooghe B, Hartstra E, Wenke D, De Houwer J, Brass M. 2016. There is more into

'doing' than 'knowing': The function of the right inferior frontal sulcus is specific for

implementing   versus   memorising   verbal   instructions.   Neuroimage.   141:350–356.

doi:10.1016/J.NEUROIMAGE.2016.07.059.

Díaz B, Cordero G, Hoogendoorn J, Sebastian-Galles N. 2022. Second-language phoneme learning

positively relates to voice recognition abilities in the native language: Evidence from behavior and

brain potentials. Front Psychol. 13:1008963. doi:10.3389/FPSYG.2022.1008963/BIBTEX.

Diedrichsen J, Balsters JH, Flavell J, Cussans E, Ramnani N. 2009. A probabilistic MR atlas of the

human   cerebellum.   Neuroimage.   46(1):39–46.   doi:10.1016/J.NEUROIMAGE.2009.01.045.

[accessed 2024 Jun 30]. https://pubmed.ncbi.nlm.nih.gov/19457380/.

Donnelly S, Kidd E. 2021. The Longitudinal Relationship Between Conversational Turn-Taking

and   Vocabulary   Growth   in   Early   Language   Development.   Child   Dev.   92(2):609–625.

doi:10.1111/CDEV.13511.   [accessed   2024   Aug   28].

https://onlinelibrary.wiley.com/doi/full/10.1111/cdev.13511.

Doughty CJ, Campbell SG, Mislevy MA, Bunting MF, Bowles AR, Koeth JT. 2010. Predicting

Near-Native Ability : The Factor Structure and Reliability of Hi-LAB. Selected proceedings of the

2008 Second Language research Forum.:10–31. http://www.lingref.com/cpp/slrf/2008/index.html.

Drijvers L, Small SL, Skipper JI. 2025. Language is widely distributed throughout the brain. Nature

Reviews Neuroscience 2025 26:3. 26(3):189–189. doi:10.1038/s41583-024-00903-0. [accessed

2025 Mar 5]. https://www.nature.com/articles/s41583-024-00903-0.

Drzezga A. 2018. The Network Degeneration Hypothesis: Spread of Neurodegenerative Patterns

Along   Neuronal   Brain   Networks.   Journal   of   Nuclear   Medicine.   59(11):1645–1648.

doi:10.2967/JNUMED.117.206300.   [accessed   2024   Sep   5].

https://jnm.snmjournals.org/content/59/11/1645.

Fedorenko E, Hsieh PJ, Nieto-Castañón A, Whitfield-Gabrieli S, Kanwisher N. 2010. New method

for fMRI investigations of language: Defining ROIs functionally in individual subjects. J

Neurophysiol.   104(2):1177–1194.

doi:10.1152/JN.00032.2010/SUPPL_FILE/DESCRIPTIONS.DOCX. [accessed 2024 Jul 12].

https://journals.physiology.org/doi/10.1152/jn.00032.2010.

Fedorenko E, Ivanova AA, Regev TI. 2024. The language network as a natural kind within the

broader landscape of the human brain. Nature Reviews Neuroscience 2024 25:5. 25(5):289–312.

doi:10.1038/s41583-024-00802-4.   [accessed   2024   Jul   14].

https://www.nature.com/articles/s41583-024-00802-4.

Fedorenko E, Varley R. 2016. Language and thought are not the same thing: evidence from

neuroimaging   and   neurological   patients.   Ann   N   Y   Acad   Sci.   1369(1):132.

doi:10.1111/NYAS.13046. [accessed 2023 May 22]. /pmc/articles/PMC4874898/.

Feng G, Ou J, Gan Z, Jia X, Meng D, Wang S, Wong PCM. 2021. Neural fingerprints underlying

individual   language   learning   profiles.   Journal   of   Neuroscience.   41(35):7372–7387.

doi:10.1523/JNEUROSCI.0415-21.2021.

Fitch WT. 2011. The evolution of syntax: An exaptationist perspective. Front Evol Neurosci.

3(DEC):18409.   doi:10.3389/FNEVO.2011.00009/BIBTEX.   [accessed   2025   Mar   14].

www.frontiersin.org.

Fitch WT. 2012. Evolutionary Developmental Biology and Human Language Evolution:

Constraints on Adaptation. Evolutionary Biology 2012 39:4. 39(4):613–637. doi:10.1007/S11692-

012-9162-Y. [accessed 2025 Mar 14]. https://link.springer.com/article/10.1007/s11692-012-9162-

y.

Fitch WT, Hauser MD, Chomsky N. 2005. The evolution of the language faculty: Clarifications

and implications. Cognition. 97(2):179–210. doi:10.1016/J.COGNITION.2005.02.005.

Fitch WT, Martins MD. 2014. Hierarchical processing in music, language, and action: Lashley

revisited. Ann N Y Acad Sci. 1316(1):87–104. doi:10.1111/NYAS.12406. [accessed 2025 Mar 14].

https://onlinelibrary.wiley.com/doi/full/10.1111/nyas.12406.

Fonseca-Mora M, Toscano-Fuentes C, Wermke K. 2011. Melodies That Help: The Relation

between Language Aptitude and Musical Intelligence. International Journal of English Studies.

22(1):101–118.

Friederici AD. 2006. Broca's Area and the Ventral Premotor Cortex in Language: Functional

Differentiation and Specificity. Cortex. 42(4):472–475. doi:10.1016/S0010-9452(08)70380-0.

Fuhrmeister P, Phillips MC, McCoach DB, Myers EB. 2023. Relationships between native and

non-native speech perception. J Exp Psychol Learn Mem Cogn. 49(7):1161–1175.

doi:10.1037/XLM0001213. [accessed 2024 Sep 6]. https://pubmed.ncbi.nlm.nih.gov/36757985/.

Gainotti G. 2017. The Differential Contributions of Conceptual Representation Format and

Language Structure to Levels of Semantic Abstraction Capacity. Neuropsychol Rev. 27(2):134–

145.   doi:10.1007/S11065-016-9339-8/METRICS.   [accessed   2024   Jul   14].

https://link.springer.com/article/10.1007/s11065-016-9339-8.

Glasser MF, Coalson TS, Robinson EC, Hacker CD, Harwell J, Yacoub E, Ugurbil K, Andersson
J, Beckmann CF, Jenkinson M, et al. 2016. A multi-modal parcellation of human cerebral cortex.
Nature 2016 536:7615. 536(7615):171–178. doi:10.1038/nature18933. [accessed 2024 Jun 30].
https://www.nature.com/articles/nature18933.

Gonzalez SL, Alvarez V, Nelson EL. 2019. Do Gross and Fine Motor Skills Differentially
Contribute to Language Outcomes? A Systematic Review. Front Psychol. 10:490791.
doi:10.3389/FPSYG.2019.02670/BIBTEX. [accessed 2024 Aug 23]. www.frontiersin.org.

Grabner G, Janke AL, Budge MM, Smith D, Pruessner J, Collins DL. 2006. Symmetric atlasing
and model based segmentation: An application to the hippocampus in older adults. Lecture Notes
in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture
Notes in Bioinformatics). 4191 LNCS-II:58–66. doi:10.1007/11866763_8.

Hagoort P. 2014. Nodes and networks in the neural architecture for language: Broca's region and
beyond. Curr Opin Neurobiol. 28:136–141. doi:10.1016/J.CONB.2014.07.013.

Hagoort P. 2017. The core and beyond in the language-ready brain. Neurosci Biobehav Rev.
81:194–204. doi:10.1016/J.NEUBIOREV.2017.01.048.

Hagoort P, Van Berkum J. 2007. Beyond the sentence given. Philosophical Transactions of the
Royal Society B: Biological Sciences. 362(1481):801. doi:10.1098/RSTB.2007.2089. [accessed
2024 Aug 28]. /pmc/articles/PMC2429998/.

Hagoort P, Özyürek A. 2024. Extending the Architecture of Language From a Multimodal
Perspective. Top Cogn Sci. 00:1–11. doi:10.1111/TOPS.12728. [accessed 2025 Mar 14].
https://onlinelibrary.wiley.com/doi/full/10.1111/tops.12728.

Hancock R, Pugh KR, Hoeft F. 2017. Neural Noise Hypothesis of Developmental Dyslexia. Trends

Cogn Sci. 21(6):434–448. doi:10.1016/J.TICS.2017.03.008. [accessed 2025 Mar 28].

https://pubmed.ncbi.nlm.nih.gov/28400089/.

Hasson U, Egidi G, Marelli M, Willems RM. 2018. Grounding the neurobiology of language in

first principles: The necessity of non-language-centric explanations for language comprehension.

Cognition. 180:135–157. doi:10.1016/J.COGNITION.2018.06.018.

Hauser MD, Chomsky N, Fitch WT. 2002. The Faculty of Language: What Is It, Who Has It, and

How Did It Evolve? Science (1979). 298(5598):1569–1579.

doi:10.1126/SCIENCE.298.5598.1569. [accessed 2024 Aug 26].

https://www.science.org/doi/10.1126/science.298.5598.1569.

Hein G, Knight RT. 2008. Superior Temporal Sulcus—It's My Area: Or Is It? J Cogn Neurosci.

20(12):2125–2136. doi:10.1162/JOCN.2008.20148. [accessed 2024 Jul 14].

https://dx.doi.org/10.1162/jocn.2008.20148.

Hickok G, Poeppel D. 2004. Dorsal and ventral streams: a framework for understanding aspects of

the functional anatomy of language. Cognition. 92(1–2):67–99.

doi:10.1016/J.COGNITION.2003.10.011.

Hickok G, Poeppel D. 2007. The cortical organization of speech processing. Nature Reviews

Neuroscience 2007 8:5. 8(5):393–402. doi:10.1038/nrn2113. [accessed 2024 Jul 14].

https://www.nature.com/articles/nrn2113.

Hintz F, McQueen J, Meyer AS. 2024. Using psychometric network analysis to examine the

components of spoken word recognition. J Cogn.

Hintz F, Voeten CC, Dobó D, Lukics KS, Lukács Á. 2024. The role of general cognitive skills in

integrating visual and linguistic information during sentence comprehension: Individual

differences across the lifespan. Sci Rep.

Horn A. 2016 Jun. HCP-MMP1.0 projected on MNI2009a GM (volumetric) in NIfTI format.

doi:10.6084/m9.figshare.3501911.v5.   https://figshare.com/articles/dataset/HCP-

MMP1_0_projected_on_MNI2009a_GM_volumetric_in_NIfTI_format/3501911.

Igelström KM, Graziano MSA. 2017. The inferior parietal lobule and temporoparietal junction: A

network   perspective.   Neuropsychologia.   105:70–83.

doi:10.1016/J.NEUROPSYCHOLOGIA.2017.01.001.

Janacsek K, Evans TM, Kiss M, Shah L, Blumenfeld H, Ullman MT. 2022. Subcortical Cognition:

The   Fruit   Below   the   Rind.   Annu   Rev   Neurosci.   45(Volume   45,   2022):361–386.

doi:10.1146/ANNUREV-NEURO-110920-013544/1.   [accessed   2025   Mar   12].

https://www.annualreviews.org/content/journals/10.1146/annurev-neuro-110920-013544.

Japee S, Holiday K, Satyshur MD, Mukai I, Ungerleider LG. 2015. A role of right middle frontal

gyrus in reorienting of attention: A case study. Front Syst Neurosci. 9(MAR):122128.

doi:10.3389/FNSYS.2015.00023/ABSTRACT. [accessed 2024 Sep 2]. www.frontiersin.org.

Jouravlev O, Mineroff Z, Blank IA, Fedorenko E. 2020. The Small and Efficient Language

Network   of   Polyglots   and   Hyper-polyglots.   Cerebral   Cortex.   31(1):62–76.

doi:10.1093/cercor/bhaa205. https://doi.org/10.1093/cercor/bhaa205.

Kellmeyer P, Ziegler W, Peschke C, Juliane E, Schnell S, Baumgaertner A, Weiller C, Saur D.

2013. Fronto-parietal dorsal and ventral pathways in the context of different linguistic

manipulations. Brain Lang. 127(2):241–250. doi:10.1016/J.BANDL.2013.09.011.

Kidd E, Donnelly S. 2020. Individual Differences in First Language Acquisition. Annu Rev

Linguist. 6:319–340. doi:10.1146/annurev-linguistics-011619-030326.

Kidd E, Donnelly S, Christiansen MH. 2018. Individual Differences in Language Acquisition and

Processing. Trends Cogn Sci. 22(2):154–169. doi:10.1016/J.TICS.2017.11.006. [accessed 2024

Aug 15]. http://www.cell.com/article/S1364661317302462/fulltext.

Kilian-Hütten N, Vroomen J, Formisano E. 2011. Brain activation during audiovisual exposure

anticipates future perception of ambiguous speech. Neuroimage. 57(4):1601–1607.

doi:10.1016/J.NEUROIMAGE.2011.05.043.   [accessed   2024   Sep   1].

https://pubmed.ncbi.nlm.nih.gov/21664279/.

Koyama MS, O'Connor D, Shehzad Z, Milham MP. 2017. Differential contributions of the middle

frontal gyrus functional connectivity to literacy and numeracy. Scientific Reports 2017 7:1. 7(1):1–

13.   doi:10.1038/s41598-017-17702-6.   [accessed   2024   Sep   2].

https://www.nature.com/articles/s41598-017-17702-6.

Krishnan A, Williams LJ, Mcintosh AR, Abdi H. 2010. Partial Least Squares (PLS) methods for

neuroimaging: A tutorial and review. doi:10.1016/j.neuroimage.2010.07.034.

Krishnan A, Williams LJ, McIntosh AR, Abdi H. 2011. Partial Least Squares (PLS) methods for

neuroimaging: A   tutorial   and   review.   Neuroimage.   56(2):455–475.

doi:10.1016/J.NEUROIMAGE.2010.07.034.

Kuhl U, Neef NE, Kraft I, Schaadt G, Dörr L, Brauer J, Czepezauer I, Müller B, Wilcke A, Kirsten

H, et al. 2020. The emergence of dyslexia in the developing brain. Neuroimage. 211:116633.

doi:10.1016/J.NEUROIMAGE.2020.116633.

Lambon Ralph MA, Jefferies E, Patterson K, Rogers TT. 2016. The neural and computational bases

of semantic cognition. Nature Reviews Neuroscience 2016 18:1. 18(1):42–55.

doi:10.1038/nrn.2016.150. [accessed 2024 Jul 14]. https://www.nature.com/articles/nrn.2016.150.

Lambon Ralph MA, Sage K, Jones RW, Mayberry EJ. 2010. Coherent concepts are computed in

the anterior temporal lobes. Proc Natl Acad Sci U S A. 107(6):2717.

doi:10.1073/PNAS.0907307107. [accessed 2024 Jul 14]. /pmc/articles/PMC2823909/.

Levinson SC, Holler J. 2014. The origin of human multi-modal communication. Philosophical

Transactions of the Royal Society B: Biological Sciences. 369(1651).

doi:10.1098/RSTB.2013.0302. [accessed 2024 Aug 5].

https://royalsocietypublishing.org/doi/10.1098/rstb.2013.0302.

Libertus K, Violi DA. 2016. Sit to talk: Relation between motor skills and language development

in infancy. Front Psychol. 7(MAR):187259. doi:10.3389/FPSYG.2016.00475/BIBTEX. [accessed

2024 Aug 23]. www.frontiersin.org.

Lutkenhoff ES, Rosenberg M, Chiang J, Zhang K, Pickard JD, Owen AM, Monti MM. 2014.

Optimized Brain Extraction for Pathological Brains (optiBET). PLoS One. 9(12):e115551.

doi:10.1371/JOURNAL.PONE.0115551. [accessed 2024 Jul 12].

https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0115551.

Lyon GR, Shaywitz SE, Bennett AS. 2003. A Definition of Dyslexia. Ann Dyslexia. 53(1):1–14.

doi:10.1002/9781118660584.ese0145.

Mahowald K, Fedorenko E. 2016. Reliable individual-level neural markers of high-level language

processing: A necessary precursor for relating neural variability to behavioral and genetic

variability. Neuroimage. 139:74–93. doi:10.1016/J.NEUROIMAGE.2016.05.073.

Majid A, Bowerman M, Kita S, Haun DBM, Levinson SC. 2004. Can language restructure

cognition? The case for space. Trends Cogn Sci. 8(3):108–114. doi:10.1016/j.tics.2004.01.003.

[accessed 2024 Aug 11]. https://pubmed.ncbi.nlm.nih.gov/15301750/.

Makris N, Goldstein JM, Kennedy D, Hodge SM, Caviness VS, Faraone S V., Tsuang MT,

Seidman LJ. 2006. Decreased volume of left and total anterior insular lobule in schizophrenia.

Schizophr Res. 83(2–3):155–171. doi:10.1016/J.SCHRES.2005.11.020. [accessed 2024 Jun 30].

https://pubmed.ncbi.nlm.nih.gov/16448806/.

Malik-Moraleda S, Ayyash D, Gallamp J, Affourtit J, Hoffmann M, Mineroff Z, Jouravlev O,

Fedorenko E. 2022 Jul 18. An investigation across 45 languages and 12 language families reveals

a universal language network. Nature Neuroscience 2022.:1–6. doi:10.1038/s41593-022-01114-5.

[accessed 2022 Jul 28]. https://www.nature.com/articles/s41593-022-01114-5.

Marchand-Krynski M-È, Morin-Moncet O, Bélanger A-M, Beauchamp MH, Leonard G. 2017.

Shared and differentiated motor skill impairments in children with dyslexia and/or attention deficit

disorder: From simple to complex sequential coordination. PLoS One. 12(5):e0177490.

McClelland JL, Mirman D, Holt LL. 2006. Are there interactive processes in speech perception?

Trends Cogn Sci. 10(8):363. doi:10.1016/J.TICS.2006.06.007. [accessed 2024 Sep 6].

/pmc/articles/PMC3523348/.

McIntosh AR, Lobaugh NJ. 2004a. Partial least squares analysis of neuroimaging data: applications

and advances. Neuroimage. 23(SUPPL. 1):S250–S263.

doi:10.1016/J.NEUROIMAGE.2004.07.020.

McIntosh AR, Lobaugh NJ. 2004b. Partial least squares analysis of neuroimaging data:

applications and advances. Neuroimage. 23(SUPPL. 1):S250–S263.

doi:10.1016/J.NEUROIMAGE.2004.07.020.

McNorgan C, Awati N, Desroches AS, Booth JR. 2014. Multimodal Lexical Processing in

Auditory Cortex Is Literacy Skill Dependent. Cerebral Cortex. 24(9):2464–2475.

doi:10.1093/CERCOR/BHT100. [accessed 2024 Aug 19].

https://dx.doi.org/10.1093/cercor/bht100.

Mihalik A, Chapman J, Adams RA, Winter NR, Ferreira FS, Shawe-Taylor J, Mourão-Miranda J.

2022. Canonical Correlation Analysis and Partial Least Squares for Identifying Brain–Behavior

Associations: A Tutorial and a Comparative Study. Biol Psychiatry Cogn Neurosci Neuroimaging.

7(11):1055–1067. doi:10.1016/J.BPSC.2022.07.012.

Munakata Y, Pfaffly J. 2004. Hebbian learning and development. Dev Sci. 7(2):141–148.

doi:10.1111/J.1467-7687.2004.00331.X. [accessed 2024 Aug 29].

https://onlinelibrary.wiley.com/doi/full/10.1111/j.1467-7687.2004.00331.x.

Nardo D, Reiterer SM. 2009. Musicality and phonetic language aptitude. In: Dogil G RS, editor.

Language Talent and Brain Activity. Berlin-NewYork: Mouton De Gruyter. p. 213–256.

Nayak S, Coleman PL, Ladányi E, Nitin R, Gustavson DE, Fisher SE, Magne CL, Gordon RL.

2022. The Musical Abilities, Pleiotropy, Language, and Environment (MAPLE) Framework for

Understanding Musicality-Language Links Across the Lifespan. Neurobiology of Language.

3(4):615–664. doi:10.1162/NOL_A_00079. [accessed 2024 Aug 15].

https://dx.doi.org/10.1162/nol_a_00079.

Overy K. 2003. Dyslexia and Music. Ann N Y Acad Sci. 999(1):497–505.

doi:10.1196/ANNALS.1284.060. [accessed 2025 Mar 28].

https://onlinelibrary.wiley.com/doi/full/10.1196/annals.1284.060.

Overy K, Nicolson RI, Fawcett AJ, Clarke EF. 2003. Dyslexia and music: measuring musical

timing skills. Dyslexia. 9(1):18–36. doi:10.1002/DYS.233. [accessed 2025 Mar 28].

https://onlinelibrary.wiley.com/doi/full/10.1002/dys.233.

Pallier C, Devauchelle AD, Dehaene S. 2011. Cortical representation of the constituent structure

of sentences. Proc Natl Acad Sci U S A. 108(6):2522–2527.

doi:10.1073/PNAS.1018711108/SUPPL_FILE/SAPP.PDF. [accessed 2024 Jul 12].

https://www.pnas.org/doi/abs/10.1073/pnas.1018711108.

Penolazzi B, Hauk O, Pulvermüller F. 2007. Early semantic context integration and lexical access

as revealed by event-related brain potentials. Biol Psychol. 74(3):374–388.

doi:10.1016/J.BIOPSYCHO.2006.09.008.

Preston JL, Molfese PJ, Frost SJ, Mencl WE, Fulbright RK, Hoeft F, Landi N, Shankweiler D,

Pugh KR. 2016. Print-Speech Convergence Predicts Future Reading Outcomes in Early Readers.

Psychol Sci. 27(1):75. doi:10.1177/0956797615611921. [accessed 2024 Sep 6].

/pmc/articles/PMC4713346/.

Price CJ. 2012. A review and synthesis of the first 20 years of PET and fMRI studies of heard

speech, spoken language and reading. Neuroimage. 62(2):816–847.

doi:10.1016/J.NEUROIMAGE.2012.04.062.

Pruim RHR, Mennes M, Buitelaar JK, Beckmann CF. 2015. Evaluation of ICA-AROMA and

alternative strategies for motion artifact removal in resting state fMRI. Neuroimage. 112:278–287.

doi:10.1016/J.NEUROIMAGE.2015.02.063.

Purpura DJ, Lonigan CJ. 2013. Informal Numeracy Skills: The Structure and Relations Among

Numbering, Relations, and Arithmetic Operations in Preschool. Am Educ Res J. 50(1):178–209.

doi:10.3102/0002831212465332/ASSET/IMAGES/LARGE/10.3102_0002831212465332-

FIG1.JPEG. [accessed 2024 Sep 6].

https://journals.sagepub.com/doi/full/10.3102/0002831212465332?casa_token=VWv3RKAdmy

UAAAAA%3A8SnfgXuCiUUvfzOgmkBjxvr77z03Y78djg3y7f0OkVPoKxZU8ffMt_7zvnTMJu

ZI4kFbGNPP1aUwhr4.

Rampinini A, Balboni I, Golestani N, Berthele R. 2024. A behavioural exploration of language

aptitude and experience, cognition and more using Graph Analysis. Brain Res. 1842:149109.

doi:10.1016/J.BRAINRES.2024.149109.

Rauschecker JP. 2012. Ventral and dorsal streams in the evolution of speech and language. Front

Evol Neurosci. 4(MAY):18154. doi:10.3389/FNEVO.2012.00007/BIBTEX. [accessed 2025 Mar

11]. www.frontiersin.org.

Rauschecker JP, Scott SK. 2009. Maps and streams in the auditory cortex: nonhuman primates

illuminate human speech processing. Nature Neuroscience 2009 12:6. 12(6):718–724.

doi:10.1038/nn.2331. [accessed 2025 Mar 11]. https://www.nature.com/articles/nn.2331.

Rauschecker JP, Scott SK. 2016. Pathways and Streams in the Auditory Cortex: An Update on

How Work in Nonhuman Primates has Contributed to Our Understanding of Human Speech

Processing. Neurobiology of Language.:287–298. doi:10.1016/B978-0-12-407794-2.00024-9.

Rolka EJ, Silverman MJ. 2015. A systematic review of music and dyslexia. Arts Psychother.

46:24–32.   doi:10.1016/j.aip.2015.09.002.

https://linkinghub.elsevier.com/retrieve/pii/S0197455615000684.

Rothmayr C, Sodian B, Hajak G, Döhnel K, Meinhardt J, Sommer M. 2011. Common and distinct

neural networks for false-belief reasoning and inhibitory control. Neuroimage. 56(3):1705–1713.

doi:10.1016/J.NEUROIMAGE.2010.12.052.

Rueckl JG, Paz-Alonso PM, Molfese PJ, Kuo WJ, Bick A, Frost SJ, Hancock R, Wu DH, Einar

Mencl W, Duñabeitia JA, et al. 2015. Universal brain signature of proficient reading: Evidence

from four contrasting languages. Proc Natl Acad Sci U S A. 112(50):15510–15515.

doi:10.1073/PNAS.1509321112/SUPPL_FILE/PNAS.201509321SI.PDF. [accessed 2024 Feb 6].

https://www.pnas.org/doi/abs/10.1073/pnas.1509321112.

Ruland SH, Palomero-Gallagher N, Hoffstaedter F, Eickhoff SB, Mohlberg H, Amunts K. 2022.

The inferior frontal sulcus: Cortical segregation, molecular architecture and function. Cortex.

153:235–256. doi:10.1016/J.CORTEX.2022.03.019.

Rutten S, Santoro R, Hervais-Adelman A, Formisano E, Golestani N. 2019. Cortical encoding of

speech enhances task-relevant acoustic information. Nature Human Behaviour 2019 3:9. 3(9):974–

987.   doi:10.1038/s41562-019-0648-9.   [accessed   2024   Sep   6].

https://www.nature.com/articles/s41562-019-0648-9.

Schell M, Zaccarella E, Friederici AD. 2017. Differential cortical contribution of syntax and

semantics: An fMRI study on two-word phrasal processing. Cortex. 96:105–120.

doi:10.1016/J.CORTEX.2017.09.002.

Schwering SC, MacDonald MC. 2020. Verbal Working Memory as Emergent from Language

Comprehension   and   Production.   Front   Hum   Neurosci.   14:488865.

doi:10.3389/FNHUM.2020.00068/BIBTEX. [accessed 2024 Aug 28]. www.frontiersin.org.

Shankweiler D, Mencl WE, Braze D, Tabor W, Pugh KR, Fulbright RK. 2008. Reading Differences

and Brain: Cortical Integration of Speech and Print in Sentence Processing Varies With Reader

Skill. Dev Neuropsychol. 33(6):745–775. doi:10.1080/87565640802418688. [accessed 2024 Sep

12]. https://www.tandfonline.com/doi/abs/10.1080/87565640802418688.

Simmons FR, Singleton C. 2008. Do weak phonological representations impact on arithmetic

development? A review of research into arithmetic and dyslexia. Dyslexia. 14(2):77–94.

doi:10.1002/DYS.341.   [accessed   2024   Aug   16].

https://onlinelibrary.wiley.com/doi/full/10.1002/dys.341.

Singleton D. 2017. Language aptitude: Desirable trait or acquirable attribute? Studies in Second
Language Learning and Teaching. 7(1):89–103. doi:10.14746/ssllt.2017.7.1.5.

Sydnor VJ, Larsen B, Bassett DS, Alexander-Bloch A, Fair DA, Liston C, Mackey AP, Milham
MP, Pines A, Roalf DR, et al. 2021. Neurodevelopment of the association cortices: Patterns,
mechanisms, and implications for psychopathology. Neuron. 109(18):2820–2846.
doi:10.1016/J.NEURON.2021.06.016.

Turker S, Hartwigsen G. 2021. Exploring the neurobiology of reading through non-invasive brain
stimulation: A review. Cortex. 141:497–521. doi:10.1016/J.CORTEX.2021.05.001.

Turker S, Kuhnke P, Eickhoff SB, Caspers S, Hartwigsen G. 2023. Cortical, Subcortical, and
Cerebellar Contributions to Language Processing: A Meta-Analytic Review of 403 Neuroimaging
Experiments. Psychol Bull. 149(11–12):699–723. doi:10.1037/BUL0000403.

Turker S, Reiterer SM. 2021. Brain, musicality, and language aptitude: A complex interplay. Annu
Rev   Appl   Linguist.   41:95–107.   doi:10.1017/S0267190520000148.
https://www.cambridge.org/core/product/identifier/S0267190520000148/type/journal_article.

Turker S, Reiterer SM, Seither-Preisler A, Schneider P. 2017. "When Music Speaks": Auditory
Cortex Morphology as a Neuroanatomical Marker of Language Aptitude and Musicality. Front
Psychol.   8:2096.   doi:10.3389/fpsyg.2017.02096.
http://journal.frontiersin.org/article/10.3389/fpsyg.2017.02096/full.

Udry I, Berthele R. 2021. The smart, the motivated and the self-confident: The role of language
aptitude, cognition, and affective variables in early instructed foreign language learning. In:
Berthele R, Udry I, editors. Individual Differences in Early Instructed Language Learning: The

Role of Language Aptitude, Cognition, and Motivation. Berlin: Language Science Press. (EuroSla

Studies Series). p. 71–90. https://doi.org/10.5281/zenodo.5464749.

Ullman MT. 2004. Contributions of memory circuits to language: The declarative/procedural

model. Cognition. 92(1–2):231–270. doi:10.1016/j.cognition.2003.10.008.

Ullman MT. 2016. Chapter 76 - The Declarative/Procedural Model: A Neurobiological Model of

Language Learning, Knowledge, and Use. In: Hickok G, Small SL, editors. Neurobiology of

Language.   San   Diego: Academic   Press.   p.   953–968.

https://www.sciencedirect.com/science/article/pii/B9780124077942000766.

Valian V. 2016. Putting together bilingualism and executive function. Linguist Approaches Biling.

6(5):565–574.

van Viersen S, Altani A, de Jong PF, Protopapas A. 2024 Apr 3. Between-word processing and

text-level skills contributing to fluent reading of (non)word lists and text. Read Writ.:1–27.

doi:10.1007/S11145-024-10533-8/TABLES/7.   [accessed   2025   Mar   17].

https://link.springer.com/article/10.1007/s11145-024-10533-8.

Wen Z. 2012. Foreign language aptitude. ELT Journal. 66(2):233–235. doi:10.1093/elt/ccr068.

Wingfield A, Grossman M. 2006. Language and the aging brain: Patterns of neural compensation

revealed   by   functional   brain   imaging.   J   Neurophysiol.   96(6):2830–2839.

doi:10.1152/JN.00628.2006/ASSET/IMAGES/LARGE/Z9K0120678070004.JPEG.   [accessed

2025 Mar 28]. https://journals.physiology.org/doi/10.1152/jn.00628.2006.

Wong PCM, Ettlinger M, Sheppard JP, Gunasekera GM, Dhar S. 2010. Neuroanatomical

characteristics and speech perception in noise in older adults. Ear Hear. 31(4):471–479.

doi:10.1097/AUD.0B013E3181D709C2. [accessed 2025 Mar 28]. https://journals.lww.com/ear-

hearing/fulltext/2010/08000/neuroanatomical_characteristics_and_speech.3.aspx.

Zaretsky E, Kraljevic JK, Core C, Lencek M. 2009. Literacy predictors and early reading and

spelling skills as a factor of orthography: Cross-linguistic evidence. Writ Lang Lit. 12(1):52–81.

doi:10.1075/WLL.12.1.03ZAR.   [accessed   2025   Mar   17].   https://www.jbe-

platform.com/content/journals/10.1075/wll.12.1.03zar.

Zhou J, Gennatas ED, Kramer JH, Miller BL, Seeley WW. 2012. Predicting Regional

Neurodegeneration from the Healthy Brain Functional Connectome. Neuron. 73(6):1216–1227.

doi:10.1016/J.NEURON.2012.03.004/ATTACHMENT/289733FF-E836-47EB-AA8C-

AA74A23CCAC0/MMC1.PDF.   [accessed   2024   Sep   5].

http://www.cell.com/article/S0896627312002279/fulltext.

1221