Responses to conflicting binocular stimuli in mouse primary visual cortex

Abbreviated title:

V1 responses to conflicting binocular stimuli

Authors

Montgomery, Daniel P.[1]; Bowen, Daniel A.[2,3]; Wu, Jin[4]; Bear, Mark F.[3]; Gaier, Eric D.[2,3,5*]

Affiliations

[1] Center for Neuroscience Research, Children's National Hospital, Washington, DC

[2] Department of Ophthalmology, Boston Children's Hospital, Boston, MA

[3] Picower Institute for Learning and Memory, Massachusetts Institute of Technology, Cambridge, MA

[4] Brandeis Neuroscience Graduate Program, Brandeis University, Waltham, MA

[5] Harvard Medical School, Boston, MA

*Corresponding Author

Correspondence

Eric D. Gaier MD PhD

Boston Children's Hospital

300 Longwood Ave

Boston, MA 02115

Phone: 617-355-6401

Email: eric.gaier@childrens.harvard.edu

Pages: 32 pages of text (not including figures)

Figures: 9 main figures, 2 supplemental figures

Tables: 1

Word counts:

 Abstract: 219

 Introduction: 605

 Results: 2398

 Discussion: 2041

 Materials and methods: 2723

Key words:

Binocular vision, interocular suppression, binocular rivalry

Conflict of Interest

No relevant conflicting relationship exists for any author with regards to this work. DPM, DAB,

JW: none. MFB: Luminopia (equity). EDG: Luminopia (equity, royalties), Stoke Therapeutics

(consultant), Neurofieldz (consultant).

Acknowledgements:

Financial Support: NIH R01 EY023037 (MFB), NIH K08 EY030164 (EDG), Massachusetts Lions Eye

Research Fund (EDG)

IMPACT STATEMENT

Distinct forms of binocular conflict engage separate circuit mechanisms in mouse

primary visual cortex, revealing how interocular disparities shape population activity through

feedforward and disinhibitory processes.

ABSTRACT

Binocular vision requires that the brain integrate input from both eyes to form a unified percept. Small interocular differences support depth perception (stereopsis), while larger disparities can cause diplopia or binocular rivalry. The neural mechanisms by which early visual circuits process concordant versus conflicting binocular signals remain incompletely understood, particularly in the case of large disparities. Here, we used visually evoked potential (VEP) recordings, unit recordings, and 2-photon calcium imaging in the binocular region of mouse primary visual cortex (bV1) to examine how distinct forms of binocular disparity engage local circuits. Using a dichoptic display, we found that interocular phase disparities reduced VEP magnitude through decreased neuronal firing early in the response (40-80 ms after stimulus onset). In contrast, orientation disparities also decreased VEP magnitude, but via increased firing later in the response (100-200 ms). This late activity was enhanced in both regular-spiking (putative excitatory) and fast-spiking (putative parvalbumin-positive inhibitory) units. In contrast, calcium imaging revealed that somatostatin-positive interneurons were suppressed during orientation conflict. These findings suggest that phase differences suppress bV1 responses via feedforward mechanisms, while orientation disparities prolong activity through disinhibition mediated by somatostatin-positive interneurons. Our results identify distinct circuit pathways recruited by different forms of binocular conflict, clarify how early visual cortex contributes to binocular integration, and provide a foundation for investigating perceptual suppression and rivalry.

INTRODUCTION

Visual processing of information coming from the two eyes requires the integration of conflicting binocular signals. Under physiologic conditions, disparities between the images projected onto each retina tend to be modest, and the brain can effectively fuse the two images into a single percept. Processing of spatial image disparities, within a range, enables the perception of depth (stereopsis) (Parker 2007). How the brain processes conflicting visual signals has been studied since the foundational work of Hubel and Wiesel (Hubel and Wiesel 1962). In mammals, signals from the two eyes largely first converge in primary visual cortex (V1) where monocularly responsive neurons with retinotopically aligned receptive fields synapse onto binocular neurons. Many binocular neurons are modulated by modest spatial differences between the inputs from each eye, a phenomenon known as disparity-tuning. Disparity-tuned neurons are thought to serve stereopsis (Hubel and Wiesel 1970, Hubel and Livingstone 1987, Cumming and DeAngelis 2001). In mice, even relatively large spatial disparities can be integrated by neurons in V1 (Scholl, Burge et al. 2013, Scholl, Pattadkal et al. 2017, La Chioma, Bonhoeffer et al. 2020), corresponding to the short viewing distance at which mice tend to interact with objects (Samonds, Choi et al. 2019).

Although disparity-tuned neurons in V1 have been extensively studied under conditions of modest spatial offset, less is known about how cortical circuits respond when binocular input is strongly conflicting. In strabismus, a clinical condition defined by eye misalignment, large interocular disparities exceed the capacity for binocular integration and preclude stereopsis (Harrad, Sengpiel et al. 1996). When disparities between the images viewed by each eye are too great, fusion breaks down into diplopia (double vision) and binocular rivalry (an alternating, bistable percept). These outcomes can arise from a range of interocular differences in visual input, suggesting that different types of conflict may engage distinct cortical mechanisms. However, how early visual circuits respond to qualitatively different forms of interocular conflict remains poorly understood.

We sought to study how mammalian V1 processes large interocular image disparities using the tractability of the mouse visual system to identify the populations of neurons involved. Our goal was to identify how distinct forms of interocular conflict are represented at the level of local circuits in primary visual cortex, providing a critical step toward understanding the neural basis of perceptual suppression. Using a combination of visually evoked potentials (VEPs), unit recordings, and 2-photon (2P) calcium imaging, we characterized cortical responses to two types of interocular stimulus disparities in mice. The first—orientation-matched grating stimuli with an interocular phase difference—mimics the magnitude of spatial disparity that tends to give rise to diplopia in humans. The second—orthogonally oriented grating stimuli—employs stimulus combinations often used to study binocular rivalry in humans (Blake and Logothetis 2002) and non-human primates (Leopold and Logothetis 1996). Although both types of disparity led to reductions in VEP magnitude, they did so via markedly different mechanisms: phase offset stimuli suppressed the early negative component of the VEP through decreased

early evoked firing, while orthogonal stimuli suppressed the late positive component through
prolonged spiking activity. These stimulus-specific patterns were evident across cortical layers
and in the responses of both regular-spiking (RS, putative excitatory) and fast-spiking (FS,
putative parvalbumin-positive inhibitory) neurons. Furthermore, 2P calcium imaging revealed
that a separate class of interneurons expressing somatostatin (SOM+) exhibited decreased
activity in response to orthogonal stimuli. These findings demonstrate that V1 circuits in mice
exhibit distinct patterns of activity in response to different forms of interocular conflict. This
divergence is reflected in anatomically and temporally specific activity patterns across
excitatory and inhibitory cell types, providing mechanistic insight into how early visual cortex
processes discordant binocular input.
123
RESULTS
Interocular phase or orientation disparities differentially reduce VEP magnitude
We first sought to determine the effect of presenting disparate stimuli to each eye on the
cortical responses of mice. We implanted mice with tungsten microelectrodes in layer 4 (L4) of
bV1 (**Figure 1A**) and measured VEPs in awake, head-fixed mice using phase-reversing sinusoidal
grating stimuli presented on a passive 3D display (**Figure 1B**). Reciprocally polarized lenses
placed in front of each eye enabled the independent presentation of binocularly discordant
stimuli. For all recordings, the eye contralateral to the recording electrode viewed full contrast
stimuli at a 45° orientation. The ipsilateral eye viewed stimuli ranging from 0% contrast (i.e.
grey screen) to 50% contrast. Ipsilateral eye stimuli were either in phase (concordant) or 180°
out of phase (phase offset) relative to the contralateral eye stimulus (**Figure 1C**). We found that
the VEP magnitudes (peak negative to peak positive voltage) elicited by concordant stimuli
were larger than those elicited by monocular (0% ipsilateral eye contrast) or phase offset
stimuli (**Figure 1D**). Notably, a statistically significant difference between concordant and phase
offset stimuli emerged at an ipsilateral eye contrast as low as 20%.
Next, we looked at how orientation differences between the two eyes affects bV1
responses. To test this, we used the same dichoptic display to present stimuli that were either
at the same orientation for each eye (concordant) or rotated 90° (orthogonal). As before, the
contralateral eye viewed a full contrast stimulus while the ipsilateral eye viewed stimuli ranging
from 0% to 50% contrast (**Figure 1E**). Again, a substantial contrast-dependent difference in VEP
magnitude between concordant and orthogonal ipsilateral eye stimuli emerged, with a
statistically significant difference between concordant and orthogonal stimuli at just 10%
ipsilateral eye contrast (**Figure 1F**). Relative to monocular stimuli, we also observed a robust
increase in VEP magnitude for concordant stimuli, and decrease in VEP magnitude for
orthogonal stimuli. These findings suggest that even low contrast engagement of the inherently
weaker ipsilateral eye is sufficient to modulate cortical responses in the context of interocular
stimulus disparities, particularly in the case of rivalry.

To better understand which features of the VEP were modulated by these disparities,
we separately analyzed the negative and positive components of the waveform. For phase
offset stimuli, the reduction in VEP magnitude was primarily due to a significant attenuation of
the negative peak (**Figure 2A**), while changes in the positive peak were smaller and more
variable (**Figure 2B**). In contrast, orthogonal stimuli led to a substantial reduction in the positive
component of the VEP, with more modest effects on the negative component (**Figure 2C-D**).
These results suggest that distinct features of the VEP are differentially modulated by
interocular phase and orientation disparities.
Because we saw more robust reductions in the VEP for orthogonal stimulus orientations
than for phase offset disparities, we next sought to further characterize the features of VEP
responses elicited by this stimulus combination. First, we examined how sensitive this effect
was to graded orientation disparities between the eyes. Fixing the contralateral eye grating
stimulus at 100% contrast at a 45° orientation and the ipsilateral eye contrast at 50%, we varied
the grating orientation of the ipsilateral eye stimulus in 5° increments (from 0° to 30°, plus a 90°
condition relative to the contralateral eye) (**Figure 3A**). Interocular orientation differences of as
little at 10° showed a statistically significant decrease relative to the 0° (concordant) condition
(**Figure 3B**), indicating that the VEP is highly sensitive to interocular orientation disparity.
Next, we examined the orientation disparity effect at higher spatial frequencies. We
again presented orthogonal stimulus orientations to the eyes, with the contralateral eye fixed
at 100% contrast, and the ipsilateral eye ranging from 0% to 50% contrast. Stimuli were
presented at either 0.2 or 0.4 cycles per degree (cpd) (**Figure 3C**). At 0.2 cpd we again observed
a strong reduction in VEP magnitude relative to the 0% contrast baseline, but no change was
observed for 0.4 cpd (**Figure 3D**). This suggests that rivalrous stimulus effects on the VEP are
spatial frequency-dependent.
Finally, we asked whether the reduction in VEP magnitude observed for orthogonal
binocular stimuli could be accounted for by interocular interactions alone, or whether similar
effects might also be elicited by overlapping stimuli within a single eye. To test this, we
presented a full contrast grating to the contralateral eye, and added an overlapping orthogonal
grating to either the same eye (monocular condition) or to the opposite eye (binocular
condition), with contrast ranging from 0% to 50% (**Figure 3E**). In both cases, the superimposed
gratings created a plaid-like pattern, differing only in whether the component gratings were
presented to one or both eyes. While VEP magnitude decreased with increasing contrast in
both conditions, the reduction was more sensitive to contrast in the binocular condition (**Figure
3F**). These results suggest that interocular presentation contributes additional suppression
beyond that observed for monocular cross-oriented stimuli, consistent with a role for binocular
interactions in shaping V1 responses to discordant inputs.
187
Changes in the VEP are driven by an early decrease in V1 firing with phase disparity, and
prolongation of firing with orientation disparity

While VEPs are a useful method for measuring summed neuronal responses within V1, one
cannot infer what circuitry underpinnings may explain differential, stimulus-dependent VEP
waveform changes. To better answer this question, we used 64-channel laminar probes to
measure neuronal activity across all layers of V1 (**Figure 4A**). Mice viewed dichoptic phase
reversing grating stimuli as described previously (see **Figure 1B**). The eye contralateral to the
recording electrode viewed full contrast stimuli at a fixed orientation, while the ipsilateral eye
viewed grey screen (monocular condition), an in-phase grating stimulus at the same angle
(concordant condition), an out-of-phase grating stimulus at the same angle (phase offset
condition), or an orthogonal grating (orthogonal condition, **Figure 4B**). In order to test all
conditions within each animal, ipsilateral eye stimuli were presented only at 50% contrast for
each dichoptic condition.
Recording VEPs across all channels enabled us to calculate a current source density plot
for each mouse. The electrode contact corresponding to L4 was identified for each mouse
based on the earliest current sink in the current source density (CSD) plots (**Figure 4C**, see
Methods). Consistent with our results using unipolar Tungsten electrodes (**Figure 1**), we
observed a reduction in L4 VEP magnitude for the phase offset and orthogonal conditions
relative to concordant stimuli (**Figure 4D-E**). We also found a statistically significant decrease
relative to monocular stimuli, which was previously only observed for orthogonal stimuli. Again,
the reduction in VEP magnitude in the phase offset and discordant conditions were caused by
distinct changes in VEP waveform; phase offset stimuli primarily decreased the VEP negativity,
whereas discordant stimuli primarily decreased the VEP positivity (**Figure 4F-G**).
To determine how changes in different components of the VEP relate to changes in
neuronal spiking in bV1, we next analyzed unit activity across cortical layers. Recordings from
individual mice were aligned based on the channel that showed the earliest L4 current sink, and
the multiunit activity envelope (MUAe) was calculated for each electrode contact (see
Methods). Approximate layer boundaries were assigned based on the distance from the L4 sink
and the L5 peak in MUAe (Senzai, Fernandez-Ruiz et al. 2019) (**Figure 5A**). To account for the
variability in spiking between layers, we z-scored the MUAe for each channel based on its
average across the entire recording session. We observed a strong visually evoked response
across all layers of V1 when mice viewed the grating monocularly through the contralateral eye
(**Figure 5B**). This response peaked around 60-80 ms after stimulus presentation and MUAe
returned to near-baseline levels by ~100 ms. To compare how activity across layers was
affected by the different stimulus conditions, we then assessed MUAe when the ipsilateral eye
viewed a concordant, phase offset, or orthogonal grating stimulus (**Figure 5C**). MUAe changed
across most layers of V1 for all three conditions (**Figure 5D**). Relative to monocular stimuli,
concordant and orthogonal stimuli increased the response in the early period following the
phase reversal (roughly corresponding in time with the negative component of the VEP),
whereas phase offset stimuli elicited a reduced response in this period (**Figure 5E-F**).
Unexpectedly, the orthogonal condition prolonged firing well beyond the visually evoked

response of the other conditions (**Figure 5E,G**). This prolonged period of firing persisted
through 200 ms after the phase reversal, corresponding to the timing of the VEP positivity in
response monocular and concordant stimuli (**Fig 4**). Thus, the reduction in the positive
component of the VEP waveform by the orthogonal condition may reflect prolongation of
activity.
234
Both regular-spiking and fast-spiking units show prolonged activity with orthogonal stimuli
Based on our finding that dichoptic orthogonal grating stimuli elicit prolonged MUA
responses, we hypothesized that this persistent activity could result in a reduction in inhibitory
drive. Specifically, we reasoned that if inhibitory interneuronal activity was diminished during
this period, excitatory neurons could remain active for a longer period. To explore this
possibility, we first examined parvalbumin-positive (PV+) interneurons, the most abundant
inhibitory cell type in cortex and key mediators of fast, perisomatic inhibition onto pyramidal
neurons (Tremblay, Lee et al. 2016). Although PV+ interneurons often exhibit similar stimulus
tuning to pyramidal neurons (Niell and Stryker 2008), we hypothesized that this relationship
might be disrupted during binocular conflict. If disinhibition contributes to the prolonged firing
observed for orthogonal stimuli, it might be reflected in altered PV+ cell activity during this
window. To test this, we classified single units as fast-spiking (FS, putative PV+ interneurons) or
regular-spiking (RS, putative pyramidal) based on trough-to-peak latency (**Figure 6A**), and
compared their responses across conditions (**Figure 6B**). RS units exhibited a clear early peak in
firing (40-80 ms after stimulus onset) that was elevated for concordant and orthogonal stimuli,
and reduced for phase offset stimuli (**Figure 6C-D**). Similar to the MUA results, a prolonged
increase in RS activity during the 100-200 ms window was observed only for orthogonal stimuli
(**Figure 6C, E**). However, FS units showed a strikingly similar pattern of responses (**Figure 6F-H**),
suggesting that PV+ interneurons are not selectively disengaged during the late phase of
orthogonal stimulation. Given this parallel between putative excitatory and PV+ cell activity, we
next considered whether the structure of these responses varied by cortical layer, which could
offer further insight into how binocular conflict is processed across the bV1 circuit.
To explore this, we analyzed the laminar distribution of RS and FS unit activity across
conditions to determine whether stimulus-dependent differences in firing rates were uniformly
expressed throughout cortex or varied by layer. Single units were sorted by depth and assigned
to L2/3, L4, L5, and L6 based on their location relative to L4 (RS: **Figure S1**; FS: **Figure S2**). In the
early time window (40–80 ms after stimulus onset), RS firing rates differed by both stimulus
condition and cortical layer. In particular, responses to phase-offset stimuli were reduced in
L2/3, L5, and L6 compared with other conditions, while responses in L4 were relatively
unchanged (**Figure 7A**). This observation was confirmed by a significant condition-by-layer
interaction. FS units, in contrast, showed no such interaction during the early period, with
broadly similar responses across layers and conditions (**Figure 7B**). During the late time window
(100–200 ms), RS units showed elevated firing for orthogonal stimuli across all layers (**Figure**

7C), consistent with the MUA results (**Figure 5B**). FS units also responded more strongly to
orthogonal stimuli during this period, but this effect was not uniform across lamina. Instead, it
was most pronounced in L2/3, where responses to orthogonal stimuli were significantly greater
than those to other conditions (**Figure 7D**). Together, these findings show that suppression of
RS activity by phase-offset stimuli occurs selectively outside of L4, suggesting that this effect is
not inherited from thalamic input but instead emerges within intracortical circuitry. In contrast,
the enhancement of RS and FS activity by orthogonal stimuli occurs broadly across layers for RS
units but is selectively amplified in L2/3 for FS units, pointing to a potential layer- and cell-type–
specific locus of circuit engagement during binocular orientation conflict.
277
bV1 principal cells and somatostatin-positive interneurons show opposite responses to
orthogonal dichoptic stimuli
In the context of established cortical circuitry (Tremblay, Lee et al. 2016), the similarity
in responses between RS and FS units to the different dichoptic stimulus conditions suggests
that modulation of PV+ interneurons is unlikely to mediate the persistent activity triggered by
the orthogonal condition. The next most likely candidate was somatostatin-positive (SOM+)
interneurons, which inhibit both excitatory cells and PV+ interneurons in the mammalian cortex
via feedback inhibition. We hypothesized that a decrease in SOM+ cell activity elicited by the
orthogonal condition disinhibits both PV+ and principal cells (PCs) in bV1. To compare the
responses of genetically defined PCs and SOM+ interneurons we used 2P calcium imaging. Mice
expressing GCaMP in promoter-specific neurons were head-fixed in front of a dichoptic display
viewing the same stimuli as described above (**Figure 8A,B**). Visual responses of individual
neurons were measured using a 2P microscope (**Figure 8C**). First, we genetically expressed
GCaMP6f in excitatory neurons using an EMX1-cre x GCaMP6f.tTA mouse line and measured
the responses at a depth corresponding to L2/3 (**Figure 8D**). Due to the relatively slow temporal
dynamics of GCaMP, we expected that the measured change in fluorescence (ΔF/F) response
for each condition would reflect the overall differences in firing rate that we observed in the
MUA, favoring a difference for the orthogonal condition. This is indeed what we observed.
Although there was a substantial amount of variation in response to the grating stimulus
orientation presented (**Figure 8E**), on average there was a robust increase in responses when
the orthogonally oriented grating was presented to the ipsilateral eye and with each phase
reversal (**Figure 8F-H**).
Next, we tested whether SOM+ cells might be differentially engaged by orthogonal
dichoptic stimuli. We expressed GCaMP7f in SOM+ interneurons in bV1 by injecting an AAV
expressing cre-dependent GCaMP7f into mice on a SOM-cre genetic line and imaged SOM+ cell
ΔF/F in L2/3 (**Figure 8I**). This time, we only presented mice with the monocular, concordant,
and orthogonal conditions and observed a robust and uniform increase in responses for
concordant stimuli relative to monocular stimuli and a decrease in responses for orthogonal
stimuli that grew with each phase reversal (**Figure 8J-M**). These observations draw a clear

contrast between the responses of genetically defined neuronal subtypes to reveal insights into

the circuitry underling bV1 responses to rivalrous dichoptic stimuli.

DISCUSSION

Cortical responses to concordant and phase offset stimuli can be accounted for via feedforward

mechanisms.

313   In this study, we used a combination of VEPs (**Figures 1-4**), MUA (**Figure 5**), SUA (**Figures**

6-7), and 2P calcium imaging (**Figure 8**) to investigate how population cortical responses of

mouse bV1 are modulated by different combinations of dichoptic visual stimuli. We found that

relative to monocular stimuli, concordant binocular stimuli and 180° phase offset stimuli elicit

opposing changes in the visually evoked response. Concordant stimuli enhanced the early

negative component of the VEP via a short latency increase in MUA, whereas phase offset

stimuli reduced the VEP negativity through a corresponding decrease in MUA. This elevated

response to concordant stimuli likely reflects increased drive of binocular neurons due to the

addition of inputs from the ipsilateral eye or the recruitment of the few ipsilateral-referring

neurons within bV1 (**Figure 9**). The relatively modest increase in VEP and MUA responses for

concordant relative to monocular stimuli likely reflects sublinear binocular integration in mouse

V1, consistent with prior work demonstrating that binocular inputs do not sum linearly but are

shaped by normalization mechanisms that preserve tuning and constrain response amplitude

(Longordo, To et al. 2013, Zhao, Liu et al. 2013).

327   The reduced response for phase offset stimuli can also be accounted for by known

feedforward mechanisms. Binocular simple cells in mouse V1 receive spatially aligned inputs

from both eyes that are typically tuned to the same orientation but differ in phase sensitivity.

When gratings of identical orientation but opposite phase are presented dichoptically, these

inputs can cancel due to destructive interference at the level of subthreshold membrane

potential, as observed in previous whole-cell recordings (Zhao, Liu et al. 2013). This interaction

reduces the net depolarization and consequently suppresses spiking output. Moreover, the

amplitude-dependent sublinear integration described by Longordo et al. (2013) predicts that

strong but conflicting monocular inputs, such as those generated by out-of-phase gratings, will

not sum constructively and may even inhibit one another, further diminishing the cortical

response. Together, these feedforward properties of binocular integration constrain responses

to phase-disparate stimuli and likely contribute to the observed suppression in MUA and VEP

amplitude.

340   This suppressive effect can also be interpreted in the context of spatial disparity. A 180°

phase offset in grating stimuli at 0.05 cpd, as used in our experiments, results in a spatial

disparity of 10° of visual field. The suppression of early visually evoked responses under this

condition suggests a feed-forward cortical mechanism, consistent with the sublinear integration

and cancellation described above. Importantly, this pattern may also reflect the disparity tuning

properties of individual neurons in mouse bV1. Prior studies have shown that mice exhibit

broader disparity tuning than other mammals, such as cats and primates, likely reflecting
adaptations for near-field visual processing in species that interact with objects at close range
(Scholl, Burge et al. 2013). Indeed, individual bV1 neurons in mice can show preference for
disparities of 10° or more (Samonds, Choi et al. 2019), making it plausible that the 180° phase
offset grating stimulus drives a small subpopulation of neurons but V1 on average is activated
less effectively as compared with zero-disparity (concordant) stimuli.
These observations are consistent with the disparity energy model first described by
Ohzawa, DeAngelis, and Freeman (Ohzawa, DeAngelis et al. 1990), in which monocular
receptive fields with a slight spatial offset converge to form binocular receptive fields tuned to
specific spatial or phase disparities. In this framework, responses are enhanced when the
binocular input matches a neuron's preferred disparity and attenuated when the disparity
deviates from that preferred offset, due to partial cancellation of left and right eye signals via
linear summation. Disparity tuning of neurons in mouse V1 has been demonstrated using both
single-unit recordings and calcium imaging, along with a corresponding pattern of facilitation
and attenuation for preferred and non-preferred disparities, respectively (Scholl, Burge et al.
2013, Scholl, Pattadkal et al. 2017, La Chioma, Bonhoeffer et al. 2020). Our results showing an
enhanced population response for concordant binocular stimuli relative to monocular stimuli,
and a suppressed response for 180° phase offset stimuli, are consistent with a greater
proportion of neurons tuned near 0° disparity and fewer tuned near 180°, in line with previous
findings that most mouse V1 neurons prefer in-phase stimuli (Fu, Tanabe et al. 2023). While
previous studies have characterized sublinear summation at the level of individual neurons
(Longordo, To et al. 2013, Zhao, Liu et al. 2013), our findings extend these principles to
population-scale measures and demonstrate how disparity-dependent facilitation and
suppression shape early cortical responses to dichoptic stimuli.
It is notable that the pattern of suppression of cortical responses for phase offset stimuli
is only statistically significant outside of L4 of bV1 for RS units (**Figure 7A**). Mice differ from
other mammals in that binocular inputs converge within thalamorecipient L4 in bV1 (Gordon
and Stryker 1996) . Our findings therefore suggest that disparity tuning may emerge after this
early stage of binocular processing. However, because we are measuring population-level
responses for each cortical layer we cannot rule out the possibility that a subset of neurons
within dLGN or L4 might display disparity tuning that drives the responses at subsequent stages
of processing within bV1. Further work will be required to fully understand how the cortical
circuitry of the early visual pathway gives rise to disparity tuning to support the processing of
stimuli with interocular phase differences.
380
Disinhibition of V1 responses to orthogonal binocular stimuli is mediated by SOM+ interneurons.
Unlike the responses to concordant and phase offset stimuli, the pattern of activity we
observe for binocular stimuli with conflicting stimulus orientations cannot be accounted for by
a simple feedforward model. In particular, we see a period of heightened MUA and SUA after

the initial visually evoked response (100-200 ms after stimulus presentation, **Figure 6**) which
corresponds to a flattening of the VEP positivity (**Figures 2, 4**). This increase is apparent across
cortical layers, and in the responses of both RS (mostly excitatory) and FS (mostly PV+
inhibitory) units within bV1 (**Figure 7B, C**).
Prior studies have shown that binocular integration in mouse V1 is sublinear when the
two eyes receive matched inputs, particularly at the preferred orientation, and that this
mechanism helps preserve orientation selectivity (Longordo, To et al. 2013, Zhao, Liu et al.
2013). While those studies did not address interocular orientation conflict, they suggest that
binocular responses should remain modest when summed across eyes. Our findings are
consistent with this prediction during the early response phase, but diverge in the later time
window, where we observe a robust and selective enhancement of activity for orthogonal
stimuli. This pattern suggests a shift from sublinear integration to active disinhibition.
SOM+ interneurons, which inhibit both PCs and PV+ interneurons within cortex and are
a powerful source of feedback inhibition (Tremblay, Lee et al. 2016, Del Rosario, Coletta et al.
2024), are well-situated to mediate this disinhibition. Our 2P calcium imaging results showed
that SOM+ cell activity increases for concordant binocular stimuli and decreases for stimuli with
orthogonal orientations (**Figure 8**). Based on this, we propose a model (**Figure 9**) in which SOM+
cells play a key role in shaping cortical responses to binocular stimuli with concordant or
orthogonal orientations. In this model, monocular inputs to V1 drive both PCs and PV+
interneurons; the PCs then activate SOM+ interneurons, which in turn provide feedback
inhibition to PCs and PV+ cells. This feedback inhibition keeps the visually evoked response
constrained in time and would generate the similar pattern of responses we observe for RS and
FS units. Presenting a concordant stimulus to the other eye would increase the drive onto PCs
(and PV+ cells), and therefore elicit stronger SOM+ cell activity (**Figure 8J-M**) and feedback
inhibition. In the case of orthogonal stimulus presentation, however, there is a reduction in the
activation of SOM+ cells, leading to disinhibition of PCs and PV+ cells after the initial peak of
activity. This disinhibition could arise from local connections between competing and mutually
inhibitory populations of SOM+ cells within V1, or via feedback from other cortical areas,
possibly mediated by vasoactive intestinal peptide-expressing interneurons. Further
experiments will be required to disentangle these possibilities.
In humans, presenting discordant stimuli to each eye results in a repeated switching of
perception of each stimulus, a phenomenon known as binocular rivalry (Levelt 1965).
Recordings of V1 during binocular rivalry experiments reveal modulation of activity
corresponding to the perceived stimulus (Polonsky, Blake et al. 2000, Tong and Engel 2001). In
awake monkeys presented with orthogonal grating stimuli to each eye and trained to report the
perceived orientation, many neurons within V1 with a preference for one orientation respond
only when the monkey reports perceiving that orientation, and are suppressed when the
opposing percept is dominant (Leopold and Logothetis 1996). Presenting rivalrous stimuli to
anesthetized monkeys elicits an alternating activation and suppression of ocular dominance

columns corresponding to each eye that resembles the dynamics and pattern of switching that
occurs in binocular rivalry, suggesting that rivalry within V1 is unlikely to solely rely on top-
down mechanisms (Xu, Han et al. 2016). In contrast, V1 neurons recorded from strabismic
monkeys did not show any net facilitation or suppression (Economides, Adams et al. 2021),
suggesting that interocular suppression in strabismus is likely to be mediated at a higher level
of processing.

Whether binocular rivalry occurs in mice, either at the level of perception or cortical
dynamics, remains an open question. Our findings show that discordant binocular inputs elicit
divergent responses in PV+ and SOM+ interneurons, with increased PV+ cell firing and
suppressed SOM+ cell activity relative to concordant stimuli. These cell-type-specific dynamics
suggest that mouse V1 circuits respond selectively to interocular conflict and may contribute to
early-stage processing of binocular mismatch. While not a direct readout of perceptual
suppression, these responses highlight fundamental principles of how cortical circuits handle
interocular conflict. Given that binocular rivalry in humans is modulated by GABAergic signaling
(Mentch, Spiegel et al. 2019), these findings underscore the value of the mouse as an accessible
model system for dissecting the inhibitory circuit mechanisms that govern binocular integration
and competition.
441
Integration across modalities and implications for binocular processing
Although VEPs, unit recordings, and calcium imaging each capture different aspects of
cortical activity, our findings highlight how these modalities together can clarify the temporal
and cell-type-specific mechanisms underlying binocular processing. VEPs primarily capture the
net synaptic currents across large populations of neurons and are particularly sensitive to
changes in activity over time, rather than absolute firing rates (Hayden, Finnie et al. 2023). In
this context, a reduction in VEP amplitude can arise from different underlying patterns. For
example, a lower initial peak in firing can reduce both the early negative and subsequent
positive components of the VEP, while a delayed or sustained increase in firing may decrease
the relative change in activity over time, resulting in a flattened or diminished VEP positivity.
This framework helps reconcile several of our findings. For phase-offset stimuli, the
reduction in VEP negativity aligns with decreased early MUA and SUA, suggesting a lower peak
response. In contrast, orthogonal stimuli produce a reduced VEP positivity alongside an
increase in late-phase spiking, reflecting a prolonged or redistributed firing pattern that
diminishes the temporal contrast captured by the VEP. Calcium imaging of excitatory neurons
supports this interpretation. Because calcium signals reflect integrated activity over longer
timescales, the observed increase in fluorescence in response to orthogonal stimuli is
consistent with the elevated firing rates during both early and late response windows. This
result aligns with prior findings (Kim, Chaloner et al. 2020) that VEP magnitude does not simply
reflect overall excitatory activity as measured by calcium imaging but is instead shaped by the
timing and synchrony of neuronal responses.

These modality-specific differences underscore the importance of a multimodal

approach for dissecting the dynamics of binocular integration. Our results show that different

forms of interocular conflict engage distinct circuit mechanisms within mouse V1. Phase-offset

stimuli rely on feedforward integration consistent with disparity tuning, whereas orientation

conflicts recruit mechanisms involving specific cell types and cortical layers, consistent with

disinhibition. These findings reveal how local circuitry in V1 encodes distinct forms of binocular

conflict and provide a foundation for future studies of both normal binocular function and its

disruption in developmental conditions such as strabismus and amblyopia.

MATERIALS AND METHODS

Resources availability

Lead Contact. Further information and requests for resources and reagents should be directed

to and will be fulfilled by the Lead Contact, Eric D. Gaier (eric.gaier@childrens.harvard.edu). Key

Resources are outlined in **Supplemental Table 1**.

Materials Availability. This study did not generate new unique reagents.

Data and Code Availability. The code used to analyze visually evoked potentials are available on

Github (https://github.com/danielmontgomery7/suppression-analysis-code). The datasets

generated during this study have not been deposited in a public repository but are available

from the corresponding author on request.

Experimental Design and Subject Details

All procedures adhered to the guidelines of the National Institutes of Health and were

approved by the Committee on Animal Care at Massachusetts Institute of Technology (MIT).

For local field potential and laminar probe experiments, we used male and female mice on a

C57BL/6N background (Charles River Laboratories). For principal cell calcium imaging

experiments, EMX1.Cre mice (B6.129S2-Emx1tm1(cre)Krj/J, The Jackson Laboratory,

RRID:IMSR_JAX:005628) and Ai93(TITL-GCaMP6f)-D;CaMK2a-tTA mice

(Igs7tm93.1(tetO−GCaMP6f)Hze Tg(Camk2a- tTA)1Mmay/J, The Jackson Laboratory,

RRID:IMSR_JAX:024108) were crossbred in the animal facility at MIT to produce

EMX1.GCaMP6f.tTA triple transgenic mice. All EMX1.GCaMP6f.tTA mice were fed with a sterile

doxycycline food pellet diet (200 mg/kg, Bio-Serv, Flemington, NJ, USA) until weaned and

changed to a normal diet. For SOM+ cell calcium imaging experiments, SOM-Cre mice (B6N.Cg-

Ssttm2.1(cre)Zjh/J; catalog #018973, The Jackson Laboratory; RRID:IMSR_JAX:018973). The

effects reported in this study did not differ qualitatively by sex, so both were combined.

Animals were housed in groups of 2–5 same-sex littermates after weaning at postnatal day 21

(P21). They had access to food and water ad libitum and were maintained on a 12 h light-dark

cycle.

Method Details

Surgeries. For local field potential experiments, young adult C57BL/6 mice (P31-32) were

injected with 0.1 mg/kg buprenex HCl and subcutaneously (s.c.) to provide analgesia. Induction

of anesthesia was achieved via inhalation of isoflurane (3% in oxygen) and thereafter

maintained via inhalant isoflurane (1-2% in oxygen). Before surgical incision, the head was

shaved and the scalp cleaned with povidone–iodine (10% w/v) and ethanol (70% v/v). The scalp

was resected, and the skull surface was scored with a scalpel. A steel head post was affixed to

the skull (anterior to bregma) with cyanoacrylate glue. Small burr holes were drilled above both

hemispheres of binocular V1 (3.1mm lateral of lambda). Tapered 300–500 kΩ tungsten

recording electrodes (FHC), 75µm in diameter at their widest point, were implanted in each

hemisphere, 470mm below the cortical surface. Silver wire (A-M Systems) reference electrodes

were placed over the left frontal cortex. Electrodes were secured using cyanoacrylate, and the

skull was covered with dental cement. Nonsteroidal anti-inflammatory drugs were

administered on return to the home cage (meloxicam, 1.0 mg/kg s.c.). Buprenex was

administered on post-operative day 1, and meloxicam on post-operative days 1 and 2. Signs of

infection and discomfort were carefully monitored. Mice were allowed to recover for at least

48h before head fixation.

For headplate surgeries for acute electrophysiology experiments, adult mice (P60-P80)

were anesthetized and prepared as described above. Following scalp incision, a lidocaine (1%)

solution was applied onto the periosteum, and the exposed area of skull gently scraped with a

scalpel blade. A mark on the skull was made above bV1 (AP -3.6, ML +3.1) with marker. A

custom stainless-steel head plate was attached to the skull and dental acrylic (C&B Metabond

Quick Adhesive Cement System) was used to form a 5mm x 5mm well above bV1. The well was

filled with Kwik-Sil (World Precision Instruments) which was held in place using a thin bridge of

Ortho-Jet dental cement (Lang Dental). Mice were returned to their home cage and monitored

for at least 48h prior to head fixation. On the morning of the acute recording, mice were placed

again under isoflurane anesthesia. The Ortho-Jet bridge and Kwik-Sil cover were removed, and

a ~2mm craniotomy was made around the mark previously made above bV1. Kwik-Sil was once

again applied, and the mice were removed from anesthesia and single-house for 2 hours. The

mice were head fixed, the Kwik- Sil removed, and a 64-channel laminar probe (H3, Cambridge

NeuroTech) was inserted slowly (~100µm/min) into bV1 perpendicular to the cortical surface.

Recordings were obtained and the mice were euthanized immediately thereafter.

For cranial window implantations for two-photon calcium imaging, adult

EMX1.GCaMP6f.tTA and SOM-Cre mice (P60-P80) were anesthetized and prepared as described

above. A 3mm craniotomy was made over bV1. For SOM-Cre mice, an adeno-associated virus

containing the GCaMP7f gene (pGP-AAV9-syn-FLEX-jGCaMP7f-WPRE; catalog #104488- AAV9,

Addgene) was loaded into a glass micropipette with a tip diameter of 40–50µm attached to a

Nanoject II injection system (Drummond Scientific). The micropipette was then inserted into

bV1 (AP -3.6, ML +3.1) layer 2/3 at depths of 300 and 200mm below the pial surface, and 50nL

of virus was delivered at each depth. For EMX1.GCaMP6f.tTA mice with genetic expression of

GCaMP, an image of the brain was taken during the surgery with a pin at the coordinates for

bV1 (AP -3.6, ML +3.1). On subsequent imaging days, the imaging field of view was matched to

the vasculature at the coordinates indicated by the pin to ensure that imaging was centered on

bV1. Next, a sterile 3-mm-round glass coverslip (CS-3R-0; Warner Instruments) was gently laid

on top of the exposed dura mater. The coverslip was secured with cyanoacrylate glue, and a

custom stainless steel headplate was attached to the skull. Once the glue had set, dental acrylic

(C&B Metabond Quick Adhesive Cement System) was mixed and applied throughout the

exposed skull surface. Nonsteroidal anti-inflammatory drugs were administered upon return to

the home cage (meloxicam, 1 mg/kg s.c.). Signs of infection and discomfort were carefully

monitored. Mice were allowed to recover for at least 3 weeks prior to head-fixation to allow

time for the window to clear and for the virus to be fully expressed.

Visual stimulus delivery. Prior to stimulus delivery, mice were acclimated to head restraint in

front of a gray screen for a 30-minute session on each of two consecutive days. For

electrophysiological experiments, mice viewed stimuli on a passive 3D monitor (Vizeo

E3D420VX, 120 Hz refresh rate) with reciprocally polarized lenses placed in front of each eye.

For calcium imaging experiments, mice viewed stimuli on an active 3D monitor (Acer

UMFG6AAB01, 144 Hz refresh rate) with shutter lenses (modified from NVIDIA 3D Vision 2

glasses) placed in front of each eye. Visual stimuli were generated using custom software

written in either C++ for interaction with a VSG2/2 card (Cambridge Research Systems) or

MATLAB (MathWorks) using the Psychtoolbox extension (http://psychtoolbox.org) to control

stimulus drawing and timing and to present different stimuli to each eye. Grating stimuli

spanned the full range of monitor display values between black and white, with gamma

correction to ensure constant total luminance in both gray-screen and patterned stimulus

conditions.

During recordings, mice were head fixed 25 cm from the screen and viewed grating

stimuli with a Gabor filter to limit the size of the stimuli to a 40° field of view directly in front of

each mouse. Except where noted otherwise, phase-reversing grating stimuli were presented at

1 Hz, 0.05 cycles per degree, and an orientation of 45° or 135°. Stimuli were always presented
to the left eye (contralateral to electrode or window) at 100% contrast.
We observed ~10% interocular bleed-through on both 3D display systems, such that
each eye received approximately 10% of the stimulus presented to the other eye. For VEP and
2-photon calcium imaging experiments (**Figures 1-3, 8**), this bleed-through was not corrected.
However, the impact on our results was likely minimal given that (1) the ipsilateral eye stimulus
was relatively low contrast (10-50%), and (2) ipsilateral eye stimulation generally evokes much
weaker responses in mouse V1 compared to the contralateral eye. Thus, any contribution of
bleed-through was negligible in the context of our measurements. For acute laminar recordings
(**Figures 4-7**), we implemented a correction for this crosstalk by reducing the stimulus
presented to each eye by 10% of the contrast intended for the opposite eye. This reduced
interocular bleed-through to below detectable levels by eye but resulted in a slightly reduced
maximum effective contrast range (0-95% instead of 0-100%).
For VEP recordings, stimuli were presented with the contrast of the right (ipsilateral) eye
stimulus increasing from 0% to 50% or decreasing from 50% to 0% in increments of 10%
contrast between blocks. Each block lasted 75 seconds with a 15 second interblock interval for
4 blocks per condition. For acute laminar recordings and calcium imaging experiments the right
(ipsilateral) eye stimulus was presented at 50% contrast (for Concordant, Phase Offset, and
Orthogonal conditions) or 0% contrast (for the Monocular condition). Stimuli for acute laminar
recordings were presented in pseudorandomly interleaved 100 second blocks with 30 second
interblock intervals for 5 blocks per condition, and stimuli for calcium imaging experiments
were presented in pseudorandomly interleaved 10 second blocks with 10 second interblock
intervals for 24 blocks per condition.
601
Electrophysiology recordings and analysis. Electrophysiological recordings were conducted in
awake, head-restrained mice. Recordings were amplified and digitized using the Recorder-64
system (Plexon Inc.) or the RHD Recording system (Intan Technologies). Local field potentials
were recorded from V1 with 1-kHz sampling using a 500-Hz low-pass filter. For the laminar
recordings on the Intan system, we sampled at 25 kHz and used a 0.1-Hz high-pass and a 7.5-
kHz low-pass filter.
All analyses were conducted using custom MATLAB code and the Chronux toolbox
(Bokil, Andrews et al. 2010). For laminar recordings, the raw 25-kHz data from each channel
were extracted and converted to mV. Then they were down sampled to 1000Hz and a third
order 1- to 300-Hz Butterworth filter was applied. For all data, the mean of the entire channel's
data was subtracted from each time point to account for any DC offset in the system. Next, the
data were locally detrended using the locdetrend function in the Chronux toolbox using a 0.5 s
window sliding in chunks of 0.1 s. Finally, a third-order Butterworth filter was used to notch
frequencies between 58 and 62Hz. For the multiunit activity of laminar recordings, the raw 25-
kHz data were extracted for each channel. A 60Hz, 10-dB bandwidth IIR notch filter was applied

to each channel and the median value of each channel was subtracted from that channel. The
median value across channels for each time point was subtracted from all channel's timepoints.
Visually evoked potentials were normalized by subtracting the average of the first 10 ms of
each trial from that trial. A 10-ms moving Gaussian was applied to smooth the visually evoked
potential waveform.
622

Current-source density analysis and laminar identification. Current-source density (CSD) analysis
on laminar local field potential (LFP) data in V1 was used to identify layer 4 (L4) for each mouse
and align our recordings (Mitzdorf 1985, Mitzdorf 1987, Aizenman, Kirkwood et al. 1996). The
LFP data were temporally smoothed with a 20 ms Gaussian window, then a five-point hamming
window was used to compute the CSD (Ulbert, Halgren et al. 2001). We define 0 mm as the site
of the earliest and deepest sink immediately below the superficial source. All other channels
were referenced according to that landmark. Thus, superficial channels were above 0 mm and
deep channels were below 0 mm. For the sake of reporting in Results, we broke the laminar
data into four segments roughly corresponding to each cortical layer (Senzai, Fernandez-Ruiz et
al. 2019): L2/3 between +300 and +60 µm, L4 between +60 and -80 µm, L5 between -80 and -
260 µm, and L6 between -260 and -460 µm.
634

Multiunit activity analysis. To measure multiunit activity, we calculated the multiunit activity
envelope (MUAe), which provides an instantaneous measure of the number and size of action
potentials of neurons in the vicinity of the electrode tip without the requirement of setting an
arbitrary threshold level (Legatt, Arezzo et al. 1980, Brosch, Bauer et al. 1995, Super and
Roelfsema 2005). MUAe was calculated by applying a third order bandpass (500-5000Hz)
Butterworth filter to the common median referenced data to isolate spiking unit activity. Next,
the absolute values of the data were taken (units of mV) to remove contamination from far-
field signals. Finally, a third order low pass (<250Hz) Butterworth filter was applied to smooth
the signal and the data were down sampled to 1000Hz to align with local field potential data.
MUAe was z-scored based on the average and standard deviation across all interblock intervals.
645

Single unit activity analysis. Single unit activity was calculated for the common median
referenced data using Kilosort 2.5 (Pachitariu, Sridhar et al. 2024) (Kilosort parameters: final
projection threshold = 4; spike threshold = 6; channel count = 64; clusters per channel = 4; min
spike rate = 0.02; high pass filter frequency = 300; optimization projection threshold = 10;
number of blocks = 5; sigma mask = 30; batch size = 1024; whitening range = 32). Spikes
identified by Kilosort were manually curated using Phy to isolate single units. Single units with a
trough-to-peak latency of less than 0.4 ms were classified as fast-spiking, and units with a
latency of greater than 0.4 ms were classified as regular-spiking. The spike rate (in
spikes/second) for each unit was calculated by binning spikes into 1 ms time bins for each trial,
and then multiplying the number of spikes within each bin across all 500 trials for each

condition by 2. Average RS and FS unit PSTHs were calculated by averaging the spike rates of
each class of unit. Firing rates during early and late time bins were calculated by averaging firing
rates between 40-80 ms after stimulus onset, and 100-200 ms after stimulus onset,
respectively. Z-scored SUA was calculated based on the average and standard deviation of the
firing rate of each unit across all interblock intervals.
661
In vivo two-photon calcium imaging. Three to four weeks following craniotomy surgery, mice
were habituated to the behavior restraint apparatus in front of a gray screen with the objective
lens of the two-photon microscope positioned on the head plate for 30 min for two consecutive
days before beginning their visual stimulus delivery. A Ti:sapphire laser (Coherent) was used for
imaging at a wave length of 930 nm. Photomultiplier tubes (Hamamatsu) and the objective lens
(20×, 0.95 numerical aperture, XLUMPLFLN, Olympus) were used to detect fluorescence
images. Calcium image recordings were triggered by time-locked transistor-transistor logic
pulses generated from the USB-1208fs data acquisition device (Measurement Computing) using
PrairieView and TriggerSync software (Bruker) and imaged at a frequency of ~14.9 Hz at a depth
of ~200 mm in bV1. The size of the imaging field of view was ~600 × 600 mm^2 at 512 × 512
pixels.
673
Calcium imaging analysis. Acquired time series of calcium imaging files were processed using
Suite2p (Pachitariu, Stringer et al. 2017). All recorded files were registered to stabilize shifts due
to animal movement. Regions of interest (ROIs) were automatically detected, then manually
curated based on the maximum projection of all frames. Mice were excluded from all further
analysis if they had fewer than 100 ROIs within the field of view, as this indicated excessive
noise during the recording. We extracted the fluorescence of each ROI for all time points. In line
with previous work, for each ROI we calculated the estimated true fluorescence of the ROI. This
is the measured fluorescence of the ROI minus 7/10ths of the average measured fluorescence
of the surrounding neuropil (Chen, Wardill et al. 2013). We used the average interblock gray
period to compute the response relative to gray (F_{stim}–F_{avg_gray})/F_{avg_gray}. The average population
response for each mouse was calculated by taking the average dF/F of all ROIs across all blocks.
685
Quantification And Statistical Analysis
Statistics. Throughout the results section, all data is expressed as group mean ± standard error
of the mean (SEM). Each dataset was assessed for normality and homogeneity of variance prior
to choosing a statistical approach, using Levene's, D'Agostino and Pearson, and, in the case of
small sample sizes, Shapiro-Wilk tests. One-way repeated-measures (RM) analysis-of-variance
(ANOVA) were used to compare group responses across conditions. Two-way RM ANOVA were
used to compare group responses across conditions, stimulus contrasts, or cortical layers.
Greenhouse-Geisser was used to modify the degrees of freedom of repeated-measures tests to
correct for violations of sphericity. Interactions were followed by tests of simple main effects.

Tukey or Sidak's methods were applied to compensate for multiple comparisons. Uncorrected

alpha was set to 0.05. Statistical analyses were performed with Prism 10 (GraphPad;

RRID:SCR_002798).

REFERENCES

Aizenman, C. D., A. Kirkwood and M. F. Bear (1996). "A current source density analysis of
evoked responses in slices of adult rat visual cortex: implications for the regulation of long-term
potentiation." Cereb Cortex **6**(6): 751-758.

Blake, R. and N. K. Logothetis (2002). "Visual competition." Nature Reviews Neuroscience **3**(1):
13-21.

Bokil, H., P. Andrews, J. E. Kulkarni, S. Mehta and P. P. Mitra (2010). "Chronux: a platform for
analyzing neural signals." J Neurosci Methods **192**(1): 146-151.

Brosch, M., R. Bauer and R. Eckhorn (1995). "Synchronous high-frequency oscillations in cat
area 18." Eur J Neurosci **7**(1): 86-95.

Chen, T. W., T. J. Wardill, Y. Sun, S. R. Pulver, S. L. Renninger, A. Baohan, E. R. Schreiter, R. A.
Kerr, M. B. Orger, V. Jayaraman, L. L. Looger, K. Svoboda and D. S. Kim (2013). "Ultrasensitive
fluorescent proteins for imaging neuronal activity." Nature **499**(7458): 295-300.

Cumming, B. G. and G. C. DeAngelis (2001). "The physiology of stereopsis." Annual review of
neuroscience **24**(1): 203-238.

Del Rosario, J., S. Coletta, S. H. Kim, Z. Mobille, K. Peelman, B. Williams, A. J. Otsuki, A. D. C.
Valerio, K. Worden and L. T. Blanpain (2024). "Lateral inhibition in V1 controls neural &
perceptual contrast sensitivity." bioRxiv: 2023.2011. 2010.566605.

Economides, J. R., D. L. Adams and J. C. Horton (2021). "Interocular suppression in primary
visual cortex in strabismus." Journal of Neuroscience **41**(25): 5522-5533.

Fu, J., S. Tanabe and J. Cang (2023). "Widespread and Multifaceted Binocular Integration in the
Mouse Primary Visual Cortex." J Neurosci **43**(38): 6495-6507.

Gordon, J. A. and M. P. Stryker (1996). "Experience-dependent plasticity of binocular responses
in the primary visual cortex of the mouse." J Neurosci **16**(10): 3274-3286.

Harrad, R., F. Sengpiel and C. Blakemore (1996). "Physiology of suppression in strabismic
amblyopia." The British journal of ophthalmology **80**(4): 373.

Hayden, D. J., P. S. B. Finnie, A. Thomazeau, A. Y. Li, S. F. Cooke and M. F. Bear (2023).
"Electrophysiological signatures of visual recognition memory across all layers of mouse V1." J
Neurosci.

Hubel, D. H. and M. S. Livingstone (1987). "Segregation of form, color, and stereopsis in primate
area 18." Journal of neuroscience **7**(11): 3378-3415.

Hubel, D. H. and T. N. Wiesel (1962). "Receptive fields, binocular interaction and functional
architecture in the cat's visual cortex." J Physiol **160**: 106-154.

Hubel, D. H. and T. N. Wiesel (1970). "Stereoscopic vision in macaque monkey. Cells sensitive to
binocular depth in area 18 of the macaque monkey cortex." Nature **225**(5227): 41-42.

Kim, T., F. A. Chaloner, S. F. Cooke, M. T. Harnett and M. F. Bear (2020). "Opposing Somatic and
Dendritic Expression of Stimulus-Selective Response Plasticity in Mouse Primary Visual Cortex."
Front Cell Neurosci **13**: 555.

La Chioma, A., T. Bonhoeffer and M. Hübener (2020). "Disparity sensitivity and binocular
integration in mouse visual cortex areas." Journal of Neuroscience **40**(46): 8883-8899.

Legatt, A. D., J. Arezzo and H. G. Vaughan, Jr. (1980). "Averaged multiple unit activity as an
estimate of phasic changes in local neuronal activity: effects of volume-conducted potentials." J
Neurosci Methods **2**(2): 203-217.

Leopold, D. A. and N. K. Logothetis (1996). "Activity changes in early visual cortex reflect
monkeys' percepts during binocular rivalry." Nature **379**(6565): 549-553.

Levelt, W. J. (1965). "Binocular Brightness Averaging and Contour Information." Br J Psychol **56**:
1-13.

Longordo, F., M.-S. To, K. Ikeda and G. J. Stuart (2013). "Sublinear integration underlies
binocular processing in primary visual cortex." Nature neuroscience **16**(6): 714-723.

Longordo, F., M. S. To, K. Ikeda and G. J. Stuart (2013). "Sublinear integration underlies
binocular processing in primary visual cortex." Nat Neurosci **16**(6): 714-723.

Mentch, J., A. Spiegel, C. Ricciardi and C. E. Robertson (2019). "GABAergic Inhibition Gates
Perceptual Awareness During Binocular Rivalry." J Neurosci **39**(42): 8398-8407.

Mitzdorf, U. (1985). "Current source-density method and application in cat cerebral cortex:
investigation of evoked potentials and EEG phenomena." Physiol Rev **65**(1): 37-100.

Mitzdorf, U. (1987). "Properties of the evoked potential generators: current source-density
analysis of visually evoked potentials in the cat cortex." Int J Neurosci **33**(1-2): 33-59.

Niell, C. M. and M. P. Stryker (2008). "Highly selective receptive fields in mouse visual cortex." J
Neurosci **28**(30): 7520-7536.

Ohzawa, I., G. C. DeAngelis and R. D. Freeman (1990). "Stereoscopic depth discrimination in the
visual cortex: neurons ideally suited as disparity detectors." Science **249**(4972): 1037-1041.

Pachitariu, M., S. Sridhar, J. Pennington and C. Stringer (2024). "Spike sorting with Kilosort4."
Nature Methods: 1-8.

Pachitariu, M., C. Stringer, M. Dipoppa, S. Schröder, L. F. Rossi, H. Dalgleish, M. Carandini and K.
D. Harris (2017). "Suite2p: beyond 10,000 neurons with standard two-photon microscopy."
bioRxiv: 061507.

Parker, A. J. (2007). "Binocular depth perception and the cerebral cortex." Nature Reviews
Neuroscience **8**(5): 379-391.

Polonsky, A., R. Blake, J. Braun and D. J. Heeger (2000). "Neuronal activity in human primary
visual cortex correlates with perception during binocular rivalry." Nature neuroscience **3**(11):
1153-1159.

Samonds, J. M., V. Choi and N. J. Priebe (2019). "Mice Discriminate Stereoscopic Surfaces
Without Fixating in Depth." J Neurosci **39**(41): 8024-8037.

Scholl, B., J. Burge and N. J. Priebe (2013). "Binocular integration and disparity selectivity in
mouse primary visual cortex." J Neurophysiol **109**(12): 3013-3024.

Scholl, B., J. J. Pattadkal and N. J. Priebe (2017). "Binocular Disparity Selectivity Weakened after
Monocular Deprivation in Mouse V1." J Neurosci **37**(27): 6517-6526.

Senzai, Y., A. Fernandez-Ruiz and G. Buzsaki (2019). "Layer-Specific Physiological Features and
Interlaminar Interactions in the Primary Visual Cortex of the Mouse." Neuron **101**(3): 500-513
e505.

Super, H. and P. R. Roelfsema (2005). "Chronic multiunit recordings in behaving animals:
advantages and limitations." Prog Brain Res **147**: 263-282.

Tong, F. and S. A. Engel (2001). "Interocular rivalry revealed in the human cortical blind-spot
representation." Nature **411**(6834): 195-199.

Tremblay, R., S. Lee and B. Rudy (2016). "GABAergic Interneurons in the Neocortex: From
Cellular Properties to Circuits." Neuron **91**(2): 260-292.

Ulbert, I., E. Halgren, G. Heit and G. Karmos (2001). "Multiple microelectrode-recording system
for human intracortical applications." J Neurosci Methods **106**(1): 69-79.

Xu, H., C. Han, M. Chen, P. Li, S. Zhu, Y. Fang, J. Hu, H. Ma and H. D. Lu (2016). "Rivalry-like
neural activity in primary visual cortex in anesthetized monkeys." Journal of neuroscience
36(11): 3231-3242.

Zhao, X., M. Liu and J. Cang (2013). "Sublinear binocular integration preserves orientation
selectivity in mouse visual cortex." Nat Commun **4**: 2088.

Zhao, X., M. Liu and J. Cang (2013). "Sublinear binocular integration preserves orientation
selectivity in mouse visual cortex." Nature communications **4**(1): 2088.

FIGURE LEGENDS:

Figure 1. Binocular stimuli with an interocular phase or orientation disparity elicit smaller

visually evoked potentials. (A) Schematic of mouse brain showing placement of a chronic

recording electrode in binocular V1 (V1b). **(B)** Mice viewed phase reversing sinusoidal grating

stimuli on a dichoptic display. **(C)** Phase offset experimental design. The contralateral (Contra)

eye viewed full-contrast stimuli, while the ipsilateral (Ipsi) eye viewed stimuli ranging from 0%

to 50% contrast, presented either in phase (Concordant) or out of phase (Phase Offset) with the

contra eye stimulus. **(D)** For ispi eye contrasts ≥20% contrast, concordant stimuli elicited

significantly larger VEPs than phase offset stimuli. Average VEP waveforms for each condition

shown above bars. Two-way repeated measures ANOVA (main effect of condition, $F_{(1,18)}$ =

16.4, $p<0.001$; condition × contrast, $F_{(5,90)}$ = 2.37, p = 0.045) followed by Šídák's multiple

comparisons test (concordant vs. phase offset at 0%, 10%, 20%, 30%, 40%, and 50% ipsi eye

contrast: p = 0.522, 0.126, <0.001, <0.001, <0.001, <0.001; Concordant 0% vs. 10%, 20%, 30%,

40%, and 50% contrast: p = 0.999, 0.088, 0.999, 0.110, <0.001; Phase Offset 0% vs. 10%, 20%,

30%, 40%, and 50% contrast: p = 0.955, 0.999, 0.678, 0.999, 0.999). **(E)** Orthogonal orientation

experimental design. Similar to **C**, but the ipsi eye viewed stimuli at either the same orientation

as the contra eye (Concordant) or rotated 90° (Orthogonal). **(F)** For ispi eye contrasts ≥10%

contrast, Concordant stimuli elicited larger VEPs than Orthogonal stimuli. Two-way repeated

measures ANOVA (main effect of condition, $F_{(1,23)}$ = 32.6, $p<0.001$; condition × contrast,

$F_{(5,115)}$ = 13.8, $p < 0.001$) followed by Šídák's multiple comparisons test (Concordant vs.

Orthogonal at 0%, 10%, 20%, 30%, 40%, and 50% ipsi eye contrast: p = 0.437, 0.037, <0.001,

<0.001, <0.001, <0.001; Concordant 0% vs. 10%, 20%, 30%, 40%, and 50% contrast: p = 0.968,

0.282, 0.017, 0.003, <0.001; Orthogonal 0% vs. 10%, 20%, 30%, 40%, and 50% contrast: p =

0.137, 0.003, <0.001, <0.001, <0.001). Scale bars: 200 ms and 100 μV.

Figure 2. Phase offset and orthogonal stimuli affect different components of the VEP

(A) VEP negativities for Concordant (grey) and Phase Offset (purple) stimuli at different

ipsilateral eye contrasts. Analyses performed using a two-way repeated measures ANOVA (main

effect of condition, $F_{(1,18)}$ = 14.0, p = 0.002; condition × contrast, $F_{(5,90)}$ = 2.88, p = 0.019)

followed by Šídák's multiple comparisons test (concordant vs. phase offset at 0%, 10%, 20%,

30%, 40%, and 50% ipsi eye contrast: p = 0.393, 0.800, 0.009, <0.001, 0.001, <0.001). **(B)** VEP

positivities for Concordant and Phase Offset stimuli at different ipsilateral eye contrasts. Two-

way repeated measures ANOVA (main effect of condition, $F_{(1,18)}$ = 9.42, p = 0.007; condition ×

contrast, $F_{(5,90)}$ = 1.03, p = 0.407) followed by Šídák's multiple comparisons test (concordant

vs. phase offset at 0%, 10%, 20%, 30%, 40%, and 50% ipsi eye contrast: p = >0.999, 0.029, 0.015,

0.304, 0.349, 0.176). **(C)** VEP negativities for Concordant (grey) and Orthogonal (red) stimuli.

Two-way repeated measures ANOVA (main effect of condition, $F_{(1,23)}$ = 11.8, p = 0.002;

condition × contrast, $F_{(5,115)}$ = 6.09, $p < 0.001$) followed by Šídák's multiple comparisons test

(concordant vs. discordant at 0%, 10%, 20%, 30%, 40%, and 50% ipsi eye contrast: p = 0.847,
0.586, 0.210, 0.002, 0.002, <0.001). **(D)** VEP positivities for Concordant and Orthogonal stimuli.
Two-way repeated measures ANOVA (main effect of condition, F(1,23) = 48.73, p < 0.001;
condition × contrast, F(5,115) = 15.4, p < 0.001) followed by Šídák's multiple comparisons test
(concordant vs. discordant at 0%, 10%, 20%, 30%, 40%, and 50% ipsi eye contrast: p = 0.437,
0.009, <0.001, <0.001, <0.001, <0.001).

Figure 3. Reductions in VEP magnitude for discordant stimuli are sensitive interocular
orientation differences, spatial frequency, and binocularity.
(A) Mice viewed dichoptic grating stimuli (100% contra and 50% ipsi eye contrast) with varying
interocular orientation differences (0° to 30° in 5° increments, plus an orthogonal orientation).
(B) VEP magnitudes were normalized to the baseline (0°) condition. Orientation differences of
10° or larger elicited significantly smaller VEPs. One-way repeated measures ANOVA with
Geisser-Greenhouse correction (F(3.32, 29.8) = 12.7, p < 0.001) followed by Dunnett's multiple
comparisons test (0° vs. 5°, 10°, 15°, 20°, 25°, 30°, and 90°: p = 0.086, 0.025, 0.001, 0.005,
<0.001, <0.001, 0.003). **(C)** Mice viewed discordant stimuli at higher spatial frequencies (0.2 and
0.4 cpd) with orthogonal ipsi eye stimuli (100% contra eye contrast, 0% to 50% ipsi eye
contrast). **(D)** VEP magnitude was reduced for 0.2 cpd, but not for 0.4 cpd. Two-way repeated
measures ANOVA with Geisser-Greenhouse correction (main effect of SF, F(1,20) = 22.66, p <
0.001; SF × contrast, F(2.92,58.4) = 4.69, p = 0.006) followed by Dunnett's multiple comparisons
test (0% vs. 10%, 20%, 30%, 40%, and 50% ipsi eye contrast for 0.02 cpd: p = 0.003, <0.001,
<0.001, <0.001, <0.001; for 0.04 cpd: p = 0.701, >0.999, >0.999, >0.999, >0.999). **(E)** Mice
viewed dichoptic grating stimuli (100% contra eye contrast) with a conflicting orthogonal
grating. The orthogonal grating (0% to 50% contrast) was either presented to the ipsi eye
(Binocular) or overlayed on top of the initial stimulus orientation for the contra eye
(Monocular). **(F)** VEP magnitude for Binocular and Monocular conditions decreased as
orthogonal grating contrast increased, but this decrease was sensitive to lower stimulus
contrast in the Binocular condition. Two-way repeated measures ANOVA with Geisser-
Greenhouse correction (main effect of condition, F(1,23) = 4.38, p = 0.048; condition × contrast,
F(3.48, 80.0) = 1.44, p = 0.235) followed by Dunnett's multiple comparisons test (0% vs. 10%,
20%, 30%, 40%, and 50% contrast for Monocular: p = 0.817, 0.011, <0.001, <0.001, <0.001; for
Binocular: p = 0.006, <0.001, <0.001, <0.001, <0.001). **B,D,F:** Error bars indicate SEM. Scale bars:
200 ms and 100 μV.

Figure 4. L4 VEPs measured using laminar probes reveal reductions in different components
of the VEP for phase offset and discordant stimuli.
(A) Recording setup. LFPs were recorded from bV1 using a 64-channel silicon probe. The probe
spanned all cortical layers with channel spacing of 20 μm. **(B-C)** Monocular VEPs recorded from
each electrode channel (**B**) were used to calculate current source densities (CSDs) shown in (**C**).

The earliest current sink in the CSD for each mouse was used to identify the electrode
corresponding to layer 4 (shown in red in **B**). Approximate layer boundaries shown as white
dashed lines in **C**. **(D)** Stimulus conditions. Mice viewed dichoptic phase reversing gratings (2 Hz,
0.05 cpd) as described previously. Contra eye stimuli (45°, 100% contrast) were kept constant
across conditions. Ipsi eye stimuli varied across 4 conditions: Monocular (0% contrast),
Concordant (50% contrast, 45°), Phase Offset (50% contrast, 45° with 180° phase offset), and
Orthogonal (50% contrast, 135°). L4 VEPs for each condition shown below schematic. **(E)** VEP
magnitude for the Phase Offset (ΔPhase) and Orthogonal (ΔAngle) conditions were decreased
relative to the Monocular (Monoc) and Concordant (Conc) conditions. One-way repeated
measures ANOVA with Geisser-Greenhouse correction ($F(2.20, 22.0) = 20.0$, $p < 0.001$) followed
by Tukey's multiple comparison's test. **(F)** Relative to Monocular, the negative component of
the L4 VEP was larger (more negative) for Concordant, smaller for Phase Offset, and no
different for Orthogonal stimuli. One-way repeated measures ANOVA with Geisser-Greenhouse
correction ($F(2.15, 21.5) = 24.9$, $p < 0.001$) followed by Tukey's multiple comparison's test. **(G)**
Relative to monocular, the positive peak of the layer 4 VEP was slightly decreased for Phase
Offset stimuli, and substantially diminished for Orthogonal stimuli. One-way repeated measures
ANOVA with Geisser-Greenhouse correction ($F(2.16, 21.6) = 19.7$, $p < 0.001$) followed by
Tukey's multiple comparison's test. **E-G:** *$p<0.05$, **$p<0.01$, ***$p<0.001$.
894
Figure 5. Multiunit activity across layers of V1 reveals changes in early and late firing that
differs for phase offset and orthogonal stimuli.
Multiunit activity was measured across layers of bV1 during the experiment shown in Figure 3.
(A) Plot of the multiunit activity envelope (MUAe) for each channel during a baseline period.
MUAe was greatest in L5. **(B-C)** MUAe for each condition was z-scored based on the baseline
MUAe shown in **A**. The z-scored MUAe for all conditions showed a brief short-latency peak in
activity corresponding in time to the VEP negativity. All cortical layers showed a statistically
meaningful visual response (z-score > 1.96). **(D)** The z-scored MUAe for each of the conditions
in **C** were compared with the Monocular (Monoc) condition in **B**. Concordant (Conc) stimuli
elicited a larger early peak in activity across cortical layers, Phase Offset (ΔPhase) stimuli
elicited a reduced early activity across all layers, and Orthogonal (ΔAngle) stimuli elicited an
extended period of elevated activity, particularly in L2/3-L5. **(E)** Due to the similarity in the
changes in activity observed across layers of V1, we collapsed the z-scored MUAe across all
layers to directly compare differences in activity in the early (40-80 ms, yellow) and late (100-
200 ms, blue) periods for each condition. **(F)** In the early period following each phase reversal,
Concordant and Orthogonal stimuli elicited greater activity than the Monocular condition, and
Phase Offset stimuli elicited reduced activity. One-way repeated measures ANOVA with
Geisser-Greenhouse correction ($F(1.54, 15.4) = 37.7$, $p < 0.001$) followed by Tukey's multiple
comparisons test. **(G)** In the late period following the initial peak in activity, Orthogonal stimuli
elicited increased activity relative to each of the other conditions. One-way repeated measures

ANOVA with Geisser-Greenhouse correction (F(1.28, 12.8) = 25.4, p < 0.001) followed by
Tukey's multiple comparisons test. Dashed lines in **A-D** indicate approximate borders between
cortical layers. **F** and **G**: **p<0.01, ***p<0.001.
918
Figure 6. Regular and fast spiking units in V1 show similar responses across conditions.
(A) Single units were classified as regular spiking (RS, black) or fast spiking (FS, blue) based on
their trough-to-peak width (see Methods). **(B)** Raster plots of action potentials for a
representative RS unit and an example FS unit for a subset of trials during each of the 4
stimulus conditions: Monocular (Monoc, light grey), Concordant (Conc, dark grey), Phase Offset
(ΔPhase, purple), and Orthogonal (ΔAngle, red). **(C)** PSTH for all RS units (putative principal cells,
n=275) averaged across all stimuli in each condition. **(D)** Average firing rate for RS units during
the early peak in activity (40-80 ms after stimulus presentation, yellow shaded region in **C**).
Relative to the Monocular condition, Concordant and Orthogonal stimuli showed an increase in
early RS activity, and Phase Offset stimuli showed decreased activity. One-way repeated
measures ANOVA with Geisser-Greenhouse correction (F(2.03, 557) = 33.78, p < 0.001)
followed by Tukey's multiple comparisons test. **(E)** Average firing rate for RS units during the
late period of activity (100-200 ms after stimulus presentation, blue shaded region in **C**).
Orthogonal stimuli elicited increased firing relative to the other three conditions. One-way
repeated measures ANOVA with Geisser-Greenhouse correction (F(1.88, 516) = 57.0, p < 0.001)
followed by Tukey's multiple comparisons test. **(F)** PSTH for all FS units (putative PV+ neurons,
n=52) averaged across all stimuli in each condition. **(G)** Average firing rate of FS units 40-80 ms
after stimulus presentation (yellow shaded region in **F**). Concordant and Orthogonal stimuli
elicited increased activity and Phase Offset stimuli elicited decreased activity relative to the
Monocular condition. One-way repeated measures ANOVA with Geisser-Greenhouse correction
(F(1.99, 102) = 14.9, p < 0.001) followed by Tukey's multiple comparisons test. **(H)** Average
firing rate of FS units 100-200 ms after stimulus presentation (blue shaded region in **F**).
Orthogonal stimuli elicited increased firing relative to all other conditions. One-way repeated
measures ANOVA with Geisser-Greenhouse correction (F(1.59, 81.0) = 27.6, p < 0.001) followed
by Tukey's multiple comparisons test. **D,E,G,H:** *p<0.05, **p<0.01, ***p<0.001
944
Figure 7. Laminar differences in responses of RS and FS units during early and late periods.
(A) Mean firing rates of RS units in each cortical layer during the early visually evoked response
(40–80 ms post-stimulus) under Monocular (light gray), Concordant (dark gray), Phase Offset
(purple), and Orthogonal (red) conditions. A significant interaction between stimulus condition
and layer was observed (two-way repeated measures ANOVA with Geisser-Greenhouse
correction: main effect of condition, F(2.05, 557) = 36.9, p < 0.001; layer, F(3, 271) = 8.27, p <
0.001; condition × layer, F(9, 813) = 2.54, p = 0.007), driven by reduced responses to Phase
Offset stimuli in L2/3, L5, and L6 but not in L4. **(B)** FS unit responses during the early time
window. No significant interaction between condition and layer was found (main effect of

condition, $F_{(1.99, 95.4)} = 15.9$, $p < 0.001$; layer, $F_{(3, 48)} = 1.85$, $p = 0.15$; condition × layer, $F_{(9,}$

$_{144)} = 1.03$, $p = 0.41$), indicating relatively uniform responses across layers. **(C)** RS unit firing

rates during the late response window (100-200 ms). Firing rates were significantly modulated

by condition ($F_{(1.89, 513)} = 60.7$, $p < 0.001$), with elevated responses to Orthogonal stimuli

observed across all layers. A main effect of layer was also observed ($F_{(3, 271)} = 2.66$, $p = 0.048$),

but there was no significant condition × layer interaction ($F_{(9, 813)} = 1.53$, $p = 0.13$). **(D)** FS unit

firing rates during the late time window. A significant condition-by-layer interaction was found

(main effect of condition, $F_{(2.14, 103)} = 58.9$, $p < 0.001$; layer, $F_{(3, 48)} = 0.64$, $p = 0.59$;

condition × layer, $F_{(9, 144)} = 14.5$, $p < 0.001$), driven by a pronounced increase in L2/3 FS

activity in response to Orthogonal stimuli. **A-D:** Error bars indicate SEM. Šídák's multiple

comparisons test used for post-hoc comparisons. *$p<0.05$, **$p<0.01$, ***$p<0.001$.

Figure 8. Orthogonal stimuli elicit larger calcium responses in L2/3 principal cells, but reduced
calcium responses in SST+ interneurons.
(A) Recording setup. Head-fixed mice expressing GCaMP with a cranial window above bV1
viewed phase-reversing grating stimuli on a dichoptic display. **(B)** Experimental conditions. Mice
viewed Monocular (contra only), Concordant (contra and ipsi at same phase and orientation),
Phase Offset (180° offset phase, same orientation), and Orthogonal (90° orientation difference)
stimuli. **(C)** Example FOV for a mouse expressing GCaMP6f in excitatory (Emx1+) cells imaged
200μm below pia (L2/3). Colored patches indicate ROIs for individual neurons. Scale
bar=100μm. **(D-H)** Responses of L2/3 pyramidal cells to each of the 4 conditions. **(E)** Scatter
plots showing average responses of individual cells to Concordant (black), Phase Offset (purple),
and Orthogonal (red) stimuli relative to Monocular. dF/F was calculated relative to baseline
response during interblock intervals. Dashed identity line on each plot indicates an equal
response to both conditions. **(F)** Cumulative distribution plot comparing the responses of all
neurons across conditions. One-way repeated measures ANOVA with Geisser-Greenhouse
correction ($F(2.59, 9724) = 124.2$, $p<0.001$) followed by Tukey's multiple comparisons test
(***$p<0.001$ relative to Monocular). **(G)** Plot showing population response to each stimulus
across all blocks for each condition. Shaded error bars indicate SEM. **(H)** Comparison of
responses for each condition by mouse (n=13). One-way repeated measures ANOVA with
Geisser-Greenhouse correction ($F(1.86, 22.28) = 27.2$, $p < 0.001$) followed by Tukey's multiple
comparisons test (*$p<0.05$, **$p<0.01$, ***$p<0.001$). **(I-M)** Responses of L2/3 SOM+ interneurons
expressing GCaMP7f for Monocular, Concordant, and Orthogonal stimulus conditions. **(J)**
Scatter plots showing average responses of individual cells to Concordant (black) and
Orthogonal (red) stimuli relative to Monocular. Dashed identity line on each plot indicates an
equal response to both conditions. **(K)** Cumulative distribution plot comparing the responses of
all SOM+ cells across conditions. One-way repeated measures ANOVA with Geisser-Greenhouse
correction ($F(1.23, 166.5) = 95.5$, $p<0.001$) followed by Tukey's multiple comparisons test
(***$p<0.001$ relative to Monocular). **(L)** Plot showing response of all SOM+ cells across time for
each stimulus condition. Shaded error bars indicate SEM. **(M)** Comparison of responses for each
condition by mouse (n=4). One-way repeated measures ANOVA with Geisser-Greenhouse
correction ($F(1.03, 3.11) = 15.7$, $p=0.026$) followed by Tukey's multiple comparisons test
(*$p<0.05$).
997
Figure 9. Circuit model for concordant and orthogonal responses
Proposed circuit diagram to account for responses elicited by concordant and conflicting visual
stimuli. Inputs from the contra eye (light grey box) drive excitatory principal cells (PCs) and
inhibitory PV+ interneurons. Feedback inhibition from SOM+ interneurons will constrain this
response in time. Concordant inputs to the ipsi eye (dark grey box) will recruit a separate
population of ipsi-responsive monocular PCs and PV+ cells. Ipsi and contra responsive cells are
reciprocally connected. Ipsi eye inputs with an orthogonal orientation (red box) will also drive

ipsi-responsive PCs and PV+ cells tuned to that orientation, but here there are not reciprocal

connections between PCs. Instead, lateral inhibition between populations of SOM+ cells will

reduce feedback inhibition onto PCs and PV+ cells, prolonging the evoked activity of these

neurons.

1009

Figure S1. Regular-spiking single unit responses to different stimulus conditions

Z-scored raster plots of regular-spiking (RS) single units for Monocular **(A)**, Concordant **(B)**,

Phase Offset **(C)**, and Orthogonal **(D)** stimuli. Units were assigned to L2/3, L4, L5, or L6 based on

the location of the electrode contact with the maximal single unit waveform, then sorted based

on their average activity between 40-80ms in the Monocular condition. Differences between

conditions and layers are quantified in Figure 5.

1016

1017   **Figure S2. Fast-spiking single unit responses to different stimulus conditions**

1018   Same as Figure S3, but for fast-spiking (FS) single units.

1019

1020

1021   **Supplemental Table 1: Key Resources**

1022

Reagent or resource	Source	Identifier
Experimental models: organisms/strains		
Mouse: C57BL/6N	Bred in lab; Originally from Charles River	RRID:MGI:5651595
Mouse: Emx1-Cre	Bred in lab; Originally from Jackson	RRID:IMSR_JAX:005628
Mouse: Ai93(TITL-GCaMP6f)-D;CaMK2a-tTA	Bred in lab; Originally from Jackson	RRID:IMSR_JAX:024108
Mouse: SOM-IRES-Cre (C57BL/6N)	Bred in lab; Originally from Jackson	RRID:IMSR_JAX:018973
Software and algorithms		
Raw data	This paper	
MATLAB 2023b	Mathworks	RRID:SCR_001622
Prism 10	Graphpad	RRID:SCR_002798
Kilosort 2.5	Github	https://github.com/MouseLand/Kilosort
Phy	Github	https://github.com/cortex-lab/phy

VEPStimulusSuite: visual stimulus generation and presentation suite	Github (Jeff Gavornik, Boston University)	https://github.com/jeffgavornik/VEPStimulusSuite
Analyses were performed using custom-written MATLAB scripts	This paper	https://github.com/danielmontgomery7/suppression-analysis-code

1023

1024

Figure 1

Figure 2

Figure 3

A

B

VEP Magnitude (% Baseline)

0° 5° 10° 15° 20° 25° 30° 90°
        *   **   ** *** ***   **

Interocular orientation difference

C

0.4 cpd

0.2 cpd

D

0.4 cpd:
0.2 cpd:

VEP Magnitude (% Baseline)

◆ 0.4 cpd
■ 0.2 cpd

0   10   20   30   40   50
     ** *** *** *** ***

Ipsi eye contrast (%)

E

Monoc

Binoc

F

Monoc:
Binoc:

VEP Magnitude (% Baseline)

● Monocular
● Binocular

0   10   20   30   40   50
     *   *** *** *** ***
         *   *   *** ***

Orthogonal stimulus contrast (%)

Figure 4

A

V1b

1.2 mm

20μm

Contra eye   Ipsi eye

B Monocular VEPs

L4

Time (msec)
0   100   200   300

C Monocular CSD

x10⁻³

Distance from L4 (μm)

L2/3
L4
L5
L6

Time (msec)
0   50   100   150   200

CSD (μV²/μm²)

D

Monocular   Concordant   Phase Offset   Orthogonal

L4 VEP

pos.
mag.
neg.

100 μV
100 ms

E

L4 VEP Magnitude (μV)

**   **   ***   **

Monoc   Conc   ΔPhase   ΔAngle

F

L4 VEP Negativity (μV)

***   *   ***   ***

Monoc   Conc   ΔPhase   ΔAngle

G

L4 VEP Positivity (μV)

***   ***   *   ***

Monoc   Conc   ΔPhase   ΔAngle

Figure 5

Figure 6

A

Norm. Voltage

— RS
— FS

Time (msec)

B   Example RS unit   Example FS unit

Monoc
Conc
ΔPhase
ΔAngle

Trial Number

Time (msec)   Time (msec)

C   RS units

Spike Rate (spikes/sec)

Monocular
Concordant
Phase Offset
Orthogonal

Time (msec)

D   RS: 40-80 ms

Spike Rate (z-score)

Monoc   Conc   ΔPhase   ΔAngle

E   RS: 100-200 ms

Spike Rate (z-score)

Monoc   Conc   ΔPhase   ΔAngle

F   FS units

Spike Rate (spikes/sec)

Time (msec)

G   FS: 40-80 ms

Spike Rate (z-score)

Monoc   Conc   ΔPhase   ΔAngle

H   FS: 100-200 ms

Spike Rate (z-score)

Monoc   Conc   ΔPhase   ΔAngle

Figure 7

Figure 8

Figure 9

Figure S1

Figure S2