Methanogens are associated with altered microbial production of short-chain fatty acids and human-host metabolizable energy

Blake Dirks[1,2], Taylor L. Davis[1,2], Elvis A. Carnero[3], Karen D. Corbin[3], Steven R. Smith[3], Bruce E. Rittmann[2,4] & Rosa Krajmalnik-Brown[1,4*]

[1]Biodesign Center for Health through Microbiomes, Arizona State University, Tempe, AZ, USA. [2]Biodesign Swette Center for Environmental Biotechnology, Arizona State University, Tempe, AZ, USA. [3]AdventHealth Translational Research Institute, Orlando, FL, USA. [4]School of Sustainable Engineering and the Built Environment, Arizona State University, Tempe, AZ, USA.

* Corresponding author: dr.rosy@asu.edu

Abstract

Methanogens are CH_4-producing, H_2-oxidizing archaea. Numerous studies have associated methanogens with obesity, but these results have been inconsistent. One possible link to host metabolism may be tied to the methanogens' ability to consume H_2, thus reducing the H_2 partial pressure and thermodynamically enhancing fermentation of sugars to short-chain fatty acids (SCFA) that the host can absorb. Because research linking methanogenesis to human metabolism is limited, our goal with this exploratory analysis was to investigate relationships between methanogens and other H_2-consuming (i.e., hydrogenotrophic) microorganisms, and the association of methanogens with human metabolizable energy. Using results from a tightly controlled randomized crossover feeding study with well-characterized human participants and novel continuous methane measurements, we analyzed hydrogenotroph abundance and activity, fecal and serum SCFAs, and host metabolism between high and low CH_4 producers. Methanogens were detected in about one-half of participants, and most were high CH_4 producers. We found no evidence that methanogens' consumption of H_2 to produce CH_4 affected the other hydrogenotrophs, but high CH_4 producers had greater serum propionate and host metabolizable energy irrespective of diet. High CH_4 producers also had greater gene and transcript abundance of a key enzyme of the H_2-consuming, propionate-producing succinate pathway. Finally, a network analysis revealed positive relationships between *Methanobrevibacter smithii* (the most prevalent methanogen in the human colon) and bacteria capable of degrading fiber and fermenting fiber-degradation products, thus forming a trophic chain to extract additional energy from undigested substrates. Our results show that methanogens in a microbial consortium were linked to host metabolizable energy through enhanced SCFA production and enhanced host absorption of SCFA.

INTRODUCTION

The human body hosts around 38 trillion bacterial cells, most of which reside in the colon [1]. Besides bacteria, the intestinal microbiota also includes archaea, protists, and viruses [2]. One of the important functions of the intestinal microbiota is fermentation [3]. Unmetabolized macronutrients entering the colon become substrates for bacterial hydrolysis and fermentation, which generates simple products, such as H_2, CO_2, and short-chain fatty acids (SCFAs) [4].

While SCFAs have received much attention due to their impact on host health [5], H_2 remains understudied despite its importance to the microbiome. High H_2 partial pressure inhibits bacteria from regenerating NAD^+ from NADH, which decreases the fermentation of complex substrates and impairs microbial growth [6,7]. H_2 partial pressure also influences the thermodynamics of SCFA production: Low partial pressure favors acetate and butyrate production, while high partial pressure favors propionate production [8].

The H_2 produced by fermentation can be oxidized by microbes known as hydrogenotrophs in well-known microbial processes that produce acetate, sulfide, or methane [9]. The three common groups of hydrogenotrophic microorganisms in the human colon are homoacetogenic bacteria, sulfate-reducing bacteria (SRB), and methanogenic archaea. These three groups can use H_2 as an electron donor to produce energy [9]. **Figure 1** summarizes the hydrogenotrophic groups, their substrates and products, and the fate of their products.

Figure 1. Hydrogenotrophic processes in the human colon when the macronutrient input is carbohydrate. Fiber that reaches the colon is hydrolyzed and fermented by the microbiome to produce SCFAs, H_2, and CO_2. Most SCFAs are absorbed by the host or excreted in feces. While some H_2 is released through the breath or flatus, most is metabolized in the colon by homoacetogenesis, dissimilatory sulfate reduction, or methanogenesis. Acetate produced though hydrogenotrophy is absorbed by the host or excreted in feces. CH_4 and some hydrogen sulfide produced through hydrogenotrophy are released through the breath or flatus.

Homoacetogenic bacteria, which include species in the genera *Blautia*, *Clostidium*, and *Ruminococcus*, oxidize H_2 and reduce CO_2 to make acetate using the Wood-Ljungdahl pathway [10]. The stoichiometry for homoacetogenesis is:

$$4H_2 + 2CO_2 \rightarrow CH_3COOH + 2H_2O \qquad (1)$$

Homoacetogens carrying out reaction (1) are autotrophs, which means that their carbon source is inorganic carbon, or CO_2. However, most homoacetogens are not obligate hydrogenotrophs and can ferment organic substrates, such as glucose, as the electron donor and carbon source [11].

SRB in the genera *Desulfovibrio* and *Fusobacteria* oxidize H_2 and reduce sulfate to sulfide through dissimilatory sulfate reduction [12]:

$$4H_2 + SO_4^{2-} + 2H^+ \rightarrow H_2S + 4H_2O \qquad (2)$$

Like homoacetogens, SRB can also ferment organic substrates, such as lactate and pyruvate [11].

The predominant methanogen in the human colon is *Methanobrevibacter smithii*, although *Methanosphaera stadtmanae* and *Methanomassiliicoccus* spp are sometimes detected [13]. Methanogens can be categorized into three groups: 1) hydrogenotrophic methanogens that reduce CO_2 (or formate); 2) acetoclastic methanogens that ferment acetate to CH_4 and CO_2; and 3) methylotrophic methanogens that oxidize and reduce methanol to CH_4 and CO_2 [14]. All methanogens found in human intestines so far are hydrogenotrophic [15]:

$$4H_2 + CO_2 \rightarrow CH_4 + 2H_2O \qquad (3)$$

Unlike homoacetogens and SRB, hydrogenotrophic methanogens do not have alternative metabolic pathways: They rely solely on H_2 and CO_2 for their metabolism, and detected CH_4 is always produced via a H_2-consuming process.

CH_4 also is unique among the hydrogenotrophs because it is only made by archaea and is not metabolized by the human body [16]. In contrast, H_2S is produced from other microbial sulfur metabolisms [17] and metabolized by mitochondria in host colonocytes [18]. Likewise, acetate is produced through fermentation by many bacteria [19], and it is a common substrate for any respiring bacteria [20].

Some evidence supports that methanogens have an impact on host metabolism in mouse models [11]. For example, mice inoculated with methanogens showed increased weight and adiposity despite consuming the same amount of food as controls [21,22]. However, the role of methanogens in human metabolism is complicated and controversial. Studies have found conflicting correlations between methanogens and obesity/leanness [23] and anorexia [24].

In this exploratory analysis, we investigated relationships between methanogens and the other hydrogenotrophs, along with the impact of methanogens on human-host metabolism, in samples from a tightly controlled randomized crossover feeding study with well characterized human participants [25]. Briefly, Corbin et al. [25] evaluated the microbial contribution to human-host energy balance using two distinctly different diets: the Western Diet (WD) and the Microbiome-enhancer Diet (MBD). The WD was comprised of foods that were low in fiber and resistant starch, small in particle size (such as peanut butter vs. whole nuts on the MBD), and included processed foods. The MBD, in contrast, was designed to deliver more microbial substrates to the colon by being less absorbable by the host. The MBD contained more whole foods, fiber, resistant starch, and was limited in processed foods. They found that diet altered microbiome's structure, and that the microbiome's metabolic activity contributed to host metabolizable energy (ME).

The results from Corbin et al. [25] are particularly well-suited for a deeper investigation into methanogens and their role in the microbiome and host metabolism. On the one hand, that study strictly accounted for energy input (diet), energy expenditure, and energy output (urine, feces, gas). On the other hand, they continuously measured CH_4 production during the inpatient portion of the study using a first-in-human method within a whole room calorimeter [26]. Combining the study's CH_4 measurements with its multi-omic data allows us to detect relationship among the methanogens, the other hydrogenotrophs, and the human host.

Since the hydrogenotrophs compete for H_2, we hypothesized that high methanogenic activity would lead to a lowering of homoacetogenic and sulfate-reduction activities. These changes in the microbial community would then be associated with factors related to the human's metabolism: e.g., fecal SCFA output, serum SCFA concentration, and host ME.

Methods

Overview of Clinical Study

Details of the clinical study (NCT02939703) from which the data and samples for this work were derived were previously published [25,26]. Briefly, the clinical study was approved by the AdventHealth Institutional Review Board and conducted at AdventHealth Translational Research Institute in Orlando, Florida. After a health screening, 17 participants (9 men and 8 women) completed the study, which was a randomized crossover-controlled feeding study with the WD as a control and the MBD as an intervention. This design minimized the impact of confounders as each participant served as their own control. The study period took place over 61 days. Each participant's caloric requirement was determined during the baseline period (days 1-9), and meals were prepared uniquely for each participant to maintain energy balance. Participants consumed those meals outpatient for 11 days then inpatient for the next 11 days with a >14-day washout between diet periods. During the 11-day inpatient stay for each diet, each participant's energy expenditure was measured in whole room calorimeter for 6 days and fecal samples were collected.

CH$_4$ Measurements

CH$_4$ release was measured continuously during each 6-day period that the participant was in the whole-room calorimeter. CH$_4$ concentration was measured with an off-axis integrated-cavity output spectroscopy (OA-ICOS). A detailed description and validation of the method can be found in Carnero et al. [27].

Colonic Transit Time

Colonic transit time (CTT) was measured while participants were in the whole-room calorimeter. Participants ingested a SmartPill™ (Medtronic) that sent data to a sensor worn by the participants recording temperature, pressure, and pH [26].

Host Metabolizable Energy (ME)

Briefly, host ME, the energy from the diet available to the host [28], was computed as the total energy intake minus the energy lost in the feces. Calculation details are published Corbin et al. [25].

Fecal and Serum Short Chain Fatty Acids

Fecal and serum SCFAs were quantified by targeted metabolomics (Metabolon, Inc., Mooresville, NC). Fecal SCFAs were normalized as published in Corbin et al. [25].

Quantification of Key Hydrogenotroph Genes

The abundances of the homoacetogens, SRB, and methanogens were measured using the quantitative polymerase chain reaction (qPCR) for genes that identify each hydrogenotroph: *acsB* for homoacetogens, *dsrA* for SRB, and *mcrA* for methanogens. Primers, thermocycler settings, standards, and references for each gene are summarized in **Table 1**. All qPCR assays were performed on a Thermofisher Applied Biosystems Quant Studio 3. Standards for each gene were custom IDT gBlocks gene fragments of the target genes from a representative microbe of each hydrogenotroph group. 7-point calibration curves for each assay were generated in triplicate using gene copy numbers ranging from 10^1 to 10^8 of their respective gene standards.

The qPCR values were transformed from log to exponential values, normalized to daily fecal output, giving us daily fecal copy number (gene copy numbers/day). Since each hydrogenotroph contains one gene copy per cell of their respective gene [29–31], qPCR measurements of gene copy number provided an accurate estimate of hydrogenotroph cell numbers/day in the feces.

Table 1: qPCR primer and themocycler details

Group	Gene	Primers	Thermocycler Settings	Reference	Standards from
Homoacetogens	*acsB*	ACS_f ACS_r	95°C 10 s, 52°C 20 s, 72°C 30 s, 40 cycles	[32]	*Blautia hydrogenotrophica* DSM 10507
Sulfate-reducing bacteria	*dsrA*	DSR1-F DSR-R	95°C 10 s, 60°C 60 s, 35 cycles	[33]	*Desulfovibrio piger* isolate FI11049
Methanogens	*mcrA*	mlas mcrA-rev	95°C 30 s, 55°C 45 s, 72°C 30 s, 40 cycles	[34]	*Methanobrevibacter smithii* ATCC 35061

DNA Sequencing and Sequence Processing

DNA sequencing and subsequent sequence processing were performed as previously described in Corbin et al. [25].

RNA Sequencing

Fecal-sample processing, RNA extraction, library prep, and mRNA sequencing were performed at the University of North Carolina at Chapel Hill Microbiome Core (Chapel Hill, NC, USA), which is supported by the following grants: Gastrointestinal Biology and Disease (CGIBD P30 DK034987) and the UNC Nutrition Obesity Research Center (NORC P30 DK056350). RNA was extracted using the Qiagen RNeasy PowerMicrobiome Kit (Cat No./ID: 26000-50). RNA depletion was performed using QIAseq FastSelect –5S/16S/23S Kit (Cat No./ID: 335925), and library prep was performed using QIAseq Stranded Total RNA Lib Kit (Cat No./ID: 180745). Samples were sequenced using the Illumina HiSeq 4000 PE150 platform. To avoid batching effects, fecal samples were randomized prior to nucleic acid extraction and all samples were sequenced at the same time.

RNA-Sequence Processing

RNA-sequencing outputs were quality controlled with FastQC (version 0.12.0) [35]. Adapters were trimmed using TrimGalore (version 0.6.5) [36], RNA sequences were aligned against Hg38 (GRCh38.p14) using STAR (version 2.7.11a) [37], aligned sequences were removed, and reads were then paired and annotated using HUMAnN3 (version 3.8) [38] using standard parameters. A small pseudo-count (equal to half of the lowest non-zero count) was added to any zeros and transcript abundances were centered log-ratio transformed as suggested for the analysis of compositional data [39,40].

Wilcoxon Signed Rank Test

The Wilcoxon signed rank test was used to compare the median $\log_{10}$(daily fecal copy number/day) for hydrogenotroph marker genes *mcrA* (methanogens), *acsB* (homoacetogens), and *dsrA* (SRB). Samples in which a gene was undetected were given a gene copy number of 1 and then log transformed.

Categorizing High and Low Methane Producers

Study participants were categorized as either high or low CH_4 producers. First, methane production rates were $\log_{10}$ transformed and then tested for multimodality by the excess mass test (Excess mass = 0.22, p-value < 2.2e-16) using the modetest function from the R package "multimode" (version 1.5) [41]. Next, the threshold for high CH_4 production was set at 37 mL CH_4/day using the locmodes function. Participants were evaluated on a per sample basis and categorized as a high or low methane producer irrespective of diet.

Linear Mixed Model

We used a linear mixed model (LMM) to test the relationship between high and low CH_4 producers, our primary independent variable of interest, and outcome variables. We accounted for the effects of diet, diet period, and diet sequence by including them as covariates in our LMMs. We included participant ID as a random factor to account for the non-independence of the samples. Details about dependent and independent variables, covariates, and random term used in our LMMs can be found in **Supplement Table 1**. LMMs were run using the lmer command from the R package "lmerTEST" (version 3.1-3) [42].

Prior to running each LMM, variable-distribution normality was evaluated using qqnorm and shapiro.test commands, both of which were from the R package "stats" (version 4.2.2) [43]. Non-normal data were transformed to improve normality. Fecal and serum SCFAs were square-root transformed. Because host ME was negatively skewed (or left-tailed), the ME values were reflected before they were transformed, meaning each ME value was subtracted from a constant (the maximum ME value plus 0.1) and then log transformed [44]. The Benjamini-Hochberg method was used to correct for multiple comparisons and adjusted p-values ≤0.10 were considered significant. The same analyses were completed with CH_4 as a continuous variable. The results of the continuous-variable analyses are in the Supplementary Data.

Network Analysis

Microbiome network analysis was performed using the Cytoscape (version 3.10.2) [45] plugin Conet (version 1.1.1) [46] an ensemble co-occurrence analysis tool. Conet was run with default settings using Pearson, Spearman, Mutual Information, Bray-Curtis dissimilarity, and Kullback-Leibler dissimilarity to detect associations between microbes. The Benjamini-Hochberg method was used to correct for multiple comparisons and adjusted p-values ≤0.5 were considered significant.

Results and Discussion

Homoacetogens and SRB were present in all participants, but not all participants had methanogens

We first investigated the abundance of methanogens, homoacetogens, and SRB among the trial participants. We performed qPCR targeting genes that encode enzymes in the hydrogenotrophic pathways for each group: *mcrA* for methanogens, *acsB* for homoacetogens, and *dsrA* for SRB.

We detected methanogens in only 9 participants for both diets and one participant for only the MBD (**Figure 2**). Consequently, the average $\log_{10}$(gene copy number/day) was 5.3±4.8 for methanogens. In contrast, we detected homoacetogens and SRB in all 17 participants for both diets. On average, homoacetogens were more abundant than SRB: $\log_{10}$(gene copy number/day) of 10.2±0.46 for homoacetogens versus 8.8±0.86 for SRB, but homoacetogens and SRB were more abundant than methanogens. Detecting methanogens in only some participants is consistent with past results showing that the abundance of *M. smithii* in the human gut was either high or very low [47,48]. The presence or absence of methanogens can have important implications for hydrogenotrophs and the microbiome. In mixed-culture experiments, SRB became the dominant hydrogenotroph when methanogens were inhibited [11]. In co-culture experiments, *Bacteroides thetaiotaomicron* expressed more glycoside hydrolases, enzymes that break down polysaccharides, and degraded and fermented more glycans when the methanogen *M. smithii* was present, compared to when the SRB *D. piger* was present [49].

Figure 2. Abundances of hydrogenotrophs in daily fecal copy numbers ($\log_{10}$ (gene copy/day)) measured by qPCR. Homoacetogens and SRB were detected in all participants in both diets. Methanogens were only detected in 9 participants in both diets and one participant in the MBD. Homoacetogens were more abundant than SRB and methanogens. SRB were more abundant than methanogens. Abundance medians are marked by the horizontal black bars. Comparison tests between abundance medians were performed with Wilcoxon signed rank test. *: adj. p-value <0.10, **: adj. p-value<0.05, ***: adj. p-value<0.01, ****: adj. p-value<0.0001. NS is non-significant with an adjusted p value > 0.10.

Homoacetogens and SRB did not show patterns of competition with methanogens

The absence of methanogens in a portion of the participants might lead to differences in competition for H_2 among the hydrogenotrophs. We hypothesized that, in the absence of methanogens, homoacetogens or SRB (or both) would become more abundant and active. To test this hypothesis, we compared mean methanogen, SRB, and homoacetogen daily fecal copy numbers and gene and transcript abundances between high and low CH_4 producers

High CH_4 producers had higher methanogen daily copy numbers in the feces and the abundance of *mcrA* genes and transcripts, compared to low CH_4 producers (**Figure 3A**). In contrast, homoacetogen daily copy numbers, *acsB* genes, and *acsB* transcripts were not significantly different between groups (**Figure 3B**). While SRB copy numbers were higher in high CH_4 producers, *dsrA* gene and transcript abundances were not significantly different between groups (**Figure 3C**). The trends were similar when this analysis was done using CH_4 production as a continuous variable (**Figure S1**).

These results suggest that competition for H_2 was not a factor limiting the growth and accumulation of SRB and homoacetogens. One explanation is that SRB and homoacetogens have diverse metabolisms and do not rely solely on H_2 as an electron donor [50,51]. For example, SRB can utilize lactate for sulfate respiration [17] or for fermentation in the absence of sulfate [52,53]. Homoacetogens can also ferment sugars instead of oxidizing H_2 during hydrogenotrophic homoacetogenesis [54]. Although homoacetogens can grow mixotrophically, using hydrogenotrophy and fermentation simultaneously for energy generation, an experiment evaluating the mixotrophy of *Blautia coccoides*, a homoacetogen found in the human gut, revealed that hydrogenotrophy was inhibited by glucose [55]. The high abundance of *acsB* genes relative to the low abundance of *acsB* transcripts suggests that homoacetogens may not have been relying on H_2 for electrons. A second explanation is that the electron flow to CH_4 was too small to have had a major impact on the other hydrogenotrophs' metabolic function: Electron flow from COD ingested to CH_4 was approximately 1% of the electron-equivalent intake to the large intestine (Davis et al. 2024).

Figure 3. Comparisons of mean hydrogenotroph daily fecal copy numbers, gene and transcript abundance between CH₄ groups. *A) Methanogen daily fecal copy numbers, mcrA genes, and mcrA transcripts were significantly higher in high CH₄ producers than low CH₄ producers. B) Homoacetogen daily fecal copy numbers, acsB genes, and acsB transcripts were not significantly different between CH₄ groups. C) Sulfate reducing bacteria (SRB) daily fecal copy numbers were significantly higher in high CH₄ producers than low CH₄ producers, but dsrA genes, and dsrA transcripts were not significantly different between CH₄ groups. CLR is the centered log-ratio transformation. Group means are marked by a white diamond. Group medians are marked by horizontal black line. All means comparisons were made by linear mixed model. *: adj. p-value <0.10, **: adj. p-value<0.05, ***: adj. p-value<0.01, ****: adj. p-value<0.0001. NS is non-significant with an adjusted p value > 0.10.*

High CH₄ producers had higher serum propionate concentrations than low CH₄ producers

Although the hydrogenotrophs did not appear to compete for H_2 and the flow of electrons into CH_4 was small, CH_4 production still may have had an impact on the H_2 concentration in the large intestine. Because fermentation is inhibited by high H_2 partial pressure [56], additional H_2 consumption by methanogens may still have relieved a thermodynamic inhibition on fermentation and led to greater fecal SCFA outputs and serum SCFA concentrations. Fecal SCFA were not significantly different between groups, although the means were consistently lower for the high CH_4 producers (**Figure 4A**). In contrast, serum propionate concentrations were higher in high CH_4 producers than low producers (**Figure 4B**). Total fecal SCFAs, fecal acetate, fecal propionate, and serum propionate were significantly positively associated with CH_4 when these analyses were done using CH4 as a continuous variable (**Figures S2**). Isotope studies have shown that serum propionate is mostly microbially produced in the intestine [20,57]. Therefore, the opposing relationships of fecal SCFA and serum propionate with CH_4 production suggest a faster rate of SCFA uptake in participants that produced more methane.

Figure 4. Comparisons of mean fecal SCFA output and serum SCFA concentration between CH₄ groups.** A) Total fecal SCFA, fecal acetate, fecal propionate and fecal butyrate were not significantly different between CH₄ groups. B) Serum propionate concentrations were significantly higher in high CH₄ producers than low CH₄ producers, but total serum SCFA, serum acetate, and serum butyrate were not significantly different between CH₄ groups. Fecal SCFA output and serum SCFA concentrations were square root transformed. Group means are marked by a white diamond. Group medians are marked by horizontal black line. All means comparisons were made by linear mixed model. *: adj. p-value <0.10, **: adj. p-value<0.05, ***: adj. p-value<0.01, *: adj. p-value<0.0001. NS is non-significant with an adjusted p value > 0.10.*

High CH$_4$ producers had a higher abundance of gene and transcripts for a propionate-producing pathway than low CH$_4$ producers

Given the positive association between serum propionate and CH$_4$-production rate, we investigated if propionate production could have been altered by methanogenesis. We evaluated the relationship between the CH$_4$-production rate and the abundance of key microbial genes and transcripts for microbial pathways for propionate production. The common propionate-producing pathways, the succinate and acrylate pathways [58], consume H$_2$ [59]. The succinate pathway has two variants, one using methylmalonyl-CoA decarboxylase (*Mmd*) and the other using methylmalonyl-CoA carboxyltransferase (*MMCT*). The acrylate pathway is characterized by the acryloy-CoA reductase (*Acr*) [58,60]. The abundances of the *Mmd* genes and transcripts were higher in high CH$_4$ producers (**Figure 5A**), *MMCT* genes and transcripts were not different between groups (**Figure 5B**), and *Acr* gene abundance, but not transcript abundance, was higher in high CH$_4$ producers (**Figure 5C**). Similar results were obtained when this analysis was done using methane as a continuous variable (**Figure S3**).

Miceli et al. [61] showed that methanogenic microbial communities, while maintaining CH$_4$ production rates, increased propionate production in response to increased carbohydrate availability for fermentation. We calculated the average amount of H$_2$ consumed by methanogenesis and propionate production in participants with detectable methanogen fecal copy numbers. The calculations can be found in the Supplementary Materials. We found that methanogenesis consumed on average around 0.12 e$^-$ eq /day of H$_2$, whereas we estimated that propionate production could have consumed as much as 716 e$^-$ eq/day of H$_2$, assuming all propionate was produced via the succinate pathway. Our results and calculations support the notion that electron flow to propionate production was substantially greater than to methanogenesis.

Figure 5. Comparison of gene and transcript abundance for the propionate-producing succinate (Mmd and MMCT) and acrylate (Acr) pathways between CH₄ groups. *A) Mmd (methylmalonoyl-CoA decarboxylase) gene and transcript abundance were significantly higher in high CH_4 producers than low CH_4 producers. B) MMCT (methylmalonyl-CoA carboxyltransferase) gene and transcript abundance were not significantly different between high and low CH_4 producers. C) Acr (acroyl-CoA reductase) genes were significantly higher in high CH_4 producers than low CH_4 producers, but transcripts were not significantly different between high and low CH_4 producers. CLR is centered log-ratio transformation. Group means are marked by a white diamond. Group medians are marked by horizontal black line. All means comparisons were made by linear mixed model. *: adj. p-value <0.10, **: adj. p-value<0.05, ***: adj. p-value<0.01, ****: adj. p-value<0.0001. NS is non-significant with an adjusted p value > 0.10.*

Host ME was greater in high CH₄ producers

Absorption of microbially generated SCFA can contribute up to 10% of daily caloric uptake [25,62], and our results indicate that high CH_4 production may be an indicator or biomarker of increased SCFA absorption by the host. Thus, we compared host ME between high and low CH_4 producers. High CH_4 producers had a significantly higher host ME than low producers (**Figure 6A**). When the analysis was performed with CH_4 as a continuous variable (**Figure 6B**), the relationship was significantly positive for all participants on both diets, and each diet showed a similar correlation, although all ME values were higher for the WD than the MBD. Because methanogens in the human colon comprise only around 1.2% of the microbiome [63] and rely solely on H_2 plus CO_2 or formate for energy [64,65], they neither produce nor consume SCFAs. In principle, methanogens could have increased host ME by consuming H_2 and thermodynamically accelerating fermentation to SCFAs. However, as we showed above, the electron flow to CH_4 was small relative to the potential electron flow to propionate. A more compelling explanation is that methanogens were a key component of a microbial community that enabled greater host uptake of energy from the large intestine.

Figure 6. Host ME in high and low CH₄ producers and by CH₄ production rate.** A)Host ME was significantly higher in high CH₄ producers than low CH₄ producers. B) Host ME was significantly positively associated with CH4 production. β is the standardized effect size for the linear mixed model describing the relationship between Host ME and CH₄ production. WD is the western diet. MBD is the microbiome enhancer diet. Group means are marked by a white diamond. Group medians are marked by horizontal black line. All means comparisons and variable relationships were made by linear mixed model. *: adj. p-value <0.10, **: adj. p-value<0.05, ***: adj. p-value<0.01, *: adj. p-value<0.0001. NS is non-significant with an adjusted p value > 0.10.*

Fiber-degrading bacteria and propionate-producing bacteria co-occur with *M. smithii*

Considering the positive association between host ME and CH_4-production rate, along with methanogens' inability to degrade fiber or ferment sugars, we looked for co-occurring bacteria that made the methanogenic microbial community better equipped to degrade and ferment organic substrates. We used CoNet [46], a network co-occurrence inference tool, to find bacteria strongly associated with *Methanobrevibacter smithii*, the predominant methanogen detected in our samples. After correction for multiple comparisons, 19 bacteria were positively correlated, and 8 bacteria were negatively correlated with *M. smithii* (**Figure 7**).

Bacteria positively associated with *M. smithii* were either fiber degraders -- *Bacteroides caccae* [66], *B. massiliensis* [67], *Bifidobacterium adolescentis* [68], *Coprococcus catus* [69], and *C. eutactus* [70]) -- or a diverse group of fermenters -- (*Alistipes shahii* [71], *Catenibacterium mitsuokai* [72], *C. leptum* [73], *Colinsella. stercoris* [74], *Holdemanella biformis* [75], *Odoribacter splanchnius* [76], and *Streptococcus thermophilus* [77]). Four of the 19 positively associated bacteria produce propionate via the succinate pathway (*B. caccae, B. massiliensis, O. splanchnicus,* and *Phascolarctobacterium succinatutens* [78]), which aligns with the results on propionate-producing pathways presented in **Figure 5**.

Bacteria negatively associated with *M. smithii* were fermenters -- such as *Clostridium bolteae* [79]*, C. symbiosum,* and *Flavonifactor plautii* [80] -- mucin-degraders -- such as *Ruminococcus gnavus* [81] and asaccharolytic bacteria -- or bacteria that cannot metabolize sugars, such as *Eggerthela lenta* [82]. All these bacteria, except *E. lenta,* can produce acetate and either ethanol or lactate. This ability is important because it allows these bacteria to continue fermentation despite high H_2 partial pressure by shifting production from acetate to less reduced ethanol or lactate [6,83]

We interpret that a network of fiber degraders, fermenters, propionate producers, and methanogens provided the host with more SCFA and contributed to higher host ME (**Figure 6**). **Figure 8** shows how the bacteria positively associated with *M. smithii* fit in a hypothesized trophic chain. Fiber degraders broke down polysaccharides into sugars; fermenters consumed the sugars to release acetate, butyrate, propionate, and H_2; and methanogens and propionate producers consumed the H_2 and kept fermentation thermodynamically favorable. Thus, the methanogenic microbial communities could utilize more complex substrates than non-methanogenic microbial communities; this explains how the methanogenic communities could produce more SCFAs for the host to absorb.

Figure 7. **Network from co-occurrence analysis of** **M. smithii** **and associated bacteria.** *Red lines represent negatively associated bacteria, and green lines represent positively associated bacteria. Asaccharolytic bacteria lack the ability to degrade sugars and utilize other substrates such as amino acids or cross-feed on bacterial metabolites such as lactate. All relationships were statistically significant after correcting for multiple comparisons (adj.p < 0.05).*

Figure 8. Trophic chain formed in a methanogenic gut microbiome. *The bacteria positively associated with* M. smithii *form a trophic chain that efficiently degrades and ferments fiber and effectively removes excess H_2 produced through fermentation by methanogenesis and the propionate-producing succinate pathway.*

Conclusion and Future Directions

Our results show that methanogens and methane production were present in only about one-half of the study participants. While methanogen fecal copy numbers and *mcrA* genes and transcripts were higher in high CH_4 producers, SRB and homoacetogen fecal copy numbers, genes, and transcripts had no relationship with CH_4 production. Consequently, we did not see evidence that methanogens' uptake of H_2 to produce methane affected the other hydrogenotrophs. This probably occurred because SRB and homoacetogens, unlike methanogens, do not rely on H_2 oxidation for energy generation. Methanogenesis was associated with higher host ME and serum propionate, but not with fecal SCFAs, suggesting that methanogens were linked to enhanced SCFA uptake. We also found that bacteria co-occurring with methanogens are well-suited to degrade and ferment fiber, as well as consuming H_2 through propionate production. Taken altogether, our results add important missing mechanistic insights into the relationship between methanogens and obesity. Methanogens appear to be part of a microbial consortium capable of enhanced energy extraction and absorption.

Future research should focus on investigating whether methane or methanogens affect absorption of SCFAs in the colon. As we show, the impact that methanogens have on host metabolism may not be in removing H_2, but rather by influencing other physiological parameters, such as SCFA production and possibly SCFA absorption. Alternatively, methanogens might simply constitute a detectable signal for enhanced energy extraction. Understanding the interactions of methanogens with the human host and other microorganisms in the human large intestine could lead to dietary or other means to modulate methanogen activity in ways that improve the host's metabolic health. Evaluating the presence and activity of methanogens that accumulate on the intestinal epithelium would provide important information that cannot be obtained solely from fecal samples. Additionally, future experiments that involve *in vitro* microbial communities of homoacetogens, SRB, and methanogens would allow for direct measurements of all hydrogenotrophic activities for a wide range of conditions; this would provide a higher resolution picture of the structural and functional relationships among the hydrogenotrophs.

References

1. Sender R, Fuchs S, Milo R. Revised Estimates for the Number of Human and Bacteria Cells in the Body. *PLoS Biol* 2016;**14**:e1002533.

2. Lee CJ, Sears CL, Maruthur N. Gut microbiome and its role in obesity and insulin resistance. *Ann N Y Acad Sci* 2020;**1461**:37–52.

3. Fan Y, Pedersen O. Gut microbiota in human metabolic health and disease. *Nat Rev Microbiol* 2021;**19**:55–71.

4. Bernalier-Donadille A. Fermentative metabolism by the human gut microbiota. *Gastroentérologie Clin Biol* 2010;**34**:S16–22.

5. Hee B Van Der, Wells JM, van der Hee B *et al.* Microbial Regulation of Host Physiology by Short-chain Fatty Acids. *Trends Microbiol* 2021;**29**:700–12.

6. Wolin MJ, Miller TL. Bioconversion of organic carbon to ch4 and co2. *Geomicrobiol J* 1987;**5**:239–59.

7. Thauer RK, Jungermann K, Decker K. Energy Conservation in Chemotrophic Anaerobic Bacteria. *Bacteriol Rev* 1977;**41**:809–809.

8. Janssen PH. Influence of hydrogen on rumen methane formation and fermentation balances through microbial growth kinetics and fermentation thermodynamics. *Anim Feed Sci Technol* 2010;**160**:1–22.

9. Kelly WJ, Mackie RI, Attwood GT *et al.* Hydrogen and formate production and utilisation in the rumen and the human colon. *Anim Microbiome* 2022;**4**, DOI: 10.1186/s42523-022-00174-z.

10. Karekar S, Stefanini R, Ahring B. Homo-Acetogens: Their Metabolism and Competitive Relationship with Hydrogenotrophic Methanogens. *Microorganisms* 2022;**10**:1–22.

11. Smith NW, Shorten PR, Altermann EH *et al.* Hydrogen cross-feeders of the human gastrointestinal tract. *Gut Microbes* 2019;**10**:270–88.

12. Kushkevych I, Cejnar J, Treml J *et al.* Recent Advances in Metabolic Pathways of Sulfate Reduction in Intestinal Bacteria. *Cells* 2020;**9**:1–16.

13. Borrel G, Brugère JF, Gribaldo S *et al.* The host-associated archaeome. *Nat Rev Microbiol* 2020;**18**:622–36.

14. Nakamura N, Lin HC, McSweeney CS *et al.* Mechanisms of microbial hydrogen disposal in the human colon and implications for health and disease. *Annu Rev Food Sci Technol* 2010;**1**:363–95.

15. Lyu Z, Shao N, Akinyemi T *et al.* Methanogenesis. *Curr Biol* 2018;**28**:R727–32.

16. Chaudhary PP, Conway PL, Schlundt J. Methanogens in humans: potentially beneficial or harmful for health. *Appl Microbiol Biotechnol* 2018;**102**:3095–104.

17. Barton LL, Ritz NL, Fauque GD *et al.* Sulfur Cycling and the Intestinal Microbiome. *Dig Dis Sci* 2017;**62**:2241–57.

18. Bouillaud F, Blachier F. Mitochondria and Sulfide: A Very Old Story of Poisoning, Feeding, and Signaling? *Antioxid Redox Signal* 2011;**15**:379–91.

19. Hosmer J, Mcewan AG, Kappler U. Bacterial acetate metabolism and its influence on human epithelia. *Emerg Top Life Sci* 2023;**0**:1–13.

20. den Besten G, Lange K, Havinga R *et al.* Gut-derived short-chain fatty acids are vividly assimilated into host carbohydrates and lipids. *Am J Physiol - Gastrointest Liver Physiol* 2013;**305**:900–10.

21. Mathur R, Kim G, Morales W *et al.* Intestinal Methanobrevibacter smithii but not total bacteria is related to diet-induced weight gain in rats. *Obesity* 2013;**21**:748–54.

22. Samuel BS, Gordon JI. A humanized gnotobiotic mouse model of host-archaeal-bacterial mutualism. *Proc Natl Acad Sci U S A* 2006;**103**:10011–6.

23. Crovesy L, Masterson D, Rosado EL. Profile of the gut microbiota of adults with obesity: a systematic review. *Eur J Clin Nutr* 2020;**74**:1251–62.

24. Singer-Englar T, Barlow G, Mathur R. Obesity, diabetes, and the gut microbiome: an updated review. *Expert Rev Gastroenterol Hepatol* 2019;**13**:3–15.

25. Corbin KD, Carnero EA, Dirks B *et al.* Host-diet-gut microbiome interactions influence human energy balance: a randomized clinical trial. *Nat Commun* 2023;**14**:3161.

26. Corbin KD, Krajmalnik-Brown R, Carnero EA *et al.* Integrative and quantitative bioenergetics: Design of a study to assess the impact of the gut microbiome on host energy balance. *Contemp Clin trials Commun* 2020;**19**:100646.

27. Carnero EA, Bock CP, Liu Y *et al.* Measurement of 24-h continuous human CH4 release in a whole room indirect calorimeter. *J Appl Physiol* 2023;**134**:766–76.

28. Elia M, Cummings JH. Physiological aspects of energy metabolism and gastrointestinal effects of carbohydrates. *Eur J Clin Nutr* 2007;**61**:S40–74.

29. Håvelsrud OE, Haverkamp THA, Kristensen T *et al.* A metagenomic study of methanotrophic microorganisms in Coal Oil Point seep sediments. *BMC Microbiol* 2011;**11**:221.

30. Yao Y, Fu B, Han D *et al.* Reduction, evolutionary pattern and positive selection of genes encoding formate dehydrogenase in Wood–Ljungdahl pathway of gastrointestinal acetogens suggests their adaptation to formate-rich habitats. *Environ Microbiol Rep* 2023:129–41.

31. Jiang L, Zheng Y, Peng X *et al.* Vertical distribution and diversity of sulfate-reducing prokaryotes in the Pearl River estuarine sediments, Southern China. *FEMS Microbiol Ecol* 2009;**70**:249–62.

32. Gagen EJ, Denman SE, Padmanabha J *et al.* Functional gene analysis suggests different acetogen populations in the bovine rumen and tammar wallaby forestomach. *Appl Environ Microbiol* 2010;**76**:7785–95.

33. Leloup J, Loy A, Knab NJ *et al.* Diversity and abundance of sulfate-reducing microorganisms in the sulfate and methane zones of a marine sediment, Black Sea. *Environ Microbiol* 2007;**9**:131–42.

34. Steinberg LM, Regan JM. Phylogenetic comparison of the methanogenic communities from an acidic, oligotrophic fen and an anaerobic digester treating municipal wastewater sludge. *Appl Environ Microbiol* 2008;**74**:6663–71.

35. Andrews S. FastQC: a quality control tool for high throughput sequence data. 2010.

36. Krueger F, James F, Ewels P *et al.* Trim Galore. 2021, DOI: 10.5281/zenodo.5127899.

37. Dobin A, Davis CA, Schlesinger F *et al.* STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* 2013;**29**:15–21.

38. Beghini F, McIver LJ, Blanco-Míguez A *et al.* Integrating taxonomic, functional, and strain-level profiling of diverse microbial communities with bioBakery 3. *Elife* 2021;**10**:1–42.

39. Greenacre M. Compositional data analysis. *Annu Rev Stat Its Appl* 2021;**8**:271–99.

40. Quinn TP, Erb I, Gloor G *et al.* A field guide for the compositional analysis of any-omics data. *Gigascience* 2019;**8**:1–14.

41. Ameijeiras-Alonso J, Crujeiras RM, Rodríguez-Casal A. Multimode: An r package for mode assessment. *J Stat Softw* 2021;**97**:1–32.

42. Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest Package: Tests in Linear Mixed Effects Models. *J Stat Softw* 2017;**82**:1–26.

43. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing. 2021.

44. Watthanacheewakul L. Transformations for Left Skewed Data. *Lect Notes Eng Comput Sci* 2021;**2242**:101–6.

45. Shannon P, Markiel A, Ozier O *et al.* Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks. *Genome Res* 2003;**13**:2498–504.

46. Faust K, Raes J. CoNet app: Inference of biological association networks using Cytoscape . *F1000Research* 2016;**5**:1–17.

47. Lahti L, Salojärvi J, Salonen A *et al.* Tipping elements in the human intestinal ecosystem. *Nat Commun* 2014;**5**:1–10.

48. Hackstein JHP, Van Alen TA. Fecal methanogens and vertebrate evolution. *Evolution (N Y)* 1996;**50**:559–72.

49. Samuel BS, Gordon JI. A humanized gnotobiotic mouse model of host-archaeal-bacterial mutualism. *Proc Natl Acad Sci U S A* 2006;**103**:10011–6.

50. Culp EJ, Goodman AL. Cross-feeding in the gut microbiome: Ecology and mechanisms. *Cell Host Microbe* 2023;**31**:485–99.

51. Hillesland KL, Stahl DA. Rapid evolution of stability and productivity at the origin of a microbial mutualism. *Proc Natl Acad Sci U S A* 2010;**107**:2124–9.

52. Plugge CM, Zhang W, Scholten JCM *et al.* Metabolic flexibility of sulfate-reducing bacteria. *Front Microbiol* 2011;**2**:1–8.

53. Noguera DR, Brusseau GA, Rittmann BE *et al.* A unified model describing the role of hydrogen in the growth of Desulfovibrio vulgaris under different environmental conditions. *Biotechnol Bioeng* 1998;**59**:732–46.

54. Schuchmann K, Müller V. Energetics and application of heterotrophy in acetogenic bacteria. *Appl Environ Microbiol* 2016;**82**:4056–69.

55. Liu C, Li J, Zhang Y *et al.* Influence of glucose fermentation on CO2 assimilation to acetate in homoacetogen Blautia coccoides GA-1. *J Ind Microbiol Biotechnol* 2015;**42**:1217–24.

56. Angenent LT, Karim K, Al-Dahhan MH *et al.* Production of bioenergy and biochemicals from industrial and agricultural wastewater. *Trends Biotechnol* 2004;**22**:477–85.

57. Boets E, Gomand S V., Deroover L *et al.* Systemic availability and metabolism of colonic-derived short-chain fatty acids in healthy subjects: a stable isotope study. *J Physiol* 2017;**595**:541–55.

58. Louis P, Flint HJ. Formation of propionate and butyrate by the human colonic microbiota. *Environ Microbiol* 2017;**19**:29–41.

59. Frolova MS, Suvorova IA, Iablokov SN *et al.* Genomic reconstruction of short-chain fatty acid production by the human gut microbiota. *Front Mol Biosci* 2022;**9**:1–25.

60. Frolova MS, Suvorova IA, Iablokov SN *et al.* Genomic reconstruction of short-chain fatty acid production by the human gut microbiota. *Front Mol Biosci* 2022;**9**:1–25.

61. Miceli JF, Torres CI, Krajmalnik-Brown R. Shifting the balance of fermentation products between hydrogen and volatile fatty acids: microbial community structure and function. Stams A (ed.). *FEMS Microbiol Ecol* 2016;**92**:fiw195.

62. Bergman EN. Energy contributions of volatile fatty acids from the gastrointestinal tract in various species. *Physiol Rev* 1990;**70**:567–90.

63. Hoegenauer C, Hammer HF, Mahnert A *et al.* Methanogenic archaea in the human gastrointestinal tract. *Nat Rev Gastroenterol Hepatol* 2022;**19**:805–13.

64. Samuel BS, Hansen EE, Manchester JK *et al.* Genomic and metabolic adaptations of Methanobrevibacter smithii to the human gut. *Proc Natl Acad Sci U S A* 2007;**104**:10643–8.

65. Catlett JL, Carr S, Cashman M *et al.* Metabolic Synergy between Human Symbionts Bacteroides and Methanobrevibacter. Claesen J (ed.). *Microbiol Spectr* 2022;**10**, DOI: 10.1128/spectrum.01067-22.

66. Wang Y, Lapointe G. Arabinogalactan utilization by bifidobacterium longum subsp. Longum ncc 2705 and bacteroides caccae atcc 43185 in monoculture and coculture. *Microorganisms* 2020;**8**:1–16.

67. Fenner L, Roux V, Mallet MN *et al.* Bacteroides massiliensis sp. nov., isolated from blood culture of a newborn. *Int J Syst Evol Microbiol* 2005;**55**:1335–7.

68. Milani C, Andrea Lugli G, Duranti S *et al.* Bifidobacteria exhibit social behavior through carbohydrate resource sharing in the gut. *Sci Rep* 2015;**5**:1–14.

69. Louis P, Solvang M, Duncan SH *et al.* Dietary fibre complexity and its influence on functional groups of the human gut microbiota. *Proc Nutr Soc* 2021;**80**:386–97.

70. Alessi AM, Gray V, Farquharson FM *et al.* β-Glucan is a major growth substrate for human gut bacteria related to Coprococcus eutactus. *Environ Microbiol* 2020;**22**:2150–64.

71. Rautio M, Eerola E, Väisänen-Tunkelrott ML *et al.* Reclassification of Bacteroides putredinis (Weinberg et al., 1937) in a new genus Alistipes gen. nov., as Alistipes putredinis comb. nov., and description of Alistipes finegoldii sp. nov., from human sources. *Syst Appl Microbiol* 2003;**26**:182–8.

72. Kageyama A, Benno Y. Catenibacterium mitsuokai gen. nov., sp. nov., a gram-positive anaerobic bacterium isolated from human faeces. *Int J Syst Evol Microbiol* 2000;**50**:1595–9.

73. Moore WEC, Johnson JL, Holdeman L V. Emendation of Bacteroidaceae and Butyrivibrio and descriptions of Desulfomonas gen. nov. and ten new species in the genera Desulfomonas, Butyrivibrio, Eubacterium, Clostridium, and Ruminococcus. *Int J Syst Bacteriol* 1976;**26**:238–52.

74. Kageyama A, Benno Y. Emendation of genus Collinsella and proposal of Collinsella stercoris sp. nov. and Collinsella intestinalis sp. nov. *Int J Syst Evol Microbiol* 2000;**50**:1767–74.

75. De Maesschalck C, Van Immerseel F, Eeckhaut V *et al.* Faecalicoccus acidiformans gen. nov., Sp. nov., Isolated from the chicken caecum, And reclassification of streptococcus pleomorphus (barnes et al. 1977), Eubacterium biforme (eggerth 1935) and eubacterium cylindroides (cato et al. 1974) as faecalicoccus p. *Int J Syst Evol Microbiol* 2014;**64**:3877–84.

76. Göker M, Gronow S, Zeytun A *et al.* Complete genome sequence of odoribacter splanchnicus type strain (1651/6 T). *Stand Genomic Sci* 2011;**4**:200–9.

77. Iyer R, Tomar SK, Uma Maheswari T *et al.* Streptococcus thermophilus strains: Multifunctional lactic acid bacteria. *Int Dairy J* 2010;**20**:133–41.

78. Watanabe Y, Nagai F, Morotomi M. Characterization of Phascolarctobacterium succinatutens sp. Nov., an asaccharolytic, succinate-utilizing bacterium isolated from human feces. *Appl Environ Microbiol* 2012;**78**:511–8.

79. Verlag F, Song Y, Liu C *et al.* Clostridium bolteae sp. nov., isolated from human sources. *Syst Appl Microbiol* 2003;**26**:84–9.

80. Sakamoto M, Iino T, Yuki M *et al.* Lawsonibacter asaccharolyticus gen. nov., sp. nov., a butyrateproducing bacterium isolated from human faeces. *Int J Syst Evol Microbiol* 2018;**68**:2074–81.

81. Crost EH, Le Gall G, Laverde-Gomez JA *et al.* Mechanistic insights into the cross-feeding of Ruminococcus gnavus and Ruminococcus bromii on host and dietary carbohydrates. *Front Microbiol* 2018;**9**:1–13.

82. Harris SC, Devendran S, Méndez- García C *et al.* Bile acid oxidation by Eggerthella lenta strains C592 and DSM 2243 T. *Gut Microbes* 2018;**9**:523–39.

83. Ungerfeld EM. Metabolic Hydrogen Flows in Rumen Fermentation: Principles and

Possibilities of Interventions. *Front Microbiol* 2020;**11**, DOI: 10.3389/fmicb.2020.00589.