Precise calcium-to-spike inference using biophysical generative models

Gerard Joey Broussard[1], Giovanni Diana[2], Francisco J. Urra Quiroz[2], B. Semihcan Sermet[2], Nelson Rebola[3], Laura A. Lynch[1], David A. DiGregorio[2,3,] *, Samuel S.-H. Wang[1, *]

[1]Neuroscience Institute, Washington Road, Princeton University, Princeton, New Jersey USA 08540.
[2]Institut Pasteur, Université Paris Cité, CNRS UMR 3571, Synapse and Circuit Dynamics Laboratory, Paris, France.
[3]L'Institut du Cerveau et de la Moelle épinière, CNRS UMR 7225, Paris, France
[4]Department of Physiology and Biophysics, University of Colorado School of Medicine, Aurora, Colorado USA 80045.

*co-senior authors

Correspondence: sswang@princeton.edu; david.digregorio@cuasnchutz.edu

Acknowledgements: The authors thank Marissa Applegate, Kiri Couchman, and Minglu Wang for collaboration and technical assistance in early stages of this project, and Yan Zhang and Loren Looger for discussion and for providing a jGCaMP8f construct prior to publication. S.W. was supported by NIH R01 NS045193 and U19 NS104648. S.W. and D.D. were supported by NIH R21 EY026434. G.J.B. was supported by the Warren Alpert Foundation and the Brain & Behavior Research Foundation. The laboratory of D.D. was also supported by the Fondation pour la Recherche Médicale (FRM EQU202003010555), Fondation pour l'Audition (FPA-RD-2018-8), and the Agence Nationale de la Recherche (ANR-17-CE16-0019, and ANR-18-CE16-0018, ANR-19-CE16 0019-02, ANR-21-CE16-0036-01).

Abstract:

The intramolecular dynamics of fluorescent indicators of neural activity can distort the accurate estimate of action potential ("spike") times. In order to develop a more accurate spike inference algorithm we characterized the kinetic responses to calcium of three popular indicator proteins, GCaMP6f, jGCaMP7f, and jGCaMP8f, using in vitro stopped-flow and brain slice recordings. jGCaMP8f showed a use-dependent slowing of fluorescence responses that caused existing inference methods to generate numerous false positives. From these data we developed a multistate model of GCaMP and used it to create Bayesian Sequential Monte Carlo (Biophys$_{SMC}$) and machine learning (Biophys$_{ML}$) inference methods that reduced false positives substantially. This biophysical method dramatically improved spike time accuracy, detecting individual spikes with a median uncertainty of 4 milliseconds, a performance level that reached the theoretical limit and is twice as accurate as any previous method. Our framework thus highlights advantages of physical model-based approaches over model-free algorithms.

Introduction

Genetically encoded calcium indicators (GECIs), as exemplified by the GCaMP family of sensors (Tian et al., 2009; Akerboom et al., 2012; Chen et al., 2013; Dana et al., 2019; Zhang et al., 2023), have transformed the study of neural populations by enabling simultaneous recordings from large ensembles of neurons (Demas et al., 2021). These sensors convert rapid changes in intracellular calcium concentration, which accompany neuronal action potentials, into measurable fluorescence signals. However, the temporal resolution of GECIs is limited by their slower kinetics compared to the rapid voltage changes associated with neuronal spikes. This mismatch limits the temporal precision with which the neural dynamics underlying behavior can be accurately estimated (Xiao et al., 2017; Li et al., 2019; Wei et al., 2020).

A variety of computational strategies have been developed to address the challenge of inferring spikes from fluorescence signals. These methods can be broadly categorized as either supervised or unsupervised. Supervised approaches leverage training data where both fluorescence and spike times are known to train machine learning models to recognize statistical regularities mapping observed signals to underlying spiking activity (Rupprecht et al., 2021; Zhou et al., 2023). Unsupervised approaches, by contrast, do not require labeled data and instead use principled assumptions about how spikes generate fluorescence based on the statistical properties of fluorescence time series to infer the underlying "latent" spike trains (e.g. Vogelstein et al., 2010; Pnevmatikakis et al., 2013; Deneux et al., 2016; Friedrich et al., 2017; Sebastian et al., 2017; Pachitariu et al., 2018). The performance of both categories is enhanced if the relationship between spike occurrences and the resultant fluorescence responses is fundamentally linear. When this linearity assumption holds, unsupervised methods can utilize simple, computationally efficient models (in particular linear deconvolution; Friedrich et al., 2017), while supervised methods benefit from stable, direct mappings that allow for robust training and generalization.

However, the spike-to-fluorescence relationship often deviates from the linear ideal. Classic examples include saturation of indicator fluorescence at high calcium concentrations (Regehr and Atluri, 1995) and slowing of calcium dynamics due to calcium buffering by high-affinity indicators (Sabatini and Regehr, 1998). Thirdly, GECI responses to free calcium changes can be supralinear (Rose et al., 2014). Such nonlinearities pose potential challenges to both unsupervised and supervised methods. Popular unsupervised methods relying on low-complexity models such as linear convolution or generative frameworks reliant upon a static decay process would be expected to exhibit systematic errors dictated by the nature of the nonlinearity. Supervised methods, meanwhile, may fail to capture the complexity of novel, nonlinear regimes that differ substantially from the conditions under which they were trained (Seroussi and Zeitouni, 2022; Sariyildiz et al., 2023). Consequently, nonlinearities in the spike-to-fluorescence transformation have the potential to degrade the accuracy and reliability of inferred spike statistics.

The recently developed jGCaMP8 sensors represent a promising advance due to their faster rise times and improved single-spike signal-to-noise ratios (SNR) (Zhang et al., 2023; Diana et al., 2024), but little is known about what new non-linearities they introduce. Past generation indicators have been reported to exhibit supralinear responses arising from closely occurring spikes (Chen et al., 2013; Éltes et al., 2019; Wei et al., 2020; Huang et al., 2021;

Schoenfeld et al., 2021). Neither models designed to capture this supralinearity (Deneux et al.,
2016; Wei et al., 2020) nor models that assume linear characteristics of the newer probes
(Zhang et al., 2023) are able to extract more accurate spiking dynamics using jGCaMP8 than
they do from previous-generation sensors.
To address these gaps, we first comprehensively characterized the dynamics of
GCaMP6f, jGCaMP7f, and jGCaMP8f responses. We identified novel nonlinear properties of
these responses, including a pronounced use-dependent slowing by jGCaMP8f during periods
of heightened activity. We developed a biophysical model of all three GCaMPs, which we used
as the core of two new spike inference approaches: a sequential Monte Carlo (SMC) method to
model latent spike-time sequences, and a machine learning-based strategy trained on synthetic
data generated by the SMC method. Both approaches identify spike times with higher accuracy
and temporal precision than any currently used technique.
95

Results

Biophysical characterization of GCaMPs
jGCaMP8 variants have been reported to possess improved kinetic properties and linear
responses to action potentials compared to earlier GCaMP versions (Zhang et al., 2023). To
precisely quantify these improvements, we expressed GCaMP6f, jGCaMP7f, or jGCaMP8f in
cerebellar granule cells using adeno-associated viral vectors. We performed high-speed (0.5-1
kHz) two-photon linescans to measure fluorescence transients in single boutons (parallel fibers,
PFs; **Figure 1A-C**) in response to extracellularly-evoked action potentials. jGCaMP8f exhibited
the fastest rise and decay times (**Figure 1C**), and a fractional per-spike fluorescence change of
0.59 ± 0.07 (n = 32 boutons, **Figure 1D**), larger than the other probes tested (GCaMP6f, 0.29 ±
0.04, n = 14 boutons; jGCaMP7f, 0.34 ± 0.05, n = 27; the small-molecule indicator Oregon
Green BAPTA-5N, OGB-5N, 0.39 ± 0.04, n = 11; mean ± SEM; ANOVA, p = 0.0041). jGCaMP8f
showed a signal-to-noise ratio of 5.6 ± 0.1 (**Figure 1E**), again higher than the other probes
(GCaMP6f, 1.4 ± 0.2; jGCaMP7f, 1.3 ± 0.2; jGCaMP8f, 5.6 ± 0.1; mean ± SEM; ANOVA, p =
6.7e-10).
The rise time of jGCaMP8f was 3.1 ± 0.2 ms, approaching that of OGB-5N (0.3 ± 0.1
ms), which has near diffusion-limited calcium on-binding kinetics (DiGregorio et al., 1999)
(**Figure 1F**) and was faster than all other probes (GCaMP6f, 18.1 ± 1.6 ms; jGCaMP7f, 13.7 ±
0.7 ms;; mean ± SEM; ANOVA, p = 6.7e-29; significant Tukey HSD post hoc comparisons: p =
1.1e-21, GCaMP6f vs. OGB5N; p = 5.4e-4, GCaMP6f vs. jGCaMP7f; p = 3.9e-23, GCaMP6f
vs. jGCaMP8f; p = 8.7e-18, jGCaMP7f vs. OGB-5N; p = 1.5e-19, jGCaMP7f vs. jGCaMP8f).
117

118 **Figure 1: Nonlinear response characteristics of fast GCaMPs in a controlled, ex**
119 **vivo setting.**

120
121 (**A**) Schematic of experimental approach showing sites of extracellular stimulation and imaged
122 regions (highlighted in yellow). (**B**) Representative traces obtained across sensors for single-
123 spike responses, with mean overlaid. (**C**) Traces from **B** averaged and normalized to compare
124 kinetic properties of GCaMPs with the low affinity synthetic indicator, OGB-5N. (**D-F**) Peak
125 response amplitude, signal-to-noise ratio, and 10-90% rise time for individual boutons. Each
126 symbol represents an average across trials in one bouton. (**G**) averaged responses to one
127 stimulus or two stimuli spaced 10 ms apart. (**H**) Relative amplitude of responses to a second
128 stimulus spaced at various time intervals from the first stimulus. Each point represents the

average response for one bouton. (**I**) Responses to 1 stimulus (left), 10 stimuli at 100 Hz
overlaid to the single stimulus response (center), and normalized to the amplitude of a single-
stimulus response to demonstrate prolonged decay (right). (**J**) Half-decay times as a function of
stimulus number. (**K-L**) Convolution of a single-stimulus response with the times of stimuli to
demonstrate effect of short-term amplitude non-linearity in GCaMP6f responses in **K** and
cumulative signal arising from many stimuli for jGCaMP8f in **L**. Stars indicate statistical
significance as: *p<0.05, **p<0.01, ***p<0.001.
136
Recent reports of GCaMP6 indicators suggest a nonlinear relationship between the
number of APs and the corresponding fluorescence response (Wei et al., 2020; Huang et al.,
2021; Zhang et al., 2023). To quantify the time-dependence of this nonlinearity across fast
indicator variants, we explored signals evoked by pairs of spikes delivered at defined time
intervals. GCaMP6f and GCaMP7f showed short-term supralinearity that decreased to a linear
summation for intervals greater than 100 ms (**Figure 1G, H**) (GCaMP6f, 3.1 ± 0.7 at 10 ms and
1.5 ± 0.3 at 100 ms fold increase, n = 7 boutons; jGCaMP7f, 1.9 ± 0.2 at 10 ms and 1.1 ± 0.1 at
100 ms, n = 14 boutons; mean ± SEM; two-way mixed ANOVA, $F(3,60)$ = 7.2, p = 3.3e-4 main
effect of indicator, with significant one-way ANOVA at 10 ms, F(3,32) = 6.4, p = 0.0016;
significant Tukey HSD post hoc comparisons: p = 0.01, GCaMP6f vs. OGB-5N). In contrast, the
jGCaMP8f responses were near-linear in their response amplitude to spikes at all tested
intervals (**Figure 1G, H**) (testing against a null distribution centered on 1; one-sample t-test, p =
0.32).
To examine responses to bursts of activity, we measured bouton fluorescence in
response to 1, 2, and 10 AP stimuli delivered at 100 Hz. GCaMP6f responses showed a strong
supralinearity, in which a 10-pulse train evoked signals approximately 20-fold larger than a
single pulse (**Figure 1G**, top), consistent with positive cooperativity. In contrast, jGCaMP8f
responses under the same comparison resulted in only a 3-fold larger response (**Figure 1F**,
bottom). These findings were broadly consistent with indicator saturation and low apparent
cooperativity (**Figure 1F**, bottom).
Unexpectedly, all three tested GCaMPs showed a slowing of the decay of fluorescence
transients driven by ten spikes compared with a single spike (**Figure 1I**) (GCaMP6f, 78 ± 5 ms 1
pulse vs. 130 ± 9 ms 10 pulse, n = 6 boutons; jGCaMP7f, 77 ± 5 ms 1 pulse vs. 126 ± 7 ms 10
pulse, n = 22 boutons; jGCaMP8f, 41 ± 3 ms 1 pulse vs. 123 ± 12 ms 10 pulse, n = 14 boutons;
mean ± SEM; two-way mixed ANOVA, $F(1,78)$ = 72.2, p = 1.0e-12 main effect of time). This
use-dependent effect was particularly pronounced for jGCaMP8f (~3-fold slowing) when
compared to the older fast sensor variants (~1.5-fold slowing) (**Figure 1J, K**). Further analysis
revealed that a two-component exponential was required to achieve acceptable fits to the decay
even for single-spike responses. Both kinetic components tended to increase in duration with
the number of spikes (**Supplementary Figure 1**).
To explore the impact of nonlinear properties of the old and new GCaMP variants, we
compared experimentally-measured spike responses to a linear model. The model was
calculated by convolving a kernel fitted to a single spike response with the spike train used to
generate the observed data. In the paired-spike protocol, the GCaMP6f kernel underestimated
the response to a 10 ms inter-spike interval. The estimate improved as the intervals lengthen
and the nonlinearity decreased (**Figure 1J**). When jGCaMP8f boutons were exposed to a 10 Hz

Poisson stimulus train, the kernel estimate diverged increasingly from the real data (**Figure 1K**).
Overall, these analyses reveal that the GCaMP6f amplitude nonlinearity is dependent on spike
timing, while the use-dependent slowing of jGCaMP8f signals is dependent on spike rate. These
findings indicate activity regimes where linear methods for deconvolution are expected to be
unsuccessful: short inter-spike intervals for older GCaMPs, and high spike rates for jGCaMP8f.
178
179   **Supplementary Figure 1: jGCaMP8f response decay is captured by two kinetic**
180   **components.**

181
(**A**) Representative single-bouton responses to 1, 2, and 10 stimuli overlaid and aligned by first
stimulus time. (**B**) Responses normalized and aligned to last spike. (**C**) Example fits to dual
sums of exponentials showing fast and slow components of decay. (**D**) Dependence of fast (top)
and slow (middle) time constants on the number of stimuli, and relative contribution of the slow
component (τ_2) to the overall time course (bottom). Individual bouton values shown as low
linewidth traces with group means shown as thicker line and SEM as error bars, n = 6 boutons
and 3 granule cell somata.
189

190 **Supplementary Figure 2: The impact of calcium clamp design on observed**
191 **stopped-flow dynamics.**

192
(**A-B**) Schematic of stopped-flow buffer systems using (**A**) a slow buffer (EGTA) and containing
unbuffered calcium in the calcium-containing bolus (**A**) and (**B**) a system with a fast buffer
(BAPTA) and fully buffered calcium in the calcium-containing bolus. (**C**) Simulations of the
EGTA-based system presented in (**A**) show uncontrolled free calcium concentrations in the first
few milliseconds of large calcium steps, but (**D**) BAPTA-based experiments do not. (**E**)
Measurements of OGB-1 responses (left) reflect uncontrolled responses that are matched by
simulations of the buffer system (right). (**F**) OGB-1 data and simulations in BAPTA-based
buffers. (**G**) Artifactual fast on-response to jGCaMP8f in EGTA-based buffer system. (**H**) True
jGCaMP8f response in BAPTA-based buffer system.
202

Supplementary Figure 3: Magnesium affects the properties of jGCaMP8f much more than older GCaMPs.

(**A**) Emission spectra of GCaMP variants in standard protein purification buffer (KMOPS, left) and under intracellular conditions that include magnesium (2 mM Mg^{2+}). (**B**) Fluorimeter counts at 490 nm under calcium-free (left) or saturating calcium (right) conditions. jGCaMP8f, but not other variants, showed strong magnesium-dependence of fluorescence emission under low-calcium conditions. No indicator showed this effect at saturating calcium. (**C**) Quantification of the weak dependence of GECI dynamic range R$_f$ (saturating calcium divided by calcium-free fluorescence output) on temperature (left) or ionic strength (center) compared to the effect seen with magnesium (right). (**D**) Dynamic range of GCaMP variants at mammalian near-physiological conditions. (**E**) Schematic of stopped-flow experiment to measure responses to rising steps of calcium concentration. (**F**) Half-rise times to calcium steps were accelerated most by magnesium for jGCaMP8f, less so for GCaMP6f, and not at all for jGCaMP7f.

Figure 2: Stopped-flow fluorimetry of GCaMP dynamic responses.

(**A**) Schematic of stopped-flow experiment and rising steps of calcium concentration. (**B**) Step-down protocols for decreasing calcium. (**C**) Responses to steps up from 0 μM free calcium. (**D**)

Responses to steps down from 10 µM free calcium. (**E**) Half-rise times for GCaMP6f,
jGCaMP7f, and jGCaMP8f as a function of calcium step size. Shaded gray region indicates the
physiological range of intracellular signals. (**F**) Comparison of brain slice presynaptic bouton
signals with stopped-flow rise times for the three indicator variants. (**G**) Rising phase responses
on logarithmic time scales. Symbols indicate the time points plotted in (**H**). (**H**) Hill plots of rising
and steady-state responses to steps in calcium. Colored numbers indicate Hill coefficients. (**I**)
Comparison of brain slice amplitude nonlinearity at 10 ms inter-stimulus response with respect
to Hill coefficients. (**J**) Representative stopped-flow traces with y-axis values presented on a log
scale for jGCaMP8f (left) and GCaMP6f (right). Sums of exponentials for each indicator
demonstrate the biphasic response and the greater contrast in speed of kinetic components
seen in jGCaMP8f. (**K**) Quantification of the factor of slowing seen in GCaMP fluorescence
decay in response to 1 and 10 stimuli in brain slices. (**L**) Comparison of brain slice slowing to
accessible slowing (see main text and methods). Error bars in all panels represent the
interquartile range of the distribution.
236
To gain insight into the mechanistic basis for GCaMP's use-dependent nonlinearities, we
measured GCaMP fluorescence responses to various calcium concentration steps directly using
stopped-flow fluorimetry on purified protein (**Figure 2A, B)**. We first optimized stopped-flow
buffer conditions to closely match intracellular ionic composition, including magnesium, which
we found to be necessary to reproduce the kinetics and dynamic range of the GCaMP
responses observed in living cells (**Supplementary Figure 2, 3**). The rise and decay kinetics of
jGCaMP8f were faster than 6f and 7f for all calcium concentration steps in the range of 0-10 µM
(**Figure 2C, D**). Within the range of calcium increase expected in response to a small number of
spikes (shaded region, **Figure 2E**), the rise time of spike-evoked calcium transients for each
GECI variant correlated with stopped-flow rise times (**Figure 2E, F**). Similarly, the paired-pulse
facilitation of a GCaMP's fluorescent response in neurons closely tracked cooperativity
measures derived from analysis of the early stopped-flow response (**Figure 2G-I**). These
experiments show that our stopped-flow measurements capture key features of action potential-
evoked calcium responses in neurons.
Previous studies showed that activation of GCaMPs by calcium occurs via two
pathways, one fast and one slow (Sun et al., 2013; Badura et al., 2014). Closer inspection of the
binding and unbinding kinetics across GCaMP sensors revealed fast and slow phases of
fluorescence increases. Responses of GCaMP6f and jGCaMP7f arose predominantly from a
slow component whose speed increased with the size of the calcium step. In contrast,
jGCaMP8f's fluorescence developed predominantly through a fast component. While both
kinetic components contributed to fluorescence changes in all GECIs, jGCaMP8f was
distinguished by having a faster and larger contribution from the fast component at all
concentrations tested (**Figure 2J**, **Supplementary Figure 4**).
The presence of two time constants in the rising phase of the fluorescence response is
consistent with entrance into at least two distinct fluorescence states. The presence of multiple
fluorescence states could account for fast and slow decay rates. We compared the time
constants governing this fast and slow increase to the use-dependent slowing effect observed in
our ex vivo (i.e. brain slice) recordings. We found a linear relationship between our ex vivo
observations and a metric we term accessible slowing which represents a tradeoff between the

speed of entrance into the fast versus slow state and stability of the slow state (**Figure 2K, L**).
These observations support the conclusion that large calcium increases push GCaMPs into a
stable fluorescent state from which they are slow to exit.
269
270 **Supplementary Figure 4:** Biphasic responses of GCaMPs to increasing calcium steps.

271
(**A**) Schematic of stopped-flow experiment and rising calcium-step protocol. (**B**) Example
analysis of fit to a sum of two exponentials. (**C**) Monoexponential rising kinetics for Oregon
Green BAPTA-1 (OGB-1) and biexponential kinetics for GCaMP variants. (**D**) Dependence of
fast and slow components on amplitude of calcium step. Upper plots show rate constants
represented in the response at the indicated step. In the lower plots, vertical bars show relative
amplitude of fast (dark) and slow (light) components. (**E**) Schematic of decreasing-calcium step

protocol. (**F**) Fluorescence responses to decreasing calcium steps. (**G**) Breakdown of kinetic
components contributing to the decay of fluorescence as in (**D**). Due to the low number of steps
considered, analysis for all variants is included in this single plot. (**H**) Monoexponential decay for
OGB-1 and more complex responses for GCaMP variants.
282
283

 Supplementary Figure 5: Selection of GCaMP biophysical parameters using in
 vitro and ex vivo datasets.

(**A**) Distributions of parameters on fits from randomized seeds with GCaMP model parameters
at figure left and residuals from best fits to stopped-flow and ex vivo experiments (paired pulse,

1-2-10 pulse, and Poisson train) based on optimizing cell parameters. GCaMP parameter sets

were sorted by hierarchical clustering and then by lowest summed residuals within each cluster.

This procedure allowed qualitative assessment of parameters that lay in different local minima

of the parameter space. (**B**) Fitted data examples for cluster 1, which was used for all further

modeling. (**C**) Fitted data for cluster 5. (**D**) Fitted data for cluster 9. In paired-pulse and Poisson

train data, vertical lines indicate the times of action potentials. Expanded timescale of Poisson

responses at right. Neither cluster 5 or 9 captured the differential slowing across the Poisson

responses.

297 **Figure 3: A biophysical model for GCaMP response.**

298
299 (**A**) Schematic GCaMP model depicting probe components, including circularly permuted green
300 fluorescent protein (cp-GFP), the N- and C-lobes of the calcium binding-calmodulin, and the
301 binding peptide (BP). Black arrows indicate key pathways to fluorescence with states
302 highlighted for their role in model function. (**B**) Schematic diagram of physiological response of a
303 cellular structure containing endogenous calcium buffer, GCaMP, and intracellular and plasma
304 membrane calcium extrusion (pump) mechanisms. (**C**) Fits of the GCaMP model to the three

GCaMP variants with different parameters for each variant. (**D**) Fits of the cell model to
physiological data with fixed GCaMP model parameters. Vertical lines indicate action potential
(spike) times. (**E**) Occupancy of the fast-fluorescent, slow-fluorescent, and slow non-fluorescent
states of GCaMP indicating a mechanism for use-dependent slowing of fluorescence decay,
becoming most apparent after 10 action potentials. (**F**) Predictions on simulated ground-truth
data for an excitatory neuron. The data consists of a single spike (black) or a 10 Hz Poisson
train of spikes (gray) (top panel). Lower panels show the average posterior spike probability
(solid lines) and spikes times from a single trajectory (triangles) derived by our SMC method
employing either the biophysical model (Biophys$_{SMC}$, green) or a linear model (Linear$_{SMC}$, pink)
as its generative engine. Vertical lines indicate action potential (spike) times. (**G**) As in (**F**) with
simulated ground truth for an inhibitory cell (30 Hz Poisson spike rate, calcium increase per
spike reduced by 2, dynamic range reduced by a factor of 2).
317

Development of a biophysical model of GCaMP
We built a biophysical model of GCaMP that included parallel fast and slow kinetic
pathways to multiple fluorescence states. Multiple kinetic pathways are known to coexist in
calmodulin, the calcium-sensing domain of GCaMP, which semi-independently binds calcium
via its N-lobe (fast-binding) and C-lobe (slow-binding) (Linse et al., 1991; Faas et al., 2011),
each subsequently capable of interacting with a target peptide to induce fluorescence (Johnson
et al., 1996; Brown et al., 1997; Sun et al., 2013; Lai et al., 2015). We reasoned that such a
model structure might phenomenologically account for both the *in vitro* and *ex vivo* observations
(**Figure 3A**). In order to estimate the kinetic rate constants of the GCaMP model we fit it to
experimental data. We took advantage of a neuron type (cerebellar granule cells) with well-
characterized calcium buffering and handling (Rebola et al., 2019). The cell model used here
included endogenous buffering, calcium extrusion mechanisms, and calcium exchange from
intracellular stores (**Figure 3B**) and could reproduce the fast dynamics of previous buffer
models (Rebola et al., 2019), but we also added a second compartment to account for a slow
decay that was not used to constrain previous models. By jointly fitting the stopped-flow data
and *ex vivo* responses, we derived the final GCaMP model parameters for each variant, which
were then fixed for all subsequent analyses (**Supplementary Figure 5; Supplementary Figure
6A, B**). This approach allowed us to reproduce observed dynamics from all experimental
systems and GCaMP variants, despite the idiosyncrasies observed across the sensors (**Figure
3C, D; Supplementary Figure 6C-F**).
Further analyses of the GCaMP model states (**Figure 3A; Supplementary Figure 6A**)
revealed additional insights into how parameterization led to different operational modes. When
separating the fluorescent GCaMP model states into those that depended on C-lobe binding
(slow fluorescent) and those that did not (fast fluorescent), we observed that jGCaMP8f alone
showed stable occupancy of the fast fluorescent states in both stopped-flow and ex vivo fits
(yellow traces **Supplementary Figure 6C-F**). The slowing of jGCaMP8f responses following
spike bursts, in contrast, was driven by increased occupancy of a slow, fluorescent and non-
fluorescent state (**Figure 3E**, blue and gray, respectively).
346

Supplementary Figure 6: A full biophysical state analysis of GCaMP response.

(**A**) Full biophysical state model of GCaMP response. Specific states referred to as "fast fluorescent", "slow fluorescent", and "slow, non-fluorescent" have centers color-coded as in model state traces below. (**B**) Best-fit parameter values for GCaMP6f, jGCaMP7f, and jGCaMP8f. (**C**) Model fits to 0.5 µM increasing step of calcium (top), with model state broken

down by occupancy. Note that jGCaMP8f is distinguished by its near-exclusive use of the fast
fluorescent states. (**D**) As in (**C**) for decreasing step. jGCaMP8f is distinguished by fast,
persistent entry into the slow, non-fluorescent state. (**E**) Model fits to fluorescence responses in
presynaptic boutons of granule cells. (**F**) State occupancy for the fluorescence responses
shown in (**E**). Fast responses arise in part from high occupancy of a slow, non-fluorescent state
at basal calcium for jGCaMP8f which efficiently transitions to fluorescence on elevation of
calcium.
361
362   **Supplementary Figure 7: Generative cell-based model of fluorescence time series**

363
364   Graphical representation of the generative SMC model described in the main text. White circles
365   denote unknown variables, gray circles denote measurements and bare variables are fixed prior
366   hyperparameters. Plates denote groups of variables. Note that GCaMP model parameters are
367   included, but fixed for the SMC approach.

Figure 4: Improved action potential inference from generative model-based methods.

(**A**) A generative model for imputing action potential times based on iterative fitting. The biophysical GCaMP model was held fixed while cell parameters were varied. Responses to action potential sequences were generated to fit cell parameters (Biophys$_{SMC}$). After fitting was complete, the Biophys$_{SMC}$ model was then used to train a machine learning based model. (**B, C**)

375 Imputed spike probabilities using a variety of unsupervised and supervised models. Values of r

376 indicate Pearson correlation between the time course of ground truth and that of the imputed

377 probabilities for data filtered at 20 ms. Vertical lines indicate true spike times and tick marks

378 indicate imputed spikes. (**B**) shows data from an excitatory cell while (**C**) shows data from an

379 inhibitory cell. (**D**) Pearson correlations for different filtering time constants. (**E**) Discrete spike

380 estimate statistics. Left panels shows the average time between the nearest imputed and

381 ground truth spikes while the right shows the normalized number of spikes that were over or

382 underestimated. Individual points are calculated over 8-second epochs covering 1000 data

383 points randomly selected from the full dataset.(**E**) Timing and detection accuracy of the different

384 imputation models. Left, distribution of per-spike timing error for the various methods. Smoothed

385 histograms and box/whisker plots summarize all errors. Right, epoch-wise recall of the various

386 methods indicating the percent of ground truth spikes recovered.

388 **Model-based insights enable improved methods of spike detection**

389   We next leveraged the GCaMP model to implement a generative spike inference

390 algorithm. Inspired by recent successes of a Bayesian autoregressive-model based approach

391 (Diana et al. 2024), we adopted the same Sequential Monte Carlo (SMC) framework in which

392 time-invariant parameters (such as those governing cell calcium handling) and latent states

393 (such as spike times) are jointly estimated in an iterative manner to estimate spike times

394 (**Supplementary Figure 7**).

395   As an initial test of this approach, we generated ground truth simulations using the

396 GCaMP model and attempted inference using our SMC framework with either the GCaMP

397 model (Biophys$_{SMC}$ hereafter) or a linear autoregressive model (Linear$_{SMC}$) as the generative

398 model (**Figure 3F, G**).Using this approach, Linear$_{SMC}$ produced a progressive increase in the

399 number of false spike predictions during periods of elevated spiking for both cell types,

400 particularly in the decay phase of the fluorescence. This is likely due to the use-dependent

401 slowing of jGCaMP8f fluorescence. In contrast, Biophys$_{SMC}$, which accommodated all

402 nonlinearities, made consistent, accurate predictions of spike times across the simulation

403 epoch. The Biophys$_{SMC}$ was able to reproduce the input spike train with higher accuracy and

404 precision in the context of simulations designed to mimic different cell types, including both

405 excitatory and inhibitory cells.

406   The Biophys$_{SMC}$ approach to spike inference is computationally time-consuming due to

407 the SMC sampling. Therefore we also used the GCaMP model to create a synthetic training

408 dataset to train a machine learning algorithm (Rupprecht 2021). Specifically, we generated

409 synthetic fluorescence data for training using cell parameters derived from the Biophys$_{SMC}$ along

410 with randomly generated spikes, noise, and baseline drift designed to mimic the statistics of the

411 dataset targeted for inference (**Figure 4A**). In this way, we obtained two methods for spike

412 inference: an unsupervised, generative Bayesian approach (Biophys$_{SMC}$), and a computationally

413 faster machine-learning approach (Biophys$_{ML}$) which was generated by using the outputs of the

414 unsupervised approach to train a supervised-learning neural network.

415   We benchmarked our spike inference methods against state-of-the-art supervised and

416 unsupervised approaches for spike inference, using excitatory and inhibitory cell responses

417 from the jGCaMP8f ground truth dataset (Zhang et al., 2023). We used 154 18-second epochs

418 of firing in 37 excitatory neurons and 3 inhibitory neurons, containing 8,161 ground truth action

potentials, with a median of 34 (IQR, 16-69) spikes per epoch. For a generative method
comparison, we used MLspike, an established algorithm based on an alternative generative
model of calcium-fluorescence dynamics (Deneux et al., 2016). For supervised-method
comparisons, we used the CASCADE (Rupprecht et al., 2021) and ENS[2] (Zhou et al., 2023)
global-trained networks, which have been reported to outperform other methods on a variety of
benchmark tests.
We first examined the ability of our methods to estimate continuous changes in activity.
Imputed spike predictions and ground truth spikes were smoothed with Gaussian filters of
various widths. $Biophys_{SMC}$ and $Biophys_{ML}$ model predictions were tightly locked to true spike
patterns for both excitatory and inhibitory classes (**Figure 4B, C**), as indicated by the higher
correlations between our method predictions and ground truth at all filter widths, particularly at
ten and twenty milliseconds. Results with CASCADE were not improved by training the network
on the jGCaMP8f dataset itself (**Supplementary Figure 8**).
We quantified the ability of various methods to track changes in firing using the Pearson
correlation between continuous spike probability, filtered on various timescales, and the
smoothed time course of ground truth (**Figure 4D**). For excitatory neurons (n = 142 epochs), a
mixed effects model shows that correlation depends on the method used (method and filter
width as fixed effects, $F_{(4,4935)}$ = 171.1, p = 2e-137). Method-wise comparisons demonstrate
that $Biophys_{SMC}$ outperforms MLspike (p = 6.5e-62, d = 1.1), CASCADE (p = 9.0e-47, d = 0.8),
and ENS[2] (p = 2.3e-82, d = 1.0). Likewise, $Biophys_{ML}$ outperformed MLspike (p = 3.9e-56, d =
1.1), CASCADE (p = 9.8e-42, d = 0.3), and ENS[2] (p = 9.4e-76, d = 0.6). $Biophys_{ML}$
outperformed CASCADE at filter widths up to 30 milliseconds, ENS[2] up to 76 milliseconds, and
MLspike up to 100 milliseconds, the widest filter tested.
For the inhibitory cell dataset (n = 12 epochs), a similarly-structured mixed effect model
again showed that correlation depends on method ($F_{(4,385)}$ = 14.7, p = 3.6e-11) with improved
performance with $Biphys_{SMC}$ (vs. MLspike p = 4.1e-9, d = 1.9; vs. CASCADE p = 3.3e-4, d = 1.0;
vs. ENS[2] p = 0.005, d = 1.1) and $Biophys_{ML}$ (vs. MLspike p = 1.4e-10, d = 2.0; vs. CASCADE p
= 3.3e-5, d = 1.2; vs. ENS[2] p = 8.2e-4, d = 1.4). Together, these comparisons demonstrate the
superior ability of $Biophys_{SMC}$ and BiophysML to estimate continuous spike probability on a
variety of timescales and in different cell types.
Next, we calculated how well the different methods were able to capture discrete spike
times. First, we matched imputed spikes to the nearest ground-truth spike within a 10 ms
interval to avoid double-counting of spikes and penalize imprecise imputed spikes
(**Supplementary Figure 9**). Our biophysical methods consistently found ground truth events at
a higher rate than other methods as assessed by sensitivity (also called recall, defined as true
positives as a fraction of all ground truth events) (**Figure 4E** left, **Table 1**; repeated measures
ANOVA, $Biophys_{SMC}$ vs. MLspike p = 4.0e-9, d = 1.1; vs. CASCADE p = 4.0e-9, d = 0.6; vs.
ENS[2] p = 4.0e-9, d = 1.0; $Biophys_{ML}$ vs. MLspike p = 4.0e-9, d = 1.1; vs. CASCADE p = 4.0e-9,
d = 0.7; vs. ENS[2] p = 4.0e-9, d = 1.0).
We further tested the methods using a standard measure of predictive performance, the
F-score. The F-score is the harmonic mean of precision (true positives as a fraction of all
imputed events) and recall and has a maximum value of 1 thus allowing assessment of
performance considering both false negatives as well as false positives. Taking this broader
assessment into account, we found that the biophysical methods again outperform the other

methods (**Table 1**; repeated measures ANOVA, Biophys$_{SMC}$ vs. MLspike p = 3.8e-9, d = 1.1; vs.
CASCADE p = 3.8e-9, d = 0.8; vs. ENS^2 p = 3.8e-9, d = 0.9; Biophys$_{ML}$ vs. MLspike p = 3.8e-9,
d = 0.9; vs. CASCADE p = 3.8e-9, d = 0.6; vs. ENS^2 p = 3.8e-9, d = 0.6). Thus, our
biophysically-based methods provide unprecedented temporal accuracy and precision
compared to current state-of-the-art approaches to spike predictions based on jGCaMP8f
imaging datasets.
Finally, we tested the ability of our methods to estimate individual spike times. We first
calculated the time between imputed spikes and the nearest matched ground-truth spike in a
100 ms interval to assess bias and uncertainty in spike time predictions (**Figure 4E**, right,
Supplementary Figure 10). On an epoch-by-epoch basis, both of the supervised methods
predicted spike times that tended to precede ground-truth times (CASCADE, 7.45 ms; ENS^2,
7.67 ms); MLspike showed a smaller bias (4.93 ms). In contrast, both biophysical methods
removed this bias (**Table 1**; repeated measures ANOVA, Biophys$_{SMC}$ vs. MLspike p = 6.0e-8, d
= 0.5; vs. CASCADE p = 6.0e-8, d = 0.8; vs. ENS^2 p = 6.0e-8, d = 0.6; Biophys$_{ML}$ vs. MLspike p
= 1.8e-6, d = 0.4; vs. CASCADE p = 6.0e-8, d = 0.9; vs. ENS^2 p = 1.8e-7, d = 0.5).
The biophysical methods also reported spike times with lower uncertainty (Biophys$_{SMC}$
median 3.8, IQR 1.8-7.3 ms; Biophys$_{ML}$ median 5.8, IQR 2.7-10.3 ms) than the other supervised
methods (CASCADE median 8.2, IQR 3.5-18.2 ms; ENS^2 median 8.3, IQR 3.8-16.6 ms) and
superior epoch-wise statistics than all other methods (repeated measures ANOVA, Biophys$_{SMC}$
vs. MLspike p = 0.01, d = 0.3; vs. CASCADE p = 6.0e-8, d = 0.5; vs. ENS^2 p = 1.2e-7, d = 0.5;
Biophys$_{ML}$ vs. MLspike p = 0.4, d = 0.1; vs. CASCADE p = 6.0e-8, d = 0.5; vs. ENS^2 p = 1.5e-5,
d = 0.4). In summary, the generative biophysical model-based methods outperformed existing
in-class and out-of-class alternatives, exceeding previous limits in the accuracy of GCaMP-
based spike decoding.
487
488

Table 1. Single-spike classification metrics for calcium-based decoding methods.
Median performance across epochs for a database of 37 excitatory cells with ground-truth
electrical recordings and jGCaMP8f fluorescence signals. Values reported as median (IQR).

	Precision	Recall	F-Score	Accuracy	Mean bias (ms)	Uncertainty (ms)
Biophys$_{SMC}$	0.84 (0.75, 0.92)	0.88 (0.71, 0.95)	0.83 (0.71, 0.91)	0.71 (0.55, 0.83)	1.9 (-2.9, 3.0)	3.8 (1.8, 7.3)
MLspike	0.70 (0.53, 0.93)	0.56 (0.43, 0.68)	0.60 (0.51, 0.70)	0.43 (0.34, 0.54)	-2.0 (-0.6, 0.9)	3.2 (1.5, 6.2)
Biophys$_{ML}$	0.76 (0.68, 0.84)	0.78 (0.68, 0.89)	0.75 (0.70, 0.80)	0.61 (0.54, 0.67)	0.0 (-2.9, 3.0)	5.8 (2.7, 10.3)
CASCADE	0.69 (0.63, 0.78)	0.70 (0.60, 0.79)	0.69 (0.61, 0.74)	0.52 (0.45, 0.59)	-7.4 (-11.2, -3.2)	8.2 (3.5, 18.2)
ENS^2	0.71 (0.65, 0.86)	0.57 (0.46, 0.69)	0.65 (0.56, 0.74)	0.48 (0.39, 0.58)	-6.6 (-12.3, -2.3)	8.3 (3.8, 16.5)

**Supplementary Figure 8: Training on jGCaMP8f data does not improve CASCADE
performance**.

(**A**) Imputed spike probabilities using the CASCADE Global Exc model or one trained on the
jGCaMP8f dataset itself. Values of r indicate Pearson correlation between the time course of
ground truth and that of the imputed probabilities for data filtered at 20 ms. Vertical lines indicate
true spike times and tick marks indicate imputed spikes. (**B**) Pearson correlations for different
filtering time constants.

Supplementary Figure 9: Statistics of discrete spike estimate comparisons to

ground truth spike times

(**A, B**) Continuous and discrete spike predictions across methods. The top two traces (black)

show raw fluorescence and Gaussian-convolved (10 ms) ground truth spike times. Ground truth

spike times are indicated as vertical tick marks. Discrete imputed spikes for each method are
shown above the corresponding continuous prediction trace, sorted into true positives, false
positives, and false negatives. An epoch was selected from a representative cell which
produced high signal-to-noise ratio (SNR) fluorescence changes in response to an action
potential (**A**) or low SNR (**B**). (**C**) Columns show the proportions of imputed spikes that were
scored as true positive (tp), false positive (fp), or false negative (fn) for each method with a
comparison of all imputed spikes produced by each method (tp + fp) compared to total number
of ground truth (gt) spikes from the 37 putative excitatory cells in the Janelia dataset. (**D-F**)
Precision (tp/(tp + fn), **D**), Recall (tp/(tp + fp), **E**), and F-score (harmonic mean of recall and
precision, **F**) as a function of the SNR of each recorded epoch. The black line in each plot
indicates a regression fit to the data with r^2 indicating the amount of variation explained by SNR
and p indicating significance of the t-statistic comparing a linear to a constant model. Of all
methods, only MLspike shows reduced performance at the SNR levels tested across this
dataset. In particular, the precision (propensity to find false negatives) was increased in low
SNR conditions.
522
523 **Supplementary Figure 10: Comparison of matched and unmatched temporal**
524 **differences across methods.**

525
526 (**A**) Bias and uncertainty between imputed and ground truth spikes matched with a 100 ms.
527 Each plot is a kernel density estimate over a boxplot of indicated metric for each imputed spike.
528 (**B**) As **A** for all imputed spikes compared to the nearest ground truth spike. Note that the
529 supervised methods show increased bias and uncertainty when removing double-counted
530 imputed spikes by using matched statistics.
531

Discussion

In this study, we identified and characterized nonlinear distortions arising from spike
activity patterns in fluorescence signals from three fast GCaMP variants, GCaMP6f, jGCaMP7f,
and jGCaMP8f. We discovered a novel use-dependent slowing of the fluorescence decay during

periods of sustained activity. Using these empirical observations, we constructed a biophysical
model which served as the foundation of an unsupervised (Biophys$_{SMC}$) and a supervised
(Biophys$_{ML}$) spike inference method. These biophysically-based inference methods extracted
spikes with greater precision than existing methods. The advantage of biophysical model-based
approaches was most apparent under high-bandwidth conditions.
Our spike-decoding framework was composed of three basic parts. First, we made
recordings in a minimal tissue-based system (acute slices) where paired fluorescent/spike
recordings were as easy as possible to make. Second, we used in vitro recordings where all
time-varying parameters of experiments were known precisely, save the GCaMP response
itself. Finally, we used these recordings to jointly constrain models of the GCaMP response per
se and to suggest an appropriate generative model of the cellular response.
Our study clarifies how response nonlinearities can differ between indicators. Earlier
GCaMPs (6f and 7f) showed a facilitation-like nonlinearity in their fluorescence response lasting
tens of milliseconds, amplifying signals from closely timed spikes as previously noted (Éltes et
al., 2019; Wei et al., 2020; Huang et al., 2021; Schoenfeld et al., 2021). We found an additional,
novel nonlinearity: a persistent use-dependence that builds progressively during periods of
heightened activity. This second nonlinearity was particularly apparent for jGCaMP8f, whose
half-decay responses were slowed by up to three-fold. We showed that this effect is problematic
for unsupervised methods that assume a linear mapping of spikes to fluorescence. In the linear
SMC approach we used as a benchmark (PGBAR, (Diana et al., 2024)), elongated decay
dynamics drove spurious predictions as the hysteretic effect forced the latent states to match
the fluorescence dynamics (**Figure 3G, H**). This linear SMC, as well as MLspike (Deneux et al.,
2016), were particularly poor spike predictors in the context of low SNR, high spike-rate data
from inhibitory cells (**Figure 3H**; **Figure 4C**).
By leveraging a biophysically-based understanding of sensor behavior, we developed
two complementary, high-performance spike inference methods. Initially, we applied the
generative model directly within an unsupervised SMC framework to substantially improve
accuracy, albeit with a computational cost inherent to Monte Carlo-based approaches
(Vogelstein et al., 2010; Speiser et al., 2017). To address this limitation, we then used our
biophysical model to produce large, synthetic datasets to train a supervised neural network that
performed more quickly.
To our knowledge, our study is the first to report improvements in a machine learning
training using a generative physical model approach. Because the biophysical model captures
important features of GCaMP responses, we were able to use it as the engine for exploring how
GCaMPs would react under different cellular and activity contexts. We achieved substantial
improvements in spike time estimates by training our model on simulated spike patterns that
explored a wide range of timing and patterns. The size and diversity of our training may explain
why Biophys$_{ML}$ outperformed CASCADE trained on the jGCaMP8f data itself. While large, the
jGCaMP8f excitatory cell dataset still falls within a range over which increasing training set size
would be expected to improve performance (Rupprecht et al., 2021; Zhou et al., 2023). Indeed,
limited training data is a fundamental hurdle for any machine learning approach, a problem that
is particularly acute in the context of labor-intensive datasets such as paired
optical/electrophysiology considered here.

However, training of supervised neural networks, even on extensive, heterogeneous
datasets, was insufficient to guarantee accurate inference for sensors with previously unseen
properties. Both ENS[2] and CASCADE, which rank highly on established benchmarks, exhibited
temporal biases and overestimated rise times for jGCaMP8f (**Figure 4**). Notably, in contrast to a
published report (Zhou et al., 2023), CASCADE outperformed ENS[2] on all aspects of inference
in the jGCaMP8f dataset. This result highlights that observations made with neural networks
trained on broad datasets do not generalize in instances where indicator properties differ
substantially from those in the training dataset. Taken together, these results suggest that for
both current and future calcium indicator proteins, building an effective supervised inference
approach requires a thorough exploration of relevant dynamic regimes using either labor-
intensive ground truth or the generative approach we describe here.
In summary, this study elucidates the biophysical mechanisms underlying GECI
fluorescence responses and offers practical solutions for more accurate and reliable neural
activity detection. As a final note, our approach may serve as a generalizable roadmap for
linking fluorescence measurements to underlying neural activity. For example, the steps we
introduce here are likely to generalize to other GECI variants—including red-shifted (Inoue et
al., 2015, 2019; Dana et al., 2016) and chemigenetic sensors (Deo et al., 2021; Farrants et al.,
2024). Appropriate state and sensor models may allow extension to other genetically encoded
sensors including those designed to detect neuromodulator release (e.g., Patriarchi et al., 2018;
Sun et al., 2018) and voltage change (e.g., Evans et al., 2023; Platisa et al., 2023).

Methods

Animals.

601   The following strains were used for physiological experiments: C57Bl/6J (Jackson
602   Laboratories stock 000664); CB6F1 (BalbC and C57Bl/6J F1); and Gabra6 (B6;129P2-
603   Gabra6tm2(cre)Wwis/Mmucd). Both males and females were used and were between 40 and
604   130 days old. Animals were used in accordance with Princeton University's Institutional Animal
605   Care and Use Committee and Institut Pasteur's CEEA-Paris1 approved protocols.
606

Protein purification.

GCaMP variants were cloned into the pET28b protein expression vector using
GCaMP6f, 7f, and 7s backbones generously provided by Douglas Kim (Janelia) and Addgene
(104488). Plasmids were transformed into IPTG-inducible BL21(DE3) competent E. coli
(MilliporeSigma 694504). Starter cultures of transformed cells were grown in 5 ml LB medium
with 50 mg/L kanamycin at 37°C for 4 hours. Cultures were then added to 1 L LB medium with
50 mg/L kanamycin and grown at 37°C until the OD at 600 nm reached 1.0. 1 mM IPTG was
added and the incubation temperature dropped to 25°C overnight for optimal protein expression.
Liquid cultures were centrifuged at 6,000 relative centrifugal force (rcf) for 10 minutes.
Pellets were resuspended in 25 mM Tris-HCl, 500 mM NaCl, 20 mM imidazole, pH 8, with 1 mM
PMSF for protease inhibition. Cells were mechanically lysed through an Emulsiflex homogenizer
(ATA Scientific). Cell debris was removed through centrifugation at 13,000 rcf for 45 minutes at
4°C and the supernatant bound to nickel-NTA agarose (ThermoScientific 25215). Protein was
eluted with 25 mM Tris-HCl, 500 mM NaCl, 500 mM imidazole, pH 8. Proteins were further

concentrated using 10 kDa MWCO centrifugal filters (MilliporeSigma UFC901008) and desalted
(GE Healthcare 17085101) into 130 mM KMOPS (in mM:100 KCl, 30 MOPS, pH 7.2), or a
buffer mimicking intracellular conditions (in mM: 125 KCl, 14.5 KOH, 30 NaCl, 0.7 Mg2Cl, 10
HEPES) termed internal buffer as modified from (Tran et al., 2018). Protein concentrations were
determined by absorption spectrophotometry using alkali denaturation in 0.1M NaOH. Under
these conditions, mature GCaMP chromophore exhibits a standard molar absorption coefficient
of 44,000M-1cm-1 at 447 nm, allowing direct recovery of GCaMP concentration using Beer's
law (Ward, 2005; Barnett et al., 2017).
629
630   **Protein steady-state and kinetic characterization.**
631   Excitation and emission spectra were measured on a PTI Quantamaster 800 spectro-
fluorometer (Horiba Jobin Yvon Inc., Edison, NJ, USA) with data collection under control of Felix
GX software. For calcium-dependent fluorescence, steady-state measurements were made at
20°C using 0.15 μM of purified protein suspended in zero-Ca2+ buffer or high-Ca2+ buffer from
a calcium calibration kit (ThermoFisher C3008MP). The zero-Ca2+ buffer was reciprocally
diluted with the high-Ca2+ buffer to reach free Ca2+ concentrations between 0.01 and 10 μM.
Hill coefficient and Kd were estimated by fitting steady state data to the Hill equation.
638   The fluorometer was also used to prepare reagents for stopped-flow kinetic
639   measurements. Nominally Ca2+-free buffers included KMOPS or internal buffer supplemented
640   by 2 mM BAPTA (Invitrogen B1204). Increasing amounts of 1 M CaCl2 were added to create
641   free calcium steps of 0 nM, 50 nM, and increasing by approximately 2x free calcium in ten steps
642   to 10 μM free calcium. Indicators were diluted to 175 nM in 0.1mM solution buffered by 0.1 mM
643   BAPTA (see design considerations in **Supplementary Figure 2**).
644   Off-response kinetics buffers were designed to have a starting free calcium
645   concentration of 10 μM in 100 μM BAPTA and a final nominal concentration of 0 nM or 50 nM
646   free calcium when mixed to Ca2+-free buffer. Proteins were diluted to 35 nM in the 10 μM free
647   calcium buffer. In all cases, fura-2 (Molecular Probes F6799) and fura-4F (Invitrogen F14174)
648   were used to calibrate free calcium concentrations.
649   On-response and off-response kinetics were measured using an Applied Photophysics
650   SX20 stopped-flow spectrometer. Excitation light was provided by a xenon arc lamp directed
651   through a monochromator set to a 4 nm passband centered on 480 nm, while emissions were
652   collected through a 550 nm longpass filter (Newport FR-OG550). Dead time in the range of 0.6-
653   1.5 ms was estimated by extrapolation from measurements using OGB-1 (Life Technologies,
654   O6806). Measurements were made at 37°C with 60 μl per reaction chamber.
655
656   **Adeno-associated virus constructs.**
657   Adeno-associated viruses (AAVs) were prepared by cloning the GCaMP variants into
658   pAAV backbones (GenScript, Piscataway, NJ). Plasmids were incorporated into AAV serotype 1
659   (AAV2/1) and serotype DJ (AAV DJ) by the Princeton Neuroscience Institute Viral
660   Neuroengineering Laboratory.
661   **Stereotactic viral Injection.**
662   Viral vectors were injected stereotactically into the cerebellar vermis. Mice were deeply
663   anesthetized by intraperitoneal injections of 1.5% ketamine (Mérial) and 0.05% xylazine (Bayer).
664   2% Xylocaine (Newpharma) was applied to the cranial incision. 100 nl vermal (6.5 mm caudal to

665 bregma, lateral 0.2 mm, ventral 3.6 mm and 3.4 mm) and 100 nl Crus I (6 mm caudal to
666 bregma, lateral 3 mm, ventral 0.5 mm) injections were infused over one minute. To allow for
667 optimal transgene expression, mice were imaged 2 to 4 weeks after injection.
668

Slice preparation.
Animals were euthanized by rapid decapitation. The brains were quickly removed and
placed in an ice-cold solution containing (in mM) 2.5 KCl, 0.5 CaCl2, 4 MgCl2, 1.25 NaH2PO4,
24 NaHCO3, 25 glucose, 230 sucrose, and 0.5 ascorbic acid. The buffer was bubbled with
95% O2 and 5% CO2. A Leica VT1200S vibratome was used to prepare coronal slices (200
µm), which were incubated at 32 °C for 30 min in (in mM): 85 NaCl, 2.5 KCl, 0.5 CaCl2, 4 MgCl2,
1.25 NaH2PO4, 24 NaHCO3, 25 glucose, 75 sucrose, and 0.5 ascorbic acid. The external
recording solution contained (in mM) 125 NaCl, 2.5 KCl, 1.5 or 2.0 CaCl2, 1.5 or 1.0 MgCl2,
1.25 NaH2PO4, 25 NaHCO3, 25 glucose, and 0.5 ascorbic acid. Slices were maintained at room
temperature for up to 6 h. All slice recordings were performed at 36–38°C.
679

Imaging in cerebellar brain slices.
Parallel fiber boutons expressing fluorescent virus were imaged with an Ultima two-
photon scanning scanhead (Bruker Nano Surfaces Division, Middleton, WI, USA) that was
mounted on an Olympus BX61W1 microscope equipped with a water-immersion objective
(60×/1.1-NA; Olympus Optical, Tokyo, Japan) and infrared Dodt-gradient contrast. Two-photon
excitation was performed with a pulsed Ti:Sapphire laser (DeepSee, Spectra-Physics, France)
tuned to 920 nm. Extracellular parallel fiber axonal stimulation was performed with a constant
voltage stimulator (Digitimer Ltd, Letchworth Garden City, UK) and a patch pipette (typically with
a tip resistance of 4–6 MΩ) filled with artificial cerebrospinal fluid and placed in the molecular
layer adjacent to labeled parallel fibers. The fluorescence of single cerebellar granule cells was
monitored using circular line scans along the cytoplasm of the GC somata or straight line scans
across individual parallel fiber boutons. Typical line rates were ~1 kHz. To determine threshold
fluorescence, extracellular stimulation intensity was increased until clear fluorescence
responses were observed after a train of 20–60 60-µs pulses at 100 or 300 Hz. To ensure action
potential initiation for each stimulus pulse, the stimulus voltage was increased by 5 V above
threshold.
696

Biophysical and cell models.
698   We modeled GCaMP fluorescence as arising from semi-independent ("hemi-
699 independent") (Lai et al., 2015) calcium binding at two lobes, each of which can subsequently
700 bind a target peptide to become fluorescent. Formally, each lobe transitions between an
701 unbound and a calcium-bound state, which is governed by first-order binding and unbinding
702 reactions:
703

$$\frac{d(\text{bound}_{\text{lobe}})}{dt} = k_{\text{on,lobe}}\,\text{unbound}_{\text{lobe}} - k_{\text{off,lobe}}\,\text{bound}_{\text{lobe}}$$
704   eqn. 1
705

706 where $k_{\text{on,lobe}}$ and $k_{\text{off,lobe}}$ are rate constants for binding and unbinding at each lobe. We
707 defined:

$$k_{\text{on,lobe}} = \frac{k_{\text{off,lobe}}}{\left(K_{a,\text{lobe}}\right)^{H_{\text{lobe}}}}$$

709 eqn. 2

711 where $K_{a,\text{lobe}}$ is an apparent dissociation constant and H_{lobe} is a Hill coefficient describing
712 cooperative binding. Once a lobe is bound with calcium, it can transition to a fluorescent state
713 by binding the target peptide. This step similarly follows first-order kinetics:

$$\frac{d(\text{fluorescent})}{dt} = k_{\text{on,peptide,lobe}}\,\text{bound}_{\text{lobe}} - k_{\text{off,peptide,lobe}}\,\text{fluorescent}$$

715 eqn. 3

717 All fluorescent states (i.e., those in which a lobe is calcium-bound and has the peptide bound)
718 contribute equally to total fluorescence. For stopped-flow simulations, model fluorescence was
719 normalized between 0 (occupancy of fluorescent states at zero calcium) and 1 (occupancy at
720 saturating calcium). For cell-based simulations, fluorescence was scaled to a dimensionless
721 quantity F by setting to 1 at 0 calcium and the parameter R_f – the dynamic range of the indicator
722 – at saturating calcium level. $\frac{\Delta F}{F}$ was then calculated as:

$$\frac{\Delta F}{F} = \frac{F\left(\text{Ca}_{\text{current}}\right) - F\left(\text{Ca}_{\text{basal}}\right)}{F\left(\text{Ca}_{\text{basal}}\right)}$$

724 eqn. 4

726 where $F\left(\text{Ca}_{\text{current}}\right)$ is the instantaneous normalized fluorescence at the current free-calcium
727 level, and $F\left(\text{Ca}_{\text{basal}}\right)$ is the steady-state fluorescence at basal calcium.
728  To simulate intracellular calcium transients (e.g., after action potentials), we extended
729 the model from Rebola et al. 2019 (Rebola et al., 2019) by including an internal calcium
730 reservoir with influx and efflux governed by Michaelis–Menten dynamics. Because cerebellar
731 granule cells have high concentrations of fast, low affinity endogenous buffers, we linearized the
732 buffer's binding capacity κ_B (Neher and Augustine, 1992) to reduce computational cost.
733 Specifically, each small increment of free calcium Ca_{in} was instantly equilibrated with
734 endogenous buffer:

$$\frac{d(\text{BCa})}{dt} = \frac{\text{Ca}_{\text{in}}\,\kappa_B}{\kappa_B + 1}$$

736 eqn. 5

738 where BCa is the portion of endogenous buffer bound to calcium, a ratio of approximately 40 for
739 our standard granule cell model. This linearization preserves the overall calcium balance while
740 allowing faster numerical integration of high buffer concentration simulations.
741  For obtaining full GCaMP model parameterization we globally fit all traces obtained from
742 stopped flow for each variant by gradient descent from 1500 quasi-randomly selected initial
743 points in parameter space. GCaMP parameters from each run were then fixed and best fits
744 were obtained across our ex vivo datasets by gradient descent over cell-specific parameters.

745   The GCaMP model parameters which minimized errors across experimental contexts were then
746   selected and fixed for each GCaMP variant.
747

Bayesian and machine-learning spike inference algorithms.

In order to infer spike times from fluorescence time series using the biophysical model of
jGCaMP8f described in the main text we adopted the Bayesian framework developed by Diana
et al, 2024 based on sequential Monte Carlo methods (particle Gibbs with ancestor sampling,
PGAS; (Lindsten et al., 2014; Diana et al., 2024) which enabled us to estimate simultaneously
the posterior distributions of spike times and the parameters of the cell model, including the
GCaMP concentration, calcium extrusion rates, peak amplitude of the calcium flux due to single
action potentials and the dynamic range of the probe. Single point estimates of model
parameters and spike times were obtained by using maximum-a-posteriori estimates.
We compared the performance of our method with MLSpike (Deneux et al., 2016) which
employs a different model for the calcium indicator dynamics and baseline fluorescence. Spike
inference in MLspike is carried out by discretizing all continuous dynamical variables in the
model and applying the Viterbi algorithm to extract maximum-a-posteriori estimates of spike
times. One of the weaknesses of this approach is the difficulty of estimating the model
parameters, for which Deneux et al provide ad hoc calibration schemes. We directly addressed
this issue using the PGAS sampling algorithm.
We compared our method with state-of-the-art supervised machine-learning models:
CASCADE (Rupprecht et al., 2021) and ENS[2] (Zhou et al., 2023). The two approaches use
different network architectures. CASCADE returns the spike probability at each time point by
taking as input a local time window by using a trained convolutional network while ENS[2] uses
the U-net applied on fixed-length segments of the original calcium trace.
769

Analysis of discrete action potential imputation.

Continuous prediction traces were converted to discrete imputed spike times using the
approach indicated by each published method. For Biophys$_{SMC}$, spike predictions were
calculated using the maximum a-posteriori of the posterior distribution of the spike state over
time. For Biophys$_{ML}$, spike predictions were calculated by optimally fitting a set of Gaussian
kernels with prior width and height determined by approximating the smoothing filter applied to
ground truth for model training as in (Rupprecht et al., 2021).
To avoid double-counting of imputed spikes for a given ground truth spike, we employed
the linear assignment algorithm to identify unique imputed/ground truth pairs. We used the
matlab function matchpairs at a cost of unassignment as specified in the main text to constrain
time interval between such pairs as specified in the main text.True positives were then defined
as imputed spikes with unique ground truth matches, with unmatched ground truth spikes giving
the false negatives, and unmatched imputed spikes being false positives. Total true positives,
false positives, and ground-truth spikes within epochs for each cell to form the distributions for
F-score, precision, and recall/specificity.
For each epoch of ground-truth recording, the bias in each imputed spike time was
calculated as the interval to the nearest ground-truth action potential while uncertainty was
defined as the absolute value of this quantity. The measures were taken to either the nearest

ground truth spike using a nearest neighbors search (unmatched) or to the nearest matched
spike.
790

Statistical presentation.

792   Box and whisker plots comprise a box indicating the median and 25-75th percentile
793 range and whiskers indicating the shorter of 1.5 times the interquartile range or the extreme
794 data point. Descriptive statistics were calculated using median-based methods, and the
795 estimated standard deviation was defined as the median absolute deviation divided by 0.6745.
796

References

Akerboom J et al. (2012) Optimization of a GCaMP calcium indicator for neural activity imaging.
J Neurosci Off J Soc Neurosci 32:13819–13840.
Badura A, Sun XR, Giovannucci A, Lynch LA, Wang SS-H (2014) Fast calcium sensor proteins
for monitoring neural activity. Neurophotonics 1:025008.
Barnett LM, Hughes TE, Drobizhev M (2017) Deciphering the molecular mechanism responsible
for GCaMP6m's Ca2+-dependent change in fluorescence. PLOS ONE 12:e0170934.
Brown SE, Martin SR, Bayley PM (1997) Kinetic Control of the Dissociation Pathway of
Calmodulin-Peptide Complexes*. J Biol Chem 272:3389–3397.
Chen T-W, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A, Schreiter ER, Kerr RA,
Orger MB, Jayaraman V, Looger LL, Svoboda K, Kim DS (2013) Ultrasensitive
fluorescent proteins for imaging neuronal activity. Nature 499:295–300.
Dana H, Mohar B, Sun Y, Narayan S, Gordus A, Hasseman JP, Tsegaye G, Holt GT, Hu A,
Walpita D, Patel R, Macklin JJ, Bargmann CI, Ahrens MB, Schreiter ER, Jayaraman V,
Looger LL, Svoboda K, Kim DS (2016) Sensitive red protein calcium indicators for
imaging neural activity. eLife 5:e12727.
Dana H, Sun Y, Mohar B, Hulse BK, Kerlin AM, Hasseman JP, Tsegaye G, Tsang A, Wong A,
Patel R, Macklin JJ, Chen Y, Konnerth A, Jayaraman V, Looger LL, Schreiter ER,
Svoboda K, Kim DS (2019) High-performance calcium sensors for imaging activity in
neuronal populations and microcompartments. Nat Methods 16:649–657.
Demas J, Manley J, Tejera F, Kim H, Traub FM, Chen B, Vaziri A (2021) High-Speed, Cortex-
Wide Volumetric Recording of Neuroactivity at Cellular Resolution using Light Beads
Microscopy. bioRxiv:2021.02.21.432164.
Deneux T, Kaszas A, Szalay G, Katona G, Lakner T, Grinvald A, Rózsa B, Vanzetta I (2016)
Accurate spike estimation from noisy calcium signals for ultrafast three-dimensional
imaging of large neuronal populations in vivo. Nat Commun 7:12190.
Deo C, Abdelfattah AS, Bhargava HK, Berro AJ, Falco N, Farrants H, Moeyaert B, Chupanova
M, Lavis LD, Schreiter ER (2021) The HaloTag as a general scaffold for far-red tunable
chemigenetic indicators. Nat Chem Biol 17:718–723.
Diana G, Sermet BS, DiGregorio DA (2024) High frequency spike inference with particle Gibbs
sampling. eLife 13.
DiGregorio DA, Peskoff A, Vergara JL (1999) Measurement of Action Potential-Induced
Presynaptic Calcium Domains at a Cultured Neuromuscular Junction. J Neurosci
19:7846–7859.
Éltes T, Szoboszlay M, Kerti-Szigeti K, Nusser Z (2019) Improved spike inference accuracy by
estimating the peak amplitude of unitary [Ca2+] transients in weakly GCaMP6f-
expressing hippocampal pyramidal cells. J Physiol 597:2925–2947.
Evans SW et al. (2023) A positively tuned voltage indicator for extended electrical recordings in

the brain. Nat Methods 20:1104–1113.

Faas GC, Raghavachari S, Lisman JE, Mody I (2011) Calmodulin as a direct detector of Ca2+
signals. Nat Neurosci 14:301–304.

Farrants H et al. (2024) A modular chemigenetic calcium indicator for multiplexed in vivo
functional imaging. Nat Methods 21:1916–1925.

Friedrich J, Zhou P, Paninski L (2017) Fast online deconvolution of calcium imaging data. PLOS
Comput Biol 13:e1005423.

Huang L, Ledochowitsch P, Knoblich U, Lecoq J, Murphy GJ, Reid RC, de Vries SE, Koch C,
Zeng H, Buice MA, Waters J, Li L (2021) Relationship between simultaneously recorded
spiking activity and fluorescence signal in GCaMP6 transgenic mice Westbrook GL,
Svoboda K, Higley M, Sabatini BL, eds. eLife 10:e51675.

Inoue M et al. (2019) Rational Engineering of XCaMPs, a Multicolor GECI Suite for In Vivo
Imaging of Complex Brain Circuit Dynamics. Cell 177:1346-1360.e24.

Inoue M, Takeuchi A, Horigane S, Ohkura M, Gengyo-Ando K, Fujii H, Kamijo S, Takemoto-
Kimura S, Kano M, Nakai J, Kitamura K, Bito H (2015) Rational design of a high-affinity,
fast, red calcium indicator R-CaMP2. Nat Methods 12:64–70.

Johnson JD, Snyder C, Walsh M, Flynn M (1996) Effects of Myosin Light Chain Kinase and
Peptides on Ca2+ Exchange with the N- and C-terminal Ca2+ Binding Sites of
Calmodulin (∗). J Biol Chem 271:761–767.

Lai M, Brun D, Edelstein SJ, Novère NL (2015) Modulation of Calmodulin Lobes by Different
Targets: An Allosteric Model with Hemiconcerted Conformational Transitions. PLOS
Comput Biol 11:e1004063.

Li P, Geng X, Jiang H, Caccavano A, Vicini S, Wu J (2019) Measuring Sharp Waves and
Oscillatory Population Activity With the Genetically Encoded Calcium Indicator
GCaMP6f. Front Cell Neurosci 13.

Lindsten F, Jordan MI, Schön TB (2014) Particle Gibbs with Ancestor Sampling. J Mach Learn
Res 15:2145–2184.

Linse S, Helmersson A, Forsén S (1991) Calcium binding to calmodulin and its globular
domains. J Biol Chem 266:8050–8054.

Pachitariu M, Stringer C, Harris KD (2018) Robustness of Spike Deconvolution for Neuronal
Calcium Imaging. J Neurosci 38:7976–7985.

Patriarchi T, Cho JR, Merten K, Howe MW, Marley A, Xiong W-H, Folk RW, Broussard GJ,
Liang R, Jang MJ, Zhong H, Dombeck D, Zastrow M von, Nimmerjahn A, Gradinaru V,
Williams JT, Tian L (2018) Ultrafast neuronal imaging of dopamine dynamics with
designed genetically encoded sensors. Science 360.

Platisa J, Ye X, Ahrens AM, Liu C, Chen IA, Davison IG, Tian L, Pieribone VA, Chen JL (2023)
High-speed low-light in vivo two-photon voltage imaging of large neuronal populations.
Nat Methods 20:1095.

Pnevmatikakis EA, Merel J, Pakman A, Paninski L (2013) Bayesian spike inference from
calcium imaging data.

Rebola N, Reva M, Kirizs T, Szoboszlay M, Lőrincz A, Moneron G, Nusser Z, DiGregorio DA
(2019) Distinct Nanoscale Calcium Channel and Synaptic Vesicle Topographies
Contribute to the Diversity of Synaptic Function. Neuron 104:693-710.e9.

Regehr WG, Atluri PP (1995) Calcium transients in cerebellar granule cell presynaptic terminals.
Biophys J 68:2156–2170.

Rose T, Goltstein PM, Portugues R, Griesbeck O (2014) Putting a finishing touch on GECIs.
Front Mol Neurosci 7.

Rupprecht P, Carta S, Hoffmann A, Echizen M, Blot A, Kwan AC, Dan Y, Hofer SB, Kitamura K,
Helmchen F, Friedrich RW (2021) A database and deep learning toolbox for noise-
optimized, generalized spike inference from calcium imaging. Nat Neurosci 24:1324–
1337.

Sabatini BL, Regehr WG (1998) Optical measurement of presynaptic calcium currents. Biophys
J 74:1549–1563.
Sariyildiz MB, Kalantidis Y, Alahari K, Larlus D (2023) No Reason for No Supervision: Improved
Generalization in Supervised Models.
Schoenfeld G, Carta S, Rupprecht P, Ayaz A, Helmchen F (2021) In Vivo Calcium Imaging of
CA3 Pyramidal Neuron Populations in Adult Mouse Hippocampus. eNeuro 8.
Sebastian J, Mari Ganesh Kumar M, Sreekar YS, Rikhye RV, Sur M, Murthy HA (2017)
GDspike: An accurate spike estimation algorithm from noisy calcium fluorescence
signals. 2017 IEEE Int Conf Acoust Speech Signal Process ICASSP:1043–1047.
Seroussi I, Zeitouni O (2022) Lower Bounds on the Generalization Error of Nonlinear Learning
Models.
Speiser A, Yan J, Archer EW, Buesing L, Turaga SC, Macke JH (2017) Fast amortized
inference of neural activity from calcium imaging data with variational autoencoders. In:
Advances in Neural Information Processing Systems. Curran Associates, Inc.
Sun F et al. (2018) A Genetically Encoded Fluorescent Sensor Enables Rapid and Specific
Detection of Dopamine in Flies, Fish, and Mice. Cell 174:481-496.e19.
Sun XR, Badura A, Pacheco DA, Lynch LA, Schneider ER, Taylor MP, Hogue IB, Enquist LW,
Murthy M, Wang SS-H (2013) Fast GCaMPs for improved tracking of neuronal activity.
Nat Commun 4:2170.
Tian L, Hires SA, Mao T, Huber D, Chiappe ME, Chalasani SH, Petreanu L, Akerboom J,
McKinney SA, Schreiter ER, Bargmann CI, Jayaraman V, Svoboda K, Looger LL (2009)
Imaging neural activity in worms, flies and mice with improved GCaMP calcium
indicators. Nat Methods 6:875.
Tran V, Park MCH, Stricker C (2018) An improved measurement of the Ca2+-binding affinity of
fluorescent Ca2+ indicators. Cell Calcium 71:86–94.
Vogelstein JT, Packer AM, Machado TA, Sippy T, Babadi B, Yuste R, Paninski L (2010) Fast
Nonnegative Deconvolution for Spike Train Inference From Population Calcium Imaging.
J Neurophysiol 104:3691–3704.
Ward WW (2005) Biochemical and Physical Properties of Green Fluorescent Protein. In: Green
Fluorescent Protein, pp 39–65. John Wiley & Sons, Ltd.
Wei Z, Lin B-J, Chen T-W, Daie K, Svoboda K, Druckmann S (2020) A comparison of neuronal
population dynamics measured with calcium imaging and electrophysiology. PLOS
Comput Biol 16:e1008198.
Xiao D, Vanni MP, Mitelut CC, Chan AW, LeDue JM, Xie Y, Chen AC, Swindale NV, Murphy TH
(2017) Mapping cortical mesoscopic networks of single spiking cortical or sub-cortical
neurons Kleinfeld D, ed. eLife 6:e19976.
Zhang Y et al. (2023) Fast and sensitive GCaMP calcium indicators for imaging neural
populations. Nature 615:884–891.
Zhou Z, Yip HM, Tsimring K, Sur M, Ip JPK, Tin C (2023) Effective and efficient neural networks
for spike inference from in vivo calcium imaging. Cell Rep Methods 3:100462.
926