Fluorescence-barcoded cell lines stably expressing membrane-anchored influenza neuraminidases

Joel Finney[1]*, Masayuki Kuraoka[2], Shengli Song[3], Akiko Watanabe[2], Xiaoe Liang[2], Dongmei Liao[2], M. Anthony Moody[2,4,5], Emmanuel B. Walter[4,5], Stephen C. Harrison[1,6]*, Garnett Kelsoe[2,3,5]

[1]Laboratory of Molecular Medicine, Children's Hospital, Harvard Medical School, Boston, MA 02115, USA
[2]Department of Integrative Immunobiology, Duke University, Durham, NC 27710, USA
[3]Department of Surgery, Duke University, Durham, NC 27710, USA
[4]Department of Pediatrics, Duke University School of Medicine, Durham, NC 27710, USA
[5]Duke Human Vaccine Institute, Duke University, Durham, NC 27710, USA
[6]Howard Hughes Medical Institute, Boston, MA 02115, USA

*Corresponding authors

Email: jfinney@crystal.harvard.edu, harrison@crystal.harvard.edu

Mailing address for both corresponding authors:
Harvard Medical School
250 Longwood Ave., SGMB 123
Boston, MA 02115

ABSTRACT

The discovery of broadly protective antibodies to the influenza virus neuraminidase (NA) has raised interest in NA as a vaccine target. However, recombinant, solubilized tetrameric NA ectodomains are often challenging to express and isolate, hindering the study of anti-NA humoral responses. To address this obstacle, we established a panel of 22 non-adherent cell lines stably expressing native, historical N1, N2, N3, N9, and NB NAs anchored on the cell surface. The cell lines are barcoded with fluorescent proteins, enabling high-throughput, 16-plex analyses of antibody binding with commonly available flow cytometers. The cell lines were at least as efficient as a Luminex multiplex binding assay at identifying NA antibodies from a library of unselected clonal IgGs derived from human memory B cells. The membrane-anchored NAs are catalytically active and are compatible with established small-molecule catalytic activity assays. NA-expressing K530 cell lines therefore represent a useful tool for studying NA immunity and evaluating influenza vaccine efficacy.

INTRODUCTION

Influenza A and B viruses (IAVs and IBVs) are substantial public health burdens. Two A subtypes (H1N1 and H3N2) and one B lineage (Victoria) currently circulate among humans (1). Another B lineage (Yamagata) that recently circulated may have gone extinct (2, 3). Additional flu strains circulating in animal populations can also cause zoonosis (4, 5). Seasonal flu vaccines offer important protection against disease, but they elicit narrow immunity against the proteins of the viral strains included in the vaccine, necessitating annual vaccine updates as pressure from herd immunity selects for escape mutations (6-9). Thus, the goal of many next-generation flu vaccines is to elicit antibody (Ab)-mediated protection against serologically divergent viral strains (10, 11). Much of the flu vaccine research of the last 15 years has focused on Ab responses to hemagglutinin (HA), the major glycoprotein on the virion surface, which mediates attachment to sialylated cellular receptors and catalyzes membrane fusion to release the viral genome into the host cell. The viral neuraminidase (NA), a less abundant glycoprotein whose function is to cleave sialic acids from cellular receptors to enable viral egress, is also an immune system target (12). The recent discovery of protective Abs that recognize conserved epitopes on NAs from divergent IAVs and IBVs has renewed interest in designing vaccines that target NA (13-18).

Recombinant, full-length, solubilized NA ectodomains are frequently challenging to express, impeding study of NA immunity. The typical expression construct for recombinant NA excludes the N-terminal stalk and transmembrane regions, instead substituting a tetramerization domain from another protein (18-20). The yield of the resultant chimeric protein is 10- to 100-fold lower than that routinely achieved for recombinant HA ectodomains (20). Full-length NAs have been expressed in the native, membrane-anchored form on enveloped virus-like particles (14, 15), but expression and purification of virus-like particles requires access to specialized equipment, which may be a barrier for some laboratories. Others have expressed full-length

NAs on transiently transfected, mammalian cell lines (14, 21), but these transfected cell lines

are neither practical for multiplexed analysis of Ab binding breadth nor easily shared between

laboratories.

We have used panels of fluorescence-barcoded K530 cell lines stably expressing

membrane-anchored HAs or major histocompatibility complex (MHC) proteins to determine the

binding breadth and avidity of relevant Abs in a flow cytometry assay (22-24). These non-

adherent cell lines are easy to culture, have short doubling times, and can be used to study Ab

binding to up to 16 different antigens simultaneously (23). The barcodes and fluorescence

intensity of Ab binding can be detected using 405 nm, 488 nm, and 633 nm excitation lasers

and standard emission filter sets available on most flow cytometers, with good inter-assay

reproducibility (23). Here, we have adapted the barcoded cell lines to express native NAs from

historical isolates of influenza A and B viruses that have circulated in human or animal

populations.

RESULTS

Standard NA monoclonal Abs brightly and specifically label K530-NA cell lines

77   We selected monoclonal, fluorescently barcoded K530 cell lines (23) stably expressing

native, full-length NAs representing 22 historical isolates of IAV or IBV (Table I). Another

barcoded cell line expressing no NA controls for binding specificity. Up to 16 cell lines can be

pooled for a single assay, with two different NA options for some of the barcodes. Collectively,

the cell lines express NAs from the N1, N2, N3, and N9 subtypes of IAV, including N1s

spanning 44 years (1977-2021) of human isolates and N2s spanning 65 years (1957-2021) of

human isolates. NAs representing the Victoria and Yamagata lineages of IBV are also included.

84   In a flow cytometry assay, recombinant IgGs (rIgGs) representing well-characterized,

monoclonal Abs (mAbs) against the NA catalytic site (13-15, 18) brightly and specifically labeled

the K530-NA cells, but not control cells that expressed no NA (Fig. 1). As reported, 1G01 (15,

18), FNI17 (14) and DA03E17 (15) bound NAs of diverse subtypes from phylogenetic groups 1

and 2 (Fig. 1). FNI17 and DA03E17 also bound NAs from both lineages of IBV, as described

(14, 15). Although 1G01 was reported to recognize NAs from some strains of IBV, 1G01 binds

IBV NAs much more weakly than IAV NAs, and even 50 μg/ml of 1G01 cannot neutralize

B/Brisbane/60/2008 or B/Phuket/3073/2013 virus in vitro (14, 15, 18). Thus, our observation that

1G01 did not label K530 cells expressing NB.BN08 or NB.PK13 (Fig. 1) agrees with prior

reports. In contrast, as expected, 1G05 mAb (13) avidly bound NAs from both IBV lineages, but

did not bind IAV NAs (Fig. 1). S1V2-72, a control IgG that specifically binds influenza

hemagglutinin (22), did not bind K530-NA cells.

96   Standard mAb CD6 was reported to bind N1.CA09 with high affinity (21). We observed

tight binding of CD6 to N1.CA09-expressing K530 cells (Fig. 1). We also discovered that CD6

avidly bound N1.MI15, a descendant of N1.CA09, and weakly bound a later descendant,

N1.SYD21. CD6 also recognized N1.VN04, which was not detected in an earlier report using

glutaraldehyde-fixed, transiently transfected 293T cells expressing membrane-anchored

N1.VN04 (21). However, our observation that CD6 binds N1.VN04 is consistent with its modest

neutralizing activity toward A/Vietnam/1203/2004 (H5N1) virus (21). Because the CD6 epitope

evenly spans adjacent protomers of the NA homotetramer (21), bright labeling of NA-expressing

K530 cells by CD6 implies that the NA is in its native, multimeric form.

K530-NA cells perform as well as a Luminex binding assay for high-throughput discovery

of NA Abs

To test whether K530-NA cells can be used to identify new NA Abs, we sorted individual

memory B (Bmem) cells from peripheral blood mononuclear cells of a healthy, teenaged donor

(T3) 14 days after immunization with the 2019-2020 seasonal influenza vaccine (for an example

of the sorting strategy, see Supp. Fig 1A) (25). During sorting, Bmem cells were not selected for

binding to any antigen. From these Bmem cell cultures, we obtained 1,847 clonal IgG-containing

culture supernatants, which we screened for NA-binding activity both by Luminex assay (using

tetrameric NA head constructs representing N1s, N2s, and NBs) and by multiplex flow

cytometry (using the nine K530-NA cell lines we had generated at the time of this experiment).

The Luminex screen identified five clonal IgGs that avidly bound NA: T3-P4A8 and T3-P10F5,

which bound N1s from new pandemic H1N1 strains; T3-P17F8 and T3-P27B8, which bound

N2.HK14; and T3-P15C5, which bound NBs from both IBV lineages (Fig. 2A). The Luminex

screen also identified 11 samples with weak or borderline binding to NA. The flow cytometry

screen with K530-NA cells (Fig. 2B) also identified the same five IgGs with avid NA binding, plus

two IgGs with modest NA binding: T3-P20D5, which cross-reacted with numerous N1s, N2s,

and NB; and T3-P28D3, which bound N2.BJ89, an NA not included in the Luminex screen. A

third IgG, T3-P35E4, cross-reacted with N1s, N2s, and NB, but also weakly bound K530 cells

that did not express NA; this sample may be auto- or polyreactive. The K530-NA cells also

identified several IgGs with weak/borderline NA binding. Thus, for high-throughput identification

of NA Abs, K530-NA cells perform at least as well as a Luminex assay using soluble NA heads;

the K530 cells also have the advantage of potentially identifying Abs to the NA stalk domain.

We expressed rIgGs representing NA mAbs identified from donor T3 and two other

teenaged donors (Supp. Fig. 1B-1C), and tested the binding breadth of these rIgGs using all 22

K530-NA cell lines (Fig. 3). The rIgGs recapitulated the binding profiles observed for the

corresponding culture supernatant IgGs in the initial Luminex or K530-based screens (Fig. 3).

Several of the rIgGs also bound additional NAs not included in the initial screens. Each rIgG

bound only one NA subtype, except T3-P20D5, which bound various N1s, N2s, N3.JS21, and

NB.BN08. T3-P20D5 bound these NAs weakly, but specifically: when we set the sensitivity of

the flow cytometer's phycoerythrin (PE) detector to a level that caused the signal from higher-

affinity standard mAbs to approach or exceed the detector limit, T3-P20D5 clearly labeled some

K530-NA cell lines, but not others (Supp. Fig 2A). It is unclear why T3-P20D5 did not bind the

tetrameric NA heads used in the Luminex screen (Supp. Fig. 2B); one possibility is that T3-

P20D5 recognizes an epitope in the NA stalk domain.

K530-NA cell lines have NA catalytic activity

In the NA-Star activity assay, which measures inhibition of sialoside cleavage by active-

site directed antibodies, K530 cell lines expressing N1, N2, N3, N9, or NB each had NA activity

≥100-fold greater than the activity present in control K530 cells expressing no NA (Fig. 4A-4C).

Control mAb S1V2-72 did not inhibit the NA activity of these cell lines, but NA catalytic site-

binding IgGs 1G01 or 1G05 inhibited the NA activity in a dose-dependent manner, with half-

maximal inhibitory concentrations (IC_{50}) comparable to reported values (Fig. 4, Table 2) (13, 15,

18). MAbs T3-P10F5, T3-P27B8, and T1-P30E7 did not significantly inhibit the catalytic activity

(Fig. 4C) of K530-NA cell lines to which the corresponding mAbs bound tightly (Fig. 3), so these

mAbs do not substantially cover the catalytic site, unlike 1G01 and 1G05 (13, 18).

DISCUSSION

Recombinant, soluble NA ectodomains are often challenging to isolate in quantities

sufficient for studies of NA Ab responses. A few alternate approaches to NA expression have

been described, to which we now add display of native, full-length NAs anchored to the surface

of fluorescently barcoded cell lines. These stable cell lines are simple to culture and easy to

share among labs. Inclusion of the native transmembrane and stalk domains of NA should

enable the NA on K530-NA lines to assume the native, tetrameric conformation (26, 27).

Indeed, that K530-NA cell lines have robust catalytic activity and are avidly and specifically

bound by well-characterized, standard mAbs implies that the membrane-bound NAs have the

proper antigenic structure (13-15, 18, 21, 28).

An advantage of K530-NA cells is their display of full-length NA, i.e., both head and stalk

domains. In principle, inclusion of the stalk domain enables discovery of anti-stalk Abs, which

have not yet been reported, and which would be invisible in experiments that use only the NA

head domain. MAb T3-P20D5 may be one such Ab; alternatively, it may bind some other

epitope not represented well on tetrameric, head-only NAs.

ACKNOWLEDGMENT

The research was supported by NIAID grant number P01 AI089618. We thank all members of

the consortium participating in this program project for discussions throughout the course of the

research reported here. We are grateful for the excellent technical support of Amanda Foreman,

Wenli Zhang, Steven Slater, Stephanie H. Smith and Darren J. Morrow at Duke University. This

research was conducted in part using equipment and services provided by the Harvard Medical

School Immunology Flow Cytometry Core Facility, the Duke University DNA Analysis Facility,

and the Duke Human Vaccine Institute Research Flow Cytometry Shared Resource Facility.

SCH is an Investigator in the Howard Hughes Medical Institute.

 FIGURE LEGENDS

FIG 1. Standard NA mAbs brightly and specifically label K530-NA cell lines. Shown are

flow cytometry histograms depicting the binding of recombinant IgG versions of standard NA

mAbs to K530 cell lines expressing membrane-anchored NAs. Each row corresponds to a

monoclonal cell line stably expressing a single type of NA. Each column corresponds to the

binding profile for a single mAb. MAbs were incubated with pooled cell lines comprising Option 1

or Option 2, and the resultant data were concatenated into a single figure.

FIG. 2. K530-NA cell lines perform as well as a Luminex assay for identifying NA mAbs.

A) Luminex assay results showing the median fluorescence intensity (MFI) of antigen binding for

clonal IgG-containing culture supernatants from Bmem isolated from donor T3. Bovine serum

albumin (BSA), insulin, keyhole limpet hemocyanin (KLH), chicken ovalbumin (OVA), and

tetanus toxoid (TT) serve as binding-specificity controls. Short, red, horizontal lines in each

column denote the limit of detection (LOD), which was calculated as six standard deviations

above the mean signal produced by cultures containing no B cells. B) Results of a high-

throughput, flow cytometry-based screen using K530-NA cell lines. Each symbol depicts the

geometric mean fluorescence intensity (geoMFI) of clonal IgG binding to a K530-NA cell line.

The dashed, horizontal line denotes the threshold for binding. Each symbol in (A) and (B)

represents a single, clonal IgG-containing culture supernatant. Clonal IgGs of interest are

denoted with uniquely colored symbols.

FIG. 3. Use of K530-NA cells to determine the binding breadth of newly identified NA

mAbs. Shown are flow cytometry histograms depicting the binding of recombinant IgG versions

of NA mAbs from donors T1, T2, or T3 to K530 cell lines expressing membrane-anchored NAs,

as in Fig. 1.

FIG. 4. NA catalytic activity of K530-NA cell lines. Serially diluted control or NA catalytic site-

binding IgGs were incubated with selected K530 cell lines for 30 min, then the NA catalytic

activity was determined by NA-Star assay. A) Catalytic activity of K530 cell lines expressing IAV

NAs, in the presence of control IgG (S1V2-72) or inhibitory IgG (1G01). B) Catalytic activity of

K530 cells expressing IBV NA, in the presence of control IgG or inhibitory IgG (1G05). C)

Catalytic activity of K530-NA cell lines in the presence of control IgG, inhibitory IgG, or IgGs that

tightly bind the corresponding NAs, but have unknown epitopes (T3-P10F5, T3-P27B8, T1-

P30E7). The lack of inhibition by the latter IgGs indicates that these mAbs do not bind the

catalytic pocket of NA. Error bars represent mean ± S.D.

Supplemental Fig 1. Isolation of IgGs from human memory B (Bmem) cells. A) Flow

cytometry plots showing the sorting strategy for isolation of Bmem. The plots are from donor T1;

the same strategy was used for sorting from other donors' samples. Inset labels denote the

identity of the gated population and its frequency within the parent population. B-C) Luminex

assay results showing the antigen-binding profiles of clonal IgG-containing culture supernatants

from Bmem cells isolated from donors T1 (B) or T2 (C). Short, red, horizontal lines in each

column denote the limit of detection, calculated as six standard deviations above the mean

signal produced by cultures containing no B cells. Clonal IgGs of interest are denoted with

uniquely colored symbols. IgGs denoted by red "X"s in (B) correspond to samples for which

binding to influenza hemagglutinin (HA) was ≥100-fold higher than binding to NA (data not

shown). These samples were considered to be HA-specific IgGs rather than NA-specific IgGs.

Supplemental Fig. 2. MAb T3-P20D5 specifically labels K530-NA cells. A) Shown are flow

cytometry histograms depicting the binding of recombinant IgG versions of NA mAbs to K530

cell lines expressing membrane-anchored NAs, as in Fig 1. MAbs were incubated with pooled

cell lines comprising Option 1. The detector for phycoerythrin (PE) fluorescence, used for

measuring IgG binding, was set at a higher voltage than for a typical experiment. As a result,

the signal for some samples (e.g. CD6 binding to N1.CA09) exceeds the detector's upper limit,

while the signal for T3-P20D5's binding to certain NAs is clearly distinguishable from its lack of

binding to the control K530 cell line that expresses no NA. B) Luminex binding assay results for

the IgG-containing culture supernatant from sample T3-P20D5. The sample contains a

significant quantity of IgG, but binding to recombinant, tetrameric NA heads is below the LOD

(calculated as in Fig. 2A).

MATERIALS AND METHODS

Study approvals and volunteers

The study procedures, informed consent, and data collection documents were reviewed and

approved by the Duke Health Institutional Review Board (Pro00020561, initial approval 2010).

We enrolled three donors (T1, T2, T3) between the ages of 13 yo and 18 yo. Each donor had a

documented history of receiving seasonal influenza vaccines ≥3 times prior to the 2019-2020

northern hemisphere flu season. Written informed consent was obtained from each donor's

parent and assent obtained for each participant. Each donor was administered the 2019-2020

quadrivalent inactivated influenza vaccine, and blood was collected 15 d (T1, T2) or 14 d (T3)

post-vaccination. Blood samples were processed into PBMCs and plasma and then aliquoted

and stored in liquid nitrogen vapor phase or at –80°C, respectively, for future analysis.

Cell line culture

Unless otherwise noted, mammalian cell lines were maintained in static cultures at 37°C with

5% CO_2 in a humidified incubator, and culture reagents were from Gibco. The MEC-147 cell line

(manuscript in preparation), a derivative of MS40L-low feeder cells (*Mus musculus*)(29, 30) that

stably expresses human IL-2, IL-4, IL-21, and BAFF, was expanded from frozen aliquots in

Iscove's Modified Dulbecco's Medium (IMDM) containing 10% HyClone FBS (Cytiva), 2-

mercaptoethanol (55 μM), penicillin (100 units/ml), and streptomycin (100 μg/ml). Lenti-X 293T

cells (*Homo sapiens*, Takara) were cultured in Dulbecco's Modified Eagle Medium (DMEM) plus

10% FBS, penicillin, streptomycin, HEPES (10 mM), sodium pyruvate (1 mM), and 1× MEM

non-essential amino acids. Expi293F cells (*Homo sapiens*; Thermo Fisher) were cultured in

Expi293 Expression Medium plus penicillin and streptomycin, at 8% CO_2 with shaking. K530-

derived cell lines (*Homo sapiens*) (23) were initially cultured in RPMI-1640 medium plus 10%

FBS, 2-mercaptoethanol, penicillin, streptomycin, HEPES, sodium pyruvate, and MEM

nonessential amino acids, but IMDM plus 10% FBS, penicillin, and streptomycin was later

chosen as the standard growth medium. High Five cells (BTI-TN-5B1-4; *Trichoplusia ni*; Thermo

Fisher) were maintained in ESF 921 medium (Expression Systems) at 28°C in spinner flasks in

air. Cell lines were not subject to authentication.

Expression and purification of recombinant, tetrameric NA heads

Recombinant NAs were soluble, head-only, tetrameric ectodomains (20). NAs were expressed

by infection of insect cells with recombinant baculovirus as described (20, 31-34). In brief, a

pFastBac vector was modified to encode a secretion signal peptide, an N-terminal His$_8$ tag, an

HRV3C protease cleavage site, a tetrabrachion (*Staphylothermus marinus*) tetramerization tag,

a thrombin cleave site, and the globular head domain of NA. The resulting baculoviruses

produce tetrameric NA heads. Supernatant from recombinant baculovirus-infected High Five

cells was harvested 72 h post-infection and clarified by centrifugation. Proteins were purified by

adsorption to cobalt-nitrilotriacetic acid (Co-NTA) agarose resin (Takara), followed by a wash in

buffer A (10 mM Tris, 150 mM NaCl, pH 7.5) plus 5 mM imidazole, elution in buffer A plus 350

mM imidazole (pH 8), and gel filtration chromatography on a Superdex 200 column (GE

Healthcare) in buffer A.

Bmem sorting and culture

PBMCs in RPMI medium plus 10% FBS were incubated with irrelevant mouse IgG1 (MG1K;

Rockland) to block nonspecific binding and then labeled with fluorochrome-conjugated mAbs.

The following human surface antigen-specific mAbs, purchased from BD Biosciences,

BioLegend, or Thermo Scientific, were used: anti-human IgM-fluorescein isothiocyanate (FITC)

(MHM-88), anti-CD3-PE-Cy5 (UCHT1), anti-CD14-Tri (TuK4), anti-CD16-PE-Cy5 (3G8), anti-

CD19-PE-Cy7 (HIB19), anti-IgG-allophycocyanin (APC) (G18-145), anti-IgD-PE (IA6-2), anti-

CD27-BV421 (M-T271), and anti-CD24-BV510 (ML5) antibodies. Labeled cells were sorted

using a FACSAria II with Diva software (BD Biosciences). Flow cytometric data were analyzed

with FlowJo software (Tree Star, Inc.). Total Bmem (CD19$^+$ Dump$^-$ CD27$^+$ CD24$^+$) (Supp. Fig.

1A) were identified as described (25, 30, 35). Surface IgD, IgM, and IgG expression were also

determined, but were not considered for sorting. Doublets were excluded from cell sorting by

forward scatter area (FSC-A) versus FSC height (FSC-H) gating. Cells positive for 7-

aminoactinomycin D (7-AAD) (BD Bioscience) or for CD3, CD14, or CD16 expression were also

excluded.

Sorted single Bmem were expanded in the presence of feeder cells as described (25,

30, 35), with some modifications. MEC-147 feeder cells (manuscript in preparation) were used

instead of MS40L-low feeder cells and exogenous cytokines. Single Bmem were sorted directly

into 96-well plates containing feeder cells and 200 µl growth medium per well. After seven days

of co-culture, 100 µl of spent medium was removed from each well and replaced with 200 µl of

fresh growth medium. On culture days ~14, ~17, and ~21, two-thirds of the spent medium from

each well was replaced with an equal volume of fresh growth medium. On culture day 25,

culture supernatants were harvested to screen the secreted clonal IgGs. Expanded clonal B

cells were frozen at –80°C for V(D)J sequence analysis.

Luminex multiplex binding assay

The specificities and avidities of clonal IgG antibodies in culture supernatants were determined

in a multiplex bead assay (Luminex Corp.) as described (25, 36) with modifications. Culture

supernatants were diluted in Luminex assay buffer (phosphate-buffered saline [PBS] containing

1% bovine serum albumin [BSA], 0.05% NaN$_3$, and 0.05% Tween 20) with 1% milk and

incubated for 2 h with the mixture of antigen-coupled microsphere beads in 96-well filter-

bottom plates (Millipore). After washing three times with assay buffer, the beads were incubated

for 1 h with PE-conjugated mouse anti-human IgG antibodies (JDC-10; Southern Biotech).

After three washes, the beads were resuspended in assay buffer and the plates read on a Bio-

Plex 3D suspension array system (Bio-Rad). Antigens and controls included BSA, mouse anti-

human Ig(κ) (SB81a; Southern Biotech), mouse anti-human Ig(λ) (JDC-12; Southern Biotech),

mouse anti-human IgG (Jackson ImmunoResearch), tetanus toxoid from *Clostridium tetani* (List

Biological Laboratories), keyhole limpet hemocyanin (KLH; Sigma), ovalbumin (OVA; Sigma),

insulin (Sigma), and a panel of recombinant, tetrameric, head-only neuraminidase constructs

representing N1 A/California/07/2009; N1 A/Michigan/45/2015, X-275; N2 A/Aichi/2/1968; N2

A/Hong Kong/4801/2014, X263B; B/Phuket/3073/2013; B/Brisbane/60/2008.

Ab *V(D)J* rearrangement amplification and analysis

Rearranged V(D)J gene sequences were obtained from cultures of clonally expanded human

Bmem as described (37). V(D)J rearrangements were identified with Cloanalyst (38) and

IMGT/V-QUEST (39).

Recombinant IgG expression and purification

AbVec2.0-IGHG1, AbVec2.1-IGLC2-MscI, and AbVec1.1-IGKC plasmids were gifts from Hedda

Wardemann (RRID:Addgene_80795; RRID:Addgene_80796; RRID:Addgene_80797)(40).

Synthetic DNAs encoding Ab heavy-or light-chain variable domains were cloned into these

expression vectors, which harbor the constant regions of human IgG1, Igκ, or Igλ. IgGs were

produced by transient transfection of Expi293F cells with the Expifectamine 293 transfection kit

(Thermo Fisher), according to the manufacturer's instructions. Five days post-transfection,

supernatants were harvested, clarified by low-speed centrifugation, mixed 1:1 with Protein A

binding buffer, and incubated overnight with Pierce Protein A agarose resin (Thermo Fisher).

The resin was collected in a chromatography column, washed with binding buffer, eluted in

Pierce IgG Elution Buffer (Thermo Fisher), neutralized by one-tenth volume of 1M Tris (pH 9),

and dialyzed into PBS plus 0.1% sodium azide. IgG concentrations were determined with a

NanoDrop spectrophotometer (Thermo Fisher).

Generation of NA-expressing K530 cell lines

pMD2.G (Addgene plasmid #12259) and psPAX2 (Addgene plasmid #12260) were gifts from

Didier Trono. Codon-optimized, NA-encoding DNA sequences were cloned into the pLB-EXIP

lentiviral transfer plasmid (23). Influenza strain designations and abbreviated names are shown

in Table 1. Transfer plasmid, pMD2.G and psPAX2 were co-transfected into Lenti-X 293T cells

to produce lentivirus. NA-expressing lentiviruses were used to transduce K530 cell lines

expressing unique combinations of EBFP2, mTurquoise2, mNeonGreen, and mCardinal (23).

Transduced K530 cells were cultured for one week, then labeled with 1G01 (18) or DA03E17

(15) human IgG1 (2 μg/ml), washed, and labeled with goat anti-human IgG-PE (Southern

Biotech; 2 μg/ml). Individual cells expressing high levels of NA were identified by flow cytometry

and then sorted into 96-well plates containing growth medium. Subclones were expanded for

~10 days, then re-analyzed by flow cytometry to identify subclones with uniform, high

expression of NA. For each cell line, one rapidly growing subclone with high NA expression was

selected for use in all subsequent experiments. Aside from selecting subclones with high NA

expression, no effort was made to select cell lines with comparable expression levels of NA. For

long-term storage, K530 cell lines suspended in 90% FBS plus 10% DMSO and cryo-preserved

in liquid nitrogen.

Flow cytometry analysis of rIgG binding to K530 cell lines

Pooled K530-NA cell lines were thawed from cryopreserved aliquots and expanded in culture for

≥3 days. Pooled K530-NA cells were incubated at room temperature (RT) for 25-30 min with 2

μg/ml rIgGs diluted in IMDM plus 10% FBS. Alternately, cells were incubated with culture

supernatants containing clonal human IgG (diluted to 1:20 final). After washing, cells were

labeled with 2 μg/ml PE-conjugated goat anti-human IgG (Southern Biotech) for 20-30 min at

RT. Cells were then washed, stained with propidium iodide to identify dead cells, and analyzed

with a BD FACSymphony A5 flow cytometer.

NA-Star assay

NA activity was determined with the NA-Star Influenza Neuraminidase Inhibitor Resistance

Detection Kit (Invitrogen). K530-NA cells were suspended in calcium-containing assay buffer

(Hank's Balanced Salt Solution [Gibco] plus 0.5% BSA) and dispensed (1×10^5 cells/25 μl/well)

into opaque, white, 96-well microtiter plates (Thermo Fisher). Recombinant IgGs were serially

diluted in assay buffer and added (25 μl/well) to the cells, mixed, and incubated at room

temperature for 30 min. NA-Star substrate was diluted 1:1,000 in assay buffer, added to each

well (10 μl/well), and incubated with the cell-antibody mix for 20 min at room temperature. After

adding NA-Star Accelerator solution (60 μl/well), chemiluminescence was detected immediately

with a Tecan Spark plate reader, using an integration time of 1 s per well. Inhibition of NA

activity was calculated as the percentage of residual NA activity relative to wells containing no

inhibitor.

FOOTNOTES

Abbreviations: Ab, antibody; Bmem, memory B cell; IAV, influenza A virus; IBV, influenza B virus; HA, hemagglutinin; NA, neuraminidase; rIgG, recombinant IgG; BSA, bovine serum albumin; KLH, keyhole limpet hemocyanin; OVA, chicken ovalbumin; TT, tetanus toxoid; mAb, monoclonal antibody

Data availability

V(D)J sequences for mAbs from donors T1, T2, and T3 are available at GenBank (www.ncbi.nlm.nih.gov/Genbank), accession numbers PQ818731-PQ818752.

Declaration of competing interests

EBW has received research funding from Pfizer, Moderna, Seqirus, Najit Technologies, and Clinetic for the conduct of clinical research studies. He has also received support as an advisor to Vaxcyte and Pfizer, as a consultant to ILiAD Biotechnologies, and as DSMB member for Shionogi. The other authors have no competing interests to declare.

CRediT Statement: Joel Finney: Conceptualization, Investigation, Visualization, Writing – Original Draft, Writing – Review and Editing; **Masayuki Kuraoka**: Investigation, Writing – Review and Editing; **Shengli Song**: Resources, Writing – Review and Editing; **Akiko Watanabe**: Investigation**; Xiaoe Liang**: Investigation**; Dongmei Liao**: Investigation**; M. Anthony Moody**: Resources, Writing – Review and Editing; Funding Acquisition**; Emmanuel B. Walter**: Resources**; Stephen C. Harrison**: Writing – Review and Editing; Funding Acquisition**; Garnett Kelsoe**: Conceptualization, Supervision, Writing – Review and Editing; Funding Acquisition

ICMJE criteria for authorship

All authors attest they meet the ICMJE criteria for authorship.

REFERENCES

1. F. Krammer *et al.*, Influenza. *Nat Rev Dis Primers* **4**, 3 (2018).
2. S. Caini *et al.*, Probable extinction of influenza B/Yamagata and its public health implications: a systematic literature review and assessment of global surveillance databases. *Lancet Microbe* **5**, 100851 (2024).
3. A. S. Monto, M. Zambon, J. P. Weir, The End of B/Yamagata Influenza Transmission - Transitioning from Quadrivalent Vaccines. *N Engl J Med* **390**, 1256-1258 (2024).
4. T. Watanabe, S. Watanabe, E. A. Maher, G. Neumann, Y. Kawaoka, Pandemic potential of avian influenza A (H7N9) viruses. *Trends Microbiol* **22**, 623-631 (2014).
5. Y. T. Li, M. Linster, I. H. Mendenhall, Y. C. F. Su, G. J. D. Smith, Avian influenza viruses in humans: lessons from past outbreaks. *Br Med Bull* **132**, 81-95 (2019).
6. J. W. Drake, Rates of spontaneous mutation among RNA viruses. *Proc Natl Acad Sci U S A* **90**, 4171-4175 (1993).
7. S. Cobey, S. E. Hensley, Immune history and influenza virus susceptibility. *Curr Opin Virol* **22**, 105-111 (2017).
8. F. Krammer, P. Palese, Advances in the development of influenza virus vaccines. *Nat Rev Drug Discov* **14**, 167-182 (2015).
9. C. Gerdil, The annual production cycle for influenza vaccine. *Vaccine* **21**, 1776-1779 (2003).
10. R. Nachbagauer *et al.*, A chimeric hemagglutinin-based universal influenza virus vaccine approach induces broad and long-lasting immunity in a randomized, placebo-controlled phase I trial. *Nat Med* **27**, 106-114 (2021).
11. C. P. Arevalo *et al.*, A multivalent nucleoside-modified mRNA vaccine against all known influenza virus subtypes. *Science* **378**, 899-904 (2022).
12. A. S. Monto *et al.*, Antibody to Influenza Virus Neuraminidase: An Independent Correlate of Protection. *J Infect Dis* **212**, 1191-1199 (2015).
13. A. Madsen *et al.*, Human Antibodies Targeting Influenza B Virus Neuraminidase Active Site Are Broadly Protective. *Immunity* **53**, 852-863 e857 (2020).
14. C. Momont *et al.*, A pan-influenza antibody inhibiting neuraminidase via receptor mimicry. *Nature* **618**, 590-597 (2023).
15. A. Yasuhara *et al.*, A broadly protective human monoclonal antibody targeting the sialidase activity of influenza A and B virus neuraminidases. *Nat Commun* **13**, 6602 (2022).
16. J. Lederhofer *et al.*, Protective human monoclonal antibodies target conserved sites of vulnerability on the underside of influenza virus neuraminidase. *Immunity* **57**, 574-586 e577 (2024).
17. R. Lei *et al.*, Leveraging vaccination-induced protective antibodies to define conserved epitopes on influenza N2 neuraminidase. *Immunity* **56**, 2621-2634 e2626 (2023).
18. D. Stadlbauer *et al.*, Broadly protective human antibodies that target the active site of influenza virus neuraminidase. *Science* **366**, 499-504 (2019).
19. X. Xu, X. Zhu, R. A. Dwek, J. Stevens, I. A. Wilson, Structural characterization of the 1918 influenza virus H1N1 neuraminidase. *J Virol* **82**, 10493-10501 (2008).
20. I. Margine, P. Palese, F. Krammer, Expression of functional recombinant hemagglutinin and neuraminidase proteins from the novel H7N9 influenza virus using the baculovirus expression system. *J Vis Exp* 10.3791/51112, e51112 (2013).
21. H. Wan *et al.*, Structural characterization of a protective epitope spanning A(H1N1)pdm09 influenza virus neuraminidase monomers. *Nat Commun* **6**, 6114 (2015).
22. J. Finney *et al.*, Protective human antibodies against a conserved epitope in pre- and postfusion influenza hemagglutinin. *Proc Natl Acad Sci U S A* **121**, e2316964120 (2024).

23. S. Song *et al.*, A cell-based multiplex immunoassay platform using fluorescent protein-barcoded
reporter cell lines. *Commun Biol* **4**, 1338 (2021).
24. S. Song *et al.*, Allo-Specific Humoral Responses: New Methods for Screening Donor-Specific
Antibody and Characterization of HLA-Specific Memory B Cells. *Front Immunol* **12**, 705140
(2021).
25. K. R. McCarthy *et al.*, Memory B Cells that Cross-React with Group 1 and Group 2 Influenza A
Viruses Are Abundant in Adult Human Repertoires. *Immunity* **48**, 174-184 e179 (2018).
26. D. V. da Silva, J. Nordholm, U. Madjo, A. Pfeiffer, R. Daniels, Assembly of subtype 1 influenza
neuraminidase is driven by both the transmembrane and head domains. *J Biol Chem* **288**, 644-
653 (2013).
27. M. Dai *et al.*, Identification of Residues That Affect Oligomerization and/or Enzymatic Activity of
Influenza Virus H5N1 Neuraminidase Proteins. *J Virol* **90**, 9457-9470 (2016).
28. D. J. Bucher, E. D. Kilbourne, A 2 (N2) neuraminidase of the X-7 influenza virus recombinant:
determination of molecular size and subunit composition of the active unit. *J Virol* **10**, 60-66
(1972).
29. X. M. Luo *et al.*, Engineering human hematopoietic stem/progenitor cells to produce a broadly
neutralizing anti-HIV antibody after in vitro maturation to human B lymphocytes. *Blood* **113**,
1422-1431 (2009).
30. K. Y. Su, A. Watanabe, C. H. Yeh, G. Kelsoe, M. Kuraoka, Efficient Culture of Human Naive and
Memory B Cells for Use as APCs. *J Immunol* **197**, 4163-4176 (2016).
31. J. R. Whittle *et al.*, Broadly neutralizing human antibody that recognizes the receptor-binding
pocket of influenza virus hemagglutinin. *Proc Natl Acad Sci U S A* **108**, 14216-14221 (2011).
32. H. Xu *et al.*, Key mutations stabilize antigen-binding conformation during affinity maturation of a
broadly neutralizing influenza antibody lineage. *Proteins* **83**, 771-780 (2015).
33. A. G. Schmidt *et al.*, Preconfiguration of the antigen-binding site during affinity maturation of a
broadly neutralizing influenza virus antibody. *Proc Natl Acad Sci U S A* **110**, 264-269 (2013).
34. D. D. Raymond *et al.*, Influenza immunization elicits antibodies specific for an egg-adapted
vaccine strain. *Nat Med* **22**, 1465-1469 (2016).
35. A. Watanabe *et al.*, Antibodies to a Conserved Influenza Head Interface Epitope Protect by an
IgG Subtype-Dependent Mechanism. *Cell* **177**, 1124-1135 e1116 (2019).
36. A. Watanabe *et al.*, Self-tolerance curtails the B cell repertoire to microbial epitopes. *JCI Insight*
4 (2019).
37. M. Kuraoka *et al.*, Infant Antibody Repertoires during the First Two Years of Influenza
Vaccination. *mBio* **13**, e0254622 (2022).
38. T. B. Kepler, Reconstructing a B-cell clonal lineage. I. Statistical inference of unobserved
ancestors. *F1000Res* **2**, 103 (2013).
39. X. Brochet, M. P. Lefranc, V. Giudicelli, IMGT/V-QUEST: the highly customized and integrated
system for IG and TR standardized V-J and V-D-J sequence analysis. *Nucleic Acids Res* **36**, W503-
508 (2008).
40. T. Tiller *et al.*, Efficient generation of monoclonal antibodies from single human B cells by single
cell RT-PCR and expression vector cloning. *J Immunol Methods* **329**, 112-124 (2008).

Table 1. List of fluorescence-barcoded cell lines expressing recombinant, membrane-anchored NAs

Fluorescent barcode[a]	Option 1		Option 2	
	Source of NA	Abbreviated NA name	Source of NA	Abbreviated NA name
0000	–	–	–	–
0001	A/Solomon Islands/03/2006 (H1N1)	N1.SI06	A/Vietnam/1204/2004 (H5N1)	N1.VN04
0010	A/Sydney/5/2021 (H1N1)	N1.SYD21	–	–
0011	A/Hong Kong/4801/2014 (H3N2)	N2.HK14	–	–
0100	A/Hong Kong/1/1968 (H3N2)	N2.HK68	–	–
0101	A/USSR/90/1977 (H1N1)	N1.USSR77	A/Japan/305/1957 (H2N2)	N2.JP57
0110	A/Michigan/45/2015 (H1N1)	N1.MI15	A/Hong Kong/1144/1999 (H3N2)	N2.HK99
0111	A/Bilthoven/1761/1976 (H3N2)	N2.BH76	A/Darwin/9/2021 (H3N2)	N2.DW21
1000	A/Kansas/14/2017 (H3N2)	N2.KS17	A/Jiangsu/428/2021 (H10N3)	N3.JS21
1001	A/Memphis/4/1987 (H1N1)	N1.ME87	–	–
1010	B/Brisbane/60/2008 (Victoria)	NB.BN08	–	–
1011	A/California/07/2009 (H1N1)	N1.CA09	–	–
1100	A/Brisbane/8/1996 (H3N2)	N2.BN96	A/Shanghai/02/2013 (H7N9)	N9.SH13
1101	A/Beijing/353/1989 (H3N2)	N2.BJ89	–	–
1110	A/Perth/16/2009 (H3N2)	N2.PE09	A/tern/Australia/G70C/1975 (H11N9)	N9.AU75
1111	B/Phuket/3073/2013 (Yamagata)	NB.PK13	–	–

Notes:

[a] Expression of a fluorescent protein is indicated with a "1", whereas "0" indicates the protein is not expressed. The "ones" place (0001) denotes expression of eBFP2, the "tens" place (0010) denotes mTurquoise, the "hundreds" place (0100) denotes mNeonGreen, and the "thousands" place (1000) denotes mCardinal.

Table 2. IC$_{50}$ values for inhibition of sialidase activity from K530-NA cell lines

NA strain	1G01 IC$_{50}$ (µg/ml)	1G05 IC$_{50}$ (µg/ml)
N1.CA09	5.4	-
N1.MI15	2.6	-
N1.SYD21	1.1	-
N1.VN04	0.9	-
N2.JP57	1.6	-
N2.HK68	1.0	-
N2.HK99	4.5	-
N2.HK14	0.7	-
N2.DW21	>50	-
N3.JS21	0.9	-
N9.AU75	5.5	-
N9.SH13	3.6	-
NB.BN08	-	0.3
NB.PK13	-	0.4

Fig. 1

Human IgG-PE

Fig. 2

A

Donor T3: Luminex screen

n = 1,847 IgGs

- P4A8
- P10F5
- P15C5
- P17F8
- P27B8

B

Donor T3: Flow cytometric screen with K530-NA cells

n = 1,847 IgGs

- P4A8
- P10F5
- P15C5
- P17F8
- P20D5
- P27B8
- P28D3
- P35E4

Fig. 3

Fig. 4