1

Recessive but damaging alleles of muscle-specific ribosomal protein gene *RPL3L* drive neonatal dilated cardiomyopathy

Michael R. Murphy[1,2,#], Mythily Ganapathi[3], Teresa M. Lee[4], Joshua M. Fisher[4], Megha V. Patel[5,6], Parul Jayakar[7], Amanda Buchanan[8], Rajesh K. Soni[9], Yue Yin[1,2,10], Feiyue Yang[1,2], Muredach P. Reilly[1], Wendy K. Chung[11], & Xuebing Wu[1,2,#]

[1]Division of Cardiology, Department of Medicine, Columbia University Irving Medical Center, New York, NY 10032, USA

[2]Department of Systems Biology, Columbia University Irving Medical Center, New York, NY 10032, USA

[3]Department of Pathology and Cell Biology, Columbia University Irving Medical Center, New York, NY 10032, USA

[4]Department of Pediatrics, Columbia University Medical Center Irving Medical Center, New York, NY 10032, USA

[5]Department of Pediatrics, Nicklaus Children's Hospital, Miami, FL 33155, USA

[6]Current: Children's Wisconsin, Medical College of Wisconsin, Milwaukee, WI 53226, USA

[7]Division of Genetics and Metabolism, Nicklaus Children's Hospital, Miami, FL 33155, USA

[8]Illumina Inc., San Diego, CA, USA

[9]Proteomics and Macromolecular Crystallography Shared Resource, Herbert Irving Comprehensive Cancer Center, Columbia University, New York, NY 10032, USA

[10]Barnard College, Columbia University, New York, NY 10027, USA

[11]Department of Pediatrics, Boston Children's Hospital, Harvard Medical School, Boston, MA 02115, USA

[#]Correspondence to: mrm2267@cumc.columbia.edu (M.R.M); xuebing.wu@columbia.edu (X.W.)

Abstract

The heart employs a specialized ribosome in its muscle cells to translate genetic information into
proteins, a fundamental adaptation with an elusive physiological role. Its significance is
underscored by the discovery of neonatal patients suffering from often fatal heart failure caused
by rare compound heterozygous variants in RPL3L, a muscle-specific ribosomal protein that
replaces the ubiquitous RPL3 in cardiac ribosomes. *RPL3L*-linked heart failure represents the only
known human disease arising from mutations in tissue-specific ribosomes, yet the underlying
pathogenetic mechanisms remain poorly understood despite an increasing number of reported
cases. While the autosomal recessive inheritance pattern suggests a loss-of-function mechanism,
Rpl3l-knockout mice display only mild phenotypes, attributed to up-regulation of the ubiquitous
Rpl3. Interestingly, living human knockouts of *RPL3L* have been identified. Here, we report two
new cases of *RPL3L*-linked severe neonatal heart failure and uncover an unusual pathogenetic
mechanism through integrated analyses of population genetic data, patient cardiac tissue, and
isogenic cells expressing *RPL3L* variants. Our findings demonstrate that patient hearts lack
sufficient RPL3 compensation. Moreover, contrary to a simple loss-of-function mechanism often
associated with autosomal recessive diseases, *RPL3L*-linked disease is driven by a combination of
gain-of-toxicity and loss-of-function. Most patients carry a recurrent toxic missense variant
alongside a non-recurrent loss-of-function variant. The non-recurrent variants trigger partial
compensation of *RPL3* similar to *Rpl3l*-knockout mice. In contrast, the recurrent missense variants
exhibit increased affinity for the *RPL3/RPL3L* chaperone GRWD1 and 60S biogenesis factors,
sequester 28S rRNA in the nucleus, disrupt ribosome biogenesis, and trigger severe cellular toxicity
that extends beyond the loss of ribosomes. These findings elucidate the pathogenetic mechanisms
underlying muscle-specific ribosome dysfunction in neonatal heart failure, providing critical
insights for genetic screening and therapeutic development. Our findings also suggest that gain-
of-toxicity mechanisms may be more widespread in autosomal recessive diseases, especially for
those involving genes with paralogs.

Main

The heart is a remarkable organ, beating continuously throughout a human lifetime with a minimal need to replenish its muscle cells (cardiomyocytes)[1]. This extraordinary durability is attributed to many unique adaptations in cardiomyocytes. One of the most fundamental yet least understood adaptations is that cardiomyocytes use a specialized ribosome to decode genetic information and synthesize proteins[2-4]. Although ribosomes in all cells use the same genetic code, the composition of the ribosome itself—a delicate molecular machine comprising 80 ribosomal proteins and 4 ribosomal RNAs (rRNAs)—can differ across cell types[5-11]. The most striking variation occurs in cardiomyocytes. During vertebrate heart development, the ubiquitous core ribosomal protein L3 (RPL3), which makes the closest contact of any protein to the ribosome's RNA-based catalytic center[12], is gradually replaced by its paralog, RPL3-like (RPL3L)[2, 13]. RPL3L shares 74% amino acid identity with RPL3[2, 3] and eventually becomes the dominant form in adult human left ventricular cardiomyocytes (95%)[14], to a lesser extent in skeletal muscle (60%), and barely detectable in non-muscle cells[15]. Interestingly, this switch is reversed during adult muscle growth (hypertrophy) following injury or atrophy[3, 16, 17]. The physiological role of this highly conserved and tightly regulated ribosomal switching remains poorly understood, highlighting a crucial gap in our understanding of cardiac biology at the most fundamental level.

The essential role of RPL3L and the specialized ribosome in the heart is underscored by an increasing number of severe neonatal dilated cardiomyopathy (DCM) cases linked to biallelic variants in *RPL3L*[18-23]. This has been designated as a new DCM subtype, Dilated Cardiomyopathy-2D (CMD2D, OMIM# 619371). It stands out as most known DCM genes encode proteins associated with the sarcomere[24-26], the contractile unit of the muscle. *RPL3L*-linked DCM was the first—and remains the only—known human disease caused by mutations in tissue-specific ribosomes. Among seven independent affected families, individuals with mutations in both *RPL3L* alleles consistently develop severe DCM, often resulting in fatal heart failure[18-23]. In contrast, heterozygous parents and siblings carrying a single mutant allele remain unaffected with the early onset CMD2D. While the autosomal recessive inheritance pattern suggests a loss-of-function of these alleles and RPL3L deficiency as the likely cause of DCM, *Rpl3l* knockout (KO) mice generated by multiple labs display only mild cardiac dysfunction[4, 16, 27, 28]. This is unexpected given the 96% homology between human and mouse RPL3L proteins and that the highly regulated RPL3-RPL3L switch is also conserved in mice[16].

The discrepancy between early onset fatal disease in humans and mild phenotypes in mice suggests either species-specific differences in heart biology or that the DCM-causing autosomal recessive *RPL3L* mutations have effects beyond simple loss-of-function, rendering them unsuitable for modeling with KO mice. While rarely reported, autosomal recessive mutations can cause disease via toxic gain-of-function[29]. Notably, in all *Rpl3l* KO mouse models, regardless how they were generated—whether through a frameshifting deletion of the entire exon 2[4], short frameshifting deletions in exon 5[16, 27], or poly(A) insertion in exon 1[28]—there is a consistent up-regulation of the ubiquitous paralog Rpl3. This compensatory increase in Rpl3 likely explains the absence of a severe cardiac phenotype in mice, as Rpl3 is expected to largely fulfill the functional role of Rpl3l. The molecular mechanism underlying the *Rpl3* compensation remains unclear.

Whether a lack of *RPL3* compensation in human underlie the severe DCM symptoms remains
unexplored. A deeper understanding of the pathogenetic mechanism of *RPL3L*-linked DCM,
including whether *RPL3* compensation occurs and how to activate the compensation, may inform
future management and the development of new therapeutics for these patients.

To date, more than ten *RPL3L* mutations have been associated with neonatal DCM[18-23], with new
variants identified in every affected family. The relative importance of these variants in disease
remains unclear, as no functional characterizations have been performed. Here, we report two
new cases of *RPL3L*-linked DCM that along with previous cases highlight the existence of two
hotspot variants, suggestive of allele-specific pathogenetic mechanisms. We systematically
validated these hypotheses using patient tissue and isogenic cell lines expressing various *RPL3L*
variants.

Hotspot G27D and D308N/V are potential driver mutations

We identified two additional unrelated cases of *RPL3L*-associated early onset CMD2D, increasing
the total number of affected families to nine (Supplemental Information). Similar to previously
reported cases[18-23], both of the two families are associated with compound heterozygous variants,
i.e., each affected individual carry two different variants on separate alleles.
115
Interestingly, both cases share D308N as one of the variants: D308N/R58Q and
D308N/T340Nfs*25 (c.1080dup), respectively. A review of all the cases of *RPL3L*-linked DCM
reveals that while mutations are distributed across nearly the entire RPL3L protein (**Fig. 1a-b**), two
notable hotspots emerge: glycine at position 27 (G27) and aspartic acid at position 308 (D308) (**Fig.**
1a). Specifically, the G27D mutation is present in three of the nine families, while mutations at
D308 (D308N and D308V) appear in four additional families. These nine families originate from six
different countries and are presumed to be unrelated. For example, the three families carrying the
G27D variant are located in the United States, Spain, and China.
124
Although all *RPL3L* variants implicated in DCM are exceedingly rare in the general population ($<10^{-4}$,
gnomAD[30]), the two hotspot variants (G27D and D308N/V) have even lower allele frequencies
($<10^{-5}$) compared to other DCM-linked *RPL3L* variants (**Fig. 1c**), indicating stronger negative
selection and greater deleterious effects. Indeed, G27D and D308N/V are predicted to be highly
pathogenic by AlphaMissense[31], a state-of-the-art deep learning model that builds on the protein
structure prediction tool AlphaFold2[32]. The low allele frequencies of these mutations in the
general population further argue against the hypothesis that the recurrence of G27D and D308N/V
in patients is due to increased mutability of the DNA bases. The recurrent occurrence of these
ultra-rare missense mutations in a substantial proportion of DCM families strongly challenges a
loss-of-function mechanism, particularly given the high conservation of much of the protein and
the fact that neither G27 nor D308 is among the residues known to be critical for RPL3/RPL3L's
role in the ribosome. Together, G27D and D308N/V account for 78% (7/9) of *RPL3L*-linked DCM
families, positioning them as the primary driver mutations in the disease.

Notably, the ten missense variants can be divided into two groups when considering allele
frequency and pathogenicity predicted by AlphaMissense (**Fig. 1c**). In addition to the hotspot
mutations G27D and D308N/V, the high-pathogenicity group includes the V231F mutation,
another extremely rare (~10^{-6}) mutation in the population and predicted to be highly pathogenic
by AlphaMissense. It is possible that with more cases to be discovered, V231F may also be found
to be a hotspot mutation similar to G27D and D308N/V in the same group. Variants in the low-
pathogenicity group are more frequent (still very rare, 10^{-5} to 10^{-4}) and predicted to be less
pathogenic (**Fig. 1c**). Interestingly, with the exception of the homozygous R116H mutation
(identified in a consanguineous population[21]), each affected individual carries one mutation from
the high-pathogenicity group and another mutation from the low-pathogenicity group (linked by
dashed lines).

Consistent with the absence of a severe cardiac phenotype in *Rpl3l* KO mice, live human knockouts
of *RPL3L* have been identified in large-scale genetic studies, including gnomAD[30], All of Us[33], and
UK Biobank[34] (**Fig. 1d**). One individual carries a premature stop codon at arginine 226 in both
alleles (R226X), truncating nearly half of the protein (full-length RPL3L is 407 amino acids), a
mutation almost certainly resulting in loss-of-function. Additionally, a total of 17 individuals are
homozygous for a splice donor variant previously shown to cause exon 9 deletion, which removes
a highly conserved 40-amino-acid region in RPL3L[35]. Although this splice donor variant is
associated with an increased risk of atrial fibrillation (OR: 1.50) in a genome-wide association
study[35], homozygous individuals carrying this variant appear to be alive and do not exhibit early-
onset severe DCM. There are also nine putative knockout humans in the Regeneron Genetics
Center Million Exome (RGC-ME) database[36] although detailed information about these variants is
not available. The lack of severe cardiac dysfunction in both putative *RPL3L* KO humans and *Rpl3l*
KO mice strongly suggests that the missense *RPL3L* mutations identified in DCM patients are not
simply loss-of-function variants.

In summary, genetic evidence indicates that hotspot missense mutations at G27 and D308 in
RPL3L likely cause DCM through a mechanism other than loss-of-function, despite an autosomal
recessive inheritance pattern.

Ribosome defects in explanted patient hearts

To investigate the pathogenic mechanisms of *RPL3L* mutations in DCM, we initially focused on a
patient carrying the hotspot D308N allele and the T189M missense allele. This patient, diagnosed
and treated at our center, underwent heart transplantation and is alive to date[18]. Using explanted
heart ventricular tissues, we extracted total RNA to perform RNA-seq and compared the results to
age-matched control healthy heart ventricular tissues.

Unexpectedly, we observed an abnormality when the total RNA was subjected to Bioanalyzer
electrophoresis for quality check (**Fig. 2a**). Specifically, there is a drastic reduction in the 28S rRNA
peak relative to the 18S peak. The 28S rRNA is the scaffold of the 60S large subunit of the ribosome.
RPL3L (and the paralog RPL3) is a core component of the 60S subunit and plays a vital role in 60S
ribosome biogenesis via its interaction with the 28S rRNA[37]. The loss of 28S rRNA indicates a defect

in 60S ribosome biogenesis in patient cells, presumably due to the D308N/T189M mutations in
RPL3L.

The RNA-seq data revealed widespread alterations in gene expression, including 1,639 significantly
up-regulated genes and 2,361 significantly down-regulated genes (**Fig. 2b**). Pathway analyses
confirmed up-regulation of gene signatures associated with systolic heart failure and apoptosis,
and interestingly, a massive up-regulation of genes associated with fetal cardiac myeloid cells,
consistent with increased inflammation in failing hearts[38] (**Fig. 2c**).

In line with impaired ribosome biogenesis, many ribosomal protein genes are differentially
expressed, with more down-regulated than up-regulated (**Fig. 2b**). *RPL3L* is down-regulated by
approximately two-fold, although not significantly (adjusted P = 0.6). Both D308N and T189M
alleles are expressed at a similar level as indicated by sequencing read coverage (**Fig. 2d**).

Surprisingly, unlike the robust increase of *Rpl3* mRNA and protein in *Rpl3l* KO mice, there was no
compensatory up-regulation of *RPL3* mRNA in the patient's heart tissue (**Fig. 2b**). Instead, *RPL3* is
slightly down-regulated (by 24%, adjusted P=0.66) like many other ribosomal protein genes.

To confirm the lack of *RPL3* compensation in the DCM heart tissue, we also examined protein
expression using mass spectrometry. Indeed, when normalized to the total abundance of 40S
ribosomal proteins, there is a significant decrease in RPL3 abundance with no change in RPL3L (**Fig.
2e**). Consistent with a 60S biogenesis defect, there is a mild but significant decrease in total 60S
ribosomal protein level relative to that of the 40S. Similar to the RNA-seq data (**Fig. 2d**), the D308N
allele is expressed at a comparable level as the T189M allele (**Fig. 2f**).

In the absence of compensatory RPL3 up-regulation, there will be a shortage of functional 60S
ribosomes in patient cells, as neither functional RPL3L nor RPL3 is present. Given the critical role
of ribosomes in cellular function, the loss of functional ribosomes likely contributes to
cardiomyocyte death and the progression of DCM. While caveats remain (see Discussion), the lack
of RPL3 compensation observed in human DCM patients underscores a critical divergence from
the *Rpl3l* KO mouse model and suggests that at least one of the two missense *RPL3L* variants
(D308N and T189M) are unlikely to function as simple loss-of-function alleles.

In summary, analysis of explanted heart tissues from DCM patients demonstrates a defect in 60S
ribosome biogenesis and a lack of RPL3 compensation. These defects likely contribute to the loss
of functional ribosomes, cardiomyocyte death, and the development of heart failure in *RPL3L*-
linked DCM.

Hotspot variants gain toxic function to drive ribosome biogenesis defects and cellular toxicity

It remains unclear whether the ribosome biogenesis defect observed in explanted patient tissue
is caused by the hotspot D308N mutation or the accompanying T189M allele. Both alleles are
expressed at comparable levels with no significant allele imbalance (**Fig. 2d** and **Fig. 2F**).

To determine the individual impact of these alleles, we generated isogenic cell lines expressing
various RPL3L variants in the absence of RPL3 (**Fig. 3a**). These cell lines were derived from human
AC16 cells, ventricular cardiomyocyte-like cells commonly used to study cardiac gene expression
and function[39], including the regulation of ribosomes and translation in cardiac hypertrophy[40].
While AC16 cells do not fully recapitulate primary cardiomyocytes, our results described below
demonstrate their suitability for studying ribosome-related processes. Of note, *RPL3L* is not
expressed in iPSC-derived cardiomyocytes (iPSC-CMs) (**Extended Data Fig. 1**), consistent with the
notion that iPSC-CMs resemble immature fetal cardiomyocytes[41, 42], while *RPL3L* expression is
restricted to mature cardiomyocytes.

Wild-type AC16 cells naturally express only *RPL3*, not *RPL3L*. To switch to *RPL3L* variants, we first
introduced a doxycycline (dox)-inducible shRNA targeting *RPL3* to deplete RPL3 in an inducible
manner. This cell line (shRPL3-only) serves as a critical baseline control for the cellular defect and
toxicity caused by the loss-of-function of ribosomal proteins. Using shRPL3-only cells, we further
integrated a second dox-inducible transgene expressing HA-tagged *RPL3L*, either the wild type or
variants (**Fig. 3a**). In addition to DCM-linked *RPL3L* variants, we also included a common SNP
(rs34265469) predicted to result in a benign missense variant (P291L). This SNP is the most
common missense variant in *RPL3L*, with an allele frequency of 4.26% (TOPMed) and is
homozygous in at least 715 living individuals. This isogenic system enabled allele-specific switching
from *RPL3* to *RPL3L* at physiological level upon dox treatment (**Fig. 3b-c**).

Using these isogenic cell lines, we found that the loss of the 28S rRNA observed in the explanted
patient heart tissue (**Fig. 2a**) was faithfully recapitulated in cells expressing the D308N allele, but
not in those expressing the T189M allele (**Fig. 3d**). Notably, the loss of 28S rRNA in D308N cells is
much stronger than that of the shRPL3-only cells (**Fig. 3d**). Given that the only difference between
these two cell lines is the expression of the D308N variant, the stronger defect indicates toxic gain-
of-function of the D308N variant. Additionally, the weaker effect in the shRPL3-only cells suggests
that ribosome loss alone is insufficient to fully account for the defects observed in patients (**Fig.
2a**).

Consistent with 60S ribosomal subunit biogenesis defects, sucrose gradient sedimentation
revealed a dramatic reduction in 60S but not 40S ribosomal subunits in D308N-expressing cells
compared to WT, SNP, or T189M cells (**Fig. 3e**). Accordingly, monosomes (80S) and translating
polysomes (poly-ribosomes), both of which contain 60S, are also depleted in D308N-expressing
cells (**Fig. 3e**). Notably, the loss of polysomes in D308N-expressing cells is comparable to cells not
expressing any *RPL3L* variant (shRPL3-only) (**Fig. 3e**), suggesting the D308N variant of RPL3L almost
completely blocks ribosome biogenesis.

In line with a severe loss of ribosomes, D308N-expressing cells exhibited a severe growth defect
(**Fig. 3f**, note that the y-axis is on a logarithmic scale). Expressing D308N for 10 days led to a 40-
fold reduction in cell counts (compared to RPL3L-WT cells, **Fig. 3f**, green vs. red). Interestingly,
cells expressing the D308N variant exhibit a 6.4-fold greater growth defect compared to cells
lacking RPL3L expression (shRPL3-only) (green vs. blue, **Fig. 3f**), again suggesting toxic gain-of-
function. Of note, the control cell line expressing RPL3 (shScramble) grew faster than cells

expressing wild type RPL3L (RPL3L-WT), mirroring a previous mouse study showing that Rpl3l
expression is associated with reduced protein synthesis and slower muscle cell growth[3].
Interestingly, cells expressing the T189M variant of *RPL3L* grew faster than cells expressing the
wild type *RPL3L*, which may be attributed to a higher expression of *RPL3* in this cell line (**Fig. 3b**,
also see below the discussion on *RPL3* compensation). The lack of deleterious effects associated
with the T189M variant aligns with its low pathogenicity prediction by AlphaMissense (**Fig. 1c**) and
is further supported by the presence of a living homozygous T189M individual in the population
genotyped in TOPMed. Taken together, these results indicate that the D308N variant is a toxic
gain-of-function variant and the primary driver of the disease in the patient carrying D308N and
T189M.

Next, we investigated the other hotspot variant G27D, together with the R161W allele found in
the same affected individual[18]. Similar to the D308N variant, cells expressing the G27D variant
grew substantially slower than those expressing the wild-type *RPL3L* allele (33-fold fewer cells by
day 10, **Extended Data Fig. 2a**), whereas cells expressing the R161W variant grew slightly faster
than cells expressing the WT RPL3L, similar to T189M. Consistent with a growth defect, G27D but
not R161W-expressing cells are depleted of 60S and 80S ribosomes as well as polysomes
(**Extended Data Fig. 2b**). These findings confirm that the hotspot variants G27D and D308N, but
not their co-inherited non-hotspot variants T189M and R161W, are responsible for ribosome
biogenesis defects and cellular toxicity.

Hotspot variants mislocalize and alter interactions with ribosome biogenesis factors

To further explore the molecular basis of the toxic gain-of-function of the hotspot variants, we
examined the subcellular localization of RPL3L variants using fluorescence imaging of the HA tag.
In contrast to the predominantly cytoplasmic localization of the wild-type RPL3L and the common
SNP variant expected from a ribosomal protein, the hotspot variants D308N and G27D both
exhibited nearly exclusive nuclear localization (**Fig. 4a**, green, quantified in **Fig. 4b**). Given the role
of RPL3/RPL3L in 60S ribosome biogenesis, this mislocalization of the hotspot variants likely disrupt
60S ribosome assembly. Supporting this, fluorescent probes for 28S rRNA revealed a significant
decrease in cytoplasmic 28S rRNA levels in D308N or G27D-expressing cells relative to the nucleus
(**Fig. 4a**, red, quantified in **Fig. 4c**). Notably, this 28S rRNA localization defect was not observed in
cells not expressing any RPL3L variant (shRPL3-only), consistent with a toxic gain-of-function of
both hotspot variants.

To gain biochemical insights into the toxicity associated with the hotspot variants, we profiled the
interactomes of RPL3L variants by using immunoprecipitation against HA-tag followed by mass
spectrometry in triplicates. Given the exclusive nuclear localization of the hotspot variants, we
used nuclear lysate as input for all samples. Interestingly, principal component analysis (PCA) of
the proteomics data separates the two hotspot variants from the rest in the first principal
component (PC1), suggesting dramatic rewiring of the interactome for hotspot variants (**Fig. 4d**).
The second principal component (PC2) separates the three nuclear variants (G27D, D308N, and
R161W) from the two cytoplasmic variants (T189M and WT). Interestingly, the top two proteins
driving the separation of hotspot variants from other variants both exhibit increased affinity for

the hotspot variants, supporting the concept of toxic gain-of-function effects (**Fig. 4e**). These two
proteins, GRWD1 and C7ORF50, although less studied in humans, have yeast homologs (Rrb1 and
Rbp95, respectively) that directly interact with yeast Rpl3 during 60S ribosome biogenesis. GRWD1
serves as the dedicated chaperone for RPL3 (and likely RPL3L as well), binding RPL3 co-
translationally to prevent its aggregation and degradation, ensuring its proper delivery to the
assembly site on nucleolar pre-60S particles[43-45]. The other protein, C7ORF50, interacts with RPL3
both genetically and physically, binding adjacent to RPL3 on the 28S rRNA during early pre-60S
ribosome biogenesis[46-48]. Several other 60S biogenesis factors, including RRP15[49], EBNA1BP2[50],
DDX24[51], NOP56[47], and MRTO4[52] also exhibited enrichment in the nuclear pulldown of RPL3L
variants (**Extended Data Fig. 3**). The sequestration of these essential 60S ribosome biogenesis
factors by hotspot variants likely underlies the 60S biogenesis defect and contributes to the toxic
gain-of-function effects associated with these variants.

In summary, our subcellular localization analysis and nuclear protein interactome profiling using
isogenic cell lines reveal that the hotspot variants D308N and G27D mislocalize to the nucleus,
leading to nuclear retention of 28S rRNA. This toxic effect is likely driven by their enhanced binding
to the dedicated RPL3/RPL3L chaperone GRWD1 and 60S ribosome biogenesis factors, including
C7ORF50.

Non-hotspot variants drive post-transcriptional *RPL3* compensatory up-regulation

Intriguingly, the non-hotspot R161W allele, identified in the same affected individual as the
hotspot G27D, also mislocalizes to the nucleus (**Fig. 4a-b**). However, unlike the hotspot variants
and consistent with its lower predicted pathogenicity (**Fig. 1c**), R161W does not cause nuclear
sequestration of 28S rRNA (**Fig. 5a/c**), nor does it lead to a ribosome biogenesis defect (**Extended
Data Fig. 2a**) or cell toxicity (**Extended Data Fig. 2b**). Western blot analysis of the polysome
fractions confirmed the absence of R161W from these ribosomes (**Extended Data Fig. 4**).

To investigate how ribosomes can still function in the absence of cytoplasmic RPL3L, we examined
the expression of RPL3, which was expected to be silenced by an shRNA induced alongside the
R161W allele upon dox treatment (**Fig. 3a**). Unexpectedly, while *RPL3* remained silenced in cells
expressing most *RPL3L* alleles, its expression was almost fully restored in cells expressing the
R161W allele, and to a lesser extent in those expressing the T189M allele, at both the mRNA (**Fig.
5a**) and protein (**Fig. 5b**) levels. Importantly, there was no significant change in the expression of
the *RPL3*-targeting shRNA (**Extended Data Fig. 5**), ruling out the possibility that *RPL3* recovery was
due to loss of shRNA-mediated knockdown in cells expressing R161W or T189M. These results
suggest that, similar to *Rpl3l*-KO mice, the expression of the non-hotspot *RPL3L* alleles R161W and
T189M leads to compensatory up-regulation of *RPL3*, suggesting that these alleles are functionally
equivalent to a knockout of *RPL3L*. Supporting a model in which compound heterozygotes carry a
toxic hotspot variant alongside a loss-of-function variant, frameshifting variants were identified in
three of the seven families with hotspot variants (**Fig. 1a**).

To elucidate the mechanism behind this *RPL3* compensation, we first examined the transcriptional
activity at the *RPL3* promoter using chromatin immunoprecipitation (ChIP) against RNA

polymerase II (Pol II). No significant differences in transcriptional activity were observed between
cells expressing G27D and R161W (**Fig. 5c**). Additionally, qPCR measurements of nascent
transcription from intronic RNA showed no significant differences (**Extended Data Fig. 6**), ruling
out transcriptional up-regulation as the mechanism of *RPL3* compensation.

We then assessed the stability of *RPL3* mRNA by inhibiting global transcription with the RNA Pol II
inhibitor α-amanitin and measuring the decay of *RPL3* mRNA over time. Compared to cells
expressing the G27D allele, *RPL3* mRNA was significantly more stable in cells expressing the
R161W allele (**Fig. 5d**, P < 0.05). Consistent with a post-transcriptional mechanism, the increase in
steady-state *RPL3* level was primarily due to increased cytoplasmic *RPL3*, with no significant
change in the nuclear fraction (**Fig. 5e**).

In summary, unlike the two hotspot alleles, G27D and D308N, which exhibit a gain-of-toxic
function, the non-hotspot alleles R161W and T189M resemble a knockout effect, promoting
compensatory upregulation of *RPL3* by stabilizing its mRNA.

Impaired protein synthesis in engineered compound heterozygous cells

RPL3L-linked DCM patients often carry a combination of a hotspot allele and a non-hotspot allele,
such as the D308N/T189M and G27D/R161W pairs we have characterized. In our analysis of
explanted heart tissue from patients carrying the D308N/T189M alleles, both alleles were
detected at the RNA (**Fig. 2d**) and protein levels (**Fig. 2f**). Given the opposing effects of the hotspot
and non-hotspot alleles on ribosome biogenesis and cell growth (**Fig. 3-5**), we sought to explore
the impact of these alleles when both are present in the same cell, as is common in most DCM
patients.

To address this, we introduced the G27D allele into cells expressing the R161W allele, creating a
model of compound heterozygous cells that mirror those found in DCM patients. We chose R161W
over T189M because R161W triggers a stronger compensatory up-regulation of *RPL3* (**Fig. 5a-b**).
Using a puromycin incorporation assay to measure nascent protein synthesis, we observed a
significant reduction in protein synthesis rates in the compound heterozygous cells compared to
those expressing R161W alone, though the rate was still higher than in cells expressing G27D alone
(**Fig. 6a-b**). This reduced protein synthesis rate aligns with the observed detrimental effects of
compound heterozygous *RPL3L* alleles in humans.

Interestingly, *RPL3* compensation remained unaffected in the compound heterozygous cells
compared to cells expressing R161W alone (**Fig. 6c**). This suggests that the compensatory up-
regulation of *RPL3* is not sufficient to overcome the toxic effects of the G27D allele.

Discussion

In this study, we reported two new cases of severe neonatal heart failure caused by mutations in
the muscle-specific ribosomal protein RPL3L and uncovered a complex mechanism by which these
mutations cause disease. Our study revealed a pattern hidden in the large number of variants
found in patients. Specifically, a small number of recurrent, toxic variants co-occur with likely loss-

of-function variants in each affected individual. Our study suggests an unexpected pathogenetic
mechanism—autosomal recessive mutations exerting gain-of-toxic function other than merely
loss-of-function effects—underlie the severe heart defects in humans and may also explain the
lack of phenotype in *Rpl3l* KO mice. Here we discuss the significance of our discovery as well as
questions remain to be addressed.

DCM is a leading cause of heart failure, with nearly half of all cases—and two-thirds of pediatric
cases—remaining of unknown etiology[24, 25, 53, 54]. This underscores the urgent need to identify and
understand novel pathogenic mechanisms to improve management and treatment. The advent of
accessible human exome sequencing and whole-genome sequencing has led to a renaissance in
the discovery of causal disease genes for DCM, including in pathways not previously known to have
specific cardiac function[55]. This itself has resulted in a lag between causal gene discovery and
research underlying disease mechanisms[56]. In the case of *RPL3L*, there has been no functional
studies to understand how mutations on the cardiac ribosome cause DCM. Here, we address this
gap by experimentally characterizing multiple *RPL3L* mutations, uncovering their impact on
ribosome biogenesis, protein synthesis, and cell viability.

Our findings suggest that rather than a simple loss-of-function mechanism, these autosomal
recessive missense mutations in *RPL3L* cause disease via a combination of gain-of-toxicity and loss-
of-function mechanism, supported by both genetic and biochemical evidence. Hotspot variants in
rare diseases are more commonly associated with toxic effects or even dominant-negative traits[57-
59]. Despite their rarity in the general population, missense mutations G27D and D308N/V occur in
seven out of nine unrelated DCM patient families. Mechanistically, hotspot variants of RPL3L
mislocalize to the nucleus and disrupt ribosome biogenesis when expressed in human cells,
recapitulating the 28S rRNA loss observed in explanted patient hearts carrying the same mutation.
Mislocalization of hotspot variants also caused 28S rRNA retention in the nucleus, a defect not
seen with RPL3 knockdown, indicating toxicity beyond loss-of-function. This toxic effect is likely
caused by enhanced binding to GRWD1 and other ribosome biogenesis factors. Ribosome
biogenesis defects result in the loss of global protein synthesis and reduced cell viability, again
beyond the simple loss of ribosomes. Future studies, especially *in vivo* studies will be needed to
elucidate how these molecular and cellular defects lead to organ level failure.

Autosomal recessive mutations in Mendelian disorders are typically considered loss-of-function
mutations[60]. Rarely do they cause disease via gain-of-toxicity or dominant-negative mechanisms.
The only prior example known is the A673V missense mutation in the β-amyloid (Aβ) precursor
protein (APP), associated with familial Alzheimer's disease[29]. In this case, the mutant allele in the
homozygous state enhances Aβ peptide production, promoting the formation of amyloid fibrils,
the primary trigger of Alzheimer's disease. In heterozygous individuals, however, the coexistence
of wild-type and mutant Aβ peptides inhibits amyloidogenesis relative to either mutant or wild-
type Aβ peptide alone. This inhibition is likely due to conformational incompatibility between the
wild-type and mutant Aβ peptides. For *RPL3L*-linked heart failure, there are several possibilities
for why heterozygous members of families carrying the recurrent and toxic variant (e.g., G27D and
D308N/V) have not developed severe disease. The presence of the wild type RPL3L protein will
likely reduce the toxicity and result in incomplete penetrance or much delayed onset of the disease.

One plausible mitigation mechanism is that these RPL3L hotspot variants form aggregates similar
to Aβ, with wild-type RPL3L in heterozygous cells preventing such aggregation and mitigating the
pathogenic effects.

Compensation by the paralog RPL3, triggered by RPL3L loss-of-function, likely explains why severe
disease arises only from a combined gain-of-toxicity and loss-of-function mechanism, rather than
loss-of-function alone. Genetic compensation by paralogs is well-documented, although the
underlying molecular mechanisms often remain unclear[61-63]. The increased expression of *Rpl3*
observed in *Rpl3l* KO mice has been proposed to explain the absence of severe cardiac phenotypes
in these animals, but the molecular basis of this compensation is not yet understood. In this study,
we demonstrate that non-hotspot missense mutations, such as R161W, also up-regulate *RPL3*
expression in human cells. This up-regulation occurs through stabilization of cytoplasmic *RPL3*
mRNA without transcriptional activation. However, the *RPL3* compensation induced by the R161W
variant does not fully counteract the toxic effects of the G27D variant when co-expressed in
human cells. These findings suggest that enhancing *RPL3* compensation alone may be insufficient
as a therapeutic strategy for patients carrying toxic *RPL3L* mutations.

We observed no significant *RPL3* compensation in failing hearts explanted from human DCM
patients with *RPL3L* mutations, in stark contrast to the robust up-regulation of *Rpl3* observed in
Rpl3l KO mice. This difference may help explain the discrepancy in disease manifestation between
humans and mice. However, our analysis has several limitations. The comparison was made
between DCM patient ventricular tissues and three age-matched healthy controls, which showed
considerable variation in *RPL3* expression. Moreover, these controls were from individuals without
cardiomyopathy or other cardiac conditions. Since the RPL3/RPL3L switch is known to be regulated
during muscle hypertrophy, it remains challenging to determine whether changes in *RPL3*
expression—or the lack thereof—are a cause or consequence of DCM. Future studies, including
mouse models carrying human-specific mutations, will be essential to confirm these findings and
further elucidate the underlying mechanisms.

In conclusion, our systematic genetic and functional analyses provide new insights into the
pathogenesis of heart failure caused by mutations in muscle-specific ribosomes, uncovering a
novel mechanism involving gain-of-toxic effects of recessive mutations. These findings lay the
groundwork for understanding other *RPL3L* variants and offer a foundation for developing
targeted therapies. Our findings also suggest that gain-of-toxicity mechanisms may be more
widespread in autosomal recessive diseases involving genes with paralogs.

Acknowledgments

We thank members of the Wu lab and the Cardiometabolic Genomics Program for discussion and
comments on the manuscript. This work is supported by NIH/NHLBI grant 1R01HL171664-01 and
Pershing Square Foundation MIND Prize (X.W.) and a postdoctoral fellowship from American Heart
Association (M.R.M). This research was funded in part through the NIH/NCI Cancer Center Support
Grant P30CA013696 and used the Genomics and High Throughput Screening Shared Resource and

CCTI Flow Cytometry Core. The CCTI Flow Cytometry Core is supported in part by the Office of the
Director, National Institutes of Health under awards S10OD030282 and S10OD020056.

Author contributions

X.W. and M.R.M. conceived the functional study and wrote the manuscript with input from all
authors. M.R.M. performed the majority of experiments. M.G. coordinated case reports and
sample collection. T.M.L. and J.M.F. identified and characterized the D308N/T340Nfs*26 case.
M.V.P., A.B., and P.J. identified and characterized the D308N/R58Q case. R.K.S. generated the
proteomics data. Y.Y. assisted in imaging. W.K.C. contributed the D308N/T189M tissue sample.
W.K.C., M.P.R., and F.Y. edited the manuscript and provided critical feedback.

Declaration of interests

459 X.W. is a member of the Scientific Advisory Board for Epitor Therapeutics. W.K.C. serves on the
460 Board of Directors at Prime Medicine.

Materials and Methods

Cell Culture, Drugs and Transfection

AC16 cells were cultured at 37°C, 5% CO_2 in low-glucose DMEM (Fisher; 11-885-084)
supplemented with 10% FBS. HEK293T cells were cultured in DMEM with 10% FBS. For
immunofluorescence assays, 2 x 10^5 cells were plated in 6 well plates and incubated for 24 hours.
For stable cell generation, EZ-Tet-RPL3-shRNA was co-transfected with CMV-dR8.91 and MD2.G
plasmids for 72 hours in HEK293T cells to produce virus. The media was centrifuged for 5 minutes
at RT, filtered using 0.45µm Syringe Filter Unit (Fisher), and 200µl viral media was added to 2 x 10^5
AC16 cells for 48 hours prior to selection using 200µg/ml Hygromycin B until control cells (non-
transduced) were dead. For stable generation of pLenti-RPL3L, procedure was followed as above
except transduction was carried out on EZ-Tet-RPL3-shRNA/AC16 cells and selection was carried
out by sorting for BV421-positive cells. For mRNA stability assay, AC16 cells were treated with
50mg/ml a-amanitin for the indicated time points.

Plasmids

To generate plasmid expressing shRNA against RPL3, a reported high efficiency shRNA sequence
for RPL3 was identified (Sigma) and ~70nt single stranded oligos were ordered from IDT and
annealed by heating to 95°C and cooled at a rate of 5°C min^{-1}. Double-stranded oligos had two 5'
overhangs corresponding to NheI and EcoRI sites, respectively. dsDNA oligos were ligated to EZ-
Tet-pLKO-Hygro (Addgene; 85972) using Quick Ligation Kit (NEB), producing Tet-inducible RPL3
shRNA vector. The pLenti-RPL3L plasmid was derived from pLentiRNACRISPR_007 (Addgene;
138149). A gene fragment containing the wild-type human RPL3L sequence with a C-terminal HA
tag was delivered on pUC57 (Genewiz). PCR products for wild-type RPL3L-HA derived from pUC57,
the pLentiRNACRISPR_007 backbone, and EBFP (unpublished in-house plasmid) with overlapping
ends were assembled via Gibson cloning using NEBuilder HiFi DNA Assembly Master Mix (NEB),

| 487 | producing Dox-inducible RPL3L and constitutive EBFP expression. To obtain RPL3L mutation |

producing Dox-inducible RPL3L and constitutive EBFP expression. To obtain RPL3L mutation plasmids, primers (Supplementary Table) were designed against pUC57-RPL3L-HA with single nucleotide substitutions in the 5' region of the forward primer sequence. Mutagenesis was performed using Site Directed Mutagenesis Kit (NEB) as per the manufacturer's instructions. pLenti-RPL3L mutations were derived similarly to the wild-type as above.

Immunofluorescence and FISH

15,000 cells were seeded on Lab-Tek II (Thermo) chamber slides and grown overnight. Cells were washed once with PBS and fixed in 300µl PFA at room temperature for 15 minutes. Fixed wells were washed 3x with PBS for 3 minutes, permeabilized in 0.5% Triton X-100 in PBS for 15 minutes, washed 3x, and incubated in 3% BSA in PBS for 1 hour to block. Cells were then incubated in 3% BSA containing primary antibody overnight at 4^{o}C, washed 3x in PBS, then incubated for 1 hour at room temperature in secondary antibody protected from light. After washing 3x, cells were incubated in 1:1000 DAPI solution for 5 minutes, washed 3x, then imaged using a Nikon Eclipse Ti Series confocal microscope equipped with 60x lens performed at Columbia University's Confocal and Specialized Microscopy Core. RPL3L was detected using anti-HA (Thermo; 2-2.2.14). Secondary staining was accomplished using goat anti-rabbit IgG (H+L) Alexa Fluor 488 (Invitrogen) and goat anti-mouse IgG (H+L) Alexa Fluor 594 (Invitrogen). For rRNA FISH, cells were fixed as above but permeabilized with 70% ethanol for at least 1 hr. Custom designed 28S rRNA FISH probes (Supplementary Table) were designed conjugated to CAL Fluor® Red 590 dye (Stellaris) and incubated with fixed samples for 16 hours in the dark. After washing using wash buffers A + B (Stellaris), samples were incubated with primary anti-HA antibody as above.

Polysome fractionation

Sucrose gradient density ultracentrifugation was used to separate the monosomal and polysomal ribosomal fractions. For human cell experiments, AC16 cells were grown in 15cm dish were lysed by adding 800µl of ice-cold lysis buffer (20mM HEPES, 150mM KCl, 10mM MgCl$_2$), plus 0.5% NP-40, 1x EDTA-free Protease Inhibitor Cocktail (PIC; Roche)), 2.5mM DTT, 25 U ml^{-1} Turbo DNase I (Invitrogen)), 10U SuperaseIN (Thermo), 2mM Vanadyl Ribonucleoside Complex (NEB) and 100ug/ml CHX (Sigma) directly to the plate and scraping. Lysed material was transferred to 1.5ml centrifuge tube then triturated 10x with 23-gauge syringe. Samples were incubated in 4ºC for 30 minutes with rotation to complete lysis, then centrifuged at 14,000g for 7 minutes. Supernatant was transferred to new tube and 700ul of lysate was loaded onto a 10-50% sucrose gradient, before ultracentrifugation at 38,000rpm for 2 hours. The sucrose gradient was made by combining 10% and 50% sucrose solutions in lysis buffer, plus 2.5mM DTT, 160U SuperaseIN, and 100µg/ml CHX. Both gradient formation and fractionation was performed on a Biocomp Gradient Profiler (Biocomp). For protein extraction post-fractionation, 1 volume of TCA to 4 volumes of protein sample was added, incubated for 10 minutes on ice, then span at 14,000rpm for 5 minutes at 4^{o}C. Pellet was washed twice with ice-cold acetone and dried at 95^{o}C for 5 minutes. For SDS-PAGE, 4x Sample reducing agent (GenScript) diluted 1:1 in RIPA was added to dry pellet and heated for an additional 5 minutes at 95^{o}C.

Western blotting

Western blot analysis was performed after SDS-PAGE using standard protocols. The antibodies used were anti-HA (Thermo; 2-2.2.14), anti-RPL3 (Sigma; 3365), anti-RPL28 (Invitrogen; 62192), anti-RPS3 (Thermo; 2G7H4), anti-GAPDH (Invitrogen; PA1988), and anti-Vinculin (CST; E1E9V).

RNA extraction/analysis and qPCR

Total RNA was isolated using NucleoSpin RNA Plus kit (Macharey-Nagel) including use of gDNA removal column. For rRNA analysis, 500ng RNA was ran on 2100 Bioanalyzer (Agilent) at the Columbia University's Molecular Pathology Core. For qPCR, cDNA synthesis was performed using 500ng RNA with the SuperScript IV Reverse Transcriptase (Invitrogen) as per the manufacturers protocol with oligo(dT) primers. Real-time PCR was carried out using PowerUp SYBR Green Master Mix using 1:3 diluted cDNA. For qRT-PCR, triplicates were performed with 0.5µM primer and 1µl cDNA per reaction in the QuantStudio 7 Flex Real-Time PCR System. Ct values obtained were plotted using GAPDH mRNA as loading control. Quantifications were performed as per the MIQE guidelines. Statistical tests were performed with Prism 9 and two-sided Student's t-test were used to calculate the P values. Data are expressed as mean ± s.d. for the number of replicates indicated, with $P < 0.01$ considered significant. All primer sequences are displayed in Supplementary Table.

Nuclear/Cytoplasmic RNA Extraction Protocol

Cells were washed with ice-cold PBS, scraped into 1.7ml tubes, and pelleted at 500 g for 5 minutes at 4°C. Cytoplasmic RNA was extracted by lysing cells in Cytoplasmic Lysis Buffer (40 mM HEPES-NaOH pH 7.5, 160 mM KCl, 10 mM $MgCl_2$, 0.5% NP-40, 0.5% glycerol, and SuperaseIn) for 30 minutes at 4°C with rotation, followed by centrifugation at 2000 g for 2 minutes to separate the cytoplasmic supernatant. Nuclear pellets were washed in Cytoplasmic Lysis Buffer (without detergent), centrifuged at 500 g for 5 minutes, and subjected to a wash with Nuclear Lysis Buffer (10 mM Tris-HCl pH 8.0, 1.5 mM KCl, 2.5 mM $MgCl_2$, 5% glycerol, 0.5% Triton X-100, 0.5% deoxycholate, and 20U SuperaseIN) for 15 minutes at 4°C. Final washes were performed with Cytoplasmic Lysis Buffer (without detergent), and nuclei were pelleted at 2000 g for 2 minutes. Both cytoplasmic and nuclear RNA were extracted as above.

Co-immunoprecipitation

To prepare magnetic bead-antibody complex, 50µl of Dynabeads Protein G (Invitrogen) was washed 3x with lysis buffer then incubated with 5µl anti-HA (Thermo; 2-2.2.14) in 400µl lysis buffer for 1 hour at RT with constant rotation. Magnetic bead-antibody complex was captured using Magnetic Separation Rack (NEB), washed 2x with cytoplasmic lysis buffer (20mM Tris-HCl, pH 7.5, 150mM KCl, 10% glycerol and 0.1% NP-40) then 3x with lysis buffer. 2x 15cm plates at 80% confluency were grown, cells were washed twice in ice-cold PBS and 400µl of ice-cold lysis buffer was dripped onto the plate and scraped, incubated in microcentrifuged tubes after triturating 10x with 23-gauge syringe, and centrifuged at 14,000g for 7 minutes. The supernatant was added to bead-antibody complex and incubated with rotation at 4°C overnight. Next day, bead-antibody-antigen complex was washed 3x for 10 minutes at 4°C with ice-cold lysis buffer. Protein was eluted by incubation with 40µl pH 3 glycine and heating at 70°C with shaking for 10 minutes, then neutralized with 1/10 volume pH 8.0 Tris.

Proteomics

Explanted frozen patient tissue was homogenized with pestle and mortar under liquid nitrogen and lysed in RIPA for mass spectrometric analysis by the Proteomics core facility at Columbia. PASEF[64] based proteomics was employed. Briefly, proteins were denatured in 0.5% sodium deoxycholate (SDC) buffer containing 100 mM Tris-HCl (pH 8.5). Samples were boiled for 20 minutes at 60°C with agitation (1000 rpm). Protein reduction and alkylation of cysteine residues were performed using 10 mM tris(2-carboxyethyl)phosphine (TCEP) and 40 mM chloroacetamide (CAA) at 45°C for 15 minutes. Following sonication in a water bath and cooling to room temperature, and cleanup with the SP3 method[65]. Proteins were digested overnight with a LysC/trypsin mix at a 1:50 enzyme-to-protein ratio at 37°C and 1400 rpm. After digestion, peptides were acidified with 1% trifluoroacetic acid (TFA), vortexed, and desalted using StageTip technology with SDB-RPS (styrene-divinylbenzene reversed-phase sulfonate) resin. The cleaned peptides were dried using a vacuum concentrator and resuspended in 10 μl of LC buffer (3% acetonitrile, 0.1% formic acid). Peptide concentrations were measured using a NanoDrop spectrophotometer, and 150 ng of each sample was subjected to PASEF analysis on a timsTOF Pro2 mass spectrometer. For liquid chromatography with tandem mass spectrometry (LC-MS/MS), peptides were separated within 87 min at a flow rate of 300 nl/min on a reversed-phase C18 column with an integrated CaptiveSpray Emitter (25 cm x 75μm, 1.6 μm, IonOptics). Mobile phases A and B were with 0.1% formic acid in water and 0.1% formic acid in ACN. The fraction of B was linearly increased from 2 to 23% within 60 min, followed by an increase to 35% within 7 min and a further increase to 90% before re-equilibration. The timsTOFPro2 mass spectrometer was operated in Parallel Accumulation Serial Fragmentation (PASEF) mode with the following parameters: a mass range of 100 to 1700 m/z, an ion mobility range of 1/K0 0.7 to 1.3 V·s/cm², a ramp time of 200 ms, and a duty cycle locked to 100%. Additional settings included a capillary voltage of 1600 V, dry gas flow rate of 3 L/min, and a dry temperature of 180°C. PASEF data acquisition was configured for 10 MS/MS frames with a total duty cycle of 2.27 seconds, targeting a charge range of 0–5, an active exclusion duration of 0.4 minutes, a target intensity of 20,000, and an intensity threshold of 2500. Collision-induced dissociation (CID) was performed with a collision energy of 59 eV. A polygon filter was applied in the m/z and ion mobility dimensions to prioritize features likely representing peptide precursors and exclude singly charged background ions. For LC-MS/MS data analysis, acquired PASEF raw files were analyzed using the MaxQuant environment V.2.2.0.0 and Andromeda for database searches at default settings with a few modifications[66]. The default is used for the first search tolerance and main search tolerance (20 ppm and 4.5 ppm, respectively). MaxQuant was set up to search with the reference human proteome database downloaded from UniProt. MaxQuant performed the search trypsin digestion with up to 2 missed cleavages. Peptide, site, and protein false discovery rates (FDR) were all set to 1% with a minimum of 1 peptide needed for identification; label-free quantitation (LFQ) was performed with a minimum ratio count of 1. The following modifications were used for protein identification and quantification: Carbamidomethylation of cysteine residues (+57.021 Da) was set as static modifications, while the oxidation of methionine residues (+15.995 Da), and deamidation (+0.984) on asparagine were set as a variable modification. Results obtained from MaxQuant, protein groups table was further used for data analysis. Custom fasta files containing mutant D308N peptides were analyzed on MaxQuant for peptide abundance relative to wildtype peptide abundance. For nuclear IP of cell

lines, MaxQuant outputs were analyzed using Differential Expression of Proteins (DEP) package in R.

Chromatin Immunoprecipitation (ChIP)

Cells were crosslinked with 1% formaldehyde in the culture medium for 15 minutes at room temperature, followed by quenching with 125 mM glycine for 5 minutes. Cells were pelleted and washed. The pellet was resuspended in RIPA buffer with protease inhibitor cocktail (Roche), incubated on ice, and sonicated to shear chromatin (~1 kb fragments). The lysate was centrifuged at 12,000 x g for 10 minutes at 4 °C, and the volume was increased with dilution buffer (165 mM NaCl, 0.01% SDS, 1.1% Triton X-100, 1.2 mM EDTA, 16.7 mM Tris–HCl, pH 8.0). Chromatin was incubated with 5 µg of anti-RNAPII (Santa Cruz; F-12) for overnight at 4 °C. Chromatin-Antibody mix was incubated with pre-washed Dynabeads and incubated at RT for 45 minutes. Beads were washed 5x in dilution buffer and eluted in genomic DNA extraction kit (Macharey-Nagel) lysis buffer, resuspended in 50 µL of water. PCR was performed using purified DNA and compared to input samples. Two-sided Student's t-tests were carried out to calculate P value.

Nuclear localization calculation

Nuclear localization was calculated as a ratio of the immunofluorescent co-localization signal of anti-HA with nuclear regions compared to anti-HA signal with cytoplasmic regions, determined as cellular regions lacking in DAPI signal. The same calculation was performed for 28S rRNA. All calculations were carried out with ImageJ. Cytoplasmic/Nuclear signal ratios for each cell were compiled and underwent one-way ANOVA with Dunnett correction for multiple comparisons across conditions.

Tissue RNA-seq

Explanted frozen patient heart tissue was homogenized using pestle and mortar under liquid nitrogen. Homogenized tissue was submerged in TRIzol and extracted using Direct-zol kit (Zymo Research). RNA integrity was analyzed with Bioanalyzer electrophoresis. Samples underwent poly(A) RNA pulldown and libraries were constructed using Illumina TruSeq chemistry. Libraries were sequenced at Columbia Sulzberger Genome Center with NovaSeq 6000 (Illumina). RTA (Illumina) was used for base calling and bcl2fastq2 (version 2.19) for converting BCL to fastq format. Gene expression was quantified using kallisto with a customized transcriptome index (human MANE select). RNA-seq data for three age-matched control samples were downloaded from European Nucleotide Archive (ENA project PRJEB26969, sample ID: 5818sTS, 5828sTS, and 5836sTS) and analyzed similarly. DESeq2 was used for differential gene expression analysis.

References

1. Bergmann, O. et al. Dynamics of Cell Generation and Turnover in the Human Heart. Cell 161, 1566-1575 (2015).
2. Van Raay, T.J., Connors, T.D., Klinger, K.W., Landes, G.M. & Burn, T.C. A novel ribosomal protein L3-like gene (RPL3L) maps to the autosomal dominant polycystic kidney disease gene region. Genomics 37, 172-176 (1996).

3. Chaillou, T., Zhang, X. & McCarthy, J.J. Expression of Muscle-Specific Ribosomal Protein L3-Like Impairs Myotube Growth. J Cell Physiol 231, 1894-1902 (2016).

4. Shiraishi, C. et al. RPL3L-containing ribosomes determine translation elongation dynamics required for cardiac function. Nat Commun 14, 2131 (2023).

5. Gay, D.M., Lund, A.H. & Jansson, M.D. Translational control through ribosome heterogeneity and functional specialization. Trends Biochem Sci 47, 66-81 (2022).

6. Genuth, N.R. et al. A stem cell roadmap of ribosome heterogeneity reveals a function for RPL10A in mesoderm production. Nat Commun 13, 5491 (2022).

7. Genuth, N.R. & Barna, M. The Discovery of Ribosome Heterogeneity and Its Implications for Gene Regulation and Organismal Life. Mol Cell 71, 364-374 (2018).

8. Simsek, D. et al. The Mammalian Ribo-interactome Reveals Ribosome Functional Diversity and Heterogeneity. Cell 169, 1051-1065 e1018 (2017).

9. Shi, Z. et al. Heterogeneous Ribosomes Preferentially Translate Distinct Subpools of mRNAs Genome-wide. Mol Cell 67, 71-83 e77 (2017).

10. Li, H. et al. A male germ-cell-specific ribosome controls male fertility. Nature 612, 725-731 (2022).

11. Zou, Q. et al. Proteostasis regulated by testis-specific ribosomal protein RPL39L maintains mouse spermatogenesis. iScience 24, 103396 (2021).

12. Meskauskas, A., Harger, J.W., Jacobs, K.L. & Dinman, J.D. Decreased peptidyltransferase activity correlates with increased programmed -1 ribosomal frameshifting and viral maintenance defects in the yeast Saccharomyces cerevisiae. RNA 9, 982-992 (2003).

13. Burn, T.C. et al. Generation of a transcriptional map for a 700-kb region surrounding the polycystic kidney disease type 1 (PKD1) and tuberous sclerosis type 2 (TSC2) disease genes on human chromosome 16p3.3. Genome Res 6, 525-537 (1996).

14. Litvinukova, M. et al. Cells of the adult human heart. Nature 588, 466-472 (2020).

15. Consortium, G.T. The GTEx Consortium atlas of genetic regulatory effects across human tissues. Science 369, 1318-1330 (2020).

16. Grimes, K.M. et al. Rpl3l gene deletion in mice reduces heart weight over time. Frontiers in Physiology 14 (2023).

17. Milenkovic, I. et al. RPL3L-containing ribosomes modulate mitochondrial activity in the mammalian heart. bioRxiv, 2021.2012.2004.471171 (2021).

18. Ganapathi, M. et al. Bi-allelic missense disease-causing variants in RPL3L associate neonatal dilated cardiomyopathy with muscle-specific ribosome biogenesis. Hum Genet 139, 1443-1454 (2020).

19. Das, B.B., Gajula, V., Arya, S. & Taylor, M.B. Compound Heterozygous Missense Variants in RPL3L Genes Associated with Severe Forms of Dilated Cardiomyopathy: A Case Report and Literature Review. Children (Basel) 9 (2022).

20. Nannapaneni, H. et al. Further Evidence of Autosomal Recessive Inheritance of RPL3L Pathogenic Variants with Rapidly Progressive Neonatal Dilated Cardiomyopathy. J Cardiovasc Dev Dis 9 (2022).

21. Al-Hassnan, Z.N. et al. Categorized Genetic Analysis in Childhood-Onset Cardiomyopathy. Circ Genom Precis Med 13, 504-514 (2020).

22.   Bajpai, A.K. et al. Exploring the Regulation and Function of Rpl3l in the Development of Early-Onset Dilated Cardiomyopathy and Congestive Heart Failure Using Systems Genetics Approach. Genes (Basel) 15 (2023).

23.   Yang, Q. et al. Novel compound heterozygous variants in the RPL3L gene causing dilated cardiomyopathy type-2D: a case report and literature review. BMC Med Genomics 16, 127 (2023).

24.   McNally, E.M. & Mestroni, L. Dilated Cardiomyopathy: Genetic Determinants and Mechanisms. Circ Res 121, 731-748 (2017).

25.   Rosenbaum, A.N., Agre, K.E. & Pereira, N.L. Genetics of dilated cardiomyopathy: practical implications for heart failure management. Nat Rev Cardiol 17, 286-297 (2020).

26.   Kamisago, M. et al. Mutations in sarcomere protein genes as a cause of dilated cardiomyopathy. N Engl J Med 343, 1688-1696 (2000).

27.   Milenkovic, I. et al. Dynamic interplay between RPL3- and RPL3L-containing ribosomes modulates mitochondrial activity in the mammalian heart. Nucleic Acids Res 51, 5301-5324 (2023).

28.   Brown, L.P. & McCarthy, J.J., Vol. PhD (University of Kentucky, 2021).

29.   Di Fede, G. et al. A recessive mutation in the APP gene with dominant-negative effect on amyloidogenesis. Science 323, 1473-1477 (2009).

30.   Chen, S. et al. A genomic mutational constraint map using variation in 76,156 human genomes. Nature 625, 92-100 (2024).

31.   Cheng, J. et al. Accurate proteome-wide missense variant effect prediction with AlphaMissense. Science 381, eadg7492 (2023).

32.   Jumper, J. et al. Highly accurate protein structure prediction with AlphaFold. Nature 596, 583-589 (2021).

33.   Bick, A.G. et al. Genomic data in the All of Us Research Program. Nature 627, 340-346 (2024).

34.   Bycroft, C. et al. The UK Biobank resource with deep phenotyping and genomic data. Nature 562, 203-209 (2018).

35.   Thorolfsdottir, R.B. et al. Coding variants in RPL3L and MYZAP increase risk of atrial fibrillation. Commun Biol 1, 68 (2018).

36.   Sun, K.Y. et al. A deep catalogue of protein-coding variation in 983,578 individuals. Nature 631, 583-592 (2024).

37.   Dorner, K., Ruggeri, C., Zemp, I. & Kutay, U. Ribosome biogenesis factors-from names to functions. EMBO J 42, e112699 (2023).

38.   Adamo, L., Rocha-Resende, C., Prabhu, S.D. & Mann, D.L. Reappraising the role of inflammation in heart failure. Nat Rev Cardiol 17, 269-285 (2020).

39.   Davidson, M.M. et al. Novel cell lines derived from adult human ventricular cardiomyocytes. J Mol Cell Cardiol 39, 133-147 (2005).

40.   Hedaya, O.M. et al. Secondary structures that regulate mRNA translation provide insights for ASO-mediated modulation of cardiac hypertrophy. Nat Commun 14, 6166 (2023).

41.   Guo, Y. & Pu, W.T. Cardiomyocyte Maturation: New Phase in Development. Circ Res 126, 1086-1106 (2020).

42. Thomas, D., Cunningham, N.J., Shenoy, S. & Wu, J.C. Human-induced pluripotent stem cells
in cardiovascular research: current approaches in cardiac differentiation, maturation
strategies, and scalable production. Cardiovasc Res 118, 20-36 (2022).

43. Pillet, B. et al. Dedicated chaperones coordinate co-translational regulation of ribosomal
protein production with ribosome assembly to preserve proteostasis. Elife 11 (2022).

44. Iouk, T.L., Aitchison, J.D., Maguire, S. & Wozniak, R.W. Rrb1p, a yeast nuclear WD-repeat
protein involved in the regulation of ribosome biosynthesis. Mol Cell Biol 21, 1260-1271
(2001).

45. Schaper, S. et al. A yeast homolog of chromatin assembly factor 1 is involved in early
ribosome assembly. Curr Biol 11, 1885-1890 (2001).

46. Bhutada, P. et al. Rbp95 binds to 25S rRNA helix H95 and cooperates with the Npa1
complex during early pre-60S particle maturation. Nucleic Acids Res 50, 10053-10077
(2022).

47. Sailer, C. et al. A comprehensive landscape of 60S ribosome biogenesis factors. Cell Rep 38,
110353 (2022).

48. Ismail, S. et al. Emergence of the primordial pre-60S from the 90S pre-ribosome. Cell Rep
39, 110640 (2022).

49. De Marchis, M.L., Giorgi, A., Schinina, M.E., Bozzoni, I. & Fatica, A. Rrp15p, a novel
component of pre-ribosomal particles required for 60S ribosome subunit maturation. RNA
11, 495-502 (2005).

50. Tafforeau, L. et al. The complexity of human ribosome biogenesis revealed by systematic
nucleolar screening of Pre-rRNA processing factors. Mol Cell 51, 539-551 (2013).

51. Bruning, L. et al. RNA helicases mediate structural transitions and compositional changes
in pre-ribosomal complexes. Nat Commun 9, 5383 (2018).

52. Rodriguez-Mateos, M. et al. Role and dynamics of the ribosomal protein P0 and its related
trans-acting factor Mrt4 during ribosome assembly in Saccharomyces cerevisiae. Nucleic
Acids Res 37, 7519-7532 (2009).

53. Lee, T.M. et al. Pediatric Cardiomyopathies. Circ Res 121, 855-873 (2017).

54. Towbin, J.A. et al. Incidence, causes, and outcomes of dilated cardiomyopathy in children.
JAMA 296, 1867-1876 (2006).

55. Lipov, A. et al. Exploring the complex spectrum of dominance and recessiveness in genetic
cardiomyopathies. Nat Cardiovasc Res 2, 1078-1094 (2023).

56. Bamshad, M.J., Nickerson, D.A. & Chong, J.X. Mendelian Gene Discovery: Fast and Furious
with No End in Sight. Am J Hum Genet 105, 448-455 (2019).

57. Young, L.C. et al. Destabilizing NF1 variants act in a dominant negative manner through
neurofibromin dimerization. Proc Natl Acad Sci U S A 120, e2208960120 (2023).

58. Utsumi, T. et al. Exclusive Characteristics of the p.E555K Dominant-Negative Variant in
Autosomal Dominant E47 Deficiency. J Clin Immunol 44, 167 (2024).

59. Boettcher, S. et al. A dominant-negative effect drives selection of TP53 missense mutations
in myeloid malignancies. Science 365, 599-604 (2019).

60. Gerasimavicius, L., Livesey, B.J. & Marsh, J.A. Loss-of-function, gain-of-function and
dominant-negative mutations have profoundly different effects on protein structure. Nat
Commun 13, 3895 (2022).

61.   El-Brolosy, M.A. et al. Genetic compensation triggered by mutant mRNA degradation.
784   Nature 568, 193-197 (2019).
785   62.   Rossi, A. et al. Genetic compensation induced by deleterious mutations but not gene
786   knockdowns. Nature 524, 230-233 (2015).
787   63.   El-Brolosy, M.A. & Stainier, D.Y.R. Genetic compensation: A phenomenon in search of
788   mechanisms. PLoS Genet 13, e1006780 (2017).
789   64.   Meier, F. et al. Online Parallel Accumulation-Serial Fragmentation (PASEF) with a Novel
790   Trapped Ion Mobility Mass Spectrometer. Mol Cell Proteomics 17, 2534-2545 (2018).
791   65.   Soni, R.K. Protocol for deep proteomic profiling of formalin-fixed paraffin-embedded
792   specimens using a spectral library-free approach. STAR Protoc 4, 102381 (2023).
793   66.   Cox, J. et al. Andromeda: A Peptide Search Engine Integrated into the MaxQuant
794   Environment. Journal of Proteome Research 10, 1794-1805 (2011).
795

Figure legends

Figure 1. Hotspot G27D and D308N/V are potential driver mutations in *RPL3L*-linked DCM. a,
Schematic of neonatal DCM-causing mutations across RPL3L protein. Each circle (missense) or
rectangle (frameshift) represents a mutation. Mutations are colored by family and compound
heterozygous mutations found in the same family are linked by dashed lines. Mutations found in
the two new cases reported here are highlighted with a star. Protein domains in RPL3L are also
labeled. The two hotspot mutations, G27D and D308N/V, are emphasized in bold. b, Positions of
the variants (highlighted in red) mapped onto the AlphaFold-predicted 3D structure of RPL3L. c,
RPL3L missense variants clustered into two groups by their allele frequency (gnomAD) and
predicted pathogenicity (AlphaMissense). Variants are colored and linked as in a. d, Putative
human *RPL3L* KO in the general population (gnomAD). LOF: loss-of-function.
807
Figure 2. Ribosome defects in explanted patient hearts. a, Bioanalyzer electrophoresis of total
RNA isolated from explanted patient heart ventricular tissue and a control sample (human AC16
cells). b, Differential gene expression analysis comparing explanted heart tissue (N=2)and age-
matched healthy control heart ventricular tissue (N=3). Genes encoding 60S or 40S ribosomal
proteins are colored red or blue, respectively. *RPL3* and *RPL3L* are highlighted by yellow circles. c,
Cumulative distribution function plots for three significant gene signatures. Red indicates genes of
interest and black indicates all other genes. The median log2 fold change (log2fc) and Kolmogorov–
Smirnov test P values are also shown. d, RNA-seq read coverage (BAM file visualized in IGV genome
browser) showing that both D308N and T189M alleles are equally expressed. e, Mass
spectrometry quantification of RPL3 (left), RPL3L (center), and total 60S proteins (right) relative
total 40S proteins in D308N/T189M patient ventricular tissue (N=2) and age-matched control
tissue (N=3). ** P < 0.0001. f, Intensity of the peptide containing the D308 residue in the T189M
variant and the peptide containing the N308 residue in the D308N variant normalized to the total
intensity in RPL3L. ns: not significant.

Figure 3. Hotspot variants disrupt ribosome biogenesis and impairs cell viability. a, Isogenic
AC16 human cardiomyocyte-like cell lines were generated to simultaneously knockdown (KD) RPL3

and overexpress HA-tagged RPL3L (or its variants) via a dox-inducible promoter. **b**, RPL3 knockdown after dox treatment. Total RNA was isolated from indicated cell lines and analyzed for RPL3 levels. shScramble: cells expressing only a control scrambled shRNA. shRPL3: cells expressing only the RPL3 targeting shRNA without RPL3L or its variants. RPL3L-WT/SNP/T189M/D308N: cells with RPL3 KD and expresses either wild type or the indicated variant of RPL3L. **c**, Efficient expression of RPL3L variants in AC16 cells after 120 hours Dox treatment. Total RNA isolated from indicated cells was analyzed for RPL3L levels (N = 3 independent experiments). **d**, Bioanalyzer electrophoresis on total RNA from indicated cell lines was quantified by ImageJ. **e**, Loss of translation capacity in shRPL3 and D308N cells, but not control or T189M cells. RPL3+ (shScramble) and RPL3L-expressing cells were separated by ultracentrifugation and fractionated in the presence of cycloheximide (N = 3). **f**, Cell growth defects in D308N and shRPL3 cells. AC16-derived cells were seeded at equal density and counted every 2 days (N = 4). The y-axis is shown on a logarithmic scale. The percentage values represent the cell density at day 10 relative to the RPL3L-WT cell line. **b ,c, f ,** Data are mean ± s.d. *: $P < 0.05$.

Figure 4. Hotspot variants mislocalize and alter interactions with ribosome biogenesis factors. **a**, Cells were stained for anti-HA antibody (RPL3L and variants) and 28S rRNA FISH probes and analyzed by 60x confocal microscopy. Scale bar = 50μm. **b**, Violin plot quantification of cytoplasmic and nuclear abundances of HA (RPL3L) signals for indicated cell lines, measured per cell (N = 26-41 cells). **c**, same as b but for 28S rRNA. One-way ANOVA was used for statistical analysis. **d**, PCA analysis of the proteins co-immunoprecipitated with RPL3L variants. N=3. **e**, Enrichment of GRWD1 and C7ORF50 in proteins co-immunoprecipitated with the hotspot variants D308N and G27D compared to other variants. Peptide intensities are normalized by the median of all samples and then log2-transformed. Error bars represent standard deviation. ns, not significant; ** $P < 0.001$; *** $P < 0.0001$; **** $P < 0.00001$.

Figure 5. Non-hotspot variants drive post-transcriptional *RPL3* compensatory up-regulation. **a**, Compensatory increase of *RPL3* mRNA in R161W cells. DCM-associated *RPL3L* variants were expressed for 120 hours in doxycycline and total RNA was extracted and analyzed for *RPL3* mRNA levels (N = 3 independent experiments). **b**, Western blotting shows compensatory increase of RPL3 protein levels in R161W cells. Cells were treated as in (a) and whole-cell lysates were immunoblotted with indicated antibodies (N = 3 independent experiments). **c**, No difference in transcription activity in *RPL3* promoter assayed by ChIP-qPCR. Multiple primer pairs were used for each region. An intergenic region upstream of *RPL3L* was used as a negative control of non-transcribed region (N = 3). **d**, *RPL3* mRNA half-life is enhanced in R161W cells. *RPL3L*-expressing variants were incubated with 50μg/ml RNA Polymerase II inhibitor α-amanitin for indicated timepoints prior to analysis for *RPL3* mRNA relative to GAPDH, normalized to 0 hour (N = 3). **e**, Compensatory *RPL3* mRNA increase occurs entirely in the cytoplasm in R161W cells. RNA underwent subcellular fractionation for amplification of *RPL3* levels relative to GAPDH (N = 4). **a,c,d,e**, Data are mean ± s.d. ns, not significant; * $P < 0.01$; *** $P < 0.0001$; **** $P < 0.00001$.

Figure 6. Impaired protein synthesis in engineered compound heterozygous cells. **a**, Cells stably integrated with G27D, R161W, or both G27D and R161W were induced with dox for 144hrs. Puromycin was added to 1uM final concentration for 30 minutes before cells were harvested for

868   Western blotting (N = 4). **b**, Quantification of puromycin incorporation. **c**, Quantification of RPL3
869   protein abundance. Data are mean ± s.d. ns, not significant; * P < 0.01.
870

871   **Extended Data Figure 1. iPSC cardiomyocytes (iPSC-CMs) do not express *RPL3L*.** Data was
872   analyzed from RNA-seq performed by Pozo et al (2022) comparing iPSC-CMs to fetal and adult
873   human heart.
874

875   **Extended Data Figure 2. The hotspot G27D variant mirrors the effects of D308N. a**, Cell growth
876   defect for G27D and R161W, as in **Fig. 3f. b**, Polysome fraction for G27D and R161W, as in **Fig. 3e**.
877   Data are mean ± s.d.
878

879   **Extended Data Figure 3. Enrichment of 60S ribosome biogenesis factors in proteins co-**
880   **immunoprecipitated with nuclear RPL3L variants.** Peptide intensities are normalized by the
881   median of all samples and then log2-transformed. Error bars represent standard deviation.
882

883   **Extended Data Figure 4. R161W does not incorporate into polysomes.** Monosome and
884   polysomal fractions from **Fig. 5** were isolated by TCA precipitation and immunoblotted for HA-tag.
885   Input (pre-ultracentrifugation) samples were loaded as controls.
886

887   **Extended Data Figure 5. No change in the expression of the *RPL3*-targeting shRNA in *RPL3L*-**
888   **expressing cells**. A custom annotation was used to detect shRNA hairpin from RNA-seq data
889   obtained from *RPL3L* variant-expressing cell lines. RPM, Reads per million (N = 3 biological
890   replicates). Error bars represent standard deviation. ns, not significant.
891

892   **Extended Data Figure 6. R161W does not induce higher *RPL3* intronic RNA as a proxy of**
893   **transcription levels**. Total RNA was isolated and cDNA was synthesized using random primers.
894   Primers targeting intron 3 and intron 9 of *RPL3* were used for amplification (N = 3). Error bars
895   represent standard deviation. ns, not significant.
896

Figure 1

a

b

c

d

Putative LOF variant	Allele frequency	Homozygous humans
R226X (stop-gain)	1.67x10^{-5}	1 (gnomAD)
c.1167+1G>A (splice donor)	2.49x10^{-3}	9 (gnomAD)
	1.70x10^{-3}	2 (AllofUs)
	3.41x10^{-3}	6 (UK Biobank)

Figure 3

Figure 4

Figure 5

Figure 6

Extended Data Figure 1

Extended Data Figure 2

a

b

Extended Data Figure 3

Extended Data Figure 4

Extended Data Figure 5

Extended Data Figure 6