High-quality genome assemblies of 152 root commensal bacteria from the model legume *Lotus japonicus*

Adrián Gómez-Repollés[1*], Eber Villa-Rodriguez[1*], Shaun Ferguson[1†], Simona Radutoiu[1†]

[1] Department of Molecular Biology and Genetics, Aarhus University, Universitetsbyen 81, Building 1872, DK-8000 Aarhus C, Denmark

[*]These authors contributed equally to this work

[†]Corresponding authors. E-mail address: ferguson@mbg.au.dk; radutoiu@mbg.au.dk

Abstract

Bacterial culture collections represent a valuable tool for mechanistic understanding of microbiome assemblies and are increasingly used to assemble tailored synthetic communities to characterize their microbe-microbe interactions and those with the environment. Given the size of these collections, short-read sequencing is primarily used to capture the encoded genetic information. Whilst sufficient for many microbiome studies, this approach is not amenable for understanding bacterial genome evolution or detailed genetic analyses at the entire genome level. Here we report the assembly of 152 full bacterial genomes from the *Lj*-SPHERE, the *Lotus japonicus* collection of root commensals. We performed long-read sequencing using Oxford Nanopore technology and used this together with pre-existing Illumina sequences to *de novo* assemble these into high quality genomes with improved contiguity and quality. These genomes now provide a solid platform for detailed, mechanistic understanding of microbiome assembly, dynamics and evolution in plants.

Background & Summary

In recent decades the importance of the plant microbiota has been well established, with burgeoning research efforts aiming to extend understanding of plant microbiomes ('the plant's second genome') for improvement of plant health, survival, and growth[1,2]. The soil environment in particular contains some of the greatest biological diversity known, and the rhizosphere (influenced by plant root exudates) contains up to 10^{11} microbes per gram of plant root[1]. However, this also means that within the rhizosphere, microbes exist in incredibly complex and dynamic communities[2,3]. To help disentangle the complexity of host-microbe and microbe-microbe interactions in these settings, reductionist experimental approaches such as the use of Synthetic Communities (SynComs) are increasingly being utilized[4-7].

SynComs consist of previously isolated bacteria, spanning diverse bacterial taxa, that covers the natural taxonomic diversity and composition from a particular environment[4,8]. These bacteria can be cultured under controlled conditions, to investigate characteristics and dynamics within the bacterial community and/or community-host interactions[9]. The application of bacterial SynComs across various research areas has yielded interesting discoveries in, for example, bioremediation[10,11], disease treatment[12-14], plant growth promotion[15,16], or plant biotic[17,18], and abiotic stress mitigation[19].

Lotus japonicus is a model legume that is widely used for understanding rhizobia-legume interactions[20,21]. The *Lotus japonicus* culture collection[22], also called *Lj*-SPHERE, was established by isolation of commensal bacteria from the root compartment of *L. japonicus* grown in Cologne agricultural soil. The isolates were whole genome sequenced and assembled using Illumina short reads, followed by taxonomic identification using amplicon sequencing of the 16S ribosomal gene marker[22]. The previous assemblies existed only as low-

quality draft genomes to enable 16S rDNA amplicon-based abundance profiling of the

community.

Here, we aimed to improve the *Lj*-SPHERE low-quality drafts to obtain contiguous and

annotated high-quality genomes as reference material for future studies. For that, we focused

on a subset of 177 taxonomically diverse genomes that were previously determined as unique

through a one-to-one whole genome nucleotide similarity comparison. Genomes with

similarity values of 99.99% or higher were clustered together, and within each cluster (n=41),

one genome was chosen as a representative. The isolates, for the 177 genomes, were re-

grown and processed for DNA extraction, library preparation, and long-read sequencing

using the Oxford Nanopore Technology (ONT) MinION sequencing platform.

We obtained ONT long-read sequences for all the targeted isolates, allowing us to re-

assemble these genomes using a hybrid assembly approach by combining the original

Illumina reads with newly produced long reads (Figure 1a). Inspection of the available

Illumina sequences revealed that for four isolates novel Illumina sequencing was required to

reach the necessary quality for the input data. The assembly pipeline (Figure 1a & 1b)

provided us with 152 high-quality annotated bacterial genomes, where a large number of

genomes were generated automatically (n=148) and only a small number of genomes

required manual curation for completion (LjRoot30, LjRoot166, LjRoot111 and

LjRoot112A) (Figure 1c). To ensure high quality assemblies, we implemented genome

validation thresholds based on the National Center for Biotechnology Information (NCBI)

Prokaryotic Genome Annotation Standards. These thresholds were defined as minimum

completeness of 80%, less than 5% genomic contamination and, at least one complete copy

of the required ribosomal genes (5S, 16S and 23S). The overall statistics performed on the

final genomes (N50 and L50), revealed that 152 genomes were successfully assembled with

good contiguity and high quality (Figure 2; Supplementary Table 1).

The hybrid assembled genomes were taxonomically classified using GTDB-tk[23], that was

utilized to provide a revised taxonomic classification and an estimate of the closest available

references supported by nucleotide-wise comparisons. Inspection of the distribution of

taxonomic identification across the 152 genomes revealed a majority of Proteobacteria

(n=125), followed by other phyla, such as Actinobacteriota, Bacteroidota, and Firmicutes

(Figure 2). At the family level, we identified 18 distinct taxa and a predominance of

Burkholderiaceae   (n=58),   Rhizobiaceae   (n=23),   Pseudomonadaceae   (n=17),   and

Caulobacteraceae (n=11). Furthermore, the results revealed a large overlap between the

previous 16S ribosomal gene marker and the newly produced taxonomy at the genus level. A

small number of genomes underwent a shift from their previous identified genus (n=27),

although the majority of them were subsequently labeled as a sister genus (n=15). A

considerable number of genomes were identified reliably at the species level (n=58), vastly

improving the taxonomic assignment available for the *Lj*-SPHERE collection.

To ensure the highest quality genomes, we identified the closest available references from

NCBI based on taxonomic classification and used these to scaffold the hybrid-assembled

genomes (Figure 1d). This revealed that out of 152 genomes, only 35 of these could use the

closest reference(s) from NCBI as a scaffolding reference. The remainder of our genomes had

a lower fragmentation rate and higher overall genome quality compared to the relevant NCBI

references, therefore the NCBI references could not be used. This demonstrates the high

quality of the reassembled *Lj*-SPHERE genomes reported here, which are publicly available

for the scientific community at the European Nucleotide Archive (ENA).

Overall, we have produced 152 high-quality annotated genomes of commensal bacteria from

the roots of *L. japonicus*, substantially improving this resource for comprehensive

bioinformatic analyses, including metagenomic analysis, gene cluster identification,

comparative genomics, and genome mining. This also creates the necessary foundation for

future experimental studies such as those focused on bacterial evolution and whole genome

mutagenesis, where a high-quality genome is required.

Materials & Methods

Acquisition of previous genomic data and Illumina sequencing reads

The genome assemblies for the *Lj*-SPHERE were downloaded from the *At*-SPHERE website

(https://www.at-sphere.com/). The 16S ribosomal gene taxonomic identification of the

downloaded genomes was obtained from the Supplementary Data previously published[22].

The corresponding pre-existing Illumina reads, in fastq format, were downloaded from the

European Nucleotide Archive or ENA (Project: PRJEB37696). Inspection of the paired end

reads revealed four isolates (LjRoot112A, LjRoot204, LjRoot205 and LjRoot206) that

contained an inadequate number or quality of Illumina reads, therefore were re-sequenced.

The extracted DNA (described below) was sent to Novogene (Munich, Germany) for library

preparation and paired end Illumina sequencing (150 bp; Novaseq 6000).

Subset of Illumina genomes

Pairwise comparisons of nucleotide similarity were computed between the Illumina genomes

using FastANI (v1.33)[24]. The genomes with similarity values of 99.99% ANI or higher were

clustered together whereas the genome with similarity values below this threshold were

considered unique entities. A representative candidate for each genome cluster was selected

based on the optimal balance between the contig number and the genome length, while

prioritizing the selection of largely contiguous genomes. The Illumina reads of the non-

selected cluster candidates were merged to increase assembly coverage for the representative

isolate. The resulting list, consisting of the cluster representative isolates along with the

unique isolates, were selected for long read sequencing.

High Molecular Weight DNA extraction

The subset of selected genomes of previously isolated, purified and stored *L. japonicus*

strains[22] were streaked from glycerol stocks onto 0.3% Tryptic Soy Agar (TSB) plates.

Pellets were prepared by re-suspending a full loop of bacterial biomass into 500 μl of sterile

distilled water and centrifuging at 9000g for 2 min. High Molecular Weight (HMW) DNA

was extracted from the pellets as detailed previously[25], with modifications for the lysis step.

Briefly, pellets were resuspended in 500 μl lysis buffer A (10 mg/ml lysozyme, 100 mM Tris-

HCl pH 8.0, 50 mM EDTA pH 8.0) and incubated for 30 min at 37°C. Then, 300 μl of lysis

buffer B (1.3 M NaCl, 100 mM Tris-HCl pH 8.0, 50 mM EDTA pH 8.0 and 4% SDS), 4 μl of

RNase A (20 mg/ml), and 8 μl of proteinase L (20 mg/ml) were added to the mix and

incubated for 20 min at 60°C. Protein and polysaccharide precipitation was achieved by

adding 0.3x volumes of 5 M potassium acetate and incubating at 4°C for 10 min. The mix

was centrifuged at 12,000g for 20 min and the supernatant was transferred to a new low-bind

Eppendorf tube. DNA was recovered by adding 1 ml of binding buffer (20% PEG 8000 and 3

M NaCl) and 100 μl of AMPure XP beads by using a magnetic rack. Finally, DNA was

washed twice with 70% ethanol, resuspended in 10 mM Tris-HCl pH 8.0, and

fluorometrically quantified using Qubit™ dsDNA HS kit. Samples were stored at 4°C until

use.

ONT platform sequencing

Diluted DNA samples (50 ng) were used as input for library preparation. Genomic DNA

libraries were prepared using the Rapid Barcoding Kit 96 V14 (SQK-RBK114.96) and

sequenced on R9.4.1 flow cells using the ONT MinION sequencing device[26]. Each round was

processed and monitored using the ONT MinKNOW software (v21.11.9).

Genome reconstruction using a hybrid assembly approach

ONT fast5 files were base called and demultiplexed into fastq files using Guppy (v6.1.1). The

ONT reads (FASTQ) together with the Illumina reads (FASTQ), were supplied to an in-house

pipeline to perform the hybrid assembly and subsequent annotation of the resulting bacterial

genomes. Our pipeline, *HA.py* (v1.2), implemented using gwf (v1.7.2) and partly adapted

from a previously published pipeline[27], started with quality control (QC) using NanoPlot

(v1.36.1)[28] and FastQC (v0.11.9) for the ONT and Illumina reads, respectively. The QC

results were assessed before proceeding. Illumina reads that passed the QC were corrected for

sequencing errors using SPAdes (v3.15.4)[29], then passed to LoRDEC (v0.9)[30] to correct

sequencing errors in the ONT reads. Corrected long reads were passed to Flye (v2.9-b1768)[31]

to generate a draft assembly using only the ONT reads. The three sources of information,

corrected short reads, corrected long reads, and ONT draft assembly, were used as input to

perform the hybrid assembly using unicycler (v0.5.0)[32]. The resulting genome was cleaned of

duplicated contigs, or contigs within larger contigs, using Mashtree (v1.2.0)[33], an in-house R

script, and seqtk (v1.3-r106).

Genome quality was checked using Quast (v5.0.2)[34], BUSCO (v5.3.2)[35], and CheckM

(v1.2.0)[36]. The BUSCO validation required complete and single genes to be above 90%, and

duplicated, fragmented, and missing genes below 5%. Moreover, CheckM validation required

completeness above 90% and contamination below 5%. If the hybrid assembly passed the

validations, it was then annotated using the Prokaryotic Genome Annotation Pipeline, or

PGAP, from the NCBI[37].

The PGAP annotation was used to validate the quality of the genome. Genomes were deemed

successful if they contained the following: a gene ratio above 80%, less than 20%

pseudogenes, at least 20 tRNAs, and at least one copy of the 5S, 16S, and 23S rRNA

subunits. For this validation the identified 5S, 16S, and 23S rRNA sequence lengths were

allowed ≤10% deviation compared to the *Escherichia coli* rRNA subunits (5S: 120 bp, 16S:

1542bp, and 23S: 2904 bp). Finally, contigs containing no genes or only pseudogenes were

considered spurious and therefore discarded from the genome and the annotation.

Manual curation

Manual curation of unsuccessful genome assemblies was performed depending on the cause

of the failure. First, when CheckM reported contamination greater than 5%, we used an

auxiliary pipeline to search for contamination within the original Illumina genomes and, in

parallel, within the ONT-based draft assemblies (an intermediate product from the hybrid

assembly pipeline). The auxiliary pipeline identified the 16S ribosomal gene using Blastn

(v2.12.0+)[38], removed sequence duplication using seqkit (v2.2.0)[39], then extracted the V3-V4

and V5-V7 subunits using QIIME2 (v2022.11.1)[40]. The SILVA database (v138)[41] was

downloaded and converted into 16S, V3-V4 and V5-V7 QIIME2 format databases and used

for taxonomical classification. The taxonomic results were used to identify sources of

contamination originating from the Illumina and/or ONT input reads.

When BUSCO encountered discrepancies between the *Lj*-SPHERE 16S ribosomal gene and

the hybrid assembly taxonomies, we utilized the aforementioned 16S, V3-V4 and V5-V7

taxonomic identifications, checked for concordance with the CheckM identification and then

updated the *Lj*-SPHERE taxonomy accordingly. The hybrid assembly was then re-run using

the corrected taxonomic information, enabling successful validation.

Taxonomic classification and phylogeny

The resulting hybrid assembled genomes were taxonomically classified using the GTDB-Tk

toolkit (v2.1.1)[23], and the resulting taxonomy was compared with the updated 16S taxonomic

identification (Manually curated SILVA database) to evaluate the degree of congruence and

account for potential discrepancies. GTDB-Tk was then utilized to determine the taxonomy

of the genomes to species level, or if not possible, to unidentified species level. A

phylogenetic tree was constructed based on the 20 validated bacterial cores genes (VBCG)

determined and utilised by the VCBG pipeline (v1.3)[42].These sequences were then fed into

RAxML (v8.2.12)[43] to construct a maximum likelihood phylogenetic tree with 100 bootstrap

replicates. The resulting phylogeny was plotted using the R package ggtree (v3.10.0).

Genome categories for ENA upload

GTDB-Tk toolkit[23] taxonomic identification results were used to identify the closest genome

references. The software placed each genome on the bacteria and archaea tree of life,

reporting the closest and the second closest reference, and a clade comprised by comparable

references. Hybrid assembled genomes consisting of more than one contig were considered

for scaffolding. The appropriate scaffold reference was selected sequentially along the

closest, second closest or clade references, considering for the latter only references with ANI

values above 80% similarity. Once selected, the references were downloaded from GenBank

or RefSeq and their quality (N50, L50, number of contigs, and genome length) was assessed

using the available metadata and Quast (v5.0.2)[34]. Among the downloaded references, only

the genomes with better quality metrics than our hybrid assembled genomes were used for

genome scaffolding. Hybrid assembled genomes were scaffolded using RagTag (v2.1.0)[44].

Finally, we manually inspected the resulting scaffolded genomes and determined which ENA

category (contig, scaffold, or chromosome) was most appropriate for data submission.

Genome assemblies with one contig were directly included in the chromosome category,

whereas genomes that were not scaffolded were defined as contig category.

Data records

225   The improved set of genomes, their corresponding ONT reads, and the Illumina reads for the

226   four Illumina re-sequenced isolates are available at the European Nucleotide Archive (ENA)

227   under the project PRJEB75483.

Technical validation

Bacterial re-growth, DNA extraction and library preparation

231   Isolates of *Lj*-SPHERE were cultured in sterile conditions on 0.3% TSB medium. They were

232   checked for purity based on single colony morphology and handled using aseptic techniques.

233   DNA extractions were carried out in a specialised laboratory environment and all equipment

234   was sterilised before use.

Genome assembly

236   The technical validation of the genomes consisted of:

237   1. Quality control of raw reads.

238   2. Taxonomy verification using multiple software.

3. Annotation validation following the NCBI Prokaryotic Genome Annotation Standards (https://www.ncbi.nlm.nih.gov/refseq/annotation_prok/standards/).

4. Assembly redundancy and spurious contig removal.

5. Genome scaffolding, when applicable.

Code availability

The in-house pipeline, HA.py (v1.2), the scaffolding pipeline, scaffolding.py (v1.0), and the auxiliary pipeline, extract16Sregion.py (v1.0), including all the scripts and Anaconda environments used for the generation of this dataset, can be found on GitHub (https://github.com/adriangeerre/HybridAssembly).

Acknowledgments

This work was supported by the project Molecular Mechanisms and Dynamics of Plant–Microbe Interactions at the Root–Soil Interface (InRoot), supported by the Novo Nordisk Foundation (grant number NNF19SA0059362) (https://ccrp.vcl.ncsu.edu/content/inroot).

Author contributions

S.F conceived and designed the study. E.V.R performed the microbiology, DNA extraction, and sequencing library preparation. S.F and E.V.R performed the ONT sequencing. A.G-R designed and executed the computational methodology, including data collection, processing, hybrid assembly, annotation, and verification of the genomes. A.G-R wrote the first draft of the manuscript with contributions from E.V.R. S.R. and S.F supervised the study and revised

261   the manuscript. All authors contributed to reviewing the draft versions of the manuscript. S.R

262   provided the funding and resources to conduct the study.

264   **Competing interests**

265   The authors declare no competing interests.

References

Berendsen, R. L., Pieterse, C. M. & Bakker, P. A. The rhizosphere microbiome and plant health. *Trends in Plant Science* **17**, 478-486 (2012).

Compant, S. *et al.* Harnessing the plant microbiome for sustainable crop production. *Nature Reviews Microbiology*, 1-15 (2024).

Philippot, L., Raaijmakers, J. M., Lemanceau, P. & Van Der Putten, W. H. Going back to the roots: the microbial ecology of the rhizosphere. *Nature Reviews Microbiology* **11**, 789-799 (2013).

Jing, J., Garbeva, P., Raaijmakers, J. M. & Medema, M. H. Strategies for tailoring functional microbial synthetic communities. *The ISME journal*, wrae049 (2024).

Schäfer, M. *et al.* Metabolic interaction models recapitulate leaf microbiota ecology. *Science* **381**, eadf5121 (2023).

Vannier, N. *et al.* Genome-resolved metatranscriptomics reveals conserved root colonization determinants in a synthetic microbiota. *Nat. Commun.* **14**, 8274 (2023).

Vorholt, J. A., Vogel, C., Carlström, C. I. & Müller, D. B. Establishing causality: opportunities of synthetic communities for plant microbiome research. *Cell Host Microbe* **22**, 142-155 (2017).

Gallardo-Navarro, O., Aguilar-Salinas, B., Rocha, J. & Olmedo-Álvarez, G. Higher-order interactions and emergent properties of microbial communities: The power of synthetic ecology. *Heliyon* **10** (2024).

Shayanthan, A., Ordoñez, P. A. C. & Oresnik, I. J. The role of synthetic microbial communities (SynCom) in sustainable agriculture. *Frontiers in Agronomy* **4**, 896307 (2022).

10   Ibrar, M. & Zhang, H. Construction of a hydrocarbon-degrading consortium and characterization of two new lipopeptides biosurfactants. *Sci. Total Environ.* **714**, 136400 (2020).

11   Vargas-Suárez, M., Fernández-Cruz, V. & Loza-Tavera, H. Biodegradation of polyacrylic and polyester polyurethane coatings by enriched microbial communities. *Applied Microbiology and Biotechnology* **103**, 3225-3236 (2019).

12   Lengfelder, I. *et al.* Complex bacterial consortia reprogram the colitogenic activity of *Enterococcus faecalis* in a gnotobiotic mouse model of chronic, immune-mediated colitis. *Front. Immunol.* **10**, 1420 (2019).

13   Li, Y. *et al.* Identification of trypsin-degrading commensals in the large intestine. *Nature* **609**, 582-589 (2022).

14   Weiss, A. S. *et al.* In vitro interaction network of a synthetic gut bacterial community. *The ISME journal* **16**, 1095-1109 (2022).

15   Kaur, S. *et al.* Synthetic community improves crop performance and alters rhizosphere microbial communities. *Journal of Sustainable Agriculture and Environment* **1**, 118-131 (2022).

16   Liu, H. *et al.* Effective colonisation by a bacterial synthetic community promotes plant growth and alters soil microbial community. *Journal of Sustainable Agriculture and Environment* **1**, 30-42 (2022).

17   Li, Z. *et al.* A simplified synthetic community rescues *Astragalus mongholicus* from root rot disease by activating plant-induced systemic resistance. *Microbiome* **9**, 1-20 (2021).

18   Pradhan, S., Tyagi, R. & Sharma, S. Combating biotic stresses in plants by synthetic microbial communities: Principles, applications and challenges. *J. Appl. Microbiol.* **133**, 2742-2759 (2022).

19   Schmitz, L. *et al.* Synthetic bacterial community derived from a desert rhizosphere

confers salt stress resilience to tomato in the presence of a soil microbiome. *The ISME*

Journal **16**, 1907-1920 (2022).

20   Handberg, K. & Stougaard, J. *Lotus japonicus*, an autogamous, diploid legume

species for classical and molecular genetics. *Plant J.* **2**, 487-496,

doi:https://doi.org/10.1111/j.1365-313X.1992.00487.x (1992).

21   Sato, S. & Tabata, S. *Lotus japonicus* as a platform for legume research. *Curr. Opin.*

Plant Biol. **9**, 128-132 (2006).

22   Wippel, K. *et al.* Host preference and invasiveness of commensal bacteria in the *Lotus*

and *Arabidopsis* root microbiota. *Nature Microbiology* **6**, 1150-1162,

doi:10.1038/s41564-021-00941-9 (2021).

23   Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk v2: memory

friendly classification with the genome taxonomy database. *Bioinformatics* **38**, 5315-

5316 (2022).

24   Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High

throughput ANI analysis of 90K prokaryotic genomes reveals clear species

boundaries. *Nat. Commun.* **9**, 5114 (2018).

25   Jones, A. *et al.* High-molecular weight DNA extraction, clean-up and size selection

for long-read sequencing. *PLoS One* **16**, e0253830 (2021).

26   Jain, M., Olsen, H. E., Paten, B. & Akeson, M. The Oxford Nanopore MinION:

delivery of nanopore sequencing to the genomics community. *Genome Biol.* **17**, 1-11

(2016).

27   Perry, B. J. *et al.* Complete genome sequences of *Trifolium* spp. inoculant strains

Rhizobium leguminosarum sv. *trifolii* TA1 and CC275e: resources for genomic study

of the *Rhizobium-Trifolium* symbiosis. *Mol. Plant-Microbe Interact.* **34**, 131-134

(2021).

28   De Coster, W. & Rademakers, R. NanoPack2: population-scale evaluation of long-

read sequencing data. *Bioinformatics* **39**, btad311 (2023).

29   Prjibelski, A., Antipov, D., Meleshko, D., Lapidus, A. & Korobeynikov, A. Using

SPAdes de novo assembler. *Current Protocols in Bioinformatics* **70**, e102 (2020).

30   Salmela, L. & Rivals, E. LoRDEC: accurate and efficient long read error correction.

Bioinformatics **30**, 3506-3514 (2014).

31   Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P. A. Assembly of long, error-prone

reads using repeat graphs. *Nat. Biotechnol.* **37**, 540-546 (2019).

32   Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Unicycler: resolving bacterial

genome assemblies from short and long sequencing reads. *PLoS Computational*

Biology **13**, e1005595 (2017).

33   Katz, L. S. *et al.* Mashtree: a rapid comparison of whole genome sequence files.

Journal of Open Source Software **4** (2019).

34   Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: quality assessment tool

for genome assemblies. *Bioinformatics* **29**, 1072-1075 (2013).

35   Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A. & Zdobnov, E. M. BUSCO

update: novel and streamlined workflows along with broader and deeper phylogenetic

coverage for scoring of eukaryotic, prokaryotic, and viral genomes. *Molecular*

Biology and Evolution **38**, 4647-4654 (2021).

36   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W.

CheckM: assessing the quality of microbial genomes recovered from isolates, single

cells, and metagenomes. *Genome Res.* **25**, 1043-1055 (2015).

37   Tatusova, T. *et al.* NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res.*

44, 6614-6624 (2016).

38   Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**,

1-9 (2009).

39   Shen, W., Le, S., Li, Y. & Hu, F. SeqKit: a cross-platform and ultrafast toolkit for

FASTA/Q file manipulation. *PLoS One* **11**, e0163962 (2016).

40   Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data

science using QIIME 2. *Nat. Biotechnol.* **37**, 852-857 (2019).

41   Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data

processing and web-based tools. *Nucleic Acids Res.* **41**, D590-D596 (2012).

42   Tian, R. & Imanian, B. VBCG: 20 validated bacterial core genes for phylogenomic

analysis with high fidelity and resolution. *Microbiome* **11**, 247 (2023).

43   Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis

of large phylogenies. *Bioinformatics* **30**, 1312-1313 (2014).

44   Alonge, M. *et al.* Automated assembly scaffolding using RagTag elevates a new

tomato system for high-throughput genome editing. *Genome Biol.* **23**, 258 (2022).

Additional information

Supplementary information:

Supplementary Table 1: Assembly statistics of Illumina and Hybrid assembled strains

Figure 1: Schematic overview of the hybrid assembly pipeline and downstream workflow. a) Hybrid assembly pipeline starting from the input sequencing data and finishing with a hybrid assembly draft. b) Computational workflow to validate and obtain a final hybrid assembly. c) Manual curation workflow including 16S taxonomic identification and reclassification. d) Genome scaffolding workflow. Across panels, the colour pattern represents: input (blue), software (orange), computational step (yellow), validation (purple), output (bright green), category (light grey), and passed/failed (green/red).

Figure 2: A maximum likelihood phylogenetic tree was constructed using concatenated alignments of 20 validated bacterial core genes (VBCG) across the 152 improved *Lj*-SPHERE genomes. The numbers at branch nodes represent 100 bootstrap replicates. The bar represents the number of substitutions per nucleotide site. Phylum is represented by four different symbols before the branch nodes. Branches are grouped as taxonomic families by coloured rectangles and tree tips. The first heatmap (green) displays the percentage of genome length covered by the N50 metric given the total length. The second heatmap (yellow/maroon) displays the percentage of genome contigs by the L50 given the total

403   number of contigs. Both heatmaps contain two columns, the first is Illumina *Lj*-SPHERE

404   genomes and the second is the hybrid assembled set of improved genomes. For both scales

405   (Norm. N50 & L50), darker values determine higher quality metrics.

Phylum
- ● Proteobacteria
- ▲ Actinobacteriota
- ◆ Firmicutes
- ■ Bacteroidota

Family
- ● Bacillaceae
- ● Beijerinckiaceae
- ● Brucellaceae
- ● Burkholderiaceae
- ● Caulobacteraceae
- ● Devosiaceae
- ● Flavobacteriaceae
- ● Intrasporangiaceae
- ● Microbacteriaceae
- ● Micrococcaceae
- ● Mycobacteriaceae
- ● Paenibacillaceae
- ● Pseudomonadaceae
- ● Rhizobiaceae
- ● Rhodanobacteraceae
- ● Sphingomonadaceae
- ● Xanthobacteraceae
- ● Xanthomonadaceae

Norm. N50

Norm. L50