Lower cortical activation and altered functional connectivity characterize passive auditory spatial attention in ASD

Sergio Osorio[1,2], Jasmine Tan[1,2], Grace Levine[1,2], Seppo P Ahlfors[1,3], Steven Graham[2], Fahimeh Mamashli[1,2,3], Sheraz Khan[1,3], Robert M Joseph[4], Zein Nayal[1,2], Ainsley Losh[2], Stephanie Pawlyszyn[2], Nicole M McGuiggan[2], Matti S Hämäläinen[1,3*], Hari Bharadwaj[5**], Tal Kenet[1,2**]

[1] Athinoula A. Martinos Center for Biomedical Imaging, Massachusetts General Hospital, Boston, MA, USA
[2] Department of Neurology, Harvard Medical School, Massachusetts General Hospital, Boston, MA, USA
[3] Department of Radiology, Harvard Medical School, Massachusetts General Hospital, Boston, MA, USA
[4] Department of Anatomy and Neurobiology, Boston University School of Medicine, Boston, MA, USA
[5] Department of Communication Science and Disorders, University of Pittsburgh, Pittsburgh, PA, USA
* Now at Department of Neuroscience and Biomedical Engineering, School of Science, Aalto University, Espoo, Finland
** These authors contributed equally to the manuscript

Abstract

Autism Spectrum Disorder (ASD) is a developmental condition characterized by difficulties in social interaction, communication and sensory processing. The ability to orient towards sounds is a key component of social interactions, yet auditory spatial attention remains relatively understudied in ASD, despite some prior research indicating differences in this domain. Here, we investigate the neural signatures associated with passive auditory spatial attention in children with ASD (n = 21, ages 6-17) relative to age- and IQ-matched typically developing (TD) children (n = 31), using source-localized magnetoencephalography (MEG). Participants listened passively, while watching a silenced movie, to non-social auditory stimuli designed to either remain lateralized to one hemifield (*stay* trials), or to change in location from one side to the contralateral hemifield (*jump* trials). Linear mixed effect modeling showed lower cortical activation in the auditory cortex in the ASD group in response to *jump* trials, relative to the TD group. Additionally, functional connectivity analyses showed higher alpha-band functional connectivity in the ASD group between auditory cortex seeds and frontoparietal regions known

to be recruited during auditory spatial attention. Right prefrontal alpha-band connectivity

estimates were associated with worse behaviorally assessed auditory processing scores and

increased ASD symptomatology. Our results align with the hypothesis that auditory spatial

attention generally, and specifically orientation to sounds even when experienced passively,

differs in ASD individuals.

Introduction

Autism Spectrum Disorder (ASD) is a developmental condition characterized by a complex set

of behavioral and sensory symptoms, the core of which is differences in social communication

(Ben-Sasson et al., 2019; Grzadzinski et al., 2013; Haesen et al., 2011; Hedger et al., 2020; O.

Miguel et al., 2017). Among its sensory manifestations, differences in processing of auditory

stimuli have been consistently documented among individuals with ASD (Alho et al., 2021;

Alho, Khan, et al., 2023; Alho, Samuelsson, et al., 2023; Bharadwaj et al., 2022; DePape et al.,

2012; Gonçalves & Monteiro, 2023; Haesen et al., 2011; Mamashli et al., 2017; Marco et al.,

2011; O'Connor, 2012; Port et al., 2015, 2017; Schafer et al., 2020).

Auditory spatial attention, i.e. our ability to perceive and track the location of sounds in

space, is a complex phenomenon involving several stages of sensory and cognitive processing.

For a sound to be successfully located in space, sensory encoding of acoustic features,

specifically the relative timing and amplitude differences between ears, needs to be integrated

with efficient selective attention and object recognition processes.(Fritz et al., 2007; Kayser et

al., 2005; van der Heijden et al., 2019). This integration allows for environmentally adequate

behavioral responses to be generated. In addition to primary and secondary auditory areas in the

superior temporal gyrus (STG), fMRI research has shown that locating a sound in space recruits

a frontoparietal network of cortical regions (Degerman et al., 2006; Deouell et al., 2007; Haist et

al., 2005; Kong et al., 2014; Smith et al., 2009; Zatorre et al., 1999). These regions include the

ventrolateral prefrontal cortex (vlPFC), the dorsolateral prefrontal cortex (dlPFC), the precentral

sulcus and gyrus, the central sulcus, the postcentral sulcus and gyrus, and the inferior and

superior temporal lobes (IPL/SPL). Unsurprisingly, these areas overlap with well-defined

functional networks observed in resting state fMRI data, including the ventral and dorsal

attention networks, and the frontoparietal executive control network (Hames et al., 2016;

Sadaghiani & Kleinschmidt, 2016; Supekar et al., 2013).

Research to date has shown that during auditory spatial attention tasks, individuals with

ASD show poorer behavioral performance, measured as higher reaction times and reduced

accuracy rates, when required to localize sounds in space (Soskey et al., 2017; Teder-Sälejärvi et

al., 2005). While previous studies using electroencephalography (EEG) and

magnetoencephalography (MEG) have established a range of differences in early sensory

encoding of auditory stimuli in ASD (Bruneau et al., 1999; Jansson-Verkasalo et al., 2005;

Orekhova et al., 2009; Roberts et al., 2010; Russo et al., 2009), these have been mostly

documented in the context of general auditory processing. One study that directly investigated

neural signatures in the context of auditory spatial attention in individuals with ASD showed that

poorer behavioral performance in ASD individuals during auditory spatial attention tasks is

accompanied by a lower amplitude of the N1 and the P3 components, two Event-Related

Potentials (ERP) associated with early sensory encoding and attentional processing,

correspondingly (Teder-Sälejärvi et al., 2005). Another study that also focused on auditory

spatial attention in ASD found that individuals with an ASD diagnosis showed an absence of the

Object-Related Negativity, an ERP component thought to reflect auditory segregation, as well as

a reduced amplitude of the P400, an attention-dependent ERP response thought to reflect

perceptual decision making (Lodhia et al., 2018). Notably, despite its importance for everyday

life and communication, auditory spatial attention remains understudied in ASD.

To address this gap in the literature, we investigated cortical responses in ASD and

Typically Developing (TD) individuals, ages 6 to 17, to direct manipulations of spatial

lateralization cues in sound. Using source-localized MEG signals, we focused on evoked

responses and functional connectivity from the auditory cortex to characterize the cortical

signatures of differences in passive auditory spatial orienting in ASD. We also conducted

correlation analyses to investigate possible associations between brain function and clinically

relevant behavioral measures of ASD.

Methods

Participants

Thirty-one typically developing (TD, mean age = 13.1 years, SD = 3.4, range = 6-17), and

twenty-one children diagnosed with autism spectrum disorder (ASD, mean age = 13.5, SD = 2.7,

range 7-17) participated in this study (table 1). All participants met an IQ cutoff of full-scale IQ

$\geq$ 70, as measured by the Kaufman Brief Intelligence Test II (KBIT, Kaufman & Kaufman,

2004). Participants were right-handed except two ASD individuals, determined using the Dean

Questionnaire (Dean, 1988). All participants had normal hearing, defined as pure tone thresholds

better than or equal to 15 dB in both ears at octave frequencies between 250 Hz and 8 kHz, tested

in a soundproof room using an audiometer. Informed consent was provided by parents or legal

guardians, in addition to participants' assent, according to protocols approved by the Institutional

Review Board of the Massachusetts General Hospital.

All ASD individuals had a formal clinical diagnosis and met criteria on the diagnostic

Autism Diagnosis Observation Schedule, Version 2, (ADOS-2, Lord et al., 2000), administered

by trained research personnel with demonstrated inter-rater reliability, the diagnostic Social

Communication Questionnaire, Lifetime Version (SCQ Lifetime, Rutter et al., 2003), and

quantitative Social Responsiveness Scale (SRS, (Constantino & Gruber, 2012)). All participants

were also administered the Sensory Profile Questionnaire (ASPS) (Brown & Dunn, 2002). ASD

participants who met ADOS-2 criteria but had a borderline ADOS-2 score, did not meet a cutoff

of ≥ 15 on the SCQ, or a total t-score < 59 on the SRS, were further evaluated by expert clinician

(RMJ) to confirm the ASD diagnosis. A diagnosis of Fragile X syndrome, tuberous sclerosis, and

other known risk factors, such as gestation for less than 36 weeks, were criteria for exclusion

from participation in the study.

All TD subjects were below SCQ thresholds. Parental report was used to rule out the

possibility of any neurological/psychiatric condition (including psychotic episodes, seizures,

clinical EEG abnormalities, Tourette syndrome, or any other tics), or substance abuse in the last

six months. Participants with Tourette syndrome or tic disorders were excluded from both

groups. Due to high comorbidity with ASD, participants with a diagnosis of ADHD, depression,

anxiety or language delay were excluded from the TD group, but not from the ASD group.

| | ASD (N = 21, 4 females) | | | TD (N = 31, 4 females) | | | |
	N	Mean (SD)	Range	N	Mean (SD)	Range	p-value
Age	21	13.5 (2.7)	7-17	31	13.1 (3.4)	6-17	0.640
NVIQ	17	102.8 (15.1)	59-130	20	112.4 (16.4)	83-147	0.083
VIQ	17	98.8 (19.1)	61-139	20	114.2 (13.8)	86-139	0.009
ADOS	21	11.2 (3.1)	7-18	-	-	-	-
SCQ	19	18.0 (8.9)	0-31	30	3.4 (3.2)	0-12	< 0.001
SRS	20	71.9 (16.0)	40-95	29	43.4 (6.0)	34-62	< 0.001
ASPS	20	12.2 (4.1)	5-20	30	17.2 (2.5)	12-20	< 0.001

Table 1. Characterization of participants. P-values are shown for independent samples t-tests for the difference of means across groups. NVIQ (non-verbal IQ), VIQ (Verbal IQ), ADOS (Autism Diagnostic Observation Schedule), SCQ (Social Communication Questionnaire), SRS (Social Responsiveness Scale), ASPS (Auditory Sensory Profile Score).

Stimuli and procedures

The auditory stimuli consisted of 1200 ms broadband (0.5-2.5 kHz) noise bursts (Figure 1a), presented in 200 trials. Stimuli were designed with a 400 microsecond Interaural Time Delay (ITD) between the stereo channels, thus generating a left- or right-lateralized percept depending on the direction of the ITD. In each trial, the broadband noise was amplitude modulated at either 25 Hz or 43 Hz (Ahlfors et al., 2024). To manipulate the ITD, discontinuities of ± 400 µs were introduced in the noise carriers in each channel 550 ms after stimulus onset. Crucially, the discontinuities were designed to coincide with a trough of the amplitude modulation to render them monaurally imperceptible. In half the trials, the 400- µs discontinuities occurred in the same direction in both ears, thus holding the ITD constant for the full duration of the trial (Figure 1b). For the other half of the trials, carrier discontinuities were introduced in opposite directions, thus switching the ITD polarity and creating a percept of a spatial jump in the location of the sound from one hemifield to the contralateral side midway through the trial (Figure 1c).

The subjects listened passively to the auditory stimuli while watching a silent movie with no subtitles. Participants were explicitly asked to ignore the auditory stimuli. After rejection of bad epochs (see below for details), the mean total number of trials was 177 per participant. The mean number of trials across groups did not differ significantly (ASD jump = 181.6, SD = 20.2; ASD stay = 180.0, SD = 22.6; TD jump = 175.5, SD = 19.8, TD stay = 176.2, SD = 19.2). The

140 interstimulus interval was uniformly distributed between 300 and 400 ms. Stimuli were

141 presented at 65 dB-SPL via MEG compatible earphones.

Figure 1. Stimuli and schematic representation of conditions. **a.** Spectrogram (top) and soundwave (bottom) of the auditory stimuli in the stay condition. The red shaded area in the bottom panel represents the time window from which panels in b and c (top) were extracted. **b.** For stay events, the directionality of the ITD was kept constant between the two stereo channels (top). Stimuli were presented with either left- or right-leading ITD (bottom), with equal probability. **c.** For jump events, the polarity of the ITD was reversed at 550ms (dotted line, top). The sound was initially presented with either left- or right-leading ITD, with equal probability. The ITD reversal created a percept of a spatial jump in the lateralization of the sound, right-to-left (R2L) or left-to-right (L2R) side (bottom).

MEG Data acquisition

MEG data was acquired using a 306-channel Vector-View Magnetoencephalography system with

204 planar gradiometer and 102 magnetometer sensors (MEGIN Oy, Finland) inside a

magnetically shielded room (IMEDCO, Switzerland). Signals were bandpass filtered between

0.1 and 330 Hz and sampled at the rate of 1,000, or 3,000 Hz and then downsampled to 1000

Hz. Head position was recorded during data acquisition using four head position indicator (HPI)

coils attached to the scalp. The location of the HPI coils, the nasion, preauricular points, and

additional points in the scalp surface were digitized using a Fastrack digitizer (Polhemus)

system. The digitization information was used to co-register the MEG and MRI data for source

modeling analyses. Electrocardiography (ECG) and electro-oculography (EOG) electrodes were

used to detect heartbeat, saccades, and blinks. Five minutes of empty-room data were recorded at

the end of the session to measure the instrumentation and environmental noise levels in the MEG

sensors on the data acquisition date.

Structural MRI

T1-weigthed magnetization-prepared rapid gradient echo (MPRAGE) structural images were

acquired using a 3T Siemens MRI scanner and a 32-channel head coil. Reconstruction of cortical

surface, cortical volume and anatomical parcellation was conducted using FreeSurfer (Dale et al.,

1999; Fischl, Sereno, & Dale, 1999).

MEG data preprocessing

MEG data was preprocessed and analyzed using MNE-Python (Gramfort et al., 2013). Noisy

MEG channels were visually identified and excluded from further analyses. Temporal Signal

Space Separation (tSSS, Taulu & Simola, 2006) was implemented to compensate for head

movements and minimize noise from sources outside of the virtual sphere circumscribing the

MEG sensors. Next, data was bandpass filtered between 0.1 and 144 Hz, and notch filters were

applied at 60 Hz and 120 Hz to remove line noise and its harmonics. Signal Space Projection

(SSP, Uusitalo & Ilmoniemi, 1997) was used to remove cardiac and oculomotor artifacts by

computing projections from the bandpass filtered (1-35 Hz) signal. The data were then epoched

from -400 ms to 1400 ms with respect to sound onset and detrended to remove slow drifts.

Epochs were excluded if the signal exceeded a peak-to-peak threshold of 4000 fT for

magnetometer channels and 5000 fT/cm for planar gradiometer channels.

Source localization

The planar gradiometer signals were used to calculate source estimates on the FreeSurfer-

reconstructed cortical surfaces for each individual. The forward solution was computed using a

single-compartment boundary element model (BEM, Hamalainen & Sarvas, 1989). Inner skull

surface triangulations were generated from MRI structural data using the watershed algorithm.

The cortical current distributions were obtained using the dynamic Statistical Parametric

Mapping (dSPM, Dale et al., 2000) with a loose orientation constraint of 0.2 and depth weighting

of 0.8 (Lin et al., 2006). To reduce the contribution of instrumentation and environmental

artifacts to the MEG signals, a noise covariance matrix was obtained from empty room

recordings. The noise covariance matrix was used to construct the inverse operator (Hämäläinen

& Ilmoniemi, 1994) with a regularization parameter set to 0.1, and to noise-normalize the source

estimates (Dale et al., 2000).

Regions of interest

Regions of interest (ROIs) in auditory cortex were identified for each subject's anatomical

surface reconstruction by visually examining the time course activations in response to ITD

reversals. ROIs corresponding to peak earliest activations were then manually delineated for

each individual participant, in both the right and the left hemisphere. Figure 2 shows the overlap

probability for left and right auditory ROIs across all subjects. ROI size was not significantly

different across groups (ASD mean = 63.1 vertices, TD mean = 64.5 vertices, t = -0.21, p = 0.83.

Table S1, Supplemental Materials). Evoked responses were computed by extracting the mean

dSPM activation across all vertices within each auditory ROI.

In addition to auditory areas, studies of spatial auditory attention using fMRI reported the

recruitment of a frontoparietal network during auditory spatial attention tasks. Areas within this

network include the dorsolateral Prefrontal Cortex (dlPFC), the ventrolateral Prefrontal Cortex

(vlPFC), the precentral/central sulci, the pre/postcentral gyri, the Superior Parietal Lobe (SPL)

and the Inferior Parietal Lobe (IFL). These target regions were therefore grouped into frontal,

central and parietal ROIs for subsequent functional connectivity analyses (Table S2,

supplemental materials), using Freesurfer cortical parcellations (Destrieux et al., 2010).

Figure 2. Anatomical overlap across all subjects for the manually drawn, left and right auditory ROIs. These ROIs were identified on the basis of cortical activations in response to ITD reversals in the Jumps condition. Overlap is presented in a common cortical space (Freesurfer's fsaverage cortical surface).

Functional connectivity analyses

We computed seed-based functional connectivity for each condition (*stay* and *jump*) from the left

and right auditory cortices with the regions of interest outlined above, including the dlPFC and

vlPFC (superior, middle and inferior prefrontal sulci, middle and inferior prefrontal gyri), central

regions (precentral and central sulci, precentral and postcentral gyri) and parietal regions

(postcentral and intraparietal sulci, superior and inferior parietal sulci), bilaterally. The seed time

courses for functional connectivity analyses corresponded to the auditory cortical activations in

response to *jump* trials. Functional connectivity was quantified using the Phase Lag Index (PLI),

a measure of phase synchronization that penalizes zero-lag estimates, thus effectively controlling

for spurious connectivity due to field spread of MEG signals (Stam et al., 2007). Data was

resampled to 250 Hz, and the PLI was computed for each trial between the right and left seed

ROI time course and the time course at each vertex, using the wavelet method (spanning four

cycles) between 4 and 30 Hz, for a 2000-ms window between -400 and 1400 ms. PLI estimates

in the jump condition were then normalized with respect to the *stay* condition to obtain a *z*-

scored PLI value, hereafter referred to as zPLI (see supplementary materials). The zPLI estimate

is a *z* value which reflects the extent to which connectivity during *jump* trials deviates with

respect to the baseline condition (i.e. *stay*), while additionally controlling for the non-normal

distribution of connectivity estimates and the difference in trial number across subjects and

conditions (Khan et al., 2013; Maris et al., 2007).

Cluster-based permutation statistics within anatomical ROIs

Clusters of vertices showing a significant difference in connectivity across groups within the

target regions of interest were identified using cluster-based permutation statistics (Maris &

Oostenveld, 2007). This test was conducted for the alpha band (8 – 12 Hz) by averaging

functional connectivity estimates over the time dimension between 550 and 850 ms, a time

window corresponding to the onset and offset of brain activation to *jump* trials. This statistical

test was implemented for each seed (left and right) and target ROI (prefrontal, central and

parietal) separately, using a one-way ANOVA with a cluster-forming threshold of $F = 10.54$

(corresponding to a p value = 1e-06) and 1000 permutations. Clusters were identified based on

spatial adjacency. To achieve this, data from each participant was morphed to a FreeSurfer

average cortical representation with 10,242 vertices per hemisphere (Fischl, Sereno, Tootell, et

al., 1999). Cluster size was then estimated for each surviving cluster and results were filtered to

keep only the top 5% biggest clusters. Finally, clusters were sorted by their statistical

significance, which was obtained by summing F values at each vertex within the formed cluster,

and only the most significant cluster per seed and anatomical ROI was considered for further

analyses. As a post-hoc control analysis, we repeated the same functional connectivity analysis

pipeline using Coherence, and repeated the same cluster-based statistical analyses using the

coherence metrics instead of the PLI metrics. This was done to ensure the robustness of cluster

identification approach. This resulted in two ROIs that showed significant group difference

across both functional connectivity metrics, and those are the ones reported in the results section.

Cluster based permutation statistics were implemented using MNE-python (Gramfort et al.,

2013).

Linear mixed effect modelling

Brain responses to *jump* trials were investigated using Linear mixed effect (LME) modeling with

the LME4 package (Bates et al., 2015) in R version 2023.6.1. The predicted variable was the

brain response to *jump* trials, quantified as the Area Under the Curve (AUC) at the activation

post jump onset. For this, the onset and offset latencies of brain activations with respect to peak

latencies in response to *jump* trials were visually identified and recorded for each subject. This

time window did not significantly differ across groups (t=1.70, p =0.091. Table S3,

Supplementary Materials). The AUC for *stay* trials was obtained using the same subject-specific

time window as for *jump* trials. Fixed effects included group (ASD, TD), condition (*stay*, *jump*),

and hemisphere (left, right). To control for the within-subject nature of the task, a random effect

for subject was included.

Brain-behavior correlations

Spearman correlation was estimated between normalized functional connectivity estimates

extracted from the two surviving statistically significant ROIs that showed significant group

differences (see Results) and the two quantitative behavioral measures: the Social

Responsiveness Scale (SRS) total t score, a quantitative measure of Autism severity, and the

Auditory Sensory Profile Score (ASPS), a quantitative measure of auditory sensory processing in

ASD. The ASPS was quantified as the average of the first five questions of the auditory section

of the Sensory Profile (parent report), which measures auditory processing atypicality in terms of

sensitivity to and distraction by noise. These questions are categorically different from questions

6-8, which are related to social communication (e.g., ignoring what parents say or not responding

to their name being called, question 6-7) and to enjoying or seeking to produce strange noises

question 8), rather than to more basic auditory processing. Multiple comparisons were addressed

by adjusting *p* values using the False Discovery Rate (FDR).

Results

Evoked responses in the auditory cortex

Both ASD and TD individuals showed stronger activations during *jump* trials compared

to *stay* trials (i.e., a jump-evoked response). AUC values between onset and offset of peak

activations during *jump* trials for both conditions are presented in Table 1.

	AUC - Jump				AUC - Stay			
	Left		Right		Left		Right	
	Mean	*SD*	*Mean*	*SD*	*Mean*	*SD*	*Mean*	*SD*
TD	1267.09	835.24	1321.34	864.35	538.77	451.13	532.25	382.52
ASD	845.09	730.18	950.64	434.52	430.63	326.76	503.19	301.30

Table 1. Area Under the Curve (AUC) for TD and ASD individuals at peak between onset and offset of brain activations to *jump* trials.

Figure 3 shows the mean source time course for ASD and TD individuals in left and right auditory ROIs. Mixed effect modeling confirmed a significant effect of group ($F(194.10) = 7.84$, $p = 0.006$), condition ($F(165.53) = 52.31$, $p = 1.67e-11$) as well as a statistically significant interaction between group and condition ($F(165.53) = 3.97$, $p = 0.048$). There was no significant effect of hemisphere in predicting the AUC in brain activations ($F(165.53) = 0.47$, $p = 0.49$). Post-hoc pairwise comparisons showed that, within each group, mean AUC values for *jump* trials were significantly higher than *stay* trials (ASD: $t(171) = 3.39$, $p = 9.0e-04$, TD: $t(171) = 7.26$, $p < 1.0e-4$, insets in Figures 3a and 3b). Across groups, AUC values during *jump* trials were significantly higher for the TD group compared to the ASD group ($t(185) = 3.39$, $p = 9.0e-05$) in both left (Figure 3c) and right (Figure 3d) auditory ROIs. Maximum Likelihood Estimation analyses confirmed that the model outperformed a null (intercept-only) model in predicting evoked responses to jumps $\chi^2(7) = 62.78$, $p = 4.19e-12$, BIC $= 3209.95$, $R^2_{conditional} = 0.32$, $R^2_{marginal} = 0.24$).

Activations in response to ITD reversals peaked on average at 734 ms in the left auditory cortex (SD = 52 ms) and at 724 ms in the right auditory cortex (SD = 48 ms) in the ASD group. In the TD group, the left hemisphere responses peaked at 742 ms (SD = 50 ms) whereas and the right hemisphere responses peaked at 739 ms (SD = 51 ms). There was no statistically significant difference in peak latency across the two groups.

Functional connectivity between auditory areas and frontoparietal ROIs

Next, we investigated whether the spatial jump was also associated with different alpha-band

functional connectivity dynamics, using zPLI, which corresponds to the PLI estimates for the

jump condition normalized by PLI for the *stay* condition. The seed time courses were the

bilateral activations observed in left and right auditory cortices within the manually outlined

ROIs, and the target regions were the prefrontal, central and parietal anatomical ROIs defined on

the basis of previously reported fMRI findings (see Methods and Supplemental Materials). Based

on our prior studies where across-group differences were modulated by functional connectivity

in the alpha (8-12 Hz) band (Alho, Khan, et al., 2023; Alho, Samuelsson, et al., 2023; Khan et

al., 2013; Mamashli et al., 2021), here too we chose to focus on zPLI estimates in the alpha

frequency between 550 and 850 ms, a time window corresponding to the onset-to-offset of the

evoked response to the jump event.

Figure 3. Evoked activations in auditory cortical ROIs. **a.** and **b.** Activations in the left (**a**) and right (**b**) auditory cortices for the TD (green) and ASD (purple) groups in response to jump (solid lines) and stay (dotted lines) trials. Solid vertical line at time zero represents sound onset. Dotted vertical line at 0.55 s represents the onset of and ITD reversal in the jump condition. Gray shades indicate the 95% confidence interval. **c.** and **d.** Mean AUC values for *jump* trials for the TD (green) and ASD (purple) groups. Whiskers indicate the Standard Error of the Mean (SEM). *** p < 0.001, ** p < 0.01, * p < 0.05.

Non-parametric cluster-based permutations tests showed two statistically significant clusters

where alpha-band connectivity differed between the two groups. The first cluster ($n_{vertices}$ = 16, F

= 44.29, p = 0.0010, Figure 4a, left panel) showed significant connectivity differences across

groups between the left auditory seed and dorsolateral portions of the prefrontal cortex (dlPFC).

Within this cluster, alpha-band connectivity was significantly lower during *jump* trials compared

to *stay* trials (median zPLI = -0.037, SD = 0.022) for TD individuals (t (30) = -10.28, p = 1.23e-

10, Figure 4b, left),), but higher during *jump* trials compared to *stay* trials (median zPLI = 0.021,

SD = 0.027) for the ASD group (t (20) = -2.71, p = 0.015, Figure 4b, right). Additionally,

prefrontal zPLI estimates significantly differed across groups (t(50) = -8.25, p = 1.23e-10, Figure

4b).

The second cluster ($n_{vertices}$ = 17, F = 67.25, p = 0.0010, Figure 4c, left panel) showed significant

connectivity differences across groups between the left auditory seed and portions of the left

superior postcentral sulcus and superior parietal gyrus. Within this cluster, alpha-band

connectivity was significantly lower during *jump* trials compared to *stay* trials (median zPLI = -

0.06, SD = 0.02) for TD individuals (t (30) = -16.38, p = 1.49e-15, Figure 4d, left), but zPLI

estimates during *jump* trials did not significantly differ from the *stay* condition (t(20) = 1.34, p =

0.19) for the ASD group (median zPLI = 0.01, SD = 0.02, Figure 4d, right).). Parietal functional

connectivity within this parietal cluster was also significantly different across groups (t(50) = -

11.19, p = 1.44e-14, Figure 4d). All reported *p* values from comparisons of MEG measures

between the two groups are FDR-adjusted.

Figure 4. Functional connectivity for the two statistically significant clusters during *jump* trials, normalized by *stay* trials. **a**. and **c**. Anatomical location of the statistically significant clusters in prefrontal (**a**) and the superior parietal regions (**c**). **b**. and **d**. Median alpha-band connectivity estimates obtained from the dlPFC (**b**) and SPL clusters (**d**) for the TD (green) and ASD (purple) groups. Whiskers represent the Standard Error (SE). Asterisks on top of horizontal lines denote significant differences across groups. Asterisks below horizontal lines denote a

Correlations between brain measures and behavioral measures

To test whether the MEG measures correspond to relevant ASD traits, we correlated the

magnitude of the evoked responses, as well as estimates of normalized functional connectivity

(zPLI) with behaviorally assessed phenotypic measures of ASD severity (SRS t-score) and

auditory processing, (the ASPS score). In the TD group, we did not find significant associations

between any of the behavioral measures and the evoked responses. Similarly, neither behavioral

measure was significantly correlated with zPLI in the TD group.

For the ASD group, higher alpha band functional connectivity in the prefrontal cluster showed

positive correlation with autism severity, as measured by the SRS t-score (r(18) = 0.51, p = 0.021

uncorrected, q = 0.08), although this correlation did not survive multiple comparisons correction.

Prefrontal connectivity in the alpha band was also negatively correlated with the auditory

processing component of the Sensory Profile Questionnaire in the ASD group (r(18) = - 0.63, p =

0.003, q = 0.025). The correlations between the behavioral measures and zPLI are shown in

Figure 5. To check for any possible effects of age, we computed post-hoc correlations between

age and functional connectivity estimates, and between age and cortical activations. No

significant correlation was found between age and these two MEG measures of brain function,

meaning age was not a factor in the response patterns and observed group differences.

Figure 5. Brain-behavior correlations. **a**. and **b**. Correlation between alpha-band connectivity estimates obtained from the prefrontal cluster and scores in the Social Responsiveness Scale, SRS (**a**) and Auditory Sensory Profile Scores, ASPS (**b**) in the TD group (green). **c**. and **d**. Correlation between alpha-band connectivity in the prefrontal cluster and SRS (**c**) and ASPS scores (**d**) in the ASD group (purple). q values show the adjusted p values after FDR correction.

Discussion

The aim of this study was to investigate passive auditory spatial attention in ASD. Specifically,

we examined cortical activations and alpha-band functional connectivity in individuals with ASD

and TD individuals in response to changes in sound location (i.e., *jump* trials) under passive

listening conditions. ASD individuals showed lower cortical activations in cortical auditory

regions and altered alpha frequency band functional connectivity between auditory and frontal

areas. Additionally, functional connectivity estimates for the ASD group were significantly

correlated with behavioral measures of ASD severity. Our findings provide insight into the

differential neural processing of spatial auditory information in ASD and TD groups.

ASD individuals show reduced auditory activation during jump trials compared to TDs

The fact that both groups showed cortical responses to *jump* trials confirms that both ASD and

TD individuals were sensitive to acoustic cues indicative of a change in the location of a sound

in space. This would be in line with findings which suggest that despite relative differences in

performance, both groups can behaviorally locate sounds in space (Lodhia et al., 2018; Soskey et

al., 2017; Teder-Sälejärvi et al., 2005). While both groups detected changes in localization,

auditory cortical activations were significantly lower in the ASD group, which may be indicative

of altered encoding of perceived acoustic features that are important to map sounds in space.

These results are in line with previous electrophysiological research, which has shown a smaller

amplitude of early and mid-latency auditory components during spatial (Lodhia et al., 2018;

Teder-Sälejärvi et al., 2005) and non-spatial auditory processing tasks (Bruneau et al., 1999;

Jansson-Verkasalo et al., 2005; Orekhova et al., 2009; Port et al., 2015; Russo et al., 2009;

Soskey et al., 2017). They are also in line with postmortem human evidence in ASD of

malformed brainstem circuits (i.e., superior olive) that are thought to be the first nuclei along the

ascending auditory pathway to process binaural cues, such as interaural time delays (Kulesza et

al., 2011). The significant group-by-condition interaction, where the ASD group exhibited a

smaller magnitude of activation during *jump* trials, further supports the notion that individuals

with ASD exhibit atypical auditory processing, potentially resulting from altered encoding or

binding of relevant acoustic features (Bharadwaj et al., 2022; Gonçalves & Monteiro, 2023;

Hames et al., 2016; Keehn et al., 2010, 2017; Lodhia et al., 2018; Marco et al., 2011).

ASD and TD Groups show opposite connectivity patterns in response to jump trials

Our functional connectivity analyses revealed significant differences in temporo-frontal and

temporo-parietal alpha-band connectivity between ASD and TD groups, with the ASD group

showing stronger connectivity during *jump* trials relative to the TD group. These results suggest

that the ASD group may lack the typical modulation of connectivity in response to sensory

changes. This could reflect a disruption in the dynamic adjustment of attentional networks

required for processing auditory spatial shifts, especially given that this is a passive paradigm,

where attention was presumably directed away from the sounds. Atypical functional connectivity

involving the auditory cortex, measured with EEG or MEG, has been documented in ASD, by

our group and others (Alho, Khan, et al., 2023; Alho, Samuelsson, et al., 2023; Demopoulos et

al., 2024; Mamashli et al., 2017). Long-range alpha-band functional connectivity has been

previously proposed as a neurophysiological marker of attentional modulation via coordination

of temporal windows of neural excitability across cortical space (Klimesch et al., 2007; Palva &

Palva, 2011; Sadaghiani & Kleinschmidt, 2016). Higher alpha-band connectivity in the ASD

group could therefore be associated with atypical synchronization of attentional networks

involved in sensory integration and cognitive control, which are often reported as dysregulated in

ASD. The lower connectivity observed in the TD group could reflect a more typical response to

the auditory shift, suggesting more efficient attentional allocation and neural processing.

Although not related to functional connectivity, previous studies on auditory spatial attention in

ASD have reported lower amplitude of mid-lantency ERP components associated with attention

and executive function, namely the P300 and P400 (Lodhia et al., 2018; Teder-Sälejärvi et al.,

2005). This lends additional support to the idea that, on top of sensory encoding, attentional

processes also differ in ASD during execution of auditory spatial attention tasks.

Prefrontal alpha connectivity in ASD correlates with auditory processing and autism severity

We found that higher temporo-parietal alpha band connectivity was positively correlated with

higher manifestations of ASD symptoms as measured by the Social Responsiveness Scale (SRS).

Although this correlation did not survive correction for multiple comparisons, the trend is

consistent with other studies showing a correlation between functional connectivity of auditory

cortex with both cerebral and cerebellar ROIs and ASD symptoms severity (Alho, Samuelsson,

et al., 2023; Sharda et al., 2018; Wilson et al., 2022).

Additionally, higher alpha-band connectivity between auditory and the prefrontal regions

was negatively correlated with behaviorally assessed auditory processing symptoms, as

measured by the Auditory Sensory Profile Score (ASPS) in the ASD group, potentially indicating

that higher connectivity is linked to poorer auditory integration of spatial cues. This is also in

line with previous work in our group, showing that increased connectivity between auditory and

parietal regions is associated with poorer auditory processing skills, as measured by the ASPS

(Alho, Khan, et al., 2023).

The finding of higher alpha-band functional connectivity in ASD during spatial attention

tasks is in line with similar findings observed in resting state which propose alpha-band

hyperconnectivity as a characteristic feature of ASD (Demopoulos et al., 2024; Orekhova et al.,

2014; Wang et al., 2020). These findings highlight the potential impact of prefrontal and parietal

regions on auditory processing in ASD.

Limitations

This study has two limitations that should be addressed in future research. First, the relatively

small sample sizes of the ASD and TD groups limit the power of the findings. While the findings

align well with prior studies, future studies with larger cohorts will be important for further

confirming the observed group differences in cortical activations and connectivity patterns, as

well as to potentially uncover additional differences. Second, because our results were obtained

during passive listening conditions, we lack task-relevant behavioral correlates that allow us to

rule out the possibility that participants with ASD actively engaged auditory attention differently

from the TD participants. Future studies with an active spatial auditory attention task and

behavioral assessments of spatial auditory attention will be needed to elucidate any behavioral

differences between the groups, and links between these differences and cortical measures.

Conclusion

The results presented here provide novel insights into the neural mechanisms underlying auditory

spatial attention in ASD. We found that ASD was associated with lower auditory cortical

activations and altered alpha-band functional connectivity between auditory cortex and frontal

and parietal regions, in response to shifts in the spatial location of sounds. The

neurophysiological measures were correlated with measures of ASD symptoms extent and

behaviorally assessed auditory processing, in the ASD group, thus confirming their relevance to

the ASD phenotype. Future directions include the evaluation of these metrics with an active,

rather than passive, auditory spatial attention paradigm, and extending it to larger samples as

well as to related conditions, so as to elucidate the specificity of the results to ASD.

Conflict of interest

The authors declare no conflict of interest.

References

Ahlfors, S. P., Graham, S., Bharadwaj, H., Mamashli, F., Khan, S., Joseph, R. M., Losh, A.,
476   Pawlyszyn, S., McGuiggan, N. M., Vangel, M., Hämäläinen, M. S., & Kenet, T. (2024). No
477   Differences in Auditory Steady-State Responses in Children with Autism Spectrum
478   Disorder and Typically Developing Children. Journal of Autism and Developmental
479   Disorders, 54(5), 1947–1960. https://doi.org/10.1007/s10803-023-05907-w

480   Alho, J., Bharadwaj, H., Khan, S., Mamashli, F., Perrachione, T. K., Losh, A., McGuiggan, N.
481   M., Joseph, R. M., Hämäläinen, M. S., & Kenet, T. (2021). Altered maturation and atypical
482   cortical processing of spoken sentences in autism spectrum disorder. *Progress in*
483   *Neurobiology*, *203*. https://doi.org/10.1016/j.pneurobio.2021.102077

484   Alho, J., Khan, S., Mamashli, F., Perrachione, T. K., Losh, A., McGuiggan, N. M., Graham, S.,
485   Nayal, Z., Joseph, R. M., Hämäläinen, M. S., Bharadwaj, H., & Kenet, T. (2023). Atypical
486   cortical processing of bottom-up speech binding cues in children with autism spectrum
487   disorders. *NeuroImage: Clinical*, *37*. https://doi.org/10.1016/j.nicl.2023.103336

488   Alho, J., Samuelsson, J. G., Khan, S., Mamashli, F., Bharadwaj, H., Losh, A., McGuiggan, N.
489   M., Graham, S., Nayal, Z., Perrachione, T. K., Joseph, R. M., Stoodley, C. J., Hämäläinen,
490   M. S., & Kenet, T. (2023). Both stronger and weaker cerebro-cerebellar functional
491   connectivity patterns during processing of spoken sentences in autism spectrum disorder.
492   *Human Brain Mapping*, *44*(17), 5810–5827. https://doi.org/10.1002/hbm.26478

493   Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2015). Fitting linear mixed-effects
494   models using lme4. *Journal of Statistical Software*. https://doi.org/10.18637/jss.v067.i01

495   Ben-Sasson, A., Gal, E., Fluss, R., Katz-Zetler, N., & Cermak, S. A. (2019). Update of a Meta-
496   analysis of Sensory Symptoms in ASD: A New Decade of Research. *Journal of Autism and*
497   *Developmental Disorders*, *49*(12), 4974–4996. https://doi.org/10.1007/s10803-019-04180-0

498   Bharadwaj, H., Mamashli, F., Khan, S., Singh, R., Joseph, R. M., Losh, A., Pawlyszyn, S.,
499   McGuiggan, N. M., Graham, S., Hämäläinen, M. S., & Kenet, T. (2022). Cortical signatures
500   of auditory object binding in children with autism spectrum disorder are anomalous in

concordance with behavior and diagnosis. *PLoS Biology*, *20*(2). https://doi.org/10.1371/journal.pbio.3001541

Brown, C., & Dunn, W. (2002). *Adolescent/adult sensory profile*. Pearson.

Bruneau, N., Roux, S., Louis Adrien, J., & Barthelemy, C. (1999). Auditory associative cortex dysfunction in children with autism: evidence from late auditory evoked potentials (N1 wave-T complex). *Clinical Neurophysiology*, *110*, 1927–1934. www.elsevier.com/locate/clinph

Constantino, J. N., & Gruber, C. P. (2012). *Social responsiveness scale: SRS-2*. Western psychological services.

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). *Cortical Surface-Based Analysis I. Segmentation and Surface Reconstruction*. http://www.idealibrary.com

Dale, A. M., Liu, A. K., Fischl, B. R., Buckner, R. L., Belliveau, J. W., Lewine, J. D., & Halgren, E. (2000). Dynamic statistical parametric mapping: Combining fMRI and MEG for high-resolution imaging of cortical activity. *Neuron*, *26*, 55–67.

Dean, R. S. (1988). Lateral preference schedule. In *Lateral preference schedule*. Psychological Assessment Resources Incorporated.

Degerman, A., Rinne, T., Salmi, J., Salonen, O., & Alho, K. (2006). Selective attention to sound location or pitch studied with fMRI. *Brain Research*, *1077*(1), 123–134. https://doi.org/10.1016/j.brainres.2006.01.025

Demopoulos, C., Jesson, X., Rae Gerdes, M., Jurigova, B. G., Hinkley, L. B., Ranasinghe, K. G., Desai, S., Honma, S., Mizuiri, D., Findlay, A., Nagarajan, S. S., & Marco, E. J. (2024). Global MEG resting state functional connectivity in children with autism and sensory processing dysfunction. *BioRxiv*. https://doi.org/10.1101/2024.01.26.577499

Deouell, L. Y., Heller, A. S., Malach, R., D'Esposito, M., & Knight, R. T. (2007). Cerebral Responses to Change in Spatial Location of Unattended Sounds. *Neuron*, *55*(6), 985–996. https://doi.org/10.1016/j.neuron.2007.08.019

DePape, A. M. R., Hall, G. B. C., Tillmann, B., & Trainor, L. J. (2012). Auditory Processing in High-Functioning Adolescents with Autism Spectrum Disorder. *PLoS ONE*, *7*(9). https://doi.org/10.1371/journal.pone.0044084

Destrieux, C., Fischl, B., Dale, A., & Halgren, E. (2010). Automatic parcellation of human cortical gyri and sulci using standard anatomical nomenclature. *NeuroImage*, *53*(1), 1–15. https://doi.org/10.1016/j.neuroimage.2010.06.010

Fischl, B., Sereno, M. I., & Dale, A. M. (1999). *Cortical Surface-Based Analysis II: Inflation, Flattening, and a Surface-Based Coordinate System*. http://www.idealibrary.com

Fischl, B., Sereno, M. I., Tootell, R. B. H., & Dale, A. M. (1999). High-resolution intersubject averaging and a coordinate system for the cortical surface. *Human Brain Mapping*, *8*(4), 272–284. https://doi.org/10.1002/(SICI)1097-0193(1999)8:4<272::AID-HBM10>3.0.CO;2-4

Fritz, J. B., Elhilali, M., David, S. V., & Shamma, S. A. (2007). Auditory attention - focusing the searchlight on sound. In *Current Opinion in Neurobiology* (Vol. 17, Issue 4, pp. 437–455). https://doi.org/10.1016/j.conb.2007.07.011

Ghanbari, Y., Bloy, L., Christopher Edgar, J., Blaskey, L., Verma, R., & Roberts, T. P. L. (2015). Joint analysis of band-specific functional connectivity and signal complexity in autism. *Journal of Autism and Developmental Disorders*, *45*(2), 444–460. https://doi.org/10.1007/s10803-013-1915-7

Gonçalves, A. M., & Monteiro, P. (2023). Autism Spectrum Disorder and auditory sensory alterations: a systematic review on the integrity of cognitive and neuronal functions related to auditory processing. In *Journal of Neural Transmission* (Vol. 130, Issue 3, pp. 325–408). Springer. https://doi.org/10.1007/s00702-023-02595-9

Gramfort, A., Luessi, M., Larson, E., Engemann, D. A., Strohmeier, D., Brodbeck, C., Goj, R., Jas, M., Brooks, T., Parkkonen, L., & Hämäläinen, M. (2013). MEG and EEG data analysis with MNE-Python. *Frontiers in Neuroscience*, *7 DEC*. https://doi.org/10.3389/fnins.2013.00267

Grzadzinski, R., Huerta, M., & Lord, C. (2013). DSM-5 and autism spectrum disorders (ASDs): An opportunity for identifying ASD subtypes. In *Molecular Autism* (Vol. 4, Issue 1). https://doi.org/10.1186/2040-2392-4-12

Haesen, B., Boets, B., & Wagemans, J. (2011). A review of behavioural and electrophysiological studies on auditory processing and speech perception in autism spectrum disorders. In *Research in Autism Spectrum Disorders* (Vol. 5, Issue 2, pp. 701–714). https://doi.org/10.1016/j.rasd.2010.11.006

Haist, F., Adamo, M., Westerfield, M., Courchesne, E., & Townsend, J. (2005). The Functional Neuroanatomy of Spatial Attention in Autism Spectrum Disorder. In *DEVELOPMENTAL NEUROPSYCHOLOGY* (Vol. 27, Issue 3).

Hämäläinen, M. S., & Ilmoniemi, R. J. (1994). Interpreting magnetic fields of the brain: minimum norm estimates. Medical & Biological Engineering & Computing, 32(1), 35–42. https://doi.org/10.1007/bf02512476

Hämäläinen, M. S., & Sarvas, J. (1989). Realistic Conductivity Geometry Model of the Human Head for Interpretation of Neuromagnetic Data. In *IEEE TRANSACTIONS ON BIOMEDICAL ENGINEERING* (Vol. 36, Issue 2).

Hames, E. C., Rajmohan, R., Fang, D., Anderson, R., Baker, M., Richman, D. M., & O. Boyle,
  M. (2016). Attentional Networks in Adolescents with High-functioning Autism: An fMRI
  Investigation. *The Open Neuroimaging Journal*, *10*(1), 102–110.
  https://doi.org/10.2174/1874440001610010102

Hedger, N., Dubey, I., & Chakrabarti, B. (2020). Social orienting and social seeking behaviors in
  ASD. A meta analytic investigation. In *Neuroscience and Biobehavioral Reviews* (Vol. 119,
  pp. 376–395). Elsevier Ltd. https://doi.org/10.1016/j.neubiorev.2020.10.003

Jansson-Verkasalo, E., Kujala, T., Jussila, K., Mattila, M. L., Moilanen, I., Näätänen, R.,
  Suominen, K., & Korpilahti, P. (2005). Similarities in the phenotype of the auditory neural
  substrate in children with Asperger syndrome and their parents. *European Journal of
  Neuroscience*, *22*(4), 986–990. https://doi.org/10.1111/j.1460-9568.2005.04216.x

Kana, R. K., Uddin, L. Q., Kenet, T., Chugani, D., & Müller, R. A. (2014). Brain connectivity in
  autism. In *Frontiers in Human Neuroscience* (Vol. 8, Issue JUNE). Frontiers Media S. A.
  https://doi.org/10.3389/fnhum.2014.00349

Kaufman, A. S., & Kaufman, N. L. (2004). *Kaufman Brief Intelligence Test* (Second Edition).
  Pearson Assessment US.

Kayser, C., Petkov, C. I., Lippert, M., & Logothetis, N. K. (2005). Mechanisms for allocating
  auditory attention: An auditory saliency map. *Current Biology*, *15*(21), 1943–1947.
  https://doi.org/10.1016/j.cub.2005.09.040

Keehn, B., Lincoln, A. J., Müller, R. A., & Townsend, J. (2010). Attentional networks in children
  and adolescents with autism spectrum disorder. *Journal of Child Psychology and
  Psychiatry, and Allied Disciplines*, *51*(11), 1251–1259. https://doi.org/10.1111/j.1469-
  7610.2010.02257.x

Keehn, B., Westerfield, M., Müller, R. A., & Townsend, J. (2017). Autism, Attention, and Alpha
  Oscillations: An Electrophysiological Study of Attentional Capture. *Biological Psychiatry:
  Cognitive Neuroscience and Neuroimaging*, *2*(6), 528–536.
  https://doi.org/10.1016/j.bpsc.2017.06.006

Khan, S., Gramfort, A., Shetty, N. R., Kitzbichler, M. G., Ganesan, S., Moran, J. M., Lee, S. M.,
  Gabrieli, J. D. E., Tager-Flusberg, H. B., Joseph, R. M., Herbert, M. R., Hämäläinen, M. S.,
  & Kenet, T. (2013). Local and long-range functional connectivity is reduced in concert in
  autism spectrum disorders. *Proceedings of the National Academy of Sciences of the United
  States of America*, *110*(8), 3107–3112. https://doi.org/10.1073/pnas.1214533110

Klimesch, W., Sauseng, P., & Hanslmayr, S. (2007). EEG alpha oscillations: The inhibition-
  timing hypothesis. *Brain Research Reviews*, *53*(1), 63–88.
  https://doi.org/10.1016/j.brainrev.2006.06.003

Kong, L., Michalka, S. W., Rosen, M. L., Sheremata, S. L., Swisher, J. D., Shinn-Cunningham, B. G., & Somers, D. C. (2014). Auditory spatial attention representations in the human cerebral cortex. *Cerebral Cortex*, *24*(3), 773–784. https://doi.org/10.1093/cercor/bhs359

Kulesza, R. J., Lukose, R., & Stevens, L. V. (2011). Malformation of the human superior olive in autistic spectrum disorders. Brain Research, 1367, 360–371. https://doi.org/10.1016/j.brainres.2010.10.015

Lin, F. H., Witzel, T., Ahlfors, S. P., Stufflebeam, S. M., Belliveau, J. W., & Hämäläinen, M. S. (2006). Assessing and improving the spatial accuracy in MEG source localization by depth-weighted minimum-norm estimates. *NeuroImage*, *31*(1), 160–171. https://doi.org/10.1016/j.neuroimage.2005.11.054

Lodhia, V., Hautus, M. J., Johnson, B. W., & Brock, J. (2018). Atypical brain responses to auditory spatial cues in adults with autism spectrum disorder. *European Journal of Neuroscience*, *47*(6), 682–689. https://doi.org/10.1111/ejn.13694

Lord, C., Risi, S., Lambrecht, L., Cook, E. H. Jr., Leventhal, B. L., DiLavore, P. C., Pickles, A., & Rutter, M. (2000). The Autism diagnostic observation schedule - Generic: A standard measure of social and communication deficits associated with the spectrum of autism. *Journal of Autism and Developmental Disorders*, *30*(3).

Mamashli, F., Khan, S., Bharadwaj, H., Michmizos, K., Ganesan, S., Garel, K. L. A., Ali Hashmi, J., Herbert, M. R., Hämäläinen, M., & Kenet, T. (2017). Auditory processing in noise is associated with complex patterns of disrupted functional connectivity in autism spectrum disorder. *Autism Research*, *10*(4), 631–647. https://doi.org/10.1002/aur.1714

Mamashli, F., Kozhemiako, N., Khan, S., Nunes, A. S., McGuiggan, N. M., Losh, A., Joseph, R. M., Ahveninen, J., Doesburg, S. M., Hämäläinen, M. S., & Kenet, T. (2021). Children with autism spectrum disorder show altered functional connectivity and abnormal maturation trajectories in response to inverted faces. *Autism Research*, *14*(6), 1101–1114. https://doi.org/10.1002/aur.2497

Marco, E. J., Hinkley, L. B. N., Hill, S. S., & Nagarajan, S. S. (2011). *Sensory Processing in Autism: A Review of Neurophysiologic Findings*.

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG-and MEG-data. *The Journal of Neuroscience Methods*, *164*(1), 177–190. https://doi.org/10.1016/j.jneumeth.2007.03.024

Maris, E., Schoffelen, J. M., & Fries, P. (2007). Nonparametric statistical testing of coherence differences. *Journal of Neuroscience Methods*, *163*(1), 161–175. https://doi.org/10.1016/j.jneumeth.2007.02.011

O'Connor, K. (2012). Auditory processing in autism spectrum disorder: A review. In
Neuroscience and Biobehavioral Reviews (Vol. 36, Issue 2, pp. 836–854).
https://doi.org/10.1016/j.neubiorev.2011.11.008

Orekhova, E. V., Elsabbagh, M., Jones, E. J., Dawson, G., Charman, T., Baron-Cohen, S.,
Bedford, R., Bolton, P., Fernandes, J., Ganea, N., Garwood, H., Gliga, T., Hudry, K.,
Murias, M., Ribeiro, H., Tucker, L., Volein, A., & Webb, S. J. (2014). EEG hyper-
connectivity in high-risk infants is associated with later autism. *Journal of*
Neurodevelopmental Disorders, *6*(1). https://doi.org/10.1186/1866-1955-6-40

Orekhova, E. V., Stroganova, T. A., Prokofiev, A. O., Nygren, G., Gillberg, C., & Elam, M.
(2009). The right hemisphere fails to respond to temporal novelty in autism: Evidence from
an ERP study. *Clinical Neurophysiology*, *120*(3), 520–529.
https://doi.org/10.1016/j.clinph.2008.12.034

Palva, S., & Palva, J. M. (2011). Functional roles of alpha-band phase synchronization in local
and large-scale cortical networks. In *Frontiers in Psychology* (Vol. 2, Issue SEP).
https://doi.org/10.3389/fpsyg.2011.00204

Port, R. G., Anwar, A. R., Ku, M., Carlson, G. C., Siegel, S. J., & Roberts, T. P. L. (2015).
Prospective MEG Biomarkers in ASD: Pre-Clinical Evidence and Clinical Promise of
Electrophysiological Signatures.

Port, R. G., Gaetz, W., Bloy, L., Wang, D. J., Blaskey, L., Kuschner, E. S., Levy, S. E., Brodkin,
E. S., & Roberts, T. P. L. (2017). Exploring the relationship between cortical GABA
concentrations, auditory gamma-band responses and development in ASD: Evidence for an
altered maturational trajectory in ASD. *Autism Research*, *10*(4), 593–607.
https://doi.org/10.1002/aur.1686

Roberts, T. P. L., Khan, S. Y., Rey, M., Monroe, J. F., Cannon, K., Blaskey, L., Woldoff, S.,
Qasmieh, S., Gandal, M., Schmidt, G. L., Zarnow, D. M., Levy, S. E., & Edgar, J. C. (2010).
MEG detection of delayed auditory evoked responses in autism spectrum disorders:
Towards an imaging biomarker for autism. *Autism Research*, *3*(1), 8–18.
https://doi.org/10.1002/aur.111

Russo, N., Zecker, S., Trommer, B., Chen, J., & Kraus, N. (2009). Effects of background noise
on cortical encoding of speech in autism spectrum disorders. *Journal of Autism and*
Developmental Disorders, *39*(8), 1185–1196. https://doi.org/10.1007/s10803-009-0737-0

Rutter, M., Bailey, A., & Lord, C. (2003). *Scq. The Social Communication Questionnaire.* (Vol.
5). Western Psychological Services.

Sadaghiani, S., & Kleinschmidt, A. (2016). Brain Networks and α-Oscillations: Structural and
Functional Foundations of Cognitive Control. *Trends in Cognitive Sciences*, *20*(11), 805–
817. https://doi.org/10.1016/j.tics.2016.09.004

Schafer, E. C., Mathews, L., Gopal, K., Canale, E., Creech, A., Manning, J., & Kaiser, K. (2020). Behavioral Auditory Processing in Children and Young Adults with Autism Spectrum Disorder. *Journal of the American Academy of Audiology*, *31*(9), 680–689. https://doi.org/10.1055/s-0040-1717138

Smith, D. V, Davis, B., Niu, K., Healy, E. W., Bonilha, L., Fridriksson, J., Morgan, P. S., & Rorden, C. (2009). *Spatial Attention Evokes Similar Activation Patterns for Visual and Auditory Stimuli*. http://mitprc.silverchair.com/jocn/article-pdf/22/2/347/1769157/jocn.2009.21241.pdf

Soskey, L. N., Allen, P. D., & Bennetto, L. (2017). Auditory spatial attention to speech and complex non-speech sounds in children with autism spectrum disorder. *Autism Research*, *10*(8), 1405–1416. https://doi.org/10.1002/aur.1790

Stam, C. J., Nolte, G., & Daffertshofer, A. (2007). Phase lag index: Assessment of functional connectivity from multi channel EEG and MEG with diminished bias from common sources. *Human Brain Mapping*, *28*(11), 1178–1193. https://doi.org/10.1002/hbm.20346

Supekar, K., Uddin, L. Q., Khouzam, A., Phillips, J., Gaillard, W. D., Kenworthy, L. E., Yerys, B. E., Vaidya, C. J., & Menon, V. (2013). Brain Hyperconnectivity in Children with Autism and its Links to Social Deficits. *Cell Reports*, *5*(3), 738–747. https://doi.org/10.1016/j.celrep.2013.10.001

Taulu, S., & Simola, J. (2006). Spatiotemporal signal space separation method for rejecting nearby interference in MEG measurements. *Physics in Medicine and Biology*, *51*(7), 1759–1768. https://doi.org/10.1088/0031-9155/51/7/008

Teder-Sälejärvi, W. A., Pierce, K. L., Courchesne, E., & Hillyard, S. A. (2005). Auditory spatial localization and attention deficits in autistic adults. *Cognitive Brain Research*, *23*(2–3), 221–234. https://doi.org/10.1016/j.cogbrainres.2004.10.021

Tesche, C., Uusitalo, M., Ilmoniemi ayb, R., Huotilainen, M., & Kajola, M. (1995). Signal-space projections of MEG data characterize both distributed and well-localized neuronal sources. In *Electroencephalography and clinical Neurophysiology* (Vol. 95).

Uusitalo, M. A., & Ilmoniemi, ~ R J. (1997). Communication Signal-space projection method for separating MEG or EEG into components. In *Biol. Eng. & Cornput* (Vol. 35).

van der Heijden, K., Rauschecker, J. P., de Gelder, B., & Formisano, E. (2019). Cortical mechanisms of spatial hearing. In *Nature Reviews Neuroscience* (Vol. 20, Issue 10, pp. 609–623). Nature Publishing Group. https://doi.org/10.1038/s41583-019-0206-5

Wang, J., Wang, X., Wang, X., Zhang, H., Zhou, Y., Chen, L., Li, Y., & Wu, L. (2020). Increased EEG coherence in long-distance and short-distance connectivity in children with autism spectrum disorders. *Brain and Behavior*, *10*(10). https://doi.org/10.1002/brb3.1796

710   Zatorre, R. J., Mondor, T. A., & Evans, A. C. (1999). *Auditory Attention to Space and Frequency*
711   *Activates Similar Cerebral Systems*. http://www.idealibrary.com
712