Iron retention coupled with trade-offs in localized symbiotic effects confers tolerance to combined iron deficiency and drought in soybean

Md Rokibul Hasan, Asha Thapa, Ahmad H. Kabir[*]

[1]School of Sciences, University of Louisiana at Monroe, Monroe, LA 71209, USA

[*]Corresponding author: kabir@ulm.edu

ABSTRACT

Iron (Fe) and water availability are closely interlinked, with deficiencies in both adversely affecting soybean growth. However, the strategies employed by soybean to tolerate such conditions remain poorly understood. This study elucidates the interactions of physiological adjustments, host factors, and microbial associations in soybean using multi-omics approaches in Clark (tolerant) and Arisoy (sensitive) genotypes exposed to Fe deficiency and drought. Clark demonstrated stress resilience by maintaining osmotic balance, nutritional status, and photosynthetic efficiency, whereas these were severely affected in Arisoy. Furthermore, Fe retention in Clark, accompanied by the upregulation of ferritin-like proteins, may mitigate oxidative stress by reducing Fenton reactions. Interestingly, the elevated jasmonic acid and salicylic acid in the roots of Clark, compared to the lower levels in Arisoy, may contribute to differential physiological adjustments for stress adaptations. RNA-seq analysis revealed 818 and 500 upregulated, along with 931 and 361 downregulated genes, in roots of Clark and Arisoy, respectively, under stress. We observed the upregulation of symbiotic genes, such as *Chalcone-flavonone isomerase* 1 and *SWEET10*, accompanied by increased rhizosphere siderophore and root flavonoid in Clark. This indicates a significant microbial role in driving differential stress tolerance. Amplicon sequencing revealed distinct root and nodule microbiome dynamics, with Clark recruiting beneficial microbes such as *Variovorax* and *Paecilomyces*, known for promoting plant growth, whereas Arisoy displayed an opposite pattern. In addition, Clark maintained *Bradyrhizobium* and tissue nitrogen status, supported by ammonium retention and induction of *Ammonium transporter* 1 in the roots. Furthermore, *in vitro* compatibility between *V. paradoxus* and P. *lilacinus* suggests a synergistic interaction, with their localized signals benefiting Clark. Remarkably, enriched microbiomes significantly improved growth parameters, accompanied by elevated rhizosphere siderophore in sensitive genotypes under stress. This study is the first to uncover mechanisms of dual stress tolerance in soybean that may advance breeding and biofertilizer strategies for climate resilience.

Keywords: Abiotic stresses, enriched microbiome, iron homeostasis, legume, siderophore, symbiosis.

 Running head: Soybean tolerance to combined Fe deficiency and drought
48

1. Introduction

Climate change poses an increasing threat to global agricultural sustainability by intensifying abiotic stresses, including drought and nutrient deficiencies. Drought significantly reduces crop growth and productivity by impairing critical physiological processes (Qiao et al., 2024). This stress arises from insufficient soil moisture, which not only restricts water availability but also limits the uptake of essential minerals by plant roots (Li et al., 2023). Iron (Fe) deficiency, a prevalent nutritional disorder in plants, results from low bioavailability in alkaline or calcareous soils, where high pH restricts Fe solubility and uptake (Li and Lan, 2017; Kabir et al., 2012). Fe deficiency not only affects plant health but also disrupts photosynthetic efficiency, gene expression, and metabolic pathways (Liang, 2022), leading to symptoms like interveinal chlorosis in young leaves (Li et al., 2021). Drought further exacerbates this condition by lowering soil moisture and Fe solubility, which is particularly critical in legumes, where Fe plays a key role in nitrogen fixation (Lindström and Mousavi, 2020; Briat et al., 2015). Drought disrupts nutrient uptake and increases reactive oxygen species (ROS), thus hindering plant growth and survival (Hussain et al., 2018). Therefore, understanding the interplay between Fe deficiency and drought responses in legume crops is crucial for devising strategies to improve resilience under complex stress conditions.

Plants utilize two main strategies to absorb Fe from the soil: Strategy I (non-graminaceous species) and Strategy II (graminaceous species) (Liang, 2022; Gill and Tuteja, 2010). Plants using Strategy I possess several mechanisms to enhance Fe uptake under Fe deficiency. One such adaptation involves increasing proton extrusion through *ATPase 1*, which acidifies the rhizosphere and improves Fe solubility (Kabir et al., 2012). Additionally, ferric-chelate reductase (FCR) enzymes convert Fe(III) into the more bioavailable Fe(II) form (Kobayashi and Nishizawa, 2012; Morrissey and Guerinot, 2009). Plants further upregulate Fe transporter proteins to facilitate the uptake of bioavailable Fe into root cells (Kroh and Pilon, 2019). Plants adapt to drought by maintaining cellular hydration and minimizing damage through strategies such as modifying root architecture to enhance water uptake and closing stomata to reduce transpiration (Kumar et al., 2021; D'Oria et al., 2022). Furthermore, the ammonium transporter located on the plasma membrane actively transports ammonium ions from nitrogen-fixing bacteria in nodules of legume crops into the plant cell (Hao et al., 2020). At the molecular level, plants undergo dynamic changes involving the activation of stress-responsive genes and signaling pathways. Several genes play a direct role in stress mitigation, including those encoding dehydrins (Hassan et al., 2015) and enzymes involved in osmolyte biosynthesis (Ghosh et al., 2021), which contribute to water retention. Furthermore, abscisic acid (ABA) signaling plays a central role by regulating stomatal closure, conserving water, and activating a cascade of stress-responsive genes under drought conditions (Chaves et al., 2003; Ali et al., 2020). Jasmonic acid (JA) protects plants by regulating stomatal closure by boosting antioxidant defenses and promoting osmotic adjustment to reduce cellular damage (Wasternack and Song, 2017; Dhakarey et al., 2016). In drought-stressed plants, JA coordinates with other hormones like ABA and salicylic acid (SA) to regulate water loss and activate stress-responsive genes (Rai et al., 2024; Yang et al., 2019). However, the physiological strategies and gene activation mechanisms triggered in plants exposed to combined Fe deficiency and drought are not yet well understood.

While plants possess intrinsic mechanisms to mitigate abiotic stress, these are often insufficient without the support of beneficial microbes. Sugars and secondary metabolites such as flavonoids play crucial roles as host determinants influencing microbial recruitment, particularly under stress conditions (Loo et al., 2024; Lidoy et al., 2023). These compounds act as chemical signals in the rhizosphere, shaping microbial communities and promoting beneficial interactions that enhance plant resilience. The shifts in the belowground microbiome underlying abiotic stresses in plants involve changes in both the relative abundance and composition of microbial groups (Trivedi et al., 2022). Under Fe-limiting conditions, rhizosphere microbes, including bacteria (*Pseudomonas, Variovorax*) and fungi (*Trichoderma*), secrete siderophores that solubilize Fe(III) for plant uptake (Kabir and Bennetzen, 2024; Wang et al., 2024; Crowley et al., 2006). A recent study revealed the enrichment of *Variovorax*, Shinella, and *Chaetomium* in the roots of pea plants under alkaline stress (Thapa et al., 2024a). Drought significantly influences not only microbial diversity but also the composition of microbial communities in the rhizosphere and plant parts (Salamon et al., 2020; Naylor and Coleman-Derr 2018). However, the dynamics of microbial communities and their synergistic relationships in response to combined stresses in legumes require further research.

Soybean (*Glycine max*), one of the most important legume crops globally, plays a vital role in food, feed, and biofuel production (Longley et al., 2020). Its productivity is highly susceptible to Fe deficiency (Zocchi et al., 2017) and drought (Du et al., 2024), with their interactive effects more severely impairing key physiological processes and further reducing yield. Despite advances in understanding individual stress responses, the interplay of molecular pathways and microbial dynamics under combined Fe deficiency and drought remains unexplored in soybean. This study addresses this gap by integrating transcriptomic, physiological, and microbiome analyses to uncover synergistic mechanisms underlying combined stress resilience in soybean. Furthermore, we investigated the interactions and persistence of enriched microbial consortia and their synergistic interactions that induce resilience in sensitive genotypes of soybeans to Fe deficiency and drought. This novel study not only sheds light on the candidate genes and dynamics of soybean-microbe interactions under dual stresses but also lays the foundation for innovative strategies to enhance soybean resilience in the face of combined Fe deficiency and drought stress in the soil.

2. Materials and methods
Plant cultivation technique:
The seeds of contrasting genotypes of soybean (tolerant: Clark, sensitive: Arisoy, AG58XF3) to the combined Fe deficiency and drought were collected from USDA-GRIN and LSU AgCenter. Seeds were surface sterilized by immersing them in a 4% sodium hypochlorite solution for 5 min, followed by three rinses with sterile water, and then germinated in a tray. After 3 d, uniform, and healthy seedlings were transplanted into 500 g soil pots. The pots contained a 1:2 mixture of natural soil (soybean field) and commercial potting mix and were used for two treatments: control (soil without lime, pH ~6.5) and Fe- drought+ (soil amended with 4.5g $NaHCO_3$ and 3.0g $CaCO_3$, pH ~7.8). The physicochemical characteristics of the soil are detailed in Supplementary Table S1. During the cultivation period, $NaHCO_3$ was added weekly to maintain the soil pH at 7.8 making Fe unavailable to plants (Kabir et al., 2012). In addition, drought conditions were induced two weeks after transplanting. Soil moisture content was monitored daily to maintain water levels at 75% for control plants and 25% for drought-stressed plants. To study the impact

of microbial absence on plant stress responses, autoclaved vermiculite was used in pots for
cultivation in a separate experiment. The induction of Fe deficiency and drought stress was
performed as described previously; however, the amounts of $NaHCO_3$ and $CaCO_3$ were reduced
to 3.0g and 2.0g, respectively, to maintain the pH at ~7.8. For the microbial consortia study,
Variovorax paradoxus and *Paecilomyces lilacinus* were inoculated into the soil at a concentration
of 1×10^9 CFU per gram. The plants were grown in a randomized complete block design with 9
replications (3 plants per pot) per treatment in the greenhouse at 25°C, with a photoperiod of 10
h light/14 h darkness and a light intensity of 250 μmol m^{-2} s^{-1}. The plants were cultivated for 4
weeks before data collection.

Determination of morphological and photosynthetic parameters:
A digital caliper was used to measure the height and length of the shoot and root, respectively.
The dry weights of the roots and shoots were then measured after they had been dried for 3d at
75°C in an electric oven. Also, nodules were carefully collected from soybean roots by gently
washing the roots to remove soil, followed by manual detachment of the nodules using sterilized
forceps. A handheld chlorophyll meter (AMTAST, United States) and a FluorPen FP 110 (Photon
Systems Instruments, Czech Republic) were used to measure the chlorophyll score and Fv/Fm
(maximal photochemical efficiency of PSII) on the uppermost fully developed leaves at three
distinct sites. The leaves were exposed to darkness for one hour before data collection to prepare
them for OJIP analysis. Furthermore, the relative water content (RWC) of fully expanded
trifoliate leaves was measured from the tip of the main stem (Schonfeld et al., 1988). Briefly, the
fresh weight (FW) of the leaves was recorded immediately after excision. After soaking the
leaves in distilled water for a full day at room temperature, they were thoroughly blotted dry
using tissue paper, and the turgid weight (TW) was determined. The leaf samples were oven-
dried for 72 h at 75°C to estimate their dry weight (DW). RWC was calculated using the
formula: RWC (%) = (FW – DW) / (TW – DW) x 100. Furthermore, young leaves and roots
taken from plants were subjected to nutrient analysis. After being separated, the roots were rinsed
with deionized water, immersed in a 0.1 mM $CaSO_4$ solution for ten min, and then cleaned under
running tap water. In addition, deionized water was used to wash the young leaves individually.
After being wrapped in envelopes, each sample was dried for 3d at 75°C in an electric oven.
Elemental concentrations were determined using inductively coupled plasma mass spectrometry
(ICP-MS) at the Agricultural and Environmental Services Laboratories, University of Georgia,
following standard protocols.

Analysis of ROS and Fenton reaction rate:
For H_2O_2 determination, root samples were processed as previously described (Alexieva et al.,
(2001). Briefly, the root samples were mixed with a 0.1% trichloroacetic acid solution and finely
ground into a powder using a mortar and pestle. The homogenized extract was centrifuged at
10,000 rpm for 15 min to remove cellular residues. The resulting supernatant was supplemented
with 1 M potassium iodide and 10 mM phosphate buffer (pH 7.0), and the mixture was incubated
in darkness for 60 min. The optical density of the solution was subsequently measured at 390 nm
using a spectrophotometer (ThermoFisher, Waltham, USA). The concentration of H_2O_2 was
quantified using a standard curve. Furthermore, hydroxyl radicals (•OH) in the roots were
determined a previously described (Coudray and Favier 2000). Briefly, fresh root tissues were
harvested, thoroughly washed with deionized water, and homogenized in ice-cold sodium
phosphate buffer (50 mM, pH 7.4) containing 1 mM EDTA. The homogenate was centrifuged at

$12,000 \times g$ for 10 min at 4°C, and the supernatant was collected. The reaction was initiated by
incubating the supernatant with 5 mM salicylic acid in sodium phosphate buffer at 37°C for 30
min in the dark, allowing for the formation of 2,3-DHBA (2,3-dihydroxybenzoic acid). The
reaction was stopped by adding cold ethanol (95% v/v), followed by centrifugation to remove
debris. The concentration of 2,3-DHBA in the supernatant was quantified spectrophotometrically
by measuring absorbance at 510 nm. A standard curve prepared with known concentrations of
2,3-DHBA was used for quantification. Furthermore, the Fenton reaction rate for root samples
was determined by the concentration of Fe and as H_2O_2 using the following formula: Fenton rate
(µmol/s) $= k \times \{Fe\ (µM)\} \times \{H_2O_2\ (µM)$, where $k = 0.01\ µM^{-1}S^{-1}$ at pH 7 and 25°C.

Ferric chelate reductase activity in roots:
Ferric   chelate   reductase   activity   in   roots   was   determined   using   sodium
bathophenanthrolinedisulfonic acid (Na□-BPDS) as the Fe^{2+}-specific chelator (Waters et al.,
2002). Root tissues from the two genotypes were first rinsed thoroughly with deionized water,
and root tips were cut and kept on ice. Approximately 100 mg of root tissue was washed in 0.2
mM $CaSO_4$ solution for 5 min and subsequently transferred into 2 mL microcentrifuge tubes
containing an assay solution consisting of 10 mM $CaSO_4$, 5 mM MES buffer (pH 5.5), 0.1 mM
Fe-EDTA, and 0.3 mM Na□-BPDS. A separate tube containing the assay solution without root
tissue served as the control. Both the sample and control tubes were incubated in a shaking water
bath at 23°C for 1 hour at $3000 \times g$ under dark conditions. Afterward, 1 mL of the assay solution
was transferred to a cuvette, and the absorbance was recorded at 535 nm using a
spectrophotometer. The ferric reductase activity was calculated based on the molar extinction
coefficient of Fe(II)-BPDS ($22.14\ mM^{-1}\ cm^{-1}$).

Phytohormone analysis in roots
The amounts of phytohormones in the roots were measured with Liquid Chromatography-Mass
Spectrometry (LC-MS). To put it briefly, 200 mg of fresh root tissue was crushed in liquid
nitrogen and then extracted using an 80:20 (v/v) methanol: water solution that contained 0.1 g/L
butylated hydroxytoluene and 0.1% formic acid. The samples were processed in a TissueLyser II
(Qiagen, USA) at 30 Hz for 30 sec while being held in pre-frozen bead beater adapters. The
samples were then incubated on a shaker at 5,000 rpm and 4°C for 16 h following a
centrifugation for 20 sec at 4°C. The supernatant was then moved to autosampler vials for LC-
MS analysis after they had been vortexed and centrifuged for 10 min at 4°C at 12,000 rpm. Each
phytohormone standard (Sigma-Aldrich, USA) was made in a solution containing 0.1% formic
acid and 80% methanol.

RNA-seq analysis in roots:
RNA-seq analysis was performed on root samples to investigate gene modulation under
combined Fe deficiency and drought. The roots were carefully cleaned by washing twice with
sterile ice-cold phosphate-buffered saline, including 10 sec of vortexing, followed by two rinses
with ice-cold sterile water. The cleaned root tissues were ground into a fine powder using a pre-
chilled mortar and pestle with liquid nitrogen. Total RNA was extracted from the powdered roots
using the SV Total RNA Isolation System (Promega, USA). For RNA-seq library preparation, 1
µg of RNA with an RNA Integrity Number (RIN) above 8 was used. The libraries were prepared
using the KAPA HiFi HotStart Library Amplification Kit (Kapa Biosystems, USA) and
sequenced on an Illumina NovaSeq 6000 platform at the RTSF Genomics Core Facility,

Michigan State University. Of the total reads generated, 92.5% passed quality control filters, with a quality score of ≥ Q30.

Bioinformatics analysis of raw FastQ files was conducted using Partek Flow genomic analysis software (Partek, St. Louis, MO, USA). Adaptor sequences and low-quality reads were removed using Trimmomatic (Bolger et al., 2014) to obtain high-quality clean reads. The clean reads were aligned to the soybean reference genome using HISAT2 (v2.0.5) (Kim et al., 2015). Gene read counts were quantified with HTSeq (Anders et al., 2015). Differentially expressed genes (DEGs) were identified using the limma v3.19 R package (Ritchie et al., 2015), with FPKM (fragments per kilobase of exon per million mapped fragments) used for expression quantification. DEGs were defined as those with an adjusted P-value < 0.05 and a fold change ≥ 2, determined using the false discovery rate (FDR) method. MA plots for DEGs were created using DESeq2, and heatmaps were generated using pheatmap in the R platform. Gene enrichment analysis was performed using the ShinyGO 0.80, which provides functional annotation and pathway analysis specific to soybean genomic datasets (Ge et al., 2020).

Determination of siderophore in the rhizosphere:
Siderophore levels in rhizosphere soil were measured using the chrome azurol S (CAS) assay (Himpsl and Mobley 2019). Briefly, rhizosphere soil was collected by removing tightly adhering soil from the root surface. The soil was homogenized in 80% methanol and centrifuged at 10,000 rpm for 15 min. A 500 µl aliquot of the supernatant was mixed with an equal volume (500 µl) of CAS solution and incubated at room temperature for 5 min. Absorbance was measured at 630 nm, with a blank containing 1 ml of CAS reagent serving as the reference. The siderophore content per gram of soil was determined using the formula: % siderophore unit = [(A_ref - A_sample) / A_ref] × 100, where A_ref is the absorbance of the CAS reagent alone, and A_sample is the absorbance of the CAS reagent mixed with the soil sample.

Determination of flavonoid in the roots:
The total flavonoid content in roots was assessed using spectrophotometric method (Piyanete et al. 2009). Briefly, 0.5 mL of a 70% ethanol extract derived from the roots was combined with 2 mL of distilled water and 0.15 mL of a 5% $NaNO_2$ solution. This mixture was incubated for 6 min, after which 0.15 mL of a 10% $AlCl_3$ solution was introduced, and then incubated for an additional 6 min. Subsequently, 2 mL of a 4% NaOH solution was added. The final volume was adjusted to 5 mL with methanol, and the mixture was mixed thoroughly. Following a 15 min incubation, the absorbance was recorded at 510 nm using a spectrophotometer. The total flavonoid concentration was calculated in milligrams per gram of dry weight, utilizing a quercetin standard curve for the determination.

Amplicon sequencing of 16S and ITS microbial community analysis in the roots:
Microbial populations in root samples were examined using Illumina amplicon sequencing, which targets the ITS region for fungus and the 16S rRNA gene for bacteria, where appropriate. The samples were vortexed in sterile phosphate-buffered saline (PBS) for 10 sec, followed by two rinses with sterile water. DNA was extracted from about 0.2 g of cleaned root tissue using the Wizard® Genomic DNA Purification Kit (Promega Corporation, USA). RNase and Proteinase K treatments were used during the extraction process to get rid of RNA and protein impurities, respectively. The primer pairs 341F (CCTACGGGNGGCWGCAG) and 805R

(GACTACHVGGGTATCTAATCC) and ITS3 (GATGAAGAACGYAGYRAA) and ITS4 (TCCTCCGCTTATTGATATGC) were used to create the amplicon libraries for the 16S rRNA and ITS genes, respectively. Sequencing was performed on the Illumina Novaseq 6000 platform (PE250). Raw sequencing reads were processed and trimmed using Cutadapt (Martin, 2011), followed by analysis with the DADA2 pipeline (Callahan et al., 2016). Reads associated with mitochondria or chloroplasts were excluded from the dataset. Taxonomic assignment of amplicon sequence variants (ASVs) was performed using the UNITE database (Nilsson et al., 2019) for ITS regions and SILVA database version 138 (Quast et al., 2013) for 16S rRNA sequences. Furthermore, microbial community composition was analyzed by calculating Bray-Curtis distance matrices from Hellinger-transformed data, followed by Principal Coordinate Analysis (PCoA) using the phyloseq and vegan packages in R. Amplicon sequence variants (ASVs) were classified at various taxonomic levels, and comprehensive analyses including alpha and beta diversity assessments, and relative abundance were performed using non-parametric Kruskal-Wallis tests determined at P < 0.05 (McMurdie and Holmes, 2013).

Nanopore 16S bacterial community analysis in the nodules:
The isolated nodules were thoroughly washed with running tap water, then surface sterilized in a 3% sodium hypochlorite solution for 3 min, and finally rinsed three times with sterilized distilled water. Total DNA extraction was carried out using the CTAB procedure, as previously described (Clarke, 2009). For PCR experiments, 5 ng of DNA from each sample was combined with 16S primers 27F (AGAGTTTGATCMTGGCTCAG) and 1492R (CGGTTACCTTGTTACGACTT) to amplify the nearly full-length bacterial 16S rRNA gene. The PCR was performed under the following thermal cycling conditions: initial denaturation at 95°C for 2 min, followed by 30 cycles of denaturation at 95°C for 30 sec, annealing at 60°C for 30 sec, and extension at 72°C for 2 min. This was followed by a final extension at 72°C for 10 min and a hold at 4°C. Amplicons from each sample were ligated to pooled barcoded reads with the 16S Barcoding Kit 24 V14 (SQK-16S114-24, Oxford Nanopore Technologies, Oxford, UK) for library creation. The sequencing was done on a MinION portable sequencer (Oxford Nanopore Technologies) with a flow cell (FLO-FLG114). The MinION™ sequencing data (FASTQ files) were processed using the EPI2ME software from Oxford Nanopore Technologies, resulting in pass reads with an average quality score >9. The EPI2ME Fastq Barcoding method (Oxford Nanopore Technologies) was used to reduce adapter and barcode sequences. Principal coordinate analysis, alpha diversity, and relative abundance of the bacterial community were calculated using the R programs phyloseq and vegan. Non-parametric Kruskal-Wallis tests were used to detect variations in taxonomic abundance at a 5% significance level (McMurdie and Holmes 2013).

Determination of total soluble sugar in roots:
The phenol-sulfuric acid method was employed as previously described with some modifications (Dey et al., 1990). Briefly, 100 mg of fresh root tissue was extracted in 10 mL of ethanol and incubated at 60°C for 1 hour. The extract was transferred to a 25 mL volumetric flask, and the residue was re-extracted with ethanol. The combined extracts were diluted to 25 mL with ethanol. A 1 mL aliquot of the extract was mixed with 1 mL of 5% phenol solution in a thick-walled test tube, followed by the addition of 5 mL of concentrated sulfuric acid. The reaction mixture was thoroughly mixed and allowed to cool to room temperature. The absorbance of the resulting solution was measured at 485 nm using a spectrophotometer. Total soluble sugar

concentration was quantified against a glucose standard curve and expressed as mg of glucose equivalents per gram of fresh weight (mg g^{-1} FW).

Determination of ammonium in roots:
Ammonia concentrations in root tissues were determined using Nessler's reagent spectrophotometric method (Jeong et al., 2013). Approximately 100 mg of fresh root tissue was homogenized in 5 mL of 2% (w/v) potassium chloride solution, and the supernatant was collected after centrifugation at 12,000 rpm for 10 min. One milliliter of the extract was mixed with 1 mL of Nessler's reagent and incubated at room temperature for 15 min. The absorbance of the reaction mixture was measured at 420 nm using a spectrophotometer. Ammonia content was quantified against a standard curve prepared with known concentrations of ammonium sulfate and expressed per gram of fresh weight (µg g^{-1}).

Molecular detection of *V. paradoxus* and *P. lilacinus* in roots:
The colonization efficiency of *V. paradoxus* and *P. lilacinus* in the roots was assessed using DNA-based qPCR. Briefly, root samples were washed twice with sterile phosphate-buffered saline, vortexed, and then rinsed twice with sterile water. DNA was extracted from approximately 0.2 g of root tissue using the cetyl trimethylammonium bromide (CTAB) method (Clarke, 2009). The extracted DNA was quantified using a NanoDrop ND-1000 Spectrophotometer (Wilmington, USA) and standardized by diluting all DNA samples to equal concentrations. The qPCR analysis was conducted using a CFX96 Touch Real-Time PCR Detection System (Bio-Rad, USA), with specific primers for *V. paradoxus* (forward: CAATCGTGGGGGATAACGC, reverse: GGCCGCTCCATTCGCGCA) and *P. lilacinus* (forward: CTC AGT TGC CTC GGC GGG AA, reverse: GTG CAA CTC AGA GAA GAA ATT CCG). The soybean *β-actin* (forward: GAGCTATGAATTGCCTGATGG, reverse: CGTITCATGAATTCCAGTAGC) that served as the internal control for relative quantification using 2-$^{\Delta\Delta}$CT method (Livak and Schmittgen 2001).

Microbial co-culture
We collected *P. lilacinus* (ATCC 10843) and *V. paradoxus* (B-1908) from the USDA Microbial Germplasm as representative strains of the genera *Paecilomyces* and *Variovorax*, respectively (Supplementary Fig. S2). We have tested the compatibility and interactions of *P. lilacinus* and *V. paradoxus* on nutrient agar in a series of *in vitro* time-course experiments. The microbial species (5 µl inoculum) were cultivated in three distinct combinations: *P. lilacinus* alone, *V. paradoxus* alone, and *P. lilacinus* and *V. paradoxus* for seven d. The growth of the microbial species was monitored for up to 7 d of culture using ImageJ software (National Institutes of Health, USA). The colony radius was measured at three different points and averaged to determine the growth of microbes.

Split-root assay in Clark
Split-root experiments were conducted on autoclaved soil consisting of a 1:2 mixture of natural field soil and commercial potting mix. The plants were grown for 2 weeks before performing the split-root assay in soil pots. This assay was conducted using a single pot with a partitioned split-root assembly (compartment), with some modifications (Thilakarathna and Cope 2021). The taproot of each seedling was severed at the midpoint, and the lateral roots were evenly distributed between two compartments within the same pot, which were filled with a mixture of

natural field soil and commercial potting mix in a 1:2 ratio. A solid plastic barrier was used to separate the compartments. Microbial inoculants (*V. paradoxus* and *P. lilacinus* at a concentration of $1\times10\square$ CFU per gram) were applied to one compartment of the split-root system, while the other compartment remained uninoculated, depending on the treatment (Fig. 7C). The treatments included three conditions: SR1: control/control (-/-), SR2: control/microbiome (-/+), and SR3: microbiome/microbiome (+/+). The split-root plants were then cultivated under these conditions for an additional four weeks before data collection.

Data analysis:
The statistical significance of each variable was evaluated using a combination of Student's t-test for pairwise comparisons and analysis of variance (ANOVA) for evaluating differences across multiple groups, where appropriate. Where significant differences were detected by ANOVA, Duncan's Multiple Range Test (DMRT) was employed as a post hoc analysis to identify specific group differences. All statistical analyses were conducted using SPSS Statistics 20.0 (IBM), ensuring a minimum of three biological replicates. Graphical representations were generated using the R package ggplot2.

3. Results
Changes in morphological and physiological attributes:
We observed distinct morphological and physiological responses in Clark and Arisoy under combined Fe deficiency and drought compared to untreated controls (Fig. 1A-1M). The combined stress caused a significant decrease in the number of nodules in both genotypes, with the decrease being more pronounced in Arisoy than in Clark (Fig. 1B). Both the root length and root dry weight were significantly declined due to combined Fe deficiency and drought in both genotypes relative to controls, with Clark showing a moderate reduction and Arisoy experiencing a more pronounced decline (Fig. 1C-1D). Shoot height and shoot dry weight showed no significant change in Clark but decreased significantly in Arisoy subjected to combined Fe deficiency and drought compared to controls (Fig. 1E-1F). Similarly, leaf relative water content (RWC) and SPAD score did not change significantly in Clark, whereas Arisoy showed a significant decrease in RWC and SPAD score under combined Fe deficiency and drought relative to controls (Fig. 1G-1H). Photosynthetic efficiency (Fv/Fm) was significantly impaired under Fe-drought+ conditions in both genotypes in contrast to untreated controls, with a less pronounced decline in Clark and a more severe reduction in Arisoy (Fig. 1I). In this study, H_2O_2 concentration showed no significant change in Clark but increased significantly in the roots of Arisoy under combined Fe deficiency and drought relative to controls (Fig. 1K). We found that Fe-drought+ did not significantly alter hydroxyl radical (•OH) levels in the roots. However, in Arisoy, Fe-drought stress led to a significant increase in •OH levels ($p < 0.01$) under concurrent Fe deficiency and drought relative to controls (Fig. 1K). The Fenton reaction rate showed contrasting responses in the two genotypes under Fe-drought+ conditions. In Clark, the Fenton reaction rate significantly decreased while Arisoy exhibited a significant increase in the Fenton reaction rate in the roots due to combined Fe deficiency and drought compared to controls (Fig. 1L). In Clark, ferric chelate reductase activity significantly increased, whereas, in Arisoy, it showed a marked decline in the roots under Fe-drought+ conditions compared to the controls (Fig. 1M).

We also cultivated a set of plants in sterile vermiculite conditions to study the effect of combined Fe deficiency and drought on growth parameters. Under Fe-Drought+ conditions, both Clark and

Arisoy cultivars showed significant reductions in SPAD, shoot height, shoot fresh weight, shoot dry weight, and leaf RWC compared to the control (Supplementary Table S2). In Clark, stem diameter, number of leaves, and root length showed no significant differences between treatments. In Arisoy, all parameters, including stem diameter, number of leaves, and root length, showed significant differences between the control and Fe-drought+ conditions (Supplementary Table S2).

Changes in nutritional status and symbiotic metabolites:
Significant changes in nutrient content were observed in both roots and leaves of Clark and Arosy exposed to combined Fe deficiency and drought (Table 1). In roots, Fe, S and N contents remained unchanged, while Mn content significantly decreased subjected to Fe deficiency and drought relative to untreated controls. In Clark leaves, no significant changes were observed for Fe, Zn, or N due to stress while S and Mn showed a significant decrease (Table 1). In contrast, Arisoy exhibited more pronounced changes in both roots and leaves. In the roots, Fe, Mn, and N contents significantly decreased, while Zn and S showed no significant changes in response to combined Fe deficiency and drought. Furthermore, Fe, Mn, S, and N contents showed a significant decline in the leaves of Arisoy due to stress, but Zn levels remained unchanged (Table 1). We also found that flavonoid content significantly increased, while soluble sugar and ammonium contents showed no significant changes between control and Fe-drought+ conditions in the roots of Clark. On the flip side, Arisoy exhibited a significant decrease in flavonoid, soluble sugar, and ammonium contents under Fe-Drought+ conditions compared to the control (Table 2).

Transcriptional changes in the roots:
Principal component analysis (PCA) of transcriptomic data revealed distinct clustering between control and Fe-drought stress treatments for both Clark and Arisoy genotypes, indicating significant transcriptional reprogramming (Fig. 2A and 2B). In Clark, the first two principal components explained 57.5% and 27.2% of the variance, respectively, while in Arisoy, they accounted for 53.8% and 24.6% of the variance. Differential gene expression analysis identified a greater number of differentially expressed genes (DEGs) in Clark compared to Arisoy. Specifically, Clark exhibited 818 upregulated and 931 downregulated genes under Fe-drought+ conditions (Fig. 2C), while Arisoy showed 500 upregulated and 361 downregulated genes in the roots (Fig. 2D). The heatmap illustrates the differential expression of stress-responsive genes in two soybean genotypes, Clark and Arisoy, under combined Fe deficiency and drought stress compared to control conditions (Fig. 2E-G; Supplementary Table S3). In RNA-seq analysis, Clark exhibited higher expression levels of key genes associated with stress tolerance, such as *Glyma.20G241500* (*Chalcone-flavone isomerase 1*), *Glyma.08G059700* (*Sugar transporter 1*), *Glyma.04G198400* (*Bidirectional sugar transporter SWEET10*), *Glyma.05G035500* (*EamA-like transporter*) and *Glyma.15G154100* (*2-oxoglutarate*) while Arisoy shows comparatively lower expression or downregulation of these genes in the roots in response to combined Fe deficiency and drought (Fig. 2E). The expression levels of genes related to nutrient uptake and transport including *Glyma.10G146600* (Fe-regulated protein 3) and *Glyma.10G167800* (*Ammonium transporter 1;2*), *Glyma.07G006500* (*Sulfate transporter 3;4*) and *Glyma.17G119400* (*Stabilizer of Fe transporter SufD*) were upregulated in Clark but showed no significant changes in Arisoy.

However, *Glyma.07G113500* (*ATPase 11*) was significantly upregulated in both Clark and Arisoy due to combined Fe deficiency and drought (Fig. 2F). Further, genes related to osmotic adjustment such as *Glyma.17G027400* (Drought-induced-19 protein) showed significantly higher expression in both cultivars, with slightly elevated levels in Arisoy due to combined Fe deficiency and drought. In contrast, *Glyma.11G106900* (Dehydration-associated protein) was significantly upregulated in Clark while Arisoy showed a non-significant higher expression in the root (Fig. 2G). In addition, the expression levels of genes associated with ROS detoxification such as *Glyma.19G091400* (Ferritin-like family protein), *Glyma.08G013800* (*Glutathione peroxidase 6*), *Glyma.07G032600* (*Glutaredoxin 4*) and *Glyma.13G172300* (*Peroxidase 19*) exhibited significantly higher expression in Clark while Arisoy showed no significant changes in the roots due to combined Fe deficiency and drought (Fig. 2H).

The functional categorization and gene enrichment analysis showed distinct biological processes and KEGG pathways in Clark and Arisoy soybean genotypes under combined Fe deficiency and drought stress (Fig. 3A-3B). In Clark, the most enriched biological processes include detection of visible light, protein-chromophore linkage, polysaccharide catabolic process, response to water, carbohydrate metabolic process, oxidation-reduction process, and response to oxidative stress (Fig. 3A). For KEGG pathways, Clark shows significant enrichment in linoleic acid metabolism, alpha-linolenic acid metabolism, glycosphingolipid biosynthesis, circadian rhythm, starch and sucrose metabolism, and biosynthesis of secondary metabolites. In Arisoy, enriched biological processes include triglyceride biosynthetic process, detection of visible light, protein-chromophore linkage, glycerol ether metabolic process, polysaccharide catabolic process, and cell redox homeostasis. Furthermore, the enriched KEGG pathways in Arisoy include steroid biosynthesis, circadian rhythm, starch and sucrose metabolism, and glycerolipid metabolism (Fig. 3B).

Changes in phytohormone levels in the roots:
The phytohormonal profiles in the roots of Clark and Arisoy showed differential patterns in response to dual Fe deficiency and drought. Indole-3-acetic acid (IAA) levels significantly decreased under Fe-drought+ stress in both Clark and Arisoy due to Fe deficiency and drought relative to untreated controls (Fig. 4A). In contrast, abscisic acid (ABA) levels were significantly elevated in both genotypes under Fe-drought+ stress compared to controls (Fig. 4B). Jasmonic acid (JA) levels showed a significant increase in the roots of Clark but showed no significant changes in the Arisoy in response to combined Fe deficiency and drought compared to controls (Fig. 4C). The salicylic acid (SA) levels responded differently to Fe-drought+ in the contrasting soybean genotypes. In Clark, SA levels significantly increased under combined Fe deficiency and drought compared to controls. Conversely, in Arisoy, SA levels significantly decreased in such circumstances (Fig. 4D).

Shift in microbial dynamics in contrasting genotypes:
We studied the effects of combined Fe deficiency and drought on the rhizosphere siderophore, microbial diversity, and community composition in the roots and nodules of Clark and Arisoy soybean genotypes. In Clark, Fe-drought+ stress significantly increased siderophore levels in the rhizosphere; however, no significant change was observed in Arisoy under the same conditions (Fig. 5A). PCA analysis showed clear clustering of root 16S microbial communities between control and Fe-drought+ treatments in both genotypes (Fig. 5B). Alpha diversity metrics showed

no significant changes in observed richness and Shannon diversity index in Clark under combined Fe deficiency and drought compared to controls. However, observed richness (p = 0.025) significantly decreased in Arisoy while Shannon diversity remained unaffected in Arisoy (Fig. 5C). The induction of Fe deficiency and drought in soil significantly altered the relative abundance of several genera in the root bacterial microbiomes. In Clark, the abundance of *Variovorax*, *Streptomyces*, and *Pararhizobium* increased significantly due to combined Fe deficiency and drought (Fig. 5D). In Arisoy, the abundance of *Streptomyces* and *Cellulomonas* significantly increased, while *Variovorax* and *Novosphingobium* significantly decreased in the roots in response to combined Fe deficiency and drought (Fig. 5E). The bacterial genera that showed no significant shift in relative abundance in the roots between control and Fe-drought+ conditions are as follows: Clark (*Shinella, Pseudomonas, Mycobacterium, Methylophilus, Herbaspirillum, Asticcacaulis*) and Arisoy (*Shinella, Pararhizobium, Mycobacterium, Methylophilus, Flavobacterium, Cellvibrio*). In this study, the nodule microbiomes were dominated by *Bradyrhizobium* in both genotypes under both conditions (Fig. 5F-5G). In Clark, *Bradyrhizobium* showed no significant changes, while in Arisoy, it significantly declined in relative abundance in the roots due to combined Fe deficiency and drought (Fig. 5F-5G). Furthermore, soybean root nodules from both the Clark and Arisoy cultivars exhibit distinct color differences under combined Fe deficiency and drought stress (Supplementary Fig. S1). In the control condition, the nodules displayed a vibrant pink color, whereas, under stress conditions, they showed a faded pink hue in both genotypes. However, the extent of color fading was less pronounced in Clark and more severe in Arisoy (Supplementary Fig. S1).

In this study, the fungal community composition in the roots showed distinct clustering between the control and Fe-drought+ treatments in both genotypes, indicating shifts in fungal community composition under stress conditions (Fig. 6A). Alpha diversity measures (Panel B) showed no significant change in observed richness for either genotype under Fe-drought+ conditions (Fig. 6B). However, the Shannon diversity index significantly decreased in the roots of Clark under Fe-drought+ conditions, while no significant change was observed for Arisoy (Fig. 6B). Relative abundance analysis highlighted specific genera that were significantly impacted by Fe-drought+ conditions. In Clark roots, *Pseudohyphophila* and *Paecilomyces* significantly increased in abundance, while other genera, including *Trichocladium, Pseudoallescheria, Paracremonium, Humicola, Cladosporium, Chaetomium* and *Arnium,* showed no significant changes (Fig. 6C). In Arisoy roots, *Cladosporium* significantly increased, and *Paecilomyces* significantly decreased in abundance (Fig. 6D). Genera such as *Scedosporium, Pseudohyphophila, Pseudoallescheria, Podospora, Fusarium, Chaetomium, Ascospheara,* and *Arnium* remained unaffected by the Fe-drought+ treatment in the roots of Arisoy (Fig. 6D).

We further illustrated ASV counts for major beneficial microbes (*Variovorax* and *Paecilomyces*) across genotypes (Clark and Arisoy) differentially shifted due to combined Fe deficiency and drought (Fig. 6E). For *Variovorax*, the ASV number increased substantially in Clark under Fe-drought+ conditions (18,108 ASVs) compared to the control (9,348 ASVs). In contrast, Arisoy showed a substantial decrease in *Variovorax* between Fe-drought+ (5,284 ASVs) and control (13,378 ASVs) conditions. For *Paecilomyces*, the ASV count increased dramatically in Clark under Fe-drought+ conditions (493,233 ASVs) compared to the controls (4,528 ASVs). However, in Arisoy, *Paecilomyces* decreased under Fe-drought+ conditions (1,771 ASVs) compared to the control (4,788 ASVs), although the increase was less pronounced than in Clark (Fig. 6E).

Interactions of *P. lilacinus* and *V. paradoxus* and changes in plant growth parameters in split-root plants:

We found that there was no significant difference in colony size in *P. lilacinus* and *V. paradoxus* between single and co-culture conditions at all time points (Day 3, Day 5, and Day 7). In contrast, for *V. paradoxus*, the colony size in co-culture was significantly larger compared to single culture on all time points, indicating enhanced growth in the presence of *P. lilacinus* (Fig. 7A-7B). Furthermore, the split-root experiment was designed to evaluate the localized and systemic effects of enriched microbiomes on plant growth and physiology in Clark. Visual observations revealed that plants in the SR3 treatment (microbiome/microbiome) exhibited the most robust growth, with larger shoots and healthier root systems compared to SR1 (control/control) and SR2 (control/microbiome) systems (Fig. 7C-7D). In this study, SR1 exhibited significantly lower SPAD values compared to SR3, while SR2 showed intermediate values that were not significantly different from either SR1 or SR3 (Fig. 7E). However, shoot height and shoot dry weight remained stable across all treatments (Fig. 7F-7G). The root length and root dry weight showed significant differences between the compartments in SR2, while these root features showed no significant differences between the compartments in SR1 and SR3 (Fig. 7H-7I). Furthermore, the rhizosphere siderophore production showed significant differences between the compartments in SR2, with Com 2 having higher siderophore levels than Com 1. However, no significant differences were observed between the compartments in SR1 and SR3 (Fig. 7J).

Effects of enriched microbiomes on sensitive soybean genotypes

The combined Fe deficiency and drought caused a significant decline in morphological and physiological traits of two sensitive genotypes (Arisoy and AG58XF3) of soybean (Fig. 8A-8F). We observed that inoculants consisting of an enriched microbiomes (*V. paradoxus* and *P. lilacinus*) significantly increased leaf SPAD, plant height, plant biomass, leaf RWC, and root nodule number under combined stress compared to Fe-Drought+ conditions in both genotypes. Plants without stress but inoculated with the enriched microbiomes exhibited similar trends in these parameters to those of the controls in both Arisoy and AG58XF3 (Fig. 8A-8F). In Arisoy, rhizosphere siderophore showed no significant changes due to combined stresses but AG58XF3 showed a significant increase compared to controls (Fig. 8G). Interestingly, the addition of enriched microbiomes to the soil, whether in the presence or absence of combined stress, resulted in a significant increase in rhizosphere siderophore levels compared to Fe-Drought+ conditions. Siderophore levels were higher in the absence of stress than in its presence in both genotypes (Fig. 8G). Furthermore, flavonoid levels in the roots showed a significant decrease in both genotypes when plants were subjected to combined Fe deficiency and drought compared to untreated controls (Fig. 8H). However, flavonoid content in the roots of Arisoy showed a significant increase due to microbiome inoculation under stress, whereas no such increase was observed in AG58XF3 when compared to plants grown solely under stress. Both Arisoy and AG58XF3 inoculated with enriched microbiomes under control conditions exhibited flavonoid levels similar to those of the untreated controls (Fig. 8H).

4. Discussion

Fe and water status are entwined in plants. Hence stress resilience in plants to combined Fe deficiency and drought stress involves a complex interplay of physiological, molecular, and

microbial associations (Santos et al., 2023; Kanwar et al., 2021). Our findings reveal the
importance of tissue-level Fe retention, hormonal signaling pathways, and the enrichment of
specific beneficial microbes in mitigating the adverse effects of dual stresses in soybean. This
study uniquely integrates transcriptomic, phytohormonal, and microbial analyses to offer a
comprehensive understanding of the mechanisms involved in improving soybean resilience
under combined iron deficiency and drought stress.

Regulation of Fe homeostasis and ROS scavenging underlying stress tolerance:
This study reveals significant genotype-specific variations in stress resilience between Clark and
Arisoy under combined Fe deficiency and drought. Clark exhibited moderate reductions in root
length and biomass while maintaining shoot biomass and leaf RWC under stress. Variations in
RWC are closely linked to the genotypic background, which affects the ability of plants to
tolerate drought stress (Zhang et al., 2023; Kabir et al., 2015). Therefore, the physiological
adjustments, such as sustained water retention supported by stable photosynthetic efficiency
might have been effective in mitigating the impact of dual stress in Clark. This aligns with the
previous findings linking chlorophyll stability and photosynthetic efficiency to withstand abiotic
stress (Therby-Vale et al., 2022; Wang et al., 2018). The physiological adjustments of Clark to
are supported by transcriptomic data, which reveal the upregulation of genes associated with
osmotic adjustment. For instance, the enhanced expression of drought-induced-19 protein
(*Glyma.17G027400*) and dehydration-associated protein (*Glyma.11G106900*) in Clark may play
a critical role in maintaining cellular integrity under drought. The overexpression of drought-
induced genes improved drought and salt tolerance in various plants (Wu et al., 2022; Wu et al.,
2020). These proteins also play a role in ABA-dependent signaling pathways and regulate
stomatal aperture, The Fe homeostasis in stress resilience has been well-documented, thereby
influencing plant water status during drought conditions (Wu et al., 2020). In another study,
DREB (dehydration-responsive element-binding protein) mutants showed increased sensitivity to
drought in rice (Wang et al., 2022).

A lack of water or minerals in soil often triggers the imbalance of nutrient status and
overproduction of ROS in plants (Barzana et al., 2021; Jiang and Zhang, 2002). In this study, the
upregulation of the genes related to dehydration protection in Clark is consistent with the
maintenance of mineral status, such as Fe and other nutrients, under combined stresses. The
improved Fe status in Clark was further supported by the induction of *Glyma.10G146600* (Fe-
regulated protein 3) in the roots. Fe regulatory proteins (IRPs) maintain Fe homeostasis in
eukaryotic cells by binding to Fe-responsive elements in target mRNAs and regulating their
stability (Zhang et al., 2014). Fe availability is crucial for enzymatic functions and
photosynthetic machinery (Connorton et al., 2017; Kobayashi and Nishizawa, 2012).
Furthermore, elevated ATPase and FCR activities under Fe deficiency are closely linked to
enhanced Fe acquisition, forming a key component of the Strategy I response in plants (Kabir et
al., 2012; Santi and Schmidt, 2009). In this study, the upregulation of *ATPase 11*, responsible for
proton extrusion (H^+), in both Clark and Arisoy suggests an adaptive response to promote
rhizosphere acidification for mobilizing soluble Fe and other minerals. Interestingly, we also
found a significant upregulation of *Sulfate transporter 3;4* and *Stabilizer of Fe transporter SufD*
in Clark when cultivated in dual Fe deficiency and drought. SufD is a key component of the
SUF (S utilization factor) machinery essential for Fe-S cluster assembly and Fe acquisition in
plants (Bai et al., 2018). S plays a key role in maintaining the homeostasis of micronutrients such

as Fe, Cu, Zn, and Mn primarily through S-derived organic ligands that facilitate metal uptake and translocation (Chorianopoulou and Bouranis 2022). Furthermore, Fe deficiency increases the demand for S uptake and assimilation (Astolfi et al., 2021). These findings align with improved S status in the root, suggesting that retaining Fe and other micronutrients helps soybean plants cope with combined Fe deficiency and drought.

The ability of Clark to limit stress-induced damage was supported by reduced Fenton reactions in the roots. This aligns with the upregulation of Ferritin-like family protein in the roots of Clark exposed to simultaneous Fe deficiency and drought. Fe and ferritin are vital for drought tolerance as they support antioxidant enzymes in neutralizing ROS during stress (Zhang et al., 2023; Ndou et al., 2023). Ferritin, a Fe-storage protein, also acts as a central player to prevent the excessive accumulation of free Fe that could lead to ROS production (Briat et al., 2010). This aligns with our findings, as the lower ROS such as H_2O_2 and •OH levels in Clark further support enhanced capacity in Clark to mitigate oxidative stress in concurrent stresses. In contrast, dual stress imposed a greater oxidative burden on Arisoy, possibly exacerbated by impaired water transport and stress signaling pathways. Additionally, differences in oxidative stress responses, particularly those associated with the Fenton reaction, may contribute to the contrasting resilience observed in Clark and Arisoy. The Fenton reaction is a chemical process in which Fe(II) reacts with H_2O_2 to generate highly reactive •OH in plants (Smirnoff and Arnaud, 2019; Richards et al., 2015). This was likely due to its enhanced FCR activity and effective Fe utilization, which limited the availability of free Fe for ROS generation in Clar under stress. Reduced Fenton activity in Clark not only minimized oxidative damage but also allowed for better allocation of Fe to essential metabolic processes. These findings align with the role of Fe homeostasis and ROS scavenging in plant stress tolerance (Taheri et al., 2022; Sun et al., 2017). The reduction in H_2O_2 and •OH levels was correlated with the upregulation of several ROS-related genes such as *Peroxidase 19*, *Glutathione peroxidase* and *Glutaredoxin 4* in Clark. Peroxidase genes mainly scavenge H_2O_2 resulting in the inhibition of •OH generations through the Fenton reaction in plants under abiotic stresses (Zhang et al., 2022; Kidwai et al., 2020). Transgenic plants expressing glutathione peroxidase genes (GPX) exhibited increased tolerance to drought, and oxidative damage in several plant species (Hou et al., 2024; Zhang et al., 2019). Furthermore, glutaredoxin protein plays a vital role in the assembly and maintenance of Fe-S clusters, which are essential for ROS detoxification and various metabolic processes (Couturier et al., 2015). *Glutaredoxin* knockouts in plants have been shown to impair stress tolerance and increase oxidative damage (Laporte et al., 2012). As a result, the better adaptability of Clark to Fe deficiency and drought is at least partially attributed to the expression of antioxidant and ROS-scavenging mechanisms, which help limit stress-induced cellular damage.

Changes in hormonal balance and symbiotic determinants in the roots
Significant hormonal changes were observed in Clark and Arisoy under the conditions of combined Fe deficiency and drought. Both genotypes showed reduced IAA, probably reallocating resources from growth to stress adaptation. A decline in auxin levels under drought and nutrient deficiency showed the conservation of energy and resources for survival rather than growth (Kurepa and Smalle, 2022; Peleg and Blumwald, 2011). Interestingly, IAA reduction is often linked to a shift in the balance of phytohormones, favoring ABA and ethylene pathways in plants exposed to drought (Mubarik et al., 2021; Gupta et al., 2017). In this study, the induction of ABA in both genotypes suggests a general adaptive response of plants to regulate stomatal

closure and limit water loss under drought (Munemasa et al., 2015). Interestingly, JA levels were significantly elevated in Clark under Fe deficiency and drought, while Arisoy exhibited no changes. The interplay between ABA and JA is not conclusive and depends on the type of stress and plant species. However, JA accumulation is essential for ABA accumulation following drought treatment in *Arabidopsis* (de Ollas et al., 2015). Similarly, increased JA levels promote ABA production in rice under drought stress (Kim et al., 2009; Li et al., 2019) suggesting that JA acts upstream of ABA in the regulatory pathway. Hence, the elevation of JA levels suggests that it may be one of the drivers of stress responses in Clark under combined Fe deficiency and drought. Additionally, Clark showed an increase in SA levels, which is known to trigger antioxidant genes and reduce ROS accumulation during drought stress in plants (Jahan et al., 2023). These findings emphasize the role of hormonal interactions, particularly JA and SA induction in Clark, in enhancing tolerance to combined Fe deficiency and drought.

In our RNA-seq analysis, genes associated with symbiotic interactions, including *Chalcone-flavonone isomerase*, *2-Oxoglutarate*, *Sugar transporter protein 1*, and *Bidirectional sugar transporter SWEET10*, showed higher expression levels in the Clark genotype compared to Arisoy. Flavonoids, including chalcones and flavones, are key host determinants that regulate microbial recruitment and gene expression, thereby promoting symbiotic interactions (Tohge et al., 2017; Kang et al., 2014; Hassan and Mathesius, 2012). Additionally, sucrose and hexoses released into the rhizosphere serve as energy sources, signaling molecules to attract beneficial microbes (Kryukov et al., 2021). Particularly, Sugar transporter protein 1 and *Sugar efflux transporter SWEET10* direct carbon flow to the rhizosphere (Chen et al., 2012) and support the energy demands of nitrogen-fixing symbionts (Breia et al., 2021; Kryukov et al., 2021). Also, the upregulation of the *EamA-like transporter gene* suggests the stability of Clark in nitrogen assimilation, as this transporter is known to contribute to nutrient exchange and signaling during symbiotic interactions in legumes (Garcia et al., 2023; Banasiak et al., 2021; Guefrachi et al., 2015). Taken together, our findings suggest the strong connection of symbiotic benefits as a key factor in improving resilience to combined Fe deficiency and drought stress in soybean. Consistently, our results showed that stress resilience in Clark is completely disrupted in sterile conditions, as indicated by reduced growth and photosynthetic attributes under Fe-drought+ conditions.

Shift in microbial recruitment underlying stress resilience:
To further elucidate microbial associations with differential stress resilience, we performed 16S and ITS sequencing on roots and nodules. The microbial community dynamics in the nodule and roots also demonstrated genotype-specific dynamics in the microbiome. Microbial diversity acts as a crucial mediator between plants and their surrounding environment (Zhang et al., 2021). In this study, the significant increase in rhizosphere siderophore in Clark under Fe deficiency and drought suggests enhanced microbial activity aimed at improving Fe solubility and availability. Siderophore production is a known adaptive response of rhizosphere microbes to Fe-limiting environments (Kabir and Bennetzen 2024). Hence, the lack of a similar response in Arisoy may indicate either limited microbial activity or insufficient recruitment of siderophore-producing microbes, which could partially explain its reduced tolerance to combined stress. This is supported by the stability of microbial abundance and diversity in Clark under stress. However, Arisoy exhibited a significant decline in bacterial abundance, indicating reduced microbial adaptability. This discrepancy in the bacterial community might contribute to the observed

differences in stress tolerance between the genotypes. Furthermore, microbial diversity is linked
to plant fitness, nutrient cycling, and stress mitigation (Naylor et al., 2017).

Our study demonstrates how contrasting genotypes recruit and sustain beneficial microbes in response to Fe deficiency and drought. Several bacterial genera such as *Variovorax*, *Streptomyces*, *Pararhizobium*, and *Novosphingobium* were altered in the roots of Clark. The increase in *Variovorax* in Clark is particularly promising, given its well-documented role in nutrient solubilization and phytohormone modulation (Jiang et al., 2012; Sun et al., 2018). Also, the differential shift in *Streptomyces* between contrasting genotypes suggests a potential genotype-specific recruitment or interaction mechanism to stress. Previous studies reveal that drought-induced Fe limitation in sorghum roots disrupts Fe homeostasis, enriching *Streptomyces* in the rhizosphere (Xu et al., 2021). They also found that the addition of exogenous Fe disrupts this enrichment and diminishes the plant growth-promoting effects of *Streptomyces*. Additionally, drought-responsive plant metabolism improves Monoderm bacterial lineages, including *Actinobacteria*, *Chloroflexi*, and *Firmicutes* (Xu and Coleman-Derr, 2019). The disappearance of *Streptomyces* in the roots under combined Fe deficiency and drought may be associated with the improved Fe status in Clark, aligning with findings in drought-exposed sorghum (Xu et al., 2021). Furthermore, Clark may possess a genotype-specific ability to maintain Fe homeostasis, reducing the need for *Streptomyces* enrichment while the stress conditions could exacerbate Fe scarcity, intensifying the recruitment of *Streptomyces* in Arisoy.

As expected, *Bradyrhizobium* dominated the nodule microbiomes of both soybean genotypes. However, its significant decline in Arisoy under stress indicates disrupted symbiotic efficiency. This could result in reduced N availability, reflected in the N status of roots and shoots, whereas Clark exhibited no such changes. Particularly, drought directly affects key processes involved in nodulation by reducing soil moisture, which limits the mobility of nitrogen-fixing bacteria and impairs their ability to colonize legume roots (Lumactud et al., 2023; Larrainzar et al., 2007). Although Clark exhibited a reduction in nodule number, the abundance of *Bradyrhizobium* within the nodules remained stable and active, indicated by the pink-red color of the cross-sections under combined Fe deficiency and drought. The pink-red interior color of root nodules due to the presence of leghemoglobin may be associated with high rates of nitrogen fixation (Jiang et al., 2021; Küçük & Cevheri, 2014). Moreover, the efficiency of Clark to maintain *Bradyrhizobium* supports its symbiotic stability, aligning with ammonium retention coupled with the upregulation of *Ammonium transporter 1* and N status. Similar studies have shown that Rhizobia-legume symbiosis is crucial for mitigating drought stress and enhancing nitrogen fixation in *Vicia sativa* and *Pisum sativum* (Álvarez-Aragón et al., 2023). Hence, the stress resilience in Clark probably stems from active *Bradyrhizobium*, enabling ammonium production, which is subsequently imported into the roots via *Ammonium transporter 1*. In ITS analysis, genotypic differences in fungal communities in the root were evident with *Paecilomyces* significantly enriching in Clark but not in Arisoy. *P. variotii* functions as an elicitor, stimulating the growth of *Arabidopsis thaliana* by inducing auxin accumulation and improving disease resistance through the activation of the SA signaling pathway (Lu et al., 2019). Similarly, tomato plants inoculated with *P. lilacinus* have been shown to exhibit enhanced growth parameters and increased flavonoid content (Musa et al., 2023). Taken together, the differential microbial dynamics observed in Clark and Arisoy underline the critical role of the root-associated microbiomes in differential stress tolerance to Fe deficiency and drought. The ability of Clark to

781 recruit *Variovorax* and *Paecilomyces* may pose the potential of leveraging enriched microbiomes
782 as a key trait for selection to improve tolerance in soybean in response to combined Fe
783 deficiency and drought.
784
Interactions of enriched microbiomes, signaling pathways, and effects on inducing stress
resilience:
Microbial compatibility is essential for the success of microbial consortia in promoting plant
growth (Prigigallo et al., 2023). Microbial co-cultures can significantly alter the growth and
metabolism often mediated by diverse metabolites produced by participating organisms
(Bertrand et al., 2014; Wolter and Hoffman, 2014). In this study, the increased colony size of *V.*
paradoxus in the presence of *P. lilacinus,* suggests a beneficial effect of *P. lilacinus* on *V.*
paradoxus. Such interactions are commonly observed in microbial communities, where one
species facilitates the growth of another through metabolic complementation (Liu et al., 2024;
Mendes et al., 2013). Furthermore, the lack of significant variation in the colony size of *P.*
lilacinus across co-culture conditions suggests that *P. lilacinus* is either unaffected by or tolerant
of the presence of *V. paradoxus*, maintaining its growth independently of any potential microbial
interactions. The interactions between *V. paradoxus* and *P. lilacinus* with other microbes need
further investigation. However, positive growth-promoting interactions are common among
microbes, particularly between strains with different carbon consumption profiles (Kehe et al.,
2020). Studies have shown that fungal-bacterial interactions can promote plant growth and
inhibit plant pathogens (Santoyo et al., 2021; Artursson et al., 2005). Therefore, the compatible
interactions between *P. lilacinus* and *V. paradoxus* suggest their synergistic coexistence that may
offer potential benefits and stress resilience in soybean plants
804
We further performed the split-root assay using sterile soil that provided valuable insights into
the localized effects of enriched microbiomes on soybean growth. Plants (Clark) subjected to the
SR3 treatment (microbiome/microbiome) exhibited slightly healthier shoots (non-significant)
and healthier root systems along with the increased leaf chlorophyll content and rhizosphere
siderophore compared to SR1 (control/control) and SR2 (control/microbiome). These findings
emphasize the local effects of enriched microbiomes in promoting soybean growth. The
involvement of local signal-promoting growth and disease resistance was also reported in
sorghum (Kabir et al., 2024) and tomato (Lucas et al., 2018). However, studies also reported that
systemic colonization by beneficial microbes may optimize physiological traits, including
photosynthetic efficiency and biomass accumulation (Chi et al., 2005; Harman et al., 2021). In
this study, the effects of *P. lilacinus* and *V. paradoxus* inoculum are context-dependent in
soybean, with localized signals dominating in compartmentalized systems, while systemic
colonization optimizes physiological traits under more uniform conditions.
818
Building on the compatibility of *P. lilacinus* and *V. paradoxus*, we validated their persistence and
impact on sensitive genotypes to combined Fe deficiency and drought stress. Microbial
inoculants have gained significant attention for their potential to enhance plant tolerance to
abiotic stresses (Gonçalves et al., 2023; Khan et al., 2023). Our study found that microbiomes
enriched in Clark (tolerant) improved plant growth and stress resilience in genotypes sensitive to
Fe deficiency and drought. Several studies have demonstrated the beneficial effects of diverse
microbiomes in mitigating single abiotic stresses in plants, including but not limited to
Burkholderia phytofirmans for drought in wheat (Naveed et al., 2014), *Azospirillum brasilense*

for drought in Arabidopsis (Cohen et al., (2015), and *Trichoderma harzianum* for Fe deficiency in pea (Thapa et al., 2024b). One of the key observations in our study was the effectiveness of the enriched microbiome in Arisoy, even when cultivated under combined Fe deficiency and drought, in nexus with elevated rhizosphere siderophore and root flavonoid levels. This suggests a synergistic interaction between the microbial inoculants, which form symbiotic associations with host plants, leading to stress mitigation in soybean under combined Fe deficiency and drought. However, AG58XF3 may possess metabolites other than flavonoids that facilitate symbiotic trade-offs with the enriched microbiomes. Host plants release a diverse array of metabolites to facilitate and promote symbiotic interactions with microbial partners (Hacquard and Martin, 2024). Hence, it is essential to study the genetic interplay of plants and their microbiomes along with environmental heterogeneity.

Conclusion

This study embodies the first detailed investigation into the complex interplay of physiological, molecular, and microbial factors improving stress tolerance in soybean plants exposed to combined Fe deficiency and drought. The ability of Clark to combat stress is associated with nutrient and redox balance, along with Fe retention and physiological adjustment, possibly driven by elevated jasmonic and salicylic acid in the roots. The upregulation of symbiotic genes and increased levels of siderophores and root flavonoids in Clark were correlated with improved nutrient status and microbial enrichment, predominantly by *Variovorax* and *Paecilomyces* in the endosphere. Furthermore, the compatibility and synergistic effects of enriched microbiomes, particularly in sensitive genotypes, highlight the potential of microbial inoculants to enhance stress resilience in soybean. Taken together, this study illustrates how host determinants and enriched microbiomes can be utilized in breeding programs and biofertilizer formulations to boost soybean productivity under challenging environmental conditions. By linking host determinants with microbial consortia, these findings lay the groundwork for innovative strategies in breeding and biofertilizer formulations with the potential to improve stress resilience in soybean and other legume crops under changing climatic conditions.

Acknowledgment

We express our gratitude to the Genomics Core of Michigan State University and CD Genomics. This research was supported by a startup grant (5SFAES-293007) and Elizabeth and Hayden Cutler's Endowed Professorship in Biotechnology from the University of Louisiana at Monroe. We also thank Mr. Timothy McMahan for his assistance with Nanopore Sequencing.

Conflict of interest

We have no conflict of interest.

Data availability

The data supporting the findings of this study are available in the article and its supplementary information files. Also, the sequencing data used and described in this study has been uploaded to NCBI under the Bioproject accession numbers: PRJNA1204368 (RNA-seq in root) and PRJNA1204372 (16S in root), PRJNA1204386 (ITS in roots) and PRJNA1204446 (16S in nodule).

Author contributions

MRH was involved in experimental design, plant cultivation, RNA extraction, RNA-seq data analysis, interpretation, 16S data analysis, split-root assay, microbial co-culture experiments, and prepared the manuscript draft. AT participated in plant phenotyping, performed biochemical assays, and extracted nodule DNA. AHK provided overall supervision, guidance in experimental design, data interpretation, and manuscript revision.

References

Álvarez-Aragón, R., Palacios, J. M., & Ramírez-Parra, E. (2023). Rhizobial symbiosis promotes drought tolerance in *Vicia sativa* and *Pisum sativum*. Environmental and Experimental Botany, 208, 105268.

Artursson, V. (2005). Bacterial-fungal interactions highlighted using microbiomics: Potential application for plant growth enhancement (Doctoral thesis, Swedish University of Agricultural Sciences). Acta Universitatis Agriculturae Sueciae, (2005:127). Department of Microbiology, Swedish University of Agricultural Sciences. https://res.slu.se/id/publ/13194

Astolfi, S., Celletti, S., Vigani, G., Mimmo, T., & Cesco, S. (2021). Interaction Between Sulfur and Iron in Plants. Frontiers in plant science, 12, 670308.

Bai, Y., Chen, T., Happe, T., Lu, Y., & Sawyer, A. (2018). Iron-sulphur cluster biogenesis via the SUF pathway. Metallomics, 10(8), 1038–1052.

Banasiak, J., Jamruszka, T., Murray, J. D., & Jasiński, M. (2021). A roadmap of plant membrane transporters in arbuscular mycorrhizal and legume-rhizobium symbioses. Plant physiology, 187(4), 2071–2091.

Barnard, R. L., Osborne, C. A., and Firestone, M. K. (2013). Responses of soil bacterial and fungal communities to extreme desiccation and rewetting. ISME J. 7, 2229–2241.

Barzana, G., Rios, J. J., Lopez-Zaplana, A., Nicolas-Espinosa, J., Yepes-Molina, L., Garcia-Ibañez, P., & Carvajal, M. (2021). Interrelations of nutrient and water transporters in plants under abiotic stress. Physiologia plantarum, 171(4), 595–619.

Bertrand, S., Bohni, N., Schnee, S., Schumpp, O., Gindro, K., & Wolfender, J. L. (2014). Metabolite induction via microorganism co-culture: a potential way to enhance chemical diversity for drug discovery. Biotechnology advances, 32(6), 1180–1204.

Breia R, Conde A, Badim H, Fortes AM, Gerós H, Granell A. Plant SWEETs: from sugar transport to plant-pathogen interaction and more unexpected physiological roles. Plant Physiol. 2021;186(2):836-852.

Briat JF, Ravet K, Arnaud N, Duc C, Boucherez J, Touraine B, Cellier F, Gaymard F. New insights into ferritin synthesis and function highlight a link between iron homeostasis and oxidative stress in plants. Ann Bot. 2010:105(5):811-22.

Briat, J. F., Dubos, C., & Gaymard, F. (2015). Iron nutrition, biomass production, and plant product quality. *Trends Plant Sci.* 20(1), 33–40.

Chen LQ, Qu XQ, Hou BH, Sosso D, Osorio S, Fernie AR, Frommer WB (2012) Sucrose efflux mediated by SWEET proteins as a key step for phloem transport. Science 335: 207–211

Chi, F., Shen, S. H., Cheng, H. P., Jing, Y. X., Yanni, Y. G., & Dazzo, F. B. (2005). Ascending migration of endophytic rhizobia, from roots to leaves, inside rice plants and assessment of benefits to rice growth physiology. Applied and environmental microbiology, 71(11), 7271–7278.

Chorianopoulou, S. N., & Bouranis, D. L. (2022). The Role of Sulfur in Agronomic Biofortification with Essential Micronutrients. Plants (Basel, Switzerland), 11(15), 1979.

Cohen, A. C., Bottini, R., Pontin, M., Berli, F. J., Moreno, D., Boccanlandro, H., et al., (2015). *Azospirillum brasilense* ameliorates the response of Arabidopsis thaliana to drought mainly via enhancement of ABA levels. Physiologia Planta. 153 (1), 79–90.

Connorton JM, Balk J, Rodríguez-Celma J. Iron homeostasis in plants - a brief overview. Metallomics. 2017;9(7):813-823.

Coudray, C., & Favier, A. (2000). Determination of salicylate hydroxylation products as an *in vivo* oxidative stress marker. Free radical biology & medicine, 29(11), 1064–1070.

Couturier, J., Przybyla-Toscano, J., Roret, T., Didierjean, C., and Rouhier, N. (2015). The roles of glutaredoxins ligating Fe-S clusters: Sensing, transfer or repair functions? *Biochimica et Biophysica Act.* 1853(6), 1513–1527.

Crowley D.E. Microbial siderophores in the plant rhizosphere. In: Barton L.L., Abadia J., editors. Fe nutrition in plants and rhizospheric microorganisms. Springer; Dordrecht, The Netherlands: 2006. pp. 169–198.

de Ollas, C., Arbona, V., & Gómez-Cadenas, A. (2015). Jasmonoyl isoleucine accumulation is needed for abscisic acid build-up in roots of Arabidopsis under water stress conditions. Plant, cell & environment, 38(10), 2157–2170.

Dhakarey R, Kodackattumannil Peethambaran P, Riemann M. (2016) Functional analysis of jasmonates in rice through mutant approaches. Plants (Basel). 2016;5(1):15.

D'Oria, A., Courbet, G., Billiot, B., Jing, L., Pluchon, S., Arkoun, M., Maillard, A., Roux, C. P., Trouverie, J., Etienne, P., Diquélou, S., and Ourry, A. (2022). Drought specifically downregulates mineral nutrition: Plant ionomic content and associated gene expression. *Plant Direct* 6(8), e402.

Du, X., Zhang, X., Chen, X., Jin, W., Huang, Z., & Kong, L. (2024). Drought stress reduces the photosynthetic source of subtending leaves and the transit sink function of podshells, leading to reduced seed weight in soybean plants. Frontiers in plant science, 15, 1337544.

Garcia, K., Cloghessy, K., Cooney, D. R., Shelley, B., Chakraborty, S., Kafle, A., Busidan, A., Sonawala, U., Collier, R., Jayaraman, D., Ané, J. M., & Pilot, G. (2023). The putative transporter *MtUMAMIT14* participates in nodule formation in *Medicago truncatula*. Scientific reports, 13(1), 804.

Ge SX, Jung D, Yao R. ShinyGO: a graphical gene-set enrichment tool for animals and plants. Bioinformatics. 2020;36(8):2628-2629.

Ghosh UK, Islam MN, Siddiqui MN, Khan MAR. Understanding the roles of osmolytes for acclimatizing plants to changing environment: a review of potential mechanism. Plant Signal Behav. 2021;16(8):1913306.

Gill, S. S., & Tuteja, N. (2010). Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. Plant physiology and biochemistry 48(12), 909–930.

Gonçalves, O. S., Creevey, C. J., & Santana, M. F. (2023). Designing a synthetic microbial community through genome metabolic modeling to enhance plant-microbe interaction. Environmental microbiome, 18(1), 81.

Guefrachi, I., Pierre, O., Timchenko, T., Alunni, B., Barrière, Q., Czernic, P., Villaécija-Aguilar, J. A., Verly, C., Bourge, M., Fardoux, J., Mars, M., Kondorosi, E., Giraud, E., & Mergaert, P. (2015). *Bradyrhizobium BclA* is a peptide transporter required for bacterial differentiation in symbiosis with aeschynomene lLegumes. Molecular plant-microbe interactions, 28(11), 1155–1166.

Gupta, A., Hisano, H., Hojo, Y., Matsuura, T., Ikeda, Y., Mori, I. C., & Senthil-Kumar, M. (2017). Global profiling of phytohormone dynamics during combined drought and pathogen

stress in Arabidopsis thaliana reveals ABA and JA as major regulators. Scientific reports, 7(1), 4017.

Hacquard, S., & Martin, F. M. (2024). The chemical language of plant-microbe-microbe associations: an introduction to a Virtual Issue. The New phytologist, 244(3), 739–742.

Hao DL, Zhou JY, Yang SY, Qi W, Yang KJ, Su YH. Function and regulation of ammonium transporters in plants. Int J Mol Sci. 2020;21(10):3557.

Harman, G., Khadka, R., Doni, F., & Uphoff, N. (2021). Benefits to Plant Health and Productivity From Enhancing Plant Microbial Symbionts. Frontiers in plant science, 11, 610065.

Hartmann, M., Brunner, I., Hagedorn, F., Bardgett, R. D., Stierli, B., Herzog, C., et al., (2017). A decade of irrigation transforms the soil microbiome of a semi-arid pine forest. Mol. Ecol. 26, 1190–1206.

Hassan, N. M., El-Bastawisy, Z. M., El-Sayed, A. K., Ebeed, H. T., & Nemat Alla, M. M. (2015). Roles of dehydrin genes in wheat tolerance to drought stress. Journal of advanced research, 6(2), 179–188.

Hassan, S., & Mathesius, U. (2012). The role of flavonoids in root-rhizosphere signalling: opportunities and challenges for improving plant-microbe interactions. Journal of experimental botany, 63(9),

Hou, K., Cao, L., Li, W., Fang, Z. H., Sun, D., Guo, Z., & Zhang, L. (2024). Overexpression of *Rhodiola crenulata* glutathione peroxidase 5 increases cold tolerance and enhances the pharmaceutical value of the hairy roots. Gene, 917, 148467.

Hussain HA, Hussain S, Khaliq A, Ashraf U, Anjum SA, Men S, Wang L. Chilling and drought stresses in crop plants: implications, cross talk, and potential management opportunities. Front Plant Sci. 2018:9:393.

Jahan, S., Tamta, S., Shankhdhar, S. C., & Shankhdhar, D. (2023). Salicylic acid potential to reversing drought-induced oxidative stress in *Bacopa monnieri* (L.) through enhancement of bioactive compound (Bacoside-A) and antioxidants including physio-biochemical attributes. South African Journal of Botany, 161, 617–626.

Jeong, H., Park, J., & Kim, H. (2013). Determination of NH☐☐ in environmental water with interfering substances using the modified Nessler method. Journal of Chemistry, 2013, Article ID 359217. https://doi.org/10.1155/2013/359217

Jiang, F., Chen, L., Belimov, A. A., Shaposhnikov, A. I., Gong, F., Meng, X., Hartung, W., Jeschke, D. W., Davies, W. J., & Dodd, I. C. (2012). Multiple impacts of the plant growth-promoting rhizobacterium *Variovorax paradoxus* 5C-2 on nutrient and ABA relations of Pisum sativum. Journal of experimental botany, 63(18), 6421–6430.

Jiang, M., & Zhang, J. (2002). Water stress-induced abscisic acid accumulation triggers the increased generation of reactive oxygen species and up-regulates the activities of antioxidant enzymes in maize leaves. Journal of experimental botany, 53(379), 2401–2410.

Jiang, S., Jardinaud, M. F., Gao, J., Pecrix, Y., Wen, J., Mysore, K., Xu, P., Sanchez-Canizares, C., Ruan, Y., Li, Q., Zhu, M., Li, F., Wang, E., Poole, P. S., Gamas, P., & Murray, J. D. (2021). NIN-like protein transcription factors regulate leghemoglobin genes in legume nodules. Science (New York, N.Y.), 374(6567), 625–628.

Kabir, A. H., and Bennetzen, J. L. (2024). Molecular insights into the mutualism that induces Fe deficiency tolerance in sorghum inoculated with *Trichoderma harzianum. Microbiol. Res.* 281, 127630.

Kabir, A. H., Paltridge, N. G., Able, A. J., Paull, J. G., and Stangoulis, J. C. (2012). Natural variation for Fe-efficiency is associated with upregulation of Strategy I mechanisms and enhanced citrate and ethylene synthesis in *Pisum sativum* L. Planta, 235(6), 1409–1419.

Kabir, A. H., Rahman, M. M., Haider, S. A., & Paul, N. K. (2015). Mechanisms associated with differential tolerance to Fe deficiency in okra (*Abelmoschus esculentus* Moench). Environmental and Experimental Botany, 112, 16–26.

Kabir, A. H., Thapa, A., Hasan, M. R., & Parvej, M. R. (2024). Local signal from Trichoderma afroharzianum T22 induces host transcriptome and endophytic microbiome leading to growth promotion in sorghum. Journal of experimental botany, 75(22), 7107–7126.

Kang, J. H., McRoberts, J., Shi, F., Moreno, J. E., Jones, A. D., & Howe, G. A. (2014). The flavonoid biosynthetic enzyme chalcone isomerase modulates terpenoid production in glandular trichomes of tomato. Plant physiology, 164(3), 1161–1174.

Kanwar, P., Baby, D., & Bauer, P. (2021). Interconnection of iron and osmotic stress signalling in plants: is FIT a regulatory hub to cross-connect abscisic acid responses? Plant biology 23 Suppl 1, 31–38.

Kehe, J., Ortiz, A., Kulesa, A., Gore, J., Blainey, P. C., & Friedman, J. (2021). Positive interactions are common among culturable bacteria. Science advances, 7(45), eabi7159.

Khan A. L. (2023). The phytomicrobiome: solving plant stress tolerance under climate change. Frontiers in plant science, 14, 1219366.

Kidwai, M., Ahmad, I. Z., & Chakrabarty, D. (2020). Class III peroxidase: an indispensable enzyme for biotic/abiotic stress tolerance and a potent candidate for crop improvement. Plant cell reports, 39(11), 1381–1393.

Kim, E. H., Kim, Y. S., Park, S. H., Koo, Y. J., Choi, Y. D., Chung, Y. Y., Lee, I. J., & Kim, J. K. (2009). Methyl jasmonate reduces grain yield by mediating stress signals to alter spikelet development in rice. Plant physiology, 149(4), 1751–1760.

Kobayashi, T., and Nishizawa, N. K. (2012). Fe uptake, translocation, and regulation in higher plants. *Ann. Rev. Plant Biol.* 63, 131–152.

Kroh, G.E., and Pilon, M. (2019). Connecting the negatives and positives of plant Fe homeostasis. *New Phytol.* 223(3), 1052-1055.

Kryukov, A. A., Gorbunova, A. O., Kudriashova, T. R., Yakhin, O. I., Lubyanov, A. A., Malikov, U. M., Shishova, M. F., Kozhemyakov, A. P., & Yurkov, A. P. (2021). Sugar transporters of the SWEET family and their role in arbuscular mycorrhiza. Vavilovskii zhurnal genetiki i selektsii, 25(7), 754–760.

Küçük, C., & Cevheri, C. (2014). Nodulation study of natural forage legume in semiarid region, Turkey. Pakistan journal of biological sciences 17(4), 535–539.

Kumar, M., Kumar Patel, M., Kumar, N., Bajpai, A. B., and Siddique, K. H. M. (2021). Metabolomics and molecular approaches reveal drought stress tolerance in plants. *Int. J. Mol. Sci.* 22(17), 9108.

Kurepa, J., & Smalle, J. A. (2022). Auxin/cytokinin antagonistic control of the shoot/root growth ratio and its relevance for adaptation to drought and nutrient deficiency stresses. International journal of molecular sciences, 23(4), 1933.

Laporte D, Olate E, Salinas P, Salazar M, Jordana X, Holuigue L. Glutaredoxin *GRXS13* plays a key role in protection against photooxidative stress in *Arabidopsis*. J Exp Bot. 2012;63(1):503-515.

Larrainzar, E., Wienkoop, S., Weckwerth, W., Ladrera, R., Arrese-Igor, C., & González, E. M. (2007). Medicago truncatula root nodule proteome analysis reveals differential plant and bacteroid responses to drought stress. Plant physiology, 144(3), 1495–1507.

Li J, Cao X, Jia X, et al., Iron deficiency leads to chlorosis through impacting chlorophyll synthesis and nitrogen metabolism in *Areca catechu* L. Front Plant Sci. 2021;12:710093.

Li, P., Yang, H., Wang, L., Liu, H., Huo, H., Zhang, C., Liu, A., Zhu, A., Hu, J., Lin, Y., & Liu, L. (2019). Physiological and transcriptome analyses reveal short-term responses and formation of memory under drought stress in rice. Frontiers in genetics, 10, 55.

Li, S., Yang, L., Huang, X., Zou, Z., Zhang, M., Guo, W., Addo-Danso, S. D., and Zhou, L. (2023). Mineral nutrient uptake, accumulation, and distribution in *Cunninghamia lanceolata* in response to drought stress. *Plants* 12(11), 2140.

Li, W., and Lan, P. (2017). The understanding of the plant Fe deficiency responses in strategy I plants and the role of ethylene in this process by omics approaches. *Front Plant Sci.* 8, 40.

Liang G. (2022). Fe uptake, signaling, and sensing in plants. *Plant Comm.* 3(5), 100349.

Lidoy J, Berrio E, García M, España-Luque L, Pozo MJ, López-Ráez JA. (2023). Flavonoids promote *Rhizophagus irregularis* spore germination and tomato root colonization: A target for sustainable agriculture. *Front Plant Sci.* 5, 1094194.

Lindström K, Mousavi SA. Effectiveness of nitrogen fixation in rhizobia. Microb Biotechnol. 2020;13(5):1314-1335.

Liu ZJ, Shan LL, Chen XF. Identification and growth-promoting effect of *Paecilomyces lilacinus* a biocontrol fungi for walnut rot disease. PLoS One. 2024;19(12):e0314160.

Longley R, Noel ZA, Benucci GMN, Chilvers MI, Trail F, Bonito G. Crop management impacts the soybean (*Glycine max*) microbiome. Front Microbiol. 2020;11:1116.

Loo, E. P., Durán, P., Pang, T. Y., Westhoff, P., Deng, C., Durán, C., Lercher, M., Garrido-Oter, R., & Frommer, W. B. (2024). Sugar transporters spatially organize microbiota colonization along the longitudinal root axis of *Arabidopsis*. Cell host & microbe, 32(4), 543–556.e6.

Lucas, M., Balbín-Suárez, A., Smalla, K., & Vetterlein, D. (2018). Root growth, function and rhizosphere microbiome analyses show local rather than systemic effects in apple plant response to replant disease soil. PloS one, 13(10), e0204922.

Lucena, C., Romera, F. J., García, M. J., Alcántara, E., and Pérez-Vicente, R. (2015). Ethylene participates in the regulation of Fe deficiency responses in strategy I plants and in rice. *Front Plant Sci.* 6, 1056.

Lumactud, R. A., Dollete, D., Liyanage, D. K., Szczyglowski, K., Hill, B., & Thilakarathna, M. S. (2023). The effect of drought stress on nodulation, plant growth, and nitrogen fixation in soybean during early plant growth. Journal of Agronomy and Crop Science, 209(3), 345–354.

Mendes, R., Garbeva, P., & Raaijmakers, J. M. (2013). The rhizosphere microbiome: significance of plant beneficial, plant pathogenic, and human pathogenic microorganisms. FEMS microbiology reviews, 37(5), 634–663.

Morrissey, J., and Guerinot, M. L. (2009). Fe uptake and transport in plants: the good, the bad, and the ionome. *Chem Rev.* 109(10), 4553–4567.

Mubarik, M. S., Khan, S. H., Sajjad, M., Raza, A., Hafeez, M. B., Yasmeen, T., Rizwan, M., Ali, S., & Arif, M. S. (2021). A manipulative interplay between positive and negative regulators of phytohormones: A way forward for improving drought tolerance in plants. Physiologia plantarum, 172(2), 1269–1290.

Munemasa S, Hauser F, Park J, Waadt R, Brandt B, Schroeder JI. Mechanisms of abscisic acid-mediated control of stomatal aperture. Curr Opin Plant Biol. 2015;28:154-162.

Musa, M., Jan, F. G., Hamayun, M., Jan, G., Khan, S. A., Rehman, G., Ali, S., & Lee, I. J. (2023). An endophytic fungal isolate *Paecilomyces lilacinus* produces bioactive secondary metabolites and promotes growth *of Solanum lycopersicum* under heavy metal stress. Agronomy, 13(3), 883.

Naveed, M., Hussain, M. B., Zahir, Z. A., Mitter, B., Sessitsch, A. (2014). Drought stress amelioration in wheat through inoculation with *Burkholderia phytofirmans* strain PsJN. Plant Growth Regul. 73, 121–131.

Naylor, D., & Coleman-Derr, D. (2018). Drought stress and root-associated bacterial Communities. Frontiers in plant science, 8, 2223.

Naylor, D., DeGraaf, S., Purdom, E., & Coleman-Derr, D. (2017). Drought and host selection influence bacterial community dynamics in the grass root microbiome. The ISME journal, 11(12), 2691–2704.

Ndou N, Rakgotho T, Nkuna M, Doumbia IZ, Mulaudzi T, Ajayi RF. Green synthesis of iron oxide (hematite) nanoparticles and their influence on *Sorghum bicolor* growth under drought stress. Plants 2023:12(7):1425.

Piyanete, C., Meechai, P. and Nakbanpotecc, W. Antioxidant activities and phenolic contents of extracts from *Salvinia olesta* and *Eichorniacrassipes*. Res. J. Biol. Sci. 4, 1113–1117 (2009).

Peleg, Z., & Blumwald, E. (2011). Hormone balance and abiotic stress tolerance in crop plants. Current opinion in plant biology, 14(3), 290–295.

Prigigallo, M. I., Staropoli, A., Vinale, F., & Bubici, G. (2023). Interactions between plant-beneficial microorganisms in a consortium: *Streptomyces microflavus* and *Trichoderma harzianum*. Microbial biotechnology, 16(12), 2292–2312.

Qiao M, Hong C, Jiao Y, Hou S, Gao H. Impacts of drought on photosynthesis in major food crops and the related mechanisms of plant responses to drought. Plants (Basel). 2024;13(13):1808.

Rai, G. K., Khanday, D. M., Choudhary, S. M., Kumar, P., Kumari, S., Martínez-Andújar, C., Martínez-Melgarejo, P. A., Rai, P. K., & Pérez-Alfocea, F. (2024). Unlocking nature's stress buster: Abscisic acid's crucial role in defending plants against abiotic stress. Plant Stress, 11, 100359.

Richards, S. L., Wilkins, K. A., Swarbreck, S. M., Anderson, A. A., Habib, N., Smith, A. G., McAinsh, M., & Davies, J. M. (2015). The hydroxyl radical in plants: from seed to seed. Journal of experimental botany, 66(1), 37–46.

Saini, A., Mapolelo, D. T., Chahal, H. K., Johnson, M. K., & Outten, F. W. (2010). SufD and SufC ATPase activity are required for iron acquisition during in vivo Fe-S cluster formation on SufB. Biochemistry, 49(43), 9402–9412.

Salamon, S., Mikołajczak, K., Błaszczyk, L., Ratajczak, K., & Sulewska, H. (2020). Changes in root-associated fungal communities in *Triticum aestivum* ssp. spelta L. and *Triticum aestivum* ssp. vulgare L. under drought stress and in various soil processing. PloS one, 15(10), e0240037.

Santi, S., & Schmidt, W. (2009). Dissecting iron deficiency-induced proton extrusion in Arabidopsis roots. The New phytologist, 183(4), 1072–1084.

Santos, C. S., Sousa, C., Bagheri, M., Pinho, S., & Vasconcelos, M. W. (2023). Predicting iron deficiency and oxidative stress in *Glycine max* through Fourier transform infrared spectroscopy in a time-course experiment. Plant and Soil, 496(1), 161–177.

Santoyo, G., Guzmán-Guzmán, P., Parra-Cota, F. I., de los Santos-Villalobos, S., Orozco-Mosqueda, M. del C., & Glick, B. R. (2021). Plant growth stimulation by microbial consortia. Agronomy, 11(2), 219.

Sun SL, Yang WL, Fang WW, Zhao YX, Guo L, Dai YJ. The plant growth-promoting Rhizobacterium *Variovorax boronicumulans* CGMCC 4969 regulates the level of indole-3-acetic acid synthesized from indole-3-acetonitrile. Appl Environ Microbiol. 2018;84(16):e00298-18.

Sun, C., Wu, T., Zhai, L., Li, D., Zhang, X., Xu, X., Ma, H., Wang, Y., & Han, Z. (2016). Reactive oxygen species function to mediate the Fe deficiency response in an Fe-efficient apple genotype: an early response mechanism for enhancing reactive oxygen production. Frontiers in plant science, 7, 1726.

Taheri P. (2022). Crosstalk of nitro-oxidative stress and iron in plant immunity. Free radical biology & medicine, 191, 137–149.

Thapa A, Hasan MR, Kabir AH. 2024a. Transcriptional reprogramming and microbiome dynamics in garden pea exposed to high pH stress during vegetative stage. BioRxiv, https://doi.org/10.1101/2024.10.05.616821

Thapa A, Hasan MR, Kabir AH. 2024b. *Trichoderma afroharzianum* T22 induces rhizobia and flavonoid-driven symbiosis to promote iron deficiency tolerance in garden pea. BioRxiv, https://doi.org/10.1101/2024.07.11.603139

Therby-Vale, R., Lacombe, B., Rhee, S. Y., Nussaume, L., & Rouached, H. (2022). Mineral nutrient signaling controls photosynthesis: focus on iron deficiency-induced chlorosis. Trends in plant science, 27(5), 502–509.

Tohge, T., de Souza, L. P., & Fernie, A. R. (2017). Current understanding of the pathways of flavonoid biosynthesis in model and crop plants. Journal of experimental botany, 68(15), 4013–4028.

Trivedi, P., Batista, B. D., Bazany, K. E., & Singh, B. K. (2022). Plant-microbiome interactions under a changing world: responses, consequences and perspectives. The New phytologist, 234(6), 1951–1959.

Wang Z, Li G, Sun H, et al., Effects of drought stress on photosynthesis and photosynthetic electron transport chain in young apple tree leaves. Biol Open. 2018;7(11):bio035279.

Wang, H., Lu, S., Guan, X., Jiang, Y., Wang, B., Hua, J., & Zou, B. (2022). Dehydration-responsive element binding protein 1C, 1E, and 1G promote stress tolerance to chilling, heat, drought, and salt in rice. Frontiers in plant science, 13, 851731.

Wang, N., Wang, T., Chen, Y., Wang, M., Lu, Q., Wang, K., Dou, Z., Chi, Z., Qiu, W., Dai, J., Niu, L., Cui, J., Wei, Z., Zhang, F., Kümmerli, R., & Zuo, Y. (2024). Microbiome convergence enables siderophore-secreting-rhizobacteria to improve iron nutrition and yield of peanut intercropped with maize. Nature communications, 15(1), 839.

Wasternack, C., & Song, S. (2017). Jasmonates: biosynthesis, metabolism, and signaling by proteins activating and repressing transcription. Journal of experimental botany, 68(6), 1303–1321.

Waters BM, Blevins DG, Eide DJ (2002) Characterization of FRO1, a pea ferric-chelate reductase involved in root iron acquisition. Plant Physiol 129:85–94

Wolter, D. J., & Hoffman, L. R. (2014). Strength in diversity. Cell host & microbe, 16(4), 427–429.

Wu, C., Lin, M., Chen, F., Chen, J., Liu, S., Yan, H., & Xiang, Y. (2022). Homologous Drought-Induced 19 Proteins, PtDi19-2 and PtDi19-7, Enhance drought tolerance in transgenic plants. International journal of molecular sciences, 23(6), 3371.

Wu, M., Liu, H., Gao, Y., Shi, Y., Pan, F., & Xiang, Y. (2020). The moso bamboo drought-induced 19 protein PheDi19-8 functions oppositely to its interacting partner, PheCDPK22, to modulate drought stress tolerance. Plant science 299, 110605.

Xu L, Coleman-Derr D. Causes and consequences of a conserved bacterial root microbiome response to drought stress. Curr. Opin. Microbiol. 2019:49:1–6.

Xu, F., Wang, K., Yuan, W., Xu, W., Shuang, L., Kronzucker, H. J., Chen, G., Miao, R., Zhang, M., Ding, M., Xiao, L., Kai, L., Zhang, J., & Zhu, Y. (2019). Overexpression of rice aquaporin OsPIP1;2 improves yield by enhancing mesophyll CO_2 conductance and phloem sucrose transport. Journal of experimental botany, 70(2), 671–681.

Xu, L., Dong, Z., Chiniquy, D., Pierroz, G., Deng, S., Gao, C., Diamond, S., Simmons, T., Wipf, H. M.-L., Caddell, D., Varoquaux, N., Madera, M. A., Hutmacher, R., Deutschbauer, A., Dahlberg, J. A., Guerinot, M. L., Purdom, E., Banfield, J. F., Taylor, J. W., ... Coleman-Derr, D. (2021). Genome-resolved metagenomics reveals role of iron metabolism in drought-induced rhizosphere microbiome dynamics. Nature Communications, 12(3209).

Yang, J., Duan, G., Li, C., Liu, L., Han, G., Zhang, Y., & Wang, C. (2019). The crosstalks between jasmonic acid and other plant hormone signaling highlight the involvement of jasmonic acid as a core component in plant response to biotic and abiotic stresses. Frontiers in plant science, 10, 1349.

Yuste, J. C., Peñuelas, J., Estiarte, M., Garcia-Mas, J., Mattana, S., Ogaya, R., et al., (2011). Drought-resistant fungi control soil organic matter decomposition and its response to temperature. Glob. Change Biol. 17, 1475–1486.

Zhang H, Song J, Dong F, Li Y, Ge S, Wei B, Liu Y. Multiple roles of wheat ferritin genes during stress treatment and *TaFER5D-1* as a positive regulator in response to drought and salt tolerance. Plant Physiol Biochem. 2023:202:107921.

Zhang Y, Xu J, Li R, Ge Y, Li Y, Li R. Plants' response to abiotic stress: mechanisms and strategies. Int J Mol Sci. 2023;24(13):10915.

Zhang, D. L., Ghosh, M. C., & Rouault, T. A. (2014). The physiological functions of iron regulatory proteins in iron homeostasis - an update. Frontiers in pharmacology, 5, 124.

Zhang, J., Cook, J., Nearing, J. T., Zhang, J., Raudonis, R., Glick, B. R., Langille, M. G. I., & Cheng, Z. (2021). Harnessing the plant microbiome to promote the growth of agricultural crops. Microbiological research, 245, 126690.

Zhang, L., Wu, M., Teng, Y., Jia, S., Yu, D., Wei, T., Chen, C., & Song, W. (2019). Overexpression of the glutathione peroxidase 5 (*RcGPX5*) gene from *Rhodiola crenulata* increases drought tolerance in *Salvia miltiorrhiza*. Frontiers in plant science, 9, 1950.

Zhang, Y., Yang, L., Zhang, M., Yang, J., Cui, J., Hu, H., & Xu, J. (2022). CfAPX, a cytosolic ascorbate peroxidase gene from *Cryptomeria fortunei*, confers tolerance to abiotic stress in transgenic Arabidopsis. Plant physiology and biochemistry, 172, 167–179.

Zocchi, G., De Nisi, P., Dell'Orto, M., Espen, L., & Gallina, P. M. (2007). Iron deficiency differently affects metabolic responses in soybean roots. Journal of experimental botany, 58(5), 993–1000.

Table 1. ICP-MS analysis of nutrient content in soybean roots and leaves under control and Fe-Drought+ conditions. Different letters denote significant differences between control and Fe-Drought+ conditions ($P < 0.05$, t-test). Data are presented as means ± SD from three independent biological samples ($n = 3$).

Elements	Genotypes	Root		Leaf	
		Control	Fe-Drought+	Control	Fe-Drought+
Fe (ppm)	Clark	160.6±19.7[a]	142.1±15.01[a]	55.6±19.89[a]	52.3±15.46[a]
	Arisoy	195.6±9.5[a]	59.3±10.4[b]	63.6±13.01[a]	34.6±10.36[b]
Zn (ppm)	Clark	25.7±4.93[a]	40.7±12.61[a]	33.0±1.22[a]	26.2±4.75[a]
	Arisoy	26.9±7.48[a]	38.5±11.42[a]	26.8±6.90[a]	39.5±28.74[a]
Mn (ppm)	Clark	72.0±9.75[a]	61.3±10.64[b]	41.1±4.38[a]	37.1±6.54[a]
	Arisoy	64.7±12.98[a]	55.8±15.23[b]	37.9±1.75[a]	22.5±3.98[b]
S (%)	Clark	0.2±0.02[a]	0.2±0.05[a]	0.5±0.03[a]	0.2±0.03[b]
	Arisoy	0.2±0.01[a]	0.2±0.02[a]	0.6±0.12[a]	0.2±0.06[b]
N (%)	Clark	3.3±0.33[a]	3.5±0.54[a]	3.4±0.05[a]	3.7±0.54[a]
	Arisoy	4.8±0.34[a]	2.3±0.44[b]	3.5±0.24[a]	2.2±0.1[b]

Table 2. Changes in flavonoid content, total soluble sugar and ammonium content in the roots of soybean grown in control and Fe-Drought+ conditions. Different letters indicate significant differences between the control and Fe-Drought+ conditions at a $P < 0.05$ level, based on *t*-test. The data represents the means ± SD of three independent biological samples ($n = 3$).

Genotypes	Flavonoid (mg g^{-1} DW)		Soluble sugar (mg g^{-1} FW)		Ammonium (µg g^{-1} FW)	
	Control	Fe-Drought+	Control	Fe-Drought+	Control	Fe-Drought+
Clark	12.1±0.68[a]	17.6±0.86[b]	11.2±3.36[a]	10.7±3.28[a]	31.0±7.13[a]	29.4±7.14[a]
Arisoy	14.4±0.8[a]	10.2±1.15[b]	10.7±3.40[a]	6.3±1.43[b]	30.0±6.05[a]	20.2±4.41[b]

Fig. 1. Growth, physiological, and biochemical responses of Clark and Arisoy genotypes under control and Fe-Drought+ conditions. A) aboveground phenotypes, B) number of nodules, C) root length, D) root dry weight, E) shoot height, F) shoot dry weight, G) leaf relative water content (RWC), H) SPAD chlorophyll content, I) photosynthetic efficiency (Fv/Fm), J) hydrogen peroxide (H_2O_2) in roots, K) hydroxyl radical (•OH) in roots, L) Fenton reaction rate in roots and M) ferric chelate reductase activity in roots. Data are presented as means ± SD ($n = 3$), with

significance levels indicated by *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, and ns (not significant), based on a *t*-test.

Fig. 2. RNA-seq analysis in the roots of Clark and Arisoy under control and Fe-Drought+ conditions. (A, B) principal component analysis (PCA) in Clark (A) and Arisoy (B), volcano plots depicting differential gene expression in Clark (C) and Arisoy (D) with upregulated genes in red, downregulated genes in blue, and non-significant genes shown in gray and Heatmaps showing DEGs (log2 fold, P <0.05) related to key functional categories: symbiotic association (E), nutrient uptake and transport (F), osmotic adjustment (G), and ROS detoxification (H). Data represent three biological replicates, with color scales indicating the log2-fold changes in expression levels.

Fig. 3. Functional enrichment analysis of biological processes and KEGG pathways in Clark and Arisoy under Fe-Drought+ stress. Enriched biological processes and KEGG pathways in Clark (A) and enriched biological processes and KEGG pathways in Arisoy (B). Data were analyzed using ShinyGO (V. 0.81), with fold enrichment and FDR significance shown on the plots.

Fig. 4. Hormonal profiles in the roots of Clark and Arisoy genotypes under control and Fe-Drought+ conditions. A) Indole-3-acetic acid (IAA), B) Abscisic acid (ABA), C) Jasmonic acid (JA), and D) Salicylic acid (SA). Data are presented as means ± SD of three biological replicates. Pink and green boxplots represent control and Fe-Drought+ conditions, respectively. Significant differences between treatments are indicated ($^*P < 0.05$, $^{**}P < 0.01$; ns = not significant), based on a *t*-test.

Fig. 5. Siderophore levels and bacterial (16S) community composition and diversity in Clark and Arisoy under control and Fe-Drought+ conditions. Rhizosphere siderophore (A), principal component analysis (B), alpha diversity measures (C), the relative abundance of bacterial genera in the roots of Clark (D) and Arisoy (E), the relative abundance of bacterial genera in the nodules of Clark (F) and Arisoy (G). Asterisks (*) indicate significant differences based on the Kruskal-Wallis test ($P < 0.05$) based on three replicates ($n = 3$) per sample group.

Fig. 6. Fungal (ITS) community composition and diversity in the roots of Clark and Arisoy under control and Fe-Drought+ conditions. Principal component analysis (B), alpha diversity measures (B), the relative abundance of fungal genera in the roots of Clark (C) and Arisoy (D), and ASV abundance of *Variovorax* and *Paecilomyces* (E). Asterisks (*) indicate significant differences based on the Kruskal-Wallis test ($P < 0.05$) based on three replicates ($n = 3$) per sample group.

Fig. 7. Interactions of enriched microbiomes and their effects on morpho-physiological responses in Clark under split-root systems. Colony growth of *Paecilomyces lilacinus* and *Variovorax paradoxus* in nutrient agar (A), colony size of *P. lilacinus* and *V. paradoxus* over time (B), diagram and establishment of split-root (SR) system (C), aboveground and belowground phenotype of split-root plants (D), split-root assay: SPAD chlorophyll content (E), shoot height

(F), shoot dry weight (G), root length (H), root dry weight (I), and rhizosphere siderophore production (J). Letters indicate significant differences ($P < 0.05$, $n = 3$) between treatments based on ANOVA.

Fig. 8. Effect of enriched microbiomes on soybean genotypes (Arisoy and AG58XF3): aboveground phenotype (A), leaf SPAD value (B), plant height (C), plant biomass (D), leaf relative water content (E), root nodule number (F), rhizosphere siderophore (G) and root flavonoid content (H). The treatments include T1: Control, T2: Fe-Drought+, T3: Fe-Drought+ Microbiome, and T4: Microbiome+. Different letters indicate significant differences among treatments within each genotype ($P < 0.05$). Data are presented as means ± SD from three independent replicates. The enriched microbiomes included *V. paradoxus* and P. *lilacinus*.

Fig. 9. A model of stress resilience in Clark exposed to Fe deficiency and drought. The model illustrates the mechanisms underlying stress resilience, including enhanced Fe and N retention mediated by root-localized symbiotic interactions with *Bradyrhizobium*, *Paecilomyces*, and *Variovorax*. Key transporters such as *Ammonium transporter 1;2*, *EamA-like transporter*, *Sugar transporter 1*, and *SWEET10* facilitate nutrient exchange and carbon flow. Fe homeostasis is maintained through the activity of ferric chelate reductase (FCR), *Fe-regulated protein 3*, and siderophore-mediated Fe acquisition. Reactive oxygen species (ROS) are scavenged by *Peroxidase 19*, *Glutaredoxin 4*, and *Glutathione peroxidase 6*, while ferritin provides additional protection by inhibiting Fenton reactions. Flavonoids and sugars produced in the roots further support microbial recruitment and hormonal modulation involving ABA, JA, and SA, collectively enhancing physiological and stress responses. This integrated system highlights the synergy between microbial symbiosis and plant metabolic pathways in resilience to combined Fe deficiency and drought stress in soybean.