Strengthening Medial Olivocochlear Feedback Reduces the Developmental Impact of Early Noise Exposure

Authors: Valeria C. Castagna[1,2,3], Luis E. Boero[1,2,3], Mariano N. Di Guilmi[1], Paul A. Fuchs[4,5], Amanda M. Lauer[4,5,6], Ana Belén Elgoyhen[1,2] and Maria Eugenia Gomez-Casati[2]

Affiliations:

[1]Instituto de Investigaciones en Ingeniería Genética y Biología Molecular, Dr. Héctor N. Torres, Consejo Nacional de Investigaciones Científicas y Técnicas, 1428 Buenos Aires, Argentina.

[2]Instituto de Farmacología, Facultad de Medicina, Universidad de Buenos Aires, 1121 Buenos Aires, Argentina and Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina.

[3]Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Fisiología y Biología Molecular y Celular, 1428 Buenos Aires, Argentina.

[4]Department of Otolaryngology-Head and Neck Surgery, Johns Hopkins University School of Medicine; Baltimore, Maryland 21205, United States.

[5]Department of Neuroscience, Johns Hopkins University School of Medicine, Baltimore, Maryland 21205, United States.

[6]Center for Functional Anatomy and Evolution, Johns Hopkins University, School of Medicine, Baltimore, Maryland 21205, United States.

Corresponding Author: María Eugenia Gómez-Casati, Ph.D.

Instituto de Farmacología. Facultad de Medicina. Universidad de Buenos Aires.

Address: Paraguay 2155. C1121ABG, Buenos Aires, Argentina.

Phone: 54-11-52853350

26   E-mail: mgomezcasati@fmed.uba.ar, megomezcasati@gmail.com

Conflict of Interest: The authors declare no competing financial or non-financial interest.

Acknowledgements: This research was supported by Agencia Nacional de Promoción Científica y Técnica (Argentina) to M.E.G-C and NIH Grant R01 DC001508 (P.A.F, A.M.L, A.B.E and M.E.G-C) and the David M. Rubenstein Fund for Hearing Research.

Author Contributions: V.C.C, L.E.B, M.N.D-G, P.A.F, A.M.L, A.B.E and M.E.G-C designed research; V.C.C performed experiments; V.C.C analyzed data. M.E.G-C wrote the paper. All authors discussed the results, read, edited, and commented on the manuscript.

ABSTRACT

The early onset of peripheral deafness significantly disrupts the proper development of the auditory system. Likewise, exposure to loud noise during early development produces a similar disruptive effect. Before hearing onset in altricial mammals, cochlear inner hair cells exhibit spontaneous electrical activity that activates primary afferents and propagates into the central nervous system. This activity is crucial for auditory system maturation and is modulated by the medial olivocochlear efferent (MOC) feedback via α9α10 nicotinic cholinergic receptors present in inner hair cells. In adults, these receptors are restricted to outer hair cells, where they mediate feedback from the MOC system to regulate cochlear amplification. Although the MOC system's protective role to prevent noise-induced hearing loss in adulthood is well-established, its influence during early developmental stages -especially in response to exposure to loud noise- remains largely unexplored. In this study, we examined the effect of the MOC system in the ears of sound-exposed mice during the early postnatal developmental period, by using genetically modified mice - α9 knockout and α9 knock-in models - with either absent or enhanced cholinergic activity. Our findings reveal that both increased and absent olivocochlear activity result in altered auditory sensitivity at the onset of hearing, along with long–range alterations in the number and morphology of ribbon synapses. Furthermore, exposure to loud noise during early development causes long-lasting damage in both wild-type and α9 knockout mice, which lack functional MOC feedback, with effects persisting into adulthood. In contrast, mice with enhanced cholinergic activity were protected from noise-induced damage, with no long-term effects on auditory function. These results highlight the increased susceptibility of the auditory system during early postnatal development. Moreover, they indicate that an enhanced MOC feedback shields the auditory system from noise damage during this period.

INTRODUCTION

The development of the cochlea and the central auditory system is a complex and highly coordinated process, requiring precise timing of cellular and molecular events. During the early postnatal period, several critical changes take place that refine the auditory system's ability to process sound information [1]. Noise exposure during this developmental period can interfere with these processes, impairing the structural organization and signal processing capabilities of the mammalian auditory system and causing lasting damage to cochlear structures and neural circuits [2–6]. Early acoustic exposure affects neuronal sensitivity, innervation, frequency tuning and response characteristics of neurons at multiple levels of the auditory pathway [2,7–11]. Such disruptions can lead to permanent auditory deficits, including reduced sensitivity and impaired sound processing, emphasizing the importance of safeguarding the auditory system during this developmental window.

Cochlear inner hair cells (IHCs) and afferent neurons in altricial mammals exhibit spontaneous electrical activity that propagates into the central nervous system (CNS) prior to the onset of hearing, which occurs around postnatal day 12 (P12) in mice [12–17]. This spontaneous firing is thought to be crucial for promoting neuronal survival in peripheral and central auditory structures, driving their maturation, and shaping their synaptic connections through activity-dependent mechanisms [18–23]. In this postnatal developmental period, medial olivocochlear (MOC) neurons descending from the superior olivary complex in the brainstem make direct synaptic contact with immature IHCs, modulating their spontaneous activity. These MOC neurons form axosomatic synapses through α9α10 nicotinic cholinergic receptors (nAChRs), coupled to Ca^{2+}-dependent SK2 channels, resulting in hyperpolarization of IHCs [24–26]. This transient efferent input is thought to shape burst firing patterns, guiding the normal functional development of IHCs' synaptic machinery and ensuring the establishment of bilateral neural representation in the developing CNS [22,23,27–29]. As IHCs mature, they lose direct efferent contacts and no

longer respond to acetylcholine (ACh). At this stage, MOC fibers extend across the tunnel of Corti

and begin forming synapses with OHCs to regulate cochlear amplification [29–34]. The rodent

olivocochlear system matures rapidly within the first 2–3 postnatal weeks, with adult-like cochlear

innervation patterns established by the end of the first month [30,35–37]. ACh-induced responses

in OHCs emerge by P6, shifting from a nicotinic-like current to an outward potassium current

around the onset of hearing, even before efferent synapses are fully developed [38,39]. While the

protective role of the MOC system in preventing noise-induced hearing loss in adulthood is well-

documented [40–44], its impact during early developmental stages, particularly in response to

acoustic exposure, remains largely unexplored. Thus, the extent to which ascending and

descending pathways interact with the acoustic environment to shape activity patterns and

instruct neuronal circuit formation remain largely unknown.

This study explores how the strength of the cholinergic MOC input and early noise

exposure affect auditory system development, focusing on the largely unknown effects of noise

during this sensitive period. We used genetically modified mice with different levels of α9α10

nAChR activity: α9 knock-out mice (Chrna9 KO; α9KO), which lacks cholinergic transmission

between efferent neurons and hair cells [45], and α9 knock-in mice (Chrna9L9'T; α9KI) carrying

an α9 point mutation that leads to an enhanced MOC effect [42]. We show that animals genetically

engineered to have differences in MOC modulation strength affect the onset of hearing and alter

the number and morphology of ribbon synapses in mature IHCs, indicating the importance of

normal activity of the cholinergic component of the efferent pathway during postnatal auditory

development. Moreover, early-age noise exposure to loud noise produces lasting alterations in

auditory sensitivity in wild type (WT), with these effects being more pronounced compared to

sound exposure in adulthood [44]. In WT mice, adult noise exposure does not lead to permanent

threshold elevations and results in comparatively minor synaptic loss, highlighting the increased

susceptibility during postnatal development [44]. In α9KO mice, which lack efferent inhibition,

early-age exposure resulted in a persistent loss of auditory nerve synapses, emphasizing the

essential role of the MOC pathway. Conversely, in α9KI mice, where cholinergic efferent inhibition

is enhanced, early-age noise exposure does not lead to elevated auditory thresholds or synaptic

loss. This protective effect occurs during a critical developmental window when the auditory

system is still maturing and olivocochlear terminals are just beginning to establish synapses with

OHCs [30,31]. Thus, our work contributes to the evidence that cochlear efferent modulation during

postnatal development is essential for establishing normal auditory function. Additionally, we

demonstrate that early-age noise exposure can interfere with normal cochlear development and

have long-term consequences for function, which can be prevented by enhanced inhibitory input

from the MOC system.

RESULTS

Onset of auditory perception in mice with varying levels of efferent inhibition

In this study we used mice with distinct levels of α9α10 nAChR activity: 1) WT, with normal

MOC function; 2) α9KO, which lacks cholinergic transmission between MOC neurons and hair

cells [45], and 3) α9KI mice carrying an α9 nAChR subunit point mutation that leads to enhanced

responses to ACh [42]. To explore the impact of early-age noise exposure, we first identified the

optimal developmental stage for introducing such exposure, by assessing the onset of hearing

under control (unexposed) conditions in these mouse models. This was done by tracking auditory

brainstem responses (ABRs), the sound-evoked potentials generated by neuronal circuits in the

ascending auditory pathway, at different postnatal ages across the three groups. Individual ABR

waves reflect the activation of auditory periphery and brainstem processing relay stations within

the first ~7 ms after sound stimulus onset [46–51]. Wave 1 represents the synchrony in firing of

spiral ganglion neurons and is highly correlated with the number of synapses between IHCs and

auditory nerve fibers [46,52,53]. Measurements were taken daily from P12 to P16, as well as P22

and P75, in anesthetized mice. Interestingly, we found that ABR onset occurred earlier in α9KI

mice, with enhanced MOC function, compared to WT mice with normal MOC synapses and was

delayed in α9KO mice lacking α9α10 nAChR activity (Figure 1a and b). Representative ABR

traces recorded at P14 in response to a 16 kHz stimulus across varying intensities illustrate

differences in ABR thresholds among the three mouse groups with distinct levels of α9α10 nAChR

activity (Figure 1a). At this developmental stage, α9KI mice exhibited lower ABR thresholds, and

more distinct upward peaks at suprathreshold intensities compared to WT mice. In contrast, α9KO

mice showed higher ABR thresholds relative to WT mice (Figure 1a). Figure 1b shows the mean

ABR thresholds measured at test frequencies of 11.33, 16, and 22.65 kHz from P12 to P75 across

the three groups of mice. A comparison between α9KI and WT mice revealed that α9KI mice

displayed measurable ABR responses as early as P13 (Kruskal-Wallis, followed by Dunn's test,

degree of freedom [df]= 2, p =0.01 at 22.65 kHz; Figure 1b) and achieved mature threshold levels

earlier than their WT littermates (Kruskal-Wallis, followed by Dunn's test, p = 0.02 at 16 and 22.65

kHz at P14; Figure 1b). In contrast, ABR onset was delayed by a day in α9KO compared to WT

mice (Kruskal-Wallis followed by Dunn's test, df = 2, p = 0.04 at 11.33 and 16 kHz; Figure 1b),

indicating that cochlear maturation is slower in the absence of pre-hearing efferent modulation.

To summarize, at P13, no ABR responses were observed in WT (0 out of 9) or α9KO (0 out of 7)

mice in response to tone pips at 80 dB SPL across 11.33, 16, and 22.65 kHz. In contrast, 6 out

of 9 α9KI mice exhibited ABR responses at P13. By P14, ABR responses were present in 77.7%

of WT mice at 11.33 and 22.65 kHz and in 100% at 16 kHz. Meanwhile, 85% of α9KO mice

responded at all frequencies, but with higher thresholds than WT (Figure 1b). All α9KI mice

exhibited robust responses at all tested frequencies by P14. By P15, 100% of WT and 91% of

α9KO mice showed ABR responses.

Tone pip-evoked ABR wave 1 amplitudes were then analyzed at a suprathreshold level of

80 dB SPL at the different test frequencies in the three groups of mice with varying levels of α9α10

nAChR activity (Figure 2a, b). As illustrated in the representative traces in Figure 2a, measurable

ABR wave 1 amplitudes at P13 were only observed in α9KI mice. At P14, the number of

discernible upward peaks was fewer in WT and α9KO mice compared to α9KI mice. Up to P15,

response peaks at 80 dB SPL were slightly larger and more distinct in α9KI mice, reflecting

enhanced auditory responses during this postnatal developmental window. However, starting at

P16, α9KI mice exhibited smaller ABR wave 1 amplitudes, with significant differences observed

at P75 at 16 and 22.65 kHz compared to WT mice (one way ANOVA followed by Holm-Sidak test,

df=2, p=0.03 at both frequencies; Figure 2b). In α9KO mice, wave 1 amplitudes remained smaller

than those of WT and α9KI mice until P16, with the difference being significant at 22.65 kHz (one

way ANOVA followed by Holm-Sidak test, df = 2, p = 0.02; Figure 2b). These findings align with

the ABR threshold results, indicating that the strength of efferent cholinergic inhibition to hair cells

contributes to the timely development of auditory function. The reduced ABR wave 1 amplitudes

obtained in mature mice with either enhanced or null cholinergic activity indicates that proper

MOC modulation during development might be crucial for the correct formation of auditory nerve

synapses. Analysis of ABR wave 1 latencies at different postnatal ages showed no significant

differences between genotypes (Figure 2c). While wave 1 latencies were consistently longer

across all groups at P14, they shortened to levels typical of mature mice by P15.

OHC function was assessed through distortion-product otoacoustic emissions (DPOAE)

at the different postnatal ages. In a healthy cochlea, two close pure tones produce distortions due

to OHC nonlinearities, which are amplified by OHC electromotility and detected with a microphone

in the external ear canal [54,55]. DPOAE responses followed a similar pattern to ABR thresholds,

though the differences were not statistically significant (Kruskal-Wallis, df = 2, p > 0.05 at all

frequencies; Figure 1c). At P13, DPOAE responses were nearly absent in WT (1 out of 9) and

α9KO (0 out of 7) mice across the tested frequencies. In contrast, 3 out of 9 α9KI mice showed

DPOAE responses at 16 and 22.65 kHz. By P14, DPOAE responses were observed in 75% of

WT and α9KO mice at 16 and 22.65 kHz, and in 60% of mice at 11.33 kHz. All α9KI mice exhibited

strong DPOAE responses at all tested frequencies by P14. By P15, response rates reached 100%

in both WT and α9KO mice. Note that at P22 and P75, mean ABR and DPOAE thresholds in α9KI

mice were slightly elevated by 5–10 dB compared to WT and α9KO mice (Figure 1b, c). This

elevation reached statistical significance only for ABR thresholds at P22 (Kruskal-Wallis, df = 2,

p = 0.03 at 11.33 kHz; p = 0.04 at 16 kHz and p = 0.02 at 22.65 kHz). The elevated thresholds in

α9KI mice at these mature stages suggest that this may be due to enhanced cholinergic activity

on OHCs, as previously reported [42]. Collectively, these findings indicate that the onset of

hearing is altered in mice with either absent or enhanced olivocochlear cholinergic feedback. Due

to this observation, we decided to expose WT, α9KO, and α9KI mice to loud noise at P15 to test

the effects of exposure to increased acoustic stimulation. At this stage, all three genotypes

exhibited measurable ABR waveforms, though their wave 1 thresholds differed and had not yet

reached mature levels (Figure 1 and 2).

ABR thresholds after early-age noise exposure in mice with different degree of MOC

inhibition

To investigate the impact of early-age noise exposure, we exposed WT, α9KO, and α9KI

mice to loud noise at P15. This allowed us to evaluate how acoustic experience influences

auditory function before full threshold maturation in each genotype. Mice were exposed to 1–16

kHz noise at 100 dB SPL for 1 hour at P15. Auditory responses were then assessed at 1-, 7-, and

60-days post-exposure to track functional changes over time.

Previous studies in WT mice exposed to the same noise protocol at P21 reported a

significant increase in auditory thresholds of 10–35 dB SPL one day after exposure, which

returned to baseline within a week, indicating a transient threshold shift [44]. In contrast, as shown

in Figure 3a, no changes in ABR thresholds were observed at P16, one day after noise exposure,

across frequencies of 8, 16, and 22.65 kHz in WT mice compared to unexposed controls. Remarkably, ABR thresholds at P16 were slightly lower than those measured at P15, prior to noise exposure, and showed no significant difference when compared to unexposed controls at the same age. This suggests that noise did not have an immediate effect on the maturation of auditory thresholds at this postnatal developmental stage. One week after exposure, however, ABR thresholds were elevated at all frequencies in noise-exposed WT animals, and this increase persisted for at least two months post-exposure (Mann- Whitney test, df = 1, P22: p = 0.006 at 8 kHz; p = 0.007 at 16 kHz; p = 0.002 at 22.65 kHz and P75: p = 0.04 at 8 kHz; p = 0.03 at 16 kHz and p = 0.04 at 22.65 kHz; Figure 3a). This outcome contrasts with the transient threshold elevation observed in WT mice, from the same genetic background, exposed at P21 using the same noise protocol [44], suggesting that noise exposure at this early developmental stage produces more detrimental long-term damage.

A similar trend as seen in WT mice was observed after exposing α9KO mice to noise at P15 (Figure 3b). One day after exposure, there were no changes in ABR thresholds at 8, 16 and 22.65 kHz. However, seven days post-exposure, there was an elevation in auditory thresholds across all frequencies (Mann-Whitney test, df = 1, p = 0.03 at 8 kHz; p = 0.01 at 16 kHz and p = 0.02 at 22.65 kHz), which, unlike in WT mice, returned to normal levels after two months (Mann- Whitney test, df = 1, p > 0.05 at all frequencies; Figure 3b). Notably, in α9KI mice, ABR thresholds were not affected by noise exposure at all the frequencies tested (Figure 3c), suggesting that even at this early age, the enhancement of cholinergic activity on OHCs provides protection from noise-induced cochlear threshold elevations (Mann-Whitney test, df = 1, p > 0.05 at all frequencies).

Amplitudes of cochlear sound-evoked potentials after early-age noise exposure in mice with different degree of efferent inhibition

To assess the neural activity across auditory nerve fibers, we measured ABR wave 1

amplitudes at frequencies of 8, 16, and 22.65 kHz at different ages after noise exposure in the

three mouse groups (Figure 4). One day after noise exposure, we observed a reduction in wave

1 amplitudes across all frequencies in WT mice. Further testing of WT mice one week and two

months post-exposure revealed a significant and long-lasting reduction, indicating a permanent

decrease of ABR wave 1 amplitude (Mann- Whitney test, df = 1, P22: p = 0.007 at 8 kHz; p =

0.005 at 16 kHz; p = 0.005 at 22.65 kHz and P75: p = 0.04 at 8 kHz; p = 0.002 at 16 kHz and p =

0.002 at 22.65 kHz; Figure 4a). Similar permanent reductions in wave 1 amplitude were observed

in α9KO mice after exposure to noise at 16 and 22.65 kHz (Mann-Whitney test, df = 1, P22: p =

0.04 at 16 kHz; p = 0.001 at 22.65 kHz and P75: p = 0.03 at 16 kHz and p = 0.04 at 22.65 kHz;

Figure 4b). Notably, no changes in ABR wave 1 amplitudes were observed at any age post-

exposure in mice with enhanced α9α10 nAChR activity, suggesting a preservation of cochlear

synapses after noise exposure (Mann-Whitney test, df = 1, p > 0.05 at all frequencies; Figure 4c).

DPOAE thresholds after early-age noise exposure in mice with different levels of α9α10

nAChR activity

To assess the impact of early-age noise exposure on the function of OHCs, we recorded

DPOAEs at 8, 16, and 22.65 kHz on the three groups of mice with varying levels of MOC function

(Figure 5). During early development, the auditory system undergoes significant maturation,

leading to decreasing DPOAE thresholds with age, as shown in Figure 1c. In WT mice, one day

after noise exposure at P16, there was no difference in DPOAE thresholds between the noise-

exposed group and unexposed controls (Figure 5a). However, when comparing DPOAE

thresholds at P16 to those at P15 within both groups (i.e. exposed and unexposed controls), a

clear reduction was observed (Figure 5a). This decline in thresholds underscores the natural

maturation process of the auditory system, indicating that early noise exposure does not

immediately disrupt the ongoing development of DPOAE thresholds at this early postnatal stage.

By P22, seven days after noise exposure, an increase in DPOAE thresholds was observed that

was statistically significant only at 16 kHz (Mann-Whitney, df = 1, p = 0.01; Figure 5a). Yet, two

months later, DPOAE thresholds were comparable to those of unexposed WT controls across all

tested frequencies, suggesting that early noise exposure did not result in long-term changes in

OHC function (Figure 5a).

In α9KO mice, DPOAE thresholds remained unchanged 1 day after noise exposure.

However, seven days post-exposure, significant threshold elevations were observed at 8 and 16

kHz, but not at 22.65 kHz (Mann-Whitney, df = 1, p = 0.02 at 8 kHz and p = 0.02 at 16 kHz). After

two months, recovery to unexposed levels occurred only at 8 kHz, while DPOAE thresholds

remained elevated at 16 kHz (Mann-Whitney, df = 1, p = 0.003 at 16 kHz; Figure 5b). In contrast,

mice with enhanced MOC function showed no changes in DPOAE thresholds at any post-

exposure time point (Figure 5c), suggesting that early-age noise exposure did not impair the

cochlear amplifier function in these mice with enhanced OHC inhibition.

Ribbon synapse changes after early noise exposure in mice with different levels of α9α10

nAChR activity

Histological analysis was performed on cochleae dissected and fixed at P75 in the three

groups of mice with varying levels of α9α10 nAChR activity to evaluate the long-term effect of

early noise exposure on the density and structure of IHC ribbon synapses. To visualize these

ribbon synapses, we performed immunostaining on whole-mount preparations of the organ of

Corti with antibodies against CtBP2-Ribeye, a presynaptic ribbon protein [56], and GluA2, an

AMPA-type glutamate receptor expressed at the postsynaptic afferent terminal [43,57,58]. Ribbon

synapses were identified by the colocalization of CtBP2 and GluA2 puncta at the base of the IHC,

as shown in representative confocal images of IHC synapses from apical whole mount organ of

Corti from control (unexposed) WT, α9KO, and α9KI mice (Supplementary Figure 1a) [58]. We compared the number of colocalized synaptic markers per IHC across three cochlear regions, each representing a distinct frequency range: apex (~4–12 kHz), middle turn (~12–20 kHz), base (~22–40 kHz) (Supplementary Figure 1b). In the low-frequency apical end, both unexposed α9KO and unexposed α9KI mice showed a significant decrease in synaptic density compared to WT, reduced 30.64% and 27.94%, respectively (Kruskal-Wallis, df=2, p<0.0001 for both groups; Supplementary Figure 1b, left panel). In the middle region, the reduction was significant only in α9KI mice, with a 35.97% decrease compared to WT (Kruskal-Wallis, df = 2, p < 0.0001; Supplementary Figure 1b, middle panel). Similarly, in the high-frequency basal region, synaptic puncta were significantly reduced only in α9KI mice, showing a 22.07% decrease relative to WT (Kruskal-Wallis, df = 2, p = 0.02; Supplementary Figure 1b, right panel). We further quantified the volume of CtBP2 puncta and GluA2 receptor patches in these colocalized synaptic markers (Supplementary Figure 1c, d). In the apical region, both α9KO and α9KI mice showed reduced ribbon volumes compared to WT (one way ANOVA followed by Holm-Sidak test, df = 2, p < 0.0001 for both groups; Supplementary Figure 1c, left panel). In the middle region, the reduction was significant only in α9KI mice compared to WT (one way ANOVA followed by Holm-Sidak test, df = 2, p < 0.0001; Supplementary Figure 1c, middle panel). Interestingly, in the basal region, α9KO mice had increased CtBP2 volumes, while this was smaller in α9KI mice (one way ANOVA followed by Holm-Sidak test, df = 2, p < 0.0001 for both groups; Supplementary Figure 1c, right panel). The size of the GluA2 receptor patches was significantly smaller in the apical region of α9KI mice, with no changes in α9KO mice compared to WT (one way ANOVA followed by Holm-Sidak test, df = 2, p < 0.0001; Supplementary Figure 1d, left panel). In the middle region, both α9KO and α9KI mice had smaller GluA2 volumes (one way ANOVA followed by Holm-Sidak test, df = 2, p < 0.0001, for both groups; Supplementary Figure 1d, middle panel). In the basal region, α9KO mice had larger GluA2 puncta compared to WT (one way ANOVA followed by Holm-Sidak

test, df = 2, p < 0.0001; Supplementary Figure 1d, right panel). These results suggest that

inhibitory input from cholinergic efferent fibers plays a key role in shaping synaptic density and

maintaining the long-term morphology of pre- and postsynaptic elements necessary for optimal

auditory function.

The long-term impact of early-age noise exposure on auditory nerve synapse numbers is

illustrated in Figure 6, on cochleae dissected and fixed at P75 from control and noise-exposed

mice with varying levels of α9α10 nAChR activity. Presynaptic, postsynaptic, and colocalized

synaptic puncta quantifications are shown for the apical, middle, and basal cochlear regions

(Figure 6, violin plots, right panels). In WT exposed ears, the number of CtBP2, GluA2, and

colocalized synaptic puncta significantly decreased at both the apical and basal cochlear ends

(Figure 6a). The decrease in CtBP2 puncta was more pronounced in the high-frequency basal

region, with a 37.2% decrease compared to controls (Mann-Whitney, df = 1, p < 0.0001 at the

apical and basal regions). Similarly, GluA2 postsynaptic receptors were reduced in both the apical

and basal turns in the cochlea of noise-exposed WT mice (Mann-Whitney, df = 1, p < 0.0001 at

both cochlear regions). Putative ribbon synapse counts, defined by juxtaposed CtBP2- and

GluA2-positive puncta, showed a 34.21% reduction in the apical and a 32.56% reduction in the

basal turn following acoustic trauma (Mann-Whitney, df = 1, p < 0.0001 at both cochlear regions)

(Figure 6a, right panel). In α9KO mice, there was also a reduction in the number of presynaptic,

postsynaptic, and colocalized puncta after noise exposure, depending on cochlear

frequency/location (Figure 6b). Ribbon puncta in α9KO exposed ears were reduced by up to

14.73% of the control in the apical and 24.75% in the basal cochlear ends after noise exposure

(Mann-Whitney, df = 1, p = 0.001 and p = 0.002 at the apical and basal region, respectively).

GluA2-positive postsynaptic receptor patches decreased by 24.79% in the basal turn of noise-

exposed mice (Mann-Whitney, df = 1, p = 0.004). Following noise exposure, synaptic puncta were

reduced by 26.17% in the apical cochlear region and 25.87% in the basal region (Mann-Whitney,

df = 1, p = 0.002 and p < 0.0001 at the apical and basal region, respectively) (Figure 6b, right

panel). Interestingly, after two months post-exposure, α9KI ears exhibited no significant changes

in presynaptic ribbons, postsynaptic GluA2 receptor patches, or putative synapses across all

cochlear regions (Figure 6c). Although not statistically significant, a modest increase in synaptic

density (~19% at the basal region) was observed in α9KI tissue (Figure 6c, right panel).

Finally, we quantified the volume of ribbons and AMPA receptor patches (i.e., colocalized

CtBP2 and GluA2 puncta) in unexposed and exposed mice with different levels of MOC feedback

(Figure 7). In WT mice, noise exposure resulted in significantly larger ribbon volumes in the basal

end of the cochlea compared to unexposed mice, with no corresponding changes in AMPA

receptor patch size (t-test, df = 1, p = 0.01; Figure 7a). No changes were observed in the apical

or middle cochlear regions (Figure 7a). Conversely, α9KO mice showed reduced volumes of both

ribbons and AMPA receptor patches in the basal region following noise exposure, with no

changes in the apical or middle regions (t-test, df = 1, p = 0.002 for CtbP2 volume and p = 0.003

for GluA2 volume; Figure 7b). Notably, α9KI mice showed no morphological changes, suggesting

that enhanced α9α10 nAChR activity protects synapses from noise-induced alterations (Figure

7c).

DISCUSSION

Auditory development is a gradual process that begins during embryonic stages and

continues through sensitive postnatal periods. While much is known about the detrimental effects

of noise exposure at mature stages, less is understood about its impact during this crucial

developmental window and the role of the MOC system in this context. Here, we explored how

varying levels of MOC inhibitory modulation affect the developmental consequences of early-age

noise exposure. Our findings reveal that changes in the strength of MOC inhibition impact the

timing of hearing onset and are crucial for the long-term maintenance of IHC ribbon synapses.

Moreover, loud noise exposure during postnatal developmental stages caused long-lasting

changes in auditory sensitivity in WT, highlighting the greater vulnerability of the auditory system

to noise-induced injury during early development. Notably, enhanced cholinergic activity in α9KI

mice conferred protection from early developmental noise-induced threshold shifts and cochlear

synaptopathy. This underscores the crucial protective role of the MOC system during these

vulnerable early developmental stages.

Before hearing begins, MOC synapses modulate IHCs' activity, shaping the

developmental increase in Ca^{2+} sensitivity of glutamate release at IHC ribbon synapses

[16,24,27,59]. Previous work in genetically modified mice and studies involving neonatal

transection of the olivocochlear bundle have shown that MOC modulation at these early ages is

essential for cochlear maturation, sensory map formation, and synaptic refinement along the

auditory pathway [22,23,27,29,60–62]. Our findings further demonstrate that α9α10 nAChR

activity regulates hearing onset and synaptic integrity. Enhanced α9α10 nAChR activity in α9KI

mice accelerates the onset of hearing, with sound-evoked potentials detected earlier than in mice

with normal MOC synapses. In the absence of MOC activity, as in α9KO mice, hearing onset is

delayed, with higher ABR thresholds and diminished wave 1 amplitudes compared to WT during

early development. Consistent with these findings, recent research on α9KO mice has shown that

auditory sensitivity at hearing onset is impaired in the absence of pre-hearing efferent input [28].

In addition, our work shows that both enhanced and absent MOC inhibition interfere with the

proper formation and maintenance of ribbon synapses in adult IHCs. At P75, both α9KO and α9KI

mice exhibit significant reductions in ribbon synapse density and alterations in the volume of pre-

and post-synaptic puncta compared to age-matched WT controls, underscoring the critical role of

balanced MOC activity in auditory system development and synaptic organization. In α9KI mice,

ribbon and GluA2 patch volumes were reduced, while α9KO mice showed region-specific effects:

decreased ribbon volume in the apical region without changes in GluA2 receptor size and

increased synaptic component size in the basal region. Previous studies have emphasized the

functional importance of ribbon size, showing that it is tightly regulated by presynaptic Ca^{2+}

channels, with alterations in Ca^{2+} influx impacting synaptic plasticity, neurotransmitter release,

and active zone maturation [62–65]. During early development, IHC active zones undergo

substantial structural and functional maturation, including the removal of extrasynaptic Ca^{2+}

channels and the co-localization of glutamate receptor patches with presynaptic ribbons [66–68].

This maturation refines presynaptic morphology, reduces the number of ribbons per synapse, and

establishes monosynaptic connections between IHCs and auditory nerve fibers [69–72]. Recent

work also suggests that fusion of synaptically anchored ribbons during this period is critical for

the development of mature active zones and sustained neurotransmitter release [66]. Given our

observation of reduced ribbon size in some cochlear regions of α9KI and α9KO mice, we propose

that altered MOC modulation strength may disrupt intracellular Ca^{2+} dynamics, which could impair

ribbon precursor fusion, thereby impairing active zone maturation. These findings highlight the

essential role of MOC synapses in shaping early auditory sensitivity, reinforcing previous

suggestions [27] that effective coordination between afferent and efferent inputs is important for

regulating spiking activity in the developing auditory pathway. This coordination is crucial for

proper synaptic and auditory system maturation.

Auditory function in vertebrates matures through a well-defined process, transforming the

system from high thresholds and limited frequency discrimination at the onset of hearing to a

highly sensitive and precise mechanism capable of resolving frequency differences as small as 1

Hz [73]. Our findings suggest that the WT mouse strain used in the present work aligns with this

established maturation pattern, as auditory function recordings from P12 to P75 reveal a clear

progression from an immature to a mature state. At the onset of hearing (P14 in our strain of

mice), the auditory system shows elevated ABR and DPOAE thresholds, reduced wave 1

amplitudes, and poorly defined later ABR waves, reflecting incomplete functional maturation. Over

the following days, thresholds progressively decreased, reaching mature levels by P22. This

maturation is accompanied by a steady increase in ABR wave 1 amplitudes from P15 to P22,

underscoring the postnatal refinement period necessary for achieving full auditory sensitivity.

Thus, by P15, the time when mice were exposed to high levels of noise, the maturation process

is not yet complete. A key finding of this study is that exposure to intense sound during early

developmental stages results in significantly more severe consequences compared to exposure

at a later age [44]. In WT mice, exposure to 100 dB SPL noise for 1 hour at P15 resulted in lasting

ABR threshold elevations for up to two months, accompanied by a 35% loss of ribbon synapses.

In contrast, the same exposure at P21 causes only transient threshold shifts, which returns to

baseline within seven days, with a less pronounced degree of cochlear synaptopathy [44]. This

stark difference underscores the heightened vulnerability of the auditory system during early

development and highlights the age-dependent impact of noise exposure. Previous reports have

also shown that the same scheme of intense noise exposure causes more severe damage in

juvenile animals at 4 weeks of age compared to adult animals at 16 weeks of age [74–76].

Although the underlying mechanisms that drive this heightened sensitivity are not known, we can

speculate that this increased sensitivity reflects a critical window in which MOC synaptic activity

is still maturing or being functionally reinforced. Alternatively, central mechanisms—such as

refinement of MOC neurons, completion of myelination, or strengthening of afferent drive—may

contribute to the vulnerability observed at this stage. Ongoing maturation of the auditory nerve–

lateral olivocochlear (LOC) complex may also play a role, potentially rendering the system more

susceptible to disruption by acoustic overexposure during this period. DPOAE thresholds were

also elevated seven days post-exposure in WT mice, although these thresholds recovered after

two months, suggesting that there is no loss of OHCs or alteration in their function. Given this

recovery of DPOAE thresholds but persistently elevated ABR thresholds, we can infer that the

elevation is not due to damage in the cochlear amplifier but rather to a deficiency in the afferent

pathway of sound signal transmission. In line with this, the increase in ABR thresholds was

accompanied by a reduction in wave 1 amplitudes, which also persisted for two months after noise

exposure. This study represents the first report of sustained ABR threshold elevation for such an

extended period following moderate noise exposure at 100 dB SPL for 1 hour. Similar results

were obtained in 14-day-old rats exposed to a higher noise level of 125 dB SPL for 25 minutes,

where after two months post-exposure, elevated thresholds at high frequencies and a loss of

presynaptic elements (CTBP2/Ribeye) were observed [77]. Our findings demonstrate a significant

loss of ribbon synapses in the apical and basal regions of the cochlea in WT mice following noise

exposure, which correlates with decreased ABR wave 1 amplitudes. In addition to cochlear

synaptopathy, we observed that IHCs of noise-exposed mice tend to have larger ribbons

(Ribeye/CtBP2 volume) compared to unexposed IHCs in the basal region of the cochlea. A similar

increase in the volume of presynaptic elements following acoustic trauma was observed using

both the RIBEYE-tagRFP mouse model [78] and volume electron microscopy [76]. The observed

ribbon enlargement in WT mice may be attributed to the fusion of individual ribbons after noise

exposure. In fact, recent evidence indicates an increase in multi-ribbon active zones as well as

movement suggesting unanchoring from synaptic tethers in synaptopathic ears [76,78,79]. While

the functional significance of this structural adaptation remains unclear, multi-ribbon active zones

persist into adulthood [66,67] and are a hallmark of cochlear aging [80], with larger ribbons being

associated with increased Ca^{2+}-dependent exocytotic activity [81]. Notably, noise exposure has

also been associated with enhanced exocytosis in IHCs, along with an increase in tone-burst

responses in auditory nerve fibers [82,83]. A recent study also demonstrated that intense noise

exposure (98 dB SPL for two hours) in young animals reduces the number of presynaptic ribbons

in OHCs, while increasing the volume and mobility of those that remain [84]. We can hypothesize

that ribbon volume enlargement may act as a compensatory mechanism to maintain auditory

sensitivity after synapse loss. Importantly, we did not detect changes in the size of GluA2 receptor

patches in WT mice post-noise exposure.

In α9KO mice, which lack α9α10 nAChR activity, elevated ABR and DPOAE thresholds

were observed one week after noise exposure at P15, consistent with findings from noise-

exposed α9KO mice at P21 [44]. However, in contrast to WT animals, ABR thresholds in α9KO

mice nearly fully recovered to unexposed control levels two months post-exposure. While DPOAE

thresholds returned to control levels at 8 and 22.65 kHz, they remained elevated at 16 kHz after

two months. This elevation in DPOAE thresholds at 16 kHz was accompanied by a slight, though

not statistically significant, elevation in ABR thresholds at the same frequency. The recovery of

ABR thresholds in α9KO mice, compared to the permanent threshold elevation seen in WT mice,

may be attributed to the reduced auditory sensitivity at hearing onset in α9KO mice due to the

absence of prehearing efferent modulation. While ABR thresholds return to pre-exposure values,

a persistent reduction in wave 1 amplitudes and ribbon synapse counts per IHC was observed.

Prior studies suggest that suprathreshold response deficits following noise exposure are driven

by the selective loss of high-threshold, low-spontaneous-rate auditory nerve fibers [85].

Accordingly, the lack of threshold elevation despite synaptic loss may reflect the preferential

vulnerability of this specific neuronal subpopulation. However, in α9KO ears exposed to noise,

both pre- and postsynaptic puncta showed a reduction in size, indicating noise may also involve

maladaptive alterations in both pre- and postsynaptic structures.

In contrast to the findings in WT and α9KO mice, α9KI mice, which exhibit enhanced

α9α10 nAChR activity, showed no ABR or DPOAE threshold elevations following sound exposure.

Additionally, no evidence of cochlear synaptopathy was detected in these mice. This phenomenon

had already been described in adult mice (P21) exposed to the same level of noise [44]. At birth,

rodent MOC neurons initially innervate IHCs, and by P10, they expand to innervate both IHCs

and OHCs. After the onset of hearing, their innervation becomes confined to OHCs, gradually

retracting from IHCs [30]. Our results therefore suggest that even with early developmental noise

exposure—when MOC innervation to OHCs is still developing—enhanced α9α10 nAChR

responses can effectively exert a protective effect on OHCs against functional and structural

damage associated with acoustic trauma. Importantly, our findings also reveal that mice with

enhanced α9α10 nAChR activity showed a modest increase in ribbon synapse density following

acoustic overexposure. This is consistent with previous work, which documented a more

substantial increase in synaptic density in α9KI mice exposed to the same noise protocol at P21

[44]. These results indicate that enhancement of the cholinergic component of the MOC system

may support synapse formation following noise exposure.

In summary, this work highlights three key findings: 1) the MOC system plays a crucial

role in regulating the development of ribbon synapse density and size, which is essential for

proper hearing onset; 2) exposure to loud noise during early development results in more severe

auditory consequences compared to the same exposure later in life; and 3) enhancing α9α10

nAChR responses during early development protects against noise-induced damage,

underscoring the importance of timely modulation of this receptor post-hearing onset to mitigate

developmental impairments.

Clinical implications:

Clinical evidence indicates that early acoustic exposure can adversely affect auditory

development, a concern particularly evident in neonates requiring prolonged hospitalization.

Advances in newborn hearing screening have revealed a higher-than-anticipated prevalence of

auditory neuropathy, especially among infants treated in neonatal intensive care units (NICUs),

where persistent high noise levels are common [86,87]. Our experimental findings align with these

clinical observations, showing that early-age noise exposure exacerbates auditory deficits in mice.

Importantly, enhancement of α9α10 nAChR signaling during early development attenuated noise-

induced damage, underscoring the receptor's protective role during a sensitive window of auditory

system maturation. These results offer proof-of-principle for the potential of pharmacological

strategies targeting α9α10 nAChRs to prevent developmental hearing loss associated with early

acoustic stress.

Figure 1: Auditory function in WT, α9KO, and α9KI mice across different postnatal ages.
(**a**) Representative ABR waveforms from WT (black), α9KI (red), and α9KO (blue) mice recorded at P14 in response to 16 kHz tone bursts at increasing sound pressure levels (dB SPL). Thresholds, defined by the presence of wave 1, are marked by green dashed traces. Waves 1, 2 and 3 are indicated by gray arrows. The scale bar applies to all three recordings. (**b**) Mean ABR thresholds from P12 to P75 in WT (n = 12), α9KO (n = 14), and α9KI (n = 14) mice at 11.33, 16, and 22.65 kHz. ABR waveforms first became detectable at P13 in α9KI mice, at P14 in WT mice, and between P14 and P15 in α9KO mice. (**c**) Mean DPOAE thresholds from P12 to P75 in WT (n = 12), α9KO (n = 14) and α9KI (n = 14) mice at 11.33, 16 and 22.65 kHz. DPOAE thresholds showed the same trend as ABR only at 11.33 kHz. Group means ± SEM are shown. Red asterisks represent the statistical significance of α9KI compared with WT, and blue asterisks represent the statistical significance of α9KO compared with WT (Kruskal-Wallis followed by post hoc Dunn test, *p < 0.05).

Figure 2: Suprathreshold ABR wave1 amplitudes and latencies in WT, α9KO, and α9KI mice during postnatal development. (**a**) Representative examples of ABR waveforms recorded at 16 kHz, 80 dB SPL, from WT (black), α9KI (red), and α9KO (blue) mice. Waves 1, 2 and 3 are indicated by gray arrows. Each column illustrates waveform maturation across developmental

stages (P12–P16, P22, P75) for each genotype. At P13, α9KI mice exhibit measurable responses, while distinct ABR waves emerge in WT and α9KO mice by P14. Waveform morphology stabilizes by P22 and remains consistent through P75. Scale bars apply to all traces. (**b**) ABR wave 1 amplitudes (µV) at 80 dB SPL for 11.33, 16, and 22.65 kHz across developmental ages in WT, α9KO, and α9KI mice. Amplitudes were measured as the difference between the maximum and minimum peaks. No values are shown for P12 due to the absence of discernible waves. (**c**) ABR wave 1 latencies (ms) at the same frequencies. A developmental decrease in latency is observed between P14 and P15 in all genotypes, though not statistically significant. Data represent mean ± SEM. Red asterisks denote significant differences between α9KI and WT; blue asterisks indicate significant differences between α9KO and WT (one-way ANOVA followed by Holm-Sidak test, *p < 0.05; **p < 0.01).

Figure 3: ABR wave 1 threshold measurements in control and noise-exposed WT, α9KO, and α9KI mice at different postnatal ages. ABR thresholds before and after exposure to noise at P15 in WT ($n_{control}$ = 14 and $n_{Exposed}$ = 12) (**a**), α9KO ($n_{control}$ = 11 and $n_{Exposed}$ = 11) (**b**), and α9KI ($n_{control}$ = 12 and $n_{Exposed}$ = 12) (**c**) mice at 8, 16 and 22.65 kHz. WT mice showed a significant increase in ABR thresholds 1 week and 2 months after exposure to noise, indicating a permanent threshold elevation. A recovery of ABR thresholds was observed in α9KO mice 2 months after exposure to noise. α9KI mice did not present any changes in ABR thresholds at any time after acoustic trauma. Each point represents the mean ± SEM. Asterisks represent the statistical significance (Mann-Whitney test, *p < 0.05; **p < 0.01).

Figure 4: Suprathreshold response amplitude for ABR wave 1 after P15 noise exposure in WT, α9KO, and α9KI mice at different postnatal ages compared to unexposed controls. ABR wave 1 amplitudes (in µV) at 80 dB SPL at 8, 16 and 22.65 kHz in WT ($n_{Control}$ = 14 and $n_{Exposed}$ = 12) (**a**), α9KO ($n_{Control}$ = 11 and $n_{Exposed}$ = 11) (**b**), and α9KI ($n_{Control}$ = 12 and $n_{Exposed}$ = 12) (**c**) mice at the same time points shown in Figure 3. Each panel inset displays representative ABR waveforms recorded at 80 dB SPL for the corresponding frequency, shown for all groups at P75. Scale bars (1 µV vertical, 1 ms horizontal) apply to all traces. WT and α9KO mice exhibited reduced ABR wave 1 amplitudes following noise exposure, whereas wave 1 amplitudes in α9KI mice remained unaffected. Data are presented as mean ± SEM. Statistical significance is indicated by asterisks (Mann-Whitney test, *p < 0.05; **p < 0.01).

Figure 5: **Assessment of OHC functional integrity at different postnatal ages following noise exposure at P15 in WT, α9KO, and α9KI mice.** DPOAE thresholds are shown for WT ($n_{Control}$ = 14 and $n_{Exposed}$ = 12) (**a**), α9KO ($n_{Control}$ = 11 and $n_{Exposed}$ = 11) (**b**), and α9KI ($n_{Control}$ = 12 and $n_{Exposed}$ = 12) (**c**) at 8, 16 and 22.65 kHz. WT mice exhibited a transient increase in DPOAE thresholds 1 week post-exposure at 16 kHz. α9KO mice showed increased thresholds at 8 and 16 kHz 1 week post-exposure, with sustained elevation at 16 kHz up to 2 months. α9KI mice demonstrated no changes in DPOAE thresholds after noise exposure. Data are presented as mean ± SEM. Statistical significance is indicated by asterisks (Mann-Whitney test, *p < 0.05; **p < 0.01).

Figure 6: Analysis of the degree of IHC synaptopathy two months after exposure to noise at P15. Left, Representative confocal images of IHC synapses at the apex of the cochlea from cochleae immunolabeled for presynaptic ribbons (CtBP2-red), postsynaptic receptor patches (GluA2-green) and IHCs (Myosin VIIa-blue). CtBP2 antibody also weakly stains IHC nuclei. Scale bar, 10μm. Right, Quantitative data obtained from WT, α9KO, and α9KI mice at P75. For each IHC, we analyzed the number of CtBP2 puncta, postsynaptic GluA2 receptor patches, and putative ribbon synapses. (**a**) In traumatized WT mice, there was a reduction in the number of CtBP2 puncta, GluA2 receptor patches, and putative synapse, at the apical and basal region of the cochlea (WT Control: n_{Apex} = 80 IHC, n_{Middle} = 75 IHC, n_{Base}= 60 IHC; WT Exposed: n_{Apex} = 82 IHC, n_{Middle} = 90 IHC, n_{Base} = 50 IHC). (**b**) In traumatized α9KO mice, there was a reduction in the number of prelocalized, postlocalized, and colocalized puncta depending on cochlear frequency/location (α9KO Control: n_{Apex} = 100 IHC, n_{Middle} = 84 IHC, n_{Base} = 118 IHC; α9KO Exposed: n_{Apex} = 234 IHC, n_{Middle} = 60 IHC, n_{Base} = 90 IHC). (**c**) In traumatized α9KI mice, no

588 difference was found in the number of presynaptic ribbons, postsynaptic AMPA receptors, and
589 colocalization puncta for the three regions of the cochlea (α9KI Control: n_{Apex} = 178 IHC, n_{Middle} =
590 319 IHC, n_{Base} = 198 IHC; α9KI Exposed: n_{Apex} = 196 IHC, n_{Middle} = 145 IHC, n_{Base} = 80 IHC). Violin
591 plots indicate median (solid line) and IQR (dashed lines). Asterisks denote the statistical
592 significance (Mann Whitney test, **p < 0.01; ***p < 0.001, ****p < 0.0001).

Figure 7: Volumes of presynaptic ribbons and postsynaptic receptor patches for each synaptic pair from control and noise-exposed WT, α9KO and α9KI mice. Violin plots comparing the volumes of fluorescence staining for ribbons (CtBP2 size) and AMPA receptors (GluA2 size) between control and noise-exposed mice. The quantification was made in the apex, middle and base regions of the cochlea from WT (**a**) (Control: n_{Apex} = 450 synaptic pairs, n_{Middle} = 509 synaptic pairs, n_{Base} = 203 synaptic pairs; Exposed: n_{Apex} = 497 synaptic pairs, n_{Middle} = 175 synaptic pairs, n_{Base} = 250 synaptic pairs); α9KO (**b**) (Control: n_{Apex} = 367 synaptic pairs, n_{Middle} = 187 synaptic pairs, n_{Base} = 221 synaptic pairs; Exposed: n_{Apex} = 207 synaptic pairs, n_{Middle} = 300 synaptic pairs, n_{Base} = 145 synaptic pairs) and α9KI (**c**) (Control: n_{Apex} = 247 synaptic pairs, n_{Middle} = 270 synaptic pairs, n_{Base} = 240 synaptic pairs; Exposed: n_{Apex} = 280 synaptic pairs, n_{Middle} = 310 synaptic pairs, n_{Base} = 360 synaptic pairs) mice. Violin plots indicate median (solid line) and IQR (dashed lines). Asterisks denote the statistical significance (t-test; *p < 0.05; ***p < 0.001).

SUPPLEMENTARY INFORMATION

Supplementary Figure 1. Analysis of IHC ribbon synapses in unexposed WT, α9KO, and α9KI mice. (**a**) Representative confocal images of IHC synapses at the apex of the cochlea from

cochleae immunolabeled for presynaptic ribbons (CtBP2-red) and postsynaptic receptor patches (GluA2-green) for WT (left), α9KO (middle) and α9KI (right). CtBP2 antibody also weakly stains IHC nuclei. Scale bar, 7 µm. (**b**) Number of putative ribbon synapses per IHC (i.e., colocalized CtBP2 and GluA2 puncta). WT: n_{Apex} = 80 IHC, n_{Middle} = 75 IHC, n_{Base} = 60 IHC from 14 animals; α9KO: n_{Apex} = 100 IHC, n_{Middle} = 84 IHC, n_{Base} = 118 IHC from 11 animals and α9KI: n_{Apex} = 178 IHC, n_{Middle} = 319 IHC, n_{Base} = 198 IHC from 12 animals. (**c**) Volumes of fluorescence staining for ribbons (CtBP2 size) and (**d**) AMPA receptors (GluA2 size) for each of the synaptic pairs (WT: n_{Apex} = 450 synaptic pairs, n_{Middle} = 509 synaptic pairs, n_{Base} = 203 synaptic pairs; α9KO: n_{Apex} = 367 synaptic pairs, n_{Middle} = 187 synaptic pairs, n_{Base} = 221 synaptic pairs and α9KI: n_{Apex} = 247 synaptic pairs, n_{Middle} = 270 synaptic pairs, n_{Base} = 240 synaptic pairs). Violin plots indicate median (solid line) and interquartile range (IQR, dashed lines). Asterisks denote the statistical significance (Kruskal–Wallis followed by a Dunn's post-test was used to analyze the number of synapses and one way ANOVA followed by Holm-Sidak test; *p < 0.05; **p < 0.01; ***p < 0.001, ****p < 0.0001).

METHOD DETAILS

Animals. α9KO and α9KI mice have been previously described [42,45] and were backcrossed with the congenic FVB.129P2-*Pde6bþ Tyrc-ch/AntJ* strain (https://www.jax.org/strain/004828) for seventeen generations (i.e., N-17). We maintained a consistent male-to-female ratio across all experimental groups and genotypes. No sex-related differences were observed in the data. This study included adult and pup animals aged P12 to P16, which were housed in a sound-controlled room within the animal care facility. Background sound levels in this environment were consistently below 50 dB SPL. Noise levels were measured using an electret condenser microphone (FG-23329-PO7; Knowles). Transient sounds during routine cage cleaning and daily maintenance by animal care staff were minimal, with peaks not exceeding 70 dB SPL. The methods reported in this study adhere to the ARRIVE guidelines for reporting *in vivo* animal research (https://arriveguidelines.org/), ensuring transparency and reproducibility. All procedures were conducted in accordance with institutional and ethical guidelines for animal care and use.

Auditory function tests. Physiological assessments of cochlear and central function, including auditory brainstem responses (ABRs) and distortion-product otoacoustic emissions (DPOAEs), were performed in mice anesthetized with xylazine (10 mg/kg, i.p.) and ketamine (100 mg/kg, i.p.) and placed in a soundproof chamber maintained at 32°C to counteract the drop in body temperature caused by anesthesia. ABR represents sound-evoked potentials generated by neural circuits in the ascending auditory pathways, while DPOAEs are sounds emitted by the cochlea, specifically by OHCs, in response to two pure-tone stimuli presented simultaneously. Sound stimuli were delivered through a custom acoustic system with two dynamic earphones used as sound sources (CDMG15008–03A; CUI) and an electret condenser microphone (FG-23329-PO7; Knowles) coupled to a probe tube to measure sound pressure near the eardrum. Data collection was conducted using a National Instruments PXI-based system equipped with 24-

bit input/output boards and operated through a custom LabView software interface. For ABRs,

platinum needle electrodes were placed into the skin at the dorsal midline close to the neural crest

and pinna with a ground electrode near the tail. ABR potentials were evoked with 5 ms tone pips

(0.5 ms rise-fall, with a $\cos^2$ envelope, at 40/s) delivered to the eardrum at log-spaced frequencies

from 5.6 to 45.25 kHz. The response was amplified 10,000X with a 0.3–3 kHz passband. Sound

level was raised in 5 dB steps from 20 to 80 dB sound pressure level (SPL). At each level, 1024

responses were averaged with alternated stimulus polarity. Threshold for ABR was defined as

the lowest stimulus level (<80 dB SPL) at which a repeatable wave 1 could be identified in the

response waveform. The ABR waves 1, 2 and 3 amplitudes were computed by off-line analysis

of the peak-to-peak amplitude of stored waveforms. The DPOAEs in response to two primary

tones of frequency f1 and f2 were recorded at 2f1-f2, with f2/f1=1.2, and the f2 level 10 dB lower

than the f1 level. Ear-canal sound pressure was amplified and digitally sampled at 4 µs intervals.

DPOAE threshold was defined as the lowest f2 level in which the signal to noise floor ratio is >1.

Noise exposure. Animals were exposed under anesthesia at P15 to a 1–16 kHz noise at 100 dB

SPL for 1 hour using closed-field earphones (CDMG15008–03A; CUI) as the sound source. The

exposure took place in the same acoustic chamber used for cochlear function tests. The noise

consisted of a flat-spectrum broadband signal calibrated to the target SPL immediately before

each exposure session.

Animal monitoring and health assessment. To assess the impact of repeated anesthesia on

the health and well-being of the animals, their weight and motor activity were monitored daily

throughout the auditory function recordings and noise exposure protocols. Animals were observed

during their recovery from anesthesia to ensure no adverse effects on their general condition.

Growth curves showed no significant differences in weight between animals that were

anesthetized and untreated controls animals, suggesting that anesthesia did not affect normal

growth. Furthermore, no behavioral changes or motor impairments were observed in the

anesthetized animals when compared to controls, indicating that repeated anesthesia had no

detectable impact on overall health or behavior.

Cochlear processing and immunostaining. Cochleae were perfused intralabyrinthly with 4%

paraformaldehyde (PFA) in phosphate-buffered saline (PBS), post-fixed with 4% PFA overnight,

and decalcified in 0.12 M EDTA. Cochlear tissues were microdissected and permeabilized by

freeze/thaw cycles in 30% sucrose. The dissected samples were then separated into distinct wells

based on cochlear region: apex, middle and base. Tissue samples were blocked with 5% normal

goat serum and 1% Triton X-100 in PBS for 1 h, followed by incubation in primary antibodies

(diluted in blocking buffer) at 37°C for 16 h. The primary antibodies used in this study were: 1)

anti-C-terminal binding protein 2 (mouse anti-CtBP2 IgG1; BD Biosciences, San Jose, CA;

Cat#612044, RRID:AB_399431, 1:200) to label the presynaptic ribbon, 2) anti-glutamate receptor

2 (mouse anti-GluA2 IgG2a; Millipore, Billerica, MA; Cat#MAB397, RRID:AB_11212990, 1:2000)

to label the post-synaptic receptor plaques and 3) anti-Myosin 7a (rabbit anti-MyosinVIIa, Proteus

Biosciences; Cat# 25-6790, RRID:AB_10015251, 1:200) to label the cochlear hair cells. Tissues

were then incubated with the appropriate Alexa Fluor-conjugated fluorescent secondary

antibodies (Invitrogen, Carlsbad, CA; 1:500 in blocking buffer) for 2 hours at room temperature.

Finally, tissues were mounted on microscope slides in FluorSave mounting media (Millipore,

Billerica, MA).

Confocal microscopy and image processing. For IHC synaptic counts, cochlear tissue from

the apex, middle and base regions was first imaged at low magnification to correlate each piece

with its corresponding frequency region [88]. The apical region corresponds to 4–12 kHz, the

middle region to 12–20 kHz, and the base region to 22–40 kHz. Z-stacks images were then

acquired using a Leica TCS SPE microscope with a 63X oil-immersion lens (1.5X or 4X digital

zoom). Imaging settings were consistent across all samples, ensuring no pixel saturation. For

each stack, the Z-step was set to 0.3 µm, with a pixel size of 0.11 µm in both the X and Y axes.

Each stack typically contained 10 to 20 IHCs. Image stacks were processed using Fiji software

(RRID:SCR_002285) [89], and a custom plugin was developed to automate the counting of

synaptic ribbons, glutamate receptor patches and co-localized synaptic puncta. The algorithm's

accuracy was validated by comparing automated counts to manual counts, ensuring reliable

quantification. Briefly, each channel was analyzed separately, and maximum projections were

generated to quantify the number of CtBP2 or GluA2 puncta. Additionally, a composite between

the three channels was produced to draw the different regions of interest (ROI) that correspond

to each IHC taking the myosin staining as a reference. The maximum projections from the single

channels were multiplied to generate a merged 32-bit image. Then, they were converted to binary

images after a custom thresholding procedure. An automatic counting of the number of particles

on each ROI was performed.

For quantitative volumetric synaptic analysis, different regions from the whole cochlea were

imaged and confocal z-stacks were processed in Fiji. z-stacks were deconvolved applying 10

iterations of the Richardson-Lucy algorithm in the DeconvolutionLab2 plugin [90], an algorithm

that has been used previously in cochlear whole mounts [91]. Then, the 3D Object Counter plugin

was used to perform three-dimensional image segmentation and object individualization of the

deconvoluted stacks. According to previous work [58,92,93] we reasoned that a synapse would

occupy a volume between 0.04 µm3 to 5 µm3, and then we set a size range of 20-2000 voxels,

with a voxel size of 0.11 µm x 0.11 µm x 0.2 µm. The volumes of presynaptic ribbons (CtBP2

fluorescence) and postsynaptic AMPA receptor clusters (GluA2 fluorescence) at IHC-afferent

synapses were measured using the 3D ROI Manager [94]. All exported volumes were checked to

ensure that volumes were within instrument resolution limits after threshold adjustments.

Statistical analysis

Data was analyzed using R Statistical Software (RRID:SCR_001905). Shapiro-Wilks test was

used for testing normal distribution of the residuals. In cases where the data did not follow a

normal distribution, the following tests were used: Kruskal-Wallis test followed by multiple

contrasts according to Dunn's method for comparisons between three groups (IHCs synaptic

counts in the control condition, ABR and DPOAE thresholds during development) and Mann-

Whitney U test for comparisons between two groups (ABR thresholds, DPOAE thresholds, and

IHCs synaptic counts). If data were normally distributed, a one-way ANOVA was conducted,

followed by Holm-Sidak post hoc tests for three-group comparisons (wave 1 amplitudes and

synaptic volumes in the control condition). For two-group comparisons, a t-test was used (synaptic

volumes). Statistical significance was set to $p < 0.05$.

REFERENCES

1.   Sanes DH, Walsh EJ. The development of central auditory processing. In: Rubel EW, Popper AN,
Fay RR, editors. Development of the Auditory System. New York: Springer; 1998. pp. 271–314.

2.   Sanes DH, Bao S. Tuning up the developing auditory CNS. Curr Opin Neurobiol. 2009;19: 188–199.
Available:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list
_uids=19535241

3.   Chang EF, Merzenich MM. Environmental noise retards auditory cortical development. Science.
2003;300: 498–502. doi:10.1126/science.1082163

4.   Keuroghlian AS, Knudsen EI. Adaptive auditory plasticity in developing and adult animals. Prog
Neurobiol. 2007;82: 109–121. doi:10.1016/j.pneurobio.2007.03.005

5.   Zhou X, Merzenich MM. Enduring effects of early structured noise exposure on temporal
modulation in the primary auditory cortex. Proc Natl Acad Sci U S A. 2008;105: 4423–4428.
doi:10.1073/pnas.0800009105

6.   Lauer AM, May BJ, Ay BRJM, Lauer AM, May BJ. The medial olivocochlear system attenuates the
753   developmental impact of early noise exposure. JARO - J Assoc Res Otolaryngol. 2011;12: 329–
754   343. doi:10.1007/s10162-011-0262-7

755   7.   Caminos C, Vale E, Lujan R, Martinez-Galan JR, Juiz JM. Developmental regulation and adult
756   maintenance of potassium channel proteins (Kv 1.1 and Kv 1.2) in the cochlear nucleus of the rat.
757   Brain Res. 2005;1056. doi:10.1016/j.brainres.2005.07.031

758   8.   Waguespack J, Salles FT, Kachar B, Ricci AJ. Stepwise morphological and functional maturation of
759   mechanotransduction in rat outer hair cells. J Neurosci Off J Soc Neurosci. 2007;27: 13890–
760   13902. doi:10.1523/JNEUROSCI.2159-07.2007

761   9.   Dorrn AL, Yuan K, Barker AJ, Schreiner CE, Froemke RC. Developmental sensory experience
762   balances cortical excitation and inhibition. Nature. 2010;465: 932–936. doi:10.1038/nature09119

763   10.   Sun YJ, Wu GK, Liu B-H, Li P, Zhou M, Xiao Z, et al. Fine-tuning of pre-balanced excitation and
764   inhibition during auditory cortical development. Nature. 2010;465: 927–931.
765   doi:10.1038/nature09079

766   11.   Froemke RC, Jones BJ. Development of auditory cortical synaptic receptive fields. Neurosci
767   Biobehav Rev. 2011;35: 2105–2113. doi:10.1016/j.neubiorev.2011.02.006

768   12.   Lippe WR. Rhythmic spontaneous activity in the developing avian auditory system. J Neurosci.
769   1994;14: 1486–1495. Available:
770   http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list
771   _uids=8126550

772   13.   Marcotti W, Johnson SL, Rusch A, Kros CJ. Sodium and calcium currents shape action potentials in
773   immature mouse inner hair cells. J Physiol. 2003;552: 743–761. Available:
774   http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list
775   _uids=12937295

776   14.   Sonntag M, Englitz B, Kopp-Scheinpflug C, Rübsamen R. Early Postnatal Development of
777   Spontaneous and Acoustically Evoked Discharge Activity of Principal Cells of the Medial Nucleus
778   of the Trapezoid Body: An In Vivo Study in Mice. J Neurosci. 2009;29: 9510–9520.
779   doi:10.1523/JNEUROSCI.1377-09.2009

780   15.   Tritsch NX, Bergles DE. Developmental regulation of spontaneous activity in the Mammalian
781   cochlea. J Neurosci Off J Soc Neurosci. 2010;30: 1539–1550. doi:10.1523/JNEUROSCI.3875-
782   09.2010

783   16.   Johnson SL, Eckrich T, Kuhn S, Zampini V, Franz C, Ranatunga KM, et al. Position-dependent
784   patterning of spontaneous action potentials in immature cochlear inner hair cells. Nat Neurosci.
785   2011;14: 711–717. doi:10.1038/nn.2803

786   17.   De Faveri F, Ceriani F, Marcotti W. In vivo spontaneous Ca2+ activity in the pre-hearing
787   mammalian cochlea. Nat Commun. 2025;16: 29. doi:10.1038/s41467-024-55519-w

18. Tierney TS, Russell FA, Moore DR. Susceptibility of developing cochlear nucleus neurons to
deafferentation-induced death abruptly ends just before the onset of hearing. J Comp Neurol.
1997;378: 295–306. doi:10.1002/(sici)1096-9861(19970210)378:2<295::aid-cne11>3.0.co;2-r

19. Friauf E, Lohmann C. Development of auditory brainstem circuitry. Activity-dependent and
activity-independent processes. Cell Tissue Res. 1999;297: 187–195. doi:10.1007/s004410051346

20. Mostafapour SP, Cochran SL, Del Puerto NM, Rubel EW. Patterns of cell death in mouse
anteroventral cochlear nucleus neurons after unilateral cochlea removal. J Comp Neurol.
2000;426: 561–571. doi:10.1002/1096-9861(20001030)426:4<561::aid-cne5>3.0.co;2-g

21. Leake PA, Hradek GT, Chair L, Snyder RL. Neonatal deafness results in degraded topographic
specificity of auditory nerve projections to the cochlear nucleus in cats. J Comp Neurol. 2006;497:
13–31. doi:10.1002/cne.20968

22. Clause A, Kim G, Sonntag M, Weisz CJC, Vetter DE, Rűbsamen R, et al. The precise temporal
pattern of prehearing spontaneous activity is necessary for tonotopic map refinement. Neuron.
2014;82: 822–835. doi:10.1016/j.neuron.2014.04.001

23. Di Guilmi MN, Boero LE, Castagna VC, Rodríguez-Contreras A, Wedemeyer C, Gómez-Casati ME,
et al. Strengthening of the efferent olivocochlear system leads to synaptic dysfunction and
tonotopy disruption of a central auditory nucleus. J Neurosci. 2019; 2536–18.
doi:10.1523/jneurosci.2536-18.2019

24. Glowatzki E, Fuchs PA. Cholinergic synaptic inhibition of inner hair cells in the neonatal
mammalian cochlea. Science. 2000;288: 2366–68.

25. Marcotti W, Johnson SL, Kros CJ. A transiently expressed SK current sustains and modulates
action potential activity in immature mouse inner hair cells. J Physiol. 2004. Available:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list
_uids=15331671

26. Goutman JD, Fuchs PA, Glowatzki E. Facilitating efferent inhibition of inner hair cells in the
cochlea of the neonatal rat. J Physiol. 2005;566: 49–59. doi:10.1113/jphysiol.2005.087460

27. Johnson SL, Wedemeyer C, Vetter DE, Adachi R, Holley MC, Elgoyhen AB, et al. Cholinergic
efferent synaptic transmission regulates the maturation of auditory hair cell ribbon synapses.
Open Biol. 2013;3. doi:10.1098/rsob.130163

28. Wang Y, Sanghvi M, Gribizis A, Zhang Y, Song L, Morley B, et al. Efferent feedback controls
bilateral auditory spontaneous activity. Nat Commun. 2021;12: 2449. doi:10.1038/s41467-021-
22796-8

29. Walsh E, McGee J, McFadden S, Liberman M. Long-term effects of sectioning the olivocochlear
bundle in neonatal cats. JNeurosci. 1998;18: 3859–3869.

30. Simmons DD, Mansdorf NB, Kim JH. Olivocochlear innervation of inner and outer hair cells during
postnatal maturation: evidence for a waiting period. J Comp Neurol. 1996;370: 551–562.
Available:

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list
_uids=8807454

31.   Vattino LG, Wedemeyer C, Elgoyhen AB, Katz E. Functional Postnatal Maturation of the Medial
Olivocochlear Efferent–Outer Hair Cell Synapse. J Neurosci. 2020;40: 4842–4857.
doi:10.1523/JNEUROSCI.2409-19.2020

32.   Wheeler EF, Bothwell M, Schecterson LC, von Bartheld CS. Expression of BDNF and NT-3 mRNA in
hair cells of the organ of Corti: quantitative analysis in developing rats. Hear Res. 1994;73: 46–56.

33.   Ernfors P, Kucera J, Lee KF, Loring J, Jaenisch R. Studies on the physiological role of brain-derived
neurotrophic factor and neurotrophin-3 in knockout mice. Int J Dev Biol. 1995;39: 799–807.
Available:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list
_uids=8645564

34.   Knipper M, Zimmermann U, Rohbock K, Kopschall I, Zenner HP. Synaptophysin and GAP-43
proteins in efferent fibers of the inner ear during postnatal development. Brain Res Dev Brain
Res. 1995;89: 73–86. Available:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list
_uids=8575095

35.   Simmons DD, Manson-Gieseke L, Hendrix TW, McCarter S. Reconstructions of efferent fibers in
the postnatal hamster cochlea. Hear Res. 1990;49: 127–39.

36.   Pujol R. Morphology, synaptology and electrophysiology of the developing cochlea. Acta
Otolaryngol Suppl. 1985;421: 5-9.

37.   Pujol R, Carlier E, Devigne C. Different patterns of cochlear innervation during the development
of the kitten. J Comp Neurol. 1978;177: 529–36.

38.   He DZ, Dallos P. Development of acetylcholine-induced responses in neonatal gerbil outer hair
cells. J Neurophysiol. 1999;81: 1162–1170. doi:10.1152/jn.1999.81.3.1162

39.   Dulon D, Lenoir M. Cholinergic responses in developing outer hair cells of the rat cochlea. Eur J
Neurosci. 1996;8: 1945–1952.

40.   Liberman MC. The olivocochlear efferent bundle and susceptibility of the inner ear to acoustic
injury. J Neurophysiol. 1991;65: 123–132. Available:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list
_uids=1999726

41.   Maison SF, Liberman MC. Predicting Vulnerability to Acoustic Injury with a Noninvasive Assay of
Olivocochlear Reflex Strength. J Neurosci. 2000;20: 4701–4707. doi:20/12/4701 [pii]

42.   Taranda J, Maison SF, Ballestero JA, Katz E, Savino J, Vetter DE, et al. A point mutation in the hair
cell nicotinic cholinergic receptor prolongs cochlear inhibition and enhances noise protection.
PLoS Biol. 2009;7: 71–83. doi:10.1371/journal.pbio.1000018

43.   Maison SF, Usubuchi H, Liberman MC. Efferent feedback minimizes cochlear neuropathy from
862   moderate noise exposure. J Neurosci. 2013;33: 5542–5552. doi:10.1523/jneurosci.5027-12.2013

863   44.   Boero LE, Castagna VC, Di Guilmi MN, Goutman JD, Elgoyhen AB, Gómez-Casati ME.
864   Enhancement of the medial olivocochlear system prevents hidden hearing loss. J Neurosci.
865   2018;38: 7440–7451. doi:10.1523/JNEUROSCI.0363-18.2018

866   45.   Vetter DE, Liberman MC, Mann J, Barhanin J, Boulter J, Brown MC, et al. Role of alpha9 nicotinic
867   ACh receptor subunits in the development and function of cochlear efferent innervation. Neuron.
868   1999;23: 93-103.

869   46.   Buchwald JS, Huang CM. Far-field acoustic response: Origins in the cat. Science. 1975;189: 382–
870   384. doi:10.1126/science.1145206

871   47.   Karplus M, Lee C, Cashore WJ, Oh W. The effects of brain bilirubin deposition on auditory brain
872   stem evoked responses in rats. Early Hum Dev. 1988;16: 185–194. doi:10.1016/0378-
873   3782(88)90098-9

874   48.   Shapiro SM. Acute brainstem auditory evoked potential abnormalities in jaundiced Gunn rats
875   given sulfonamide. Pediatr Res. 1988;23: 306–310. doi:10.1203/00006450-198803000-00015

876   49.   Shaw NA. The auditory evoked potential in the rat--a review. Prog Neurobiol. 1988;31: 19–45.
877   doi:10.1016/0301-0082(88)90021-4

878   50.   Melcher J, Kiang NY. Generators of the brainstem auditory evoked potential in cat. III: Identified
879   cell populations. Hear Res. 1996;93: 52–71. doi:10.1016/0378-5955(95)00200-6

880   51.   Kim SE, Turkington K, Kushmerick C, Kim JH. Central dysmyelination reduces the temporal fidelity
881   of synaptic transmission and the reliability of postsynaptic firing during high-frequency
882   stimulation. J Neurophysiol. 2013;110: 1621–1630. doi:10.1152/jn.00117.2013

883   52.   Antoli-Candela FJ, Kiang N. Unit activity underlying the N1 potential. In: Naunton R, Fernandez C,
884   editors. Evoked electrical activity in the auditory nervous system. 1978. pp. 165–191.

885   53.   Kujawa SG, Liberman MC. Adding insult to injury: cochlear nerve degeneration after "temporary"
886   noise-induced hearing loss. J Neurosci. 2009;29: 14077–14085. doi:10.1523/JNEUROSCI.2845-
887   09.2009

888   54.   Shera CA, Guinan Jr. JJ. Evoked otoacoustic emissions arise by two fundamentally different
889   mechanisms: a taxonomy for mammalian OAEs. J Acoust Soc Am. 1999;105: 782–798.
890   doi:10.1121/1.426948.

891   55.   Robles L, Ruggero MA. Mechanics of the mammalian cochlea. Physiol Rev. 2001;81: 1305–52.
892   doi:10.1152/physrev.2001.81.3.1305

893   56.   Khimich D, Nouvian R, Pujol R, Tom Dieck S, Egner A, Gundelfinger ED, et al. Hair cell synaptic
894   ribbons are essential for synchronous auditory signalling. Nature. 2005;434: 889–894.
895   doi:10.1038/nature03393.

57. Matsubara A, Laake JH, Davanger S, Usami S, Ottersen OP. Organization of AMPA receptor
subunits at a glutamate synapse: a quantitative immunogold analysis of hair cell synapses in the
rat organ of Corti. J Neurosci. 1996;16: 4457–4467.

58. Liberman LD, Wang H, Liberman MC. Opposing gradients of ribbon size and AMPA receptor
expression underlie sensitivity differences among cochlear-nerve/hair-cell synapses. J Neurosci
Off J Soc Neurosci. 2011;31: 801–8. doi:10.1523/JNEUROSCI.3389-10.2011

59. Sendin G, Bourien J, Rassendren F, Puel J-L, Nouvian R. Spatiotemporal pattern of action potential
firing in developing inner hair cells of the mouse cochlea. Proc Natl Acad Sci U S A. 2014;111:
1999–2004. doi:10.1073/pnas.1319615111

60. Hirtz JJ, Boesen M, Braun N, Deitmer JW, Kramer F, Lohr C, et al. Cav1.3 calcium channels are
required for normal development of the auditory brainstem. J Neurosci Off J Soc Neurosci.
2011;31: 8280–8294. doi:10.1523/JNEUROSCI.5098-10.2011

61. Babola TA, Li S, Wang Z, Kersbergen CJ, Elgoyhen AB, Coate TM, et al. Purinergic Signaling
Controls Spontaneous Activity in the Auditory System throughout Early Development. J Neurosci
Off J Soc Neurosci. 2021;41: 594–612. doi:10.1523/JNEUROSCI.2178-20.2020

62. Frank T, Khimich D, Neef A, Moser T. Mechanisms contributing to synaptic Ca2+ signals and their
heterogeneity in hair cells. Proc Natl Acad Sci U S A. 2009;106: 4483–4488.
doi:10.1073/pnas.0813213106

63. Sheets L, Kindt KS, Nicolson T. Presynaptic CaV1.3 channels regulate synaptic ribbon size and are
required for synaptic maintenance in sensory hair cells. J Neurosci Off J Soc Neurosci. 2012;32:
17273–17286. doi:10.1523/JNEUROSCI.3005-12.2012

64. Wong HC, Zhang Q, Beirl AJ, Petralia RS, Wang Y-X, Kindt K. Synaptic mitochondria regulate hair-
cell synapse size and function. eLife. 2019;8: e48914. doi:10.7554/eLife.48914

65. Özçete ÖD, Moser T. A sensory cell diversifies its output by varying Ca2+ influx-release coupling
among active zones. EMBO J. 2021;40: e106010. doi:10.15252/embj.2020106010

66. Michanski S, Smaluch K, Maria Steyer A, Chakrabarti R, Setz C, Oestreicher D, et al. Mapping
developmental maturation of inner hair cell ribbon synapses in the apical mouse cochlea. Proc
Natl Acad Sci U S A. 2019;116: 6415–6424. doi:10.1073/pnas.1812029116

67. Hua Y, Ding X, Wang H, Wang F, Lu Y, Neef J, et al. Electron Microscopic Reconstruction of Neural
Circuitry in the Cochlea. Cell Rep. 2021;34: 108551. doi:10.1016/j.celrep.2020.108551

68. Payne SA, Joens MS, Chung H, Skigen N, Frank A, Gattani S, et al. Maturation of Heterogeneity in
Afferent Synapse Ultrastructure in the Mouse Cochlea. Front Synaptic Neurosci. 2021;13: 678575.
doi:10.3389/fnsyn.2021.678575

69. Sobkowicz HM, Rose JE, Scott GE, Slapnick SM. Ribbon synapses in the developing intact and
cultured organ of Corti in the mouse. J Neurosci. 1982;2: 942–57.

70.   Merchan-Perez A, Liberman MC. Ultrastructural differences among afferent synapses on cochlear
932   hair cells: correlations with spontaneous discharge rate. J Comp Neurol. 1996;371: 208–221.
933   doi:10.1002/(SICI)1096-9861(19960722)371:2<208::AID-CNE2>3.0.CO;2-6

934   71.   Huang L-C, Barclay M, Lee K, Peter S, Housley GD, Thorne PR, et al. Synaptic profiles during
935   neurite extension, refinement and retraction in the developing cochlea. Neural Develop. 2012;7:
936   38. doi:10.1186/1749-8104-7-38

937   72.   Wong AB, Rutherford MA, Gabrielaitis M, Pangršič T, Göttfert F, Frank T, et al. Developmental
938   refinement of hair cell synapses tightens the coupling of Ca2+ influx to exocytosis. EMBO J.
939   2014;33: 247–264. doi:10.1002/embj.201387110

940   73.   Song L, McGee J, Walsh EJ. Frequency- and level-dependent changes in auditory brainstem
941   responses (ABRS) in developing mice. J Acoust Soc Am. 2006;119: 2242–2257.
942   doi:10.1121/1.2180533

943   74.   Fernandez KA, Jeffers PWC, Lall K, Liberman MC, Kujawa SG. Aging after noise exposure :
944   Acceleration of cochlear synaptopathy in " recovered " ears. J Neurosci. 2015;35: 7509–20.
945   doi:10.1523/JNEUROSCI.5138-14.2015

946   75.   Schrode KM, Dent ML, Lauer AM. Sources of variability in auditory brainstem response thresholds
947   in a mouse model of noise-induced hearing loss. J Acoust Soc Am. 2022;152: 3576–3582.
948   doi:10.1121/10.0016593

949   76.   Lu Y, Liu J, Li B, Wang H, Wang F, Wang S, et al. Spatial patterns of noise-induced inner hair cell
950   ribbon loss in the mouse mid-cochlea. iScience. 2024;27: 108825. doi:10.1016/j.isci.2024.108825

951   77.   Rybalko N, Chumak T, Bureš Z, Popelář J, Šuta D, Syka J. Development of the acoustic startle
952   response in rats and its change after early acoustic trauma. Behav Brain Res. 2015;286: 212–221.
953   doi:10.1016/j.bbr.2015.02.046

954   78.   Mohamad NI, Santra P, Park Y, Matthews IR, Taketa E, Chan DK. Synaptic ribbon dynamics after
955   noise exposure in the hearing cochlea. Commun Biol. 2024;7: 421. doi:10.1038/s42003-024-
956   06067-y

957   79.   Moverman DJ, Liberman LD, Kraemer S, Corfas G, Liberman MC. Ultrastructure of noise-induced
958   cochlear synaptopathy. Sci Rep. 2023;13: 19456. doi:10.1038/s41598-023-46859-6

959   80.   Stamataki S, Francis HW, Lehar M, May BJ, Ryugo DK. Synaptic alterations at inner hair cells
960   precede spiral ganglion cell loss in aging C57BL/6J mice. Hear Res. 2006;221: 104–118.
961   doi:10.1016/j.heares.2006.07.014

962   81.   Jeng JY, Ceriani F, Olt J, Brown SDM, Holley MC, Bowl MR, et al. Pathophysiological changes in
963   inner hair cell ribbon synapses in the ageing mammalian cochlea. J Physiol. 2020;598: 4339–4355.
964   doi:10.1113/JP280018

965   82.   Boero LE, Payne S, Gómez-Casati ME, Rutherford MA, Goutman JD. Noise Exposure Potentiates
966   Exocytosis From Cochlear Inner Hair Cells. Front Synaptic Neurosci. 2021;13: 740368.
967   doi:10.3389/fnsyn.2021.740368

968   83.   Suthakar K, Liberman MC. Auditory-Nerve Responses in Mice with Noise-Induced Cochlear
969   Synaptopathy. https://doi.org/101152/jn003422021. 2021 [cited 24 Nov 2021].
970   doi:10.1152/JN.00342.2021

971   84.   Wood MB, Nowak N, Mull K, Goldring A, Lehar M, Fuchs PA. Acoustic Trauma Increases Ribbon
972   Number and Size in Outer Hair Cells of the Mouse Cochlea. J Assoc Res Otolaryngol JARO.
973   2021;22: 19–31. doi:10.1007/s10162-020-00777-w

974   85.   Furman AC, Kujawa SG, Liberman MC. Noise-induced cochlear neuropathy is selective for fibers
975   with low spontaneous rates. J Neurophysiol. 2013;110: 577–586. doi:10.1152/jn.00164.2013

976   86.   Benini F, Magnavita V, Lago P, Arslan E, Pisan P. Evaluation of noise in the neonatal intensive care
977   unit. Am J Perinatol. 1996;13: 37–41. doi:10.1055/s-2007-994200

978   87.   Lahav A, Skoe E. An acoustic gap between the NICU and womb: a potential risk for compromised
979   neuroplasticity of the auditory system in preterm infants. Front Neurosci. 2014;8: 381.
980   doi:10.3389/fnins.2014.00381

981   88.   Muller M, von Hunerbein K, Hoidis S, Smolders JW. A physiological place-frequency map of the
982   cochlea in the CBA/J mouse. Hear Res. 2005;202: 63–73. Available:
983   http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list
984   _uids=15811700

985   89.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-source
986   platform for biological-image analysis. Nat Methods. 2012;9: 676–82. doi:10.1038/nmeth.2019

987   90.   Sage D, Donati L, Soulez F, Fortun D, Schmit G, Seitz A, et al. DeconvolutionLab2: An open-source
988   software for deconvolution microscopy. Methods. 2017;115: 28–41.
989   doi:10.1016/j.ymeth.2016.12.015

990   91.   Jing Z, Rutherford MA, Takago H, Frank T, Fejtova A, Khimich D, et al. Disruption of the
991   presynaptic cytomatrix protein bassoon degrades ribbon anchorage, multiquantal release, and
992   sound encoding at the hair cell afferent synapse. J Neurosci. 2013;33: 4456–4467.
993   doi:10.1523/JNEUROSCI.3491-12.2013

994   92.   Gilels F, Paquette ST, Zhang J, Rahman I, White PM. Mutation of foxo3 causes adult onset
995   auditory neuropathy and alters cochlear synapse architecture in mice. J Neurosci. 2013;33:
996   18409–18424. doi:10.1523/JNEUROSCI.2529-13.2013

997   93.   Paquette ST, Gilels F, White PM. Noise exposure modulates cochlear inner hair cell ribbon
998   volumes, correlating with changes in auditory measures in the FVB/nJ mouse. Sci Rep. 2016;6: 1–
999   13. doi:10.1038/srep25056

1000   94.   Ollion J, Cochennec J, Loll F, Escudé C, Boudier T. TANGO: A generic tool for high-throughput 3D
1001   image analysis for studying nuclear organization. Bioinformatics. 2013;29: 1840–1841.
1002   doi:10.1093/bioinformatics/btt276

1003