Strengthening Medial Olivocochlear Feedback Reduces the Developmental Impact of Early Noise Exposure

Authors: Valeria C. Castagna[1,2,3], Luis E. Boero[1,2,3], Mariano N. Di Guilmi[1], Camila Catalano Di Meo[1,2,3], Jimena A. Ballestero[2], Paul A. Fuchs[4,5], Amanda M. Lauer[4,5,6], Ana Belén Elgoyhen[1,2] and Maria Eugenia Gomez-Casati[2]

Running Title: Early Noise Protection via MOC Feedback

Affiliations:

[1]Instituto de Investigaciones en Ingeniería Genética y Biología Molecular, Dr. Héctor N. Torres, Consejo Nacional de Investigaciones Científicas y Técnicas, 1428 Buenos Aires, Argentina.

[2]Instituto de Farmacología, Facultad de Medicina, Universidad de Buenos Aires, 1121 Buenos Aires, Argentina and Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina.

[3]Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Fisiología y Biología Molecular y Celular, 1428 Buenos Aires, Argentina.

[4]Department of Otolaryngology-Head and Neck Surgery, Johns Hopkins University School of Medicine; Baltimore, Maryland 21205, United States.

[5]Department of Neuroscience, Johns Hopkins University School of Medicine, Baltimore, Maryland 21205, United States.

[6]Center for Functional Anatomy and Evolution, Johns Hopkins University, School of Medicine, Baltimore, Maryland 21205, United States.

Corresponding Author: María Eugenia Gómez-Casati, Ph.D.

Instituto de Farmacología. Facultad de Medicina. Universidad de

Buenos Aires.

Address: Paraguay 2155. C1121ABG, Buenos Aires, Argentina.

Phone: 54-11-52853350

E-mail: mgomezcasati@fmed.uba.ar, megomezcasati@gmail.com

Number of Pages: 41

Number of Figures: 8

Number of words Abstract: 250

Number of words Introduction: 660

Number of words Discussion: 1497

Conflict of Interest: The authors declare no competing financial or non-financial interest.

Acknowledgements: This research was supported by Agencia Nacional de Promoción Científica

y Técnica (Argentina) to M.E.G-C and NIH Grant R01 DC001508 (P.A.F, A.M.L, A.B.E and M.E.G-

C) and the David M. Rubenstein Fund for Hearing Research.

Author Contributions: V.C.C, L.E.B, M.N.D-G, P.A.F, A.M.L, A.B.E and M.E.G-C designed

research; V.C.C, C. C.dM and J.A.B performed experiments; V.C.C analyzed data. M.E.G-C

wrote the paper. All authors discussed the results, read, edited, and commented on the

manuscript.

Abstract

The early onset of peripheral deafness significantly alters the proper development of the auditory system. Likewise, exposure to loud noise during early development produces a similar disruptive effect. Before hearing onset in altricial mammals, cochlear inner hair cells exhibit spontaneous electrical activity that drives auditory circuit development. This activity is modulated by medial olivocochlear (MOC) efferent feedback through α9α10 nicotinic cholinergic receptors in inner hair cells. In adults, these receptors are restricted to outer hair cells, where they mediate MOC feedback to regulate cochlear amplification. Although the MOC system's protective role to prevent noise-induced hearing loss in adulthood is well-established, its influence during early developmental stages -especially in response to exposure to loud noise- remains largely unexplored. In this study, we investigated the role of MOC feedback during early postnatal development using α9 knockout (KO) and α9 knock-in (KI) mice of either sex, which respectively lack or exhibit enhanced cholinergic activity. Our findings reveal that both increased and absent olivocochlear activity result in altered auditory sensitivity at the onset of hearing, along with long–range alterations in the number and morphology of ribbon synapses. Early noise exposure caused lasting auditory damage in both wild-type and α9KO mice, with deficits persisting into adulthood. In contrast, α9KI mice were protected from noise-induced damage, with no long-term effects on auditory function. These results highlight the increased susceptibility of the auditory system during early postnatal development. Moreover, they indicate that an enhanced MOC feedback shields the auditory system from noise damage during this period.

SIGNIFICANCE STATEMENT

Early development represents a sensitive window for shaping auditory function. We show that the medial olivocochlear system is critical for establishing normal ribbon synapse density and size; key features for proper hearing onset. We also show that the developing auditory system is

73   especially vulnerable to loud noise, with early exposure causing more severe and lasting effects

74   than similar noise later in life. Notably, enhancing α9α10 nAChR receptor activity during this early

75   stage offers protection against noise-induced damage, revealing a time-sensitive opportunity to

76   safeguard auditory development.

Introduction

Cochlear and central auditory system development relies on tightly coordinated cellular

and molecular events, with critical refinements during early postnatal life to optimize sound

encoding (Sanes and Walsh, 1998). Noise exposure during this sensitive period can disrupt these

mechanisms, damaging cochlear structures and neural circuits (Chang and Merzenich, 2003;

Keuroghlian and Knudsen, 2007; Zhou and Merzenich, 2008; Sanes and Bao, 2009; Lauer et al.,

2011). Early acoustic exposure affects neuronal sensitivity, innervation, frequency tuning and

response characteristics across the auditory pathway (Caminos et al., 2005; Waguespack et al.,

2007; Sanes and Bao, 2009; Dorrn et al., 2010; Sun et al., 2010; Froemke and Jones, 2011).

Such disruptions can lead to permanent auditory deficits, emphasizing the importance of

safeguarding the developing auditory system.

In altricial mammals, inner hair cells (IHCs) exhibit spontaneous electrical activity that

propagates into the central nervous system (CNS) before hearing onset (~P12 in mice) (Lippe,

1994; Marcotti et al., 2003; Sonntag et al., 2009; Tritsch and Bergles, 2010; Johnson et al., 2011;

De Faveri et al., 2025). This activity is crucial for neuronal survival and for refining auditory circuits

(Tierney et al., 1997; Friauf and Lohmann, 1999; Mostafapour et al., 2000; Leake et al., 2006;

Clause et al., 2014; Di Guilmi et al., 2019). During this period, MOC neurons from the superior

olivary complex form transient axosomatic synapses with immature IHCs, modulating their

spontaneous firing (Glowatzki and Fuchs, 2000; Marcotti et al., 2004; Johnson et al., 2013). These

synapses, mediated by α9α10 nicotinic cholinergic receptors (nAChRs) coupled to SK2 channels,

hyperpolarize IHCs (Glowatzki and Fuchs, 2000; Marcotti et al., 2004; Goutman et al., 2005),

shaping burst patterns that guide synaptic development and ensure bilateral representation in the

CNS (Walsh et al., 1998; Johnson et al., 2013; Clause et al., 2014; Di Guilmi et al., 2019; Wang

et al., 2021). As IHCs mature, they lose direct efferent contacts and no longer respond to

acetylcholine (ACh) (Katz et al., 2004; Roux et al., 2011). At this stage, MOC fibers cross the

tunnel of Corti and innervate OHCs to regulate cochlear amplification (Wheeler et al., 1994;

Ernfors et al., 1995; Knipper et al., 1995; Simmons et al., 1996; Walsh et al., 1998; Vattino et al.,

2020). ACh-induced responses in OHCs emerge by P6, shifting from nicotinic-like to outward

potassium currents by hearing onset (Dulon and Lenoir, 1996; He and Dallos, 1999). The

olivocochlear system matures rapidly within the first 2–3 postnatal weeks, with adult-like

innervation by the end of the first month (Pujol et al., 1978; Pujol, 1985; Simmons et al., 1990,

1996). While the protective role of the MOC system in preventing noise-induced hearing loss in

adulthood is well-documented (Liberman, 1991; Maison and Liberman, 2000; Maison et al., 2002,

2013; Taranda et al., 2009; Boero et al., 2018), its impact during early developmental stages,

remains largely unexplored.

This study explores how cholinergic MOC input strength influences auditory development

and noise vulnerability during this sensitive period. We used mice with altered α9α10 nAChR

activity: α9KO mice, which lacks efferent transmission (Vetter et al., 1999), and α9KI, carrying an

α9 point mutation that leads to an enhanced MOC effect (Taranda et al., 2009). We show that

MOC strength modulates hearing onset timing and IHC ribbon synapse formation. Wild-type (WT)

mice exposed to noise at hearing onset showed greater threshold shifts and synaptic loss than

similar exposure in adulthood (Boero et al., 2018), revealing heightened developmental

vulnerability. In α9KO mice, early exposure resulted in a persistent loss of auditory nerve

synapses, emphasizing the essential role of the MOC pathway. Conversely, α9KI mice were

protected from both thresholds shifts and synaptic loss after early noise exposure. This protection

occurred during a developmental window when the auditory system is still maturing and MOC-

OHC synapses are forming (Simmons et al., 1996; Vattino et al., 2020). These findings highlight

the importance of efferent cholinergic input during cochlear development and suggest that

strengthening MOC feedback during this window can mitigate long-term effects of early acoustic

trauma.

Materials and Methods

Animals. α9KO and α9KI mice have been previously described (Vetter et al., 1999; Taranda et al., 2009) and were backcrossed with the congenic FVB.129P2-*Pde6b⁺ Tyrc-ch/AntJ* strain (https://www.jax.org/strain/004828) for seventeen generations (i.e., N-17). We maintained a consistent male-to-female ratio across all experimental groups and genotypes. No sex-related differences were observed in the data. This study included adult and pup animals aged P12 to P75, which were housed in a sound-controlled room within the animal care facility. Background sound levels in this environment were consistently below 50 dB SPL. Noise levels were measured using an electret condenser microphone (FG-23329-PO7; Knowles). Transient sounds during routine cage cleaning and daily maintenance by animal care staff were minimal, with peaks not exceeding 70 dB SPL. The methods reported in this study adhere to the ARRIVE guidelines for reporting *in vivo* animal research (https://arriveguidelines.org/), ensuring transparency and reproducibility. All procedures were conducted in accordance with institutional and ethical guidelines for animal care and use.

Auditory function tests. Physiological assessments of cochlear and central function, including auditory brainstem responses (ABRs) and distortion-product otoacoustic emissions (DPOAEs), were performed in mice anesthetized with xylazine (10 mg/kg, i.p.) and ketamine (100 mg/kg, i.p.) and placed in a soundproof chamber maintained at 32°C to counteract the drop in body temperature caused by anesthesia. ABR represents sound-evoked potentials generated by neural circuits in the ascending auditory pathways, while DPOAEs are sounds emitted by the cochlea, specifically by OHCs, in response to two pure-tone stimuli presented simultaneously. Sound stimuli were delivered through a custom acoustic system with two dynamic earphones used as sound sources (CDMG15008–03A; CUI) and an electret condenser microphone (FG-

23329-PO7; Knowles) coupled to a probe tube to measure sound pressure near the eardrum.

Data collection was conducted using a National Instruments PXI-based system equipped with 24-

bit input/output boards and operated through a custom LabView software interface. For ABRs,

platinum needle electrodes were placed into the skin at the dorsal midline close to the neural crest

and pinna with a ground electrode near the tail. ABR potentials were evoked with 5 ms tone pips

(0.5 ms rise-fall, with a $\cos^2$ envelope, at 40/s) delivered to the eardrum at log-spaced frequencies

from 5.6 to 45.25 kHz. The response was amplified 10,000X with a 0.3–3 kHz passband. Sound

level was raised in 5 dB steps from 20 to 80 dB sound pressure level (SPL). At each level, 1024

responses were averaged with alternated stimulus polarity. Threshold for ABR was defined as

the lowest stimulus level (<80 dB SPL) at which a repeatable wave 1 could be identified in the

response waveform. The ABR wave 1 amplitude was computed by off-line analysis of the peak-

to-peak amplitude of stored waveforms. The DPOAEs in response to two primary tones of

frequency f1 and f2 were recorded at 2f1-f2, with f2/f1=1.2, and the f2 level 10 dB lower than the

f1 level. Ear-canal sound pressure was amplified and digitally sampled at 4 μs intervals. DPOAE

threshold was defined as the lowest f2 level in which the signal to noise floor ratio is >1.

Noise exposure. Animals were exposed under anesthesia at P15 to broadband noise (1–16 kHz,

100 dB SPL) for 1 hour using closed-field earphones (CDMG15008–03A; CUI) as the sound

source. This broader frequency range was selected to induce significant threshold shifts in the

apical region of the cochlea, as previously described in (Boero et al., 2018). Noise exposure was

conducted in the same acoustic chamber used for cochlear function tests. The noise consisted of

a flat-spectrum broadband signal calibrated to the target SPL immediately before each exposure

session.

Animal monitoring and health assessment. To assess the impact of repeated anesthesia on

the health and well-being of the animals, their weight and motor activity were monitored daily

throughout the auditory function recordings and noise exposure protocols. Animals were observed

during their recovery from anesthesia to ensure no adverse effects on their general condition.

Growth curves showed no significant differences in weight between animals that were

anesthetized and untreated controls animals, suggesting that anesthesia did not affect normal

growth. Furthermore, no behavioral changes or motor impairments were observed in the

anesthetized animals when compared to controls, indicating that repeated anesthesia had no

detectable impact on overall health or behavior.

Cochlear processing and immunostaining. Cochleae were perfused intralabyrinthly with 4%

paraformaldehyde (PFA) in phosphate-buffered saline (PBS), post-fixed with 4% PFA overnight,

and decalcified in 0.12 M EDTA. Cochlear tissues were microdissected and permeabilized by

freeze/thaw cycles in 30% sucrose. The dissected samples were then separated into distinct wells

based on cochlear region: apex, middle and base. Tissue samples were blocked with 5% normal

goat serum and 1% Triton X-100 in PBS for 1 h, followed by incubation in primary antibodies

(diluted in blocking buffer) at 37°C for 16 h. The primary antibodies used in this study were: 1)

anti-C-terminal binding protein 2 (mouse anti-CtBP2 IgG1; BD Biosciences, San Jose, CA;

Cat#612044, RRID:AB_399431, 1:200) to label the presynaptic ribbon, 2) anti-glutamate receptor

2 (mouse anti-GluA2 IgG2a; Millipore, Billerica, MA; Cat#MAB397, RRID:AB_11212990, 1:2000)

to label the post-synaptic receptor plaques and 3) anti-Myosin 7a (rabbit anti-MyosinVIIa, Proteus

Biosciences; Cat# 25-6790, RRID:AB_10015251, 1:200) to label the cochlear hair cells. Tissues

were then incubated with the appropriate Alexa Fluor-conjugated fluorescent secondary

antibodies (Invitrogen, Carlsbad, CA; 1:500 in blocking buffer) for 2 hours at room temperature.

Finally, tissues were mounted on microscope slides in FluorSave mounting media (Millipore,

Billerica, MA).

Confocal microscopy and image processing. For IHC synaptic counts, cochlear tissue from

the apex, middle and base regions was first imaged at low magnification to correlate each piece

with its corresponding frequency region (Müller et al., 2005). The apical region corresponds to 4–

12 kHz, the middle region to 12–20 kHz, and the base region to 22–40 kHz. Z-stacks images were

then acquired using a Leica TCS SPE microscope with a 63X oil-immersion lens (1.5X or 4X

digital zoom). Imaging settings were consistent across all samples, ensuring no pixel saturation.

For each stack, the Z-step was set to 0.3 µm, with a pixel size of 0.11 µm in both the X and Y

axes. Each stack typically contained 10 to 20 IHCs. Image stacks were processed using Fiji

software (RRID:SCR_002285) (Schindelin et al., 2012), and a custom plugin was developed to

automate the counting of synaptic ribbons, glutamate receptor patches and co-localized synaptic

puncta. The algorithm's accuracy was validated by comparing automated counts to manual

counts, ensuring reliable quantification. Briefly, each channel was analyzed separately, and

maximum projections were generated to quantify the number of CtBP2 or GluA2 puncta.

Additionally, a composite between the three channels was produced to draw the different regions

of interest (ROI) that correspond to each IHC taking the myosin staining as a reference. The

maximum projections from the single channels were multiplied to generate a merged 32-bit image.

Then, they were converted to binary images after a custom thresholding procedure. An automatic

counting of the number of particles on each ROI was performed. The plugin used for synapse

quantification, along with detailed instructions, is available at:

https://github.com/vcastagna/CountsSynapses.

For quantitative volumetric synaptic analysis, different regions from the whole cochlea

were imaged and confocal z-stacks were processed in Fiji. z-stacks were deconvolved applying

10 iterations of the Richardson-Lucy algorithm in the DeconvolutionLab2 plugin (Sage et al.,

2017), an algorithm that has been used previously in cochlear whole mounts (Jing et al., 2013).

Then, the 3D Object Counter plugin was used to perform three-dimensional image segmentation

and object individualization of the deconvoluted stacks. According to previous work (Liberman et

al., 2011; Gilels et al., 2013; Paquette et al., 2016) we reasoned that a synapse would occupy a

volume between 0.04 µm3 to 5 µm3, and then we set a size range of 20-2000 voxels, with a voxel

size of 0.11 µm x 0.11 µm x 0.2 µm. The volumes of presynaptic ribbons (CtBP2 fluorescence)

and postsynaptic AMPA receptor clusters (GluA2 fluorescence) at IHC-afferent synapses were

measured using the 3D ROI Manager (Ollion et al., 2013). All exported volumes were checked to

ensure that volumes were within instrument resolution limits after threshold adjustments.

Statistical analysis. Data was analyzed using R Statistical Software (RRID:SCR_001905).

Shapiro-Wilks test was used for testing normal distribution of the residuals. In cases where the

data did not follow a normal distribution, the following tests were used: Kruskal-Wallis test followed

by multiple contrasts according to Dunn's method for comparisons between three groups (IHCs

synaptic counts in the control condition, ABR and DPOAE thresholds during development) and

Mann-Whitney U test for comparisons between two groups (ABR thresholds, DPOAE thresholds,

and IHCs synaptic counts). If data were normally distributed, a one-way ANOVA was conducted,

followed by Holm-Sidak post hoc tests for three-group comparisons (wave 1 amplitudes and

synaptic volumes in the control condition). For two-group comparisons, a t-test was used (synaptic

volumes). Statistical significance was set to $p < 0.05$.

Results

Onset of auditory perception in mice with varying levels of efferent inhibition

In this study we used mice with distinct levels of α9α10 nAChR activity: 1) WT, with normal

MOC function; 2) α9KO, which lacks cholinergic transmission between MOC neurons and hair

cells (Vetter et al., 1999), and 3) α9KI mice carrying an α9 nAChR subunit point mutation that

leads to enhanced responses to ACh (Taranda et al., 2009). To explore the impact of noise

exposure at hearing onset, we first identified the optimal developmental stage for introducing such

exposure, by assessing the onset of hearing under control (unexposed) conditions in these mouse

models. This was done by tracking auditory brainstem responses (ABRs), the sound-evoked

potentials generated by neuronal circuits in the ascending auditory pathway, at different postnatal

ages across the three groups. Individual ABR waves reflect the activation of auditory periphery

and brainstem processing relay stations within the first ~7 ms after sound stimulus onset

(Buchwald and Huang, 1975; Karplus et al., 1988; Shapiro, 1988; Shaw, 1988; Melcher and

Kiang, 1996; Kim et al., 2013). Wave 1 represents the synchrony in firing of spiral ganglion

neurons and is highly correlated with the number of synapses between IHCs and auditory nerve

fibers (Buchwald and Huang, 1975; Antoli-Candela and Kiang, 1978; Kujawa and Liberman,

2009). Measurements were taken daily from P12 to P16, as well as P22 and P75, in anesthetized

mice. Interestingly, we found that ABR onset occurred earlier in α9KI mice, with enhanced MOC

function, compared to WT mice with normal MOC synapses and was delayed in α9KO mice

lacking α9α10 nAChR activity (Figure 1a and b). Representative ABR traces recorded at P14 in

response to a 16 kHz stimulus across varying intensities illustrate differences in ABR thresholds

among the three mouse groups with distinct levels of α9α10 nAChR activity (Figure 1a). At this

developmental stage, α9KI mice exhibited lower ABR thresholds, and more distinct peaks at

suprathreshold intensities compared to WT mice. In contrast, α9KO mice showed higher ABR

thresholds relative to WT mice (Figure 1a). Figure 1b shows the mean ABR thresholds measured

at test frequencies of 11.33, 16, and 22.65 kHz from P12 to P75 across the three groups of mice.

A comparison between α9KI and WT mice revealed that α9KI mice displayed measurable ABR

responses as early as P13 (Kruskal-Wallis, followed by Dunn's test, degree of freedom [df] = 2, p

= 0.01 at 22.65 kHz; Figure 1b) and achieved mature threshold levels earlier than their WT

littermates (Kruskal-Wallis, followed by Dunn's test, df = 2, p = 0.02 at 16 and 22.65 kHz at P14;

Figure 1b). In contrast, ABR onset was delayed by a day in α9KO compared to WT mice (Kruskal-

Wallis followed by Dunn's test, df = 2, p = 0.04 at 11.33 and 16 kHz; Figure 1b), indicating that

cochlear maturation is slower in the absence of pre-hearing efferent modulation. To summarize,

at P13, no ABR responses were observed in WT (0 out of 9) or α9KO (0 out of 7) mice in response

to tone pips at 80 dB SPL across 11.33, 16, and 22.65 kHz. In contrast, 6 out of 9 α9KI mice

exhibited ABR responses at P13. By P14, ABR responses were present in 77.7% of WT mice at

11.33 and 22.65 kHz and in 100% at 16 kHz. Meanwhile, 85% of α9KO mice responded at all

frequencies, but with higher thresholds than WT (Figure 1b). All α9KI mice exhibited robust

responses at all tested frequencies by P14. By P15, 100% of WT and 91% of α9KO mice showed

ABR responses.

Tone pip-evoked ABR wave 1 amplitudes were then analyzed at a suprathreshold level of

80 dB SPL at the different test frequencies in the three groups of mice with varying levels of α9α10

nAChR activity (Figure 2a, b). As illustrated in the representative traces in Figure 2a, measurable

ABR wave 1 amplitudes at P13 were only observed in α9KI mice. At P14, the number of

discernible peaks were fewer in WT and α9KO mice compared to α9KI mice. Up to P15, response

peaks at 80 dB SPL were slightly larger and more distinct in α9KI mice, reflecting enhanced

auditory responses during this postnatal developmental window. However, starting at P16, α9KI

mice exhibited smaller ABR wave 1 amplitudes, with significant differences observed at P75 at

16 and 22.65 kHz compared to WT mice (one way ANOVA followed by Holm-Sidak test, df = 2, p

= 0.03 at both frequencies; Figure 2b). In α9KO mice, wave 1 amplitudes remained smaller than

those of WT and α9KI mice until P16, with the difference being significant at 22.65 kHz (one way

ANOVA followed by Holm-Sidak test, df = 2, p = 0.02; Figure 2b). These findings align with the

ABR threshold results, indicating that the strength of efferent cholinergic inhibition to hair cells

contributes to the timely development of auditory function. The reduced ABR wave 1 amplitudes

obtained in mature mice with either enhanced or null cholinergic activity indicates that proper

MOC modulation during development might be crucial for the correct formation of auditory nerve

synapses. Analysis of ABR wave 1 latencies at different postnatal ages showed no significant

differences between genotypes (Figure 2c). While wave 1 latencies were consistently longer

across all groups at P14, they shortened to levels typical of mature mice by P15.

OHC function was assessed through distortion-product otoacoustic emissions (DPOAE)

at the different postnatal ages. In a healthy cochlea, two close pure tones produce distortions due

to OHC nonlinearities, which are amplified by OHC electromotility and detected with a microphone

in the external ear canal (Shera and Guinan Jr., 1999; Robles and Ruggero, 2001). DPOAE

responses followed a similar pattern to ABR thresholds, though the differences were not

statistically significant (Kruskal-Wallis, df = 2, p > 0.05 at all frequencies; Figure 1c). At P13,

DPOAE responses were nearly absent in WT (1 out of 9) and α9KO (0 out of 7) mice across the

tested frequencies. In contrast, 3 out of 9 α9KI mice showed DPOAE responses at 16 and 22.65

kHz. By P14, DPOAE responses were detectable in 75% of WT and α9KO mice at 16 and 22.65

kHz, and in 60% of mice at 11.33 kHz. All α9KI mice exhibited strong DPOAE responses at all

tested frequencies by P14. By P15, response rates reached 100% in both WT and α9KO mice.

Note that at P22 and P75, mean ABR and DPOAE thresholds in α9KI mice were slightly elevated

by 5–10 dB compared to WT and α9KO mice (Figure 1b, c). This elevation reached statistical

significance only for ABR thresholds at P22 (Kruskal-Wallis, df = 2, p = 0.03 at 11.33 kHz; p =

0.04 at 16 kHz and p = 0.02 at 22.65 kHz). The elevated thresholds in α9KI mice at these mature

stages are most likely due to enhanced suppression of OHCs activity resulting from increased

cholinergic neurotransmission. This effect, previously described by Taranda et al. (2009), can be

reversed by strychnine, a known antagonist of the α9α10 nAChR. Collectively, these findings

indicate that the onset of hearing is altered in mice with either absent or enhanced olivocochlear

cholinergic feedback.

ABR thresholds after noise exposure at hearing onset in mice with different degree of MOC inhibition

To investigate the impact of noise exposure during early development, we exposed WT, α9KO, and α9KI mice to loud noise at P15. This allowed us to evaluate how acoustic experience influences auditory function before full threshold maturation in each genotype. Mice were exposed to 1–16 kHz noise at 100 dB SPL for 1 hour at P15. Auditory responses were then assessed at 1-, 7-, and 60-days post-exposure to track functional changes over time.

Previous studies in WT mice exposed to the same noise protocol at P21 showed a significant increase in auditory thresholds of 10–35 dB SPL one day after exposure, which returned to baseline within a week, indicating a transient threshold shift (Boero et al., 2018). In contrast, as shown in Figure 3a, no changes in ABR thresholds were observed at P16, one day after noise exposure, across frequencies of 8, 16, and 22.65 kHz in WT mice compared to unexposed controls. Remarkably, ABR thresholds at P16 were slightly lower than those measured at P15, prior to noise exposure, and showed no significant difference when compared to unexposed controls at the same age. This suggests that noise did not have an immediate effect on the maturation of auditory thresholds at this postnatal developmental stage. One week after exposure, however, ABR thresholds were elevated at all frequencies in noise-exposed WT animals, and this increase persisted at the same level for at least two months post-exposure (Mann- Whitney test, df = 1, P22: p = 0.006 at 8 kHz; p = 0.007 at 16 kHz; p = 0.002 at 22.65 kHz and P75: p = 0.04 at 8 kHz; p = 0.03 at 16 kHz and p = 0.04 at 22.65 kHz; Figure 3a). This outcome contrasts with the transient threshold elevation observed in WT mice from the same genetic background, exposed at P21 using the same noise protocol (Boero et al., 2018), suggesting that noise exposure at this early developmental stage may disrupt the normal maturation process, potentially leading to more profound and lasting auditory deficits.

A similar trend as seen in WT mice was observed after exposing α9KO mice to noise at

P15 (Figure 3b). One day after exposure, there were no changes in ABR thresholds at 8, 16 and

22.65 kHz. However, seven days post-exposure, there was an elevation in auditory thresholds

across all frequencies (Mann-Whitney test, df = 1, p = 0.03 at 8 kHz; p = 0.01 at 16 kHz and p =

0.02 at 22.65 kHz), which, unlike in WT mice, returned to normal levels after two months (Mann-

Whitney test, df = 1, p > 0.05 at all frequencies; Figure 3b). Notably, in α9KI mice, ABR thresholds

were not affected by noise exposure at all the frequencies tested (Figure 3c), suggesting that at

this early age, the enhancement of cholinergic activity on OHCs provides protection from noise-

induced cochlear threshold elevations (Mann-Whitney test, df = 1, p > 0.05 at all frequencies).

Amplitudes of cochlear sound-evoked potentials after noise exposure at hearing onset in

mice with different degree of efferent inhibition

To assess the neural activity across auditory nerve fibers, we measured ABR wave 1

amplitudes at frequencies of 8, 16, and 22.65 kHz at different ages after noise exposure in the

three mouse groups (Figure 4). One day after noise exposure, we observed a reduction in wave

1 amplitudes across all frequencies in WT mice. Further testing of WT mice one week and two

months post-exposure revealed a significant and long-lasting reduction, indicating a permanent

decrease of ABR wave 1 amplitude (Mann- Whitney test, df = 1, P22: p = 0.007 at 8 kHz; p =

0.005 at 16 kHz; p = 0.005 at 22.65 kHz and P75: p = 0.04 at 8 kHz; p = 0.002 at 16 kHz and p =

0.002 at 22.65 kHz; Figure 4a). Similar permanent reductions in wave 1 amplitude were observed

in α9KO mice after exposure to noise at 16 and 22.65 kHz (Mann-Whitney test, df = 1, P22: p =

0.04 at 16 kHz; p = 0.001 at 22.65 kHz and P75: p = 0.03 at 16 kHz and p = 0.04 at 22.65 kHz;

Figure 4b). Notably, no changes in ABR wave 1 amplitudes were observed at any age post-

exposure in mice with enhanced α9α10 nAChR activity, suggesting a preservation of cochlear

synapses after noise exposure (Mann-Whitney test, df = 1, p > 0.05 at all frequencies; Figure 4c).

DPOAE thresholds after noise exposure at hearing onset in mice with different levels of α9α10 nAChR activity

To assess the impact of hearing onset noise exposure on the function of OHCs, we recorded DPOAEs at 8, 16, and 22.65 kHz on the three groups of mice with varying levels of MOC function (Figure 5). During early development, the auditory system undergoes significant maturation, reflected by a progressive decrease in DPOAE thresholds with age, as shown in Figure 1c. In WT mice, one day after noise exposure at P16, there was no difference in DPOAE thresholds between the noise-exposed group and unexposed controls (Figure 5a). However, when comparing DPOAE thresholds at P16 to those at P15 within both groups (i.e. exposed and unexposed controls), a clear reduction was observed (Figure 5a). This decline in thresholds underscores the natural maturation process of the auditory system, indicating that early noise exposure does not immediately disrupt the ongoing development of DPOAE thresholds at this early postnatal stage. By P22, seven days after noise exposure, an increase in DPOAE thresholds was observed that was statistically significant only at 16 kHz (Mann-Whitney, df = 1, p = 0.01; Figure 5a). Yet, two months later, DPOAE thresholds were comparable to those of unexposed WT controls across all tested frequencies, suggesting that early noise exposure did not result in long-term changes in OHC function (Figure 5a).

In α9KO mice, DPOAE thresholds remained unchanged one day after noise exposure. However, seven days post-exposure, significant threshold elevations were observed at 8 and 16 kHz, but not at 22.65 kHz (Mann-Whitney, df = 1, p = 0.02 at 8 kHz and p = 0.02 at 16 kHz). After two months, recovery to unexposed levels occurred only at 8 kHz, while DPOAE thresholds remained elevated at 16 kHz (Mann-Whitney, df = 1, p = 0.003 at 16 kHz; Figure 5b). In contrast, mice with enhanced MOC function showed no changes in DPOAE thresholds at any post-exposure time point (Figure 5c), suggesting that noise exposure at hearing onset did not impair the cochlear amplifier function in these mice with enhanced OHC inhibition.

Ribbon synapse changes after early noise exposure in mice with different levels of α9α10

nAChR activity

Histological analysis was performed on cochleae dissected and fixed at P15 and P75 from

unexposed mice of all three genotypes, which exhibit varying levels of α9α10 nAChR activity, to

evaluate its effect of on the density and structure of IHC ribbon synapses. To visualize these

ribbon synapses, we performed immunostaining on whole-mount preparations of the organ of

Corti with antibodies against CtBP2-Ribeye, a presynaptic ribbon protein (Khimich et al., 2005),

and GluA2, an AMPA-type glutamate receptor expressed at the postsynaptic afferent terminal

(Matsubara et al., 1996; Liberman et al., 2011; Maison et al., 2013). Ribbon synapses were

identified by the colocalization of CtBP2 and GluA2 puncta at the base of the IHC, as shown in

representative confocal images of IHC synapses from apical whole mount organ of Corti from

unexposed WT, α9KO, and α9KI mice (Figure 6a) (Liberman et al., 2011). We quantified the

number of colocalized synaptic markers per IHC at both ages across three cochlear regions, each

representing a distinct frequency range: apex (~4–12 kHz), middle turn (~12–20 kHz), and base

(~22–40 kHz) (Figure 6b). At the low-frequency apical end, both unexposed α9KO and α9KI mice

showed a significant reduction in synaptic density compared to WT, with decreases of 33% and

31.18% at P15, and 30.64% and 27.94% at P75, respectively (Kruskal-Wallis, df = 2, p < 0.0001

for both groups; Figure 6b, left panel). In the middle region, synaptic density was significantly

reduced in both α9KO and α9KI mice compared to WT at P15 (by 28.3% and 35.33%,

respectively). However, at P75, only α9KI mice showed a significant reduction (35.97%) relative

to WT (Kruskal-Wallis, df = 2, p < 0.0001; Figure 6b, middle panel). Similarly, in the high-frequency

basal region, synaptic puncta were significantly reduced at P15 in both α9KO and α9KI mice (by

24.9% and 24.3%, respectively; Kruskal-Wallis, df = 2, p < 0.001 in α9KO and p < 0.01 in α9KI;

Figure 6b, right panel), whereas at P75, a significant reduction of 22.07 % was observed only in

α9KI mice (Kruskal-Wallis, df = 2, p = 0.02; Figure 6b, right panel). No significant changes in the

number of synaptic puncta were observed between P15 and P75 in WT, α9KO, or α9KI mice

across any cochlear region (Figure 6b), indicating that under unexposed conditions, alterations in

transient MOC inhibition affect the number of ribbon synapses early in development and persist

into maturity.

We further quantified the volume of CtBP2 puncta and GluA2 receptor patches within

colocalized synaptic markers at both P15 and P75 across the three groups of mice (Figure 6c, d).

In the apical region, both α9KO and α9KI mice showed reduced ribbon volumes compared to WT

at both ages (one way ANOVA followed by Holm-Sidak test, df = 2, p < 0.0001, for both groups

and ages; Figure 6c, left panel). In the middle cochlear region at P15, CtBP2 puncta were

enlarged in α9KO mice and reduced in α9KI mice compared to WT. By P75, only α9KI mice

continued to show a significant reduction relative to WT (one way ANOVA followed by Holm-Sidak

test, df = 2, p < 0.0001, for both groups and ages; Figure 6c, middle panel). In the basal region,

CtBP2 volumes were increased in α9KO mice and reduced in α9KI mice at both ages (one way

ANOVA followed by Holm-Sidak test, df = 2, p < 0.0001, for both groups and ages; Figure 6c,

right panel). GluA2 receptor patch size was significantly reduced in the apical region of α9KI mice

compared to WT at both P15 and P75, while no changes were observed in α9KO mice (one way

ANOVA followed by Holm-Sidak test, df = 2, p < 0.0001, for both ages; Figure 6d, left panel). In

the middle region, no differences in GluA2 size were observed at P15; however, by P75, both

α9KO and α9KI mice exhibited reduced GluA2 volumes compared to WT. (one way ANOVA

followed by Holm-Sidak test, df = 2, p < 0.05 in α9KO and p < 0.0001 in α9KI; Figure 6d, middle

panel). In the basal region, GluA2 puncta were significantly larger in α9KO mice compared to WT

at both P15 and P75 (one-way ANOVA followed by Holm-Sidak test, df = 2, p < 0.01 at P15 and

p < 0.0001 at P75; Figure 6d, right panel), while no differences were observed in α9KI mice.

These results suggest that inhibitory input from cholinergic efferent fibers plays a key role in

shaping synaptic density and preserving the morphology of pre- and postsynaptic elements

essential for proper auditory function.

The long-term impact of noise exposure at hearing onset on auditory nerve synapse

numbers is illustrated in Figure 7, on cochleae dissected and fixed at P75 from control and noise-

exposed mice with varying levels of α9α10 nAChR activity. Presynaptic, postsynaptic, and

colocalized synaptic puncta quantifications are shown for the apical, middle, and basal cochlear

regions (Figure 7, violin plots, right panels). In WT exposed ears, the number of CtBP2, GluA2,

and colocalized synaptic puncta significantly decreased at both the apical and basal cochlear

ends (Figure 7a). The decrease in CtBP2 puncta was more pronounced in the high-frequency

basal region, with a 37.2% decrease compared to controls (Mann-Whitney, df = 1, p < 0.0001 at

the apical and basal regions). Similarly, GluA2 postsynaptic receptors were reduced in both the

apical and basal turns in the cochlea of noise-exposed WT mice (Mann-Whitney, df = 1, p <

0.0001 at both cochlear regions). Putative ribbon synapse counts, defined by juxtaposed CtBP2-

and GluA2-positive puncta, showed a 34.21% reduction in the apical and a 32.56% reduction in

the basal turn following acoustic trauma (Mann-Whitney, df = 1, p < 0.0001 at both cochlear

regions) (Figure 7a, right panel). In α9KO mice, there was also a reduction in the number of

presynaptic, postsynaptic, and colocalized puncta after noise exposure, depending on cochlear

frequency/location (Figure 7b). Ribbon puncta in α9KO exposed ears were reduced by up to

14.73% of the control in the apical and 24.75% in the basal cochlear ends after noise exposure

(Mann-Whitney, df = 1, p = 0.001 and p = 0.002 at the apical and basal region, respectively).

GluA2-positive postsynaptic receptor patches decreased by 24.79% in the basal turn of noise-

exposed mice (Mann-Whitney, df = 1, p = 0.004). Following noise exposure, synaptic puncta were

reduced by 26.17% in the apical cochlear region and 25.87% in the basal region (Mann-Whitney,

df = 1, p = 0.002 and p < 0.0001 at the apical and basal region, respectively) (Figure 7b, right

panel). Interestingly, after two months post-exposure, α9KI ears exhibited no significant changes

in presynaptic ribbons, postsynaptic GluA2 receptor patches, or putative synapses across all

cochlear regions (Figure 7c). Although not statistically significant, a modest increase in synaptic

density (~19% at the basal region) was observed in α9KI tissue (Figure 7c, right panel).

Finally, we quantified the volume of ribbons and AMPA receptor patches (i.e., colocalized

CtBP2 and GluA2 puncta) in unexposed and exposed mice with different levels of MOC feedback

(Figure 8). In WT mice, noise exposure resulted in significantly larger ribbon volumes in the basal

end of the cochlea compared to unexposed mice, with no corresponding changes in AMPA

receptor patch size (t-test, df = 1, p = 0.01; Figure 8a). No changes were observed in the apical

or middle cochlear regions (Figure 8a). Conversely, α9KO mice showed reduced volumes of both

ribbons and AMPA receptor patches in the basal region following noise exposure, with no

changes in the apical or middle regions (t-test, df = 1, p = 0.002 for CtbP2 volume and p = 0.003

for GluA2 volume; Figure 8b). Notably, α9KI mice showed no morphological changes, suggesting

that enhanced α9α10 nAChR activity protects synapses from noise-induced alterations (Figure

8c).

Discussion

Auditory development is a gradual process that begins during embryonic stages and

continues through sensitive postnatal periods. While the detrimental effects of noise exposure in

adulthood are well documented, less is understood about its impact during development and the

role of the MOC system in this context. Here, we show that varying levels of MOC inhibition impact

the timing of hearing onset and are crucial for proper development of IHC ribbon synapses. Early

noise exposure caused lasting auditory deficits in WT, highlighting the greater developmental

vulnerability. In contrast, enhanced cholinergic activity prevented damage, underscoring the

MOC's protective role in early development.

Before hearing begins, MOC synapses modulate IHCs' activity, shaping the

developmental increase in Ca^{2+} sensitivity of glutamate release at IHC ribbon synapses

(Glowatzki and Fuchs, 2000; Johnson et al., 2011, 2013). Disruption of this input, via genetic

modification or neonatal olivocochlear bundle transection, impairs cochlear and auditory circuit

maturation (Walsh et al., 1998; Frank et al., 2009; Hirtz et al., 2011; Johnson et al., 2013; Clause

et al., 2014; Di Guilmi et al., 2019; Babola et al., 2021). Our findings demonstrate that α9α10

nAChR activity regulates hearing onset and synaptic integrity. Enhanced α9α10 nAChR activity

in α9KI mice accelerates the onset of hearing, with sound-evoked potentials detected earlier than

in WT mice. In α9KO mice, which lack MOC input, hearing onset was delayed, with elevated ABR

thresholds and diminished wave 1 amplitudes. This aligns with prior reports using Ca²⁺ imaging

showing diminished sensitivity in α9KO inferior colliculus at hearing onset (Wang et al., 2021).

Importantly, both enhanced and absent MOC inhibition disrupted IHC ribbon synapse

development, with reduced synapse density and altered volumes at P15 and P75. Despite similar

synaptic reductions, their distinct ABR phenotypes suggest differential effects on auditory nerve

fiber subtypes. The earlier hearing onset in α9KI mice could reflect preservation or accelerated

maturation of low-threshold, high-spontaneous-rate fibers, while their disruption in α9KO mice

may contribute to delayed onset. Morphologically, α9KI mice showed reduced ribbon and GluA2

patch volumes across all cochlear turns, whereas α9KO mice exhibited region-specific changes:

smaller ribbons in apical regions with unchanged GluA2 patches, and enlarged pre- and post-

synaptic components in middle/basal turns. These ribbon alterations may reflect disruption of

Ca²⁺-dependent ribbon fusion or stabilization, impairing synaptic refinement. Ribbon size and

maturation is tightly regulated by presynaptic Ca^{2+} influx (Frank et al., 2009; Sheets et al., 2012;

Wong et al., 2019; Özçete and Moser, 2021). Fusion of ribbon precursors during this period is

thought to support sustained release at mature active zones (Michanski et al., 2019). In

developing IHCs, intracellular Ca²⁺ dynamics are shaped by voltage-gated Ca²⁺ channels and by

α9α10 nAChRs, which are highly permeable to Ca²⁺ (Weisstaub et al., 2002; Gomez-Casati et

al., 2005; Moglie et al., 2018). In α9KO mice, the absence of α9α10-mediated Ca²⁺ influx leads to

increased action potential frequency and disrupted exocytosis (Johnson et al., 2013). In α9KI

mice, the L9'T mutation reduces receptor desensitization, likely prolonging Ca²⁺ entry and altering

IHC firing during the prehearing period (Taranda et al., 2009). These results highlight the MOC

synapse's role in coordinating afferent and efferent activity for normal auditory development.

Auditory sensitivity in vertebrates matures from high thresholds and poor tuning at hearing

onset to precise, low-threshold processing (Song et al., 2006). In WT mice, ABR and DPOAE

data from P12 to P75 showed this progression. At P14, thresholds were elevated and waveforms

immature, but by P22, thresholds and wave 1 amplitudes reached mature levels. Thus, at P15,

the time of noise exposure, auditory maturation was still incomplete. A key finding of this study is

that noise exposure during early development leads to more severe and lasting effects than those

reported at later stages. In WT mice, noise exposure at P15 caused lasting threshold shifts and

35% synapse loss, while the same exposure at P21 led to only transient threshold shifts and

milder synaptopathy (Boero et al., 2018). This underscores the heightened vulnerability at hearing

onset, likely due to both noise-induced damage and disruption of ongoing synaptic or hair cell

maturation. Similar age-dependent vulnerability has been reported in other juvenile models

(Fernandez et al., 2015; Jensen et al., 2015; Schrode et al., 2022; Lu et al., 2024). While the

mechanisms behind this heightened sensitivity remain unclear, it may arise from ongoing

maturation of the MOC system and central auditory pathways, including myelination and afferent

input. Developing interactions between the auditory nerve and LOC system may further increase

susceptibility to noise exposure.

In WT mice, DPOAE thresholds were elevated seven days post-exposure, reaching

statistical significance at 16 kHz, but returned to baseline by two months, suggesting preserved

or repaired OHC function. Whether this reflects true recovery, delayed maturation, or

compensation remains to be determined. In contrast, persistent ABR threshold elevation and

reduced wave 1 amplitudes suggest lasting afferent damage, likely due to cochlear synaptopathy affecting both high- and low-threshold auditory nerve fibers (Reijntjes et al., 2025). This is the first report of sustained ABR threshold elevation following moderate noise exposure, similar to lasting high-frequency shifts and presynaptic loss seen in P14 rats and noise-sensitive P15 C57BL/6J mice (Davis et al., 2003) exposed to high-intensity sound (Rybalko et al., 2015; Boero et al., 2021). Noise exposure caused significant ribbon synapse loss in apical and basal cochlear regions of WT mice, correlating with reduced ABR wave 1 amplitudes; however, despite functional deficits at 16 kHz, synaptic loss in the mid-frequency region was minimal—reflecting a known mismatch due to cochlear place-frequency map shifts after noise exposure (Müller and Smolders, 2005). In addition, IHCs of noise-exposed mice tend to have larger ribbons, especially in the basal turn. Similar post-trauma ribbon enlargement was observed in RIBEYE-tagRFP mice (Mohamad et al., 2024) and via volume EM (Lu et al., 2024), likely reflecting the fusion of individual ribbons. Multi-ribbon active zones and ribbon unanchoring from synaptic tethers have been observed in synaptopathic ears (Moverman et al., 2023; Lu et al., 2024; Mohamad et al., 2024). Though their function is unclear, these structures persist into adulthood (Michanski et al., 2019; Hua et al., 2021) and are a hallmark of cochlear aging (Stamataki et al., 2006), with larger ribbons being associated with increased exocytotic activity (Jeng et al., 2020). Notably, noise exposure has also been associated with enhanced exocytosis in IHCs, along with an increase in tone-burst responses in auditory nerve fibers (Boero et al., 2021; Suthakar and Liberman, 2021).

In α9KO mice, ABR and DPOAE thresholds were elevated one week post-exposure at P15, consistent with prior findings at P21 (Boero et al., 2018). However, unlike WT, ABR thresholds in α9KOs nearly fully recovered by two months. DPOAE thresholds also normalized except at 16 kHz, matching a slight, non-significant ABR threshold elevation at that frequency. The recovery of ABR thresholds observed in α9KO mice, unlike the persistent elevation in WT mice, may stem from reduced auditory sensitivity at hearing onset in α9KO mice, leading to less

effective noise exposure and milder impact. Despite threshold recovery, reduced wave 1

amplitudes and synapse counts suggest lasting damage to high-threshold, low-spontaneous-rate

fibers (Furman et al., 2013; Reijntjes et al., 2025). Additionally, noise-exposed α9KO ears showed

reduced sizes of pre- and postsynaptic puncta, indicating structural synaptic alterations. At P16

(one day post- exposure), ABR and DPOAE thresholds in both WT and α9KO mice were slightly

reduced compared to P15 and aligned with those of unexposed controls, suggesting that noise

had no immediate impact on auditory maturation at this stage. In contrast, Boero et al. (2021)

reported lasting damage in C57BL/6J mice one day after 120 dB SPL at P15, highlighting that

early noise effects depend on complex interactions between factors such as intensity, duration,

and genetic background, and differ from outcomes seen with noise exposure at mature stages.

588   Unlike WT and α9KO mice, α9KI mice showed no threshold shifts or synaptopathy after

noise, consistent with prior reports in mature α9KI mice (Taranda et al., 2009; Boero et al., 2018)

and recent gene therapy studies supporting protective effects of enhanced α9α10 nAChR activity

(Zhang et al., 2023; Slika et al., 2025). The protective role of the MOC system has been

recognized for decades: efferent stimulation reduces threshold shifts (Rajan, 1988; Reiter and

Liberman, 1995), while surgical de-efferentation increases vulnerability (Kujawa and Liberman,

1997). Overexpressing wild-type α9 also confers greater resistance to noise (Maison et al., 2002).

In vivo studies show that ACh evokes fast (~100ms) and slow (~10s) MOC effects on cochlear

responses (Sridhar et al., 1995, 1997; Cooper and Guinan, 2003). Fast suppression involves SK2-

mediated OHC hyperpolarization, while slow suppression likely reflects Ca^{2+} signaling and

increased OHC stiffness, both reducing cochlear vibrations (Cooper and Guinan, 2003). Evidence

suggests that efferent protection likely relies on slow, Ca^{2+}-dependent mechanisms that reduce

acoustic injury by dampening cochlear vibrations or altering OHC function via α9α10 receptor-

mediated Ca^{2+} influx (Reiter and Liberman, 1995; Maison et al., 2007). Though MOC-OHC

innervation is still maturing during early development (Simmons et al., 1996), our findings show

that enhanced α9α10 activity already confers robust protection against acoustic trauma.

Figures

Figure 1: Auditory function in WT, α9KO, and α9KI mice across different postnatal ages.
(**a**) Representative ABR waveforms from WT (black), α9KI (red), and α9KO (blue) mice recorded at P14 in response to 16 kHz tone bursts at increasing sound pressure levels (dB SPL). Thresholds, defined by the presence of wave 1, are marked by green dashed traces. Waves 1, 2 and 3 are indicated by gray arrows. The scale bar applies to all three recordings. (**b**) Mean ABR thresholds from P12 to P75 in WT (P12 to P14 n = 9 and P15 to P75 n = 14), α9KO (P12 to P14 n = 7 and P15 to P75 n = 11), and α9KI (P12 to P14 n = 9 and P15 to P75 n = 12) mice at 11.33, 16, and 22.65 kHz. ABR waveforms first became detectable at P13 in α9KI mice, at P14 in WT mice, and between P14 and P15 in α9KO mice. (**c**) Mean DPOAE thresholds from P12 to P75 in WT (P12 to P14 n = 9 and P15 to P75 n = 14), α9KO (P12 to P14 n = 7 and P15 to P75 n = 11) and α9KI (P12 to P14 n = 9 and P15 to P75 n = 12) mice at 11.33, 16 and 22.65 kHz. DPOAE thresholds showed the same trend as ABR only at 11.33 kHz. Group means ± SEM are shown. Red asterisks represent the statistical significance of α9KI compared with WT**,** and blue asterisks represent the statistical significance of α9KO compared with WT (Kruskal-Wallis followed by post hoc Dunn test, *p < 0.05).

Figure 2: Suprathreshold ABR Wave1 amplitudes and latencies in WT, α9KO, and α9KI mice during postnatal development. (**a**) Representative examples of ABR waveforms recorded at 16 kHz, 80 dB SPL, from WT (black), α9KI (red), and α9KO (blue) mice. Waves 1, 2 and 3 are indicated by gray arrows. Each column illustrates waveform maturation across developmental stages (P12–P16, P22, P75) for each genotype. At P13, α9KI mice exhibit measurable responses, while distinct ABR waves emerge in WT and α9KO mice by P14. Waveform morphology stabilizes by P22 and remains consistent through P75. Scale bars apply to all traces. (**b**) ABR wave 1 amplitudes (µV) at 80 dB SPL for 11.33, 16, and 22.65 kHz across developmental ages in WT, α9KO, and α9KI mice. Amplitudes were measured as the difference between the maximum and minimum peaks. No values are shown for P12 due to the absence of discernible waves. (**c**) ABR wave 1 latencies (ms) at the same frequencies. A developmental decrease in latency is observed between P14 and P15 in all genotypes, though not statistically significant. The number of animals per genotype is the same as in Figure 1. Data represent mean ± SEM. Red asterisks denote significant differences between α9KI and WT; blue asterisks indicate significant differences between α9KO and WT (one-way ANOVA followed by Holm-Sidak test, *p < 0.05; **p < 0.01).

Figure 3: ABR wave 1 threshold measurements in control and noise-exposed WT, α9KO, and α9KI mice at different postnatal ages. ABR thresholds before and after exposure to noise at P15 in WT ($n_{control}$ = 14 and $n_{Exposed}$ = 12) (**a**), α9KO ($n_{control}$ = 11 and $n_{Exposed}$ = 11) (**b**), and α9KI ($n_{control}$ = 12 and $n_{Exposed}$ = 12) (**c**) mice at 8, 16 and 22.65 kHz. WT mice showed a significant increase in ABR thresholds 1 week and 2 months after exposure to noise, indicating a permanent threshold elevation. A recovery of ABR thresholds was observed in α9KO mice 2 months after exposure to noise. α9KI mice did not present any changes in ABR thresholds at any time after acoustic trauma. Each point represents the mean ± SEM. Asterisks represent the statistical significance (Mann-Whitney test, *p < 0.05; **p < 0.01).

Figure 4: Suprathreshold response amplitude for ABR wave 1 after P15 noise exposure in WT, α9KO, and α9KI mice at different postnatal ages compared to unexposed controls. ABR wave 1 amplitudes (in μV) at 80 dB SPL at 8, 16 and 22.65 kHz in WT ($n_{Control}$ = 14 and $n_{Exposed}$ = 12) (**a**), α9KO ($n_{Control}$ = 11 and $n_{Exposed}$ = 11) (**b**), and α9KI ($n_{Control}$ = 12 and $n_{Exposed}$ = 12) (**c**) mice at the same time points shown in Figure 3. Each panel inset displays representative ABR waveforms recorded at 80 dB SPL for the corresponding frequency, shown for all groups at P75. Scale bars (1 μV vertical, 1 ms horizontal) apply to all traces. WT and α9KO mice exhibited reduced ABR wave 1 amplitudes following noise exposure, whereas wave 1 amplitudes in α9KI mice remained unaffected. Data are presented as mean ± SEM. Statistical significance is indicated by asterisks (Mann-Whitney test, *p < 0.05; **p < 0.01).

Figure 5: **Assessment of OHC functional integrity at different postnatal ages following noise exposure at P15 in WT, α9KO, and α9KI mice.** DPOAE thresholds are shown for WT ($n_{Control}$ = 14 and $n_{Exposed}$ = 12) (**a**), α9KO ($n_{Control}$ = 11 and $n_{Exposed}$ = 11) (**b**), and α9KI ($n_{Control}$ = 12 and $n_{Exposed}$ = 12) (**c**) at 8, 16 and 22.65 kHz. WT mice exhibited a transient increase in DPOAE thresholds 1 week post-exposure at 16 kHz. α9KO mice showed increased thresholds at 8 and 16 kHz 1 week post-exposure, with sustained elevation at 16 kHz up to 2 months. α9KI mice demonstrated no changes in DPOAE thresholds after noise exposure. Data are presented as mean ± SEM. Statistical significance is indicated by asterisks (Mann-Whitney test, *p < 0.05; **p < 0.01).

Figure 6: Analysis of IHC ribbon synapses in unexposed WT, α9KO, and α9KI mice at P15 and P75. (**a**) Representative confocal images of IHC synapses at the apex of the cochlea from cochleae immunolabeled for presynaptic ribbons (CtBP2-red) and postsynaptic receptor patches (GluA2-green) for WT (left), α9KO (middle) and α9KI (right) at P15 and P75. CtBP2 antibody also weakly stains IHC nuclei. Scale bar, 7 μm. (**b**) Quantification of putative ribbon synapses per IHC (i.e., colocalized CtBP2 and GluA2 puncta). WT P15: n_{Apex} = 182 IHC, n_{Middle} = 248, n_{Base} = 68 (6 animals); WT P75: n_{Apex} = 80 IHC, n_{Middle} = 75, n_{Base} = 60 IHC (14 animals); α9KO P15: n_{Apex} = 201 IHC, n_{Middle} = 230, n_{Base} = 164 (5 animals); α9KO P75: n_{Apex} = 100 IHC, n_{Middle} = 84, n_{Base} = 118 (11 animals); α9KI P15: n_{Apex} = 155 IHC, n_{Middle} = 284, n_{Base} = 85 (5 animals) and α9KI P75: n_{Apex} = 178 IHC, n_{Middle} = 319, n_{Base} = 198 (12 animals). (**c**) Volumes of fluorescence staining for ribbons (CtBP2 size) and (**d**) AMPA receptors (GluA2 size) for each of the synaptic pairs (WT P15: n_{Apex} = 684 synaptic pairs, n_{Middle} = 729, n_{Base} = 380; WT P75: n_{Apex} = 450, n_{Middle} = 509, n_{Base} = 203; α9KO P15: n_{Apex} = 845, n_{Middle} = 800, n_{Base} = 735; α9KO P75: n_{Apex} = 367, n_{Middle} = 187, n_{Base} = 221; α9KI P15: n_{Apex} = 428, n_{Middle} = 484, n_{Base} = 232 and α9KI P75: n_{Apex} = 247, n_{Middle} = 270, n_{Base} = 240). Violin plots indicate median (solid line) and interquartile range (IQR, dashed lines). Asterisks denote statistical significance. The number of synapses was analyzed using the Kruskal–Wallis

test followed by Dunn's post-test, while synaptic volumes were analyzed using one-way ANOVA
followed by the Holm–Sidak test.; *p < 0.05; **p < 0.01; ***p < 0.001, ****p < 0.0001).

Figure 7: Analysis of the degree of IHC synaptopathy two months after exposure to noise
at P15. Left, Representative confocal images of IHC synapses at the apex of the cochlea from
cochleae immunolabeled for presynaptic ribbons (CtBP2-red), postsynaptic receptor patches
(GluA2-green) and IHCs (Myosin VIIa-blue). CtBP2 antibody also weakly stains IHC nuclei. Scale
bar, 10μm. Right, Quantitative data obtained from WT, α9KO, and α9KI mice at P75. For each
IHC, we analyzed the number of CtBP2 puncta, postsynaptic GluA2 receptor patches, and
putative ribbon synapses. (**a**) In traumatized WT mice, there was a reduction in the number of
CtBP2 puncta, GluA2 receptor patches, and putative synapse, at the apical and basal region of
the cochlea (WT Control: n_{Apex} = 80 IHC, n_{Middle} = 75, n_{Base}= 60; WT Exposed: n_{Apex} = 82 IHC, n_{Middle}
= 90, n_{Base} = 50). (**b**) In traumatized α9KO mice, there was a reduction in the number of
prelocalized, postlocalized, and colocalized puncta depending on cochlear frequency/location
(α9KO Control: n_{Apex} = 100 IHC, n_{Middle} = 84, n_{Base} = 118; α9KO Exposed: n_{Apex} = 234 IHC, n_{Middle} =
60, n_{Base} = 90). (**c**) In traumatized α9KI mice, no difference was found in the number of presynaptic
ribbons, postsynaptic AMPA receptors, and colocalization puncta for the three regions of the
cochlea (α9KI Control: n_{Apex} = 178 IHC, n_{Middle} = 319, n_{Base} = 198; α9KI Exposed: n_{Apex} = 196 IHC,
n_{Middle} = 145, n_{Base} = 80). Violin plots indicate median (solid line) and IQR (dashed lines). Asterisks
denote the statistical significance (Mann Whitney test, **p < 0.01; ***p < 0.001, ****p < 0.0001).

Figure 8: Volumes of presynaptic ribbons and postsynaptic receptor patches for each
synaptic pair from control and noise-exposed WT, α9KO and α9KI mice. Violin plots
comparing the volumes of fluorescence staining for ribbons (CtBP2 size) and AMPA receptors
(GluA2 size) between control and noise-exposed mice. The quantification was made in the apex,
middle and base regions of the cochlea from WT (**a**) (Control: n_{Apex} = 450 synaptic pairs, n_{Middle} =
509, n_{Base} = 203; Exposed: n_{Apex} = 497 synaptic pairs, n_{Middle} = 175, n_{Base} = 250); α9KO (**b**) (Control:
n_{Apex} = 367 synaptic pairs, n_{Middle} = 187, n_{Base} = 221; Exposed: n_{Apex} = 207 synaptic pairs, n_{Middle} =
300, n_{Base} = 145) and α9KI (**c**) (Control: n_{Apex} = 247 synaptic pairs, n_{Middle} = 270, n_{Base} = 240;
Exposed: n_{Apex} = 280 synaptic pairs, n_{Middle} = 310, n_{Base} = 360) mice. Violin plots indicate median
(solid line) and IQR (dashed lines). Asterisks denote the statistical significance (t-test; *p < 0.05;
***p < 0.001).

References

Antoli-Candela FJ, Kiang N (1978) Unit activity underlying the N1 potential. In: Evoked electrical activity in the auditory nervous system (Naunton R, Fernandez C, eds), pp 165–191.

Babola TA, Li S, Wang Z, Kersbergen CJ, Elgoyhen AB, Coate TM, Bergles DE (2021) Purinergic Signaling Controls Spontaneous Activity in the Auditory System throughout Early Development. J Neurosci 41:594–612.

Boero LE, Castagna VC, Di Guilmi MN, Goutman JD, Elgoyhen AB, Gómez-Casati ME (2018) Enhancement of the medial olivocochlear system prevents hidden hearing loss. The Journal of Neuroscience 38:7440–7451 Available at: http://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.0363-18.2018.

Boero LE, Payne S, Gómez-Casati ME, Rutherford MA, Goutman JD (2021) Noise Exposure Potentiates Exocytosis From Cochlear Inner Hair Cells. Front Synaptic Neurosci 13:740368.

Buchwald JS, Huang CM (1975) Far-field acoustic response: Origins in the cat. Science 189:382–384.

Caminos C, Vale E, Lujan R, Martinez-Galan JR, Juiz JM (2005) Developmental regulation and adult maintenance of potassium channel proteins (Kv 1.1 and Kv 1.2) in the cochlear nucleus of the rat. Brain research 1056 Available at: https://pubmed.ncbi.nlm.nih.gov/16122713/ [Accessed October 16, 2024].

Chang EF, Merzenich MM (2003) Environmental noise retards auditory cortical development. Science 300:498–502.

Clause A, Kim G, Sonntag M, Weisz CJC, Vetter DE, Rűbsamen R, Kandler K (2014) The precise temporal pattern of prehearing spontaneous activity is necessary for tonotopic map refinement. Neuron 82:822–835.

Cooper NP, Guinan JJ (2003) Separate mechanical processes underlie fast and slow effects of medial olivocochlear efferent activity. J Physiol 548:307–312.

Davis RR, Kozel P, Erway LC (2003) Genetic influences in individual susceptibility to noise: a review. Noise Health 5:19–28.

De Faveri F, Ceriani F, Marcotti W (2025) In vivo spontaneous Ca2+ activity in the pre-hearing mammalian cochlea. Nat Commun 16:29 Available at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC11695946/ [Accessed April 4, 2025].

Di Guilmi MN, Boero LE, Castagna VC, Rodríguez-Contreras A, Wedemeyer C, Gómez-Casati ME, Elgoyhen AB (2019) Strengthening of the efferent olivocochlear system leads to synaptic dysfunction and tonotopy disruption of a central auditory nucleus. The Journal of Neuroscience:2536–18.

Dorrn AL, Yuan K, Barker AJ, Schreiner CE, Froemke RC (2010) Developmental sensory experience balances cortical excitation and inhibition. Nature 465:932–936.

Dulon D, Lenoir M (1996) Cholinergic responses in developing outer hair cells of the rat cochlea.
Eur J Neurosci 8:1945–1952.

Ernfors P, Kucera J, Lee KF, Loring J, Jaenisch R (1995) Studies on the physiological role of
brain-derived neurotrophic factor and neurotrophin-3 in knockout mice. Int J Dev Biol
39:799–807   Available   at:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation
&list_uids=8645564.

Fernandez KA, Jeffers PWC, Lall K, Liberman MC, Kujawa SG (2015) Aging after noise
exposure : Acceleration of cochlear synaptopathy in " recovered " ears. The Journal of
Neuroscience 35:7509–7520.

Frank T, Khimich D, Neef A, Moser T (2009) Mechanisms contributing to synaptic Ca2+ signals
and their heterogeneity in hair cells. Proc Natl Acad Sci U S A 106:4483–4488.

Friauf E, Lohmann C (1999) Development of auditory brainstem circuitry. Activity-dependent and
activity-independent processes. Cell Tissue Res 297:187–195.

Froemke RC, Jones BJ (2011) Development of auditory cortical synaptic receptive fields.
Neurosci Biobehav Rev 35:2105–2113.

Furman AC, Kujawa SG, Liberman MC (2013) Noise-induced cochlear neuropathy is selective for
fibers with low spontaneous rates. Journal of Neurophysiology 110:577–586.

Gilels F, Paquette ST, Zhang J, Rahman I, White PM (2013) Mutation of foxo3 causes adult onset
auditory neuropathy and alters cochlear synapse architecture in mice. Journal of
Neuroscience 33:18409–18424.

Glowatzki E, Fuchs PA (2000) Cholinergic synaptic inhibition of inner hair cells in the neonatal
mammalian cochlea. Science 288:2366–2368.

Gomez-Casati ME, Fuchs PA, Elgoyhen AB and, Katz E (2005) Biophysical and pharmacological
characterization of nicotinic cholinergic receptors in rat cochlear inner hair cells. Journal
of   Physiology   566:103–118   Available   at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1464719&tool=pmcentrez&re
ndertype=abstract [Accessed August 10, 2013].

Goutman JD, Fuchs PA, Glowatzki E (2005) Facilitating efferent inhibition of inner hair cells in the
cochlea of the neonatal rat. Journal of Physiology 566:49–59 Available at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1464729&tool=pmcentrez&re
ndertype=abstract [Accessed August 31, 2013].

He DZ, Dallos P (1999) Development of acetylcholine-induced responses in neonatal gerbil outer
hair cells. J Neurophysiol 81:1162–1170.

Hirtz JJ, Boesen M, Braun N, Deitmer JW, Kramer F, Lohr C, Müller B, Nothwang HG, Striessnig
J, Löhrke S, Friauf E (2011) Cav1.3 calcium channels are required for normal development
of the auditory brainstem. J Neurosci 31:8280–8294.

Hua Y, Ding X, Wang H, Wang F, Lu Y, Neef J, Gao Y, Moser T, Wu H (2021) Electron Microscopic Reconstruction of Neural Circuitry in the Cochlea. Cell Rep 34:108551.

Jeng JY, Ceriani F, Olt J, Brown SDM, Holley MC, Bowl MR, Johnson SL, Marcotti W (2020) Pathophysiological changes in inner hair cell ribbon synapses in the ageing mammalian cochlea. Journal of Physiology 598:4339–4355.

Jensen JB, Lysaght AC, Liberman MC, Qvortrup K, Stankovic KM (2015) Immediate and delayed cochlear neuropathy after noise exposure in pubescent mice. PLoS ONE 10:1–17.

Jing Z, Rutherford MA, Takago H, Frank T, Fejtova A, Khimich D, Moser T, Strenzke N (2013) Disruption of the presynaptic cytomatrix protein bassoon degrades ribbon anchorage, multiquantal release, and sound encoding at the hair cell afferent synapse. Journal of Neuroscience 33:4456–4467.

Johnson SL, Eckrich T, Kuhn S, Zampini V, Franz C, Ranatunga KM, Roberts TP, Masetto S, Knipper M, Kros CJ, Marcotti W (2011) Position-dependent patterning of spontaneous action potentials in immature cochlear inner hair cells. Nat Neurosci 14:711–717.

Johnson SL, Wedemeyer C, Vetter DE, Adachi R, Holley MC, Elgoyhen AB, Marcotti W (2013) Cholinergic efferent synaptic transmission regulates the maturation of auditory hair cell ribbon synapses. Open Biology 3.

Karplus M, Lee C, Cashore WJ, Oh W (1988) The effects of brain bilirubin deposition on auditory brain stem evoked responses in rats. Early Hum Dev 16:185–194.

Katz E, Elgoyhen AB, Gómez-Casati ME, Knipper M, Vetter DE, Fuchs P a, Glowatzki E, Gomez-Casati ME, Knipper M, Vetter DE, Fuchs P a, Glowatzki E, Bele A, Knipper M, Vetter DE, Fuchs P a (2004) Developmental regulation of nicotinic synapses on cochlear inner hair cells. J Neurosci 24:7814–7820 Available at: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15356192 [Accessed August 31, 2013].

Keuroghlian AS, Knudsen EI (2007) Adaptive auditory plasticity in developing and adult animals. Prog Neurobiol 82:109–121.

Khimich D, Nouvian R, Pujol R, Tom Dieck S, Egner A, Gundelfinger ED, Moser T (2005) Hair cell synaptic ribbons are essential for synchronous auditory signalling. Nature 434:889–894 Available at: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15829963.

Kim SE, Turkington K, Kushmerick C, Kim JH (2013) Central dysmyelination reduces the temporal fidelity of synaptic transmission and the reliability of postsynaptic firing during high-frequency stimulation. J Neurophysiol 110:1621–1630.

Knipper M, Zimmermann U, Rohbock K, Kopschall I, Zenner HP (1995) Synaptophysin and GAP-43 proteins in efferent fibers of the inner ear during postnatal development. Brain Res Dev Brain Res 89:73–86 Available at: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=8575095.

Kujawa SG, Liberman MC (1997) Conditioning-related protection from acoustic injury: Effects of
chronic deefferentation and sham surgery. Journal of Neurophysiology 78:3095–3106.

Kujawa SG, Liberman MC (2009) Adding insult to injury: cochlear nerve degeneration after
"temporary" noise-induced hearing loss. J Neurosci 29:14077–14085 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/19906956%5Cnhttp://www.pubmedcentral.nih.gov/
articlerender.fcgi?artid=PMC2812055.

Lauer AM, May BJ, Ay BRJM, Lauer AM, May BJ (2011) The medial olivocochlear system
attenuates the developmental impact of early noise exposure. JARO - Journal of the
Association for Research in Otolaryngology 12:329–343 Available at:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation
&list_uids=21347798.

Leake PA, Hradek GT, Chair L, Snyder RL (2006) Neonatal deafness results in degraded
topographic specificity of auditory nerve projections to the cochlear nucleus in cats. J
Comp Neurol 497:13–31.

Liberman LD, Wang H, Liberman MC (2011) Opposing gradients of ribbon size and AMPA
receptor expression underlie sensitivity differences among cochlear-nerve/hair-cell
synapses. The Journal of neuroscience : the official journal of the Society for
Neuroscience   31:801–808   Available   at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3290333&tool=pmcentrez&re
ndertype=abstract [Accessed August 31, 2013].

Liberman MC (1991) The olivocochlear efferent bundle and susceptibility of the inner ear to
acoustic   injury.   J   Neurophysiol   65:123–132   Available   at:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation
&list_uids=1999726.

Lippe WR (1994) Rhythmic spontaneous activity in the developing avian auditory system. J
Neurosci   14:1486–1495   Available   at:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation
&list_uids=8126550.

Lu Y, Liu J, Li B, Wang H, Wang F, Wang S, Wu H, Han H, Hua Y (2024) Spatial patterns of
noise-induced inner hair cell ribbon loss in the mouse mid-cochlea. iScience 27:108825.

Maison SF, Liberman MC (2000) Predicting Vulnerability to Acoustic Injury with a Noninvasive
Assay of Olivocochlear Reflex Strength. The Journal of neuroscience 20:4701–4707.

Maison SF, Luebke AE, Liberman MC, Zuo J (2002) Efferent protection from acoustic injury is
mediated via alpha9 nicotinic acetylcholine receptors on outer hair cells. J Neurosci
22:10838-46.

Maison SF, Parker LL, Young L, Adelman JP, Zuo J, Liberman MC (2007) Overexpression of SK2
Channels Enhances Efferent Suppression of Cochlear Responses without Enhancing
Noise   Resistance.   Journal   of   neurophysiology   97:2930   Available   at:
https://pmc.ncbi.nlm.nih.gov/articles/PMC1850627/ [Accessed June 5, 2025].

Maison SF, Usubuchi H, Liberman MC (2013) Efferent feedback minimizes cochlear neuropathy
from moderate noise exposure. Journal of Neuroscience 33:5542–5552.

Marcotti W, Johnson SL, Kros CJ (2004) A transiently expressed SK current sustains and
modulates action potential activity in immature mouse inner hair cells. J Physiol Available
at:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation
&list_uids=15331671.

Marcotti W, Johnson SL, Rusch A, Kros CJ (2003) Sodium and calcium currents shape action
potentials in immature mouse inner hair cells. J Physiol 552:743–761 Available at:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation
&list_uids=12937295.

Matsubara A, Laake JH, Davanger S, Usami S, Ottersen OP (1996) Organization of AMPA
receptor subunits at a glutamate synapse: a quantitative immunogold analysis of hair cell
synapses in the rat organ of Corti. The Journal of Neuroscience 16:4457–4467.

Melcher J, Kiang NY (1996) Generators of the brainstem auditory evoked potential in cat. III:
Identified cell populations. Hearing research 93:52–71 Available at:
https://pubmed.ncbi.nlm.nih.gov/8735068/ [Accessed November 1, 2021].

Michanski S, Smaluch K, Maria Steyer A, Chakrabarti R, Setz C, Oestreicher D, Fischer C, Möbius
W, Moser T, Vogl C, Wichmann C (2019) Mapping developmental maturation of inner hair
cell ribbon synapses in the apical mouse cochlea. Proceedings of the National Academy
of Sciences of the United States of America 116:6415–6424.

Moglie MJ, Fuchs PA, Elgoyhen AB, Goutman JD (2018) Compartmentalization of antagonistic
Ca2+ signals in developing cochlear hair cells. Proc Natl Acad Sci U S A 115:E2095–
E2104.

Mohamad NI, Santra P, Park Y, Matthews IR, Taketa E, Chan DK (2024) Synaptic ribbon
dynamics after noise exposure in the hearing cochlea. Commun Biol 7:421.

Mostafapour SP, Cochran SL, Del Puerto NM, Rubel EW (2000) Patterns of cell death in mouse
anteroventral cochlear nucleus neurons after unilateral cochlea removal. J Comp Neurol
426:561–571.

Moverman DJ, Liberman LD, Kraemer S, Corfas G, Liberman MC (2023) Ultrastructure of noise-
induced cochlear synaptopathy. Sci Rep 13:19456.

Müller M M, von Hunerbein K, Hoidis S, Smolders JW (2005) A physiological place-frequency
map of the cochlea in the CBA/J mouse. Hear Res 202:63–73 Available at:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation
&list_uids=15811700.

Müller M, Smolders JW (2005) Shift in the cochlear place-frequency map after noise damage in
the mouse. NeuroReport 16:1183–1187 Available at:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation
&list_uids=16012345.

Ollion J, Cochennec J, Loll F, Escudé C, Boudier T (2013) TANGO: A generic tool for high-
throughput 3D image analysis for studying nuclear organization. Bioinformatics 29:1840–
1841.

Özçete ÖD, Moser T (2021) A sensory cell diversifies its output by varying Ca2+ influx-release
coupling among active zones. EMBO J 40:e106010.

Paquette ST, Gilels F, White PM (2016) Noise exposure modulates cochlear inner hair cell ribbon
volumes, correlating with changes in auditory measures in the FVB/nJ mouse. Scientific
Reports 6:1–13 Available at: http://dx.doi.org/10.1038/srep25056.

Pujol R (1985) Morphology, synaptology and electrophysiology of the developing cochlea. Acta
Otolaryngol Suppl 421:5-9.

Pujol R, Carlier E, Devigne C (1978) Different patterns of cochlear innervation during the
development of the kitten. J Comp Neurol 177:529-36.

Rajan R (1988) Effect of electrical stimulation of the crossed olivocochlear bundle on temporary
threshold shifts in auditory sensitivity. I . Dependence on electrical stimulation parameters.
J Neurophysiol 60:549–568.

Reijntjes DOJ, Burke K, Paul S, Mueller U, Glowatzki E, Lauer AM (2025) Increased vulnerability
to noise exposure of low spontaneous rate type 1C spiral ganglion neuron synapses with
inner hair cells (Pre-Print). bioRxiv:2025.05.19.654865.

Reiter ER, Liberman MC (1995) Efferent-mediated protection from acoustic overexposure:
relation to slow effects of olivocochlear stimulation. Journal of neurophysiology 73:506–
514.

Robles L, Ruggero MA (2001) Mechanics of the mammalian cochlea. Physiological reviews
81:1305–1352.

Roux I, Wersinger E, McIntosh JM, Fuchs P a, Glowatzki E (2011) Onset of cholinergic efferent
synaptic function in sensory hair cells of the rat cochlea. The Journal of neuroscience : the
official journal of the Society for Neuroscience 31:15092–15101 Available at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3213862&tool=pmcentrez&re
ndertype=abstract [Accessed August 10, 2013].

Rybalko N, Chumak T, Bureš Z, Popelář J, Šuta D, Syka J (2015) Development of the acoustic
startle response in rats and its change after early acoustic trauma. Behav Brain Res
286:212–221.

Sage D, Donati L, Soulez F, Fortun D, Schmit G, Seitz A, Guiet R, Vonesch C, Unser M (2017)
DeconvolutionLab2: An open-source software for deconvolution microscopy. Methods
115:28–41 Available at: http://dx.doi.org/10.1016/j.ymeth.2016.12.015.

Sanes DH, Bao S (2009) Tuning up the developing auditory CNS. Curr Opin Neurobiol 19:188–
199   Available   at:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation
&list_uids=19535241.

Sanes DH, Walsh EJ (1998) The development of central auditory processing. In: Development
947   of the Auditory System (Rubel EW, Popper AN, Fay RR, eds), pp 271–314. New York:
948   Springer.

949   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden
950   C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P,
951   Cardona A (2012) Fiji: an open-source platform for biological-image analysis. Nature
952   methods 9:676–682.

953   Schrode KM, Dent ML, Lauer AM (2022) Sources of variability in auditory brainstem response
954   thresholds in a mouse model of noise-induced hearing loss. J Acoust Soc Am 152:3576–
955   3582 Available at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9756347/ [Accessed
956   May 16, 2024].

957   Shapiro SM (1988) Acute brainstem auditory evoked potential abnormalities in jaundiced Gunn
958   rats given sulfonamide. Pediatr Res 23:306–310.

959   Shaw NA (1988) The auditory evoked potential in the rat--a review. Prog Neurobiol 31:19–45.

960   Sheets L, Kindt KS, Nicolson T (2012) Presynaptic CaV1.3 channels regulate synaptic ribbon size
961   and are required for synaptic maintenance in sensory hair cells. J Neurosci 32:17273–
962   17286.

963   Shera CA, Guinan Jr. JJ (1999) Evoked otoacoustic emissions arise by two fundamentally
964   different mechanisms: a taxonomy for mammalian OAEs. J Acoust Soc Am 105:782–798.

965   Simmons DD, Mansdorf NB, Kim JH (1996) Olivocochlear innervation of inner and outer hair cells
966   during postnatal maturation: evidence for a waiting period. J Comp Neurol 370:551–562
967   Available   at:
968   http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation
969   &list_uids=8807454.

970   Simmons DD, Manson-Gieseke L, Hendrix TW, McCarter S (1990) Reconstructions of efferent
971   fibers in the postnatal hamster cochlea. Hear Res 49:127-39.

972   Slika E, Fuchs PA, Wood MB (2025) Virally mediated enhancement of efferent inhibition reduces
973   acoustic trauma in wild-type murine cochleas. Mol Ther Methods Clin Dev 33:101455
974   Available at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC11999434/ [Accessed May
975   29, 2025].

976   Song L, McGee J, Walsh EJ (2006) Frequency- and level-dependent changes in auditory
977   brainstem responses (ABRS) in developing mice. J Acoust Soc Am 119:2242–2257.

978   Sonntag M, Englitz B, Kopp-Scheinpflug C, Rübsamen R (2009) Early Postnatal Development of
979   Spontaneous and Acoustically Evoked Discharge Activity of Principal Cells of the Medial
980   Nucleus of the Trapezoid Body: An In Vivo Study in Mice. J Neurosci 29:9510–9520
981   Available at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6666529/ [Accessed October
982   14, 2024].

Sridhar TS, Brown MC, Sewell WF (1997) Unique postsynaptic signaling at the hair cell efferent synapse permits calcium to evoke changes on two time scales. Journal of Neuroscience 17:428–437.

Sridhar TS, Liberman MC, Brown MC, Sewell WF (1995) A novel cholinergic "slow effect" of efferent stimulation on cochlear potentials in the guinea pig. J Neurosci 15:3667–3678.

Stamataki S, Francis HW, Lehar M, May BJ, Ryugo DK (2006) Synaptic alterations at inner hair cells precede spiral ganglion cell loss in aging C57BL/6J mice. Hearing Research 221:104–118 Available at: http://www.ncbi.nlm.nih.gov/pubmed/17005343 [Accessed August 31, 2013].

Sun YJ, Wu GK, Liu B-H, Li P, Zhou M, Xiao Z, Tao HW, Zhang LI (2010) Fine-tuning of pre-balanced excitation and inhibition during auditory cortical development. Nature 465:927–931.

Suthakar K, Liberman MC (2021) Auditory-Nerve Responses in Mice with Noise-Induced Cochlear Synaptopathy. https://doi.org/101152/jn003422021 Available at: https://journals.physiology.org/doi/abs/10.1152/jn.00342.2021 [Accessed November 24, 2021].

Taranda J, Maison SF, Ballestero JA, Katz E, Savino J, Vetter DE, Boulter J, Liberman MC, Fuchs PA, Elgoyhen AB (2009) A point mutation in the hair cell nicotinic cholinergic receptor prolongs cochlear inhibition and enhances noise protection. PLoS Biology 7:71–83.

Tierney TS, Russell FA, Moore DR (1997) Susceptibility of developing cochlear nucleus neurons to deafferentation-induced death abruptly ends just before the onset of hearing. J Comp Neurol 378:295–306.

Tritsch NX, Bergles DE (2010) Developmental regulation of spontaneous activity in the Mammalian cochlea. J Neurosci 30:1539–1550.

Vattino LG, Wedemeyer C, Elgoyhen AB, Katz E (2020) Functional Postnatal Maturation of the Medial Olivocochlear Efferent–Outer Hair Cell Synapse. Journal of Neuroscience 40:4842–4857.

Vetter DE, Liberman MC, Mann J, Barhanin J, Boulter J, Brown MC, Saffiote-Kolman J, Heinemann SF, Elgoyhen AB (1999) Role of alpha9 nicotinic ACh receptor subunits in the development and function of cochlear efferent innervation. Neuron 23:93-103.

Waguespack J, Salles FT, Kachar B, Ricci AJ (2007) Stepwise morphological and functional maturation of mechanotransduction in rat outer hair cells. J Neurosci 27:13890–13902.

Walsh E, McGee J, McFadden S, Liberman M (1998) Long-term effects of sectioning the olivocochlear bundle in neonatal cats. JNeurosci 18:3859–3869.

Wang Y, Sanghvi M, Gribizis A, Zhang Y, Song L, Morley B, Barson DG, Santos-Sacchi J, Navaratnam D, Crair M (2021) Efferent feedback controls bilateral auditory spontaneous activity. Nat Commun 12:2449 Available at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8079389/ [Accessed March 7, 2024].

Weisstaub N, Vetter DE, Elgoyhen AB, Katz E (2002) The alpha9alpha10 nicotinic acetylcholine
receptor is permeable to and is modulated by divalent cations. Hear Res 167:122-35.

Wheeler EF, Bothwell M, Schecterson LC, von Bartheld CS (1994) Expression of BDNF and NT-
3 mRNA in hair cells of the organ of Corti: quantitative analysis in developing rats. Hear
Res 73:46–56.

Wong HC, Zhang Q, Beirl AJ, Petralia RS, Wang Y-X, Kindt K (2019) Synaptic mitochondria
regulate hair-cell synapse size and function. eLife 8:e48914 Available at:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6879205/ [Accessed December 19, 2024].

Zhang Y, Hiel H, Vincent PFY, Wood MB, Elgoyhen AB, Chien W, Lauer A, Fuchs PA (2023)
Engineering olivocochlear inhibition to reduce acoustic trauma. Mol Ther Methods Clin
Dev 29:17–31.

Zhou X, Merzenich MM (2008) Enduring effects of early structured noise exposure on temporal
modulation in the primary auditory cortex. Proc Natl Acad Sci U S A 105:4423–4428
Available at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2393777/ [Accessed May 13,
2024].

1036
1037