Early Noise Exposure and Changes in Medial Olivocochlear Strength Alters Auditory Pathway Development

Authors: Valeria C. Castagna[1,2], Luis E. Boero[1,2], Mariano N. Di Guilmi[1], Paul A. Fuchs[3], Amanda M. Lauer[3], Ana Belén Elgoyhen[1,2] and Maria Eugenia Gomez-Casati[2]

Affiliations: [1]Instituto de Investigaciones en Ingeniería Genética y Biología Molecular, Dr. Héctor N. Torres, Consejo Nacional de Investigaciones Científicas y Técnicas, 1428 Buenos Aires, Argentina.

[2]Instituto de Farmacología, Facultad de Medicina, Universidad de Buenos Aires, 1121 Buenos Aires, Argentina and Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina.

[3]Departments of Otolaryngology-HNS, Neuroscience, and Functional Anatomy and Evolution, Johns Hopkins University, School of Medicine, Baltimore, Maryland 21205, United States.

Corresponding Author: María Eugenia Gómez-Casati, Ph.D.

Instituto de Farmacología. Facultad de Medicina. Universidad de Buenos Aires.

Address: Paraguay 2155. C1121ABG, Buenos Aires, Argentina.

Phone: 54-11-52853350

E-mail: mgomezcasati@fmed.uba.ar, megomezcasati@gmail.com

Conflict of Interest: The authors declare no competing financial or non-financial interest.

26   **Acknowledgements:** This research was supported by Agencia Nacional de Promoción Científica

27   y Técnica (Argentina) to M.E.G-C and NIH Grant R01 DC001508 (A.M.L and A.B.E).

29   **Author Contributions:** V.C.C, L.E.B, M.N.D-G, P.A.F, A.M.L, A.B.E and M.E.G-C designed

30   research; V.C.C performed experiments; V.C.C analyzed data. M.E.G-C wrote the paper. All

31   authors discussed the results, read, edited, and commented on the manuscript.

ABSTRACT

Periodic spontaneous activity is a key feature of developing sensory systems, essential for the formation and refinement of neural circuits. Before the onset of hearing in altricial mammals, cochlear inner hair cells exhibit spontaneous electrical activity that activates primary afferents to propagate into the central nervous system. This activity, modulated by the medial olivocochlear efferent feedback via α9α10 nicotinic cholinergic receptors present in inner hair cells, is crucial for auditory system maturation. In this study, we examined the impact of sound exposure at levels below those causing hair cell damage during the critical developmental period by using genetically modified mice - α9 knockout and α9 knock-in models - with either absent or enhanced cholinergic activity. We further assessed how varying levels of medial olivocochlear feedback interact with early-age noise exposure to affect auditory development. Our findings reveal that both increased and absent olivocochlear activity result in altered auditory sensitivity at the onset of hearing and affect ribbon synapse number and morphology. Additionally, early exposure to loud noise caused long lasting changes in the inner ear of wild-type and α9 knockout mice, underscoring the heightened vulnerability during early development. In contrast, mice with enhanced cholinergic activity were protected from these changes. Overall, this work highlights the critical role of medial olivocochlear modulation in proper auditory system development and indicates that early noise exposure can interfere with normal cochlear maturation, resulting in pronounced long-term effects.

INTRODUCTION

Cochlear inner hair cells (IHCs) and afferent neurons in altricial mammals exhibit spontaneous electrical activity that propagates into the central nervous system (CNS) prior to the onset of hearing, which occurs around postnatal day 12 (P12) (1–5). This spontaneous firing is thought to be crucial for promoting neuronal survival in early auditory centers, driving their maturation, and shaping their synaptic connections through activity-dependent mechanisms (6–11). In the mouse cochlea, spontaneous activity in IHCs arises at late embryonic stages, triggered by ATP release from adjacent inner supporting cells and relies on interactions between depolarizing Ca^{2+} currents and repolarizing K^+ currents (2, 4, 12–14).

Before the onset of hearing, IHCs' spontaneous activity is modulated by transient efferent feedback from medial olivocochlear (MOC) neurons descending from the superior olivary complex in the brainstem. These MOC endings form axosomatic synaptic contacts with immature IHCs via α9α10 nicotinic cholinergic receptors (nAChRs), coupled to Ca^{2+}-dependent SK2 channels, resulting in hyperpolarization of IHCs (15–17). This transient efferent innervation is thought to shape burst firing patterns, guiding the normal functional development of IHCs' synaptic machinery and ensuring the establishment of bilateral neural representation in the developing CNS (10, 11, 18, 19). As IHCs mature, they lose direct efferent contacts and no longer respond to acetylcholine (ACh). Thus, in adult mice MOC fibers synapse exclusively with OHCs (20, 21). The MOC reflex in adulthood is thought to regulate hearing's dynamic range (22), enhances signal detection in noisy environments (23–25), contributes to selective attention (26), provides protection from noise-induced hearing damage (27–31) and helps slow cochlear aging (32, 33).

During the postnatal period around the onset of hearing, the cochlea and central auditory system undergo significant structural and functional changes with ongoing refinement continuing at different levels of the auditory pathway. While maturation at the auditory cortex level is typically reached around P40, certain structures, such as the endbulbs of Held in the brainstem, continue

to refine their synaptic architecture beyond this time frame, up to approximately postnatal week 9

(34–38). Several studies have shown that sound exposure during development significantly

contributes to the functional shaping of the structural organization and signal processing

capabilities of the mammalian auditory system (34, 39–41). Early acoustic exposure influences

neuronal sensitivity, innervation, frequency tuning and response characteristics of neurons at

multiple levels of the auditory pathway (35–37, 42–44). Nevertheless, the extent to which

ascending and descending pathways interact with the acoustic environment to shape activity

patterns and instruct neuronal circuit formation remain largely unknown. In this study, we queried

the effect of the strength of the cholinergic component of the MOC system on auditory system

development. We used genetically modified mice with varying levels of α9α10 nAChR activity: α9

knock-out mice (Chrna9 KO; α9KO), which lacks cholinergic transmission between efferent

neurons and hair cells (45), and α9 knock-in mice (Chrna9L9'T; α9KI) carrying an α9 point

mutation that leads to enhanced MOC activity (29). In addition, we explored the effect of exposure

to sound levels below those causing hair cell damage or permanent threshold shifts during this

critical period of auditory system development. We demonstrate that alterations in MOC

modulation results in changes in auditory sensitivity at the onset of hearing and affect the number

and morphology of ribbon synapses of mature IHCs. Moreover, early-age exposure to loud noise

produced permanent alterations in auditory sensitivity in wild type (WT) and α9KO mice, indicating

the importance of normal patterns of sound-driven activity during the initial stages of auditory

development. Notably, the enhancement of cholinergic activity effectively prevented elevations in

auditory thresholds and cochlear synaptopathy following noise exposure, even during the early

stages of hearing development when the auditory system is still maturing and olivocochlear

terminals have recently formed synapses with OHCs (20, 21). Thus, our work contributes to the

notion that cochlear efferent modulation during critical developmental periods is essential for

establishing proper auditory circuitry. Furthermore, we demonstrate that early noise exposure can

interfere with normal cochlear development, leading to more pronounced long-term

consequences.

RESULTS

Onset of auditory perception in mice with different degrees of efferent inhibition

We tracked the onset and development of auditory brainstem responses (ABRs) and distortion-

product otoacoustic emissions (DPOAE) at different postnatal ages in mice with distinct levels of

α9α10 nAChR activity: 1) WT, with normal MOC function; 2) α9KO, which lacks cholinergic

transmission between MOC neurons and hair cells (45), and 3) α9KI mice carrying an α9 nAChR

subunit point mutation that leads to enhanced responses to ACh (29). Individual ABR waves

reflect the activation of auditory periphery and brainstem processing relay stations within the first

~7 ms after sound stimulus onset (waves 1 to 5). Wave 1 represents the synchrony in firing of

spiral ganglion neurons and is highly correlated with the number of synapses between IHCs and

auditory nerve fibers, while waves 2-5 represent responses from the ascending auditory pathway

(46–51). DPOAEs, sounds recorded from the external auditory canal, assess OHC function. In a

healthy cochlea, two close pure tones produce distortions due to OHC nonlinearities, which are

amplified by OHC electromotility and detected in the external ear canal (52, 53). Measurements

were obtained in anesthetized mice between P12 to P75. Figures 1a and b illustrate that ABR

onset, defined as the lowest sound level (<80 dB SPL) at which a discernible wave 1 was

observed, occurred earlier in α9KI mice with enhanced MOC function compared to WT mice with

normal MOC synapses, and was delayed in α9KO mice lacking α9α10 nAChR activity.

Representative ABR traces recorded at P14 in response to a 16 kHz stimulus across varying

intensities illustrate differences in ABR thresholds among the three mouse groups with distinct

levels of α9α10 nAChR activity (Figure 1a). At this developmental stage, α9KI mice exhibited

lower ABR thresholds, and more distinct upward peaks at suprathreshold intensities compared to

WT mice. In contrast, α9KO mice showed higher ABR thresholds relative to WT mice (Figure 1a).

Figure 1b shows the mean ABR thresholds measured at test frequencies of 11.33, 16, and 22.65

kHz from P12 to P75 across the three groups of mice. Statistical analysis revealed significant

differences among the groups (Kruskal-Wallis, degree of freedom [df] = 2, p = 0.03 at 11.33 kHz;

p = 0.02 at 16 kHz and p = 0.01 at 22.65 kHz). α9KI mice exhibited measurable ABR responses

as early as P13 and achieved mature threshold levels earlier than their WT littermates (Figure

1b). In contrast, ABR onset was significantly delayed in α9KO mice, particularly at 11.33 and 16

kHz (Kruskal-Wallis, df = 2, p = 0.04 at 11.33 and 16 kHz; Figure 1b) indicating that cochlear

maturation is slower in the absence of pre-hearing efferent modulation. At P13, no ABR responses

were observed in WT (0/9) or α9KO (0/7) mice in response to tone pips at 80 dB SPL across

11.33, 16, and 22.65 kHz. In contrast, 6 out of 9 α9KI mice exhibited ABR responses at P13. By

P14, ABR responses were present in 77.7% of WT mice at 11.33 and 22.65 kHz and in 100% at

16 kHz. Meanwhile, 85% of α9KO mice showed responses across all frequencies, although with

elevated thresholds compared to WT (Figure 1b and Supplementary Figure 1a). All α9KI mice

demonstrated robust responses at all tested frequencies at P14. By P15, response rates reached

100% in WT mice and 91% in α9KO mice. DPOAE responses exhibited a similar trend as ABR

thresholds only at 11.33 kHz, although the changes were not statistically significant (Kruskal-

Walis, df = 2, p > 0.05 at all frequencies; Figure 1c). Note that at P22 and P75, mean ABR and

DPOAE thresholds in α9KI mice were slightly elevated by 5–10 dB compared to WT and α9KO

mice (Figure 1b, c). This elevation reached statistical significance only for ABR thresholds at P22

(Kruskal-Wallis, df = 2, p = 0.03 at 11.33 kHz; p = 0.04 at 16 kHz and p = 0.02 at 22.65 kHz). The

elevated thresholds in α9KI mice at these "mature stages" suggest that this may be due to

enhanced cholinergic activity on OHCs, indicating cholinergic MOC effects under resting

conditions as previously reported (29). Collectively, these findings indicate that auditory sensitivity

is altered at hearing onset in mice with either absent or enhanced cholinergic activity.

Suprathreshold response amplitude for ABR waves 1, 2 and 3 in mice with different levels

of α9α10 nAChR activity

ABR individual waves indicate the activation of successive auditory processing centers and can

serve as a functional hearing assessment to identify retro-cochlear abnormalities contributing to

hearing loss (47–49). Tone pip-evoked ABR waveform amplitudes were analyzed at a

suprathreshold level of 80 dB SPL for peaks corresponding to the auditory nerve (wave 1),

cochlear nucleus (wave 2), and superior olivary complex (SOC) (wave 3) (50, 51) at different test

frequencies in mice with varying levels of MOC strength.

During the early postnatal period, ABR amplitudes were smaller, and the number of discernible

upward peaks was fewer in WT and α9KO mice compared to α9KI mice (see representative ABR

traces from P13 to P15, Supplementary Figure 1a). Up to P15, response peaks at 80 dB SPL

were slightly larger and more distinct in α9KI mice, reflecting their enhanced auditory responses

during this critical developmental window. Analysis of ABR wave 1 latencies across different

postnatal ages revealed no significant differences between genotypes (Supplementary Figure

1b). At P14, wave 1 latencies were consistently longer across all groups. By P15, latencies

stabilized to levels characteristic of mature mice, coinciding with the rapid auditory maturation

phase. ABR wave 1 amplitudes increased progressively with postnatal age, reaching a plateau at

P22 across all genotypes and frequencies (11.33, 16, and 22.65 kHz; Figure 2a). In WT mice,

this increase exhibited a steadier progression compared to that observed in mice with either null

or enhanced α9α10 nAChR activity. Notably, ABR wave 1 amplitudes in α9KI mice tended to be

slightly larger than those in WT mice up to P15, although these differences were not statistically

significant (one-way ANOVA, df = 2, $p > 0.05$ at all frequencies; Figure 2a). After P15, wave 1

amplitudes in WT mice tended to surpass those of α9KI mice, with significant differences

observed at P75 at 16 and 22.65 kHz (one way ANOVA followed by Holm-Sidak test, df=2, p=0.03

at both frequencies; Figure 2a). In α9KO mice, wave 1 amplitudes remained smaller than those

of WT and α9KI mice until P16, with the difference being significant only at 22.65 kHz (one way

ANOVA followed by Holm-Sidak test, df = 2, p = 0.02; Figure 2a). These findings align with the

auditory threshold results, suggesting that the strength of transient MOC efferent innervation to

IHCs contributes to the timely development of auditory function. ABR wave 2 amplitudes

increased linearly across postnatal ages, stabilizing at P22 in all groups, with no significant

differences observed across genotypes at any frequency or age (one way ANOVA, df = 2, p >

0.05 at all frequencies; Figure 2b). Conversely, ABR wave 3 amplitudes showed marked

reductions in α9KI mice at most postnatal ages, with significant decreases at P22 and P75 for 16

kHz (one way ANOVA followed by Holm-Sidak test, df = 2, p = 0.005 at P22 and p = 0.01 at P75;

Figure 2c). In α9KO mice, reduced wave 3 amplitudes were observed at P15, specifically at 16

kHz (one way ANOVA followed by Holm-Sidak test, df=2, p=0.04; Figure 2c). These findings

indicate that both disruption and enhancement of the MOC synapses influence the maturation of

cochlear synapses and the development of extra peripheral nuclei in the auditory pathway.

ABR thresholds after early-age noise exposure in mice with different degrees of efferent

inhibition

To investigate the impact of early-age noise exposure on auditory development, we exposed WT,

α9KO, and α9KI mice to loud noise at P15. At this stage, all three genotypes exhibited measurable

ABR waveforms, though their wave 1 thresholds differed and had not yet reached mature levels

(Figure 1). This allowed us to evaluate how early noise exposure influences auditory function

before threshold maturation in each genotype. Mice were exposed to 1–16 kHz noise at 100 dB

SPL for 1 hour at P15. Auditory responses were then assessed at 1, 7, and 60 days post-exposure

to track functional changes over time.

Previous studies in WT mice exposed to the same noise protocol at P21 reported significant

auditory threshold shifts of 10–35 dB SPL one day after exposure, with thresholds returning to baseline within a week, indicating a transient threshold shift (31). In contrast, as shown in Figure 3a, no changes in ABR thresholds were observed one day after noise exposure at P16 across frequencies of 8, 16, and 22.65 kHz in WT mice compared to unexposed controls. Remarkably, ABR thresholds at P16 were lower than those measured at P15, prior to noise exposure, suggesting that noise had no immediate impact on the maturation of auditory thresholds at this early postnatal developmental stage. One week after exposure, however, ABR thresholds were elevated at all frequencies in noise-exposed WT animals, and this increase persisted for at least two months post-exposure (Mann- Whitney test, df = 1, P22: p = 0.006 at 8 kHz; p = 0.007 at 16 kHz; p = 0.002 at 22.65 kHz and P75: p = 0.04 at 8 kHz; p = 0.03 at 16 kHz and p = 0.04 at 22.65 kHz; Figure 3a). This outcome contrasts with the transient threshold elevation observed in WT mice, from the same genetic background, exposed at P21 using the same noise protocol (31), suggesting that noise exposure at this early developmental stage produces a more detrimental long-term damage.

A similar trend as seen in WT mice was observed after exposing α9KO mice to noise at P15 (Figure 3b). One day after exposure, there were no changes in ABR thresholds at 8, 16 and 22.65 kHz. However, seven days post-exposure, there was an elevation in auditory thresholds across all frequencies (Mann-Whitney test, df = 1, p = 0.03 at 8 kHz; p = 0.01 at 16 kHz and p = 0.02 at 22.65 kHz), which, unlike in WT mice, returned to normal levels after two months (Mann-Whitney test, df = 1, p > 0.05 at all frequencies; Figure 3b). Notably, in α9KI mice, ABR thresholds were not affected by noise exposure at all the frequencies tested (Figure 3c), suggesting that even at this early age, the enhancement of cholinergic activity on OHCs provides protection from noise-induced cochlear threshold elevations (Mann-Whitney test, df = 1, p > 0.05 at all frequencies).

Amplitudes of cochlear sound-evoked potentials after early-age noise exposure in mice

with different degrees of efferent inhibition

To assess the neural activity across auditory nerve fibers, we measured ABR wave 1 amplitudes

at frequencies of 8, 16, and 22.65 kHz at different ages after noise exposure in the three mouse

groups (Figure 4). One day after noise exposure, we observed a reduction in wave 1 amplitudes

across all frequencies in WT mice. Further testing WT mice one week and two months post-

exposure revealed a significant and lasting reduction, indicating a permanent decrease of ABR

wave 1 amplitude (Mann- Whitney test, df = 1, P22: p = 0.007 at 8 kHz; p = 0.005 at 16 kHz; p =

0.005 at 22.65 kHz and P75: p = 0.04 at 8 kHz; p = 0.002 at 16 kHz and p = 0.002 at 22.65 kHz;

Figure 4a). Similar permanent reductions in wave 1 amplitude were observed in α9KO mice after

exposure to noise at 16 and 22.65 kHz (Mann-Whitney test, df = 1, P22: p = 0.04 at 16 kHz; p =

0.001 at 22.65 kHz and P75: p = 0.03 at 16 kHz and p = 0.04 at 22.65 kHz; Figure 4b). Notably,

no changes in ABR wave 1 amplitudes were observed at any stage post-exposure in mice with

enhanced α9α10 nAChR activity, suggesting a preservation of cochlear synapses after noise

exposure (Mann-Whitney test, df = 1, p > 0.05 at all frequencies; Figure 4c).

DPOAE thresholds after early-age noise exposure in mice with different levels of α9α10

nAChR activity

To assess the impact of early-age noise exposure on the function of OHCs, we recorded DPOAE

at 8, 16, and 22.65 kHz on the three groups of mice with varying levels of MOC function (Figure

5). During early development, the auditory system undergoes significant maturation, leading to

decreasing DPOAE thresholds with age, as shown in Figure 1c. In WT mice, one day after noise

exposure at P16, there was no difference in DPOAE thresholds between the noise-exposed group

and unexposed controls (Figure 5a). Importantly, when comparing DPOAE thresholds at P16 to

those at P15 within both groups, a clear reduction was observed (Figure 5a). This decline in

thresholds underscores the natural maturation process of the auditory system, indicating that

early noise exposure does not immediately disrupt the ongoing development of DPOAE

thresholds at this early postnatal stage. By P22, seven days after noise exposure, an increase in

DPOAE thresholds was observed that was statistically significant only at 16 kHz (Mann-Whitney,

df = 1, p = 0.01; Figure 5a). Yet, two months later, DPOAE thresholds were comparable to those

of unexposed WT controls across all tested frequencies, suggesting that early noise exposure did

not result in long-term changes in OHC function (Figure 5a).

In α9KO mice, DPOAE thresholds remained unchanged 1 day after noise exposure. However,

significant threshold elevations were observed at 8 and 16 kHz, but not at 22.65 kHz, seven days

post-exposure (Mann-Whitney, df = 1, p = 0.02 at 8 kHz and p = 0.02 at 16 kHz). After two months,

recovery to unexposed levels occurred only at 8 kHz, while DPOAE thresholds remained elevated

at 16 kHz (Mann-Whitney, df = 1, p = 0.003 at 16 kHz; Figure 5b). In contrast, mice with enhanced

MOC function showed no changes in DPOAE thresholds at any time point post-exposure (Figure

5c), suggesting that early-age noise exposure did not impair cochlear amplifier function.

Ribbon synapse changes after early noise exposure in mice with different levels of α9α10

nAChR activity

Following the completion of functional testing, histological analysis was performed on cochleae

dissected and fixed at P75 from all experimental groups to evaluate the density and structure of

IHC ribbon synapses. To visualize these ribbon synapses, we performed immunostaining on

whole-mount preparations of the organ of Corti with antibodies against CtBP2-Ribeye, a

presynaptic ribbon protein (54), and GluA2, an AMPA-type glutamate receptor expressed at the

postsynaptic afferent terminal (30, 55, 56). Ribbon synapses were identified by the colocalization

of CtBP2 and GluA2 puncta at the base of the IHC (56). We first compared the number of

colocalized synaptic markers per IHC in unexposed WT, α9KO, and α9KI mice across three

cochlear locations - apical, medial, and basal - in each cochlear whole-mount (Figure 6). In the

low-frequency apical end, both α9KO and α9KI mice exhibited a significant decrease in synaptic

density, with reductions of 30.64% and 27.94%, respectively (Kruskal-Wallis, df=2, p<0.0001 for

both groups; Figure 6a, left panel). In the medial region, the reduction was significant only in α9KI

mice, with a decrease of 35.97% compared to WT (Kruskal-Wallis, df = 2, p < 0.0001; Figure 6a,

middle panel). Similarly, in the high-frequency basal region, colocalized synaptic puncta were

significantly reduced only in α9KI mice, with a 22.07% decrease relative to WT (Kruskal-Wallis,

df = 2, p = 0.02; Figure 6a, right panel).

We further quantified the volume of CtBP2 puncta and GluA2 receptor patches in these

colocalized synaptic markers (Figure 6b, c). In the apical region, both α9KO and α9KI mice

showed a reduction in ribbon volumes (one way ANOVA followed by Holm-Sidak test, df = 2, p <

0.0001 for both groups; Figure 6b, left panel). In the middle region, the decrease was significant

only in α9KI ears in comparison with α9KO ears (one way ANOVA followed by Holm-Sidak test,

df = 2, p < 0.0001; Figure 6b, middle panel). Interestingly, in the basal region, an increase in

CtBP2 volume was detected in α9KO mice, while a decrease was observed in α9KI mice (one

way ANOVA followed by Holm-Sidak test, df = 2, p < 0.0001 for both groups; Figure 6b, right

panel). The size of the GluA2 receptor patches was significantly reduced in α9KI mice in the apical

region, with no significant changes in α9KO mice compared to WT (one way ANOVA followed by

Holm-Sidak test, df = 2, p < 0.0001; Figure 6c, left panel). In the medial region, a reduction in

GluA2 volume was observed in both α9KO and α9KI mice (one way ANOVA followed by Holm-

Sidak test, df = 2, p < 0.0001, for both groups; Figure 6c, middle panel). In the basal region, α9KO

mice exhibited an increase in GluA2 size relative to WT (one way ANOVA followed by Holm-Sidak

test, df = 2, p < 0.0001; Figure 6c, right panel). These results suggest that inhibitory input from

cholinergic efferent fibers to hair cells plays a key role in shaping the synaptic density and

maintaining the proper morphology of pre- and postsynaptic elements critical for optimal auditory

function.

Figure 7 illustrates the impact of early-age noise exposure on auditory nerve synapse numbers

across the different experimental groups. Presynaptic, postsynaptic, and colocalized synaptic

counts were conducted in the apical, medial, and basal region of the cochlea. In WT exposed

ears, the number of presynaptic, postsynaptic, and colocalized synaptic puncta significantly

decreased at both the apical and basal cochlear ends (Figure 7a). The reduction in CtBP2 puncta

was more pronounced in the high-frequency basal region, showing a 37.2% decrease compared

to controls (Mann-Whitney, $df = 1$, $p < 0.0001$ at the apical and basal regions). Similarly, GluA2

postsynaptic receptors were reduced at both the apical and basal turns in the cochlea of

traumatized WT mice (Mann-Whitney, $df = 1$, $p < 0.0001$ at both cochlear regions). Putative ribbon

synapse counts, defined by juxtaposed CtBP2- and GluA2-positive puncta, showed a 34.21%

reduction in the apical and a 32.56% reduction in the basal turn following acoustic trauma (Mann-

Whitney, $df = 1$, $p < 0.0001$ at both cochlear regions; Figure 7a, right panel). In α9KO mice, there

was also a reduction in the number of presynaptic, postsynaptic, and colocalized puncta

depending on cochlear frequency/location after noise exposure (Figure 7b). Ribbon puncta in

α9KO exposed ears were reduced by up to 14.73% of the control in the apical and 24.75% in the

basal cochlear ends after noise exposure (Mann-Whitney, $df = 1$, $p = 0.001$ and $p = 0.002$ at the

apical and basal region, respectively). GluA2-positive postsynaptic receptor patches decreased

by 24.79% in the basal turn of noise-exposed mice (Mann-Whitney, $df = 1$, $p = 0.004$). Following

noise exposure, synaptic puncta were reduced by 26.17% in the apical cochlear region and

25.87% in the basal region (Mann-Whitney, $df = 1$, $p = 0.002$ and $p < 0.0001$ at the apical and

basal region, respectively; Figure 7b, right panel). Interestingly, after two months post-exposure,

α9KI ears exhibited no significant changes in presynaptic ribbons, postsynaptic GluA2 receptor

patches, or putative synapses across all cochlear regions. Although not statistically significant, a

modest increase in synaptic density (~19% at the basal region) was observed in α9KI tissue

(Figure 7c, right panel).

We finally quantified the volume of ribbons and AMPA receptor patches at ribbon synapses (i.e.,

colocalized CtBP2 and GluA2 puncta) from unexposed and exposed mice with different levels of

MOC function (Figure 8). In WT mice, noise exposure led to significantly higher ribbon volumes

in the basal end of the cochlea compared to unexposed mice, with no corresponding changes in

AMPA receptor patch size (t-test, df = 1, p = 0.01; Figure 8 a). No changes were detected in the

apical and medial regions of the cochlea (Figure 8 a). Conversely, in α9KO mice, both ribbon and

AMPA receptor volumes were reduced in the basal region following noise exposure, while

volumes remained unchanged in the apical and medial regions (t-test, df = 1, p = 0.002 for CtbP2

volume and p = 0.003 for GluA2 volume; Figure 8 b). Interestingly, α9KI mice showed no

morphological alterations, indicating that enhanced α9α10 nAChR activity protects synapses from

noise-induced changes (Figure 8 c).

DISCUSSION

This study examined the influence of MOC inhibitory modulation in auditory maturation and

assessed the effects of early loud sound exposure on developmental outcomes. Our findings

highlight the importance of the MOC system in the proper establishment of IHC ribbon synapses

and reveal that changes in the strength of MOC inhibition impact the timing of hearing onset.

Moreover, loud noise exposure during critical developmental stages caused lasting changes in

auditory sensitivity in WT, highlighting the greater vulnerability of the auditory system to noise-

induced injury during early development. Notably, enhanced cholinergic activity as seen in α9KI

mice conferred protection from noise-induced threshold shifts and cochlear synaptopathy. This

underscores the crucial protective role of the MOC system during these vulnerable early

developmental stages.

Before hearing begins, MOC fibers form direct synapses with IHCs, with cholinergic inhibition

playing a key role in regulating the developmental increase in Ca^{2+} sensitivity of glutamate release

at IHC ribbon synapses (18). This regulation is accomplished by precisely modulating IHC action

potential activity over a period of immature development (5, 15, 18, 57). It is hypothesized that

the maintenance of activity patterns and fidelity in the transmission of these signals are essential

for the proper establishment of the sensory map and the refinement of synaptic connections along

the auditory pathway (10, 11, 58–61).

The current findings highlight the important role of α9α10 nAChR-mediated activity in auditory

function development and ribbon synapse maturation. Enhanced α9α10 nAChR activity in α9KI

mice accelerates the onset of hearing, with sound-evoked potentials detected earlier than in mice

with normal MOC synapses. Conversely, the absence of MOC activity, as observed in α9KO mice,

delays the onset and development of hearing. Consistent with these findings, recent research on

α9KO mice has shown that auditory sensitivity at hearing onset is impaired in the absence of pre-

hearing efferent input (19). In addition, our work shows that both enhanced and absent MOC

inhibition disrupt the normal maturation of ribbon synapses in adult IHCs. Specifically, α9KO and

α9KI mice exhibit significant reductions in ribbon synapse density and notable alterations in the

volume of pre- and post-synaptic puncta compared to WT mice at P75. This underscores the

critical role of balanced MOC activity in auditory system development and synaptic organization.

In α9KI mice, enhanced MOC function is associated with reductions in both ribbon size and GluA2

receptor patch volume. Conversely, α9KO mice show region-specific effects: in the apical region

of the cochlea, ribbon volume decreases without changes in GluA2 receptor size, whereas in the

basal region, both CtBP2 volume and GluA2 receptor size increase. These observations suggest

that altered MOC modulation has differential impacts on synaptic structures along the cochlear

tonotopic axis. Previous studies have emphasized the functional importance of ribbon size,

showing that it is tightly regulated by presynaptic Ca^{2+} channels, with alterations in Ca^{2+} influx

impacting synaptic plasticity, neurotransmitter release, and active zone maturation (62–65).

During early development, IHC active zones undergo substantial structural and functional

maturation, including the removal of extrasynaptic Ca^{2+} channels and the co-localization of

glutamate receptor patches with presynaptic ribbons (66–68). This maturation refines presynaptic

morphology, reduces the number of ribbons per synapse, and establishes monosynaptic

connections between IHCs and auditory nerve fibers (69–72). Recent work also suggests that

fusion of synaptically anchored ribbons during this period is critical for the development of mature

active zones and sustained neurotransmitter release (66). Given our observation of reduced

ribbon size in some cochlear regions of α9KI and α9KO mice, we propose that altered MOC

modulation strength may disrupt intracellular Ca^{2+} dynamics, which could impair ribbon precursor

fusion, thereby impairing active zone maturation. These findings highlight the essential role of

MOC synapses in shaping early auditory sensitivity and emphasize the importance of coordinated

activity between afferent and efferent pathways in regulating spiking activity during this critical

developmental window. This coordination is crucial for proper synaptic and auditory system

maturation.

Auditory function in vertebrates matures through a well-defined process, transforming the system

from high thresholds and limited frequency detection at the onset of hearing to a highly sensitive

and precise mechanism capable of resolving frequency differences as small as 1 Hz (73). Our

findings suggest that the WT mouse strain used in the present work aligns with this established

maturation pattern, as auditory function recordings from P12 to P75 reveal a clear progression

from an immature to a mature state. At the onset of hearing (P14 in our strain of mice), the auditory

system shows elevated ABR and DPOAE thresholds, reduced wave 1 amplitudes, and poorly

defined later ABR waves, reflecting incomplete functional maturation. Over the following days,

thresholds progressively decreased, reaching mature levels by P22. This maturation is

accompanied by a steady increase in ABR wave 1 amplitudes from P15 to P22, underscoring the

critical refinement period necessary for achieving full auditory sensitivity. Thus, by P15, the time

when mice were exposed to high levels of noise, the maturation process is not yet complete. A

key finding of this study is that exposure to intense sound during early developmental stages

results in significantly more severe consequences compared to exposure at a later age (31). In

WT mice, noise exposure at 100 dB SPL for 1 hour at P15 caused persistent ABR threshold

elevations lasting up to two months. In contrast, the same exposure at P21 led to only transient

threshold shifts, which resolved within seven days (31). This stark difference underscores the

heightened vulnerability of the auditory system during early development and highlights the age-

dependent impact of noise exposure. Previous reports have also shown that the same scheme of

intense noise exposure causes more severe damage in juvenile animals at 4 weeks of age

compared to adult animals at 16 weeks of age (74–76). DPOAE thresholds were also elevated

seven days post-exposure in WT mice, although these thresholds recovered after two months,

suggesting that there is no loss of OHCs or alteration in their function. Given this recovery of

DPOAE thresholds but persistently elevated ABR thresholds, we can infer that the elevation is

not due to damage in the cochlear amplifier but rather to a deficiency in the afferent pathway of

sound signal transmission. In line with this, the increases in ABR thresholds were accompanied

by a reduction in the wave 1 amplitude, which also persisted for two months following the

exposure to noise. This study represents the first report of sustained ABR threshold elevation for

such an extended period following moderate noise exposure at 100 dB SPL for 1 hour. Similar

results were obtained in 14-day-old rats exposed to a higher noise level of 125 dB SPL for 25

minutes, where after two months post-exposure, elevated thresholds at high frequencies and a

loss of presynaptic elements (CTBP2/Ribeye) were observed (77). In another study involving 10-

day-old rats exposed to intense sounds of 105 dB SPL for two hours, a loss of presynaptic

elements was also observed after 8 weeks. However, auditory thresholds did not show changes

(78). Our findings demonstrate a significant loss of ribbon synapses in the apical and basal

regions of the cochlea in WT mice following noise exposure. In addition to cochlear synaptopathy,

we observed that IHCs of noise-exposed mice tend to have larger ribbons (Ribeye/CtBP2 volume)

than those of the unexposed IHCs at the basal region of the cochlea. A similar increase in the

volume of presynaptic elements following acoustic trauma was observed using both the RIBEYE-

tagRFP mouse model (79) and volume electron microscopy (76). The observed ribbon

enlargement in WT mice may be attributed to the fusion of individual ribbons after noise exposure.

In fact, recent evidence indicates an increase in multi-ribbon active zones as well as movement

suggesting unanchoring from synaptic tethers in synaptopathic ears (76, 79, 80). While the

functional significance of this structural adaptation remains unclear, multi-ribbon active zones

persist into adulthood (66, 67) and are a hallmark of cochlear aging (81), with larger ribbons being

associated with increased Ca^{2+}-dependent exocytotic activity (82). Notably, noise exposure has

also been associated with enhanced exocytosis in IHCs, along with an increase in tone-burst

responses in auditory nerve fibers (83, 84). In addition, studies have shown that intense noise

exposure (98 dB SPL for two hours) in young animals reduces the number of presynaptic ribbons

in OHCs, while increasing the volume and mobility of the remaining ribbons (85). We can

hypothesize that ribbon volume enlargement may act as a compensatory mechanism to maintain

auditory sensitivity after synapse loss. Importantly, we did not detect changes in the size of GluA2

receptor patches in WT mice post-noise exposure.

In α9KO mice, which lack α9α10 nAChR activity, elevated ABR and DPOAE thresholds were

observed one week after noise exposure at P15, consistent with findings from noise-exposed

α9KO mice at P21 (31). However, in contrast to WT animals, ABR thresholds in α9KO mice nearly

fully recovered to unexposed control levels two months post-exposure. While DPOAE thresholds

returned to control levels at 8 and 22.65 kHz, they remained elevated at 16 kHz after two months.

This elevation in DPOAE thresholds at 16 kHz was accompanied by a slight, though not

statistically significant, elevation in ABR thresholds at the same frequency. The recovery of ABR

thresholds in α9KO mice, compared to the permanent threshold elevation seen in WT mice, may

be attributed to the reduced auditory sensitivity at hearing onset in α9KO mice due to the absence

of prehearing efferent modulation. While ABR thresholds return to pre-exposure values, a

decrease in wave 1 amplitudes was observed, accompanied by a reduction in the number of

ribbon synapses per IHC. Previous work in guinea pigs has suggested that deficits observed post-

noise exposure in suprathreshold stimulus encoding are likely due to the preferential impact of

noise exposure on high-threshold, low spontaneous firing rate neurons (86). Therefore, we can

hypothesize that noise exposure may have specifically affected this particular population of fibers,

and thus, despite some degree of synapse loss, changes in ABR thresholds were not observed.

However, in α9KO ears exposed to noise, both pre- and postsynaptic puncta showed a reduction

in size, indicating noise may also involve maladaptive alterations in both pre- and postsynaptic

structures.

In contrast to the findings in WT and α9KO mice, α9KI mice, which exhibit enhanced α9α10

nAChR activity, showed no ABR or DPOAE threshold elevations following sound exposure.

Additionally, no evidence of cochlear synaptopathy was detected in these mice. This phenomenon

had already been described in adult mice (P21) exposed to the same level of noise (31). At birth,

rodent MOC neurons initially innervate IHCs, and by P10, they expand to innervate both IHCs

and OHCs. After the onset of hearing, their innervation becomes confined to OHCs, gradually

retracting from IHCs (20). Our results therefore suggest that even with early developmental noise

exposure—when MOC innervation to OHCs is still developing—enhanced α9α10 nAChR

responses can effectively exert a protective effect on OHCs against functional and structural

damage associated with acoustic trauma. Importantly, our findings also reveal that mice with

enhanced MOC function showed a modest increase in ribbon synapse density following acoustic

overexposure. This is consistent with previous work, which documented a more substantial

increase in synaptic density in α9KI mice exposed to the same noise protocol at P21 (31). These

results indicate that MOC system enhancement may support synapse formation following noise

exposure.

In summary, this work highlights three key findings: 1) the MOC system plays a crucial role in

regulating the development of ribbon synapse density and size, which is essential for proper

hearing onset; 2) early exposure to loud noise results in more severe auditory consequences; and

3) enhancing $\alpha 9\alpha 10$ nAChR responses during early development protects against noise-induced

damage, underscoring the importance of timely modulation of this receptor post-hearing onset to

mitigate developmental impairments.

 FIGURES

Figure 1: Auditory function in WT, α9KO, and α9KI mice across different postnatal ages.
(**a**) Representative ABR waveforms from WT (black), α9KI (red), and α9KO (blue) mice recorded
at P14 in response to 16 kHz tone bursts at increasing sound pressure levels (dB SPL).
Thresholds, defined by the presence of wave 1, are marked by green dashed traces. The scale
bar applies to all three recordings. (**b**) Mean ABR thresholds from P12 to P75 in WT (n = 12),
α9KO (n = 14), and α9KI (n = 14) mice at 11.33, 16, and 22.65 kHz. ABR waveforms first became
detectable at P13 in α9KI mice, at P14 in WT mice, and between P14 and P15 in α9KO mice. (**c**)
Mean DPOAE thresholds from P12 to P75 in WT (n = 12), α9KO (n = 14) and α9KI (n = 14) mice
at 11.33, 16 and 22.65 kHz. DPOAE thresholds showed the same trend as ABR only at 11.33
kHz. Group means ± SEM are shown. Red asterisks represent the statistical significance of α9KI
compared with WT, and blue asterisks represent the statical significance of α9KO compared with
WT (Kruskal-Wallis followed by post hoc Dunn test, *p < 0.05).

Figure 2: Amplitudes of ABR waves 1, 2, and 3 in WT, α9KO, and α9KI mice across postnatal ages. ABR wave amplitudes (in μV) for waves 1 (**a**), 2 (**b**), and 3 (**c**) in WT, α9KO, and α9KI mice across postnatal ages, measured at 80 dB SPL for frequencies of 11.33, 16, and 22.65 kHz. Amplitudes were determined by calculating the distance between the maximum and minimum peaks. Data points represent the mean ± SEM for WT (n = 12), α9KO (n = 14), and α9KI (n = 14) mice. Red asterisks indicate significant differences between α9KI and WT, while blue asterisks indicate significant differences between α9KO and WT (one-way ANOVA followed by Holm-Sidak test, *p < 0.05; **p < 0.01).

Figure 3: ABR wave 1 threshold measurements in control and noise-exposed WT, α9KO, and α9KI mice at different postnatal ages. ABR thresholds before and after exposure to noise at P15 in WT ($n_{control}$ = 14 and $n_{Exposed}$ = 12) (**a**), α9KO ($n_{control}$ = 11 and $n_{Exposed}$ = 11) (**b**), and α9KI ($n_{control}$ = 12 and $n_{Exposed}$ = 12) (**c**) mice at 8, 16 and 22.65 kHz. WT mice showed a significant increase in ABR thresholds 1 week and 2 months after exposure to noise, indicating a permanent threshold elevation. A recovery of ABR thresholds was observed in α9KO mice 2 months after exposure to noise. α9KI mice did not present any changes in ABR thresholds at any time after acoustic trauma. Each point represents the mean ± SEM. Asterisks represent the statistical significance (Mann-Whitney test, *p < 0.05; **p < 0.01).

516
517
Figure 4: Suprathreshold response amplitude for ABR wave 1 after P15 noise exposure in
WT, α9KO, and α9KI mice at different postnatal ages compared to unexposed controls.
ABR wave 1 amplitudes (in μV) at 80 dB SPL at 8, 16 and 22.65 kHz in WT ($n_{Control}$ = 14 and
$n_{Exposed}$ = 12) (**a**), α9KO ($n_{Control}$ = 11 and $n_{Exposed}$ = 11) (**b**), and α9KI ($n_{Control}$ = 12 and $n_{Exposed}$ = 12)
(**c**) mice at the same time points shown in Figure 3. Each panel inset displays representative ABR
waveforms recorded at 80 dB SPL for the corresponding frequency, shown for all groups at P75.
Scale bars (1 μV vertical, 1 ms horizontal) apply to all traces. WT and α9KO mice exhibited
reduced ABR wave 1 amplitudes following noise exposure, whereas wave 1 amplitudes in α9KI
mice remained unaffected. Data are presented as mean ± SEM. Statistical significance is
indicated by asterisks (Mann-Whitney test, *p < 0.05; **p < 0.01).

Figure 5: **Assessment of OHC functional integrity at different postnatal ages following noise exposure at P15 in WT, α9KO, and α9KI mice.** DPOAE thresholds are shown for WT ($n_{Control}$ = 14 and $n_{Exposed}$ = 12) (**a**), α9KO ($n_{Control}$ = 11 and $n_{Exposed}$ = 11) (**b**), and α9KI ($n_{Control}$ = 12 and $n_{Exposed}$ = 12) (**c**) at 8, 16 and 22.65 kHz. WT mice exhibited a transient increase in DPOAE thresholds 1 week post-exposure at 16 kHz. α9KO mice showed increased thresholds at 8 and 16 kHz 1 week post-exposure, with sustained elevation at 16 kHz up to 2 months. α9KI mice demonstrated no changes in DPOAE thresholds after noise exposure. Data are presented as mean ± SEM. Statistical significance is indicated by asterisks (Mann-Whitney test, *p < 0.05; **p < 0.01).

Figure 6: Analysis of IHC ribbon synapses in unexposed WT, α9KO, and α9KI mice.
Quantitative data obtained from unexposed mice at P75 at the apical (left panel), medial (middle panel) and basal cochlear region (right panel). (**a**) Number of putative ribbon synapses per IHC (i.e., colocalized CtBP2 and GluA2 puncta). WT: $n_{Apical} = 80$ IHC, $n_{Medial} = 75$ IHC, $n_{Basal} = 60$ IHC from 14 animals; α9KO: $n_{Apical} = 100$ IHC, $n_{Medial} = 84$ IHC, $n_{Basal} = 118$ IHC from 11 animals and α9KI: $n_{Apical} = 178$ IHC, $n_{Medial} = 319$ IHC, $n_{Basal} = 198$ IHC from 12 animals. (**b**) Volumes of fluorescence staining for ribbons (CtBP2 size) and (**c**) AMPA receptors (GluA2 size) for each of the synaptic pairs (WT: $n_{Apical} = 450$ synaptic pairs, $n_{Medial} = 509$ synaptic pairs, $n_{Basal} = 203$ synaptic pairs; α9KO: $n_{Apical} = 367$ synaptic pairs, $n_{Medial} = 187$ synaptic pairs, $n_{Basal} = 221$ synaptic pairs and α9KI: $n_{Apical} = 247$ synaptic pairs, $n_{Medial} = 270$ synaptic pairs, $n_{Basal} = 240$ synaptic pairs). Violin plots indicate median (solid line) and interquartile range (IQR, dashed lines). Asterisks denote the statistical significance (Kruskal–Wallis followed by a Dunn's post-test was used to analyze the number of synapses and one way ANOVA followed by Holm-Sidak test; *p < 0.05; **p < 0.01; ***p < 0.001, ****p < 0.0001).

Figure 7: Analysis of the degree of IHC synaptopathy two months after exposure to noise at P15. Left, Representative confocal images of IHC synapses at the apical region from cochleae immunolabeled for presynaptic ribbons (CtBP2-red), postsynaptic receptor patches (GluA2-green) and IHCs (Myosin VIIa-blue). CtBP2 antibody also weakly stains IHC nuclei. Scale bar, 10μm. Right, Quantitative data obtained from WT, α9KO, and α9KI mice at P75. For each IHC, we analyzed the number of CtBP2 puncta, postsynaptic GluA2 receptor patches, and putative ribbon synapses. (**a**) In traumatized WT mice, there was a reduction in the number of CtBP2 puncta, GluA2 receptor patches, and putative synapse, at the apical and basal region of the cochlea (WT Control: n_{Apical} = 80 IHC, n_{Medial} = 75 IHC, n_{Basal} = 60 IHC; WT Exposed: n_{Apical} = 82 IHC, n_{Medial} = 90 IHC, n_{Basal} = 50 IHC). (**b**) In traumatized α9KO mice, there was a reduction in the number of prelocalized, postlocalized, and colocalized puncta depending on cochlear frequency/location (α9KO Control: n_{Apical} = 100 IHC, n_{Medial} = 84 IHC, n_{Basal} = 118 IHC; α9KO Exposed: n_{Apical} = 234 IHC, n_{Medial} = 60 IHC, n_{Basal} = 90 IHC). (**c**) In traumatized α9KI mice, no

566 difference was found in the number of presynaptic ribbons, postsynaptic AMPA receptors, and
567 colocalization puncta for the three regions of the cochlea (α9KI Control: n_{Apical} = 178 IHC, n_{Medial} =
568 319 IHC, n_{Basal} = 198 IHC; α9KI Exposed: n_{Apical} = 196 IHC, n_{Medial} = 145 IHC, n_{Basal} = 80 IHC). Violin
569 plots indicate median (solid line) and IQR (dashed lines). Asterisks denote the statistical
570 significance (Mann Whitney test, **$p < 0.01$; ***$p < 0.001$, ****$p < 0.0001$).

Figure 8: Volumes of presynaptic ribbons and postsynaptic receptor patches for each synaptic pair from control and noise-exposed WT, α9KO and α9KI mice. Violin plots comparing the volumes of fluorescence staining for ribbons (CtBP2 size) and AMPA receptors (GluA2 size) between control and noise-exposed mice. The quantification was made at the apical, medial and basal cochlear regions from WT (**a**) (Control: n_{Apical} = 450 synaptic pairs, n_{Medial} = 509 synaptic pairs, n_{Basal} = 203 synaptic pairs; Exposed: n_{Apical} = 497 synaptic pairs, n_{Medial} = 175

synaptic pairs, n_{Basal} = 250 synaptic pairs); α9KO (**b**) (Control: n_{Apical} = 367 synaptic pairs, n_{Medial} =
187 synaptic pairs, n_{Basal} = 221 synaptic pairs; Exposed: n_{Apical} = 207 synaptic pairs, n_{Medial} = 300
synaptic pairs, n_{Basal} = 145 synaptic pairs) and α9KI (**c**) (Control: n_{Apical} = 247 synaptic pairs, n_{Medial}
= 270 synaptic pairs, n_{Basal} = 240 synaptic pairs; Exposed: n_{Apical} = 280 synaptic pairs, n_{Medial} =
310 synaptic pairs, n_{Basal} = 360 synaptic pairs) mice. Violin plots indicate median (solid line) and
IQR (dashed lines). Asterisks denote the statistical significance (t-test; *$p < 0.05$; ***$p < 0.001$).

SUPPLEMENTAL INFORMATION

Supplementary Figure 1. Suprathreshold ABR waveforms and wave 1 latencies in WT,
α9KO, and α9KI mice across postnatal development. (**a**) Representative ABR traces at 16
kHz, 80 dB SPL, from WT (black), α9KI (red), and α9KO (blue) mice, recorded at P12–P16, P22,
and P75. At P13, α9KI mice exhibit measurable responses, whereas distinct ABR waves emerge
in WT and α9KO mice by P14. By P22, ABR amplitudes stabilize and remain consistent through
P75. Scale bar applies to all traces. (**b**) ABR wave 1 latencies at 11.33, 16, and 22.65 kHz for
WT, α9KI, and α9KO mice. All groups show a reduction in latency between P14 and P15, though
this change is not statistically significant. Data are shown as mean ± SEM.

METHOD DETAILS

Animals. α9KO and α9KI mice have been previously described (29, 45) and were backcrossed with the congenic FVB.129P2-*Pde6bᵖ Tyrᶜ-ᶜʰ/AntJ* strain (https://www.jax.org/strain/004828) for seventeen generations (i.e., N-17). We maintained a consistent male-to-female ratio across all experimental groups and genotypes. No sex-related differences were observed in the data. Animals were housed in a sound-controlled room within the animal care facility. Background sound levels in this environment were consistently below 50 dB SPL. Noise levels were measured using an electret condenser microphone (FG-23329-PO7; Knowles). Transient sounds during routine cage cleaning and daily maintenance by animal care staff were minimal, with peaks not exceeding 70 dB SPL. The methods reported in this study adhere to the ARRIVE guidelines for reporting *in vivo* animal research (https://arriveguidelines.org/), ensuring transparency and reproducibility. All procedures were conducted in accordance with institutional and ethical guidelines for animal care and use.

Auditory function tests. Physiological assessments of cochlear and central function, including auditory brainstem responses (ABRs) and distortion-product otoacoustic emissions (DPOAEs), were performed in mice anesthetized with xylazine (10 mg/kg, i.p.) and ketamine (100 mg/kg, i.p.) and placed in a soundproof chamber maintained at 32°C to counteract the drop in body temperature caused by anesthesia. ABR represents sound-evoked potentials generated by neural circuits in the ascending auditory pathways, while DPOAEs are sounds emitted by the cochlea, specifically by OHCs, in response to two pure-tone stimuli presented simultaneously. Sound stimuli were delivered through a custom acoustic system with two dynamic earphones used as sound sources (CDMG15008–03A; CUI) and an electret condenser microphone (FG-23329-PO7; Knowles) coupled to a probe tube to measure sound pressure near the eardrum. Data collection was conducted using a National Instruments PXI-based system equipped with 24-

bit input/output boards and operated through a custom LabView software interface. For ABRs,

platinum needle electrodes were placed into the skin at the dorsal midline close to the neural crest

and pinna with a ground electrode near the tail. ABR potentials were evoked with 5 ms tone pips

(0.5 ms rise-fall, with a $\cos^2$ envelope, at 40/s) delivered to the eardrum at log-spaced frequencies

from 5.6 to 45.25 kHz. The response was amplified 10,000X with a 0.3–3 kHz passband. Sound

level was raised in 5 dB steps from 20 to 80 dB sound pressure level (SPL). At each level, 1024

responses were averaged with alternated stimulus polarity. Threshold for ABR was defined as

the lowest stimulus level (<80 dB SPL) at which a repeatable wave 1 could be identified in the

response waveform. The ABR waves 1, 2 and 3 amplitudes were computed by off-line analysis

of the peak-to-peak amplitude of stored waveforms. The DPOAEs in response to two primary

tones of frequency f1 and f2 were recorded at 2f1-f2, with f2/f1=1.2, and the f2 level 10 dB lower

than the f1 level. Ear-canal sound pressure was amplified and digitally sampled at 4 µs intervals.

DPOAE threshold was defined as the lowest f2 level in which the signal to noise floor ratio is >1.

Noise exposure. Animals were exposed under anesthesia at P15 to a 1–16 kHz noise at 100 dB

SPL for 1 hour using closed-field earphones (CDMG15008–03A; CUI) as the sound source. The

exposure took place in the same acoustic chamber used for cochlear function tests. The noise

consisted of a flat-spectrum broadband signal calibrated to the target SPL immediately before

each exposure session.

Cochlear processing and immunostaining. Cochleae were perfused intralabyrinthly with 4%

paraformaldehyde (PFA) in phosphate-buffered saline (PBS), post-fixed with 4% PFA overnight,

and decalcified in 0.12 M EDTA. Cochlear tissues were then microdissected and permeabilized

by freeze/thawing in 30% sucrose. The microdissected pieces were blocked in 5% normal goat

serum with 1% Triton X-100 in PBS for 1 h, followed by incubation in primary antibodies (diluted

in blocking buffer) at 37°C for 16 h. The primary antibodies used in this study were: 1) anti-C-

terminal binding protein 2 (mouse anti-CtBP2 IgG1; BD Biosciences, San Jose, CA; Cat#612044,

RRID:AB_399431, 1:200) to label the presynaptic ribbon, 2) anti-glutamate receptor 2 (mouse

anti-GluA2 IgG2a; Millipore, Billerica, MA; Cat#MAB397, RRID:AB_11212990, 1:2000) to label

the post-synaptic receptor plaques and 3) anti-Myosin 7a (rabbit anti-MyosinVIIa, Proteus

Biosciences; Cat# 25-6790, RRID:AB_10015251, 1:200) to label the cochlear hair cells. Tissues

were then incubated with the appropriate Alexa Fluor-conjugated fluorescent secondary

antibodies (Invitrogen, Carlsbad, CA; 1:500 in blocking buffer) for 2 hours at room temperature.

Finally, tissues were mounted on microscope slides in FluorSave mounting media (Millipore,

Billerica, MA).

Confocal microscopy and image processing. For IHC synaptic counts, confocal z-stacks (0.3

μm step size) of the apical, medial and basal regions from each cochlea were taken using a Leica

TCS SPE microscope equipped with 63X (1.5X or 4X digital zoom) oil-immersion lens. Each stack

usually contained 10 to 20 IHCs. Image stacks were imported to Fiji software

(RRID:SCR_002285) (87) and a custom plugin was developed to automate the counting of

synaptic ribbons, glutamate receptor patches and co-localized synaptic puncta. The algorithm's

accuracy was validated by comparing automated counts to manual counts, ensuring reliable

quantification. Briefly, each channel was analyzed separately, and maximum projections were

generated to quantify the number of CtBP2 or GluA2 puncta. Additionally, a composite between

the three channels was produced to draw the different regions of interest (ROI) that correspond

to each IHC taking the myosin staining as a reference. The maximum projections from the single

channels were multiplied to generate a merged 32-bit image. Then, they were converted to binary

images after a custom thresholding procedure. An automatic counting of the number of particles

on each ROI was performed.

For quantitative volumetric synaptic analysis, different regions from the whole cochlea were

imaged and confocal z-stacks were processed in Fiji. z-stacks were deconvolved applying 10

iterations of the Richardson-Lucy algorithm in the DeconvolutionLab2 plugin (88), an algorithm

that has been used previously in cochlear whole mounts (89). Then, the 3D Object Counter plugin

was used to perform three-dimensional image segmentation and object individualization of the

deconvoluted stacks. According to previous work (56, 90, 91) we reasoned that a synapse would

occupy a volume between 0.04 µm3 to 5 µm3, and then we set a size range of 20-2000 voxels,

with a voxel size of 0.11 µm x 0.11 µm x 0.2 µm. The volumes of presynaptic ribbons (CtBP2

fluorescence) and postsynaptic AMPA receptor clusters (GluA2 fluorescence) at IHC-afferent

synapses were measured using the 3D ROI Manager (92). All exported volumes were checked

to ensure that volumes were within instrument resolution limits after threshold adjustments.

Statistical analysis

Data were analyzed using R Statistical Software (RRID:SCR_001905). Shapiro-Wilks test was

used for testing normal distribution of the residuals. In cases where the data did not follow a

normal distribution, the following tests were used: Kruskal-Wallis test followed by multiple

contrasts according to Dunn's method for comparisons between three groups (IHCs synaptic

counts in the control condition) and Mann-Whitney U test for comparisons between two groups

(ABR thresholds, DPOAE thresholds, and IHCs synaptic counts). If data were normally

distributed, a one-way ANOVA was conducted, followed by Holm-Sidak post hoc tests for three-

group comparisons (wave 1 amplitudes and synaptic volumes in the control condition). For two-

group comparisons, a t-test was used (synaptic volumes). Statistical significance was set to

$p<0.05$.

REFERENCES

1. W. R. Lippe, Rhythmic spontaneous activity in the developing avian auditory system. *J Neurosci* **14**, 1486–1495 (1994).

2. W. Marcotti, S. L. Johnson, A. Rusch, C. J. Kros, Sodium and calcium currents shape action potentials in immature mouse inner hair cells. *J Physiol* **552**, 743–761 (2003).

3. M. Sonntag, B. Englitz, C. Kopp-Scheinpflug, R. Rübsamen, Early Postnatal Development of Spontaneous and Acoustically Evoked Discharge Activity of Principal Cells of the Medial Nucleus of the Trapezoid Body: An In Vivo Study in Mice. *J. Neurosci.* **29**, 9510–9520 (2009).

4. N. X. Tritsch, D. E. Bergles, Developmental regulation of spontaneous activity in the Mammalian cochlea. *J. Neurosci. Off. J. Soc. Neurosci.* **30**, 1539–1550 (2010).

5. S. L. Johnson, *et al.*, Position-dependent patterning of spontaneous action potentials in immature cochlear inner hair cells. *Nat. Neurosci.* **14**, 711–717 (2011).

6. T. S. Tierney, F. A. Russell, D. R. Moore, Susceptibility of developing cochlear nucleus neurons to deafferentation-induced death abruptly ends just before the onset of hearing. *J. Comp. Neurol.* **378**, 295–306 (1997).

7. E. Friauf, C. Lohmann, Development of auditory brainstem circuitry. Activity-dependent and activity-independent processes. *Cell Tissue Res.* **297**, 187–195 (1999).

8. S. P. Mostafapour, S. L. Cochran, N. M. Del Puerto, E. W. Rubel, Patterns of cell death in mouse anteroventral cochlear nucleus neurons after unilateral cochlea removal. *J. Comp. Neurol.* **426**, 561–571 (2000).

9. P. A. Leake, G. T. Hradek, L. Chair, R. L. Snyder, Neonatal deafness results in degraded topographic specificity of auditory nerve projections to the cochlear nucleus in cats. *J. Comp. Neurol.* **497**, 13–31 (2006).

10. A. Clause, *et al.*, The precise temporal pattern of prehearing spontaneous activity is necessary for tonotopic map refinement. *Neuron* **82**, 822–835 (2014).

11. M. N. Di Guilmi, *et al.*, Strengthening of the efferent olivocochlear system leads to synaptic dysfunction and tonotopy disruption of a central auditory nucleus. *J. Neurosci.* 2536–18 (2019). https://doi.org/10.1523/jneurosci.2536-18.2019.

12. W. Marcotti, S. L. Johnson, M. C. Holley, C. J. Kros, Developmental changes in the expression of potassium currents of embryonic, neonatal and mature mouse inner hair cells. *J Physiol* **548**, 383–400 (2003).

13. H. C. Wang, *et al.*, Spontaneous Activity of Cochlear Hair Cells Triggered by Fluid Secretion Mechanism in Adjacent Support Cells. *Cell* **163**, 1348–1359 (2015).

14. S. L. Johnson, J. Olt, S. Cho, H. von Gersdorff, W. Marcotti, The Coupling between Ca^{2+} Channels and the Exocytotic Ca^{2+} Sensor at Hair Cell Ribbon Synapses Varies Tonotopically along the Mature Cochlea. *J. Neurosci.* **37**, 2471–2484 (2017).

15. E. Glowatzki, P. A. Fuchs, Cholinergic synaptic inhibition of inner hair cells in the neonatal
mammalian cochlea. *Science* **288**, 2366–68 (2000).

16. W. Marcotti, S. L. Johnson, C. J. Kros, A transiently expressed SK current sustains and
modulates action potential activity in immature mouse inner hair cells. *J Physiol* (2004).

17. J. D. Goutman, P. A. Fuchs, E. Glowatzki, Facilitating efferent inhibition of inner hair cells
in the cochlea of the neonatal rat. *J. Physiol.* **566**, 49–59 (2005).

18. S. L. Johnson, *et al.*, Cholinergic efferent synaptic transmission regulates the maturation of
auditory hair cell ribbon synapses. *Open Biol.* **3** (2013).

19. Y. Wang, *et al.*, Efferent feedback controls bilateral auditory spontaneous activity. *Nat.
Commun.* **12**, 2449 (2021).

20. D. D. Simmons, N. B. Mansdorf, J. H. Kim, Olivocochlear innervation of inner and outer
hair cells during postnatal maturation: evidence for a waiting period. *J Comp Neurol* **370**,
551–562 (1996).

21. L. G. Vattino, C. Wedemeyer, A. B. Elgoyhen, E. Katz, Functional Postnatal Maturation of
the Medial Olivocochlear Efferent–Outer Hair Cell Synapse. *J. Neurosci.* **40**, 4842–4857
(2020).

22. J. J. Guinan, P. Dallos, A. N. Popper, R. R. Fay, "Efferent Physiology" in *The Cochlea*, P.
Dallos, A. N. Popper, R. R. Fay, Eds. (Springer, 1996), pp. 435–502.

23. R. L. Winslow, M. B. Sachs, Effect of electrical stimulation of the crossed olivocochlear
bundle on auditory nerve response to tones in noise. *J. Neurophysiol.* **57**, 1002–1021
(1987).

24. R. L. Winslow, M. B. Sachs, Single-tone intensity discrimination based on auditory-nerve
rate responses in backgrounds of quiet, noise, and with stimulation of the crossed
olivocochlear bundle. *Hear Res* **35**, 165–189 (1988).

25. T. Kawase, M. C. Liberman, Antimasking effects of the olivocochlear reflex. I.
Enhancement of compound action potentials to masked tones. *J Neurophysiol* **70**, 2519–
2532 (1993).

26. P. H. Delano, D. Elgueda, C. M. Hamame, L. Robles, Selective attention to visual stimuli
reduces cochlear sensitivity in chinchillas. *J Neurosci* **27**, 4146–4153 (2007).

27. M. C. Liberman, The olivocochlear efferent bundle and susceptibility of the inner ear to
acoustic injury. *J Neurophysiol* **65**, 123–132 (1991).

28. S. F. Maison, M. C. Liberman, Predicting Vulnerability to Acoustic Injury with a Noninvasive
Assay of Olivocochlear Reflex Strength. *J. Neurosci.* **20**, 4701–4707 (2000).

29. J. Taranda, *et al.*, A point mutation in the hair cell nicotinic cholinergic receptor prolongs
cochlear inhibition and enhances noise protection. *PLoS Biol.* **7**, 71–83 (2009).

30.   S. F. Maison, H. Usubuchi, M. C. Liberman, Efferent feedback minimizes cochlear
766   neuropathy from moderate noise exposure. *J. Neurosci.* **33**, 5542–5552 (2013).

767   31.   L. E. Boero, *et al.*, Enhancement of the medial olivocochlear system prevents hidden
768   hearing loss. *J. Neurosci.* **38**, 7440–7451 (2018).

769   32.   C. M. Liberman, L. D. Liberman, S. F. Maison, Efferent feedback slows cochlear aging. *J.*
770   *Neurosci.* **34**, 4599–4607 (2014).

771   33.   L. E. Boero, *et al.*, Preventing presbycusis in mice with enhanced medial olivocochlear
772   feedback. *Proc Natl Acad Sci U A* **117**, 11811–11819 (2020).

773   34.   E. F. Chang, M. M. Merzenich, Environmental noise retards auditory cortical development.
774   *Science* **300**, 498–502 (2003).

775   35.   J. Waguespack, F. T. Salles, B. Kachar, A. J. Ricci, Stepwise morphological and functional
776   maturation of mechanotransduction in rat outer hair cells. *J. Neurosci. Off. J. Soc.*
777   *Neurosci.* **27**, 13890–13902 (2007).

778   36.   D. H. Sanes, S. Bao, Tuning up the developing auditory CNS. *Curr Opin Neurobiol* **19**,
779   188–199 (2009).

780   37.   R. C. Froemke, B. J. Jones, Development of auditory cortical synaptic receptive fields.
781   *Neurosci. Biobehav. Rev.* **35**, 2105–2113 (2011).

782   38.   M. Nakamura, P. Valerio, S. Bhumika, T. R. Barkat, Sequential Organization of Critical
783   Periods in the Mouse Auditory System. *Cell Rep.* **32**, 108070 (2020).

784   39.   A. S. Keuroghlian, E. I. Knudsen, Adaptive auditory plasticity in developing and adult
785   animals. *Prog. Neurobiol.* **82**, 109–121 (2007).

786   40.   X. Zhou, M. M. Merzenich, Enduring effects of early structured noise exposure on temporal
787   modulation in the primary auditory cortex. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 4423–4428
788   (2008).

789   41.   A. M. Lauer, B. J. May, B. R. J. M. Ay, A. M. Lauer, B. J. May, The medial olivocochlear
790   system attenuates the developmental impact of early noise exposure. *JARO - J. Assoc.*
791   *Res. Otolaryngol.* **12**, 329–343 (2011).

792   42.   C. Caminos, E. Vale, R. Lujan, J. R. Martinez-Galan, J. M. Juiz, Developmental regulation
793   and adult maintenance of potassium channel proteins (Kv 1.1 and Kv 1.2) in the cochlear
794   nucleus of the rat. *Brain Res.* **1056** (2005).

795   43.   A. L. Dorrn, K. Yuan, A. J. Barker, C. E. Schreiner, R. C. Froemke, Developmental sensory
796   experience balances cortical excitation and inhibition. *Nature* **465**, 932–936 (2010).

797   44.   Y. J. Sun, *et al.*, Fine-tuning of pre-balanced excitation and inhibition during auditory
798   cortical development. *Nature* **465**, 927–931 (2010).

799   45.   D. E. Vetter, *et al.*, Role of alpha9 nicotinic ACh receptor subunits in the development and
800   function of cochlear efferent innervation. *Neuron* **23**, 93-103. (1999).

46. J. S. Buchwald, C. M. Huang, Far-field acoustic response: Origins in the cat. *Science* **189**,
382–384 (1975).

47. M. Karplus, C. Lee, W. J. Cashore, W. Oh, The effects of brain bilirubin deposition on
auditory brain stem evoked responses in rats. *Early Hum. Dev.* **16**, 185–194 (1988).

48. S. M. Shapiro, Acute brainstem auditory evoked potential abnormalities in jaundiced Gunn
rats given sulfonamide. *Pediatr. Res.* **23**, 306–310 (1988).

49. N. A. Shaw, The auditory evoked potential in the rat--a review. *Prog. Neurobiol.* **31**, 19–45
(1988).

50. J. Melcher, N. Y. Kiang, Generators of the brainstem auditory evoked potential in cat. III:
Identified cell populations. *Hear. Res.* **93**, 52–71 (1996).

51. S. E. Kim, K. Turkington, C. Kushmerick, J. H. Kim, Central dysmyelination reduces the
temporal fidelity of synaptic transmission and the reliability of postsynaptic firing during
high-frequency stimulation. *J. Neurophysiol.* **110**, 1621–1630 (2013).

52. C. A. Shera, J. J. Guinan Jr., Evoked otoacoustic emissions arise by two fundamentally
different mechanisms: a taxonomy for mammalian OAEs. *J Acoust Soc Am* **105**, 782–798
(1999).

53. L. Robles, M. A. Ruggero, Mechanics of the mammalian cochlea. *Physiol. Rev.* **81**, 1305–
52 (2001).

54. D. Khimich, *et al.*, Hair cell synaptic ribbons are essential for synchronous auditory
signalling. *Nature* **434**, 889–894 (2005).

55. A. Matsubara, J. H. Laake, S. Davanger, S. Usami, O. P. Ottersen, Organization of AMPA
receptor subunits at a glutamate synapse: a quantitative immunogold analysis of hair cell
synapses in the rat organ of Corti. *J. Neurosci.* **16**, 4457–4467 (1996).

56. L. D. Liberman, H. Wang, M. C. Liberman, Opposing gradients of ribbon size and AMPA
receptor expression underlie sensitivity differences among cochlear-nerve/hair-cell
synapses. *J. Neurosci. Off. J. Soc. Neurosci.* **31**, 801–8 (2011).

57. G. Sendin, J. Bourien, F. Rassendren, J.-L. Puel, R. Nouvian, Spatiotemporal pattern of
action potential firing in developing inner hair cells of the mouse cochlea. *Proc. Natl. Acad.*
Sci. U. S. A. **111**, 1999–2004 (2014).

58. M. Youssoufian, Development of a Robust Central Auditory Synapse in Congenital
Deafness. *J. Neurophysiol.* **94**, 3168–3180 (2005).

59. R. N. Leao, *et al.*, Topographic organization in the auditory brainstem of juvenile mice is
disrupted in congenital deafness. *J. Physiol.* **571**, 563–578 (2006).

60. J. J. Hirtz, *et al.*, Cav1.3 calcium channels are required for normal development of the
auditory brainstem. *J. Neurosci. Off. J. Soc. Neurosci.* **31**, 8280–8294 (2011).

61.   T. A. Babola, *et al.*, Purinergic Signaling Controls Spontaneous Activity in the Auditory
837   System throughout Early Development. *J. Neurosci. Off. J. Soc. Neurosci.* **41**, 594–612
838   (2021).

839   62.   T. Frank, D. Khimich, A. Neef, T. Moser, Mechanisms contributing to synaptic Ca2+
840   signals and their heterogeneity in hair cells. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 4483–4488
841   (2009).

842   63.   L. Sheets, K. S. Kindt, T. Nicolson, Presynaptic CaV1.3 channels regulate synaptic ribbon
843   size and are required for synaptic maintenance in sensory hair cells. *J. Neurosci. Off. J.*
844   *Soc. Neurosci.* **32**, 17273–17286 (2012).

845   64.   H. C. Wong, *et al.*, Synaptic mitochondria regulate hair-cell synapse size and function.
846   *eLife* **8**, e48914 (2019).

847   65.   Ö. D. Özçete, T. Moser, A sensory cell diversifies its output by varying Ca2+ influx-release
848   coupling among active zones. *EMBO J.* **40**, e106010 (2021).

849   66.   S. Michanski, *et al.*, Mapping developmental maturation of inner hair cell ribbon synapses
850   in the apical mouse cochlea. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 6415–6424 (2019).

851   67.   Y. Hua, *et al.*, Electron Microscopic Reconstruction of Neural Circuitry in the Cochlea. *Cell*
852   *Rep.* **34**, 108551 (2021).

853   68.   S. A. Payne, *et al.*, Maturation of Heterogeneity in Afferent Synapse Ultrastructure in the
854   Mouse Cochlea. *Front. Synaptic Neurosci.* **13**, 678575 (2021).

855   69.   H. M. Sobkowicz, J. E. Rose, G. E. Scott, S. M. Slapnick, Ribbon synapses in the
856   developing intact and cultured organ of Corti in the mouse. *J Neurosci* **2**, 942–57. (1982).

857   70.   A. Merchan-Perez, M. C. Liberman, Ultrastructural differences among afferent synapses
858   on cochlear hair cells: correlations with spontaneous discharge rate. *J. Comp. Neurol.* **371**,
859   208–221 (1996).

860   71.   L.-C. Huang, *et al.*, Synaptic profiles during neurite extension, refinement and retraction in
861   the developing cochlea. *Neural Develop.* **7**, 38 (2012).

862   72.   A. B. Wong, *et al.*, Developmental refinement of hair cell synapses tightens the coupling of
863   Ca2+ influx to exocytosis. *EMBO J.* **33**, 247–264 (2014).

864   73.   L. Song, J. McGee, E. J. Walsh, Frequency- and level-dependent changes in auditory
865   brainstem responses (ABRS) in developing mice. *J. Acoust. Soc. Am.* **119**, 2242–2257
866   (2006).

867   74.   K. A. Fernandez, P. W. C. Jeffers, K. Lall, M. C. Liberman, S. G. Kujawa, Aging after noise
868   exposure : Acceleration of cochlear synaptopathy in " recovered " ears. *J. Neurosci.* **35**,
869   7509–20 (2015).

870   75.   K. M. Schrode, M. L. Dent, A. M. Lauer, Sources of variability in auditory brainstem
871   response thresholds in a mouse model of noise-induced hearing loss. *J. Acoust. Soc. Am.*
872   **152**, 3576–3582 (2022).

76. Y. Lu, *et al.*, Spatial patterns of noise-induced inner hair cell ribbon loss in the mouse mid-
cochlea. *iScience* **27**, 108825 (2024).

77. N. Rybalko, *et al.*, Development of the acoustic startle response in rats and its change after
early acoustic trauma. *Behav. Brain Res.* **286**, 212–221 (2015).

78. L. Shi, *et al.*, Noise-induced damage to ribbon synapses without permanent threshold
shifts in neonatal mice. *Neuroscience* **304**, 368–377 (2015).

79. N. I. Mohamad, *et al.*, Synaptic ribbon dynamics after noise exposure in the hearing
cochlea. *Commun. Biol.* **7**, 421 (2024).

80. D. J. Moverman, L. D. Liberman, S. Kraemer, G. Corfas, M. C. Liberman, Ultrastructure of
noise-induced cochlear synaptopathy. *Sci. Rep.* **13**, 19456 (2023).

81. S. Stamataki, H. W. Francis, M. Lehar, B. J. May, D. K. Ryugo, Synaptic alterations at
inner hair cells precede spiral ganglion cell loss in aging C57BL/6J mice. *Hear. Res.* **221**,
104–118 (2006).

82. J. Y. Jeng, *et al.*, Pathophysiological changes in inner hair cell ribbon synapses in the
ageing mammalian cochlea. *J. Physiol.* **598**, 4339–4355 (2020).

83. L. E. Boero, S. Payne, M. E. Gómez-Casati, M. A. Rutherford, J. D. Goutman, Noise
Exposure Potentiates Exocytosis From Cochlear Inner Hair Cells. *Front. Synaptic
Neurosci.* **13**, 740368 (2021).

84. K. Suthakar, M. C. Liberman, Auditory-Nerve Responses in Mice with Noise-Induced
Cochlear Synaptopathy. *https://doi.org/10.1152/jn.00342.2021* (2021).
https://doi.org/10.1152/JN.00342.2021.

85. M. B. Wood, *et al.*, Acoustic Trauma Increases Ribbon Number and Size in Outer Hair
Cells of the Mouse Cochlea. *J. Assoc. Res. Otolaryngol. JARO* **22**, 19–31 (2021).

86. A. C. Furman, S. G. Kujawa, M. C. Liberman, Noise-induced cochlear neuropathy is
selective for fibers with low spontaneous rates. *J. Neurophysiol.* **110**, 577–586 (2013).

87. J. Schindelin, *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat.
Methods* **9**, 676–82 (2012).

88. D. Sage, *et al.*, DeconvolutionLab2: An open-source software for deconvolution
microscopy. *Methods* **115**, 28–41 (2017).

89. Z. Jing, *et al.*, Disruption of the presynaptic cytomatrix protein bassoon degrades ribbon
anchorage, multiquantal release, and sound encoding at the hair cell afferent synapse. *J.
Neurosci.* **33**, 4456–4467 (2013).

90. F. Gilels, S. T. Paquette, J. Zhang, I. Rahman, P. M. White, Mutation of foxo3 causes adult
onset auditory neuropathy and alters cochlear synapse architecture in mice. *J. Neurosci.*
33, 18409–18424 (2013).

91. S. T. Paquette, F. Gilels, P. M. White, Noise exposure modulates cochlear inner hair cell
909   ribbon volumes, correlating with changes in auditory measures in the FVB/nJ mouse. *Sci.*
910   *Rep.* **6**, 1–13 (2016).

911   92. J. Ollion, J. Cochennec, F. Loll, C. Escudé, T. Boudier, TANGO: A generic tool for high-
912   throughput 3D image analysis for studying nuclear organization. *Bioinformatics* **29**, 1840–
913   1841 (2013).