Zero-phase-delay synchrony between interacting neural populations: implications for functional connectivity derived biomarkers

Chirag Mehra[a], Ahmad Beyh[a,b], Petroula Laiou[c,d], Pilar Garces[e], Emily JH Jones[f,g], Luke Mason[a], Jan Buitelaar[h], Mark H Johnson[i], Declan Murphy[a], Eva Loth[a], Flavio Dell'Acqua[a], Joshua B Ewen[j,k], Mark P Richardson[c], Jonathan O'Muircheartaigh[a,l]

a) Department of Forensic and Neurodevelopment Sciences, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK
b) Department of Psychiatry, Brain Health Institute, Rutgers University, Piscataway, USA
c) Department of Basic and Clinical Neuroscience, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK
d) Department of Biostatistics & Health Informatics, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK
e) Roche Pharma Research and Early Development, Neuroscience and Rare Diseases, Roche Innovation Center Basel, F. Hoffmann–La Roche Ltd., Basel, Switzerland
f) Department of Child and Adolescent Psychiatry, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK
g) MRC Centre for Developmental Neurobiology, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK
h) Department of Cognitive Neuroscience, Donders Institute for Brain, Cognition and Behaviour, Radboud University Medical Centre, Nijmegen, Netherlands.
i) Department of Psychology, University of Cambridge, Cambridge, UK
j) Developmental-Behavioral Pediatrics, Ann & Robert H. Lurie Children's Hospital of Chicago, Chicago, USA
k) Department of Pediatrics at Northwestern University Feinberg School of Medicine, Chicago, Illinois, USA
l) Department of Neuroimaging, King's College London, London, UK

Corresponding author: Chirag Mehra, chirag.1.mehra@kcl.ac.uk

Abstract

Neural populations synchronise their activity with either zero-phase-delay (activity in interacting regions occurs simultaneously) or a phase-delay (activity in one region follows the other). In electroencephalography and magnetoencephalography functional connectivity analyses, artefactual connectivity can also occur with zero-phase-delay. To minimise artefact, contemporary analyses typically exclude all zero-phase-delay interactions. However, the extent to which 'true' interactions are resultingly lost – and the impact this has on the performance of functional connectivity metrics as biomarkers – remains unknown. Here, we show that most cortico-cortical functional connectivity occurs with zero- or near-zero phase-delay, even where such connectivity is unlikely to be artefactual. Including, rather than excluding, zero-phase-delay connectivity increases the reliability, convergence with neurobiology (structure-function concordance, homotopic interhemispheric connectivity, and age-related connectivity changes), and prognostic ability of functional connectivity metrics. We find that excluding zero-phase-delay connections penalises functional connectivity strength between the strongest structurally connected regions: stronger structural connections lead to functional connections with phase-delays closer to zero, mediated by a shorter signal propagation time. Our findings challenge generally accepted assumptions that zero-phase-exclusive methods are superior to zero-phase-inclusive methods.

Keywords: functional connectivity, biomarker, electroencephalography, zero-phase-delay, zero-phase-lag, signal leakage

Introduction

Populations of neurons produce oscillatory electromagnetic activity (Bishop, 1932; Buzsáki, 2002). Through the precise synchronisation and integration of these oscillations, the brain produces cognitive function, sensory experience and behaviour (Buzsáki, 2006; Lopes da Silva, 2013; Rodriguez et al., 1999a; Singer, 1999; Uhlhaas et al., 2009; Varela et al., 2001; Womelsdorf & Fries, 2007). Studying these interactions, or functional connectivity, has provided insight into several cognitive processes (Dai et al., 2017a; Hampson et al., 2006; Kang et al., 2011) and neuropsychiatric disorders (Du et al., 2018). Indeed, metrics derived from functional connectivity analyses have been proposed as candidate diagnostic or prognostic biomarkers for brain-based conditions (Du et al., 2018; Gao et al., 2020; Gholipour et al., 2022; Gil Ávila et al., 2023; Hong et al., 2020; Plitta et al., 2015).

The averaged oscillatory electromagnetic activity of neural populations can be non-invasively measured by electroencephalography (EEG) and/or magnetoencephalography (MEG; Singer, 1999). Brain-regions are assumed to be functionally connected if the phase and/or amplitude (Engel et al., 2013; Varela et al., 2001) of their oscillations are statistically interdependent (Friston, 2011). Functionally connected brain regions can have perfectly synchronous activity (with zero-phase-delay), or activity in one region may follow the other (producing a phase-delay). However, the extent to which measured zero-phase-delay connectivity reflects 'true' versus artefactual connectivity in empirical signals remains unknown.

There is robust evidence demonstrating the presence of 'true' zero-phase-delay functional connectivity, replicated across species (Engel et al., 1991; O'Reilly & Elsabbagh, 2021; Vicente et al., 2008; Witham et al., 2007). Spike-train recordings (Engel et al., 1991) have demonstrated that most functional connections between homotopic interhemispheric regions occur with zero-phase-delay. Computational models suggest that "resonance-induced synchrony" (Gollo et al., 2014) facilitates zero-phase-delay connectivity: bidirectional information transfer between regions acts to mutually alter their dynamics into a stable zero-phase-delay pattern. Functions dependent on zero-phase-delay connectivity include maximizing the reliability of information transmission and facilitating spike-time-dependent plasticity (Gollo et al., 2014), coding and binding features of sensory objects (Gray et al., 1989; Rodriguez et al., 1999b; Singer, 1999), and others (Roelfsema et al., 1997; Steinmetz et al., 2000). Evidence from invasive recordings also shows near-zero-phase-delay connectivity (Uhlhaas et al., 2009), describing phase-delays that are much smaller than those arising from axonal conduction delays. An exemplar function of near-zero-phase-delay connectivity is neural encoding by hippocampal place cells (O'Keefe & Recce, 1993).

Artefactual functional connectivity due to signal leakage and volume conduction also occurs with zero-phase-delay (Brunner et al., 2016; Hipp et al., 2012; Nolte et al., 2004; Srinivasan et al., 2007), as spatial blurring in signal leakage (Anzolin et al., 2019; Colclough et al., 2015; Hipp et al., 2012; Srinivasan et al., 2007) and field-spread in volume conduction (Anzolin et al., 2019; Bastos & Schoffelen, 2016; Nolte et al., 2004) occur with zero- or almost-zero-time delay. This artefactual connectivity is unavoidable in EEG and MEG data. It mimics 'true' connectivity (Anzolin et al., 2019) and may

101 dominate the pattern of connectivity measurements (Hipp et al., 2012; Srinivasan et al.,
102 2007).
103
Therefore, contemporary functional connectivity analyses typically exclude zero-phase-
delay connections, prioritising the goal of limiting artefactual connectivity due to signal
leakage and volume conduction (Colclough et al., 2016; Miljevic et al., 2022). However,
the extent to which biologically relevant 'true' connectivity is also lost when excluding
zero-phase-delay connectivity is unknown. Including versus excluding zero-phase-
delay connectivity may substantially alter the conclusions drawn from otherwise
identical signals (Cociu et al., 2018), hindering progress in biomarker discovery (for
example, O'Reilly et al., 2017). No study, to our knowledge, has systematically
assessed the effect of including versus excluding zero-phase-delay connections on the
performance of functional connectivity metrics as candidate biomarkers.
114
Here, using empirical EEG data, we used a novel approach to quantify the proportion of
zero- and near-zero phase-delay connectivity when signal leakage artefact was likely
negligible. Then, we compared the effects of including versus excluding zero-phase-
delay functional connectivity on properties desired in biomarkers: high test-retest
reliability (Davis et al., 2020) and convergence with underlying biology (FDA-NIH
Biomarker Working Group, 2016). Finally, we compared the performance of functional
connectivity metrics derived from zero-phase inclusive versus exclusive methods as
prognostic biomarkers for longitudinal changes in cognitive abilities. Thus, our analyses
leveraged multiple performance criteria not trivially correlated with each other.

Methods

Participants

125
We included participants from the AIMS-2-TRIALS Longitudinal European Autism
Project, described here (Charman et al., 2017; Loth et al., 2017). Written informed
consent was obtained from all participants for being included in the study. In this
analysis, we included 153 neurotypical participants (33% female), aged 6-31 years,
from 5 study sites. For the analyses predicting longitudinal changes in spatial working
memory, participants were split into children (6-12 years, n = 41), adolescents (13-17
years, n = 50) and adults (18-31 years, n = 62), Table 1. Included participants needed at
least 20 seconds of usable eyes-closed resting-state EEG data, a good quality (defined
below) T1-weighted (T1w) magnetic resonance spectroscopy (MRI) scan image, and a
full-scale intelligence quotient $\geq$75. We excluded participants with diagnosed
neurodevelopmental conditions, intellectual disability, neurological conditions, mental
health conditions (e.g. schizophrenia, bipolar affective disorder) and significant
systemic disease (e.g. auto-immune conditions). Intelligence quotient was measured
by the Wechsler Abbreviated Scale of Intelligence, Second Edition (Wechsler, 2011).
140

Table 1 – Characteristics of included participants. This study included 153 participants aged between 6 and 31 years. For the analyses predicting longitudinal changes in spatial working memory, participants were grouped by age: children (6-12 years), adolescents (13-17 years), and adults (18-31 years).					
	6-12 yrs (n = 41)	13-17 yrs (n = 50)	18-31 yrs (n = 62)	Total (n = 153)	*p* value
Age (years)					
Mean (SD)	10.2 (1.9)	15.9 (1.4)	23.4 (3.5)	17.4 (6.0)	-
Range	6 - 12	13 - 18	18 – 31	6 – 31	
Sex					
Female	15 (37%)	20 (40%)	16 (26%)	51 (33%)	.25
Full Scale IQ					
Mean (SD)	111 (12)	108 (12)	109 (12)	109 (12)	.41
Range	76 – 139	77- 133	85 – 142	76 – 142	
Site					
London	12	7	24	43	< .001
Mannheim	7	18	3	28	
Nijmegen	13	17	8	38	
Rome	0	0	16	16	
Utrecht	9	8	11	28	

Electroencephalography methods

EEG artefact cleaning and source localisation were performed as per our group's

previous work (Garcés et al., 2022).

EEG data acquisition: Four segments of 30 second, eyes-closed EEG recordings were

captured. These short epochs were used to maximise comfort for young participants.

Data were acquired at five sites: Central Institute of Mental Health (CIMH, Mannheim,

Germany), King's College London (KCL, United Kingdom), University Nijmegen Medical

Centre (RUNMC, Netherlands), University Campus BioMedico (UCBM, Rome, Italy) and

University Medical Centre Utrecht (UMCU, Netherlands). The following EEG systems

were employed: Brainvision (CIMH, KCL, RUNMC), Biosemi (UMCU) and Micromed

(UCBM), with sampling frequencies of 5000 Hz (KCL, RUNMC), 2048 Hz (UMCU), 2000

Hz (CIMH) and 256–1000 Hz (UCBM). All sites used 10–20 layout caps, with 60–70

electrodes.

EEG artefact cleaning: Only data from the 61 electrodes commonly used across all

sites were retained. The following steps were performed: (1) electrode channels with

poor data quality were removed, (2) temporal segments with large transient artifacts

such as muscle bursts or movements were discarded, (3) independent component

analyses were applied using fastICA (Hyvärinen, 1999) to distinguish signal from noise,

(4) artefactual independent components were identified and eliminated, (5) and

channels discarded in (1) were interpolated. The common average signal across all
channels was used as the reference. If more than 10 cannels were eliminated in step 1,
the epoch was not included in subsequent analyses.
166
Prior to source localisation, initial downsampling and bandpass filtering were
performed, as per Hyvärinen (1999). Except for signals from UCBM (which were
recorded at 256 Hz and not initially downsampled), EEG signals were downsampled to
1000 Hz, with an antialiasing filter (low-pass finite impulse response filter, Kaiser
window, cutoff of 300 Hz, transition band width 100 Hz), using EEGlab (Delorme &
Makeig, 2004). Then, the entire EEG resting-state recording (all four 30 second epochs
and the segments in between them), with 10 seconds of data padding, was bandpass
filtered between 1–32 Hz with a finite impulse response filter of order 2000, using a
Hamming window, in both forward and backward directions, with detrending, using
FieldTrip (Oostenveld et al., 2011).
177
Source Reconstruction: Source reconstruction was performed using individual-
participant T1-weighted MRI images (acquisition parameters detailed below). T1-
weighted MRIs were segmented with Statistical Parametric Mapping 12 (SPM12)
software (Penny W et al., 2006) into grey matter, white matter, cerebrospinal fluid, bone,
soft tissue and air. These probabilistic images were then smoothed (5mm full width half
maximum), thresholded and resliced to produce binary masks of 2mm × 2mm × 2mm
resolution for three tissue types: brain (including grey matter, white matter and
cerebrospinal fluid), skull and scalp. These binary masks were transformed to
hexahedral meshes with FieldTrip (Oostenveld et al., 2011). All three considered tissue
types were assumed to have homogeneous and isotropic conductivity: 330 ms/m for
the brain and scalp (Gabriel et al., 1996), and an age-dependent skull conductivity of
$3.958 + 62.77 \times e^{-0.2404 \times age\,in\,years}$ ms/m, in line with the BESA (*BESA® | Brain Electrical*
Source Analysis: Home, 1995) recommended conductivity ratios. Segmentations were
visually inspected. The forward model was derived with FieldTrip and SimBio (Vorwerk
et al., 2018). For the inverse model, 1200 source locations of interest were defined in
grey matter in Montreal Neurological Institute (MNI) space, following a 3D cubic
diamond grid. Source positions were transformed from MNI space to each subject's
individual space with a nonlinear transformation using SPM12. Electrode positions were
determined by transforming standard MNI positions to subject-space with the same
transformation, then projecting to the scalp surface. Source time series were estimated
with linearly constrained minimum variance beamformer (Anzolin et al., 2019), using a
regularization of 5% of the average trace of the covariance matrix. The 1200 sources
were parcelled into 68 cortical regions of interest, as per the Desikan-Killiany atlas
(Desikan et al., 2006). Each region was comprised of between 1 and 67 sources (median
= 13). When there was more than one source per region, the representative time series
for a given region was defined as the first principal component of all the source time
series in it.
205
Cropping, down-sampling and filtering
We chose 20 second eyes-closed resting-state EEG epochs (from a maximum segment
length of 30 seconds) to produce functional connectivity adjacency matrices. This
epoch length was chosen as a compromise between maximising the number of

included participants and maximising network stability (Fraschini et al., 2016; Pullon et
al., 2020).
212
213 We then further down-sampled the signals to 250Hz. Data quality checks were
214 subsequently performed.
215
216 We treated the whole 20 second epoch as a single window. Producing multiple windows
217 has the advantage of better capturing moment-to-moment connectivity. However,
218 longer window-length better captures the 'ground truth' (Pullon et al., 2020).
219 Additionally, longer windows are needed to capture weaker connections (Pullon et al.,
220 2020) due to the noise inherent to EEG signals.
221
The signals were then filtered into the conventional frequency bands: delta (1-4 Hz),
theta (4-8 Hz), alpha (8-13 Hz), low-beta (13-20 Hz) and high-beta (20-32 Hz). High-pass
followed by low-pass filters were used in sequence, using two-pass (using the MATLAB
function filtfilt), finite impulse response filters, of order 250. To reduce signal distortion,
data padding of 500 samples (2 seconds) was used during filtering and while performing
Hilbert transformations.
228
Quantifying phase-delays
To quantify the phase-delay between two signals $x_i(t)$ and $x_j(t)$, $t = 1, ..., N$ where
N is the number of points in an EEG epoch, we first applied the Hilbert transform to
obtain the instantaneous phases $\theta_i(t), \theta_j(t)$ for the signals $x_i(t)$ and $x_j(t)$ respectively.
Then, the phase-delay distribution $\Delta\theta(t) = \theta_i(t) - \theta_j(t)$ was projected into the unit
circle. The instantaneous phase-delays across samples in an epoch were averaged to
produce the mean phase-delay throughout the epoch:
236

237
$$\tau_{i,j} = angle(\frac{1}{N}\sum_{t=1}^{N} e^{i*\Delta\theta(t)})$$

238
Zero- and ±π-phase-delay functional connectivity
Note that 'true' ±π-phase-delay functional connectivity also occurs between region-
pairs where 'true' zero-phase-delay connectivity is prevalent (Ni et al., 2009; O'Reilly &
Elsabbagh, 2021; Palmer et al., 2012), and has important functions (Hilgetag et al.,
2001; Ni et al., 2009; O'Reilly & Elsabbagh, 2021; Palmer et al., 2012). As ±π-phase-
delay functional connectivity is also excluded by zero-phase-exclusive functional
connectivity methods, for brevity, zero-phase-delay and π-phase-delay are
subsequently referred to together as 'zero-phase-delay.'
247
Investigating where the effects of signal leakage are likely negligible
We graphically examined the relationship between the proportion of zero and near-zero
phase-delay functional connectivity and Euclidean distance between region-pairs.
Artefactual connectivity due to signal leakage occurs at zero-phase-delay (Gohel et al.,
2017; Hipp et al., 2012) and its influence is a function of Euclidean distance between
sources (Bastos & Schoffelen, 2016; Gohel et al., 2017; He et al., 2019; Hipp et al.,
2012). Therefore, we explored if there were Euclidean distances between brain regions

at which the proportion of zero- and near-zero phase-delay connectivity did not vary as
a function of Euclidean distance. Such a phenomena would suggest that artefactual
connectivity due to signal leakage is likely negligible at these Euclidean distances.
258
259 Euclidean distance between region-pairs was measured from the centroids of the
260 regions, obtained from the FreeSurfer surface parcellation, measured per participant
261 using individual T1w MRI images.
262
In the absence of an established definition of 'near-zero-phase-delay' functional
connectivity, we defined near-zero-phase-delay as within < | 0.3 | radians from 0 or ±π
radians. These phase-delays comprise approximately 19% of possible phase-delays in
a unit circle (between -π and π). For comparison, the phase delay caused by signal
transmission delays along a (white matter) streamline of length 128mm (the median
streamline length in our dataset), assuming a white matter transmission speed of
4.5m/s (van Blooijs et al., 2023) and a synaptic transmission delay of 0.75 milliseconds
(Katz & Miledi, 1965), for oscillations at 10 Hz, is approximately 1.83 radians. Note that
by design, all zero-phase-exclusive methods alter the functional connectivity value
based on how close phase-delay is to 0 or π. Connectivity strength values are only
unaltered by zero-phase-exclusive methods at phase delays of ±π/2 radians, and
approach zero as phase-delay approaches 0 or ±π. At phase delays of 0.3 radians, the
measured connectivity strength of 'true' (and artefactual) functional connectivity is
decreased by 70% using the wPLI and Img COH compared to when using zero-phase-
inclusive methods.
278
Functional Connectivity Methods
We limited our selection of methods for comparison to those that characterise
functional connectivity based on phase relationships, amplitude relationships, or a
combination thereof; do not make assumptions about directionality of connectivity
based on the sign of phase difference (Stam & van Straaten, 2012); are bivariate; and
are commonly used. Thus, we selected phase locking value (PLV; Lachaux et al., 1999),
amplitude envelope correlation (AEC; Bruns et al., 2000) and coherence (Adey et al.,
1967) as zero-phase-delay inclusive methods. We selected their methodologically
corresponding zero-phase-delay exclusive methods: weighted phase lag index (wPLI,
Vinck et al., 2011), orthogonalized amplitude envelope correlation (Orth AEC; Hipp et
al., 2012) and the imaginary part of coherency (Img COH; Nolte et al., 2004),
respectively.
291
292 **Coherence and imaginary part of coherency**
293 Coherence measures the consistency of relative amplitude and phase between pairs of
294 signals at a given frequency.
295
296 Coherence is the absolute value of coherency, $C_{ij}(f)$, between signals *i* and *j*, defined as:
297

298 $$C_{ij}(f) \equiv \frac{S_{ij}(f)}{(S_{ii}(f)S_{jj}(f))^{1/2}}$$

Where $S_{ij}(f)$ is the cross spectral density of signals i and j, $S_{ii}(f)$ is the power spectral density of signal i, and $S_{jj}(f)$ is the power spectral density of signal j. We used the Welch method to calculate power spectral density, as it has the advantage of handling signal leakage effects well (Welch, 1967).

Therefore, coherence Coh_{ij}(f), is:

$$Coh_{ij}(f) \equiv \left| \frac{S_{ij}(f)}{(S_{ii}(f)S_{jj}(f))^{1/2}} \right|$$

The imaginary part of coherency only considers the imaginary component of the complex-valued cross spectral density:

$$Img\ Coh_{ij}(f) \equiv \left| \frac{Img(S_{ij}(f))}{(S_{ii}(f)S_{jj}(f))^{1/2}} \right|$$

Phase Locking value
Phase-based connectivity methods assume that two brain regions are connected if the rhythm of their oscillations are similar, largely independent of amplitude relationships.

The phase locking value between the signals $x_i(t)$ and $x_j(t)$, $t = 1, ..., N$ where N is the number of points in an epoch is defined as:

$$PLV_{ij} = \frac{1}{N} \left| \sum_{t=1}^{N} e^{i*\Delta\theta(t)} \right|$$

Where $\Delta\theta(t)$ is the difference of the instantaneous phases $\theta_i(t), \theta_j(t)$ of the signals $x_i(t), x_j(j)$. Here, we did not perform the permutation testing that is typical for the PLV (Lachaux et al., 1999), to enable comparisons with the other structural and functional connectivity methods used in this analysis.

Weighted Phase Lag Index
The weighted phase lag index is derived by multiplying the phase lag index value by a multiplier based on the amplitude of the phase-difference.

The weighted phase lag index between two signals i and j is defined as:

$$wPLI_{ij} \equiv \frac{\left| E\left\{ Im\left(S_{ij}(f) \right) \right\} \right|}{E\left\{ \left| Im\left(S_{ij}(f) \right) \right| \right\}} = \frac{\left| E\left\{ \left| Im\left(S_{ij}(f) \right) \right| sgn\left(Im\left(S_{ij}(f) \right) \right) \right\} \right|}{E\left\{ \left| Im\left(S_{ij}(f) \right) \right| \right\}},$$

where S_{ij} is the cross spectrum between the signals i and j.

Amplitude envelope correlation and orthogonalized amplitude envelope correlation

340 The amplitude envelope correlation (AEC) detects amplitude relationships between
341 signals, independent of phase relationships. The AEC between region-pairs i and j is the
342 magnitude of the Pearson correlation coefficient between the absolute values of Hilbert
343 transformed signal i and the absolute values of Hilbert transformed signal j.
344
345 In the orthogonalized AEC, pairs of signals are orthogonalized with respect to each other
346 prior to carrying out the AEC calculation. Orthogonalization was performed (on a per-
347 sample basis) in two directions (i to j, j to i), producing $i_{\perp j}$ and $j_{\perp i}$ timeseries, as per Hipp
348 and colleagues (2012). $i_{\perp j}$, for example, was derived by:
349
350 $$i_{\perp j}(t,f) = i(t,f) - i_{\parallel j}(t,f),$$
351 when
352
353 $$i_{\parallel j}(t,f) = a_{i,j}(f,T)j(t,f),$$
354
Where $a_{i,j}$ is the regression coefficient obtained by ordinary least squares describing the
linear relationship between i and j throughout the epoch (t = 20). The orthogonalized
time series were then squared and log transformed, to make the time series more
normal, as per Hipp and colleagues (2012). Then, power envelope correlations were
computed between i and $j_{\perp i}$, and j and $i_{\perp j}$ – the mean of the two correlation values was
taken to be the undirected orthogonalized connectivity value between i and j.
361
Test-retest reliability
We quantified the intrasession test-retest reliability between two epochs recorded
between 1-3 minutes of each other (n = 99). Intrasession test-retest reliability was used
to reduce the confounding effects of circadian rhythm (Fafrowicz et al., 2019) and
caffeine (Haller et al., 2013) on functional connectivity. Two measures of test-retest
reliability were used: absolute agreement and consistency. Edgewise absolute
agreement was quantified using the interclass correlation coefficient (ICC (2,1)). ICC
values were interpreted as per Koo & Li (2016): 0.9 -1.0 indicate excellent reliability,
0.75-0.9 indicate good reliability, 0.5-0.75 indicate moderate reliability and 0.0-0.5
indicate poor reliability. Negative values were given a value of 0. We quantified the
consistency of the whole adjacency matrix using Spearman's correlation coefficient, as
per van der Velde and colleagues (2019).
374
Structural Connectivity
Structural connectivity between 68 grey matter regions, as parcellated by the Desikan-
Killiany atlas (Desikan et al., 2006), was quantified by using spherical deconvolution
diffusion weighted imaging. Diffusion imaging data was available from 2 of 5 study sites
(KCL and Mannheim), comprising 50 participants (Table 2a). Spherical deconvolution
diffusion imaging was used to overcome the crossing fibres problem that is a key
limitation of traditional diffusion tensor imaging (Dell'Acqua et al., 2013). Connectivity
strength between two regions was based on the streamline count between them,
normalised to the value of 1, as per Chu and colleagues (2015).
384
T1-weighted cortical reconstruction using FreeSurfer

Signals from diffusion imaging were projected on to the cortical surface derived from
each participant's T1w MRI image. As previously done by our group (Pretzsch et al.,
2022), all T1 scans were manually inspected; those with visible anomalies or significant
movement artefacts were excluded. For each T1w MRI, models of the cortical surface
using FreeSurfer v6.0 (https://surfer.nmr.mgh.harvard.edu/) were computed. A detailed
description of these well-validated and automated procedures is provided by Fischl
(2012). In brief, FreeSurfer performs intensity normalization, skull stripping, and an
image segmentation using a connected-components algorithm. Next, it generates a
filled white matter volume for each hemisphere and, by fitting a deformable template, a
surface tessellation for each of these volumes. For each subject, this yields a mesh (of
triangular elements) for the inner (white-matter) and outer (pial) cortical surface
consisting of ~150k vertices (points in each triangular element) per hemisphere. Each of
these reconstructed surfaces was inspected visually for reconstruction errors by three
independent and experienced researchers. Scans were either accepted or manually
edited after automatic pre-processing. Manually edited images were iteratively re-pre-
processed and visually re-assessed.

Table 2a – demographics of participants who had both EEG and diffusion weighted imaging data. FSIQ: full scale intellectual quotient

Participants	KCL n = 28	Mannheim n = 22
Mean Age (SD)	23 (4.7)	15 (2.9)
Age Range	15 - 31	8 - 19
Mean FSIQ (SD)	109 (13.3)	110 (13.3)
FSIQ Range	85 – 142	77 – 133

Table 2b. Diffusion MRI acquisition parameters

	KCL	Mannheim
Scanner	GE MR750	Siemens TimTrio
Sequence	SE-EPI	SE-EPI
Field of view	256×256 mm^2	256×256 mm^2
Matrix size	128×128	128×128
Number of slices	72	72
Voxel size	$2 \times 2 \times 2$ mm^3	$2 \times 2 \times 2$ mm^3
Echo time	67 ms	102 ms
Repetition time	12 s	12 s
Flip angle	90	90
B-value	1500 s/mm^2	1500 s/mm^2
Diffusion directions	60	60
No. b=0 images	6	6

Diffusion MRI pre-processing

Diffusion MRI acquisition parameters are detailed in Table 2b. Tools from Connectome Workbench (https://www.humanconnectome.org/software) were used to re-tessellate the T1w surfaces of each participant to the common tessellation of the '32k_FS_LR' template. This was done to reduce the number of vertices in each surface to 32,492 points, reducing the computational burden of the cortical projection step.

The diffusion series of each participant was corrected for thermal noise (Veraart et al., 2016) and Gibbs ringing artefacts (Kellner et al., 2016). Due to the lack of reversed phase encoding images, the *SynB0* framework (Schilling et al., 2019) was used to generate undistorted b=0 EPIs and these were used to estimate the susceptibility distortion field using *topup* (Andersson et al., 2003). Next, *eddy* was used to correct the diffusion series for artefacts caused by frame-wise motion and eddy currents (Andersson & Sotiropoulos, 2016), intra-frame motion (Andersson et al., 2017), and signal dropout (Andersson et al., 2016), as well as susceptibility distortions using the field generated by *topup*. The anisotropic map (Dell'Acqua et al., 2014) was generated from the corrected data and used to calculate a rigid-body registration of the diffusion data to the subject's T1w structural image in FLIRT (Jenkinson et al., 2002), and this transformation was applied to the entire diffusion series with a spline interpolation to align it with the T1w image; the bvecs were rotated with the same affine matrix.

Tractography and structural connectivity

Diffusion modelling was performed using the damped Richardson-Lucy spherical deconvolution algorithm (Dell'Acqua et al., 2010) in *StarTrack* (https://mr-startrack.com), with the following parameters: fibre response α = 1.5; number of iterations = 300; amplitude threshold η = 0.0015; geometric regularisation v = 16. Fibre tracking was performed with the Euler-type algorithm according to the following parameters: minimum HMOA threshold = 0.002; maximum angle threshold = 45°; step size = 1 mm; minimum fibre length = 20 mm; maximum fibre length = 300 mm.

Each participant's resulting tractogram was then converted to a connectivity matrix, as previously done by our group (Beyh et al., 2022). In brief, each streamline was projected to its nearest neighbouring vertex along the grey-white matter boundary surface (obtained from *FreeSurfer*) with a maximum distance of 3 mm. This resulted in a vertex-wise matrix, which was collapsed into a smaller matrix based on the Desikan-Killiany cortical parcellation (Desikan et al., 2006). Several parcel-wise connectivity matrices were produced with the following edge weights: streamline count; log-transformed streamline count; hindrance modulated orientational anisotropy (HMOA; Dell'Acqua et al., 2013), a tract-specific measure of white matter microstructure; and median fibre length.

Structure-Function Concordance

Structure-function concordance was calculated on a per-participant basis (n = 50), using the spearman correlation coefficient.

Predicting age

We assessed the ability of mean strength derived from each functional connectivity method to predict current age over and above head circumference. First, we calculated the R^2 from 10-fold cross-validation using head circumference as the only predictor of current age (model 1). Then, we added mean strength this model (model 2). We used the change in R^2 (model 2 – model 1) as the quantity of interest. 147 participants had head circumference data. We excluded two outlier values (90cm from a 22-year-old and 66cm from a 13-year-old) as these likely reflected measurement inaccuracies.

We assessed if model 2 – model 1 cross-validation R^2 between methodologically corresponding zero-phase inclusive and exclusive methods (e.g. PLI versus wPLI) were significantly different using permutation testing. We created a null distribution by permuting age 1000 times for each method-pair in each frequency band.

Prognostic ability for longitudinal changes in spatial working memory

Spatial working memory was measured using the 'Find the Phone' Task (Owen et al., 1990; Sjöwall et al., 2013). In this task, multiple telephones were shown on a computer screen and 'rang' one at a time. Through trial and error, participants had to guess which phone was ringing. Participants were instructed that each phone would only ring once within a trial. The task was to remember which telephone(s) had already rung to avoid re-selecting that phone. Instances where a participant selected a phone that had

already rung, or had previously been incorrectly selected, were counted as errors. The
metric of interest was 'total participant errors.'

For each functional connectivity method, we used the weighted, normalised pathlength
in the theta band as the candidate prognostic biomarker. The theta band has been
robustly associated with working memory (Bauer et al., 2021a; Dai et al., 2017a; Fell &
Axmacher, 2011; Hyman et al., 2010; Jones & Wilson, 2005; Siapas et al., 2005). We only
looked at connectivity between region-pairs with a strong evidence-base of being
implicated in spatial working memory, including regions in the prefrontal cortex (MacKey
& Curtis, 2017; Siapas et al., 2005; Wager & Smith, 2003; Wirt & Hyman, 2017), the
posterior parietal cortex (Almeida et al., 2015; MacKey & Curtis, 2017), the anterior
cingulate cortex (Wirt & Hyman, 2017), and the entorhinal cortex (Almeida et al., 2015;
Salimi et al., 2021), as defined by the Desikan-Killiany atlas (Desikan et al., 2006). In
total, there were 14 regions of interest in each hemisphere. Shorter weighted,
normalised pathlengths reflect higher network efficiency. Global network efficiency and
pathlength have been associated with working memory performance (Langer et al.,
2013; Stanley et al., 2015).

To calculate weighted, normalised pathlength, we first created weighted, undirected
functional connectivity adjacency matrices (networks) with 28 nodes. One matrix was
created per functional connectivity method. Each node corresponded to a cortical
region of the Desikan-Killiany atlas implicated in spatial working memory, while the
weighted edges were the functional connectivity strengths between pairs of nodes.
Next, using the MATLAB brain connectivity toolbox (Rubinov et al., 2009), we inverted
network edges to lengths (edges with larger weights produce shorter lengths). Then, we
computed the *shortest possible* distance between all pairs of nodes. Finally, we
calculated the average shortest distance between all pair of nodes to produce the
weighted normalised pathlength. Formally, the shortest weighted path length d_{ij}
between any two nodes, e.g. i and j is defined as:

$$d_{i,j}^{w} = \sum_{w_{ij} \in g_{i \leftrightarrow j}} w_{ij},$$

where $gi \leftrightarrow j$ is the shortest weighted path between nodes i and j.

The weighted, average shortest pathlength (L) between node i and all other nodes of the
network is defined as:

$$L_i = \frac{\sum_{i \neq j} d_{i,j}^{w}}{(N-1)} ,$$

where N is the number of nodes (28 in this analysis) and $d_{i,j}$ is the shortest path length
between nodes i and j, when j is any node that is not i.

To calculate the normalised weighted pathlength, we used a normalisation process by
creating 500 random networks by randomising the empirical network, while preserving
the degree and strength distributions: each edge was rewired 5 times, with edge weights
randomised every 5th step. The weighted pathlength of each random network was

calculated, and then averaged across each of the 500 random networks to produce
L_{rand}. The normalised average shortest pathlength of node i was calculated as:
516

517
$$L_i' = \frac{L_i}{L_{rand}}$$

L_i' at each node was averaged across 28 nodes to produce the mean normalised,
weighted pathlength (referred to as normalised, weighted pathlength).

Using general linear models, we compared the ability of normalised, weighted
pathlength produced from each functional connectivity method to predict changes in
total participant errors between the time the EEG was recorded and a time 12-30
months in the future (mean = 19 months, SD = 3.5 months).

Penalisation Scale
We created a penalisation scale to quantify how connectivity values would be altered by
zero-phase-exclusive methods. The scale was between 0 and 1, where the most
penalizable phase differences (0 and ±π) were given a value of 1 and least penalizable
phase differences (±π/2) were given a value of 0. Penalisation scale values were
calculated as follows:

534
$$Penalisation\ value = \frac{||\text{average phase delay}| - \frac{\pi}{2}|}{\frac{\pi}{2}},$$

where the average absolute phase delay was limited between 0 and +π. To examine if
stronger structural connections were more likely to be associated with more
penalizable phase-delays, we spearman correlated the penalisation values with the
structural connectivity matrix on a per-participant basis. This was only done for region-
pairs between which there were 1 or more streamlines.

Proxy of signal transmission time
We used a proxy of signal transmission time through white matter tracts. We divided the
median length (distance) of white matter tracts between region-pairs by a proxy of
transmission speed. The median hindrance modulated orientational anisotropy (HMOA)
was used a proxy of transmission speed, as white matter microstructural integrity can
reflect myelination (Dell'Acqua et al., 2013) and correlates with nerve conduction
velocity (Heckel et al., 2015; Wang et al., 2022). Additionally, the transmission latencies
of evoked potentials (an established method of approximating transmission speed; van
Blooijs et al., 2023) negatively correlate with white matter microstructure across studies
(Evstigneev et al., 2013; Whitford et al., 2011).

Bootstrapping for mediation effects
We performed a mediation analysis to investigate if the association between structural
connectivity strength and the penalisability of the phase-difference of functional
connectivity was mediated by a proxy for white matter signal transmission time. We
tested the significance of the indirect effect using bootstrapping procedures.

Unstandardized indirect effects were computed for each of 1000 bootstrapped
samples, and the 95% confidence interval was computed by determining the indirect
effects at the 2.5th and 97.5th percentiles.

Results
Most functional connectivity occurs with zero- and near-zero phase-delay, even
when artefactual connectivity is likely negligible
We quantified the distribution of the average phase-delay of functional connectivity
between brain region-pairs (Fig. 1b). Between all region-pairs across all participants,
most measured functional connectivity occurred with mean phase-delays of either
zero-/near-zero or π-/near-π radians. Across frequency bands, alpha had the lowest
proportion of zero- and near-zero phase delay connectivity (52% of all connections),
while high-beta had the highest proportion (69% of all connections), Fig. 1b.

Fig. 1: Quantifying zero- and near-zero phase-delay functional connectivity

a) Proportion of zero- & near-zero phase-delay synchrony is static beyond ≥ 60mm Euclidean distance between region-pairs

b) Connectivity between:

a) Where the percentage of zero- and near-zero phase delay connectivity does not vary as a function of Euclidean distance, artefactual connectivity due to signal leakage is likely to be negligible. This was found to be at Euclidean distances of ≥ 60mm for all frequency bands. The number of pairs of regions per person at each minimum distance is listed above the graph. b) Histograms of the average phase-delay of functional connectivity during an epoch between all region-pairs, region-pairs > 80mm Euclidean distance apart (where signal leakage is likely negligible) and between homotopic interhemispheric region-pairs. n = 153. 8-13 Hz and 20-32 Hz are the frequency bands with the lowest and highest near-zero phase delay connectivity, respectively. *Italics*: the proportion of zero- (and ±π) and near-zero (and near ±π) connectivity. FC = functional connectivity.

In source reconstructed data, signal leakage (and not volume conduction) is thought to
be the primary contributor of artefactual zero-phase-delay connectivity (He et al.,
2019). To quantify the proportion of zero- and near-zero phase-delay functional
connectivity when artefactual connectivity due to signal leakage was likely negligible,
we used a novel approach. We leveraged the fact that artefactual connectivity due to
signal leakage occurs at zero-phase-delay (Gohel et al., 2017; Hipp et al., 2012) and
decreases with increasing Euclidean distance between sources (Bastos & Schoffelen,
2016; Gohel et al., 2017; He et al., 2019; Hipp et al., 2012). Therefore, at sufficiently
large distances where the percentage of zero- and near-zero phase-delay connectivity
does not vary as a function of Euclidean distance, artefactual connectivity due to signal
leakage is likely to be negligible. We examined this relationship graphically (Fig. 1a),
finding that the proportion of zero- and near-zero phase-delay connectivity did not vary
as a function of Euclidean distance > 60mm. As expected (Srinivasan et al., 2007), this
was true for all frequency bands. This finding is consistent with artefactual connectivity
due to signal leakage being likely negligible beyond 60mm, in keeping with previous
estimates using simulated data (Anzolin et al., 2019; Gohel et al., 2017).
586
587   When limiting our analyses to region-pairs 80mm (where zero- / near-zero phase-delay
588   connectivity was minimal) or more apart, we found that most functional connectivity
589   still occurred with zero- and near-zero phase-delay, in all frequency bands except alpha
590   (where 47% of all functional connections occurred with zero- or near-zero phase-delay,
591   Fig. 1b).
592
Finally, we looked at homotopic interhemispheric functional connectivity as an
example of connectivity known to robustly occur across species and measurement
modalities (Cauda et al., 2021; Conturo et al., 1999; Engel et al., 1991; O'Reilly &
Elsabbagh, 2021). Remarkably, 83-89% of homotopic interhemispheric connectivity
occurred with zero- or near-zero phase-delays (Fig. 1b), despite the significant
Euclidean distance between them. This proportion was greater than that of 34 randomly
selected pairs of heterotopic interhemispheric region-pairs, in each frequency band
(e.g. alpha: X^2 (1, n = 5,202) = 8.19, *p* = .0042, Supplementary Table S1).
601
Comparing the performance of zero-phase inclusive versus exclusive functional
connectivity metrics
Test-retest reliability is higher in zero-phase inclusive than exclusive connectivity
methods
In the alpha band, median edgewise ICC values were 0.20 – 0.71 across zero-phase-
inclusive methods and 0.07 – 0.17 across zero-phase-exclusive methods (n = 99, Fig. 2,

Supplementary Table S2). Absolute agreement was greatest in high-beta for all zero-
phase-inclusive methods and Img COH. It was highest in alpha for the wPLI and in low-
beta for Orth AEC. The same patterns were also found for the consistency of each
measure (Supplementary Table S3).
612
We found high absolute agreement for measured connectivity between region-pairs
with a small Euclidean distance between them using zero-phase inclusive but not
exclusive methods (Fig. 2). There are two plausible causes for this: 1) neighbouring
region-pairs reliably (structurally and functionally) connect to each other (Bullmore &
Sporns, 2012; Conturo et al., 1999; Finger et al., 2016), and this functional connectivity
occurs with zero- and near-zero phase-delays and 2) artefactual connectivity (Hipp et
al., 2012).
620
621   Absolute agreement at homotopic interhemispheric region-pairs (Fig. 2) was high in all
622   zero-phase inclusive methods, moderate with Img COH, and low with the Orth AEC and
623   wPLI.
624
625   The relative test-retest reliability quantities across methods (Colclough et al., 2016;
626   Fraschini et al., 2019) and frequency bands (van der Velde et al., 2019) is consistent
627   with previous work, despite methodological differences.

Fig. 2: edgewise absolute agreement across functional connectivity methods and frequency bands

Edgewise absolute agreement, quantified by intraclass correlation coefficient (ICC), for each functional connectivity method. In 8 - 13 Hz, for example, median ICC values were 0.20 – 0.71 across zero-phase-inclusive methods versus 0.07 – 0.17 across zero-phase-exclusive methods. High absolute agreement was seen between homotopic interhemispheric region-pairs (off-diagonal edges in the adjacency matrix) in zero-phase-inclusive but not exclusive methods. High absolute agreement was seen between adjacent regions (main diagonal) in zero-phase inclusive methods, but not in zero-phase exclusive methods. Brain regions in the matrices were ordered by lobe and their proximity to each other. The colour bar denotes the edgewise ICC value. Negative ICC values were limited at 0.

Convergence with underlying neurobiology

We assessed the extent to which metrics derived from each functional connectivity method converged with underlying neurobiology by quantifying: 1) structure-function concordance, as functional connectivity is influenced by monosynaptic structural cortico-cortical pathways (Baum et al., 2020; Chu et al., 2015; Finger et al., 2016; Fotiadis et al., 2024; Honey et al., 2009), 2) the ability of functional connectivity metrics to capture homotopic interhemispheric connectivity, 3) their ability to predict participant age.

Concordance with structural connectivity is greater in zero-phase-inclusive methods

For all zero-phase-delay inclusive methods, mean structure-function concordance was significantly greater than zero in every frequency band (n = 50, one-sample t-test or Wilcoxon signed-rank test, $\mu = 0$, $p <.0001$, Bonferroni corrected), Fig. 3. Concordance was not significantly different to zero for wPLI and orthogonalized AEC (one-sample t-test, $p > .05$). Surprisingly, concordance was significantly less than zero for Img COH (one-sample t-test or Wilcoxon signed-rank test, $p <.0001$, Bonferroni corrected), Fig. 3a-b. Concordance for each zero-phase-delay inclusive method was significantly higher than that for its methodologically corresponding zero-phase-delay exclusive method (paired-samples t-test or Wilcoxon signed-rank test, $p < .0001$, Bonferroni corrected, Fig. 3a-b). All findings were replicated when analysing each of the two sites separately, despite a difference in mean age between sites and large site effects in MRI analyses (Jovicich et al., 2006), Supplementary Table S4a-b.

Fig. 3: Structure-function concordance is higher in zero-phase inclusive versus exclusive methods

a) Structure-function concordance in the alpha band and b) in all frequency bands (n = 50). In all frequency bands, concordance was significantly greater than zero with zero-phase-inclusive methods (one-sided t-test or Wilcoxon sum-ranked test, *μ* = 0, *p* < .0001, Bonferroni corrected). Concordance was not significantly different from zero for two zero-phase-exclusive methods (wPLI and orthogonalized AEC). Surprisingly, concordance was significantly less than zero for the imaginary part of coherency. Concordance for each zero-phase-inclusive method was significantly higher than that of its methodologically corresponding zero-phase-exclusive method in each frequency band (paired samples t-test or Wilcoxon signed-rank test, *p* < .0001, Bonferroni corrected). *** concordance significantly different from zero, *p* < .0001, Bonferroni corrected. ᵂ = data not normally distributed, therefore one-sided Wilcoxon sign-rank test performed, otherwise one-sided t-tests performed. SFC = structure-function concordance.

b)

0-phase inclusive method	Freq (Hz)	Mean SFC	SD	0-phase exclusive method	Freq (Hz)	Mean SFC	SD
Coherence	1 - 4ᵂ	.098***	.043	Imaginary part of Coherency	1 - 4	-.038***	.039
	4 - 8	.108***	.040		4 - 8	-.061***	.048
	8 - 13	.105***	.039		8 - 13	-.048***	.049
	13 - 20	.106***	.045		13 - 20	-.055***	.051
	20 - 32	.113***	.047		20 - 32	-.072***	.048
PLV	1 - 4ᵂ	.129***	.038	wPLI	1 - 4	-.001	.041
	4 - 8	.128***	.037		4 - 8	-.010	.040
	8 - 13	.119***	.033		8 - 13	-.010	.039
	13 - 20	.130***	.033		13 - 20	.002	.033
	20 - 32	.132***	.037		20 - 32	.001	.044
AEC	1 - 4	.123***	.042	Orthogonalized AEC	1 - 4	.006	.035
	4 - 8	.123***	.037		4 - 8	.010	.036
	8 - 13ᵂ	.111***	.043		8 - 13	.006	.045
	13 - 20	.126***	.036		13 - 20	.008	.036
	20 - 32	.129***	.037		20 - 32	.005	.034

Homotopic interhemispheric functional connectivity is robustly present with zero-phase-inclusive but not zero-phase-exclusive methods

Homotopic interhemispheric connectivity is robustly found across species and modalities (Engel et al., 1991; Mancuso et al., 2019; O'Reilly & Elsabbagh, 2021; Schmahmann & Pandya, 2009; Vicente et al., 2008; Witham et al., 2007). We examined each functional and structural connectivity method's ability to capture this fundamental brain property. We calculated the edgewise mean functional (n = 153) and structural (n = 50) connectivity strength across all participants. On visual inspection, strong homotopic interhemispheric connectivity was found with all zero-phase-inclusive functional connectivity methods (Fig. 4a) and with structural connectivity (reflecting callosal connections, Fig. 4b). No zero-phase-exclusive methods showed prominent homotopic interhemispheric connectivity. Img COH showed significantly weaker connectivity between homotopic interhemispheric region-pairs than between non-homotopic region-pairs (Fig 4a, Supplementary Table S5), which we explored in a *post hoc* analysis below. Of note, given the relatively large Euclidean distances between them (median 57mm, IQR 21-80mm), functional connectivity between homotopic region-pairs would be minimally impacted by signal leakage.

Fig. 4: Edgewise structural and functional connectivity strengths

a) Edgewise functional connectivity averaged across participants

b) Edgewise structural connectivity averaged across participants

Mean edgewise **a)** EEG functional connectivity matrices (n = 153) and **b)** structural connectivity matrix (n = 50). Strong homotopic interhemispheric connectivity (seen at the off-diagonal edges in each adjacency matrix) was found with structural connectivity and with all zero-phase-inclusive functional connectivity methods. No zero-phase-exclusive methods showed prominent callosal connectivity. The imaginary part of coherency showed significantly weaker connectivity between homotopic interhemispheric region-pairs than between other region-pairs. Zero-phase-delay exclusive methods did not show strong connectivity between neighbouring region-pairs (main diagonal). Colour bars reflect connectivity strength, kept constant within each connectivity method; yellower colours denote higher connectivity strength. Brain regions in matrices are ordered by lobe and their proximity to each other.

The ability to predict current age is better in zero-phase-inclusive versus exclusive
methods
Given that brain functional connectivity matures with age (Collin & Van Den Heuvel,
2013; Edde et al., 2021; Fair et al., 2009), we quantified the extent to which age-related
changes are captured by each functional connectivity method. We assessed the ability
of mean strength derived from each method to predict current age over and above head
circumference (which correlates with age). First, we calculated the R^2 from 10-fold
cross-validation using head circumference as the only predictor of current age (model
1, cross-validation R^2 = .27). Then, we added mean strength to this model (model 2). The
subsequent change in R^2 (model 2 – model 1) is reported in Table 3a, n = 145. Across all
frequency bands, model 2 – model 1 R^2 was greater in a zero-phase-inclusive method
compared to its methodologically corresponding zero-phase-exclusive method;
permutation analysis showed that this was statistically significant (false discovery rate
corrected) in 10/15 comparisons.
685

Table 3a: Prediction of current age is greater in zero-phase inclusive versus exclusive methods. Values show model 2 – model 1 ten-fold cross validation R^2 (n = 145) for each functional connectivity method and frequency band. Model 1: *current age ~ head circumference* (model 1 cross-validation R^2 = .27). Model 2: *current age ~ head circumference + mean strength*.

Freq (Hz)	0-phase-inclusive method	Model 2 – 1 cross validation R^2	0-phase-exclusive method	Model 2 – 1 cross validation R^2	p_{FDR} permute
1 - 4	Coherence	.04	Imaginary part of Coherency	.02	.746
4 - 8		.09		.02	**.041**
8 - 13		.06		.05	.905
13 - 20		.07		-.02	**.020**
20 - 32		.01		-.01	.070
1 - 4	PLV	.05	wPLI	-.01	**.010**
4 - 8		.12		.02	**.015**
8 - 13		.16		.11	.101
13 - 20		.11		.01	**.041**
20 - 32		.10		.04	**.015**
1 - 4	AEC	.04	Orthogonalized AEC	.00	.285
4 - 8		.10		-.01	**.041**
8 - 13		.15		.02	**.000**
13 - 20		.14		.02	**.041**
20 - 32		.15		-.01	**.000**

Table 3b: Prediction of longitudinal changes in spatial working memory performance is greater in zero-phase inclusive versus exclusive methods. In models predicting longitudinal changes in spatial working memory performance, a significant main effect of functional connectivity derived pathlength was only found for coherence in the 6–12-year group and PLV in the 18-31-year group. For these models, and their corresponding zero-phase-exclusive methods, we report model 2 – model 1 leave-one-out cross-validation R^2. Model 1: *longitudinal change in spatial working memory ~ covariates*. Model 2: *longitudinal change in spatial working memory ~ covariates + functional connectivity derived pathlength*. *significant main effect of covariate in general linear model (*p* < .05). SWM = spatial working memory, T1 = timepoint 1, T2 = timepoint 2, PL = pathlength

Age Group	Model 1 covariates	Model 1 R^2	Covariate added in model 2:	Model 2 R^2	Model 2 – Model 1 R^2
6-12 yrs (n = 32)	Age + T1 SWM	.006	Coh PL*	.107	.101
			Img COH PL	.001	-.005
18-31 yrs (n=36)	Age + T1 SWM + T2-T1 time interval	.489	PLV PL*	.563	.074
			wPLI PL	.453	-.036

Prognostic ability for longitudinal changes in spatial working memory is higher in
zero-phase inclusive versus exclusive methods
We explored the ability of metrics derived from each functional connectivity method to
act as prognostic biomarkers (Pletcher & Pignone, 2011) for longitudinal changes in
spatial working memory ability, above and beyond current working memory abilities. We

chose spatial working memory as an outcome measure because it has highly specific
associations with functional connectivity (Dai et al., 2017b; Fox et al., 2005; Hampson
et al., 2006; Kang et al., 2011) and altered working memory is implicated in several
neuropsychiatric conditions (Alloway et al., 2009; Chai et al., 2018; Gathercole &
Alloway, 2006).
696
We found significant pathlength x age interactions in general linear models predicting
changes in spatial working memory performance, when analysing all 6-31-year-old
participants together (examples for coherence and Img COH provided in
Supplementary Tables S6-7). Therefore, children (6-12 years, n = 32), adolescents (13-17
years, n = 41) and adults (18-31 years, n = 36) were analysed separately. Model
covariates (selected using Akaike information criterion) were kept constant within each
age group.
704
A main effect of network pathlength was only found for methods sensitive to zero-
phase-delay connectivity: coherence in children (β = 3.8, SE = 1.5, p = .014; full model
$F_{3/28}$ = 4.01, p = .017) and PLV in adults (β = -2.6, SE = 1.1, p = .027; full model $F_{3/32}$ =
18.53, p = .003), Supplementary Tables S8-9. For these statistically significant findings,
we used leave-one-out cross-validation (LOOCV) to quantify model performance (Table
3b). First, we calculated the LOOCV R^2 using covariates (model 1). Then, we added
weighted, normalised pathlength to this model (model 2). Model 2 – model 1 LOOCV R^2
increased by .101 for coherence in children and .074 for PLV in adults. For comparison
in their methodologically corresponding zero-phase-exclusive methods, model 2 –
model 1 LOOCV R^2 *decreased* by .005 for Img COH and .036 for wPLI.
715

Post hoc analyses
An unexpected finding from our analyses was negative structure-function concordance
with Img COH. We theorised that this was because the strongest structural connections
had the most 'penalised' functional connections when using Img COH, based on the
following logic: stronger structural connections result in quicker (decreased) signal
transmission time between region-pairs (Westerhausen et al., 2006; Whitford et al.,
2011). Quicker transmission time means that oscillatory activity at each connected
region-pair occurs with smaller time-delays, and therefore, smaller phase-delays. Zero-
phase-exclusive methods may penalize, rather than merely exclude, functional
connectivity at near-zero-phase-delays (a salient methodological detail is that zero-
phase-exclusive methods modify the connectivity strength value based on the
magnitude of phase delay, while zero-phase-inclusive methods to do not). Hence,
stronger structural connections may lead to weaker functional connections with some
zero-phase-exclusive methods.
730
To test this, we hypothesised that: 1) the connectivity strength values of zero- and near-
zero phase-delay connections are penalised, and not merely excluded, if derived from
zero-phase-exclusive methods. 2) Stronger structural connections lead to functional
connections with more penalizable phase delays, mediated by quicker possible white
matter transmission time. We expected a partial mediation, as strong structural
connections such as the corpus callosum also produce zero-phase-delay functional
connectivity through resonance-induced synchrony. 3) Structural connections that

allow the quickest connectivity between region-pairs result in the weakest functional
connectivity if derived from Img COH.
740
Near-zero-phase-delay functional connectivity may be penalised and not merely
excluded by zero-phase-delay-exclusive methods
We inspected the relationship between absolute phase delay and functional
connectivity strength (n = 153), between region-pairs where signal leakage is likely
negligible (region-pairs ≥80mm apart), Fig. 5. With zero-phase-inclusive methods, the
median connectivity strength was significantly higher for near-zero/near-π phase-delay
connectivity compared to connectivity at other phase delays, in all frequency bands
(Fig. 5b). For Img COH and wPLI the opposite was true: median connectivity strength
was significantly lower for near-zero/near-π phase-delay connectivity compared to
connectivity at other phase delays. For orthogonalized AEC, median connectivity
strength was similar between near-zero phase-delay and other phase-delay
connectivity, in keeping with the milder form of orthogonalization we used compared to
others (Brookes et al., 2012; Hipp et al., 2012). Therefore, zero- and near-zero phase-
delay functional connectivity may be penalised, and not merely excluded, by zero-
phase-exclusive methods.

Fig. 5: When signal leakage is likely negligible, most functional connectivity occurs with near-zero-phase-delay, penalised by zero-phase-exclusive methods

b) Comparison of the median strength of zero / near-zero versus non-near-zero phase-delay functional connectivity

0-phase-inclusive-method	Freq (Hz)	Median FC strength		0-phase-exclusive-method	Freq (Hz)	Median FC strength	
		Near-zero phase delay	Other phase delay			Near-zero phase delay	Other phase delay
Coherence	1 - 4*	.52	.39	Imaginary part of Coherency	1 - 4*	.11	.13
	4 - 8*	.52	.39		4 - 8*	.12	.15
	8 - 13*	.51	.40		8 - 13*	.14	.16
	13 - 20*	.48	.38		13 - 20*	.15	.17
	20 - 32*	.46	.37		20 - 32*	.17	.19
PLV	1 - 4*	.36	.17	wPLI	1 - 4*	.06	.16
	4 - 8*	.34	.15		4 - 8*	.06	.14
	8 - 13*	.32	.16		8 - 13*	.06	.16
	13 - 20*	.28	.11		13 - 20*	.04	.10
	20 - 32*	.26	.09		20 - 32*	.04	.08
AEC	1 - 4*	.21	.10	Orth AEC	1 - 4*	.09	.08
	4 - 8*	.22	.10		4 - 8	.08	.08
	8 - 13*	.20	.10		8 - 13	.08	.07
	13 - 20*	.17	.07		13 - 20*	.06	.06
	20 - 32*	.12	.05		20 - 32*	.04	.04

a) The relationship between absolute phase-delay and functional connectivity strength (yellower colours denote higher density of scatter points), and **b)** a comparison of edge strength when functional connectivity has zero / near-zero versus non-near-zero phase delays, for region-pairs > 80mm Euclidean distance apart (where artefactual connectivity due to signal leakage is likely negligible). For zero-phase-inclusive methods, across all frequency bands, median strength was significantly higher for near-zero-phase-delay connections compared to connections with non-near-zero-phase-delays. For the imaginary part of coherency and wPLI, across all frequency bands, the median strength was significantly lower for near-zero phase-delay connectivity compared to non-near-zero-phase-delay connectivity. For orthogonalized AEC, median strength was similar between near-zero and non-near-zero phase delay connections. Zero-phase-exclusive methods modify the connectivity strength value based on the magnitude of phase delay, while zero-phase-inclusive methods to do not. Therefore, zero-phase-exclusive methods may penalise, rather than merely exclude, zero- and near-zero phase-delay connections. * p < .0001, Bonferroni corrected. FC = functional connectivity.

Stronger structural connectivity leads to functional connectivity with more penalizable phase-delays, mediated by shorter white matter transmission times

We hypothesised that the stronger a structural connection, the more likely it was to produce functional connectivity with phase-delays closer to zero, penalised by zero-phase-exclusive methods. The mean correlation between streamline count and penalisability of phase-delays was significantly greater than 0 for all frequency bands (n = 50, one-sample t-test, $\mu = 0$, p < .0001, Bonferroni corrected), Fig. 6a. Hence, stronger structural connectivity led to functional connectivity with phase-delays more penalizable by zero-phase-exclusive methods.

To test whether this relationship was mediated by quicker signal transmission time through white matter connections, we used a proxy of signal transmission time through

white matter tracts. Across all frequency bands, we found that the effect of structural
connectivity strength on the penalisability of the phase-delay of functional connectivity
was significantly partially mediated by the proxy of white matter transmission time (e.g.
proportion of the direct effect mediated by the proxy of signal transmission time was
.234 [95% confidence intervals, calculated using 1000 bootstrapped samples = .197 -
.270] in alpha, Fig. 6b-c). The mediation was strongest in higher frequency bands,
consistent with stronger structure-function concordance in higher frequency bands in
zero-phase-inclusive methods (Fig. 3b).
776

Fig. 3: Stronger structural connectivity leads to functional connectivity with phase-delays closer to 0 or ±π, mediated by a proxy of shorter white matter transmission time

a)   Correlation between structural connectivity strength and penalisation values

Freq (Hz)	Spearman's Correlation	SD
1 - 4	.092***	.039
4 - 8	.098***	.036
8 - 13	.096***	.036
13 - 20	.092***	.031
20 - 32	.095***	.042

b)

c) Mediation analysis statistics for the model in b), for each frequency band

Freq (Hz)	Total effect (x10⁻³)	Average direct effect (x10⁻³)	a	b (x10⁻⁶)	Average causal mediation effect (x10⁻³)	Proportion mediated [95% confidence interval]
1 – 4	2.97***	2.32***	-159***	-4.04***	0.64	.217 [.175 - .260]
4 – 8	2.61***	2.08***	-159***	-3.33***	0.53	.203 [.160 - 0.260]
8 – 13	3.86***	2.96***	-159***	-5.67***	0.90	.234 [.200 - .280]
13 – 20	2.64***	1.92***	-159***	-4.49***	0.71	.271 [.225 - .330]
20 – 32	2.17***	1.57***	-159***	-3.78***	0.60	.277 [.220 - .350]

a) We created a penalisation scale between 0 and 1, where phase-delays most penalised by zero-phase-exclusive methods (0 and ±π) have a value of 1 and phase-delays least penalised (±π/2) have a value of 0. The mean correlation across participants between streamline count and penalisability of phase-delays was significantly greater than 0 for all frequency bands, n = 50 (*** p <.0001, Bonferroni corrected, one-sample t-test, $\mu = 0$). **b-c)** We created a proxy of signal transmission time through white matter tracts. The effect of structural connectivity strength on the penalisabiltiy of the phase-delay of functional connectivity was partially mediated by the proxy of white matter transmission time, across all frequency bands. All effects were statistically significant (*** = p < .0001, Bonferroni corrected). SD = standard deviation.

Structural connections with shorter signal transmission time are associated with
weaker functional connectivity with some zero-phase-exclusive methods
We found that the shorter the proxy of signal transmission time through white matter
tracts, the stronger the functional connectivity value was for all zero-phase-inclusive
methods (one-sample t-test, $\mu = 0$, $p < .0001$, Bonferroni corrected), Fig. 7. This is the
expected biological relationship (Bassett & Bullmore, 2017; Bullmore & Sporns, 2012;
Cabral et al., 2014). This was also the case for the alpha band in wPLI ($p < .05$,
Bonferroni corrected) and orthogonalized AEC ($p < .001$, Bonferroni corrected), but no
other frequency bands when using these methods. Note that alpha has the smallest
proportion of near-zero-phase-delay connectivity (Fig. 1b). For Img COH, the shorter the
proxy of signal transmission time through white matter tracts, the weaker functional
connectivity strength, in all frequency bands (one-sample t-test, $\mu = 0$, $p < .0001$,
Bonferroni corrected). This is inconsistent with prior network literature (Bassett &
Bullmore, 2017; Bullmore & Sporns, 2012; Cabral et al., 2014).
791
792 Thus, the structural connections that are prioritised in brain through quick structural
793 pathways may be penalised by some zero-phase-delay-exclusive functional
794 connectivity methods.
795

b) Spearman's correlation between the proxy of white matter signal transmission time and functional connectivity strength

0-phase-inclusive-method	Freq (Hz)	Mean correlation	SD	0-phase-exclusive-method	Freq (Hz)	Mean correlation	SD
Coherence	1 - 4	-.132***	.060	Imaginary part of Coherency	1 - 4	.051***	.062
	4 - 8	-.120***	.058		4 - 8	.078***	.062
	8 - 13	-.170***	.063		8 - 13	.083***	.060
	13 - 20	-.147***	.072		13 - 20	.071***	.070
	20 - 32	-.148***	.052		20 - 32	.087***	.067
PLV	1 - 4	-.156***	.060	wPLI	1 - 4	-.007	.053
	4 - 8	-.160***	.060		4 - 8	.004	.060
	8 - 13	-.206***	.058		8 - 13	-.033*	.062
	13 - 20	-.180***	.064		13 - 20	.000	.054
	20 - 32	-.175***	.063		20 - 32	-.005	.071
AEC	1 - 4	-.155***	.056	Orthogonalized AEC	1 - 4	-.012	.056
	4 - 8	-.163***	.063		4 - 8	-.022	.063
	8 - 13	-.192***	.071		8 - 13	-.045**	.065
	13 - 20	-.173***	.067		13 - 20	-.015	.059
	20 - 32	-.172***	.067		20 - 32	-.007	.057

We created a proxy of signal transmission time through white matter tracts. The quicker (shorter) the proxy of signal transmission time through structural connections, the stronger the functional connectivity value for all zero-phase-inclusive methods (one-sample t test, $\mu = 0$, $p < .0001$, Bonferroni corrected), **a)** shown graphically for alpha and **b)** tabulated for all frequency bands (n = 50). This was also the case for alpha (but not other frequency bands) in wPLI ($p < .05$, Bonferroni corrected) and orthogonalized AEC ($p < .001$, Bonferroni corrected). For the Img COH, the quicker the proxy of structural transmission time, the weaker functional connectivity strength, in all frequency bands (one-sample t-test, $\mu = 0$, $p < .0001$, Bonferroni corrected). Such a relationship is inconsistent with prior literature. ***$p < .0001$, **$p < .001$, *$p < .05$ (Bonferroni corrected).

Discussion

Discussion

We demonstrated that most interactions between cortical region-pairs occur with zero-
and near-zero phase-delay. We used a novel approach to show that this is also the case
when artefactual connectivity due to signal leakage is likely negligible. Including zero-
and near-zero phase-delay connectivity increased test-retest reliability, convergence
with neurobiology (concordance with structural connectivity, ability to capture
homotopic interhemispheric connectivity and ability to predict chronological age) and
prognostic ability for longitudinal changes in cognitive ability. Thereby, through
converging lines of evidence, we show that functional connectivity metrics that include
zero- and near-zero phase-delay connections have more properties desired in
candidate biomarkers. We propose that the decreased structure-function
concordance, ability to capture age, and ability to predict longitudinal changes in
cognition associated with zero-phase-exclusive methods are due to their penalisation
of most 'true' cortico-cortical functional connectivity.

We investigated homotopic interhemispheric connectivity as an example of structural
and functional connectivity robustly found across species (Engel et al., 1991; Mancuso
et al., 2019; O'Reilly & Elsabbagh, 2021; Schmahmann & Pandya, 2009; Vicente et al.,
2008; Witham et al., 2007). We captured conspicuous homotopic connections using
structural connectivity and zero-phase-inclusive functional connectivity methods. We
found that they were either inconspicuous or penalised when using zero-phase-
exclusive methods. In contrast, a previous study (Hipp et al., 2012) captured
connectivity between sensory homotopic region-pairs using orthogonalized AEC.
However, given our and others' (Engel et al., 1991; O'Reilly & Elsabbagh, 2021) finding
that homotopic region-pairs predominantly interact with zero- and near-zero-phase-
delay, our findings are expected. We see the inability to capture this fundamental
feature of brain connectivity as a key weakness of zero-phase-exclusive methods.

We found that structure-function concordance was significantly greater when using
zero-phase-inclusive compared to zero-phase-exclusive methods. This is consistent
with a previous small study of 17 participants (Finger et al., 2016). Additionally, we were
surprised that with some zero-phase-exclusive methods, here Img COH, the strongest
structural connections paradoxically had the weakest functional connections. We
showed that this was because stronger structural connectivity led to functional
connectivity with phase-delays closer to zero, mediated by a proxy of shorter signal
transmission times. Then, we showed that functional connectivity at near-zero-phase-
delays had penalised strength when derived from some zero-phase-exclusive methods.
Hence, for Img COH, we found that region-pairs connected by white matter tracts with
the quickest possible signal transmission times had the weakest functional
connections. This relationship is inconsistent with previous network literature (Bassett
& Bullmore, 2017; Bullmore & Sporns, 2012; Cabral et al., 2014).

Predictive validity for longitudinal changes in spatial working memory performance, as
well as the ability to predict current age, was substantially greater when zero- and near-
zero phase-delay connections were included. Strengths of our approach included the
fact that our spatial working memory prediction findings were derived using region-pairs

(Almeida et al., 2015; Jones & Wilson, 2005; MacKey & Curtis, 2017; Salimi et al., 2021;
Siapas et al., 2005; Wager & Smith, 2003; Wirt & Hyman, 2017), the frequency band and
the network topology measure known to be implicated in spatial working memory
(Bauer et al., 2021b; Dai et al., 2017a; Fell & Axmacher, 2011; Hyman et al., 2010; Jones
& Wilson, 2005; Siapas et al., 2005), specified *a priori*. Consistent with our findings, a
small, cross-sectional prior study (Jian et al., 2017) found that including versus
excluding zero-phase-delay connectivity substantially increased the variance explained
in predicting current task state. It showed that predictive ability was not dependent on
differing characteristics between functional connectivity methods besides the inclusion
or exclusion of zero-phase-delay connectivity: it excluded zero-phase-delay
connectivity using a large-Laplacian while keeping the connectivity method constant.
853
Conversely, zero-phase-exclusive methods did demonstrate some advantages. In
agreement with others (Hipp et al., 2012; Nolte et al., 2004; Vinck et al., 2011), it
appeared that zero-phase-exclusive methods likely minimised artefactual connectivity
due to volume conduction and signal leakage: functional connectivity between
neighbouring region-pairs was minimal when using them (Fig. 4). However, valid
functional connections are also most commonly present between nearby region-pairs
(Bassett & Bullmore, 2017; Honey et al., 2009; Mehrkanoon et al., 2014; Varela et al.,
2001), and were not reliably captured by zero-phase-exclusive methods (Figs. 2 and 4).
862
Quantifying 'true' zero- and near-zero phase-delay connectivity has not been previously
achieved in empirical EEG or MEG data. Our approach to identify the distance between
region-pairs where the effects of signal leakage were likely negligible was novel.
Previous approaches have used simulated data (Anzolin et al., 2019; Gohel et al., 2017),
with findings consistent with ours. However, simulated data cannot yet accurately
inform us about the proportion of 'true' cortico-cortical functional connectivity that
occurs with zero- and near-zero phase-delay, as even the most biologically realistic
simulations (Moran et al., 2013) cannot accurately model the emergent properties of
whole-cortex connectivity (Engel et al., 1991; Mill et al., 2017; Uhlhaas et al., 2009).
Therefore, unlike simulated data approaches, our approach can illustrate that most
cortical functional connectivity occurs at zero- or near-zero phase-delay, and that
excluding it has deleterious effects on the properties of biomarkers derived from
functional connectivity metrics. While the extent of artefactual zero-phase-delay
connectivity in empirical signals is dependent on methodological choices, our findings
highlight that zero- and near-zero-phase delay synchronisation comprises a substantial
proportion of 'true' neural interactions.
879
A key strength of our functional connectivity metric performance assessment approach
was the use of convergence across performance criteria – each based on different
modalities not trivially correlated with each other. Previous attempts at assessing the
validity of zero-phase-inclusive versus zero-phase-exclusive methods have used single
measures of functional connectivity performance (Brookes et al., 2012; Colclough et
al., 2016; Hipp et al., 2012), which may lead to bias (Fraschini et al., 2019; Noble et al.,
2021). Even using (only) two modalities could be misleading: for example, both
structural connectivity and functional connectivity from zero-phase-inclusive methods
are more reliable with decreasing Euclidean distance between region-pairs (Calamante,

2019; Jbabdi & Johansen-Berg, 2011). Thus, high structure-function concordance may
reflect reliable but artefactual functional connectivity correlating with reliable and valid
structural connectivity at small distances. Therefore, our use of non-neuroimaging
markers of performance was important.
893
An additional strength was our multi-site study design; our most unexpected finding
(negative structure-function concordance with Img COH) was reproduced when
analysing each study site separately. Next, as a measure to assess convergence
against, the structural connectome has high inter- and intra-individual reliability (Chu et
al., 2015; Dennis et al., 2012). Further, to test each functional connectivity metric
property, we used multiple analysis approaches, with consistent results.
900
Is high test-retest reliability a good measure of functional connectivity metric
performance? High reliability has been used as evidence of the presence of volume
conduction or signal leakage artefact in functional connectivity analyses (Colclough et
al., 2016; Nagy et al., 2024). Further, high validity may be present with low test-retest
reliability: brain-regions functionally connected at one moment may vary from the next.
However, in the context of a candidate prognostic or diagnostic biomarker, high test-
retest reliability is crucial; it establishes the upper limit of its predictive validity
(Carmines & Zeller, 1979).
909
While we aimed to identify where the effects of signal leakage were negligible, the use of
EEG additionally introduced volume conduction artefact (Wolters et al., 2004). In
sensor space, artefactual connectivity due to volume conduction has a non-linear
relationship with scalp distance (Srinivasan et al., 2007), making it challenging to
account for. Source reconstruction aims to limit volume conduction artefact (He et al.,
2019): in source space, simulated data show that artefactual connectivity due to
volume conduction decreases with increasing Euclidean distance (Anzolin et al., 2019).
This makes our approach to determine areas of minimal signal leakage also relevant to
volume conduction. Further, EEG is a ubiquitous tool in research and clinical settings
(Mclane et al., 2015), a strength when attempting to identify candidate biomarkers.
920
To some extent, our findings are surprising given the field's move towards favouring
zero-phase-exclusive methods (Garcés et al., 2022; Hipp et al., 2012; Miljevic et al.,
2022; Nolte et al., 2004). Zero-phase-exclusive methods were introduced to identify
'true' brain interactions as opposed to trivial artefact (Nolte et al., 2004; Vinck et al.,
2011). Over the last 20+ years, they have consistently been published as improvements
on their corresponding zero-phase-inclusive methods. This position has been adopted
by several studies attempting to identify candidate diagnostic or prognostic biomarkers
(Garcés et al., 2022; Miljevic et al., 2022; Sato et al., 2023). Some authors have argued
that zero-phase connectivity in the brain is physiologically implausible (Miljevic et al.,
2022), despite robust evidence to the contrary.
931
The presence of 'true' zero-phase-delay synchrony in brain is unintuitive: action
potential transmission delays due to axonal conduction and synaptic transmission are
in the order of tens of milliseconds (Uhlhaas et al., 2009; Vicente et al., 2008). Such
time-delays would cause substantial phase-delays between connected brain regions.

Thus, zero-phase-delay functional connectivity across distant neural populations would
not be possible if the brain was a simple linear system. A leading postulated
mechanism for zero-phase-delay connectivity is "resonance-induced synchrony" (Gollo
et al., 2014): bidirectional information transfer between regions acts to mutually alter
their dynamics into a stable zero-phase-delay pattern. Key structures have empirically
shown to facilitate this, including cortico-thalamo-cortical loops (Uhlhaas et al., 2009;
Vicente et al., 2008) and the corpus callosum (Bloom & Hynd, 2005; Musgrave & Gloor,
1980; Vicente et al., 2008; however see Contreras et al., 1996). Other hypothesised
mechanisms of zero-phase-delay synchrony include entrainment (Gray & McCormick,
1996) and emergence (Kopell et al., 2000). It is possible that multiple mechanisms work
in unison (Vicente et al., 2008).
947
The debate central to our study is the extent to which measured zero- and near-zero
phase-delay connectivity in source reconstructed data is artefactual versus 'true'. We
found that most functional connectivity occurred with zero- and near-zero phase-
delays, even when signal leakage was likely negligible. Converging lines of evidence
demonstrate that including zero-phase-delay functional connectivity increases the
performance of functional connectivity metrics as biomarkers. These findings challenge
generally accepted assumptions that zero-phase-exclusive methods are superior to
zero-phase-inclusive methods.
956

Data and code availability

The LEAP Study data is available here, subject to an approved data application: https://redcap.pasteur.fr/surveys/?s=YRFF78PH89

Code will be made available here: https://github.com/cmehra-brain

Author Contributions

Chirag Mehra – Conceptualisation, Methodology, Formal analysis, Writing - Original Draft, Writing - Review & Editing, Visualization

Ahmad Beyh - Formal analysis, Writing - Original Draft, Writing - Review & Editing, Visualization

Petroula Laiou – Resources, Writing - Original Draft

Pilar Garces - Data Curation, Formal analysis, Writing - Review & Editing

Emily JH Jones – Writing - Review & Editing, Funding acquisition, Project administration

Luke Mason - Data Curation, Project administration

Jan Buitelaar – Writing - Review & Editing, Funding acquisition, Project administration

Mark H Johnson - Project administration

Declan Murphy – Funding acquisition, Project administration

Eva Loth – Writing - Review & Editing, Funding acquisition, Project administration

Flavio Dell'Acqua – Formal analysis, Writing - Review & Editing, Funding acquisition, Project administration, Supervision

Joshua B Ewen – Conceptualisation, Writing - Review & Editing, Supervision

Mark P Richardson - Writing - Review & Editing, Supervision, Funding acquisition

Jonathan O'Muircheartaigh - Writing - Review & Editing, Supervision, Funding acquisition

Funding

This work was supported by EU-AIMS (European Autism Interventions), which received support from the Innovative Medicines Initiative Joint Undertaking under grant agreement no. 115300, the resources of which are composed of financial contributions from the European Union's Seventh Framework Programme (grant FP7/2007-2013), from the European Federation of Pharmaceutical Industries and Associations companies' in-kind contributions, and from Autism Speaks. The results leading to this publication have also received funding from the Innovative Medicines Initiative 2 Joint Undertaking under grant agreement No 777394 for the project AIMS-2-TRIALS. This Joint Undertaking receives support from the European Union's Horizon 2020 research and innovation programme and EFPIA and AUTISM SPEAKS, Autistica, and SFARI. [The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.] Any views expressed are those of the author(s) and not necessarily those of the funders (IHI-JU2).

Declaration of Competing Interests

Joshua B Ewen previously consulted for Novartis. In the past 3 years, Jan K Buitelaar has been a consultant to / member of advisory board of / and/or speaker for Takeda, Medice, Angelini, Neuraxpharm and Bitsphi. He is not an employee of any of these companies,

1000   and not a stock shareholder of any of these companies. He has no other financial or
1001   material support, including expert testimony, patents, royalties. Pilar Garces is
1002   employed by Roche Innovation Center, Basel, Switzerland. No other author has
1003   competing interests to declare.

1004   Acknowledgments

1005   We are grateful to the AIMS-2-TRIALS LEAP-group for data collection and quality control
1006   procedures. We are grateful to Dr Piotr J. Franaszczuk for his expert advice on signal
1007   analysis and filtering and to Prof Dr Christine Ecker and team for pre-processing the T1-
1008   weighted MRI images.

1009

References

Adey, W. R., Elul, R., Walter, R. D., & Crandall, P. H. (1967). The cooperative behavior of neuronal populations during sleep and mental tasks. *Electroencephalography and Clinical Neurophysiology*.

Alloway, T. P., Rajendran, G., & Archibald, L. M. D. (2009). Working Memory in Children With Developmental Disorders. *Journal of Learning Disabilities*, *42*(4), 372–382. https://doi.org/10.1177/0022219409335214

Almeida, R., Barbosa, J., & Compte, A. (2015). Neural circuit basis of visuo-spatial working memory precision: a computational and behavioral study. *Journal of Neurophysiology*, *114*(3), 1806–1818. https://doi.org/10.1152/JN.00362.2015

Andersson, J. L. R., Graham, M. S., Drobnjak, I., Zhang, H., Filippini, N., & Bastiani, M. (2017). Towards a comprehensive framework for movement and distortion correction of diffusion MR images: Within volume movement. *NeuroImage*, *152*, 450–466. https://doi.org/10.1016/J.NEUROIMAGE.2017.02.085

Andersson, J. L. R., Graham, M. S., Zsoldos, E., & Sotiropoulos, S. N. (2016). Incorporating outlier detection and replacement into a non-parametric framework for movement and distortion correction of diffusion MR images. *NeuroImage*, *141*, 556–572. https://doi.org/10.1016/J.NEUROIMAGE.2016.06.058

Andersson, J. L. R., Skare, S., & Ashburner, J. (2003). How to correct susceptibility distortions in spin-echo echo-planar images: Application to diffusion tensor imaging. *NeuroImage*, *20*(2), 870–888. https://doi.org/10.1016/S1053-8119(03)00336-7

Andersson, J. L. R., & Sotiropoulos, S. N. (2016). An integrated approach to correction for off-resonance effects and subject movement in diffusion MR imaging. *NeuroImage*, *125*, 1063–1078. https://doi.org/10.1016/J.NEUROIMAGE.2015.10.019

Anzolin, A., Presti, P., Van De Steen, F., Astolfi, L., Haufe, S., & Marinazzo, D. (2019). Quantifying the Effect of Demixing Approaches on Directed Connectivity Estimated Between Reconstructed EEG Sources. *Brain Topography*, *32*(4), 655–674. https://doi.org/10.1007/S10548-019-00705-Z

Bassett, D. S., & Bullmore, E. T. (2017). Small-World Brain Networks Revisited. *The Neuroscientist : A Review Journal Bringing Neurobiology, Neurology and Psychiatry*, *23*(5), 499–516. https://doi.org/10.1177/1073858416667720

Bastos, A. M., & Schoffelen, J. M. (2016). A tutorial review of functional connectivity analysis methods and their interpretational pitfalls. *Frontiers in Systems Neuroscience*, *9*(JAN2016), 165147. https://doi.org/10.3389/FNSYS.2015.00175/BIBTEX

Bauer, M., Buckley, M. G., & Bast, T. (2021a). Individual differences in theta-band oscillations in a spatial memory network revealed by electroencephalography predict rapid place learning. *Brain and Neuroscience Advances*, *5*, 239821282110027. https://doi.org/10.1177/23982128211002725

Bauer, M., Buckley, M. G., & Bast, T. (2021b). Individual differences in theta-band oscillations in a spatial memory network revealed by electroencephalography predict rapid place learning. *Https://Doi.Org/10.1177/23982128211002725*, *5*, 239821282110027. https://doi.org/10.1177/23982128211002725

Baum, G. L., Cui, Z., Roalf, D. R., Ciric, R., Betzel, R. F., Larsen, B., Cieslak, M., Cook, P. A., Xia, C. H., Moore, T. M., Ruparel, K., Oathes, D. J., Alexander-Bloch, A. F., Shinohara, R. T., Raznahan, A., Gur, R. E., Gur, R. C., Bassett, D. S., & Satterthwaite, T. D. (2020). Development of structure–function coupling in human brain networks during youth. *Proceedings of the National Academy of Sciences of the United States of America*,

117(1), 771–778.
https://doi.org/10.1073/PNAS.1912034117/SUPPL_FILE/PNAS.1912034117.SAPP.PDF
BESA® | Brain Electrical Source Analysis: Home. (1995). https://www.besa.de/
Beyh, A., Dell'Acqua, F., Cancemi, D., De Santiago Requejo, F., ffytche, D., & Catani, M. (2022).
The medial occipital longitudinal tract supports early stage encoding of visuospatial
information. *Communications Biology 2022 5:1*, *5*(1), 1–10.
https://doi.org/10.1038/s42003-022-03265-4
Bishop, Geo. H. (1932). Cyclic changes in the excitability of the optic pathway of the rabbit.
American Journal of Physiology, *103*(1), 213–224.
https://doi.org/10.1152/AJPLEGACY.1932.103.1.213
Bloom, J. S., & Hynd, G. W. (2005). The role of the corpus callosum in interhemispheric
transfer of information: Excitation or inhibition? *Neuropsychology Review*, *15*(2), 59–71.
https://doi.org/10.1007/S11065-005-6252-Y/METRICS
Brookes, M. J., Woolrich, M. W., & Barnes, G. R. (2012). Measuring functional connectivity in
MEG: A multivariate approach insensitive to linear source leakage. *NeuroImage*, *63*(2),
910–920. https://doi.org/10.1016/J.NEUROIMAGE.2012.03.048
Brunner, C., Billinger, M., Seeber, M., Mullen, T. R., Makeig, S., Lansky, P., Cohen, M. X., &
Özkurt, T. E. (2016). Volume conduction influences scalp-based connectivity estimates.
Frontiers in Computational Neuroscience, *10*(NOV), 212359.
https://doi.org/10.3389/FNCOM.2016.00121/BIBTEX
Bruns, A., Eckhorn, R., Jokeit, H., & Ebner, A. (2000). Amplitude envelope correlation detects
coupling among incoherent brain signals. *Neuroreport*, *15*, 1509–1514.
Bullmore, E., & Sporns, O. (2012). The economy of brain network organization. *Nature*
Reviews Neuroscience 2012 13:5, *13*(5), 336–349. https://doi.org/10.1038/nrn3214
Buzsáki, G. (2002). Theta oscillations in the hippocampus. *Neuron*, *33*(3), 325–340.
https://doi.org/10.1016/S0896-6273(02)00586-X
Buzsáki, G. (2006). *Rhythms of the Brain*. Oxford Academic.
https://books.google.com/books?hl=en&lr=&id=ldz58irprjYC&oi=fnd&pg=PA4&ots=Q3
Xd56bNDP&sig=95tLOMYuIOWVdNlEQCVN5Y24sBw
Cabral, J., Luckhoo, H., Woolrich, M., Joensson, M., Mohseni, H., Baker, A., Kringelbach, M.
L., & Deco, G. (2014). Exploring mechanisms of spontaneous functional connectivity in
MEG: How delayed network interactions lead to structured amplitude envelopes of
band-pass filtered oscillations. *NeuroImage*, *90*, 423–435.
https://doi.org/10.1016/J.NEUROIMAGE.2013.11.047
Calamante, F. (2019). The Seven Deadly Sins of Measuring Brain Structural Connectivity
Using Diffusion MRI Streamlines Fibre-Tracking. *Diagnostics*, *9*(3).
https://doi.org/10.3390/DIAGNOSTICS9030115
Cauda, F., Nani, A., Liloia, D., Gelmini, G., Mancuso, L., Manuello, J., Panero, M., Duca, S.,
Zang, Y. F., & Costa, T. (2021). Interhemispheric co-alteration of brain homotopic
regions. *Brain Structure & Function*, *226*(7), 2181. https://doi.org/10.1007/S00429-021-
02318-4
Chai, W. J., Abd Hamid, A. I., & Abdullah, J. M. (2018). Working memory from the
psychological and neurosciences perspectives: A review. *Frontiers in Psychology*,
9(MAR), 327922. https://doi.org/10.3389/FPSYG.2018.00401/BIBTEX
Charman, T., Loth, E., Tillmann, J., Crawley, D., Wooldridge, C., Goyard, D., Ahmad, J.,
Auyeung, B., Ambrosino, S., Banaschewski, T., Baron-Cohen, S., Baumeister, S.,
Beckmann, C., Bölte, S., Bourgeron, T., Bours, C., Brammer, M., Brandeis, D., Brogna, C.,

… Buitelaar, J. K. (2017). The EU-AIMS Longitudinal European Autism Project (LEAP):
Clinical characterisation. *Molecular Autism*, *8*(1), 27. https://doi.org/10.1186/s13229-
017-0145-9

Chu, C. J., Tanaka, N., Diaz, J., Edlow, B. L., Wu, O., Hämäläinen, M., Stufflebeam, S., Cash, S.
S., & Kramer, M. A. (2015). EEG functional connectivity is partially predicted by
underlying white matter connectivity. *NeuroImage*, *108*, 23–33.
https://doi.org/10.1016/J.NEUROIMAGE.2014.12.033

Cociu, B. A., Das, S., Billeci, L., Jamal, W., Maharatna, K., Calderoni, S., Narzisi, A., & Muratori,
F. (2018). Multimodal functional and structural brain connectivity analysis in autism: A
preliminary integrated approach with EEG, fMRI, and DTI. *IEEE Transactions on*
Cognitive and Developmental Systems, *10*(2), 213–226.
https://doi.org/10.1109/TCDS.2017.2680408

Colclough, G. L., Brookes, M. J., Smith, S. M., & Woolrich, M. W. (2015). A symmetric
multivariate leakage correction for MEG connectomes. *Neuroimage*, *117*, 439.
https://doi.org/10.1016/J.NEUROIMAGE.2015.03.071

Colclough, G. L., Woolrich, M. W., Tewarie, P. K., Brookes, M. J., Quinn, A. J., & Smith, S. M.
(2016). How reliable are MEG resting-state connectivity metrics? *Neuroimage*, *138*, 284.
https://doi.org/10.1016/J.NEUROIMAGE.2016.05.070

Collin, G., & Van Den Heuvel, M. P. (2013). The ontogeny of the human connectome:
development and dynamic changes of brain connectivity across the life span. *The*
Neuroscientist : A Review Journal Bringing Neurobiology, Neurology and Psychiatry,
19(6), 616–628. https://doi.org/10.1177/1073858413503712

Contreras, D., Destexhe, A., Sejnowski, T. J., & Steriade, M. (1996). Control of spatiotemporal
coherence of a thalamic oscillation by corticothalamic feedback. *Science (New York,*
N.Y.), *274*(5288), 771–774. https://doi.org/10.1126/SCIENCE.274.5288.771

Conturo, T. E., Lori, N. F., Cull, T. S., Akbudak, E., Snyder, A. Z., Shimony, J. S., Mckinstry, R. C.,
Burton, H., & Raichle, M. E. (1999). Tracking neuronal fiber pathways in the living
human brain. *Proceedings of the National Academy of Sciences of the United States of*
America, *96*(18), 10422–10427. https://doi.org/10.1073/PNAS.96.18.10422

Dai, Z., De Souza, J., Lim, J., Ho, P. M., Chen, Y., Li, J., Thakor, N., Bezerianos, A., & Sun, Y.
(2017a). EEG Cortical Connectivity Analysis of Working Memory Reveals Topological
Reorganization in Theta and Alpha Bands. *Frontiers in Human Neuroscience*, *11*.
https://doi.org/10.3389/FNHUM.2017.00237

Dai, Z., De Souza, J., Lim, J., Ho, P. M., Chen, Y., Li, J., Thakor, N., Bezerianos, A., & Sun, Y.
(2017b). Eeg cortical connectivity analysis of working memory reveals topological
reorganization in theta and alpha bands. *Frontiers in Human Neuroscience*, *11*, 237.
https://doi.org/10.3389/FNHUM.2017.00237/BIBTEX

Davis, K. D., Aghaeepour, N., Ahn, A. H., Angst, M. S., Borsook, D., Brenton, A., Burczynski, M.
E., Crean, C., Edwards, R., Gaudilliere, B., Hergenroeder, G. W., Iadarola, M. J., Iyengar,
S., Jiang, Y., Kong, J. T., Mackey, S., Saab, C. Y., Sang, C. N., Scholz, J., … Pelleymounter,
M. A. (2020). Discovery and validation of biomarkers to aid the development of safe
and effective pain therapeutics: challenges and opportunities. *Nature Reviews*
Neurology 2020 16:7, *16*(7), 381–400. https://doi.org/10.1038/s41582-020-0362-2

Dell'Acqua, F., Lacerda, L., Catani, M., & Simmons, A. (2014, May). *Anisotropic Power Maps:*
A diffusion contrast to reveal low anisotropy tissues from HARDI data.
https://archive.ismrm.org/2014/0730.html

Dell'Acqua, F., Scifo, P., Rizzo, G., Catani, M., Simmons, A., Scotti, G., & Fazio, F. (2010). A modified damped Richardson–Lucy algorithm to reduce isotropic background effects in spherical deconvolution. *NeuroImage*, *49*(2), 1446–1458. https://doi.org/10.1016/J.NEUROIMAGE.2009.09.033

Dell'Acqua, F., Simmons, A., Williams, S. C. R., & Catani, M. (2013). Can spherical deconvolution provide more information than fiber orientations? Hindrance modulated orientational anisotropy, a true-tract specific index to characterize white matter diffusion. *Human Brain Mapping*, *34*(10), 2464–2483. https://doi.org/10.1002/HBM.22080

Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*, *134*(1), 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Dennis, E. L., Jahanshad, N., Toga, A. W., McMahon, K. L., de Zubicaray, G. I., Martin, N. G., Wright, M. J., & Thompson, P. M. (2012). Test-Retest Reliability of Graph Theory Measures of Structural Brain Connectivity. *Medical Image Computing and Computer-Assisted Intervention : MICCAI ... International Conference on Medical Image Computing and Computer-Assisted Intervention*, *15*(0 3), 305. https://doi.org/10.1007/978-3-642-33454-2_38

Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., Buckner, R. L., Dale, A. M., Maguire, R. P., Hyman, B. T., Albert, M. S., & Killiany, R. J. (2006). An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *NeuroImage*, *31*(3), 968–980. https://doi.org/10.1016/J.NEUROIMAGE.2006.01.021

Du, Y., Fu, Z., & Calhoun, V. D. (2018). Classification and prediction of brain disorders using functional connectivity: Promising but challenging. *Frontiers in Neuroscience*, *12*(AUG), 373866. https://doi.org/10.3389/FNINS.2018.00525/BIBTEX

Edde, M., Leroux, G., Altena, E., & Chanraud, S. (2021). Functional brain connectivity changes across the human life span: From fetal development to old age. *Journal of Neuroscience Research*, *99*(1), 236–262. https://doi.org/10.1002/JNR.24669

Engel, A. K., Gerloff, C., Hilgetag, C. C., & Nolte, G. (2013). Intrinsic coupling modes: multiscale interactions in ongoing brain activity. *Neuron*, *80*(4), 867–886. https://doi.org/10.1016/J.NEURON.2013.09.038

Engel, A. K., König, P., Kreiter, A. K., & Singer, W. (1991). Interhemispheric Synchronization of Oscillatory Neuronal Responses in Cat Visual Cortex. *Science*, *252*(5009), 1177–1179. https://doi.org/10.1126/SCIENCE.252.5009.1177

Evstigneev, V. V, Kistsen, V. V, & Bulaev, I. V. (2013). The Effect of Structural White Matter Abnormalities on the Clinical Course of Epilepsy. *Advances in Clinical and Experimental Medicine*, *4*(22), 529–537.

Fafrowicz, M., Bohaterewicz, B., Ceglarek, A., Cichocka, M., Lewandowska, K., Sikora-Wachowicz, B., Oginska, H., Beres, A., Olszewska, J., & Marek, T. (2019). Beyond the Low Frequency Fluctuations: Morning and Evening Differences in Human Brain. *Frontiers in Human Neuroscience*, *13*, 477076. https://doi.org/10.3389/FNHUM.2019.00288/BIBTEX

Fair, D. A., Cohen, A. L., Power, J. D., Dosenbach, N. U. F., Church, J. A., Miezin, F. M., Schlaggar, B. L., & Petersen, S. E. (2009). Functional Brain Networks Develop from a "Local to Distributed" Organization. *PLoS Computational Biology*, *5*(5), e1000381. https://doi.org/10.1371/journal.pcbi.1000381

Fell, J., & Axmacher, N. (2011). The role of phase synchronization in memory processes. *Nature Reviews Neuroscience 2011 12:2*, *12*(2), 105–118. https://doi.org/10.1038/nrn2979

Finger, H., Bönstrup, M., Cheng, B., Messé, A., Hilgetag, C., Thomalla, G., Gerloff, C., & König, P. (2016). Modeling of Large-Scale Functional Brain Networks Based on Structural Connectivity from DTI: Comparison with EEG Derived Phase Coupling Networks and Evaluation of Alternative Methods along the Modeling Path. *PLOS Computational Biology*, *12*(8), e1005025. https://doi.org/10.1371/JOURNAL.PCBI.1005025

Fischl, B. (2012). FreeSurfer. *NeuroImage*, *62*(2), 774–781. https://doi.org/10.1016/J.NEUROIMAGE.2012.01.021

Fotiadis, P., Parkes, L., Davis, K. A., Satterthwaite, T. D., Shinohara, R. T., & Bassett, D. S. (2024). Structure–function coupling in macroscale human brain networks. *Nature Reviews Neuroscience 2024*, 1–17. https://doi.org/10.1038/s41583-024-00846-6

Fox, M. D., Snyder, A. Z., Vincent, J. L., Corbetta, M., Van Essen, D. C., & Raichle, M. E. (2005). The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proceedings of the National Academy of Sciences of the United States of America*, *102*(27), 9673–9678. https://doi.org/10.1073/PNAS.0504136102/SUPPL_FILE/04136FIG5.PDF

Fraschini, M., Demuru, M., Crobe, A., Marrosu, F., Stam, C. J., & Hillebrand, A. (2016). The effect of epoch length on estimated EEG functional connectivity and brain network organisation. *Journal of Neural Engineering*, *13*(3). https://doi.org/10.1088/1741-2560/13/3/036015

Fraschini, M., Pani, S. M., Didaci, L., & Marcialis, G. L. (2019). Robustness of functional connectivity metrics for EEG-based personal identification over task-induced intra-class and inter-class variations. *Pattern Recognition Letters*, *125*, 49–54. https://doi.org/10.1016/J.PATREC.2019.03.025

Friston, K. J. (2011). Functional and Effective Connectivity: A Review. *Brain Connectivity*, *1*(1). https://doi.org/10.1089/brain.2011.0008

Gabriel, S., Lau, R. W., & Gabriel, C. (1996). The dielectric properties of biological tissues: II. Measurements in the frequency range 10 Hz to 20 GHz. *Physics in Medicine and Biology*, *41*(11), 2251–2269. https://doi.org/10.1088/0031-9155/41/11/002

Gao, Y., Sengupta, A., Li, M., Zu, Z., Rogers, B. P., Anderson, A. W., Ding, Z., & Gore, J. C. (2020). Functional connectivity of white matter as a biomarker of cognitive decline in Alzheimer's disease. *PLOS ONE*, *15*(10), e0240513. https://doi.org/10.1371/JOURNAL.PONE.0240513

Garcés, P., Baumeister, S., Mason, L., Chatham, C. H., Holiga, S., Dukart, J., Jones, E. J. H., Banaschewski, T., Baron-Cohen, S., Bölte, S., Buitelaar, J. K., Durston, S., Oranje, B., Persico, A. M., Beckmann, C. F., Bougeron, T., Dell'Acqua, F., Ecker, C., Moessnang, C., … Zwiers, M. P. (2022). Resting state EEG power spectrum and functional connectivity in autism: a cross-sectional analysis. *Molecular Autism 2022 13:1*, *13*(1), 1–16. https://doi.org/10.1186/S13229-022-00500-X

Gathercole, S. E., & Alloway, T. P. (2006). Practitioner review: short-term and working memory impairments in neurodevelopmental disorders: diagnosis and remedial support. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, *47*(1), 4–15. https://doi.org/10.1111/J.1469-7610.2005.01446.X

Carmines, E., & Zeller, R. (1979). Reliability and Validity Assessment. *Reliability and Validity Assessment*. https://doi.org/10.4135/9781412985642

Gholipour, T., You, X., Stufflebeam, S. M., Loew, M., Koubeissi, M. Z., Morgan, V. L., & Gaillard, W. D. (2022). Common functional connectivity alterations in focal epilepsies identified by machine learning. *Epilepsia*, *63*(3), 629. https://doi.org/10.1111/EPI.17160

Gil Ávila, C., Bott, F. S., Tiemann, L., Hohn, V. D., May, E. S., Nickel, M. M., Zebhauser, P. T., Gross, J., & Ploner, M. (2023). DISCOVER-EEG: an open, fully automated EEG pipeline for biomarker discovery in clinical neuroscience. *Scientific Data 2023 10:1*, *10*(1), 1–14. https://doi.org/10.1038/s41597-023-02525-0

Gohel, B., Lee, P., Kim, M. Y., Kim, K., & Jeong, Y. (2017). MEG Based Functional Connectivity: Application of ICA to Alleviate Signal Leakage. *IRBM*, *38*(3), 127–137. https://doi.org/10.1016/J.IRBM.2017.03.002

Gollo, L. L., Mirasso, C., Sporns, O., & Breakspear, M. (2014). Mechanisms of Zero-Lag Synchronization in Cortical Motifs. *PLOS Computational Biology*, *10*(4), e1003548. https://doi.org/10.1371/JOURNAL.PCBI.1003548

Gray, C. M., König, P., Engel, A. K., & Singer, W. (1989). Oscillatory responses in cat visual cortex exhibit inter-columnar synchronization which reflects global stimulus properties. *Nature 1989 338:6213*, *338*(6213), 334–337. https://doi.org/10.1038/338334a0

Gray, C. M., & McCormick, D. A. (1996). Chattering cells: superficial pyramidal neurons contributing to the generation of synchronous oscillations in the visual cortex. *Science (New York, N.Y.)*, *274*(5284), 109–113. https://doi.org/10.1126/SCIENCE.274.5284.109

Haller, S., Rodriguez, C., Moser, D., Toma, S., Hofmeister, J., Sinanaj, I., Van De Ville, D., Giannakopoulos, P., & Lovblad, K. O. (2013). Acute caffeine administration impact on working memory-related brain activation and functional connectivity in the elderly: a BOLD and perfusion MRI study. *Neuroscience*, *250*, 364–371. https://doi.org/10.1016/J.NEUROSCIENCE.2013.07.021

Hampson, M., Driesen, N. R., Skudlarski, P., Gore, J. C., & Constable, R. T. (2006). Brain Connectivity Related to Working Memory Performance. *The Journal of Neuroscience*, *26*(51), 13338. https://doi.org/10.1523/JNEUROSCI.3408-06.2006

He, B., Astolfi, L., Valdes-Sosa, P. A., Marinazzo, D., Palva, S. O., Benar, C. G., Michel, C. M., & Koenig, T. (2019). Electrophysiological Brain Connectivity: Theory and Implementation. *IEEE Transactions on Biomedical Engineering*, *66*(7), 2115–2137. https://doi.org/10.1109/TBME.2019.2913928

Heckel, A., Weiler, M., Xia, A., Ruetters, M., Pham, M., Bendszus, M., Heiland, S., & Baeumer, P. (2015). Peripheral Nerve Diffusion Tensor Imaging: Assessment of Axon and Myelin Sheath Integrity. *PLOS ONE*, *10*(6), e0130833. https://doi.org/10.1371/JOURNAL.PONE.0130833

Hilgetag, C. C., Théoret, H., & Pascual-Leone, A. (2001). Enhanced visual spatial attention ipsilateral to rTMS-induced 'virtual lesions' of human parietal cortex. *Nature Neuroscience 2001 4:9*, *4*(9), 953–957. https://doi.org/10.1038/nn0901-953

Hipp, J. F., Hawellek, D. J., Corbetta, M., Siegel, M., & Engel, A. K. (2012). Large-scale cortical correlation structure of spontaneous oscillatory activity. *Nature Neuroscience 2012 15:6*, *15*(6), 884–890. https://doi.org/10.1038/nn.3101

Honey, C. J., Sporns, O., Cammoun, L., Gigandet, X., Thiran, J. P., Meuli, R., & Hagmann, P. (2009). Predicting human resting-state functional connectivity from structural connectivity. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(6), 2035–2040. https://doi.org/10.1073/PNAS.0811168106/SUPPL_FILE/0811168106SI.PDF

Hong, S. J., Xu, T., Nikolaidis, A., Smallwood, J., Margulies, D. S., Bernhardt, B., Vogelstein, J., & Milham, M. P. (2020). Toward a connectivity gradient-based framework for reproducible biomarker discovery. *NeuroImage*, *223*, 117322. https://doi.org/10.1016/J.NEUROIMAGE.2020.117322

Hyman, J. M., Zilli, E. A., Paley, A. M., & Hasselmo, M. E. (2010). Working memory performance correlates with prefrontal-hippocampal theta interactions but not with prefrontal neuron firing rates. *Frontiers in Integrative Neuroscience*, *4*(MARCH 2010), 1162. https://doi.org/10.3389/NEURO.07.002.2010/BIBTEX

Hyvärinen, A. (1999). Fast and robust fixed-point algorithms for independent component analysis. *IEEE Transactions on Neural Networks*, *10*(3), 626–634. https://doi.org/10.1109/72.761722

Jbabdi, S., & Johansen-Berg, H. (2011). Tractography: Where Do We Go from Here? *Https://Home.Liebertpub.Com/Brain*, *1*(3), 169–183. https://doi.org/10.1089/BRAIN.2011.0033

Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved Optimization for the Robust and Accurate Linear Registration and Motion Correction of Brain Images. *NeuroImage*, *17*(2), 825–841. https://doi.org/10.1006/NIMG.2002.1132

Jian, W., Chen, M., & McFarland, D. J. (2017). EEG Based Zero-phase Phase-locking Value (PLV) and Effects of Spatial Filtering During Actual Movement. *Brain Research Bulletin*, *130*, 156. https://doi.org/10.1016/J.BRAINRESBULL.2017.01.023

Jones, M. W., & Wilson, M. A. (2005). Theta Rhythms Coordinate Hippocampal–Prefrontal Interactions in a Spatial Memory Task. *PLOS Biology*, *3*(12), e402. https://doi.org/10.1371/JOURNAL.PBIO.0030402

Jovicich, J., Czanner, S., Greve, D., Haley, E., Van Der Kouwe, A., Gollub, R., Kennedy, D., Schmitt, F., Brown, G., MacFall, J., Fischl, B., & Dale, A. (2006). Reliability in multi-site structural MRI studies: Effects of gradient non-linearity correction on phantom and human data. *NeuroImage*, *30*(2), 436–443. https://doi.org/10.1016/J.NEUROIMAGE.2005.09.046

Kang, S. S., Sponheim, S. R., Chafee, M. V., & MacDonald, A. W. (2011). Disrupted functional connectivity for controlled visual processing as a basis for impaired spatial working memory in schizophrenia. *Neuropsychologia*, *49*(10), 2836–2847. https://doi.org/10.1016/J.NEUROPSYCHOLOGIA.2011.06.009

Katz, B., & Miledi, R. (1965). The measurement of synaptic delay, and the time course of acetylcholine release at the neuromuscular junction. *Proceedings of the Royal Society of London. Series B. Biological Sciences*, *161*, 483–495. https://doi.org/10.1098/RSPB.1965.0016

Kellner, E., Dhital, B., Kiselev, V. G., & Reisert, M. (2016). Gibbs-ringing artifact removal based on local subvoxel-shifts. *Magnetic Resonance in Medicine*, *76*(5), 1574–1581. https://doi.org/10.1002/MRM.26054

Koo, T. K., & Li, M. Y. (2016). A Guideline of Selecting and Reporting Intraclass Correlation Coefficients for Reliability Research. *Journal of Chiropractic Medicine*, *15*(2), 155–163. https://doi.org/10.1016/J.JCM.2016.02.012

Kopell, N., Ermentrout, G. B., Whittington, M. A., & Traub, R. D. (2000). Gamma rhythms and beta rhythms have different synchronization properties. *Proceedings of the National Academy of Sciences of the United States of America*, *97*(4), 1867–1872. https://doi.org/10.1073/PNAS.97.4.1867/ASSET/F27CF424-7A7C-4906-87B7-C2D4BFFFF605/ASSETS/GRAPHIC/PQ0305260005.JPEG

Lachaux, J. P., Rodriguez, E., Martinerie, J., & Varela, F. J. (1999). Measuring phase synchrony in brain signals. *Human Brain Mapping*, *8*(4), 194. https://doi.org/10.1002/(sici)1097-0193(1999)8:4<194::aid-hbm4>3.0.co;2-c

Langer, N., von Bastian, C. C., Wirz, H., Oberauer, K., & Jäncke, L. (2013). The effects of working memory training on functional brain network efficiency. *Cortex*, *49*(9), 2424–2438. https://doi.org/10.1016/J.CORTEX.2013.01.008

Lopes da Silva, F. (2013). EEG and MEG: Relevance to neuroscience. *Neuron*, *80*(5), 1112–1128. https://doi.org/10.1016/j.neuron.2013.10.017

Loth, E., Charman, T., Mason, L., Tillmann, J., Jones, E. J. H., Wooldridge, C., Ahmad, J., Auyeung, B., Brogna, C., Ambrosino, S., Banaschewski, T., Baron-Cohen, S., Baumeister, S., Beckmann, C., Brammer, M., Brandeis, D., Bölte, S., Bourgeron, T., Bours, C., … Buitelaar, J. K. (2017). The EU-AIMS Longitudinal European Autism Project (LEAP): design and methodologies to identify and validate stratification biomarkers for autism spectrum disorders. *Molecular Autism*, *8*(1). https://doi.org/10.1186/S13229-017-0146-8

MacKey, W. E., & Curtis, C. E. (2017). Distinct contributions by frontal and parietal cortices support working memory. *Scientific Reports 2017 7:1*, *7*(1), 1–7. https://doi.org/10.1038/s41598-017-06293-x

Mancuso, L., Costa, T., Nani, A., Manuello, J., Liloia, D., Gelmini, G., Panero, M., Duca, S., & Cauda, F. (2019). The homotopic connectivity of the functional brain: a meta-analytic approach. *Scientific Reports 2019 9:1*, *9*(1), 1–19. https://doi.org/10.1038/s41598-019-40188-3

Mclane, H. C., Berkowitz, A. L., Patenaude, B. N., Mckenzie, E. D., Wolper, E., Wahlster, S., Fink, G., & Mateen, F. J. (2015). Availability, accessibility, and affordability of neurodiagnostic tests in 37 countries. *Neurology*, *85*(18), 1614. https://doi.org/10.1212/WNL.0000000000002090

Mehrkanoon, S., Breakspear, M., Britz, J., & Boonstra, T. W. (2014). Intrinsic Coupling Modes in Source-Reconstructed Electroencephalography. *Https://Home.Liebertpub.Com/Brain*, *4*(10), 812–825. https://doi.org/10.1089/BRAIN.2014.0280

Miljevic, A., Bailey, N. W., Vila-Rodriguez, F., Herring, S. E., & Fitzgerald, P. B. (2022). Electroencephalographic Connectivity: A Fundamental Guide and Checklist for Optimal Study Design and Evaluation. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, *7*(6), 546–554. https://doi.org/10.1016/J.BPSC.2021.10.017

Mill, R. D., Bagic, A., Bostan, A., Schneider, W., & Cole, M. W. (2017). Empirical validation of directed functional connectivity. *NeuroImage*, *146*, 275–287. https://doi.org/10.1016/J.NEUROIMAGE.2016.11.037

Moran, R., Pinotsis, D. A., & Friston, K. (2013). Neural masses and fields in dynamic causal modelling. *Frontiers in Computational Neuroscience*, *7*(APR 2013), 44756. https://doi.org/10.3389/FNCOM.2013.00057/BIBTEX

Musgrave, J., & Gloor, P. (1980). The role of the corpus callosum in bilateral interhemispheric synchrony of spike and wave discharge in feline generalized penicillin epilepsy. *Epilepsia*, *21*(4), 369–378. https://doi.org/10.1111/J.1528-1157.1980.TB04084.X

Nagy, P., Tóth, B., Winkler, I., & Boncz, Á. (2024). The effects of spatial leakage correction on the reliability of EEG-based functional connectivity networks. *Human Brain Mapping*, *45*(8), e26747. https://doi.org/10.1002/HBM.26747

Ni, Z., Gunraj, C., Nelson, A. J., Yeh, I. J., Castillo, G., Hoque, T., & Chen, R. (2009). Two phases of interhemispheric inhibition between motor related cortical areas and the primary

motor cortex in human. *Cerebral Cortex (New York, N.Y. : 1991)*, *19*(7), 1654–1665. https://doi.org/10.1093/CERCOR/BHN201

Noble, S., Scheinost, D., & Constable, R. T. (2021). A guide to the measurement and interpretation of fMRI test-retest reliability. *Current Opinion in Behavioral Sciences*, *40*, 27–32. https://doi.org/10.1016/J.COBEHA.2020.12.012

Nolte, G., Bai, O., Wheaton, L., Mari, Z., Vorbach, S., & Hallett, M. (2004). Identifying true brain interaction from EEG data using the imaginary part of coherency. *Clinical Neurophysiology*, *115*(10), 2292–2307. https://doi.org/10.1016/j.clinph.2004.04.029

O'Keefe, J., & Recce, M. L. (1993). Phase relationship between hippocampal place units and the EEG theta rhythm. *Hippocampus*, *3*(3), 317–330. https://doi.org/10.1002/HIPO.450030307

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2011). FieldTrip: Open Source Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Computational Intelligence and Neuroscience*, *2011*(1), 156869. https://doi.org/10.1155/2011/156869

O'Reilly, C., & Elsabbagh, M. (2021). Intracranial recordings reveal ubiquitous in-phase and in-antiphase functional connectivity between homotopic brain regions in humans. *Journal of Neuroscience Research*, *99*(3), 887–897. https://doi.org/10.1002/JNR.24748

O'Reilly, C., Lewis, J. D., & Elsabbagh, M. (2017). Is functional brain connectivity atypical in autism? A systematic review of EEG and MEG studies. *PloS One*, *12*(5). https://doi.org/10.1371/JOURNAL.PONE.0175870

Owen, A. M., Downes, J. J., Sahakian, B. J., Polkey, C. E., & Robbins, T. W. (1990). Planning and spatial working memory following frontal lobe lesions in man. *Neuropsychologia*, *28*(10), 1021–1034. https://doi.org/10.1016/0028-3932(90)90137-D

Palmer, L. M., Schulz, J. M., Murphy, S. C., Ledergerber, D., Murayama, M., & Larkum, M. E. (2012). The cellular basis of GABA(B)-mediated interhemispheric inhibition. *Science (New York, N.Y.)*, *335*(6071), 989–993. https://doi.org/10.1126/SCIENCE.1217276

Penny W, Friston K, John T. Ashburner J, Kiebel S, & Nichols T. (2006). *Statistical Parametric Mapping: The Analysis of Functional Brain Images*. https://books.google.co.uk/books/about/Statistical_Parametric_Mapping_The_Analy.html?id=G_qdEsDlkp0C&redir_esc=y

Pletcher, M. J., & Pignone, M. (2011). Evaluating the Clinical Utility of a Biomarker: A Review of Methods for Estimating Health Impact. *Circulation*, *123*(10), 1116. https://doi.org/10.1161/CIRCULATIONAHA.110.943860

Plitta, M., Barnes, K. A., Wallace, G. L., Kenworthy, L., & Martin, A. (2015). Resting-state functional connectivity predicts longitudinal change in autistic traits and adaptive functioning in autism. *Proceedings of the National Academy of Sciences of the United States of America*, *112*(48), E6699–E6706. https://doi.org/10.1073/PNAS.1510098112/ASSET/DD00959F-B1DB-4FD5-8E76-F562207A69C9/ASSETS/GRAPHIC/PNAS.1510098112FIG02.JPEG

Pretzsch, C. M., Schäfer, T., Lombardo, M. V., Warrier, V., Mann, C., Bletsch, A., Chatham, C. H., Floris, D. L., Tillmann, J., Yousaf, A., Jones, E., Charman, T., Ambrosino, S., Bourgeron, T., Dumas, G., Loth, E., Oakley, B., Buitelaar, J. K., Cliquet, F., … Ecker, C. (2022). Neurobiological Correlates of Change in Adaptive Behavior in Autism. *The American Journal of Psychiatry*, *179*(5), 336–349. https://doi.org/10.1176/APPI.AJP.21070711/ASSET/IMAGES/LARGE/APPI.AJP.21070711F3.JPEG

Pullon, R. M., Yan, L., Sleigh, J. W., & Warnaby, C. E. (2020). Granger Causality of the Electroencephalogram Reveals Abrupt Global Loss of Cortical Information Flow during Propofol-induced Loss of Responsiveness. *Anesthesiology*, *133*(4), 774. https://doi.org/10.1097/ALN.0000000000003398

Rodriguez, E., George, N., Lachaux, J. P., Martinerie, J., Renault, B., & Varela, F. J. (1999a). Perception's shadow: long-distance synchronization of human brain activity. *Nature 1999 397:6718*, *397*(6718), 430–433. https://doi.org/10.1038/17120

Rodriguez, E., George, N., Lachaux, J. P., Martinerie, J., Renault, B., & Varela, F. J. (1999b). Perception's shadow: long-distance synchronization of human brain activity. *Nature 1999 397:6718*, *397*(6718), 430–433. https://doi.org/10.1038/17120

Roelfsema, P. R., Engel, A. K., König, P., & Singer, W. (1997). Visuomotor integration is associated with zero time-lag synchronization among cortical areas. *Nature 1997 385:6612*, *385*(6612), 157–161. https://doi.org/10.1038/385157a0

Rubinov, M., Kötter, R., Hagmann, P., & Sporns, O. (2009). Brain connectivity toolbox: a collection of complex network measurements and brain connectivity datasets. *NeuroImage*, *47*, S169. https://doi.org/10.1016/S1053-8119(09)71822-1

Salimi, M., Tabasi, F., Nazari, M., Ghazvineh, S., Salimi, A., Jamaati, H., & Raoufy, M. R. (2021). The olfactory bulb modulates entorhinal cortex oscillations during spatial working memory. *Journal of Physiological Sciences*, *71*(1), 1–9. https://doi.org/10.1186/S12576-021-00805-1/FIGURES/5

Sato, J., Safar, K., Vogan, V. M., & Taylor, M. J. (2023). Functional connectivity changes during working memory in autism spectrum disorder: A two-year longitudinal MEG study. *NeuroImage: Clinical*, *37*, 103364. https://doi.org/10.1016/J.NICL.2023.103364

Schilling, K. G., Blaber, J., Huo, Y., Newton, A., Hansen, C., Nath, V., Shafer, A. T., Williams, O., Resnick, S. M., Rogers, B., Anderson, A. W., & Landman, B. A. (2019). Synthesized b0 for diffusion distortion correction (Synb0-DisCo). *Magnetic Resonance Imaging*, *64*, 62–70. https://doi.org/10.1016/J.MRI.2019.05.008

Schmahmann, J. D., & Pandya, D. N. (2009). Fiber Pathways of the Brain. *Fiber Pathways of the Brain*, 1–654. https://doi.org/10.1093/ACPROF:OSO/9780195104233.001.0001

Siapas, A. G., Lubenov, E. V., & Wilson, M. A. (2005). Prefrontal phase locking to hippocampal theta oscillations. *Neuron*, *46*(1), 141–151. https://doi.org/10.1016/j.neuron.2005.02.028

FDA-NIH Biomarker Working Group. (2016). BEST (Biomarkers, EndpointS, and other Tools) Resource. In *BEST (Biomarkers , EndpointS , and other Tools) Resource*. Food and Drug Administration (US). https://www.ncbi.nlm.nih.gov/books/NBK326791/

Singer, W. (1999). Neuronal Synchrony: A Versatile Code for the Definition of Relations? *Neuron*, *24*(1), 49–65. https://doi.org/10.1016/S0896-6273(00)80821-1

Sjöwall, D., Roth, L., Lindqvist, S., & Thorell, L. B. (2013). Multiple deficits in ADHD: executive dysfunction, delay aversion, reaction time variability, and emotional deficits. *Journal of Child Psychology and Psychiatry*, *54*(6), 619–627. https://doi.org/10.1111/JCPP.12006

Srinivasan, R., Winter, W. R., Ding, J., & Nunez, P. L. (2007). EEG and MEG coherence: measures of functional connectivity at distinct spatial scales of neocortical dynamics. *Journal of Neuroscience Methods*, *166*(1), 41. https://doi.org/10.1016/J.JNEUMETH.2007.06.026

Stam, C. J., & van Straaten, E. C. W. (2012). Go with the flow: Use of a directed phase lag index (dPLI) to characterize patterns of phase relations in a large-scale model of brain

dynamics. *NeuroImage*, *62*(3), 1415–1428.
https://doi.org/10.1016/J.NEUROIMAGE.2012.05.050

Stanley, M. L., Simpson, S. L., Dagenbach, D., Lyday, R. G., Burdette, J. H., & Laurienti, P. J. (2015). Changes in Brain Network Efficiency and Working Memory Performance in Aging. *PLoS ONE*, *10*(4). https://doi.org/10.1371/JOURNAL.PONE.0123950

Steinmetz, P. N., Roy, A., Fitzgerald, P. J., Hsiao, S. S., Johnson, K. O., & Niebur, E. (2000). Attention modulates synchronized neuronal firing in primate somatosensory cortex. *Nature 2000 404:6774*, *404*(6774), 187–190. https://doi.org/10.1038/35004588

Uhlhaas, P. J., Pipa, G., Lima, B., Melloni, L., Neuenschwander, S., Nikolić, D., & Singer, W. (2009). Neural synchrony in cortical networks: history, concept and current status. *Frontiers in Integrative Neuroscience*, *3*(JUL). https://doi.org/10.3389/NEURO.07.017.2009

van Blooijs, D., van den Boom, M. A., van der Aar, J. F., Huiskamp, G. M., Castegnaro, G., Demuru, M., Zweiphenning, W. J. E. M., van Eijsden, P., Miller, K. J., Leijten, F. S. S., & Hermes, D. (2023). Developmental trajectory of transmission speed in the human brain. *Nature Neuroscience 2023 26:4*, *26*(4), 537–541. https://doi.org/10.1038/s41593-023-01272-0

van der Velde, B., Haartsen, R., & Kemner, C. (2019). Test-retest reliability of EEG network characteristics in infants. *Brain and Behavior*, *9*(5). https://doi.org/10.1002/BRB3.1269

Varela, F., Lachaux, J. P., Rodriguez, E., & Martinerie, J. (2001). The brainweb: phase synchronization and large-scale integration. *Nature Reviews. Neuroscience*, *2*(4), 229–239. https://doi.org/10.1038/35067550

Veraart, J., Fieremans, E., & Novikov, D. S. (2016). Diffusion MRI noise mapping using random matrix theory. *Magnetic Resonance in Medicine*, *76*(5), 1582–1593. https://doi.org/10.1002/MRM.26059

Vicente, R., Gollo, L. L., Mirasso, C. R., Fischer, I., & Pipa, G. (2008). Dynamical relaying can yield zero time lag neuronal synchrony despite long conduction delays. *Proceedings of the National Academy of Sciences of the United States of America*, *105*(44), 17157–17162. https://doi.org/10.1073/PNAS.0809353105/SUPPL_FILE/0809353105SI.PDF

Vinck, M., Oostenveld, R., Van Wingerden, M., Battaglia, F., & Pennartz, C. M. A. (2011). An improved index of phase-synchronization for electrophysiological data in the presence of volume-conduction, noise and sample-size bias. *NeuroImage*, *55*(4), 1548–1565. https://doi.org/10.1016/J.NEUROIMAGE.2011.01.055

Vorwerk, J., Oostenveld, R., Piastra, M. C., Magyari, L., & Wolters, C. H. (2018). The FieldTrip-SimBio pipeline for EEG forward solutions. *BioMedical Engineering Online*, *17*(1), 1–17. https://doi.org/10.1186/S12938-018-0463-Y/FIGURES/8

Wager, T. D., & Smith, E. E. (2003). Neuroimaging studies of working memory: A meta-analysis. *Cognitive, Affective and Behavioral Neuroscience*, *3*(4), 255–274. https://doi.org/10.3758/CABN.3.4.255/METRICS

Wang, X., Luo, L., Xing, J., Wang, J., Shi, B., Li, Y. M., & Li, Y. G. (2022). Assessment of peripheral neuropathy in type 2 diabetes by diffusion tensor imaging. *Quantitative Imaging in Medicine and Surgery*, *12*(1), 395–405. https://doi.org/10.21037/QIMS-21-126

Wechsler, D. (2011). *Wechsler Abbreviated Scale of Intelligence--Second Edition*. APA PsycTests. https://psycnet.apa.org/doiLanding?doi=10.1037%2Ft15171-000

Welch, P. D. (1967). The Use of Fast Fourier Transform for the Estimation of Power Spectra: A Method Based on Time Averaging Over Short, Modified Periodograms. *IEEE*

Transactions on Audio and Electroacoustics, *15*(2), 70–73.
https://doi.org/10.1109/TAU.1967.1161901
Westerhausen, R., Kreuder, F., Woerner, W., Huster, R. J., Smit, C. M., Schweiger, E., &
Wittling, W. (2006). Interhemispheric transfer time and structural properties of the
corpus callosum. *Neuroscience Letters*, *409*(2), 140–145.
https://doi.org/10.1016/J.NEULET.2006.09.028
Whitford, T. J., Kubicki, M., Ghorashi, S., Schneiderman, J. S., Hawley, K. J., McCarley, R. W.,
Shenton, M. E., & Spencer, K. M. (2011). Predicting inter-hemispheric transfer time from
the diffusion properties of the corpus callosum in healthy individuals and schizophrenia
patients: A combined ERP and DTI study. *NeuroImage*, *54*(3), 2318–2329.
https://doi.org/10.1016/J.NEUROIMAGE.2010.10.048
Wirt, R. A., & Hyman, J. M. (2017). Integrating Spatial Working Memory and Remote
Memory: Interactions between the Medial Prefrontal Cortex and Hippocampus. *Brain*
Sciences 2017, Vol. 7, Page 43, *7*(4), 43. https://doi.org/10.3390/BRAINSCI7040043
Witham, C. L., Wang, M., & Baker, S. N. (2007). Cells in somatosensory areas show synchrony
with beta oscillations in monkey motor cortex. *The European Journal of Neuroscience*,
26(9), 2677–2686. https://doi.org/10.1111/J.1460-9568.2007.05890.X
Wolters, C. H., Anwander, A., Maess, B., MacLeod, R. S., & Friederici, A. D. (2004). The
influence of volume conduction effects on the EEG/MEG reconstruction of the sources
of the Early Left Anterior Negativity. *Conference Proceedings : ... Annual International*
Conference of the IEEE Engineering in Medicine and Biology Society. IEEE Engineering in
Medicine and Biology Society. Annual Conference, *2004*, 3569–3572.
https://doi.org/10.1109/IEMBS.2004.1404003
Womelsdorf, T., & Fries, P. (2007). The role of neuronal synchronization in selective attention.
Current Opinion in Neurobiology, *17*(2), 154–160.
https://doi.org/10.1016/J.CONB.2007.02.002
1550
1551

1552 ## Supplementary Materials

1553

Supplementary Table S1 – Comparing the percentage of near-zero-phase-delay functional connectivity between homotopic- and heterotopic- interhemispheric region-pairs. As there were 34 cortical regions per hemisphere, we randomly selected 34 heterotopic region-pairs, kept consistent across participants, to compare with the 34 homotopic region-pairs. DF = degrees of freedom.

	Percentage of near-zero-phase-delay connectivity between interhemispheric regions		Statistics		
Freq (Hz)	Homotopic (%)	Heterotopic (%)	X^2	DF	*p*
1 – 4	84	55	6.05	1	.014
4 – 8	86	58	5.44	1	.020
8 – 13	83	50	8.19	1	.004
13 – 20	87	61	4.57	1	.033
20 – 32	89	67	3.10	1	.078

1554

Supplementary Table S2 - Intrasession edgewise absolute agreement of EEG functional connectivity methods. Median edgewise absolute agreement was calculated for each edge, using the interclass correlation coefficient (2,1; ICC). n = 99.

Method	Freq (Hz)	Median ICC	IQR	Method	Freq (Hz)	Median ICC	IQR
Coherence	1 - 4	.15	.26	Imaginary part of Coherency	1 - 4	.02	.16
Coherence	4 - 8	.19	.30	Imaginary part of Coherency	4 - 8	.03	.15
Coherence	8 - 13	.20	.30	Imaginary part of Coherency	8 - 13	.07	.16
Coherence	13 - 20	.24	.31	Imaginary part of Coherency	13 - 20	.03	.16
Coherence	20 - 32	.31	.36	Imaginary part of Coherency	20 - 32	.07	.18
PLV	1 - 4	.69	.29	wPLI	1 - 4	-.01	.13
PLV	4 - 8	.75	.28	wPLI	4 - 8	.02	.14
PLV	8 - 13	.71	.31	wPLI	8 - 13	.17	.18
PLV	13 - 20	.84	.20	wPLI	13 - 20	.04	.15
PLV	20 - 32	.88	.14	wPLI	20 - 32	.09	.17
AEC	1 - 4	.37	.43	Orth AEC	1 - 4	.07	.15
AEC	4 - 8	.47	.45	Orth AEC	4 - 8	.10	.18
AEC	8 - 13	.48	.42	Orth AEC	8 - 13	.14	.20
AEC	13 - 20	.57	.42	Orth AEC	13 - 20	.16	.18
AEC	20 - 32	.71	.36	Orth AEC	20 - 32	.04	.15

1555
1556

1557

Supplementary Table S3 – Intrasession consistency of the EEG functional connectivity adjacency matrix. Zero-phase-inclusive methods had moderate to high consistency values, while zero-phase-exclusive methods had near 0 consistency values.

Method	Freq (Hz)	Median Consistency	IQR	Method	Freq (Hz)	Median Consistency	IQR
Coherence	1 - 4	.34	.16	Imaginary part of Coherency	1 - 4	.07	.09
	4 - 8	.40	.18		4 - 8	.10	.11
	8 - 13	.40	.16		8 - 13	.12	.11
	13 - 20	.40	.16		13 - 20	.11	.12
	20 - 32	.47	.18		20 - 32	.16	.16
PLV	1 - 4	.80	.10	wPLI	1 - 4	-.01	.07
	4 - 8	.84	.09		4 - 8	.01	.11
	8 - 13	.82	.09		8 - 13	.12	.16
	13 - 20	.88	.06		13 - 20	.02	.08
	20 - 32	.91	.05		20 - 32	.02	.08
AEC	1 - 4	.55	.15	Orth AEC	1 - 4	.01	.05
	4 - 8	.64	.18		4 - 8	.02	.07
	8 - 13	.60	.15		8 - 13	.04	.13
	13 - 20	.66	.14		13 - 20	.02	.10
	20 - 32	.71	.13		20 - 32	.01	.06

1558
1559

Supplementary Table S4 - Structural-function concordance findings are replicated when analysing the **a)** KCL (n = 28) and **b)** Mannheim (n = 22) study sites separately. Across all frequency bands, concordance was significantly greater than 0 with zero-phase-inclusive methods (one-sided t-test or Wilcoxon sum-ranked test, $\mu = 0$, $p < .0001$, Bonferroni corrected). Concordance was not significantly different from 0 for wPLI and orthogonalized AEC. Surprisingly, concordance was significantly less than zero for the imaginary part of coherency. Concordance for each zero-phase inclusive method was significantly higher than that for its methodologically corresponding zero-phase exclusive method (e.g. coherence versus imaginary part of coherency in 1-4 Hz), for each frequency band (paired samples t-test or Wilcoxon signed-rank test, $p < .0001$, Bonferroni corrected). *** concordance significantly different from zero, $p < .0001$, Bonferroni corrected. ᵂ = data not normally distributed, therefore one-sided Wilcoxon sign-rank test performed, otherwise one-sided t-tests performed. SFC = structure-function concordance; SD = standard deviation.

4a) SFC at KCL (n = 28)

Method	Freq (Hz)	Mean SFC	SD	Method	Freq (Hz)	Mean SFC	SD
Coherence	1 - 4ᵂ	.112***	.049	Imaginary part of Coherency	1 - 4ᵂ	-.041***	.041
Coherence	4 - 8	.119***	.043	Imaginary part of Coherency	4 - 8	-.075***	.042
Coherence	8 - 13	.112***	.035	Imaginary part of Coherency	8 - 13	-.057***	.045
Coherence	13 - 20	.116***	.053	Imaginary part of Coherency	13 - 20	-.066***	.060
Coherence	20 - 32	.124***	.050	Imaginary part of Coherency	20 - 32	-.081***	.041
PLV	1 - 4	.142***	.040	wPLI	1 - 4	.000	.049
PLV	4 - 8ᵂ	.142***	.039	wPLI	4 - 8	-.004	.040
PLV	8 - 13ᵂ	.132***	.030	wPLI	8 - 13	-.013	.040
PLV	13 - 20	.144***	.034	wPLI	13 - 20	-.007	.033
PLV	20 - 32	.146***	.040	wPLI	20 - 32	-.007	.045
AEC	1 - 4ᵂ	.137***	.046	Orthogonalized AEC	1 - 4	.007	.040
AEC	4 - 8	.133***	.038	Orthogonalized AEC	4 - 8	.006	.035
AEC	8 - 13	.120***	.047	Orthogonalized AEC	8 - 13	.001	.049
AEC	13 - 20	.140***	.036	Orthogonalized AEC	13 - 20	.006	.039
AEC	20 - 32	.141***	.039	Orthogonalized AEC	20 - 32	.000	.037

1560
1561

1562 **4b) SFC at Mannheim (n = 22)**

Method	Freq (Hz)	Mean SFC	SD	Method	Freq (Hz)	Mean SFC	SD
Coherence	1 - 4	.080***	.027	Imaginary part of Coherency	1 - 4	-.034***	.035
	4 - 8	.094***	.032		4 - 8	-.043***	.049
	8 - 13	.096***	.043		8 - 13	-.037***	.052
	13 - 20	.094***	.028		13 - 20	-.040***	.033
	20 - 32	.099***	.041		20 - 32	-.061***	.054
PLV	1 - 4	.112***	.028	wPLI	1 - 4	-.003	.030
	4 - 8	.109***	.025		4 - 8	-.018	.039
	8 - 13	.102***	.029		8 - 13	-.007	.038
	13 - 20	.113***	.024		13 - 20	.013	.030
	20 - 32	.115***	.027		20 - 32	.012	.042
AEC	1 - 4	.104***	.029	Orth AEC	1 - 4	.004	.030
	4 - 8	.109***	.030		4 - 8	.015	.038
	8 - 13	.100***	.034		8 - 13	.012	.040
	13 - 20	.108***	.028		13 - 20	.012	.031
	20 - 32	.114***	.030		20 - 32	.011	.030

1563
1564
1565

Supplementary Table S5 – Comparison of mean functional connectivity strength between homotopic region-pairs versus all other region-pairs when using the imaginary part of coherency. Statistics calculated using the Wilcoxon rank sum test, p values Bonferroni corrected.					
	Mean strength between region-pairs with imaginary part of coherency		Statistics		
Freq (Hz)	Homotopic	Non-homotopic	Z	T	p
1 – 4	.097	.137	6.63	2582291	< .0001
4 – 8	.103	.146	7.19	2584398	< .0001
8 – 13	.112	.157	7.52	2585675	< .0001
13 – 20	.119	.167	6.65	2582369	< .0001
20 – 32	.125	.177	7.64	2586111	< .0001

1566
1567

Supplementary Table S6 – General linear model illustrating a significant age x pathlength interaction in predicting longitudinal changes in spatial working memory errors, when using the imaginary part of coherency. Participants aged 6-31-years.

	Est. coeff	Std. Error	t	*p*
Intercept	-223.60	118.10	-1.89	.06
Pathlength (Img COH)	229.10	119.10	1.92	.06
Age	15.42	6.54	2.36	.02
Time 1 spatial working memory errors	-0.60	0.08	-7.35	.00
T2-T1 time interval	0.01	0.01	1.21	.23
Pathlength (Img COH) x age	**-15.80**	**6.58**	**-2.40**	**.02**

Model: *Longitudinal changes in spatial working memory errors ~ Pathlength (derived from imaginary part of coherency) + Age + time 1 spatial working memory errors + T2-T1 time interval + Pathlength (derived from imaginary part of coherency) x Age.* $R^2 = .39$, $R^2_{adj} = .36$, $F_{5/102} = 12.9$, *p* < .001.

1568

Supplementary Table S7 – General linear model illustrating a significant age x pathlength interaction in predicting longitudinal changes in spatial working memory errors, when using coherence. Participants aged 6-31-years

	Est. coeff	Std. Error	t	*p*
Intercept	-348.90	135.00	-2.59	.01
Pathlength (COH)	331.80	126.90	2.62	.01
Age	20.36	7.23	2.82	.01
T2-T1 time interval	0.01	0.01	1.45	.15
Time 1 spatial working memory errors	-0.59	0.08	-7.04	.00
Pathlength (COH) x age	**-19.48**	**6.82**	**-2.86**	**.01**

Model: *Longitudinal changes in spatial working memory errors ~ Pathlength (derived from coherence) + Age + time 1 spatial working memory errors + T2-T1 time interval + Pathlength (derived from coherence) x Age.* $R^2 = .40$, $R^2_{adj} = .36$, $F_{5/102} = 13.3$, *p* < .001.

1569
1570

1571

Supplementary Table S8 – General linear model predicting longitudinal changes in spatial working memory ability in 6-12-year-olds using coherence.				
	β	Std. Error	t	*p*
Intercept	21.1	9.9	2.1	.04
Pathlength (COH), standardised	**3.8**	**1.5**	**2.62**	**.01**
Age	-2.05	0.83	-2.46	.02
Time 1 spatial working memory errors	-0.25	0.17	-1.50	.14
Model: *Longitudinal changes in spatial working memory errors ~ Pathlength (derived from coherence, standardised) + Age + time 1 spatial working memory errors*. R^2 = .30, R^2_{adj} = .23, $F_{3/28}$ = 4.0, *p* = .017.				

1572

Supplementary Table S9 – General linear model predicting longitudinal changes in spatial working memory ability in 18-31-year-olds using PLV.				
	β	Std. Error	t	*p*
Intercept	-5.6	7.8	-0.73	.47
Pathlength (PLV), standardised	**-2.6**	**1.1**	**-2.32**	**.03**
Time 1 spatial working memory errors	-0.96	0.13	-7.15	.00
Age difference	0.02	0.01	1.58	.12
Model: *Longitudinal changes in spatial working memory errors ~ Pathlength (derived from PLV, standardised) + Age difference + time 1 spatial working memory errors*. R^2 = .63, R^2_{adj} = .60, $F_{3/32}$ = 18.5, *p* < .001.				

1573