Humidity-driven ABA depletion determines plant-pathogen competition for leaf

water

Shigetaka Yasuda[1*], Akihisa Shinozawa[2,9], Yuanjie Weng[3,4,9], Arullthevan Rajendram[1,9],

Taishi Hirase[1,9], Haruka Ishizaki[1], Ryuji Suzuki[5], Shioriko Ueda[1], Rahul Sk[6], Yumiko

Takebayashi[3], Izumi Yotsui[2], Masatsugu Toyota[5,7,8], Masanori Okamoto[3], Yusuke Saijo[1*]

[1]Graduate School of Science and Technology, Nara Institute of Science and Technology,

Ikoma, Japan. [2]Department of Bioscience, Tokyo University of Agriculture, Tokyo, Japan

[3]Center for Sustainable Resource Science, RIKEN, Yokohama, Japan. [4]United Graduate

School of Agricultural Science, Tokyo University of Agriculture and Technology, Fuchu,

Japan. [5]Department of Biochemistry and Molecular Biology, Saitama University, Saitama,

Japan. [6]The NODAI Genome Research Center (NGRC), Tokyo University of Agriculture,

Tokyo, Japan. [7]Suntory Rising Stars Encouragement Program in Life Sciences (SunRiSE),

Suntory Foundation for Life Sciences, Kyoto, Japan. [8]College of Plant Science and

Technology, Huazhong Agricultural University, Wuhan, China. [9]These authors

contributed equally: Akihisa Shinozawa, Yuanjie Weng, Arullthevan Rajendram, Taishi

Hirase

*Corresponding authors: shige-yasuda@bs.naist.jp; saijo@bs.naist.jp

Abstract

Bacterial phytopathogens, such as *Pseudomonas syringae* pv. tomato (*Pst*) DC3000,

induce water-soaked lesions in the leaf apoplast under high humidity, facilitating infection.

However, it remains largely unclear how plants regulate their resistance to restrict

bacterial infection in response to humidity. Here, we demonstrate that abscisic acid

(ABA)-catabolizing ABA 8'-hydroxylase, encoded by *CYP707A3*, plays a critical role in

this resistance in *Arabidopsis thaliana*. Elevated humidity induces *CYP707A3* expression,

which is essential for reducing ABA levels and promoting stomatal opening, thereby

limiting bacterial water-soaking and infection following leaf invasion. High humidity also

increases cytosolic Ca^{2+} levels via the Ca^{2+} channels CNGC2 and CNGC4, with partial

involvement from CNGC9, activating the calmodulin-binding transcription activator

CAMTA3 to drive *CYP707A3* induction. However, *Pst* DC3000 counteracts this defense

response using type III secretion effectors, particularly AvrPtoB, facilitating water-

soaking. These findings provide insights into the mechanisms underlying the competition

between plants and bacteria for leaf water under elevated humidity.

Introduction

Bacterial phytopathogens cause foliar diseases that threaten plant survival and crop yields

by proliferating within the leaf apoplast[1]. An important layer of defense against bacterial

infection is provided by cell surface receptor-like kinases/proteins (RLKs/RLPs),

collectively known as pattern recognition receptors (PRRs). PRRs detect microbe-

associated molecular patterns (MAMPs) and endogenous damage-associated molecular

patterns (DAMPs) to activate pattern-triggered immunity (PTI)[2-4]. Following ligand

perception, PRRs form active complexes with co-receptors to trigger intracellular

signaling cascades. This involves rapid cytosolic Ca^{2+} influx, a burst of reactive oxygen

spices (ROS), activation of mitogen activated protein kinase (MAPK) cascades,

production of defense-related phytohormones such as salicylic acid (SA), stomatal

closure, transcriptional reprogramming, and production of antibacterial compounds[2-4]. To

overcome PTI, bacterial pathogens deliver effector proteins into host tissues and cells via

type III secretion (T3S) system[1]. The bacterial phytopathogen *Pseudomonas syringae* pv.

tomato (*Pst*) DC3000 has 36 T3S effectors[5]. Recent studies have shown that *Pst* DC3000

utilizes specific effectors to manipulate the leaf apoplast environment, establishing

conditions favorable for infection[6].

High humidity promotes foliar diseases caused by bacterial pathogens through multiple

mechanisms. Initially, impaired stomatal closure facilitates bacterial invasion into leaves[7].

Once inside, bacterial pathogens induce apoplast hydration under high humidity, a critical

process for disease development known as water-soaking[8]. To induce water-soaking, *Pst*

DC3000 employs the highly conserved effectors HopM1 and AvrE, which manipulate the

host plant's abscisic acid (ABA) pathways, a phytohormone critical for water retention[9,10].

Plants typically increase ABA levels under drought conditions to promote stomatal

closure and conserve water[11]. However, under high humidity, when plants naturally

reduce ABA levels, *Pst* DC3000 stimulates ABA biosynthesis and signaling in the host

through HopM1 and AvrE, thereby enhancing water-soaking and promoting disease

progression[9,10]. Similarly, *Xanthomonas* bacteria exploit host ABA pathways through their effectors to facilitate leaf infection in rice, wheat, and Arabidopsis[12-14]. Despite recent advances in the identification and characterization of bacterial effectors involved in water-soaking, our understanding of the plant resistance mechanisms that counteract bacterial water acquisition remains limited[15,16].

In nature, high humidity often accompanies prolonged rainfall, inducing a wide range of physiological effects in plants, such as defects in cuticle formation, stomatal opening, changes in water uptake and transpiration, and leaf petiole movement (hyponasty)[7,17-19]. In Arabidopsis, high humidity rapidly induces transcriptional and post-transcriptional changes in leaves, modulating phytohormone pathways[17,19]. ABA levels are tightly regulated through the coordination of its biosynthesis and catabolism[20]. Under high humidity, ABA levels decrease via the action of *CYP707A1* and *CYP707A3*, which encode ABA 8'-hydroxylases, thereby facilitating stomatal opening[17]. Concurrently, the production and signaling of defense-promoting salicylic acid (SA) are suppressed, partly due to impaired ubiquitination of NPR1, an SA receptor and key transcriptional co-activator of SA-responsive genes, reducing its binding to target gene promoters[21]. Additionally, high humidity stimulates the biosynthesis of the gaseous phytohormone ethylene, which promotes leaf hyponasty[19]. Despite these extensive changes in phytohormone production and signaling, the mechanisms by which the modulation of host phytohormone pathways under high humidity impacts plant-pathogen interactions remain poorly understood.

In this study, we demonstrate that the high humidity-induced *CYP707A*3 plays a pivotal role in restricting bacterial water-soaking by promoting stomatal opening in leaves. High humidity elevates cytosolic Ca^{2+} concentrations via the Ca^{2+} channels CYCLIC NUCLEOTIDE-GATED CHANNEL 2 (CNGC2), CNGC4, and CNGC9, triggering *CYP707A3* induction through calmodulin-binding transcription activator 3 (CAMTA3). However, during infection, *Pst* DC3000 counteracts this signaling module and suppresses high humidity-induced transcriptional changes through its T3S effectors, with AvrPtoB playing a predominant role. These findings provide valuable insights into the mechanisms regulating plant-pathogen competition for leaf water under high humidity.

Result

***CYP707A3* is required for resistance to bacterial water-soaking under high humidity**

Upon exposure to high humidity, *CYP707A1* and *CYP707A3*, but not *CYP707A2* or *CYP707A4*, were transiently induced in wild-type (WT) leaves within 0.5 h (Fig. 1a). To investigate the role of *CYP707A1* and *CYP707A3* in bacterial resistance, we measured ABA levels and assessed bacterial water-soaking in the corresponding mutants. In WT plants, high humidity triggered a marked reduction in ABA levels within 2 h (Fig. 1b). This ABA decline was completely abolished in the *cyp707a1 cyp707a3* double mutant, which instead exhibited constitutively high ABA accumulation (Fig. 1b), indicating that these genes are required for humidity-induced ABA catabolism.

We next assessed water-soaking in *cyp707a* mutants (Supplementary Fig. 1a, b) using *Pst ΔhrcC*, a strain lacking a functional T3S system. Resistance to water-soaking was evaluated by monitoring leaf water retention following bacterial infiltration under high humidity. At 1 day post-inoculation (dpi), pronounced water-soaking was observed in *cyp707a3* and *cyp707a1 cyp707a3*, but not in WT, *cyp707a1*, or *cyp707a2* (Fig. 1c). The phenotype was more severe in *cyp707a1 cyp707a3* than in *cyp707a3*, suggesting a predominant role for *CYP707A3* with a partial contribution from *CYP707A1* in water-soaking resistance. Accordingly, we used *cyp707a1 cyp707a3* for most subsequent analyses. Under moderate humidity, however, even *cyp707a1 cyp707a3* displayed no detectable water-soaking (Fig. 1d). Consistently, *Pst* Δ*hrcC* growth was significantly higher in *cyp707a1 cyp707a3* than in WT under high humidity, but was indistinguishable under moderate humidity (Fig. 1e). These results demonstrate that *CYP707A3* is crucial for restricting water-soaking during *Pst* DC3000 infection, thereby limiting bacterial proliferation specifically under high humidity.

***CYP707A3*-mediated water-soaking resistance depends on stomatal opening**

Pst DC3000 induces stomatal closure following leaf invasion to facilitate water-soaking[9,10]. Consistent with previous reports[17], loss of *CYP707A3* and *CYP707A1* impaired high-humidity-induced stomatal opening (Fig. 2a, b). To test whether

CYP707A3-mediated bacterial resistance is linked to stomatal regulation, we introduced

the *ost2-3D* mutation, an active allele of the plasma membrane H$^+$-ATPase AHA1 that

drives constitutive stomatal opening[22], into the *cyp707a1 cyp707a3* background

(Supplementary Fig. 2a, b). As expected, the *cyp707a1 cyp707a3 ost2-3D* triple mutant

exhibited enhanced stomatal opening compared with *cyp707a1 cyp707a*3 under both high

and moderate humidity (Fig. 2a, b). Correspondingly, water-soaking was almost

completely abolished in *cyp707a1 cyp707a3 ost2-3D* following *Pst ΔhrcC* infiltration,

similar to *ost2-3D* (Fig. 2c). This suppression was accompanied by markedly reduced *Pst*

ΔhrcC growth in both *cyp707a1 cyp707a3 ost2-3D* and *ost2-3D* (Fig. 2d). These findings

indicate that *CYP707A3* confers resistance to water-soaking, at least in part, by promoting

stomatal opening under high humidity.

ABA depletion enhances bacterial resistance by restricting water-soaking

independently of SA under high humidity

SA antagonizes many ABA-dependent processes and has been reported to suppress water-

soaking under light conditions[16,23,24]. However, both SA and ABA are required for

stomatal closure during early defense responses to bacterial phytopathogens[25]. To assess

whether SA influences *CYP707A3*-mediated water-soaking resistance, we introduced the

sid2-2 mutation, which disrupts ICS1-dependent SA biosynthesis[26], into the *cyp707a1*

cyp707a3 background (Supplementary Fig. 3a, b). The *cyp707a1 cyp707a3 sid2-2* triple

mutant, but not *sid2-2*, exhibited extensive water-soaking following *Pst ΔhrcC* infiltration

(Supplementary Fig. 3c), indicating that ICS1-dependent SA biosynthesis is dispensable

for *CYP707A3*-mediated water-soaking resistance.

We next tested whether ABA depletion alone is sufficient to confer bacterial resistance

under high humidity in an SA-independent manner. To this end, we examined water-

soaking and bacterial growth following *Pst* DC3000 infiltration in the *aao3* mutant, which

is deficient in ABA biosynthesis[27], with or without the *sid2-2* mutation (Supplementary

Fig. 3d, e). Consistent with previous studies on ABA dependence[9,10], *Pst* DC3000-

induced water-soaking was markedly reduced in *aao3*, irrespective of *sid2-2* mutation

(Supplementary Fig. 3f). This reduction correlated with decreased *Pst* DC3000 growth at

3 dpi in both *aao3* and *aao3 sid2*, although the effect was less pronounced in the latter

(Supplementary Fig. 3g). Under moderate humidity, however, suppression of *Pst* DC3000

growth by the *aao3* mutation was observed only in the presence of functional *SID2*

(Supplementary Fig. S3h), as reported previously[24]. Collectively, these results

demonstrate that ABA depletion confers resistance to *Pst* DC3000 under high humidity

by restricting water-soaking development, independently of the SA-ABA antagonism

observed under moderate humidity.

CYP707A3/A1-mediated ABA depletion and PTI contribute additively to bacterial

resistance

High humidity differentially affects PTI responses triggered by flg22, a MAMP derived

from bacterial flagellin[21]. To test whether high humidity-triggered ABA depletion via

CYP707A3 and *CYP707A1* contributes to this modulation, we compared PTI marker gene

expression in WT and *cyp707a1 cyp707a3* following *Pst ΔhrcC* infiltration under

moderate and high humidity. After 24 h acclimation to the respective humidity conditions,

we measured *FRK1*, *NHL10*, and *PHI-1* expression at 1 h post-inoculation (hpi). Under

high humidity, *FRK1* induction was enhanced, *NHL10* induction was unchanged, and

PHI-1 induction was abolished (Supplementary Fig. 4a). These patterns were similar in

WT and *cyp707a1 cyp707a3*, suggesting that CYP707A3/A1-mediated ABA depletion

has limited impact on the humidity-dependent PTI gene regulation.

We next examined the relationship between *CYP707A3*-mediated water-soaking

resistance and PTI by crossing *cyp707a1 cyp707a3* with the *bak1-5 bkk1 cerk1* triple

mutant (hereafter *bbc*), which disrupts PRR activation mediated by the co-receptors

BAK1, BKK1, and CERK1[28] (Supplementary Fig. 4b, c). The *cyp707a1 cyp707a3 bbc*

quintuple mutant exhibited pronounced water-soaking following *Pst ΔhrcC* infiltration,

similar to *cyp707a1 cyp707a3* (Supplementary Fig. 4d). By contrast, *bbc* showed WT-

like or slightly enhanced resistance to water-soaking (Supplementary Fig. 4d), indicating

that humidity-triggered ABA depletion restricts water-soaking independently of

BAK1/BKK1/CERK1-dependent PTI branches. Notably, *Pst ΔhrcC* growth was

significantly higher in *cyp707a1 cyp707a3 bbc* compared to either *cyp707a1 cyp707a3*

or *bbc* (Supplementary Fig. 4e), indicating additive effects. Together, these results

indicate that CYP707A3/A1-mediated ABA depletion and BAK1/BKK1/CERK1-

dependent PTI pathways contribute additively to *Pst* DC3000 resistance under high
humidity.
194

Global transcriptome analysis reveals CAMTA binding motif enrichment in early-
phase high humidity-induced genes
To capture transcriptional reprogramming triggered by high humidity, we performed
RNA-sequencing (RNA-seq) on WT leaves exposed to moderate or high humidity for 0.5
and 5 h. In total, 2394 up-regulated and 2054 down-regulated genes were identified as
differentially expressed (DEGs: fold change > 2, FDR < 0.05) under high humidity
compared with moderate humidity (Fig. 3a and Supplementary Table 1). The DEGs
identified at 0.5 h and 5 h showed limited overlap (up-regulated: 43.5% of 0.5 h and
30.5% of 5 h; down-regulated: 30% of 0.5 h and 6.7% of 5 h) (Fig. 3a), suggesting that
distinct regulatory mechanisms are involved during early and late stages of the response.
This distinction was further supported by principal component analysis (PCA), which
revealed clear separation of transcriptomes according to both humidity conditions and
exposure durations (Fig. 3b).
208

K-means clustering of the 1000 most variable genes yielded 10 distinct clusters (Fig. 3c
and Supplementary Table 2). High humidity-induced genes were distributed across four
clusters, with *CYP707A1* and *CYP707A3* notably present in Cluster 6 (Fig. 3c). Clusters
2 and 6 exhibited rapid transcriptional upregulation within 0.5 h of treatment, Cluster 9
peaked at 0.5 h and gradually declined by 5 h while remaining elevated under high
humidity relative to moderate humidity, and Cluster 1 was primarily induced at 5 h (Fig.
3c). Gene ontology analysis highlighted over-represented biological processes in these
clusters, including 'hypoxia response' (Clusters 1 and 2), 'jasmonate (JA) response'
(Cluster 6), and 'cell wall biogenesis' (Cluster 9) (Supplementary Fig. 5). Notably, *Cis*-
regulatory motif analysis revealed significant enrichment of CAMTA binding motifs,
particularly in Clusters 2 and 6 (Fig. 3d), implicating CAMTA transcription factors as
potential key regulators of early transcriptional responses to high humidity.
221

High humidity-induced *CYP707A3* expression requires CAMTA binding motifs in
its promoter

The promoter region of *CYP707A3* contains two tandem CAMTA binding motifs (CGCG-
box), whereas the *CYP707A1* promoter lacks these elements (Fig. 4a). To assess the
contribution of these motifs to high humidity-induced *CYP707A3* expression, we
transiently expressed a GUS reporter driven by either a native *CYP707A3* promoter
(harboring both CGCG-boxes) or a modified promoter lacking these motifs (ΔCGCG) in
NahG plants (Supplementary Fig. 6a), which are depleted of SA to facilitate
Agrobacterium-mediated gene delivery. *GUS* expression driven by the native promoter
was significantly induced 0.5 h after high humidity exposure, whereas the ΔCGCG
promoter failed to confer this induction (Fig. 4b). In these plants, high humidity induction
of endogenous *CYP707A3* expression remained unaffected (Fig. 4b). These results
indicate that CAMTA binding motifs mediate, at least in part, the high humidity
responsiveness of the *CYP707A3* promoter.
236
To evaluate the physiological relevance of this regulation, we generated *cyp707a3*
transgenic plants expressing CYP707A3 fused to a C-terminal 3×FLAG tag (CYP707A3-
FLAG), driven by either the native or ΔCGCG promoter (Supplementary Fig. 6b, c).
Consistent with the GUS reporter results, high humidity-induced *CYP707A3-FLAG*
expression was markedly reduced in the ΔCGCG promoter line compared with the native
promoter line (Supplementary Fig. 6d). Following *Pst ΔhrcC* infiltration, the impaired
water-soaking resistance of *cyp707a3* was fully restored by *CYP707A3-FLAG* expressed
under the native promoter, but only partially restored under the ΔCGCG promoter
(Supplementary Fig. 6e). Together, these findings indicate that CAMTA-binding motifs
in the *CYP707A3* promoter are critical for its high humidity-induced expression and for
conferring resistance to bacterial water-soaking.
248
High humidity triggers intracellular Ca^{2+} signaling to induce *CYP707A3* expression
CAMTA transcription factors are activated upon binding to Ca^{2+}-CaM[29]. Given their
putative role in high humidity-induced transcription, we hypothesized that elevated
humidity triggers an increase in cytosolic Ca^{2+} levels. Using *35S::GCaMP3* plants
expressing the Ca^{2+} biosensor GCaMP3, we detected a rapid rise in cytosolic Ca^{2+} in leaf
cells within 10 min of high humidity exposure (Supplementary Fig. 7a, b and
Supplementary Movie 1). To test whether cytosolic Ca^{2+} influx contributes to high

humidity-induced *CYP707A3* and *CYP707A1* expression, we applied the Ca^{2+} channel inhibitors $LaCl_3$ and $GdCl_3$ prior to humidity treatment. Both inhibitors significantly suppressed *CYP707A3* induction, but did not suppress *CYP707A1* (Supplementary Fig. 7c), indicating a specific requirement for Ca^{2+} signaling in *CYP707A3* induction. Temporal expression profiling further revealed that high humidity-triggered cytosolic Ca^{2+} elevation precedes *CYP707A3* induction (Supplementary Fig. 7b, d). These findings indicate that humidity elevation triggers cytosolic Ca^{2+} influx in leaves, which in turn mediates *CYP707A3* expression.

CNGC2/4/9 play a critical role in high humidity-induced *CYP707A3* expression

CNGCs, comprising 20 members in Arabidopsis, mediate cytosolic Ca^{2+} influx in diverse physiological processes[30]. To assess their role in high humidity-triggered Ca^{2+} signaling, we examined *CYP707A3* induction in loss-of-function mutants of individual *CNGCs*. An initial screen in 2-week-old plants revealed that high humidity-induced *CYP707A3* expression was significantly reduced in *cngc2*, *cngc4*, and *cngc9* (Supplementary Fig. 8a, b). Subsequent analysis in 5-week-old plants showed that high humidity induction of both *CYP70A3* and *CYP707A1* was impaired in *cngc2* and *cngc4*, whereas only *CYP707A3* induction was reduced in *cngc9* (Fig. 4c). These results suggest that CNGC2 and CNGC4 act cooperatively to mediate Ca^{2+} influx required for high humidity-induced *CYP707A3* expression*, with partial contribution from CNGC9. Collectively, these findings suggest that CNGC2/4-mediated cytosolic Ca^{2+} elevation, together with partial support from CNGC9, is crucial for *CYP707A3* and *CYP707A1* induction in response to high humidity.

CAMTA3 drives *CYP707A3* induction, ABA depletion, and resistance to bacterial water-soaking under high humidity

Among the six CAMTA transcription factors in Arabidopsis, CAMTA3 is a central regulator of responses to diverse stimuli[31-34]. To assess its role in high humidity-induced *CYP707A3* expression, we used an inactive CAMTA3-A855V variant, which carries A855V substitution in the IQ domain that disrupts CAMTA3 function without triggering SA defense activation[35]. Introduction of *CAMTA3* or *CAMTA3-A855V* under the control of a native *CAMTA3* promoter in the *camta2 camta3* mutant background did not alter plant growth or basal SA levels (Supplementary Fig. 9a-c). *CAMTA3* expression restored

WT-like responses to high humidity, including *CYP707A3* induction, ABA depletion, and stomatal opening (Fig. 4d-f). In contrast, these responses were markedly impaired in *CAMTA3-A855V*-expressing plants (Fig. 4d-f), indicating that CAMTA3 activity is required for their induction. Consistent with the absence of CGCG motifs in the *CYP707A1* promoter, high humidity-induced *CYP707A1* expression was unaffected in either *CAMTA3-* or *CAMTA3-A855V*-expressing plants (Fig. 4d), underscoring distinct regulatory mechanisms for *CYP707A3* and *CYP707A1* expression. Moreover, the levels of SA and jasmonoyl-isoleucine (JA-Ile), the major bioactive form of JA, remained unchanged across plant genotypes and humidity treatments (Supplementary Fig. 9c). Together, these findings indicate that CAMTA3 promotes ABA depletion and stomatal opening under high humidity by inducing *CYP707A3*, without altering SA or JA accumulation.

We next tested whether CAMTA3 activity is required to suppress bacterial water-soaking and proliferation under high humidity. Plants expressing *CAMTA3-A855V*, but not *CAMTA3*, exhibited extensive water-soaking and permitted increased growth of *Pst Δhrc* under high humidity (Fig. 4g, h), whereas bacterial growth was unaffected under moderate humidity (Fig. 4i). These findings demonstrate that CAMTA3 activity is required for resistance to bacterial water-soaking and infection specifically under high humidity. This is consistent with a model in which elevated humidity triggers Ca^{2+} influx through CNGC2/4/9, activating CAMTA3 to drive *CYP707A3*-mediated resistance.

Pst DC3000 suppresses high humidity-induced *CYP707A3* expression via type III secretion effectors

Pst DC3000 enhances host ABA responses by manipulating ABA biosynthesis, signaling, and translocation to promote water-soaking under high humidity[9,10]. We tested whether *Pst* DC3000 also targets high humidity-triggered ABA depletion via *CYP707A3* and *CYP707A1*. WT leaves were inoculated with *Pst* DC3000, *Pst* Δ*hrcC*, and *Pst* Δ28E (lacking 28 T3S effectors)[36]. At 8 hpi under moderate humidity, plants were shifted to moderate or high humidity for 0.5 h before RT-qPCR analysis (Fig. 5a). At this point, bacterial populations were comparable between *Pst* DC3000 and *Pst* Δ*hrcC* (Fig. 5b), minimizing potential effects of differential bacterial growth. Under these conditions,

CYP707A1 was not induced by high humidity, even in the absence of bacterial inoculation

(Fig. 5C), likely due to desensitization from water infiltration. By contrast, *CYP707A3*

was significantly induced by high humidity in mock-inoculated leaves and in those

inoculated with *Pst* Δ*hrcC* or *Pst* Δ28E, but this induction was markedly suppressed by

Pst DC3000 (Fig. 5c). CAMTA3 protein accumulation was unaffected by inoculation

with the WT or mutant bacteria (Supplementary Fig. 10), suggesting that suppression is

not due to CAMTA3 depletion. Consistent with previous reports[9,10], *Pst* DC3000, but not

Pst Δ*hrcC* or *Pst* Δ28E, also induced *NCED3*, encoding a key ABA biosynthesis enzyme,

under these conditions (Fig. 5c). Together, these results indicate that *Pst* DC3000 employs

T3S effectors to suppress *CYP707A3* induction while promoting *NCED3* expression,

thereby maintaining elevated ABA levels under high humidity to facilitate infection.

Pst DC3000 blocks high humidity-driven plant transcriptome reprogramming via

type III secretion effectors

To further investigate how *Pst* DC3000 modulates plant responses to high humidity, we

performed RNA-seq analysis on leaves inoculated with *Pst* DC3000, *Pst* Δ*hrcC*, and *Pst*

Δ28E for 8 h, followed by a 0.5 h of exposure to moderate or high humidity. PCA revealed

a clear separation of leaf transcriptomes based on humidity conditions in non-inoculated

leaves and those inoculated with *Pst* Δ*hrcC* or *Pst* Δ28E (Fig. 5d). In contrast, this

separation was absent in leaves inoculated with *Pst* DC3000 (Fig. 5d), indicating that *Pst*

DC3000 broadly disrupts transcriptomic responses to high humidity via T3S effectors. K-

means clustering of the 2000 most variable genes resulted in 10 distinct clusters (Fig. 5e

and Supplementary Table 3). High humidity-induced genes were enriched in three

clusters, with Clusters 3 and 8 containing *CYP707A1* and *CYP707A3*, respectively (Fig.

5e). *Pst* DC3000 strongly suppressed high humidity-induced up-regulation in Clusters 3

and 8, and partially in Cluster 4 (Fig. 5e). Notably, Cluster 3 genes were upregulated

following *Pst* DC3000 inoculation under moderate humidity, irrespective of subsequent

humidity conditions (Fig. 5e). These findings indicate that *Pst* DC3000, through T3S

effectors, not only suppresses CAMTA3-regulated genes but also broadly alters early

transcriptional networks responsive to high humidity during infection.

AvrPtoB contributes to suppression of *CYP707A3*-mediated water-soaking

resistance

To identify the bacterial effector(s) responsible for suppressing *CYP707A3* induction under high humidity, we analyzed multiple effector-deletion mutants of *Pst* DC3000[36]. High humidity-induced *CYP707A3* expression was significantly suppressed in WT leaves at 8 hpi with *Pst* Δ14E (lacking 14 T3S effectors) and *Pst* Δ20E (lacking 20 T3S effectors), similar to *Pst* DC3000 (Fig. 6a), at a time when T3S-dependent bacterial growth was not evident (Fig. 5b). In contrast, *Pst* Δ28E failed to suppress *CYP707A3* induction, indicating that one or more of the remaining eight effectors (*HopK1*, *HopY1*, *HopB1*, *HopAF1*, *AvrPtoB*, *AvrPto*, *HopE1*, and *HopA1*) are required for this suppression (Fig. 6a).

Given that *in-planta* expression of *AvrPtoB* has been associated with increased ABA levels through induction of *NCED3*[37], we tested whether AvrPtoB contributes to *CYP707A3* suppression. To this end, we generated transgenic plants expressing either *AvrPtoB* or its partially redundant counterpart, *AvrPto*[38], under a DEX-inducible promoter (*DEX::AvrPtoB* and *DEX::AvrPto*; Fig. 6b, c and Supplementary Fig. 11a). Following DEX treatment, high humidity-induced *CYP707A3* expression was specifically suppressed in *DEX::AvrPtoB*, but not in *DEX::AvrPto*, despite lower protein accumulation of AvrPtoB compared to AvrPto (Fig. 6b, c and Supplementary Fig. 11b). An E3 ubiquitin ligase-deficient variant, AvrPtoB-F479A[39], retained the ability to suppress *CYP707A3* induction (Fig. 6d and Supplementary Fig. 11a), suggesting that this activity is independent of its E3 ligase function. In contrast, high humidity induction of *CYP707A1* was specifically suppressed in *AvrPto*-expressing leaves (Supplementary Fig. 11c-e). Collectively, these findings suggest that bacteria selectively target *CYP707A3* and *CYP707A1* with distinct T3S effectors, AvrPtoB and AvrPto, respectively, under high humidity.

Consistent with the predominant role for *CYP707A3* in water-soaking resistance (Fig. 1c), high humidity-triggered ABA depletion and stomatal opening were also compromised in *DEX::AvrPtoB* following DEX treatment (Fig. 6e, f). Importantly, DEX-induced *AvrPtoB* expression substantially restored water-soaking in leaves infiltrated with *Pst* DC3000 *ΔhrcC* (Fig. 6g). The results indicate that AvrPtoB plays a critical role in counteracting

CYP707A3-mediated water-soaking resistance under high humidity.

Pst DC3000 employs AvrPtoB to suppress *CYP707A3* expression and promote water-
soaking under high humidity

We next evaluated the physiological relevance of AvrPtoB and AvrPto in promoting ABA-
dependent bacterial infection under high humidity using *Pst* DC3000 mutant strain
lacking both *AvrPtoB* and *AvrPto* (*Pst* Δ*avrPtoB* Δ*avrPto*). Plants were shifted from
moderate to high humidity at 2 hpi, following conditions conventionally used in previous
studies[8-10] (Fig. 7a and Supplementary Fig. S12a). RT-qPCR analysis revealed that
CYP707A3 expression was suppressed from 6 to 24 hpi with *Pst* DC3000, despite
continuous high humidity from 2 hpi (Supplementary Fig. S12b). This suppression was
absent with *Pst* Δ28E, whereas with *Pst* Δ*avrPtoB* Δ*avrPto* it was impaired at 6 hpi
but restored at 12-24 hpi (Supplementary Fig. S12b), indicating that AvrPtoB and/or
AvrPto mediate early suppression of *CYP707A3*, while other effectors contribute to later
suppression. Consistently, ABA levels increased at 6 hpi with *Pst* DC3000, but not with
Pst Δ*avrPtoB* Δ*avrPto* or *Pst* Δ28E (Supplementary Fig S12c). This initial increase was
followed by further accumulation at 12 and 24 hpi with *Pst* DC3000 and partially with
Pst Δ*avrPtoB* Δ*avrPto*, but not with *Pst* Δ28E. Together, these findings indicate that
AvrPtoB- and/or AvrPto-mediated suppression of *CYP707A3* is essential for the initial
phase of ABA accumulation under high humidity, which is critical for subsequent water-
soaking and bacterial proliferation.

To dissect individual contributions of these effectors, we generated *Pst* DC3000 mutant
strains lacking either *AvrPtoB* (*Pst* Δ*avrPtoB*) or *AvrPto* (*Pst* Δ*avrPto*) (Fig. 7b and
Supplementary Fig. S13a, b). The two effectors did not affect each other's expression
during infection (Fig. 7b). Consistent with DEX induction (Fig. 6b, c), high humidity-
induced *CYP707A3* expression was suppressed by *Pst* DC3000 and *Pst* Δ*avrPto*, but
permitted with *Pst* Δ*avrPtoB* and *Pst* Δ*avrPtoB* Δ*avrPto* (Fig. 7b). By contrast, enhanced
CYP707A1 expression relative to *Pst* DC3000 was observed only with *Pst* Δ*avrPtoB*
Δ*avrPto* (Fig. 7b), suggesting redundancy in *CYP707A1* suppression. For *NCED3*

induction, *Pst* Δ*avrPto* resembled *Pst* DC3000, whereas *Pst* Δ*avrPtoB* and *Pst* Δ*avrPtoB* Δ*avrPto* failed to induce it (Fig. 7b), indicating a predominant role of AvrPtoB in both *CYP707A3* suppression and *NCED3* induction. Correspondingly, ABA accumulation at 6 hpi was enhanced with *Pst* DC3000 and *Pst* Δ*avrPto*, but not with *Pst* Δ*avrPtoB* or *Pst* Δ*avrPtoB* Δ*avrPto* (Fig. 7c). Together, these results indicate that AvrPtoB is the primary determinant of ABA accumulation under high humidity, acting through both *CYP707A3* suppression and *NCED3* induction.

We next assessed the impact of AvrPtoB on bacterial water-soaking and infection. Under high humidity, *Pst* DC3000 induced extensive water-soaking, which was strongly reduced in leaves infiltrated with *Pst* Δ*avrPtoB* or *Pst* Δ*avrPtoB* Δ*avrPto*, while *Pst* Δ*avrPto* retained partial activity (Supplementary Fig. 13c, d). These results indicate a predominant role for AvrPtoB, with a moderate contribution from AvrPto. Importantly, delayed suppression of *CYP707A3* by *Pst* Δ*avrPtoB* Δ*avrPto* at 12-24 hpi (Supplementary Fig. 12b) was insufficient to promote water-soaking, highlighting the importance of early ABA depletion for this virulence activity. Moreover, *Pst* Δ*avrPtoB* induced smaller water-soaked area compared to *Pst* DC3000 even in *cyp707a1 cyp707a3* leaves (Fig, 7d, e). This reduction correlated with milder disease symptoms and lower bacterial growth in both WT and *cyp707a1 cyp707a3* plants (Fig. 7f, g). These findings indicate that AvrPtoB also targets an additional, *CYP707A1/CYP707A3*-independent layer of water-soaking resistance, which may align with the additive roles of BAK1/BKK1/CERK1-mediated PTI and ABA-catabolism-mediated defenses in restricting bacterial growth under high humidity (Supplementary Fig. 4d, e).

Discussion

Terrestrial plants are constantly exposed to fluctuations in air humidity and have evolved elaborate adaptation mechanisms[17,19,40-42]. High humidity, often following prolonged rainfall, substantially exacerbates plant diseases caused by fungal and bacterial pathogens[8,43]. It is therefore plausible that plants pre-condition or activate defense responses in anticipation of these threats when sensing increased humidity. In this study, we demonstrate that the CNGC2/4/9-CAMTA3-CYP707A3 module is involved in response to elevated humidity, conferring bacterial resistance (Fig. 7h). Our findings

suggest that high humidity acts as a danger signal to enhance defense readiness. This

aligns with earlier studies that high humidity induces ROS accumulation in Arabidopsis

leaf apoplast, a hallmark of defense responses during pathogen challenge[19,21].

Additionally, high humidity increases cuticle permeability, which has been associated

with enhanced resistance to the fungal pathogen *Botrytis cinerea*[18,44,45]. Rainfall and

submergence have also been linked to strengthened disease resistance in plants[34,46]. These

findings suggest that plants possess the mechanisms linking the sensing of increased

water or humidity levels to the activation or reinforcement of defense responses. Our

findings reveal a mechanism linking humidity perception to the activation of defenses

that prevent bacterial access to leaf water.

Antagonistic interactions between ABA and SA signaling have been well

documented[23,24,47,48], yet their modulation under varying environmental conditions

remain under-explored. In Arabidopsis, *CYP707A3* overexpression enhances

transpiration rates[49]. ABA application suppresses systemic acquired resistance (SAR)

inducers acting both upstream and downstream of SA accumulation[23]. Enhanced SA

production under continuous light suppresses *Pst* DC3000-induced water-soaking

without interfering with *NCED3* induction[16]. We show that CYP707A3-mediated ABA

depletion confers water-soaking resistance independently of SA under high humidity,

whereas ABA deficiency enhances bacterial resistance in an SA-dependent manner under

moderate humidity. These observations underscore a humidity-dependent differentiation

in bacterial virulence strategies and the role of ABA depletion in bacterial resistance. The

distinct regulation of SA-ABA antagonism aligns with the decrease in SA production and

effectiveness under high humidity[21]. Notably, however, genes associated with SA defense,

such as *SARD1*, *CBP60g*, and *BCS1*[50-52], are induced under high humidity even in the

absence of pathogen challenge (Cluster 3 in Fig 5e and Supplementary Table 3),

suggesting potential coupling between humidity sensing and SA-mediated resistance,

which may be further potentiated by PRR signaling (Supplementary Fig. 4e).

To our knowledge, this study provides the first evidence that phytopathogens promote

infection by suppressing high humidity responses in host plants through effector proteins.

Our RNA-seq analyses (Fig. 5d, e) demonstrate that *Pst* DC3000 employs multiple T3S

effectors to inhibit humidity-triggered transcriptomic changes, thereby circumventing *CYP707A3*-mediated water-soaking resistance (Fig. 7h). Interestingly, *Pst* DC3000 T3S effectors also induce a subset of humidity-inducible genes, including *CYP707A1*, regardless of humidity conditions (Fig. 5e). The functional significance of their induction represents an open question for future studies. *CYP707A1* is predominantly induced in guard cells under high humidity[17] and *Pst* DC3000 elevates ABA levels in guard cells[10]. Whether bacterial *CYP707A1* induction occurs specifically in guard cells or in other tissues remains to be determined. Suppression of *CYP707A3* induction alone is sufficient to permit ABA-mediated stomatal closure, facilitating water-soaking under high humidity, as indicated by the impaired water-soaking resistance in *cyp707a3* mutants (Fig. 1c). Recent findings indicate that post-invasion stomatal reopening, rather than the initial closure to restrict bacterial entry, is critical for resisting bacterial water-soaking and infection[53], emphasizing the role of CYP707A-mediated ABA depletion as a core defense barrier limiting bacterial access to apoplastic water.

We identify AvrPtoB as a major T3S effector suppressing this defense. Genetic analyses show that among the 36 T3S effectors, AvrPtoB has a prominent role in blocking high humidity-induced *CYP707A3* expression (Fig. 7h). Earlier studies reported that AvrPtoB can target CYP707A3 and paralogs for proteasomal degradation when these proteins are transiently expressed in *Nicotiana benthamiana*[54]. By suppressing ABA catabolism and simultaneously promoting ABA biosynthesis via *NCED3* induction[37,54], AvrPtoB manipulates both arms of host ABA balance, facilitating stomatal closure and maintaining elevated ABA under high humidity. How AvrPtoB E3 ubiquitin ligase activity, known to act on membrane-localized immune receptors and transcription cofactors[55-60], contributes to suppression of *CYP707A3* and induction of *NCED3* remains to be determined.

Our work identifies CAMTA3 transcription regulator and CNGC2/4/9 Ca^{2+} channels as key regulators of humidity-induced *CYP707A3* expression, consistent with recent findings[61]. While the oomycete effector AVRblb2 has been shown to suppress CNGC activity[62], the potential targeting of CNGCs by *Pst* DC3000 effectors remains unexplored. Notably, the presence of a dominant-negative CAMTA3-A855V allele significantly impairs resistance to bacterial and fungal pathogens, SAR, as well as early transcriptional

responses shared by PTI, effector-triggered immunity and abiotic stress responses[63-65].
This underscores the critical role of CAMTA3 in promoting and/or priming defense
activation under high humidity. The disruption of *CAMTA1/2/3* is also linked to
nucleotide-biding leucine-rich repeat receptor activation leading to SA defense activation
and autoimmunity[35,66], suggesting that CAMTAs might be targeted by pathogen effectors
to counteract humidity-induced defenses. Future studies on AvrPtoB and other effectors,
including HopAM1, HopM1, and AvrE, which promote ABA responses[9,42,67], and their
interactions with the CNGC2/4/9-CAMTA3-CYP707A3 signaling module and another
water-soaking defense(s) will deepen our understanding of the interplay between
humidity-triggered defenses and effector-mediated host manipulation. This work
advances our knowledge of how plants and pathogens optimize immunity and virulence
strategies under high humidity, driven by their competition for water as a decisive factor.
522
523 **Methods**
524
Plant materials and growth conditions
Arabidopsis accession Columbia-0 (Col-0) was used as the wild-type (WT) in this study.
The *cyp707a1 cyp707a3 ost2-3D*, *cyp707a1 cyp707a3 sid2-2*, *aao3 sid2-2*, *bak1-5 bkk1*
cerk1 (*bbc*), and *cyp707a1 cyp707a3 bbc* mutants were generated by genetic crossing.
The genotypes of Arabidopsis mutants and transgenic lines were verified by PCR, Sanger-
sequencing, or chemical selection. Plant materials and genotyping primers used are listed
in Supplementary Tables 4 and 6, respectively.
532
Surface-sterilized seeds were sown on half-strength Murashige and Skoog (MS) medium
[2.3 g/L Murashige and Skoog Plant Salt Mixture (FUJIFILM Wako Pure Chemical, Cat#
392-00591), 1% (w/v) sucrose, 0.5 g/L MES, 1% (w/v) agar (FUJIFILM Wako Pure
Chemical, Cat# 016-11875), pH 5.6-5.8 adjusted by KOH] followed by stratification for
1-4 days in the dark at 4°C. Plates were placed vertically in a growth chamber [22°C, 16
h light/8 h dark (long-day) or 8 h light/16 h dark (short-day), illuminated with daylight
white fluorescent lamps (2 lamps/side)]. Seedlings were then transplanted into a soli
mixture [50% (v/v) Super Mix A (Sakata Seed), 50% (v/v) Vermiculite G20 (Nittai)] and
cultivated in either a growth chamber [22°C, 60% RH, 8 h light/16 h dark, illuminated

with daylight white fluorescent lamps (2 lamps/side)] or a growth room [22°C, 30-70%

RH, 16 h light/8 h dark, illuminated with daylight white fluorescent lamps (4

lamps/shelf)]. Plants were irrigated with either tap water or a modified Hoagland solution

[1.67 mM KNO_3, 0.67 mM $MgSO_4$, 5.21 µM KH_2PO_4, 28.3 µM KCl, 15.5 µM H_3BO_4,

3.07 µM $MnCl_2$, 0.28 µM $ZnSO_4$, 0.11 µM $CuSO_4$, 13.3 µg/L 80% molybdic acid

(FUJIFILM Wako Pure Chemical, Cat# 134-03332), 1.67 mM $Ca(NO_3)_2$, 46.3 µM

Fe(III)-EDTA (Dojindo, Cat# 345-01245)].

For general cultivation, 2-week-old seedings were transplanted and grown under long-

day conditions with irrigation using the modified Hoagland solution. For experiments

using 5-week-old plants, 2-week-old seedlings were transplanted and grown under short-

day conditions with irrigation using the modified Hoagland solution. For experiments

using 2-week-old plants, 4-day-old seedlings were transplanted and grown under long-

day conditions with irrigation using tap water.

Plasmid construction

For the generation of Arabidopsis transgenic plants, the promoter region (-2776 bp to -1

bp relative to the start codon) and the genomic complementation sequence (-2776 to

+1930 bp relative to the start codon) of *CYP707A3* were amplified from Arabidopsis WT

genomic DNA. The full-length coding sequences (excluding the stop codon) of AvrPtoB

and AvrPto were amplified from *Pst* DC3000 genomic DNA. A deletion of the CGCG

boxes in the *CYP707A3* promoter (-2220 bp to -2150 bp relative to the start codon) and

the AvrPtoB F479A substitution were introduced by overlap extension PCR. PCR

amplifications were performed using PrimeSTAR GXL DNA Polymerase (Takara Bio,

Cat# R050A). Amplified products were cloned into pENTR/D-TOPO using either the

pENTR/D-TOPO Cloning Kit (Thermo Scientific, Cat# K240020SP) or the In-Fusion

Snap Assembly Master Mix (Takara Bio, Cat# 638948). The resulting entry clones were

verified by Sanger sequencing with the BigDye Terminator v3.1 Cycle Sequencing Kit

(Applied Biosystems, Cat# 4337455) and recombined with destination vectors using

Gateway LR Clonase II Enzyme mix (Invitrogen, Cat# 11791020). The following

destination vectors were used: pGWB533[68] for GUS reporter constructs, pGWB501 (C-

3×FLAG) for *CYP707A3* complementation constructs, and pTA70001-DEST (Addgene,

Cat# 71745) for DEX-inducible constructs.

The pGWB501 (C-3×FLAG) vector was generated by cloning the 3×FLAG tag coding

sequence from pAMPAT-GW-3×FLAG[69] into the AfeI site of pGWB501[68] using In-

Fusion Snap Assembly Master Mix. For constructs encoding C-terminally 3×FLAG-

tagged AvrPtoB, AvrPto, and AvrPtoB-F479A, entry clones were recombined with

pAMPAT-GW-3×FLAG, and the resulting coding sequences were subsequently cloned

into pENTR/D-TOPO.

For the generation of Pst DC3000 mutant strains, the flanking 1 kb regions upstream and

downstream of either *avrPtoB* or *avrPto* were amplified from *Pst* DC3000 genomic DNA

by PCR using PrimeSTAR GXL DNA Polymerase. Amplified fragments were assembled

and cloned into EcoRI- and XbaI-digested pK18mobsacB using In-Fusion Snap

Assembly Master Mix. The resulting constructs were verified by Sanger sequencing with

the BigDye Terminator v3.1 Cycle Sequencing Kit.

All cloning procedures were performed according to the manufacturer's instructions.

Plasmids generated in this study and the primers for cloning are listed in Supplementary

Tables 5 and 6, respectively.

Generation of Arabidopsis transgenic plants

Expression vectors were introduced into *Agrobacterium tumefaciens* strain

GV3101::pMP90 by electroporation. Transgenic plants were generated by transforming

WT or *cyp707a3* mutant plants using the floral dip method[70]. For the *CYP707A3*

complementation experiments, T2 segregating lines were analyzed, whereas T3 or T4

homozygous lines were selected and used for all other experiments. The transgenic plants

used in this study are listed in Supplementary Table 4.

Generation of *Pst* DC3000 mutant strains

Pst DC3000 mutant strains lacking either *avrPtoB* or *avrPto* were generated by

introducing the pK18mobsacB-based constructs into *Pst* DC3000, as previously

described[71]. Deletions of *avrPtoB* and *avrPto* were verified by PCR using EmeraldAmp

MAX PCR Master Mix (Takara Bio, Cat# RR320A) and Sanger sequencing with the BigDye Terminator v3.1 Cycle Sequencing Kit.. The primers used for PCR and Sanger sequencing are listed in Supplementary Table 6.

Chemicals

The following stock solutions were prepared and stored at $-30°C$: $LaCl_3$ (FUJIFILM Wako Pure Chemical, Cat# 123-04222), 1 M in water; $GdCl_3$ (FUJIFILM Wako Pure Chemical, Cat# 078-02661), 1 M in water; Acetosyringone (FUJIFILM Wako Pure Chemical, Cat# 320-29611), 100 mM in DMSO; Dexamethasone (DEX) (FUJIFILM Wako Pure Chemical, Cat# 047-18863), 10 mM in ethanol.

Humidity treatment

High humidity treatment was performed by adding approximately 200 mL of modified Hoagland solution or tap water (for 2-week-old plants grown under long-day conditions) to a plastic container (Daiso, Cat# 4550480269726), spraying tap water onto the container walls, placing the plants inside, sealing the container with plastic wrap, and immediately returning it to the growth conditions. Moderate humidity treatment was performed by adding approximately 500 mL of modified Hoagland solution or tap water (for 2-week-old plants grown under long-day conditions) to a plastic tray (As One, Cat# 1-4618-01), placing the plants inside, and immediately returning it to the growth conditions.

For inhibitor treatments and *Pst* DC3000 inoculation experiments, the indicated solutions were infiltrated into three leaves per 5-week-old WT plant using a needleless syringe (Terumo, Cat# SS-01T). Excess solution on the leaf surfaces was gently removed using tissue paper (Elleair, Cat# DSI20703128). The infiltrated plants were maintained under their growth conditions for 24 h (for inhibitor treatments) or 8 h (for *Pst* DC3000 inoculation), followed by humidity treatment. During *Pst* DC3000 inoculation experiments, lights were kept on continuously from inoculation until sampling.

Agrobacterium-mediated transient expression

Agrobacterium-mediated transient expression was performed as previously described[72] with minor modifications. *Agrobacterium tumefaciens* strain GV3101::pMP90 carrying

the relevant expression vectors was cultured overnight at 28°C with shaking (180 rpm) in

2×YT liquid medium [1% (w/v) yeast extract, 1.6% (w/v) tryptone, 0.5% (w/v) NaCl]

supplemented with 50 μg/mL gentamicin and 100 μg/mL spectinomycin. Bacterial cells

were collected by centrifugation at 2,000 g for 10 min at 25°C, washed once with sterile

water, and resuspended in sterile water to an OD_{600} of 0.05. Acetosyringone (100 mM

stock solution) was added to the suspension to a final concentration of 100 μM.

The bacterial suspensions were infiltrated into three leaves per 5-week-old *NahG* plant

using a needleless syringe. Excess solution on the leaf surfaces was gently removed using

tissue paper. The infiltrated plants were maintained under their growth conditions for

approximately 48 h before humidity treatment.

***Pst* DC3000 inoculation assays**

The following *Pst* DC3000 strain were used in this study: WT, $\Delta hrcC$[73], $\Delta 28E$[36],

$\Delta avrPtoB \Delta avrPto$[38], $\Delta avrPtoB$, and $\Delta avrPto$. Strains were cultured overnight at 28°C

with shaking (180 rpm) in King's B liquid medium [2% (w/v) Bacto Proteose Peptone

No. 3 (Gibco, Cat# 211693), 0.15% (w/v) K_2HPO4, 1% (v/v) glycerol, 5 mM $MgSO_4$]

supplemented with 50 μg/mL rifampicin and additional antibiotics as appropriate.

Bacterial cells were collected by centrifugation at 2,000 g for 10 min at 25°C, washed

once with sterile water, and resuspended in sterile water to the OD_{600} values indicated in

figure legends.

Bacterial suspensions were infiltrated into three leaves per plant using a needleless

syringe. Excess solution on the leaf surfaces was gently removed using tissue paper.

Infiltrated plants were maintained under their respective growth conditions for the

indicated time. To ensure high humidity (~95% RH), infiltrated plants were covered with

a plastic container as described in humidity treatment.

For water-soaking assays, the abaxial sides of infiltrated leaves were photographed at 1

day post-infiltration (dpi). The number of leaves exhibiting water-soaking was recorded,

or the water-soaked area relative to total leaf area was quantified using Image J software

(https://imagej.net/ij/). For growth assays, three inoculated leaves were collected from

three different plants at the indicated dpi. Leaves were homogenized in 1 mL sterile water,

and 10 uL aliquots of serial dilutions (10^{-1} to 10^{-8}) were spotted onto NYGA medium

[0.5% (w/v) Bacto Peptone (Gibco, Cat# 211677), 0.3% (w/v) yeast extract (Nacalai

Tesque, Cat# 15838-45), 2% (v/v) glycerol. 1.5% agar (Nacalai Tesque, Cat# 01162-15)]

supplemented with 50 μg/mL rifampicin. Colony-forming units (CFUs) were determined

after 2 days of incubation at 28°C.

RT-qPCR

Total RNA was extracted from three leaves or a single shoot segment using Sepasol-RNA

I Super G (Nacalai Tesque, Cat# 09379-55), followed by DNase treatment and reverse

transcription (RT) with the PrimeScript RT reagent Kit with the gDNA Eraser (Takara,

Cat# RR047B), according to the manufacturer's instructions. Quantitative PCR (qPCR)

was performed using Power SYBR Green PCR Master Mix (Applied Biosystems, Cat#

4368702) on a Thermal Cycler Dice Real Time System III (Takara, Cat# TP950). Gene

expression levels were calculated using the ΔΔCt method, with normalization to *ACTIN2*

as an internal reference. Primers used for qPCR are listed in Supplementary Table 6.

Immunoblotting

Frozen plant tissue (three leaves per sample) was ground into a fine powder and

resuspended in protein extraction buffer [62.5 mM Tris-HCl (pH 6.8), 10% glycerol, 2%

SDS, 5% (v/v) 2-mercaptoethanol, 0.02% bromophenol blue] supplemented with 1 mM

EDTA (pH 8.0) and protease inhibitor cocktail (Roche, Cat# 11873580001). The buffer

was added at a ratio of 3 μL per mg of tissue weight. Extracts were incubated at 95°C for

5 min and subsequently centrifuged at 20,000 *g* for 10 min at 25°C, and the supernatants

were collected.

Proteins were separated on a 10% polyacrylamide gel and transferred to a PVDF

membrane (Millipore, Cat# IPVH00010). The membrane was blocked 2% (w/v) skim

milk (Nacalai Tesque, Cat# 31149-75) in TBS-T [25 mM Tris, 192 mM glycine, 0.1%

(v/v) Tween-20] for 1 h at 25°C with agitation, rinsed once with deionized water, and

once with TBS-T. The blocked membrane was incubated overnight at 4°C with the

primary antibody solution under agitation, rinsed five times with deionized water, and

washed once in TBS-T for 5 min at 25°C. Subsequently, the membrane was incubated

with the secondary antibody solution for 1 h at 25°C with agitation, rinsed five times with

deionized water, and washed once with TBS-T for 5 min at 25°C.

Chemiluminescent detection was performed using Chemi-Lumi One L (Nacalai Tesque,

Cat# 07880) and a FUSION FX imaging system (Vilber). The following antibodies were

used for immunoblotting: anti-FLAG (Sigma-Aldrich, Cat# F1804; 1:5000 dilution), anti-

GFP (MBL, Cat# 598; 1:5000 dilution), anti-mouse IgG HRP-linked (Cell Signaling

Technology, Cat# 7076; 1:5000 dilution), and anti-rabbit IgG HRP-linked (Cell Signaling

Technology, Cat# 7074; 1:5000 dilution). All antibodies were diluted in TBS-T.

Phytohormone quantification

SA, ABA, and JA-Ile were extracted and purified from freeze-dried samples as previously

described[74]. For hormone measurement, isotope-labeled internal standards were added at

the initial extraction procedure. Plant hormones were analyzed by a LC-MS/MS system

(ACQUITY UPLC I-Class PLUS FTN/Xevo TQ-S micro; Waters) equipped with a short

ODS column (TSKgel ODS-120H 2.0 mm I.D. x 5 cm, 1.9 μm; TOSHO) or a long ODS

column (TSKgel ODS-120H 2.0 mm I.D. x 15 cm, 1.9 μm; TOSHO). A binary solvent

system composed of 0.1% acetic acid (v/v) ACN (A) and 0.1% acetic acid (v/v) H_2O (B)

was used. SA, ABA, and JA-Ile were measured using the short ODS column, and

separation was performed at a flow rate of 0.2 mL/min with the following linear gradient

of solvent B: 0–2 min (1%–5%), 2–5 min (5%–15%), 5–25 min (15%–40%), 25.1–27

min (100%), and 27.1–30 min (1%). The retention times for each compound were as

follows: d_6-SA (7.39), SA (7.46), d_6-ABA (12.59), ABA (12.67), $^{13}C_6$-JA-Ile (21.64), and

JA-Ile (21.94). The MS/MS analysis was performed under the following conditions:

capillary (kV) = 1.00, source temperature (°C) = 150, desolvation temperature (°C) = 500,

cone gas flow (L/h) = 50, desolvation gas flow (L/h) = 1000. Ion mode was negative. The

MS/MS transitions (m/z) and collision energies (V) for each compound were as follows:

d_6-SA (142.1/98.0, 18V), SA (137.0/93.4, 18V), d_6-ABA (269.1/159.1, 8V), ABA

(263.1/153.1, 8V), $^{13}C_6$-JA-Ile (328.2/136.1, 20V), and JA-Ile (322.2/130.1, 20V).

Stomatal aperture measurement

Following humidity treatment, detached leaves were immediately immersed in 4% (v/v)

formaldehyde and incubated overnight at 25°C. Images of stomata on the abaxial leaf

surface were captured using a Leica DMI6000 B microscope equipped with 10× objective

lens. Stomatal aperture was quantified using Image J software.

Real-time cytosolic Ca^{2+} imaging

Real-time cytosolic Ca^{2+} imaging of whole plants expressing GCaMP3 was performed as

described previously[75,76].

RNA-sequencing and data analysis

Total RNA was extracted from three leaves and treated with DNase using the NucleoSpin

RNA Plant kit (Macherey-Nagel, Cat# 740949.50) according to the manufacturer's

instruction. RNA quality was assessed using a bioanalyzer (Agilent Technologies).

Each raw read data stored in DDBJ with accession numbers DRR545835 to DRR545846

was generated and analyzed as follows. Each RNA library was prepared from 1000 ng of

total RNA using the NEBNext Ultra II RNA Library Prep Kit for Illumina (New England

Biolabs, Cat# E7770) according to the manufacturer's instruction. PCR for library

amplification was performed for 7 cycles. The average length of the library was 350 bp,

as measured by a bioanalyzer. Concentrations were measured by Kapa Library

Quantification Kit (Kapa Biosystems, Cat# KK4824), each sample was diluted to 10 nM,

and equal volumes were mixed. Single reads of 75 bp were obtained using a NextSeq 500

system (Illumina). CLC genomics workbench version 21 (Qiagen) was used to trim raw

reads and map to the TAIR10 Arabidopsis genome (https://www.arabidopsis.org/). In

trimming, the following parameters were changed (quality limit = 0.001, number of 5′

terminal nucleotides =15, number of 3′ terminal nucleotides = 5, and minimum number

of nucleotides in reads =25) and other options were default values. In mapping, all options

were default values.

The generation and analysis of each raw read data stored in DDBJ with accession numbers

DRR545847 to DRR545870 differs from the above analysis in three points. 1) Each

library was prepared from 500 ng total RNA using the NEBNext Ultra II Directional RNA

Library Prep Kit for Illumina (New England Biolabs, Cat# 7760). PCR for library
amplification was performed for 8 cycles. Each sample was diluted to 4 nM, and equal
volumes were mixed. 2) Single reads of 100 bp were obtained using a NextSeq 1000
system (Illumina). 3) CLC genomics workbench version 24 (Qiagen) was used for reads
trimming and mapping. In trimming and mapping, the following parameters were
changed (minimum number of nucleotides in reads = 40, and strand setting = Revers). All
other methods, numbers, options are the same as above analysis.

The obtained read counts were processed and analyzed using iDEP 2.01
(https://bioinformatics.sdstate.edu/idep/) with default settings. Gene Ontology and *cis*-
regulatory enrichment analyses were performed using ShinyGO 0.81
(https://bioinformatics.sdstate.edu/go/) with default settings. For *cis*-regulatory
enrichment analysis, the AGRIS database was used as the pathway reference.

Accession numbers
The gene sequences analyzed in this study were obtained from The Arabidopsis
Information Resource (https://www.arabidopsis.org/) and The Pseudomonas Genome
Database (https://www.pseudomonas.com/), under the accession numbers listed in
Supplementary Table 7.

Statistical analysis
All data, except RNA-seq, were analyzed using GraphPad Prism 10 software
(https://www.graphpad.com/features). No statistical methods were used to predetermine
sample size, and no data were excluded from analyses. Experiments were not randomized,
and investigators were not blinded to group allocation or outcome assessment. Statistical
comparisons were performed using two-tailed Welch's t test, or one-way or two-way
ANOVA followed by Tukey's multiple comparison test, as specified in the figure legends.

Data availability
The raw RNA sequencing data have been submitted in the DDJB database under
accession numbers DRR545835 to DRR545870.

Acknowledgements

We thank Chika Tateda, Mie Matsubara, Taiga Ishihara, Natsuki Tsuchida, and Temma Takazawa (Nara Institute of Science and Technology) for technical assistance; Yuri Kanno (RIKEN CSRS) for supporting plant hormone analysis; Akira Mine (Kyoto University) for *cyp707a* seeds; Atsushi Takemiya (Yamaguchi University) for *ost2-3D* seeds; Michael F. Thomashow (Michigan State University) for *camta* seeds; Alan Collmer (Cornell University) for *Pst* DC3000 strains. We also thank the NAIST Life Science Collaboration Center (LiSCo) for supporting this study. This work was supported in part by JSPS KAKENHI Grant Numbers 21K14829 (S.Y.), 24H00565 (M.T.) and 21H02507 (Y.S.), JST ACT-X Grant Number JPMJAX22BN (S.Y.), ERATO Grant Number JPMJER2403 (M.T.), Cooperative Research Grant of the Genome Research for BioResource, NODAI Genome Research Center, Tokyo University of Agriculture (S.Y.), and Institute for Fermentation, Osaka Grant Number G-2025-2-089 (S.Y.).

Author contributions

S.Y. and Y.S. conceived the study. S.Y. designed the research, conducted most of the experiments, and analyzed the data. A.R, T.H., and S.U. performed part of the RT-qPCR and *Pst* DC3000 inoculation assay. Y.W., Y.T., and M.O. performed phytohormone quantification. H.I., R.Suzuki, and M.T performed GCaMP3 imaging. A.S., R.Sk, and I.Y. performed RNA-sequencing. S.Y. and Y.S. wrote the manuscript with input from the other authors.

Competing interests

The authors declare no competing interests.

References

1   Xin, X. F., Kvitko, B. & He, S. Y. Pseudomonas syringae: what it takes to be a pathogen. *Nat. Rev. Microbiol.* **16**, 316-328 (2018).

2   Boutrot, F. & Zipfel, C. Function, Discovery, and Exploitation of Plant Pattern Recognition Receptors for Broad-Spectrum Disease Resistance. *Annu. Rev. Phytopathol.* **55**, 257-286 (2017).

3   DeFalco, T. A. & Zipfel, C. Molecular mechanisms of early plant pattern-

triggered immune signaling. *Mol. Cell* **81**, 3449-3467 (2021).

4   Saijo, Y., Loo, E. P. & Yasuda, S. Pattern recognition receptors and signaling in
plant-microbe interactions. *Plant J.* **93**, 592-613 (2018).

5   Wei, H. L. *et al.* Pseudomonas syringae pv. tomato DC3000 Type III Secretion
Effector Polymutants Reveal an Interplay between HopAD1 and AvrPtoB. *Cell*
Host Microbe **17** (2015).

6   Roussin-Léveillée, C., Mackey, D., Ekanayake, G., Gohmann, R. & Moffett, P.
Extracellular niche establishment by plant pathogens. *Nat. Rev. Microbiol.* **22**,
360-372 (2024).

7   Panchal, S. *et al.* Regulation of Stomatal Defense by Air Relative Humidity.
Plant Physiol. **172**, 2021-2032 (2016).

8   Xin, X. F. *et al.* Bacteria establish an aqueous living space in plants crucial for
virulence. *Nature* **539**, 524-529 (2016).

9   Hu, Y. *et al.* Bacterial effectors manipulate plant abscisic acid signaling for
creation of an aqueous apoplast. *Cell Host Microbe* **30**, 518-529 (2022).

10   Roussin-Léveillée, C. *et al.* Evolutionarily conserved bacterial effectors hijack
abscisic acid signaling to induce an aqueous environment in the apoplast. *Cell*
Host Microbe **30**, 489-501 (2022).

11   Waadt, R. *et al.* Plant hormone regulation of abiotic stress responses. *Nat. Rev.*
Mol. Cell Biol. **23**, 680-694 (2022).

12   Zhang, D., Tian, C., Yin, K., Wang, W. & Qiu, J. L. Postinvasive Bacterial
Resistance Conferred by Open Stomata in Rice. *Mol. Plant Microbe Interact.* **32**,
255-266 (2019).

13   Peng, Z. *et al.* Xanthomonas translucens commandeers the host rate-limiting
step in ABA biosynthesis for disease susceptibility. *Proc. Natl Acad. Sci. USA*
116, 20938-20946 (2019).

14   You, Y. *et al.* The eINTACT system dissects bacterial exploitation of plant
osmosignalling to enhance virulence. *Nat. Plants* **9**, 128-141 (2023).

15   Wu, J. *et al.* CURLY LEAF modulates apoplast liquid water status in
Arabidopsis leaves. *Plant Physiol.* **193**, 792-808 (2023).

16   Lajeunesse, G. *et al.* Light prevents pathogen-induced aqueous
microenvironments via potentiation of salicylic acid signaling. *Nat. Commun.*
14, 713 (2023).

17   Okamoto, M. *et al.* High humidity induces abscisic acid 8'-hydroxylase in
stomata and vasculature to regulate local and systemic abscisic acid responses in
Arabidopsis. *Plant Physiol.* **149**, 825-834 (2009).

18   Kim, H., Yu, S. I., Jung, S. H., Lee, B. H. & Suh, M. C. The F-Box Protein
SAGL1 and ECERIFERUM3 Regulate Cuticular Wax Biosynthesis in Response
to Changes in Humidity in Arabidopsis. *Plant Cell* **31**, 2223-2240 (2019).
19   Jiang, Z. *et al.* Ethylene signaling modulates air humidity responses in plants.
Plant J. **117**, 653-668 (2024).
20   Nambara, E. & Marion-Poll, A. Abscisic acid biosynthesis and catabolism.
Annu. Rev. Plant Biol. **56**, 165-185 (2005).
21   Yao, L. *et al.* High air humidity dampens salicylic acid pathway and NPR1
function to promote plant disease. *EMBO J.* **42**, e113499 (2023).
22   Yamauchi, S. *et al.* The Plasma Membrane H^+-ATPase AHA1 Plays a Major
Role in Stomatal Opening in Response to Blue Light. *Plant Physiol.* **171**, 2731-
2743 (2016).
23   Yasuda, M. *et al.* Antagonistic interaction between systemic acquired resistance
and the abscisic acid-mediated abiotic stress response in Arabidopsis. *Plant Cell*
20, 1678-1692 (2008).
24   de Torres Zabala, M., Bennett, M. H., Truman, W. H. & Grant, M. R.
Antagonism between salicylic and abscisic acid reflects early host-pathogen
conflict and moulds plant defence responses. *Plant J.* **59**, 375-386 (2009).
25   Melotto, M., Zhang, L., Oblessuc, P. R. & He, S. Y. Stomatal Defense a Decade
Later. *Plant Physiol.* **174**, 561-571 (2017).
26   Rekhter, D. *et al.* Isochorismate-derived biosynthesis of the plant stress hormone
salicylic acid. *Science* **365**, 498-502 (2019).
27   Seo, M. *et al.* The Arabidopsis aldehyde oxidase 3 (AAO3) gene product
catalyzes the final step in abscisic acid biosynthesis in leaves. *Proc. Natl Acad.*
Sci. USA **97**, 12908-12913 (2000).
28   Yuan, M. *et al.* Pattern-recognition receptors are required for NLR-mediated
plant immunity. *Nature* **592**, 105-109 (2021).
29   Wang, Y. *et al*. A calmodulin-binding transcription factor links calcium signaling
to antiviral RNAi defense in plants. *Cell Host Microbe.* **29**, 1393-1406 (2021).
30   DeFalco, T. A., Moeder, W. & Yoshioka, K. Opening the Gates: Insights into
Cyclic Nucleotide-Gated Channel-Mediated Signaling. *Trends Plant Sci.* **21**,
903-906 (2016).
31   Kim, Y., Park, S., Gilmour, S. J. & Thomashow, M. F. Roles of CAMTA
transcription factors and salicylic acid in configuring the low-temperature
transcriptome and freezing tolerance of Arabidopsis. *Plant J.* **75**, 364-376
(2013).

902 32 Kidokoro, S. *et al.* Different Cold-Signaling Pathways Function in the
903   Responses to Rapid and Gradual Decreases in Temperature. *Plant Cell* **29**, 760-
904   774 (2017).

905 33 Darwish, E. *et al.* Touch signaling and thigmomorphogenesis are regulated by
906   complementary CAMTA3- and JA-dependent pathways. *Sci. Adv.* **8**, eabm2091
907   (2022).

908 34 Matsumura, M. *et al.* Mechanosensory trichome cells evoke a mechanical
909   stimuli-induced immune response in Arabidopsis thaliana. *Nat. Commun.* **13**,
910   1216 (2022).

911 35 Kim, Y. S. *et al.* CAMTA-Mediated Regulation of Salicylic Acid Immunity
912   Pathway Genes in Arabidopsis Exposed to Low Temperature and Pathogen
913   Infection. *Plant Cell* **29**, 2465-2477 (2017).

914 36 Cunnac, S. *et al.* Genetic disassembly and combinatorial reassembly identify a
915   minimal functional repertoire of type III effectors in Pseudomonas syringae.
916   *Proc. Natl Acad. Sci. USA* **108**, 2975-2980 (2011).

917 37 de Torres-Zabala, M. *et al.* Pseudomonas syringae pv. tomato hijacks the
918   Arabidopsis abscisic acid signalling pathway to cause disease. *EMBO J.* **26**,
919   1434-1443 (2007).

920 38 Lin, N. C. & Martin, G. B. An avrPto/avrPtoB mutant of Pseudomonas syringae
921   pv. tomato DC3000 does not elicit Pto-mediated resistance and is less virulent
922   on tomato. *Mol. Plant Microbe Interact.* **18**, 43-51 (2005).

923 39 Janjusevic, R., Abramovitch, R. B., Martin, G. B. & Stebbins, C. E. A bacterial
924   inhibitor of host programmed cell death defenses is an E3 ubiquitin ligase.
925   *Science* **311**, 222-226 (2006).

926 40 Mwimba, M. *et al.* Daily humidity oscillation regulates the circadian clock to
927   influence Plant Physiol.. *Nat. Commun.* **9**, 4290 (2018).

928 41 Rowe, J. *et al.* Next-generation ABACUS biosensors reveal cellular ABA
929   dynamics driving root growth at low aerial humidity. *Nat. Plants* **9**, 1103-1115
930   (2023).

931 42 Tulva, I., Koolmeister, K. & Hõrak, H. Low relative air humidity and increased
932   stomatal density independently hamper growth in young Arabidopsis. *Plant J.*
933   **119**, 2718-2736 (2024).

934 43 Romero, F. *et al.* Humidity and high temperature are important for predicting
935   fungal disease outbreaks worldwide. *New Phytol.* **234**, 1553-1556 (2022).

936 44 L'Haridon, F. *et al.* A permeable cuticle is associated with the release of reactive
937   oxygen species and induction of innate immunity. *PLoS Pathog.* **7**, e1002148

938  (2011).

939 45 Cui, F. *et al.* Dissecting Abscisic Acid Signaling Pathways Involved in Cuticle
940  Formation. *Mol. Plant* **9**, 926-938 (2016).

941 46 Hsu, F. C. *et al.* Submergence confers immunity mediated by the WRKY22
942  transcription factor in Arabidopsis. *Plant Cell* **25**, 2699-2713 (2013).

943 47 Ding, Y., Dommel, M. & Mou, Z. Abscisic acid promotes proteasome-mediated
944  degradation of the transcription coactivator NPR1 in Arabidopsis thaliana. *Plant*
945  *J.* **86**, 20-34 (2016).

946 48 Berens, M. L. *et al.* Balancing trade-offs between biotic and abiotic stress
947  responses through leaf age-dependent variation in stress hormone cross-talk.
948  *Proc. Natl Acad. Sci. USA* **116**, 2364-2373 (2019).

949 49 Umezawa, T. *et al.* CYP707A3, a major ABA 8'-hydroxylase involved in
950  dehydration and rehydration response in Arabidopsis thaliana. *Plant J.* **46**, 171-
951  182 (2006).

952 50 Zhang, Y. *et al.* Control of salicylic acid synthesis and systemic acquired
953  resistance by two members of a plant-specific family of transcription factors.
954  *Proc. Natl Acad. Sci. USA* **107**, 18220-18225 (2010).

955 51 Wang, L. *et al.* CBP60g and SARD1 play partially redundant critical roles in
956  salicylic acid signaling. *Plant J.* **67**, 1029-1041 (2011).

957 52 Zhang, B. *et al.* The mitochondrial outer membrane AAA ATPase AtOM66
958  affects cell death and pathogen resistance in Arabidopsis thaliana. *Plant J.* **80**,
959  709-727 (2014).

960 53 Kemppinen, J., Pollmeier, M., Ehonen, S., Brosché, M. & Sierla, M. Water
961  immunity overrides stomatal immunity in plant resistance to *Pseudomonas*
962  *syringae*. *Plant Physiol.* **198**, kiaf127 (2025).

963 54 Liu, Y., Mahmud, M. R., Xu, N. & Liu, J. The Pseudomonas syringae effector
964  AvrPtoB targets abscisic acid signaling pathway to promote its virulence in
965  Arabidopsis. *Phytopathol. Res.* **4**, 5 (2022).

966 55 Shan, L. *et al.* Bacterial effectors target the common signaling partner BAK1 to
967  disrupt multiple MAMP receptor-signaling complexes and impede plant
968  immunity. *Cell Host Microbe* **4**, 17-27 (2008).

969 56 Göhre, V. *et al.* Plant pattern-recognition receptor FLS2 is directed for
970  degradation by the bacterial ubiquitin ligase AvrPtoB. *Curr. Biol.* **18**, 1824-1832
971  (2008).

972 57 Gimenez-Ibanez, S. *et al.* AvrPtoB targets the LysM receptor kinase CERK1 to
973  promote bacterial virulence on plants. *Curr. Biol.* **19**, 423-429 (2009).

58   Chen, H. *et al.* A Bacterial Type III Effector Targets the Master Regulator of
Salicylic Acid Signaling, NPR1, to Subvert Plant Immunity. *Cell Host Microbe*
22, 777-788 (2017).

59   Wang, W., Liu, N., Gao, C., Rui, L. & Tang, D. The Pseudomonas Syringae
Effector AvrPtoB Associates With and Ubiquitinates Arabidopsis Exocyst
Subunit EXO70B1. *Front. Plant Sci.* **10**, 1027 (2019).

60   Wang, M. Y. *et al.* The plant immune receptor SNC1 monitors helper NLRs
targeted by a bacterial effector. *Cell Host Microbe* **31**, 1792-1803 (2023).

61   Hussain, S. *et al.* Calcium signaling triggers early high humidity responses in
Arabidopsis thaliana. *Proc. Natl Acad. Sci. USA* **121**, e2416270121 (2024).

62   Lee, S. *et al.* Oomycete effector AVRblb2 targets cyclic nucleotide-gated
channels through calcium sensors to suppress pattern-triggered immunity. *New
Phytol.* **241**, 1277-1291 (2024).

63   Jing, B. *et al.* Brush and spray: a high-throughput systemic acquired resistance
assay suitable for large-scale genetic screening. *Plant Physiol.* **157**, 973-980
(2011).

64   Nie, H. *et al.* SR1, a calmodulin-binding transcription factor, modulates plant
defense and ethylene-induced senescence by directly regulating NDR1 and
EIN3. *Plant Physiol.* **158**, 1847-1859 (2012).

65   Jacob, F. *et al.* A dominant-interfering camta3 mutation compromises primary
transcriptional outputs mediated by both cell surface and intracellular immune
receptors in Arabidopsis thaliana. *New Phytol.* **217**, 1667-1680 (2018).

66   Lolle, S. *et al.* Matching NLR Immune Receptors to Autoimmunity in camta3
Mutants Using Antimorphic NLR Alleles. *Cell Host Microbe* **21**, 518-529
(2017).

67   Goel, A. K. *et al.* The Pseudomonas syringae type III effector HopAM1
enhances virulence on water-stressed plants. *Mol. Plant Microbe Interact.* **21**,
361-370 (2008).

68   Nakagawa, T. *et al.* Improved Gateway binary vectors: high-performance
vectors for creation of fusion constructs in transgenic analysis of plants. *Biosci.
Biotechnol. Biochem.* **71**, 2095-2100 (2007).

69   Yamada, K. *et al.* Danger peptide receptor signaling in plants ensures basal
immunity upon pathogen-induced depletion of BAK1. *EMBO J.* **35**, 46-61
(2016).

70   Zhang, X., Henriques, R., Lin, S. S., Niu, Q. W. & Chua, N. H. Agrobacterium-
mediated transformation of Arabidopsis thaliana using the floral dip method.

Nat. Protoc. **1**, 641-646 (2006).

71   Matsumoto, A., Schlüter, T., Melkonian, K., Takeda, A., Nakagami, H., Mine, A.
A versatile Tn7 transposon-based bioluminescence tagging tool for quantitative
and spatial detection of bacteria in plants. *Plant Commun.* **3**, 100227 (2021).

72   Rosas-Díaz, T. *et al.* Arabidopsis NahG Plants as a Suitable and Efficient System
for Transient Expression using Agrobacterium tumefaciens. *Mol. Plant* **10**, 353-
356 (2017).

73   Wei, W. *et al.* The gene coding for the Hrp pilus structural protein is required for
type III secretion of Hrp and Avr proteins in Pseudomonas syringae pv. tomato.
Proc. Natl Acad. Sci. USA **97**, 2247-2252 (2000).

74   Sato, Y. *et al.* Temporal dynamics of N-hydroxypipecolic acid and salicylic acid
pathways in the disease response to powdery mildew in wheat. *Biochem.*
Biophys. Res. Commun. **734**, 150624 (2024).

75   Toyota, M. *et al.* Glutamate triggers long-distance, calcium-based plant defense
signaling. *Science* **361**, 1112-1115 (2018).

76   Aratani, Y., Uemura, T., Hagihara, T., Matsui, K. & Toyota, M. Green leaf
volatile sensory calcium transduction in Arabidopsis. *Nat. Commun.* **14**, 6236
(2023).

77   Okamoto, M. *et al.* CYP707A1 and CYP707A2, which encode abscisic acid 8'-
hydroxylases, are indispensable for proper control of seed dormancy and
germination in Arabidopsis. *Plant Physiol.* **141**, 97-107 (2006).

78   Kushiro, T. *et al.* The Arabidopsis cytochrome P450 CYP707A encodes ABA 8'-
hydroxylases: key enzymes in ABA catabolism. *EMBO J.* **23**, 1647-1656 (2004).

79   Wildermuth, M. C., Dewdney, J., Wu, G. & Ausubel, F. M. Isochorismate
synthase is required to synthesize salicylic acid for plant defence. *Nature* **414**,
562-565 (2001).

80   Seo, M. *et al.* Comparative studies on the Arabidopsis aldehyde oxidase (AAO)
gene family revealed a major role of AAO3 in ABA biosynthesis in seeds. *Plant*
Cell Physiol. **45**, 1694-1703 (2004).

81   Schwessinger, B. *et al.* Phosphorylation-dependent differential regulation of
plant growth, cell death, and innate immunity by the regulatory receptor-like
kinase BAK1. *PLoS Genet.* **7**, e1002046 (2011).

82   He, K. *et al.* BAK1 and BKK1 regulate brassinosteroid-dependent growth and
brassinosteroid-independent cell-death pathways. *Curr. Biol.* **17**, 1109-1115
(2007).

83   Miya, A. *et al.* CERK1, a LysM receptor kinase, is essential for chitin elicitor

| 1046 | | signaling in Arabidopsis. *Proc. Natl Acad. Sci. USA* **104**, 19613-19618 (2007). |

signaling in Arabidopsis. *Proc. Natl Acad. Sci. USA* **104**, 19613-19618 (2007).

84   Yu, X. *et al.* The Receptor Kinases BAK1/SERK4 Regulate Ca^{2+} Channel-
Mediated Cellular Homeostasis for Cell Death Containment. *Curr. Biol.* **29**,
3778-3790 (2019).

85   Clough, S. J. *et al.* The Arabidopsis dnd1 "defense, no death" gene encodes a
mutated cyclic nucleotide-gated ion channel. *Proc. Natl Acad. Sci. USA* **97**,
9323-9328 (2000).

86   Gobert, A., Park, G., Amtmann, A., Sanders, D. & Maathuis, F. J. Arabidopsis
thaliana cyclic nucleotide gated channel 3 forms a non-selective ion transporter
involved in germination and cation transport. *J. Exp. Bot.* **57**, 791-800 (2006).

87   Jurkowski, G. I. *et al.* Arabidopsis DND2, a second cyclic nucleotide-gated ion
channel gene for which mutation causes the "defense, no death" phenotype. *Mol.*
Plant Microbe Interact. **17**, 511-520 (2004).

88   Tan, Y. Q. *et al.* Three CNGC Family Members, CNGC5, CNGC6, and CNGC9,
Are Required for Constitutive Growth of Arabidopsis Root Hairs as Ca(2+)-
Permeable Channels. *Plant Commun.* **1**, 100001 (2020).

89   Tunc-Ozdemir, M. *et al.* Cyclic nucleotide gated channels 7 and 8 are essential
for male reproductive fertility. *PLoS One* **8**, e55277 (2013).

90   Yoshioka, K. *et al.* The chimeric Arabidopsis CYCLIC NUCLEOTIDE-GATED
ION CHANNEL11/12 activates multiple pathogen resistance responses. *Plant*
Cell **18**, 747-763 (2006).

91   Shih, H. W., DePew, C. L., Miller, N. D. & Monshausen, G. B. The Cyclic
Nucleotide-Gated Channel CNGC14 Regulates Root Gravitropism in
Arabidopsis thaliana. *Curr. Biol.* **25**, 3119-3125 (2015).

92   Moon, J. Y., Belloeil, C., Ianna, M. L. & Shin, R. Arabidopsis CNGC Family
Members Contribute to Heavy Metal Ion Uptake in Plants. *Int. J. Mol. Sci.* **20**,
413 (2019).

93   Tunc-Ozdemir, M. *et al.* A cyclic nucleotide-gated channel (CNGC16) in pollen
is critical for stress tolerance in pollen reproductive development. *Plant Physiol.*
161, 1010-1020 (2013).

94   Ladwig, F. *et al.* Phytosulfokine Regulates Growth in Arabidopsis through a
Response Module at the Plasma Membrane That Includes CYCLIC
NUCLEOTIDE-GATED CHANNEL17, H^+-ATPase, and BAK1. *Plant Cell* **27**,
1718-1729 (2015).

95   Gao, Q. F. *et al.* Cyclic nucleotide-gated channel 18 is an essential Ca^{2+} channel
in pollen tube tips for pollen tube guidance to ovules in Arabidopsis. *Proc. Natl*

Acad. Sci. USA **113**, 3096-3101 (2016).

96   Kugler, A., Köhler, B., Palme, K., Wolff, P. & Dietrich, P. Salt-dependent

regulation of a CNG channel subfamily in Arabidopsis. *BMC Plant Biol.* **9**, 140

(2009).

97   Lawton, K. *et al.* Systemic acquired resistance in Arabidopsis requires salicylic

acid but not ethylene. *Mol. Plant Microbe Interact.* **8**, 863-870 (1995).

98   Doherty, C. J., Van Buskirk, H. A., Myers, S. J. & Thomashow, M. F. Roles for

Arabidopsis CAMTA transcription factors in cold-regulated gene expression and

freezing tolerance. *Plant Cell* **21**, 972-984 (2009).

99   Gu, Y. & Innes, R. W. The KEEP ON GOING protein of Arabidopsis recruits the

ENHANCED DISEASE RESISTANCE1 protein to trans-Golgi network/early

endosome vesicles. *Plant Physiol.* **155**, 1827-1838 (2011).

1094

Fig. 1: *CYP707A3* is required for resistance to bacterial water-soaking under high

humidity.

a, Expression levels of *CYP707A* genes in humidity-treated leaves. Wild-type (WT) plants

were exposed to moderate humidity (MH) or high humidity (HH) for the indicated time.

Bars represent means ($n = 3$). **b**, ABA levels in humidity-treated leaves. The indicated

plants were exposed to MH or HH for 2 h. Bars represent means ($n = 3$). **c, d**, Water-

soaking in *Pst* Δ*hrcC*-inoculated leaves. The indicated plants were infiltrated with *Pst*

Δ*hrcC* ($OD_{600} = 0.02$) and maintained under MH or HH for 1 day. The number of water-

soaked leaves was pooled from two independent experiments. **e**, Bacterial growth in *Pst*

Δ*hrcC*-inoculated leaves. The indicated plants were infiltrated with *Pst* Δ*hrcC* ($OD_{600} =$

0.02) and maintained under MH or HH for 3 days. Bars represent means ($n = 6$). Different

letters indicate statistically significant differences ($P < 0.05$; two-way ANOVA followed

by Tukey's test). dpi, days post-infiltration. Experiments in (**a**) and (**e**) were repeated twice

with similar results.

1109

1110   **Fig. 2: *CYP707A3*-mediated water-soaking resistance depends on stomatal opening.**

1111   **a**, **b**, Representative images of stomata (**a**) and stomatal aperture (**b**) in humidity-treated

1112   leaves. The indicated plants were exposed to moderate humidity (MH) or high humidity

1113   (HH) for 1 h. Bars represent the median ($n = 149-264$). **c**, Water-soaking in *Pst* Δ*hrcC*-

1114   inoculated leaves. The indicated plants were infiltrated with *Pst* Δ*hrcC* ($OD_{600} = 0.02$)

and maintained under HH for 1 day. The number of water-soaked leaves was pooled from two independent experiments. **d**, Bacterial growth in *Pst ΔhrcC*-inoculated leaves. The indicated plants were infiltrated with *Pst ΔhrcC* ($OD_{600} = 0.02$) and maintained under HH for 3 days. Bars represent means ($n = 6$). Different letters indicate statistically significant differences ($P < 0.05$; two-way ANOVA followed by Tukey's test in **b**; one-way ANOVA followed by Tukey's test in **d**). WT, wild-type; dpi, days post-infiltration. Experiments in (**b**) and (**d**) were repeated twice with similar results.

Fig. 3: CAMTA binding motifs are overrepresented in early-phase high humidity-induced genes.

a-c, Transcriptome profiling of humidity-treated leaves by RNA-seq. Wild-type plants were exposed to moderate humidity (MH) or high humidity (HH) for the indicated time. **a**, Venn diagrams showing differentially expressed genes (DEGs) (fold change > 2; FDR < 0.05) at 0.5 h and 5 h after treatment. **b**, Principal component analysis (PCA) of transcriptomic data. **c**, K-means clustering of the 1000 most variable genes. **d**, Top 10 enriched *cis*-regulatory elements among high humidity-induced genes identified in the clusters shown in (**c**).

Fig. 4: CAMTA3 drives *CYP707A3* expression to promote resistance to bacterial water-soaking under high humidity.

a, Schematic diagram of the promoter regions of *CYP707A3* and *CYP707A1*, highlighting the CGCG box. **b**, *GUS* expression driven by the *CYP707A3* promoter, with or without CGCG boxes, in humidity-treated leaves. The indicated constructs were transiently expressed in *NahG* leaves via *Agrobacterium* infiltration. Infiltrated plants were maintained under moderate humidity (MH) for 24 h, then exposed to MH or high humidity (HH) for 0.5 h. Bars represent means ($n = 3$). **c, d**, Expression levels of *CYP707A3* and *CYP707A1* in humidity-treated leaves of *cngc* (**c**) and *camta3* (**d**) mutants. The indicated plants were exposed to MH or HH for 0.5 h. Bars represent means ($n = 3$). **e**, ABA levels in humidity-treated leaves. The indicated plants were exposed to MH or HH for 2 h. Bars represent means ($n = 5$). **f**, Stomatal aperture in humidity-treated leaves. The indicated plants were exposed to MH or HH for 1 h. Bars represent the median ($n = 220$-247). **g**, Water-soaking in *Pst ΔhrcC*-inoculated leaves. The indicated plants were

infiltrated with *Pst* Δ*hrcC* (OD$_{600}$ = 0.02) and maintained under HH for 1 day. The number

of water-soaked leaves was pooled from two independent experiments. **h**, **i**, Bacterial

growth in *Pst* Δ*hrcC*-inoculated leaves. The indicated plants were infiltrated with *Pst*

Δ*hrcC* (OD$_{600}$ = 0.02) and maintained under HH (**h**) or MH (**i**) for 3 days. Bars represent

means (*n* = 6). Different letters indicate statistically significant differences (*P* < 0.05; two-

way ANOVA followed by Tukey's test in **b-f**; one-way ANOVA followed by Tukey's test

in **h** and **i**). WT, wild-type; CAMTA3-A855V, a partial loss-of-function variant; dpi, days

post-infiltration. Experiments in (**b-d**), (**f**), (**h**), and (**i**) were repeated twice with similar

results.

1156

Fig. 5: *Pst* DC3000 blocks high humidity-driven plant transcriptome

reprogramming via T3S effectors.

a, Schematic overview of the experimental design for analyzing high humidity-

responsive gene expression in *Pst* DC3000-inoculated leaves. **b**, Bacterial titers in wild-

type (WT) leaves infiltrated with *Pst* DC3000 or *Pst* Δ*hrcC* (OD$_{600}$ = 0.02) at 8 hours

post-infiltration (hpi) under moderate humidity (MH). Bars represent means (*n* = 6). **c**,

Expression levels of *CYP707A3*, *CYP707A1*, and *NCED3* in *Pst* DC3000-inoculated

leaves following humidity treatment. WT plants were infiltrated with the indicated *Pst*

DC3000 strains (OD$_{600}$ = 0.02), maintained under MH for 8 h, and then exposed to MH

or high humidity (HH) for 0.5 h. Bars represent means (*n* = 3). **d**, **e**, Transcriptome

profiling of *Pst* DC3000-inoculated leaves following humidity treatment as described in

(**a**) and (**c**). **d**, Principal component analysis (PCA) of transcriptomic data. **e**, K-means-

clustering of the 2000 most variable genes. Different letters indicate statistically

significant differences (*P* < 0.05; Welch's *t*-test in **b**; two-way ANOVA followed by

Tukey's test in **c**). Experiments in (**b**) and (**c**) were repeated twice with similar results.

1172

Fig. 6: T3S effector AvrPtoB suppresses *CYP707A3*-mediated water-soaking

resistance.

a, Expression levels of *CYP707A3* in *Pst* DC3000-inoculated leaves following humidity

treatment. Wild-type (WT) plants were infiltrated with the indicated *Pst* DC3000 strains

(OD$_{600}$ = 0.02), maintained under moderate humidity (MH) for 8 h, and then exposed to

MH or high humidity (HH) for 0.5 h. Bars represent means (*n* = 3). The box highlights

the effectors retained in *Pst* Δ20E but absent in *Pst* Δ28E. **b-d**, Expression levels of

CYP707A3 in humidity-treated leaves with or without induction of *AvrPtoB* (**b**), *AvrPto*

(**c**), or *AvrPtoB-F479A* (**d**). The indicated plants were infiltrated with 0.1% ethanol

(−DEX) or 10 μM DEX (+DEX), maintained under moderate humidity (MH) for 24 h,

then exposed to MH or HH for 0.5 h. Bars represent means ($n = 3$). **e**, ABA levels in

humidity-treated leaves with or without *AvrPtoB* induction. The indicated plants were

treated with DEX as described in (**b-d**), and then exposed to MH or HH for 2 h. Bars

represent means ($n = 3$). **f**, Stomatal aperture in humidity-treated leaves with or without

AvrPtoB induction. The indicated were treated with DEX as described in (**b-d**), and then

exposed to MH or HH for 1 h. Bars represent the median ($n = 253\text{-}367$). **g**, Water-soaking

in *Pst* Δ*hrcC*-inoculated leaves with or without *AvrPtoB* induction. The indicated plants

were infiltrated with *Pst* Δ*hrcC* ($OD_{600} = 0.02$) supplemented with −DEX or +DEX and

maintained under HH for 1 day. The number of water-soaked leaves was pooled from two

independent experiments. Different letters indicate statistically significant differences (*P*

< 0.05; two-way ANOVA followed by Tukey's test). AvrPtoB-F479A, E3 ligase deficient

variant; ND, not detected. Experiments in (**a-d**) and (**f**) were repeated twice with similar

results.

1196

Fig. 7: *Pst* DC3000 employs AvrPtoB to suppress *CYP707A3* expression and promote

water-soaking under high humidity.

a, Schematic overview of the experimental design for analyzing gene expression, ABA

accumulation, water-soaking, and bacterial growth in *Pst* DC3000-inoculated leaves. **b**,

Expression levels of *CYP707A3*, *CYP707A1*, and *NCED3* in *Pst* DC3000-inoculated

leaves at 6 hours post-infiltration (hpi) under HH. Wild-type (WT) plants were infiltrated

with the indicated *Pst* DC3000 strains ($OD_{600} = 0.2$), maintained under moderate humidity

(MH) for 2 h, and then transferred to HH for 4 h. Bars represent means ($n = 3$). **c**, ABA

levels in *Pst* DC3000-inoculated leaves. WT plants were inoculated with the indicated

Pst DC3000 strains as described in (**b**). Bars represent means ($n = 6$). **d**, **e**, Representative

images (**d**) and quantification (**e**) of water-soaking in *Pst* DC3000-inoculated leaves. WT

and *cyp707a1 cyp707a3* mutant plants were infiltrated with the indicated *Pst* DC3000

strains ($OD_{600} = 0.2$), maintained under MH for 2 h, and then transferred to HH for 20-22

h. Bars represent the median ($n = 36$; pooled from two independent experiments). **f**, **g**,

Disease symptoms (**f**) and bacterial growth (**g**) in *Pst* DC3000-inoculated leaves. WT and *cyp707a1 cyp707a3* mutant plants were infiltrated with the indicated *Pst* DC3000 strains (OD_{600} = 0.0002), maintained under MH for 2 h, and then transferred to HH for 3 days. Bars represent means (n = 6). Different letters indicate statistically significant differences (P < 0.05; one-way ANOVA followed by Tukey's test in **b** and **c**; two-way ANOVA followed by Tukey's test in **e** and **g**). dpi, days post-infiltration. Experiments in (**b**), (**f**), and (**g**) were repeated twice with similar results. **g**, Working model of *CYP707A3*-mediated water-soaking resistance and its suppression by *Pst* DC3000 via T3S effector AvrPtoB under high humidity.

Fig. 1: *CYP707A3* is required for resistance to bacterial water-soaking under high humidity.
a, Expression levels of *CYP707A* genes in humidity-treated leaves. Wild-type (WT) plants were exposed to moderate humidity (MH) or high humidity (HH) for the indicated time. Bars represent means (*n* = 3). **b,** ABA levels in humidity-treated leaves. The indicated plants were exposed to MH or HH for 2 h. Bars represent means (*n* = 3). **c, d,** Water-soaking in *Pst ΔhrcC*-inoculated leaves. The indicated plants were infiltrated with *Pst ΔhrcC* (OD₆₀₀ = 0.02) and maintained under MH or HH for 1 day. The number of water-soaked leaves was pooled from two independent experiments. **e,** Bacterial growth in *Pst ΔhrcC*-inoculated leaves. The indicated plants were infiltrated with *Pst ΔhrcC* (OD₆₀₀ = 0.02) and maintained under MH or HH for 3 days. Bars represent means (*n* = 6). Different letters indicate statistically significant differences (*P* < 0.05; two-way ANOVA followed by Tukey's test). dpi, days post-infiltration. Experiments in (**a**) and (**e**) were repeated twice with similar results.

Fig. 2: *CYP707A3*-mediated water-soaking resistance depends on stomatal opening.
a, **b**, Representative images of stomata (**a**) and stomatal aperture (**b**) in humidity-treated leaves. The indicated plants were exposed to moderate humidity (MH) or high humidity (HH) for 1 h. Bars represent the median (n = 149-264). **c**, Water-soaking in *Pst ΔhrcC*-inoculated leaves. The indicated plants were infiltrated with *Pst ΔhrcC* (OD_{600} = 0.02) and maintained under HH for 1 day. The number of water-soaked leaves was pooled from two independent experiments. **d**, Bacterial growth in *Pst ΔhrcC*-inoculated leaves. The indicated plants were infiltrated with *Pst ΔhrcC* (OD_{600} = 0.02) and maintained under HH for 3 days. Bars represent means (n = 6). Different letters indicate statistically significant differences (P < 0.05; two-way ANOVA followed by Tukey's test in **b**; one-way ANOVA followed by Tukey's test in **d**). WT, wild-type; dpi, days post-infiltration. Experiments in (**b**) and (**d**) were repeated twice with similar results.

Fig. 3: CAMTA binding motifs are overrepresented in early-phase high humidity-induced genes.
a-c, Transcriptome profiling of humidity-treated leaves by RNA-seq. Wild-type plants were exposed to moderate humidity (MH) or high humidity (HH) for the indicated time. **a**, Venn diagrams showing differentially expressed genes (DEGs) (fold change > 2; FDR < 0.05) at 0.5 h and 5 h after treatment. **b**, Principal component analysis (PCA) of transcriptomic data. **c**, K-means clustering of the 1000 most variable genes. **d**, Top 10 enriched *cis*-regulatory elements among high humidity-induced genes identified in the clusters shown in (**c**).

Fig. 4: CAMTA3 drives *CYP707A3* expression to promote resistance to bacterial water-soaking under high humidity.

a, Schematic diagram of the promoter regions of *CYP707A3* and *CYP707A1*, highlighting the CGCG box. **b**, *GUS* expression driven by the *CYP707A3* promoter, with or without CGCG boxes, in humidity-treated leaves. The indicated constructs were transiently expressed in *NahG* leaves via *Agrobacterium* infiltration. Infiltrated plants were maintained under moderate humidity (MH) for 24 h, then exposed to MH or high humidity (HH) for 0.5 h. Bars represent means (n = 3). **c, d**, Expression levels of *CYP707A3* and *CYP707A1* in humidity-treated leaves of *cngc* (**c**) and *camta3* (**d**) mutants. The indicated plants were exposed to MH or HH for 0.5 h. Bars represent means (n = 3). **e**, ABA levels in humidity-treated leaves. The indicated plants were exposed to MH or HH for 2 h. Bars represent means (n = 5). **f**, Stomatal aperture in humidity-treated leaves. The indicated plants were exposed to MH or HH for 1 h. Bars represent the median (n = 220-247). **g**, Water-soaking in *Pst ΔhrcC*-inoculated leaves. The indicated plants were infiltrated with *Pst ΔhrcC* (OD_{600} = 0.02) and maintained under HH for 1 day. The number of water-soaked leaves was pooled from two independent experiments. **h, i**, Bacterial growth in *Pst ΔhrcC*-inoculated leaves. The indicated plants were infiltrated with *Pst ΔhrcC* (OD_{600} = 0.02) and maintained under HH (**h**) or MH (**i**) for 3 days. Bars represent means (n = 6). Different letters indicate statistically significant differences (P < 0.05; two-way ANOVA followed by Tukey's test in **b-f**; one-way ANOVA followed by Tukey's test in **h** and **i**). WT, wild-type; CAMTA3-A855V, a partial loss-of-function variant; dpi, days post-infiltration. Experiments in (**b-d**), (**f**), (**h**), and (**i**) were repeated twice with similar results.

Fig. 5: *Pst* DC3000 blocks high humidity-driven plant transcriptome reprogramming via T3S effectors.

a, Schematic overview of the experimental design for analyzing high humidity-responsive gene expression in *Pst* DC3000-inoculated leaves. **b**, Bacterial titers in wild-type (WT) leaves infiltrated with *Pst* DC3000 or *Pst* Δ*hrcC* (OD_{600} = 0.02) at 8 hours post-infiltration (hpi) under moderate humidity (MH). Bars represent means (*n* = 6). **c**, Expression levels of *CYP707A3*, *CYP707A1*, and *NCED3* in *Pst* DC3000-inoculated leaves following humidity treatment. WT plants were infiltrated with the indicated *Pst* DC3000 strains (OD_{600} = 0.02), maintained under MH for 8 h, and then exposed to MH or high humidity (HH) for 0.5 h. Bars represent means (*n* = 3). **d**, **e**, Transcriptome profiling of *Pst* DC3000-inoculated leaves following humidity treatment as described in (**a**) and (**c**). **d**, Principal component analysis (PCA) of transcriptomic data. **e**, K-means-clustering of the 2000 most variable genes. Different letters indicate statistically significant differences (*P* < 0.05; Welch's *t*-test in **b**; two-way ANOVA followed by Tukey's test in **c**). Experiments in (**b**) and (**c**) were repeated twice with similar results.

Fig. 6: T3S effector AvrPtoB suppresses *CYP707A3*-mediated water-soaking resistance.
a, Expression levels of *CYP707A3* in *Pst* DC3000-inoculated leaves following humidity treatment. Wild-type (WT) plants were infiltrated with the indicated *Pst* DC3000 strains (OD$_{600}$ = 0.02), maintained under moderate humidity (MH) for 8 h, and then exposed to MH or high humidity (HH) for 0.5 h. Bars represent means (*n* = 3). The box highlights the effectors retained in *Pst* Δ20E but absent in *Pst* Δ28E. **b-d**, Expression levels of *CYP707A3* in humidity-treated leaves with or without induction of *AvrPtoB* (**b**), *AvrPto* (**c**), or *AvrPtoB-F479A* (**d**). The indicated plants were infiltrated with 0.1% ethanol (−DEX) or 10 μM DEX (+DEX), maintained under moderate humidity (MH) for 24 h, then exposed to MH or HH for 0.5 h. Bars represent means (*n* = 3). **e**, ABA levels in humidity-treated leaves with or without *AvrPtoB* induction. The indicated plants were treated with DEX as described in (**b-d**), and then exposed to MH or HH for 2 h. Bars represent means (*n* = 3). **f**, Stomatal aperture in humidity-treated leaves with or without *AvrPtoB* induction. The indicated were treated with DEX as described in (**b-d**), and then exposed to MH or HH for 1 h. Bars represent the median (*n* = 253-367). **g**, Water-soaking in *Pst* Δ*hrcC*-inoculated leaves with or without *AvrPtoB* induction. The indicated plants were infiltrated with *Pst* Δ*hrcC* (OD$_{600}$ = 0.02) supplemented with −DEX or +DEX and maintained under HH for 1 day. The number of water-soaked leaves was pooled from two independent experiments. Different letters indicate statistically significant differences (*P* < 0.05; two-way ANOVA followed by Tukey's test). AvrPtoB-F479A, E3 ligase deficient variant; ND, not detected. Experiments in (**a-d**) and (**f**) were repeated twice with similar results.

Fig. 7: *Pst* DC3000 employs AvrPtoB to suppress *CYP707A3* expression and promote water-soaking under high humidity.

a, Schematic overview of the experimental design for analyzing gene expression, ABA accumulation, water-soaking, and bacterial growth in *Pst* DC3000-inoculated leaves. **b**, Expression levels of *CYP707A3*, *CYP707A1*, and *NCED3* in *Pst* DC3000-inoculated leaves at 6 hours post-infiltration (hpi) under HH. Wild-type (WT) plants were infiltrated with the indicated *Pst* DC3000 strains (OD_{600} = 0.2), maintained under moderate humidity (MH) for 2 h, and then transferred to HH for 4 h. Bars represent means (n = 3). **c**, ABA levels in *Pst* DC3000-inoculated leaves. WT plants were inoculated with the indicated *Pst* DC3000 strains as described in (**b**). Bars represent means (n = 6). **d**, **e**, Representative images (**d**) and quantification (**e**) of water-soaking in *Pst* DC3000-inoculated leaves. WT and *cyp707a1 cyp707a3* mutant plants were infiltrated with the indicated *Pst* DC3000 strains (OD_{600} = 0.2), maintained under MH for 2 h, and then transferred to HH for 20-22 h. Bars represent the median (n = 36; pooled from two independent experiments). **f**, **g**, Disease symptoms (**f**) and bacterial growth (**g**) in *Pst* DC3000-inoculated leaves. WT and *cyp707a1 cyp707a3* mutant plants were infiltrated with the indicated *Pst* DC3000 strains (OD_{600} = 0.0002), maintained under MH for 2 h, and then transferred to HH for 3 days. Bars represent means (n = 6). Different letters indicate statistically significant differences (P < 0.05; one-way ANOVA followed by Tukey's test in **b** and **c**; two-way ANOVA followed by Tukey's test in **e** and **g**). dpi, days post-infiltration. Experiments in (**b**), (**f**), and (**g**) were repeated twice with similar results. **g**, Working model of *CYP707A3*-mediated water-soaking resistance and its suppression by *Pst* DC3000 via T3S effector AvrPtoB under high humidity.

45