LAG3+ CD8+ T cell Subset Drives HR+/HER2- Breast Cancer Reduction in Bispecific

Antibody Armed Activated T Cell Therapy

AUTHORS: Robert Weldon Barnes,*,† Archana Thakur, ‡ Suna Onengut-Gumuscu,§ Lawrence

G. Lum, ‡,¶ Sepideh Dolatshahi *,†,¶

* Department of Biomedical Engineering, University of Virginia (UVA) School of Medicine,

Charlottesville, Virginia 22908

† University of Virginia Cancer Center, Charlottesville, Virginia, USA

‡ Division of Hematology & Oncology, Department of Medicine, UVA School of Medicine

§ Department of Genome Sciences, University of Virginia School of Medicine

¶ Beirne B. Carter Center for Immunology Research, UVA School of Medicine, Charlottesville,

Virginia 22908

RUNNING TITLE: LAG3+ Adoptive T Cells Effectively Reduce HR+ Breast Cancer

CORRESPONDING AUTHOR: Sepideh Dolatshahi, Email: sdolatshahi@virginia.edu, Phone:

434-243-7549

KEYWORDS

Bispecific antibody armed T cell, Checkpoint Inhibitors, Systems Immunology, Therapeutic

synergy, Mechanistic Modeling, Rational Design of Combination Therapies, LAG3, TIGIT,

HR+/HER2- Breast Cancer.

DECLARATIONS

Availability of data and material – Computational modeling algorithms (OPLS-R) and the

mechanistic model of BAT-tumor interactions (ODE model) are available on GitHub

(https://github.com/Dolatshahi-Lab). All cytotoxicity and processed flow cytometry data will be

provided as Supplemental tables. Raw flow cytometry data and CITE-seq data will be provided

upon request.

Competing interests – LGL is cofounder of Transtarget Inc, Founder of BATs, LLC, member of

SAB of Rapa Therapeutics and Tundra Targeted Therapeutics and consults for iCell Gene

Therapeutics. AT is a cofounder of Alpha Immune LLC.

Funding – This work was funded by SD startup funds, UVA Cancer Center Support Grant P30

CA044579 and LGL UVA startup funds. The UVA Comprehensive Cancer Center Support Grant

(P30CA044579) from the NCI additionally supported work performed by the Genome Analysis

and Technology Core, RRID:SCR_018883 and Biorepository and Tissue Research Facility,

RRID:SCR_022971. This research was also supported by the UVA Farrow Fellowship.

Authors contributions – RWB, SD, and LGL were responsible for the conceptualization of the

project and methodologies. RWB and SD wrote the manuscript with input from all authors. SD

and LGL provided supervision. RWB performed experiments, analyzed data, and built the

computational models. RWB, AT, SOG, LGL, and SD contributed to the design and development

of the investigation. SD was responsible for funding acquisition. All authors approved the final

version of the manuscript.

Acknowledgements – This work was presented at the 2024 Society for Immunotherapy of

Cancer (SITC) annual meeting (Abstract #718). We thank the Dolatshahi lab members and Dr.

Kristin Anderson for providing feedback.

Authors' information (optional) –

Robert W. Barnes– ORCID iD 0000-0001-6494-4033

Sepideh Dolatshahi – ORCID iD 0000-0003-0226-0933

Suna Onengut-Gumuscu – ORCID iD 0000-0002-6563-8334

Archana Thakur – ORCID iD 0000-0003-3876-9555

Lawrence G. Lum – ORCID iD 0000-0001-7561-6035

LIST OF ABBREVIATIONS – Tumor microenvironment, TME; Bispecific antibody armed T cell,

BATs; checkpoint inhibitor, CPI; hormone receptor-positive, HR+; Lymphocyte activation gene 3,

LAG3; Programmed Cell Death Protein 1, PD-1; T cell immunoreceptor with immunoglobulin and

ITIM domain, TIGIT; bispecific antibodies, BiAb; overall survival, OS; metastatic breast cancer,

MBC; ordinary differential equation, ODE; unarmed activated T cells, ATC; cellular indexing of

transcriptomes and epitopes, CITE-seq; real-time cell analysis, RTCA; orthogonalized partial

least squares regression, OPLS-R; variable importance in projection, VIP; uniform manifold

approximation and projection, UMAP; T cell receptor, TCR.

What is already known on this topic – Bispecific antibody armed T cell (BATs) therapies are

adoptive T cell therapies that can effectively reroute T cell cytotoxicity toward cancerous cells, but

lack consistent and durable anti-tumor responses. Checkpoint proteins expressed on the surface

of activated T cells dampen immune responses and can be overstimulated in solid tumors to

hamper tumor clearance by T cells. Checkpoint inhibitor drugs can improve T cell anti-tumor

response by blocking checkpoint protein signaling but are only effective if the targeted checkpoint

protein is expressed on the T cell and activated in the tumor microenvironment, highlighting an

opportunity to enhance BAT efficacy by combining treatment with synergistic CPI.

What this study adds – This study characterizes dynamic, time-resolved patterns in checkpoint protein expression by breast cancer-targeting adoptive T cells and predicts the significance of high-prevalence checkpoint proteins on T cell function. It also demonstrates the use of multivariate statistic and mathematical modeling toward rational design of targets and timing strategies for synergistic combination therapies.

How this study might affect research, practice, or policy – The output of this study provides justification for therapeutic strategies combining adoptive T cell therapies with checkpoint inhibitor drugs targeting TIGIT and LAG3 as a means of improving patient responses in HER2-/HR+ breast cancers.

ABSTRACT

Tumor clearance by T cells is impaired by insufficient tumor antigen recognition, insufficient tumor infiltration, and the immunosuppressive tumor microenvironment (TME). Although targeted T cell therapy circumvents failures in tumor antigen recognition, suppression by the TME and failure to infiltrate the tumor can hinder tumor clearance. Checkpoint inhibitors (CPI) promise to reverse T cell suppression and can be combined with bispecific antibody armed T cell (BATs) therapy to improve clinical outcomes. We hypothesize that adoptively transferred T cell function may be improved by the addition of CPI if the inhibitory pathway is functionally active. This study develops a kinetic-dynamic model of killing of hormone receptor-positive (HR+) breast cancer cells mediated by BATs using single-cell transcriptomic and temporal protein data to identify T cell phenotypes and quantify inhibitory receptor expression. LAG3, PD-1, and TIGIT were

identified as inhibitory receptors expressed by cytotoxic effector CD8 BATs upon

exposure to HR+ breast cancer cell lines. These data were combined with real-time tumor

cytotoxicity data in a multivariate statistical analysis framework to predict the relevant

contributions of T cells expressing each receptor to tumor reduction. A mechanistic

kinetic-dynamic mathematical model was developed and parametrized using protein

expression and cytotoxicity data for in silico validation of the findings of the multivariate

statistical analysis. The model corroborated the predictions of the multivariate statistical

analysis which identified LAG3+ BATs as the primary effectors, while TIGIT expression

dampened cytotoxic function. These results inform CPI selection for BATs combination

therapy and provide a framework to maximize BATs anti-tumor function.

INTRODUCTION

HR+/HER2- breast cancers represent 68% of all breast cancer diagnoses(1).

Targeted adoptive T cell therapies offer an effective approach to elicit T cell immune

responses against tumors lacking a sufficiently recognizable tumor antigen(2–6).

Bispecific antibody-armed activated T-cell (BATs) therapies utilize activated T cells armed

with bispecific antibodies (BiAb) targeting CD3 and a tumor antigen to redirect non-MHC

restricted T cell cytotoxicity toward high and low antigen expressing cancer cells(6, 7).

This strategy has shown encouraging results in reducing tumor burden, improving quality

of life, and improving median overall survival (OS) in patients with stage IV breast(8, 9),

prostate(10), and pancreatic(5) cancers and neuroblastoma(11).

In a phase I trial involving 17 heavily pretreated HER2-/HR+ metastatic breast

cancer (MBC) patients, HER2 BATs therapy achieved a median OS of 27.4 months with

59% patients exhibiting stable disease at 14.5 weeks(12). Similarly, a phase II trial with

24 HER2-/HR+ MBC patients reported a decrease in tumor markers in 56.5% of patients

and a median OS of 15.2 months following HER2 BATs infusions(13). Despite these

successes, the lack of consistent and durable anti-tumor responses highlights an

opportunity to enhance BAT efficacy by combining treatment with synergistic T cell

targeting therapies.

Checkpoint inhibitors improve T cell function against solid tumors by preventing

inhibitory signaling. While CPIs have shown efficacy in some breast cancers, response

rates in HER2-/HR+ patients remain disappointingly low (5-12%) compared to HER2+ or

TNBC patients(14–16). CPI efficacy depends on the presence of the target protein and

its complimentary ligands or receptors, suggesting that identifying reliable CPI targets in

HER2-/HR+ tumors could improve therapeutic outcomes (17, 18). By characterizing

inhibitory receptor expression dynamics and BATs function, this study investigates

whether the timing and selection of CPI targets can provide an opportunity for the

synergistic use of CPIs with BATs therapies for HR+/HER2- patients.

Computational modeling integrates diverse measurements to provide insights into

how cellular changes affect therapeutic outcomes. Network modeling has revealed

therapeutically targetable gene expression mechanisms governing T cell fate selection,

function, and exhaustion(20–23), but lacks temporal resolution and cannot account for

repeated T cell stimulation in targeted cell therapies. Existing mechanistic models of

adoptive T cell therapies have described the emergence of T cell functional states(21–

23), but do not address therapeutic dysfunction.

144  To address this gap in knowledge, we developed a dynamic ordinary differential

145 equation (ODE) model that incorporates BATs inhibitory receptor expression profiles. This

146 model predicts the cytotoxic function of each BATs subset and can serve as an *in-silico*

147 testbed to optimize the synergistic use of CPI therapies by targeting the most functional

148 receptors at their peak expression time.

149  Using single-cell transcriptomics and temporal protein and gene expression data,

150 we characterized T cell phenotype emergence and receptor expression dynamics. These

151 data, combined with tumor cytotoxicity profiles, were analyzed using a multivariate

152 statistical framework to predict the relative contributions of BATs subspecies to tumor

153 cytotoxicity.

154  Our ODE model of BATs-mediated tumor killing was constructed from matched

155 temporal protein expression and cytotoxicity data to validate our multivariate analysis

156 findings. Both our multivariate statistical analysis and kinetic-dynamic model identified the

157 LAG3+ BATs subset as a key driver of tumor reduction across multiple donors and cancer

158 cell lines. These results suggest targeting LAG3 in BATs-CPI combinatorial therapies

159 could significantly improve therapeutic outcomes in HR+/HER2- breast cancer patients.

161 **MATERIALS AND METHODS**

162 **Target cell coculture**

163 MCF-7, CAMA1, and T47D cells were acquired from ATCC. Cells were thawed in the

164 media suggested by the manufacturer. MCF-7 cells were passaged in cRPMI (RPMI1640

165 + 10%FBS). CAMA1 and T47D cells were gradually acclimated to cRPMI by subsequent

166   passages in mixed (RPMI/native) media with gradually increasing RPMI proportions.

167   Cells were passaged every 2-3 days to avoid confluence.

169 ATC generation

Activated T cells (ATC) were generated using T cells collected from donor PBMC. Blood

was collected from normal donors and PBMC were obtained by Ficoll-Paque density

gradient separation, counted, suspended in RPMI supplemented with 100 U/mL IL-2 and

20 ng/mL OKT3 to activate T cells and stimulate proliferation[24]. T cells were expanded

by refreshing the cultures with RPMI containing 100 U of IL-2/mL every 2-3 days for 14

days. The expanded T cells were harvested and cryopreserved at $100e^6$ viable cells

(vc)/mL at -160°C in LN2 until use. ATC used in this study were collected from three

donors. Donor information was anonymized prior to the experiment with the exception of

sex. Two donors were female and one donor was male.

180 HER2-targeting Bispecific Antibody Preparation

The HER2-targeting bispecific antibody (HER2bi) was generated by chemical

heteroconjugation of OKT3, a murine IgG2a anti-CD3 monoclonal antibody (Miltenyi

Biotec), and Herceptin, a humanized anti-HER2 IgG1 monoclonal antibody, using sulfo-

succimidyl 4-(N-Maleimidomethyl) cyclohexane-1-carboxylate (Sulfo-SMCC) and Traut's

reagents as described previously[25]. An SDS-PAGE analysis of the HER2bi used in this

study is shown in **Fig S1**.

188 BATs preparation

BATs were prepared as described(26). Briefly, ATC were thawed at 37°C, resuspended

in RPMI, and washed 2x in RPMI. Washed ATC were resuspended in RPMI + 50 ng/10^6

vc anti-HER2x anti-CD3 bispecific antibody (HER2Bi) and rocked on a rocking platform

for 15 minutes to allow for HER2Bi to attach to ATC to generate HER2 BATs(27). HER2

BATs were washed 3x to remove excess biab and resuspended at 10^6 vc/mL. A pictorial

description of BAT generation can be found in **Fig S2**.

Single-cell cellular indexing of transcriptomes and epitopes by sequencing (CITE-

seq) Analysis

ATC from 3 donors were thawed and split into two groups per donor. One group from

each donor was armed as described above to generate BATs. Each group was cocultured

for 1 hours with MCF7 tumor cells at a 10:1 BAT:MCF7 ratio. BATs were then collected

and measured using 10X Genomics CITE-seq technology(27). TotalSeq^{TM}-B

oligonucleotide-labeled antibodies for CD45RA, CD3, CD4, CD8, CD69, and CD25 were

included in the CITE-seq panel, and cells were stained following the manufacturer's

protocol (BioLegend). CITE-seq libraries were processed in parallel following 10X

Genomics Single Cell 3' protocol and sequenced on an Illumina NextSeq 500.

Raw protein tag and transcriptomic data from each donor were integrated into a Seurat

object using the Seurat 5.0 R package(28). Each object was filtered to remove cells with

< 500 features or > 2500 features, as well as those with < 1% or > 5% mitochondrial DNA.

Data within each object was normalized and the 3000 most variable features were

identified. All 6 objects were integrated into a single object using Seurat

FindIntegrationAnchors and IntegrateData functions. Data was then scaled, and a

principal component analysis (PCA) was conducted to identify the top 100 principal

components (PCs). A K nearest-neighbor graph was constructed using the top 25 PCs

and used to identify cell clusters. A uniform manifold approximation and projection

(UMAP) plot for visualization was created using these clusters. Differential gene

expression was used to identify T cell clusters. Briefly, a Wilcoxon Ranked Sum test was

run using Seurat's FindAllMarkers function. Genes strongly differentiating each cluster

were cross-referenced with T cell literature to identify the phenotype in question.

Real-time cell analysis (RTCA) measurements

The xCelligence RTCA monitor was used to measure tumor cell death across the course

of the assay. xCelligence plates were zeroed with RPMI prior to seeding(29). In brief, 2e4

vc target breast cancer cell lines cultured in RPMI1640 supplemented with 10% FBS were

added to each xCelligence well and allowed to adhere. Incubator conditions were

maintained at 37 °C and 5% CO_2. After 24 hours, BATs were added to each well at

predefined ratios between 0.125 – 2 BATs:Tumor. As a surrogate for cytotoxicity,

conductivity measurements were recorded continuously for 120h.

BATs coculture preparation

Time-course samples were prepared by coculturing BATs and tumor cells at an effector:

target ratio of 1:1. Flat bottom 24 well plates were seeded with MCF-7 cells in RPMI at

1.2e5 vc/mL and incubated for 24 hours. After 24h, BATs were added at a E:T as

determined by target cell counts from a sample plate. Bat-tumor cocultures were

incubated for up to 72 hours. Samples were collected from wells at 4, 24, 48, and 72

hours. Collected samples were used for flow cytometry readings.

Checkpoint inhibitor testing

To test the effects of checkpoint inhibitors on BAT function, RTCA plates were seeded

with $2x\ 10^4$ vc/well MCF7 cells, allowed to adhere, and grown for 24 hours. Conditions

and media were identical to those described above. BATs were then mixed with either

10.0, 3.2, 1.0, 0.32, 0.1, 0.032, or 0 ug/mL Tiragolumab biosimilar InVivoSIM anti-human

TIGIT or Relatlimab biosimilar InVivoSIM anti-human LAG3, incubated for 30 minutes at

37C, then transferred at a 1:1 BAT:Tumor ratio into corresponding RTCA wells. BAT-

tumor cocultures were incubated for up to 72 hours. An identical setup in 96 well plates

was performed separately to generate flow cytometry samples. Samples were collected

from wells at 4, 24, 48, and 72 hours. Collected samples were used for flow cytometry

readings.

Flow Cytometry

Flow cytometry stain panels (**Table 1**) were created by mixing fluorescence-conjugated

antibody stains targeting the ligands of interest. The following panels were generated:

Inhibitory (CD45, CD4, CD8, TIGIT, PD-1, LAG3, TIM-3, CTLA, BTLA), Stimulatory

(CD45, CD4, CD8, 4-1BB, CD27, OX40, GITR, HVEM, LIGHT, ICOS), and Coexpression

(CD45, CD4, CD8, 4-1BB, 4-1BBL, OX40, TIGIT, PD-1, LAG3). Cells were resuspended,

counted for viability, stained in 4 °C DPBS, and fixed before fluorescent intensity readings

were acquired on the NovoCyte 3005 3 laser cytometer. Collected data was compensated

and analyzed using FlowJo flow cytometry data analysis software. Cells were identified

using size and complexity characteristics identified from a T cell-only control sample. Live

cells were identified using LIVE/DEAD Fixable Violet Dead Cell stain kit. T cells were

further separated from tumor cells by either CD3 or CD45 expression. T cells were further

subset using anti-CD4 and anti-CD8 antibodies. A sample flow gating strategy is shown

in **Fig S3**. The MFIs of each target stain can be seen in **Fig S4**. Prevalence of each

receptor was calculated by percentage expression within CD4 or CD8 subsets.

Panel	Marker/ Analyte	Reagent/ Analyte Detector	Fluorochrome / Reporter	Source / Vendor	Clone	Dilution/ conc.	RRID
Inhibitory Panel	**CD3**	Anti-CD3	APC	BD	HIT3a	1:5	AB_398592
	CD4	Anti-CD4	AF700	Biolegend	A161A1	1:25	AB_2616933
	CD8	Anti-CD8	APCCY7	BD	RPA-T8	1:25	AB_396865
	CTLA4	Anti-CTLA4	PECY5	BD	BNI3	1:25	AB_396177
	BTLA	Anti-BTLA	PECY7	Biolegend	MIH26	1:25	AB_2629566
	PD-1	Anti-PD-1	BV650	BD	EH12.1	1:25	AB_2738595
	TIGIT	Anti-TIGIT	PE	Biolegend	A15153G	1:25	AB_2632729
	LAG3	Anti-LAG3	BV785	Biolegend	11C3C65	1:25	AB_2716126
	TIM-3	Anti-TIM-3	PerCPEF710	Invitrogen	F38-2E2	1:25	AB_11041821
Coexpression Panel	**CD45**	Anti-CD45	PECY5	Biolegend	HI30	1:25	AB_314397
	CD4	Anti-CD4	AF700	Biolegend	A161A1	1:25	AB_2616933
	CD8	Anti-CD8	APCCY7	BD	RPA-T8	1:25	AB_396865
	OX40	Anti-OX40	FITC	Biolegend	Ber-ACT35	1:25	AB_10661725
	4-1BBL	Anti-41BBL	PECY7	Invitrogen	4H3	1:25	AB_2688229
	PD-1	Anti-PD-1	BV650	BD	EH12.1	1:25	AB_2738595
	TIGIT	Anti-TIGIT	PE	Biolegend	A15153G	1:25	AB_2632729
	LAG3	Anti-LAG3	BV785	Biolegend	11C3C65	1:25	AB_2716126
	4-1BB	Anti-4-1BB	APC	Biolegend	4B4-1	1:25	AB_830671
Stimulatory Panel	**CD45**	Anti-CD45	BV786	Biolegend	HI30	1:25	AB_314397
	CD4	Anti-CD4	AF700	Biolegend	A1616A1	1:25	AB_2616933
	CD8	Anti-CD8	APCCY7	BD	RPA-T8	1:25	AB_396865
	OX40	Anti-OX40	FITC	Biolegend	Ber-ACT35	1:25	AB_10661725
	LIGHT	Anti-LIGHT	PECY7	Biolegend	T5-39	1:25	AB_2721659
	ICOS	Anti-ICOS	PerCPEF710	Invitrogen	ISA-3	1:25	AB_10854730
	HVEM	Anti-HVEM	APC	Biolegend	122	1:25	AB_2565252
	CD27	Anti-CD27	BV510	BD	L128	1:25	AB_2737998
	4-1BB	Anti-41BB	PE	BD	4B4-1	1:5	AB_396252
	GITR	Anti-GITR	PECY5	Biolegend	621	1:25	AB_2240646

Table 1: Targets for fluorescent antibodies included in each flow cytometry panel.

OPLS-R analysis

An orthogonalized partial least squares (OPLS)-regression model was built to predict the

impact of BATs receptor expression on tumor reduction. BATs were binned into distinct

groups by their expression of CD8 and CD4, and again by their expression of PD-1, TIGIT,

and/or LAG3. Triplicate counts were averaged within donor and against target cell line.

These counts were inputted to predict the changes in tumor cell count and the rate of

change in tumor counts as measured by RTCA. Separate models were created for the 3-

donor study and the 3-target study.

Inputs for tumor counts or rate of change compared against BATs counts were taken at

a set time interval from the time of BATs collection. Models for each study were created

with 0, 1, 4, 12, and 24-hour intervals from the time of BATs collection to increase the

robustness of model predictions. The variable importance in projection (VIP) scores for

each BATs subset were calculated(30), ranked, and artificially oriented in the direction of

their loadings on latent variable 1 (LV1). The directionality of groups with VIP scores >1

was used to predict the positive or negative impact on tumor growth for each

receptor/ligand.

ODE model construction and testing

An ODE model was constructed using mass action equations to describe changes in

tumor cell count. The following equation was used to describe tumor cell count.

287
$$\frac{dTumor}{dt} = a_{TP} * Tumor - \frac{a_{TP} * Tumor^2}{Max_{Tumor}} - \left(a_{Kill_N} * NBATs * Tumor\right)$$

288
$$- \left(a_{Kill_L} * LBATs * Tumor\right) - \left(a_{Kill_P} * TBATs * Tumor\right)$$

289
$$- \left(a_{Kill_D} * DBATs * Tumor\right)$$

The first two terms represent tumor growth in the absence of BATs, and the next four

negative terms represent the contribution of each BAT subset to the rate of tumor cell

clearance. Tumor growth rate was assumed to be sigmoidal, and was fit to data collected

from the control 0:1 BATs:Tumor condition during the RTCA experiment. NBATs (LAG3-

TIGIT- CD8s), TBATs (LAG3- TIGIT+ CD8s), LBATs (LAG3+ TIGIT- CD8s), and DBATs

(LAG3+ TIGIT+ CD8s) rate of change were determined by taking the derivative of spline

functions fit to the flow cytometry cell counts collected at 0, 4, 24, 48, and 72 hours.

The CaliPro parameter optimization package was used to determine the potential ranges

and optimal values for each species rate parameter (31). CaliPro optimizes unknown

model parameters by creating a parameter space using Latin Hypercube Sampling within

parameter-specific ranges set by the user. Each parameter was tested by software, and

parameter sets that meet a user-defined set of criteria are used to generate a new LHS

parameter space for a second iteration. This procedure continued until a large enough

portion of parameters met the acceptance criteria. The final parameter space contains a

refined range of parameters which best describe the system in question. Acceptable

parameter spaces were those that generated tumor counts within +/- 10% of the

measured values at 0, 4, 12, 24 ,36, 48, 60, and 72 hours.

RESULTS

Bispecific antibody-driven interactions generate highly cytotoxic LAG3+ effector T

cell populations after 1 hour exposure to target cells

HER2 BATs were generated by arming activated T cells from donor PBMC with a HER2-

targeting bispecific antibody (**Fig S1**) (Materials and Methods) and cocultured with HR+

breast cancer cells. To identify phenotypic changes in T cells driven by the engagement

of T cell receptor (TCR) and tumor antigen mediated by the BiAb, we profiled

transcriptional and protein expression changes in HER2 BATs and unarmed ATC after

exposure to tumor cells using CITE-seq (**Fig 1A**). The matched protein and RNA data

were integrated and pooled, and an unsupervised graph-based clustering method was

applied to generate 12 cell clusters (**Fig 1B**). CD4 and CD8 protein tags were used to

identify major T cell populations, while CD69 and CD45RA protein tags helped

differentiate between resting, dividing, and activated T cell subsets (**Fig 1B, Fig S5**).

These 12 cell subsets exhibited varying transcriptomic expression profiles of cytotoxic

factors, inhibitory receptors, costimulatory receptors, and relevant transcription factors

(**Fig 1D**). Wilcoxon ranked-sum testing was used to identify marker genes distinguishing

each group (**Fig 1C**) to assist in applying known T cell phenotypes.

The presence of the BiAb drove phenotypic changes in both CD4 and CD8 T cells (**Fig**

1A, B). Activated CD4 BATs, indicated by CD69 protein expression, switched primarily to

a pro-inflammatory T helper 1 (Th1) phenotype (**Fig 1C**, **Fig S5**). A smaller subset of

activated CD4 BATs was observed to adopt a humoral Th2 phenotype (**Fig 1B**), which

not only drives B cells to produce immunoglobulins but may also facilitate tumor

progression(32). The Th1 population was comprised primarily of BATs, which accounted

for 82% of the population, while the Th2 population was evenly split between BATs (46%)

and ATC (54%). CD8 BATs displayed progressive activation: a mid-activation group,

labeled CD8_Act, was CD45RA-CD69+, while a fully activated effector group, labeled

CD8_Eff, was CD45RA+CD69+. Additionally, two smaller BATs groups, an NK-like (NKT)

group marked by KLRD1(33) and FCER1G(34) and mucosal-associated invariant (MAIT)

group marked by KLRB1(35) and NCR3(36), were also present. The CD8_Eff group was

identified as the primary cytotoxic group, as demonstrated by high expression of

granzyme (GZMA, GZMB, GZMK, GZMH) and perforin (PRF) genes, as well as

inflammatory cytokine (IFN-γ, TNF-α) and FAS ligand expression (**Fig 1D**). The CD8_Eff

group was also enriched in EOMES and TBX21, two transcriptional regulators associated

with T cell maturation, terminal differentiation, and effector function(37), further indicating

the role of this group as an effector T cell (**Fig 1D**). TCF7, a transcriptional regulator

indicating a pre-antigen encounter T cell state(38), was downregulated in the CD8_Eff

group, while TOX, a gene associated with T cell effector function and exhaustion(39), was

upregulated (**Fig 1D**).

Strong cytotoxic potential was observed to correlate with targetable inhibitory and

costimulatory receptor expression (**Fig 1D**). The cosignaling receptors LAG3 and TIGIT,

which directly modulate BATs cytotoxicity, were enriched in the CD8_Eff group (**Fig 1E**).

While expression of PD-1 (PDCD1), TIM3 (HAVCR2), and CTLA4 was detected in the

CD8_Eff group (**Fig 1D**), their levels were low and not significantly different from those in

the total cell population.

Costimulatory receptors 4-1BB (TNFRSF9), ICOS, CD27, HVEM (TNFRSF14), and

LIGHT (TNFSF14) were observed in the CD8_Eff group (**Fig 1D**). LIGHT and ICOS

expression were upregulated in all activated BATs groups but were not significantly

upregulated in the CD8_Eff group (**Fig 1D**). CD27 was present but downregulated

compared to other CD8 groups (**Fig 1D**), indicating terminal effector differentiation(40).

Only 4-1BB expression was significantly upregulated in the CD8_Eff group (See

Discussion).

While LAG3 expression was observed throughout the CD8_Eff group, TIGIT expression

was concentrated within a subset of these cells (**Fig 1E, F**). Comparison between

$TIGIT_{High}$ and $TIGIT_{Low}$ CD8_Eff cells reveals distinct expression of regulatory genes

associated with T cell effector differentiation. $TIGIT_{High}$ CD8_Effs are enriched for EOMES

and CMC1, while $TIGIT_{Low}$ CD8_Effs are enriched for HOPX and SPRY1. All four genes

are involved in T cell effector differentiation(37, 41–43), but the absence of overlap may

indicate a potential divergence in function between $TIGIT_{High}$ and $TIGIT_{Low}$ CD8_Eff cells.

These expression patterns may indicate that highly cytotoxic CD8 BATs are more strongly

regulated by TIGIT or LAG3 after initial interaction with breast cancer tumor cells and

might benefit from CPI therapies targeting these receptors. Likewise, 4-1BB agonism may

provide costimulatory signals preferentially to a highly cytotoxic subpopulation.

Temporal profiling reveals distinct dynamics of costimulatory and inhibitory

receptor expression following tumor exposure

The emergence of therapeutically targetable cosignaling receptors as marker genes for

highly cytotoxic effector cells after 1 hour of exposure suggests promising potential for

effective combinatorial BATs therapies. To determine whether the observed expression

patterns of TIGIT, LAG3 and 4-1BB from CITE-seq data persist beyond 1 hour of

stimulation and whether transcriptomic expression patterns translate to expression of

surface proteins we analyzed surface receptor expression on BATs cocultured with the

breast cancer cell lines. The coexpression dynamics of inhibitory and stimulatory

receptors on BATs were profiled over a 72-hour coculture with HR+ tumor cells (**Fig 2A**).

Flow cytometry panels used to profile receptor expression were designed based on gene

expression values observed in **Fig 1D** and are detailed in **Table 1**.

CD4 BATs preferentially express PD-1, which coexpresses with LAG3 and TIGIT

CD4 BATs expressed PD-1, TIGIT, and LAG3 during the initial 72-hour challenge (**Fig**

2B). CD4 BATs preferentially expressed PD-1, while LAG3 and TIGIT were primarily

coexpressed on PD-1 expressing cells (**Fig 2E**). ICOS expression within the CD4 pool

gradually increased with time, while CD27 expression dipped over 24h before rebounding

(**Fig S6A**). Th1 CD4 cells expressed PD-1 significantly higher than other BATs groups in

our CITE-seq data, potentially suggesting the immediate emergence of a Th1 CD4 subset

(**Fig 1D**). Th1 CD4 cells expressed PD-1 significantly higher than other BATs groups in

our CD4 data, potentially suggesting the immediate emergence of a Th1 CD4 subset.

CD8 BATs rapidly express LAG3 and 4-1BB while gradually increasing PD-1 and TIGIT

expression

The inhibitory panel showed that CD8 BATs preferentially expressed LAG3 (**Fig 2 B, C**).

TIGIT and PD-1 were observed to a lesser extent on CD8 T cells, and primarily

coexpressed with LAG3 (**Fig 2 C, E, Fig S7**). TIM3, CTLA4, and BTLA were minimally

expressed by BATs over the 72 hours duration of the experiment. In all samples, LAG3+

CD8 BATs hit a proportional and numerical high between 24 and 48 hours after coculture,

before declining across the remaining assay duration. PD-1 and TIGIT expression

gradually increased as the assay progressed. Interestingly, samples that maintained a

higher proportion of LAG3+ CD8 T cells throughout the assay cleared a greater proportion

of tumor cells cumulatively across 72 hours. A stimulatory receptor profile demonstrated

CD8 BATs primarily expressed 4-1BB and ICOS, with 4-1BB spiking rapidly during the

first 24h before exiting the system and ICOS gradually increasing with time. CD27, which

was strongly expressed at t = 0 hours, rapidly decreased in expression during the first 24

hours, before recovering to initial expression levels during the last 48 hours of the

experiment (**Fig S6A**). Costimulatory receptor expression overlap was minimal, with the

exception of CD27 coexpression (**Fig S6B**). The presence of LAG3, TIGIT, and 4-1BB

indicates the CD8 BATs are primarily the CD8_Eff phenotype observed in our CITE-seq

data.

A coexpression panel was designed to explore overlap between costimulatory and

inhibitory receptors (**Fig 2D, F**). Expression of coinhibitory receptors reflected what was

observed on the inhibitory panel – LAG3 and PD1 expression dominated CD8 and CD4

BATs, respectively, with both PD-1 and TIGIT coexpressing with LAG3 at increasing

frequency on CD8s (**Fig 2D, F**). PD-1 and TIGIT rarely coexpressed on CD8s in the

absence of LAG3. Interestingly, 4-1BB strongly coexpressed with LAG3 on CD8s during

the first 24h (**Fig 2D, F**). LAG3 and 4-1BB presence simultaneously peaked at 24 hours,

and LAG3 and TIGIT remained the primary inhibitory receptors after 24 hours, (**Fig 2D,**

F), which suggests that 4-1BB expression or stimulation precedes T cell activation

pathways that preferentially express LAG3 and/or TIGIT.

Multivariate statistical analysis predicts opposing effects of LAG3 and TIGIT

expression on BATs-mediated tumor cell cytotoxicity rates

We hypothesized that LAG3, TIGIT, and PD-1 represent distinct anti-tumor functional

phenotypes, with their expression kinetics closely correlating with changes in BATs

function and phenotype. PD-1 was enriched in activated Th1 and Th2 CD4 subgroups but

did not associate strongly with CD8_Eff in our CITE-seq data (**Fig 1D**). LAG3 and TIGIT

correlated strongly with components of cytotoxic pathways in our CITE-seq data (**Fig 1D**)

and suggests a highly cytotoxic activated effector state in BATs. However, TIGIT was only

partially enriched within our CD8_Eff group and correlated with regulatory components

associated with effector cell maturation (**Fig 1D**). A Partial Least Squares Regression

(PLS-R) model was constructed to determine if tumor cell reduction cytotoxicity could be

predicted by receptor presence (**Fig 3A, B**). Cells were binned by CD4/CD8, TIGIT,

LAG3, and PD-1 expression, with cells expressing combinations of these receptors put in

unique bins. The prevalence of each phenotype at each timepoint was used to predict the

rate of tumor cytotoxicity measured by RTCA at 4 hours post-receptor measurement. Our

model predicted 98% of the variance seen in the rate of tumor growth and reduction, and

outperformed 1000 random permutations of the data (p <0.001) (**Fig 3A**). Examining the

top predictors of the cytotoxicity, which were defined as those with VIP >1, we observed

that CD8 BATs expressing LAG3 with or without PD-1 correlated with increased rates of

cytotoxicity, while TIGIT expression on either CD8 or CD4 cells was negatively correlated

with the rate of cytotoxicity (**Fig 3 B, C**). Cells expressing a combination of LAG3 and

TIGIT, regardless of PD-1 expression, were not predictive. Cells expressing none of the

three receptors were negatively correlated with cytotoxicity, similar to TIGIT expressing

cells, potentially suggesting a pre-activated state. In summary, our data show that the

expression of LAG3 on BATs is significantly associated with anti-tumor cytotoxicity, while

TIGIT expression dampens BATs-driven cytotoxicity (**Fig 3C, Fig S8**). Considering these

results, we predict CD8 BATs will fall into one of 4 generalized functional groups: NBATs

(LAG3-/TIGIT-), LBATs (LAG3+/TIGIT-), TBATs (LAG3-/TIGIT+), and DBATs

(LAG3+/TIGIT+), with exposure to tumor cells driving the transfer of BATs from a LAG3

and/or TIGIT negative group to a LAG3 and/or TIGIT positive group (**Fig 3D**).

Mechanistic modeling identifies LAG3 BATs as key drivers of tumor reduction

The PLS-R model indicates that the balance of BATs subsets defined by receptor

expression plays a crucial role in tumor reduction. To more directly determine the

functional impact of each subset and validate our PLS-R findings, an ODE-based dynamic

model of tumor-BATs interactions was constructed (See Methods). Surviving tumor cell

counts were extracted from RTCA data, while BATs subset counts were extracted from

receptor expression measured by the coexpression flow cytometry panels. Given the

homogeneity in CD4 population dynamics across experiments, cytotoxic CD8 cells were

selected for the development of the model. Cytotoxic potential as predicted for each

subset by the PLS-R model was used to define four functional groups in the kinetic-

dynamic BATs-tumor model. Specifically, the PLS-R model predicted LAG3 and TIGIT

expression have significant but opposing effects on the rate of tumor expansion or

survival. LAG3 expression predicts higher rates of tumor reduction, while TIGIT predicts

weaker tumor reduction. The expression of PD-1 was not predictive. Based on the above

findings, a mechanistic model of tumor-BATs interactions was constructed consisting of

tumor cells and 4 receptor-defined BATs species: LAG3-/TIGIT- BATs (NBATs),

LAG3+/TIGIT- BATs (LBATs), LAG3-/TIGIT+ BATs (TBATs), and LAG3+/TIGIT+ BATs (DBATs) (**Fig 4A**).

Our ODE model was constructed and distinct cytotoxicity rates were estimated for each BATs phenotype, accurately replicating tumor cell count dynamics over a 72-h BAT-tumor coculture. Tumor growth rates were determined by fitting experimental tumor growth data to sigmoidal curves, using maximum cell counts observed in tumor-only control conditions. Experimentally collected BATs counts were modeled using spline functions to represent dynami changes in BATs numbers over time. CaliPro, an iterative parameter optimization package, was employed to estimate the cytotoxicity parameter space. Value ranges for cytotoxicity rates for each BAT phenotype were identified to best match experimental BATs-tumor dynamics (**Fig 4B**). Finally, estimated cytotoxicity rates and the total number of tumor cells killed by each BATs subpopulation were compared as key metrics to evaluate the cytotoxic potential of each cell phenotype.

When comparing a single donor across 3 HER2-/HR+ breast cancer cell lines, LBATs emerge as the clear driver of cytotoxicity (**Table 2** and **Fig 4C**). LBATs demonstrate a larger capacity for tumor clearance both through a higher cytotoxicity rate (**Table 2**) and greater number of tumor cells removed compared to all other species (**Fig 4C**). This model prediction matches the functional potential observed in our PLS-R model, where LAG3+ BATs correlated with greater tumor reduction (**Fig 3B**), and our CITE-seq data, where LAG3+ CD8_Eff cells expressed significantly higher granzyme and perforin concentrations than other observed groups (**Fig 1D**). Both factors make the generation of these cells highly desirable therapeutically. Additionally, LAG3+ BATs were observed to proliferate strongly in all systems, indicating they can sustain an effective anti-tumor

response when generated. DBATs, which expressed both LAG3 and TIGIT, were observed to also contribute to cytotoxicity, although at a lower rate than LBATs (**Fig 4C**). NBATs were 5.75-fold less cytotoxic than LBATs, although their presence as the most populous BAT subtype meant they contributed significantly to

Parameter	Target		
	MCF7	CAMA1	T47D
Tumor growth rate	0.13	0.06	0.16
Max conductivity	1.4	3.2	1.2
LBAT Cytotoxicity Rate (Range)	1.18E-06 (1.01E-06 - 1.34E-06)		
NBAT Cytotoxicity Rate (Range)	1.95E-07 (1.31E-07 - 1.97E-07)		
DBAT Cytotoxicity Rate (Range)	9.54E-07 (6.61E-07 - 9.63E-07)		
DBAT Cytotoxicity Rate (Range)	9.54E-07 (6.61E-07 - 9.63E-07)		

Table 2: Optimal parameters and ranges determined by CaliPro meeting pass criteria for Donor 1 BATs challenged against 3 tumor cell lines.

tumor removal. TBATs demonstrated significantly lower cytotoxicity compared to either LAG3+ subgroups LBATs or DBATs and remained at low enough numbers to contribute minimally to tumor reduction.

Comparisons across donors revealed consistent patterns of cytotoxicity. While the behavior of BATs derived from different donors could not be described using identical parameter sets, individual models predicted LBATs as the most cytotoxic subset, followed by DBATs, with

Parameter	Donor		
	Donor 1	Donor 2	Donor 3
Tumor growth rate	0.0748		
Max conductivity	7.6		
LBAT Cytotoxicity Rate (Range)	1.18E-06 (1.01E-06 - 1.34E-06)	8.59E-06 (8.38E-06- 1.16E-05)	5.28E-06 (3.89E-06 - 5.95E-06)
NBAT Cytotoxicity Rate (Range)	1.95E-07 (1.31E-07 - 1.97E-07)	4.48E-08 (2.17E-08 - 4.91E-08)	4.49E-07 (5.61E-08 - 7.48E-07)
DBAT Cytotoxicity Rate (Range)	9.54E-07 (6.61E-07 - 9.63E-07)	3.71E-06 (7.90E-07 - 4.35E-06)	1.33E-06 (1.27E-06 - 5.52E-06)
TBAT Cytotoxicity Rate (Range)	3.98E-07 (3.23E-07 - 4.25E-07)	8.59E-08 (5.42E-08 - 5.74E-07)	3.36E-07 (3.34E-07 - 7.31E-07)

Table 3: Optimal parameters ranges determined by CaliPro meeting pass criteria for BATs from three donors challenged against MCF7 tumor cells**.**

NBATs and TBATs alternating as the least and second least cytotoxic subgroup (**Fig 5B-**

C, Table 3).

Notably, Donor 2, which demonstrated the lowest tumor cell clearance, relied more

heavily on its DBATs for tumor reduction. This likely resulted from a higher relative

abundance of DBATs generated during the coculture experiment compared to Donors 1

and 3.

Across all donors, LAG3 emerged as a robust indicator of effector cytotoxicity, suggesting

that the LAG3+ CD8 BATs subset plays a primary role in tumor elimination during HR+

breast cancer-HER2 targeted BATs immunotherapy. While LAG3 has known inhibitory

functions, it is known to be upregulated on activated cytotoxic T cells, and its presence

on BATs appears to indicate a functional cytotoxic cell. In contrast, TIGIT expression in

BATs cells may act either as a suppressor of cytotoxic function or an indicator of further

dysfunction. To test whether the activation of either receptor was responsible for the

differences in cytotoxicity observed in each species, the 3-donor experiment was

repeated in the presence of varying concentrations of either an anti-LAG3 Relatlimab

biosimilar or an anti-TIGIT Tiragolumab biosimilar (**Fig S9A**). The addition of anti-LAG3

generated a modest increase in cytotoxicity in Donors 1 and 3, which had higher

proportions of LAG3+ CD8 BATs (**Fig S9B**). While it is not likely that MCF7 cells express

LAG3 ligands MHC-II or FGL-1, LAG3 is known to be autoinhibitory, and thus the addition

of LAG3 may prevent this autoinhibitory action in BATs. The addition of anti-TIGIT did not

have a positive effect on BAT cytotoxicity for any donor, suggesting that TIGIT is not

activated in this system (**Fig S9C**). As MCF7 cells are not known to express TIGIT ligands

PVR, NECTIN1, or NECTIN2, the lack of efficacy is not surprising and suggests that TIGIT

presence in our system indicates less functional BATs. Future in vivo studies should test

the impact of these checkpoint blockades on BATs when in the presence of LAG3 or

TIGIT ligands to confirm these results.

DISCUSSION

Changes in receptor expression as a function of differentiation likely reflect the T cell's

capacity to mount diverse and multifunctional immune responses against perceived

threats. Understanding how these phenotypes emerge, and function can significantly

enhance our ability to engineer T cells for predictable and effective responses against

tumors or pathogens. In this study, we characterized emerging T cell subtypes in BATs

challenged by HR+ HER2- breast cancers and developed a model combining T cell

inhibitory receptor expression dynamics with cytotoxicity rates to identify associations

between targetable pathways of T cell inhibition and T cell anti-tumor cytotoxicity.

While inhibitory receptor expression is intrinsically linked with T cell stimulation, the

factors driving differential expression remain unclear(44–46). Across three donors and

when challenged with 3 HR+ breast cancer cell lines, only TIGIT, LAG3, and PD-1 were

expressed within the first 72 hours of challenge. This suggests that combinatorial

therapies pairing BATs with CPI therapies should prioritize these receptors.

Our model indicates an association between TIGIT surface expression and reduced BATs

function. Because TIGIT ligands PVR, NECTIN1, and NECTIN2 have been detected in

breast cancer and correlate with poor prognostic outcomes, the loss of function in TIGIT+

cells may result of TIGIT-ligand driven inhibition(47, 48). Blocking TIGIT may reverse the

suppressive regulatory effect and augment the cytotoxicity of TIGIT+ and LAG3+/TIGIT+

subset of BATs. However, TIGIT presence on T cells may indicate a more terminally

differentiated and exhausted state. Coexpression of TIGIT with other inhibitory receptors,

including PD-1 and LAG3, is a hallmark of T cell dysfunction in tumors(49–52). Because

we observe TIGIT expression primarily in tandem with PD-1 or LAG3, these cells may be

terminally exhausted and less capable of mediating cytotoxic activity even if TIGIT

activation or activity is blocked.

LAG3+ cells do not appear to be inhibited in our model. LAG3 expression in the absence

of other inhibitory receptors, such as TIGIT or TIM-3, is a hallmark of cytotoxic effector T

cell activation(53–55). LAG3 has been observed as primarily regulating effector functions

like cytotoxicity for tumor-targeting T cells, and thus LAG3 expression by BATs may

indicate a highly functional effector group(56, 57). These studies also observe PD-1 as

primarily responsible for inhibiting cell proliferation, and thus the lack of PD-1 on our

LAG3+ population may facilitate the proliferation observed in this population. The primary

LAG3 ligand, MHC-II, is expressed predominantly on antigen-presenting cells (APCs),

and thus should not be present in our coculture system(58). However, as breast cancers

can be enriched in APCs, including pro-tumor macrophages, LAG3+ BATs may

encounter strong inhibition when applied in vivo(59, 60). Thus, preemptive blockade of

LAG3 may ensure these primary cytotoxic cells maintain their function in patients,

regardless of APC presence. Additionally, LAG3 has been observed to inhibit T cell

cytotoxicity without stimulation(61). Thus, LAG3 blockade may improve BATs function

regardless of APC presence.

The two-cell, two-dimensional model we generated focuses exclusively on BAT changes

that may be triggered through interactions with tumor cells, other T cells, or the cell signals

produced by tumors and T cells. In vivo, the TME will contain a far more diverse array of

cells and signals, many of which may affect BAT development and expression of LAG3,

TIGIT, or related receptors. Other receptors may become more significant players in BAT

function and inhibition when BATs are exposed to the wider range of signals in the TME.

However, the direct tumor-BAT and BAT-BAT interactions our model accounts for will

play a large role in BAT development, and thus the findings here will likely still improve

BAT function.

Our investigation was limited to the dynamic changes in T cell behavior, without capturing

tumor cell ligand expression dynamics. We assumed that components governing the rate

of interaction between BATs and target cells remain constant, except for changing

cytotoxicity observed in different BATs phenotypes. Furthermore, our coculture system

was focused strictly on BiAb-mediated CD8 T cell-tumor interactions and studying signals

and secreted factors from other cells was beyond the scope of our work. These factors

can influence tumor and T cell behaviors affecting cytotoxicity. TCR-activated CD4 and

CD8 T cells are known to produce cytokines and other signaling molecules, including

interferon gamma (IFNγ) and tumor necrosis factor (TNFα)(62–64), which affect both T

cell function and tumor proliferation(65), and were highly expressed in our CD8_Eff group.

While the effect of these cytokines may be captured by the cytotoxicity rates predicted in

our model, any compounding effects of their persistent presence or increasing

concentration may have not been captured by our model. Because the goal of this project

was to characterize CD8 BAT development, CD4 BATs were not included in our

mechanistic model. While certainly relevant to tumor clearance, CD4 cells are largely

known to be cytokine producing cells, which did not fit in our BAT-Tumor interaction

model. CD4 receptor expression was largely similar across donors as well, and thus we

assume that their impact is similar across experiments. Limiting the model to CD8 T cells

might explain the donor-to-donor variability observed in subset-specific tumor clearance

rates (**Table 3**). The initial BAT population for Donor 2 had a much higher proportion of

CD4 T cells compared to Donors 1 and 3 and cleared fewer target cells (**Fig S10**). Donor

2 began with and generated fewer LAG3+ CD8s, and an increased CD4 proportion

demands fewer CD8 cells, so this observation may simply be the result of fewer cytotoxic

BATs present. However, CD8 BATs expansion was also lower with higher CD4, and

particularly PD-1+ CD4, presence, possibly indicating an inhibitory effect elicited from

CD4 BATs. The inclusion of other effector cells might further improve the model fits (**Fig

5A**) but is outside the scope of this paper.

Target cell phenotypic changes also likely change BATs-target interactions but are not

captured in our system. Cytokine presence and cell damage can change receptor

expression on target cells in a manner that may alter BATs-target interactions with time.

Inflammatory cytokines, and in particular Type I interferons, can upregulate inhibitory

receptor expression on T cells(66). Similarly, IFNγ can upregulate cognate ligands for

inhibitory receptors, such as PD-L1, on tumor cells(67). Cell damage and tumor

mutational status can also upregulate these ligands(68–70). Expression of PVR, a TIGIT

ligand, has been observed in response to DNA damage, and may be present on some or

all the target cells in our system(71). Thus, the target cells in our system may evolve to

inhibit BATs expressing certain inhibitory receptors.

Preferential generation of more cytotoxic BATs subspecies may improve tumor clearance

against HR+ breast cancers. Costimulatory signaling is a crucial part of T cell activation

and plays a role in inhibitory receptor expression intensity[72, 73]. Both our CITE-seq and

flow cytometry data indicate 4-1BB expression is associated with cytotoxic effector BATs.

4-1BB was observed to be strongly correlated with LAG3+/TIGIT+ CD8_Eff cells in our

CITE-seq data (**Fig 1D**), and 4-1BB ligand expression was strongly correlated with LAG3

and TIGIT expression for the initial 24-48h of our coculture experiment before

disappearing from our system (**Fig 2D, Fig S6A**). This may indicate that 4-1BB exists to

stimulate highly cytotoxic BATs upon their initial encounter with target cells. However, it

is unclear if 4-1BB stimulation is associated with TIGIT expression alone, or if it is also

linked to LAG3 expression. Further exploration of the impact of 4-1BB stimulation on

BATs function and inhibitory receptor expression may provide insight into what incoming

signals govern BATs function. If 4-1BB activation changes BATs function, whether to

increase or decrease cytotoxicity, utilization of 4-1BB targeting therapies may enhance

tumor reduction. A small number of 4-1BBL+ BATs were observed in our system at 4

hours, which may be indicative of a self-stimulating T cell subtype previously observed to

increase tumor kill in mouse breast cancer models[74]. However, the transient nature of

both 4-1BB and 4-1BBL makes their utilization as targets difficult.

Together, these findings provide a compelling support for the use of predictive statistical

analysis and mathematical modeling in the rational design of combination

immunotherapies. They also strongly suggest that the anti-solid tumor activity of BATs in

our phase I/II metastatic breast cancer clinical trials may be in part driven by the

dominance of a LAG3+ cytotoxic phenotype over TIGIT downregulatory activity[12, 13].

Our recent phase II trial in hormone refractory prostate cancer (HRPC) that combines the

CPI pembrolizumab with HER2 targeted BATs showed significant clinical improvement in

clinical responses and improved survival(75) when the use of CPI alone has not been

successful in clinical test for HRPC. Furthermore, our recent ongoing trial combining

pembrolizumab with HER2 BATs in selected metastatic breast cancer patients heavily

pretreated with chemotherapy led to stable disease and improved survival (unpublished).

The improvement in patient response to BATs combined with a PD-1/PD-L1 targeting

therapy bodes well for further work in this area: our model predicted PD-1 to have a lower

impact on BATs function than LAG3 or TIGIT and thus combining BATs with a LAG3 or

TIGIT blockade may improve responses to an even greater extent. Further investigations

into CPI and BATs combination strategies targeting stimulatory or inhibitory receptors to

enhance BATs effector functions will likely improve the potential of BATs therapy in

metastatic breast cancers and other solid tumors.

REFERENCES

1. Giaquinto, A. N., H. Sung, K. D. Miller, J. L. Kramer, L. A. Newman, A. Minihan, A. Jemal, and R. L. Siegel. 2022. Breast Cancer Statistics, 2022. *CA Cancer J Clin* 72: 524–541.

2. Albarrán, V., M. San Román, J. Pozas, J. Chamorro, D. I. Rosero, P. Guerrero, J. C. Calvo, C. González, C. García de Quevedo, P. Pérez de Aguado, J. Moreno, A. Cortés, and A. Soria. 2024. Adoptive T cell therapy for solid tumors: current landscape and future challenges. *Front Immunol* 15: 1352805.

3. Helmich, B. K., and R. W. Dutton. 2001. The role of adoptively transferred CD8 T cells and host cells in the control of the growth of the EG7 thymoma: factors that determine the relative effectiveness and homing properties of Tc1 and Tc2 effectors. *J Immunol* 166: 6500–6508.

4. Lee, H. J., Y.-A. Kim, C. K. Sim, S.-H. Heo, I. H. Song, H. S. Park, S. Y. Park, W. Seon Bang, I. A. Park, M. Lee, J. H. Lee, Y. S. Cho, S. Chang, J. Jung, J. Kim, S. B. Lee, S. Y. Kim, M. S. Lee, and G. Gong. 2017. Expansion of tumor-infiltrating lymphocytes and their potential for application as adoptive cell transfer therapy in human breast cancer. 8: 113345–113359.

5. Lum, L. G., A. Thakur, M. Choi, A. Deol, V. Kondadasula, D. Schalk, K. Fields, M. Dufrense, P. Philip, G. Dyson, H. D. Aon, and A. F. Shields. 2020. Clinical and immune responses to anti-CD3 x anti-EGFR bispecific antibody armed activated T cells (EGFR BATs) in pancreatic cancer patients. *Oncoimmunology* 9.

6. Grabert, R. C., L. P. Cousens, J. A. Smith, S. Olson, J. Gall, W. B. Young, P. A. Davol, and L. G. Lum. 2006. Human T Cells Armed with Her2/neu Bispecific Antibodies Divide, Are Cytotoxic, and Secrete Cytokines with Repeated Stimulation Cancer Therapy: Preclinical. *Clin Cancer Res* 12.

7. Lum, L. G., A. Thakur, Q. Liu, A. Deol, Z. Al-Kadhimi, L. Ayash, M. H. Abidi, C. Pray, E. N. Tomaszewski, P. A. Steele, D. L. Schalk, H. Yano, A. Mitchell, M. Dufresne, J. P. Uberti, and V. Ratanatharathorn. 2013. CD20-targeted T cells after stem cell transplantation for high risk and refractory non-Hodgkin's lymphoma. *Biol Blood Marrow Transplant* 19: 925–933.

8. Thakur, A., R. Rathore, S. V. Kondadasula, J. P. Uberti, V. Ratanatharathorn, and L. G. Lum. 2018. Immune T cells can transfer and boost anti-breast cancer immunity. *Oncoimmunology* 7.

9. Lum, L. G., A. Thakur, Z. Al-Kadhimi, G. A. Colvin, F. J. Cummings, R. D. Legare, D. S. Dizon, N. Kouttab, A. Maizel, W. Colaiace, Q. Liu, and R. Rathore. 2015. Targeted T-cell therapy in stage IV breast cancer: A phase i clinical trial. *Clinical Cancer Research* 21: 2305–2314.

10. Davol, P. A., J. A. Smith, N. Kouttab, G. J. Elfenbein, and L. G. Lum. 2004. Anti-CD3 × Anti-HER2 Bispecific Antibody Effectively Redirects Armed T Cells to Inhibit Tumor Development and Growth

in Hormone-Refractory Prostate Cancer–Bearing Severe Combined Immunodeficient Beige Mice. *Clin Prostate Cancer* 3: 112–121.

11. Yankelevich, M., A. Thakur, S. Modak, R. Chu, J. Taub, A. Martin, D. Schalk, A. Schienshang, S. Whitaker, K. Rea, D. W. Lee, Q. Liu, A. F. Shields, N. K. V. Cheung, and L. G. Lum. 2024. Targeting refractory/recurrent neuroblastoma and osteosarcoma with anti-CD3×anti-GD2 bispecific antibody armed T cells. *J Immunother Cancer* 12: e008744.

12. Lum, L. G., A. Thakur, Z. Al-Kadhimi, G. A. Colvin, F. J. Cummings, R. D. Legare, D. S. Dizon, N. Kouttab, A. Maizel, W. Colaiace, Q. Liu, and R. Rathore. 2015. Targeted T-cell therapy in stage IV breast cancer: A phase i clinical trial. *Clinical Cancer Research* 21: 2305–2314.

13. Lum, L. G., Z. Al-Kadhimi, A. Deol, V. Kondadasula, D. Schalk, E. Tomashewski, P. Steele, K. Fields, M. Giroux, Q. Liu, L. Flaherty, M. Simon, and A. Thakur. 2021. Phase II clinical trial using anti-CD3 × anti-HER2 bispecific antibody armed activated T cells (HER2 BATs) consolidation therapy for HER2 negative (0–2+) metastatic breast cancer. *J Immunother Cancer* 9: e002194.

14. Rugo, H. S., J. P. Delord, S. A. Im, P. A. Ott, S. A. Piha-Paul, P. L. Bedard, J. Sachdev, C. Le Tourneau, E. M. J. van Brummelen, A. Varga, R. Salgado, S. Loi, S. Saraf, D. Pietrangelo, V. Karantza, and A. R. Tan. 2018. Safety and Antitumor Activity of Pembrolizumab in Patients with Estrogen Receptor-Positive/Human Epidermal Growth Factor Receptor 2-Negative Advanced Breast Cancer. *Clin Cancer Res* 24: 2804–2811.

15. Cardoso, F., L. Jia, K. Hirshfield, and V. Karantza. 2019. KEYNOTE-756: Randomized, double-blind, phase III study of pembrolizumab vs placebo + neoadjuvant chemotherapy (CT) and adjuvant endocrine therapy (ET) for high-risk, early-stage estrogen receptor–positive human epidermal growth factor receptor 2–negative (ER+/HER2−) breast cancer (BC). *Annals of Oncology* 30: iii38.

16. Dirix, L. Y., I. Takacs, G. Jerusalem, P. Nikolinakos, H. T. Arkenau, A. Forero-Torres, R. Boccia, M. E. Lippman, R. Somer, M. Smakal, L. A. Emens, B. Hrinczenko, W. Edenfield, J. Gurtler, A. von Heydebreck, H. J. Grote, K. Chin, and E. P. Hamilton. 2018. Avelumab, an anti-PD-L1 antibody, in patients with locally advanced or metastatic breast cancer: a phase 1b JAVELIN Solid Tumor study. *Breast Cancer Res Treat* 167: 671–686.

17. Rotte, A., J. Y. Jin, and V. Lemaire. 2018. Mechanistic overview of immune checkpoints to support the rational design of their combinations in cancer immunotherapy. *Ann Oncol* 29: 71–83.

18. Goleva, E., T. Lyubchenko, L. Kraehenbuehl, M. E. Lacouture, D. Y. M. Leung, and J. A. Kern. 2021. Our current understanding of checkpoint inhibitor therapy in cancer immunotherapy. *Ann Allergy Asthma Immunol* 126: 630–638.

19. Ildefonso, G. V., and S. D. Finley. 2023. A data-driven Boolean model explains memory subsets and evolution in CD8+ T cell exhaustion. *npj Systems Biology and Applications 2023 9:1* 9: 1–13.

20. Yan, M., J. Hu, H. Yuan, L. Xu, G. Liao, Z. Jiang, J. Zhu, B. Pang, Y. Ping, Y. Zhang, Y. Xiao, and X. Li. 2021. Dynamic regulatory networks of T cell trajectory dissect transcriptional control of T cell state transition. *Mol Ther Nucleic Acids* 26: 1115–1129.

21. Bolouri, H., M. Young, J. Beilke, R. Johnson, B. Fox, L. Huang, C. C. Santini, C. M. Hill, A. R. van der V. de Vries, P. T. Shannon, A. Dervan, P. Sivakumar, M. Trotter, D. Bassett, and A. Ratushny. 2020. Integrative network modeling reveals mechanisms underlying T cell exhaustion. *Scientific Reports 2020 10:1* 10: 1–15.

22. Sontag, E. D. 2017. A Dynamic Model of Immune Responses to Antigen Presentation Predicts Different Regions of Tumor or Pathogen Elimination. *Cell Syst* 4: 231-241.e11.

23. Cess, C. G., and S. D. Finley. 2020. Data-driven analysis of a mechanistic model of CAR T cell signaling predicts effects of cell-to-cell heterogeneity. *J Theor Biol* 489: 110125.

24. Uberti, J. P., I. Joshi, M. Ueda, F. Martilotti, L. L. Sensenbrenner, and L. G. Lum. 1994. Preclinical Studies Using Immobilized OKT3 to Activate Human T Cells for Adoptive Immunotherapy: Optimal Conditions for the Proliferation and Induction of Non-MHC-Restricted Cytotoxicity. *Clin Immunol Immunopathol* 70: 234–240.

25. Sen, M., D. M. Wankowski, N. K. Garlie, R. E. Siebenlistd, D. Van Epps, A. V. LeFever, and L. G. Lum. 2001. Use of anti-CD3 x anti-HER2/neu bispecific antibody for redirecting cytotoxicity of activated T cells toward HER2/neu+ tumors. *J Hematother Stem Cell Res* 10: 247–260.

26. Sen, M., D. M. Wankowski, N. K. Garlie, R. E. Siebenlistd, D. Van Epps, A. V. LeFever, and L. G. Lum. 2004. Use of Anti-CD3 × Anti-HER2/neu Bispecific Antibody for Redirecting Cytotoxicity of Activated T Cells Toward HER2/neu+ Tumors. *https://home.liebertpub.com/scd* 10: 247–260.

27. Stoeckius, M., C. Hafemeister, W. Stephenson, B. Houck-Loomis, P. K. Chattopadhyay, H. Swerdlow, R. Satija, and P. Smibert. 2017. Simultaneous epitope and transcriptome measurement in single cells. *Nature Methods 2017 14:9* 14: 865–868.

28. Hao, Y., T. Stuart, M. H. Kowalski, S. Choudhary, P. Hoffman, A. Hartman, A. Srivastava, G. Molla, S. Madad, C. Fernandez-Granda, and R. Satija. 2023. Dictionary learning for integrative, multimodal and scalable single-cell analysis. *Nature Biotechnology 2023 42:2* 42: 293–304.

29. Thakur, A., J. Ung, E. N. Tomaszewski, A. Schienschang, T. M. LaBrie, D. L. Schalk, and L. G. Lum. 2021. Priming of pancreatic cancer cells with bispecific antibody armed activated T cells sensitizes tumors for enhanced chemoresponsiveness. *Oncoimmunology* 10.

30. Chong, I. G., and C. H. Jun. 2005. Performance of some variable
selection methods when multicollinearity is present. *Chemometrics*
and Intelligent Laboratory Systems 78: 103–112.

31. Joslyn, L. R., D. E. Kirschner, and J. J. Linderman. CaliPro: A
Calibration Protocol That Utilizes Parameter Density Estimation to
Explore Parameter Space and Calibrate Complex Biological Models. .

32. Guo, M., M. Y. R. Liu, and D. G. Brooks. 2024. Regulation and
impact of tumor-specific CD4+ T cells in cancer and immunotherapy.
Trends Immunol 45: 303–313.

33. Zheng, L., S. Qin, W. Si, A. Wang, B. Xing, R. Gao, X. Ren, L.
Wang, X. Wu, J. Zhang, N. Wu, N. Zhang, H. Zheng, H. Ouyang, K.
Chen, Z. Bu, X. Hu, J. Ji, and Z. Zhang. 2021. Pan-cancer single-cell
landscape of tumor-infiltrating T cells. *Science (1979)* 374.

34. Correia, M. P., A. Stojanovic, K. Bauer, D. Juraeva, L. O.
Tykocinski, H. M. Lorenz, B. Brors, and A. Cerwenka. 2018. Distinct
human circulating NKp30+FceRIγ+CD8+ T cell population exhibiting
high natural killer-like antitumor potential. *Proc Natl Acad Sci U S A*
115: E5980–E5989.

35. Garner, L. C., A. Amini, M. E. B. FitzPatrick, M. J. Lett, G. F.
Hess, M. Filipowicz Sinnreich, N. M. Provine, and P. Klenerman.
2023. Single-cell analysis of human MAIT cell transcriptional,
functional and clonal diversity. *Nature Immunology 2023 24:9* 24:
1565–1578.

36. Lu, B., M. Liu, J. Wang, H. Fan, D. Yang, L. Zhang, X. Gu, J.
Nie, Z. Chen, A. J. Corbett, M. J. Zhan, S. Zhang, V. L. Bryant, A. M.
Lew, J. McCluskey, H. bin Luo, J. Cui, Y. Zhang, Y. Zhan, and G. Lu.
2020. IL-17 production by tissue-resident MAIT cells is locally
induced in children with pneumonia. *Mucosal Immunol* 13: 824–835.

37. Intlekofer, A. M., N. Takemoto, E. J. Wherry, S. A. Longworth, J.
T. Northrup, V. R. Palanivel, A. C. Mullen, C. R. Gasink, S. M.
Kaech, J. D. Miller, L. Gapin, K. Ryan, A. P. Russ, T. Lindsten, J. S.
Orange, A. W. Goldrath, R. Ahmed, and S. L. Reiner. 2005. Effector
and memory CD8+ T cell fate coupled by T-bet and eomesodermin.
Nature Immunology 2005 6:12 6: 1236–1244.

38. Willinger, T., T. Freeman, M. Herbert, H. Hasegawa, A. J.
McMichael, and M. F. C. Callan. 2006. Human Naive CD8 T Cells
Down-Regulate Expression of the WNT Pathway Transcription
Factors Lymphoid Enhancer Binding Factor 1 and Transcription
Factor 7 (T Cell Factor-1) following Antigen Encounter In Vitro and
In Vivo. *The Journal of Immunology* 176: 1439–1446.

39. Khan, O., J. R. Giles, S. McDonald, S. Manne, S. F. Ngiow, K. P.
Patel, M. T. Werner, A. C. Huang, K. A. Alexander, J. E. Wu, J.
Attanasio, P. Yan, S. M. George, B. Bengsch, R. P. Staupe, G.
Donahue, W. Xu, R. K. Amaravadi, X. Xu, G. C. Karakousis, T. C.
Mitchell, L. M. Schuchter, J. Kaye, S. L. Berger, and E. J. Wherry.

2019. TOX transcriptionally and epigenetically programs CD8+ T cell exhaustion. *Nature 2019 571:7764* 571: 211–218.

40. Bullock, T. N. 2017. Stimulating CD27 to quantitatively and qualitatively shape adaptive immunity to cancer. *Curr Opin Immunol* 45: 82–88.

41. Albrecht, I., U. Niesner, M. Janke, A. Menning, C. Loddenkemper, A. A. Kühl, I. Lepenies, M. H. Lexberg, K. Westendorf, K. Hradilkova, J. Grün, A. Hamann, J. A. Epstein, H. D. Chang, K. Tokoyoda, and A. Radbruch. 2010. Persistence of effector memory Th1 cells is regulated by Hopx. *Eur J Immunol* 40: 2993–3006.

42. Shehata, H. M., S. Khan, E. Chen, P. E. Fields, R. A. Flavell, and S. Sanjabi. 2018. Lack of Sprouty 1 and 2 enhances survival of effector CD8+ T cells and yields more protective memory cells. *Proc Natl Acad Sci U S A* 115: E8939–E8947.

43. Chen, Y., J. Gao, M. Ma, K. Wang, F. Liu, F. Yang, X. Zou, Z. Cheng, and D. Wu. 2024. The potential role of CMC1 as an immunometabolic checkpoint in T cell immunity. *Oncoimmunology* 13.

44. Thommen, D. S., J. Schreiner, P. Müller, P. Herzig, A. Roller, A. Belousov, P. Umana, P. Pisa, C. Klein, M. Bacac, O. S. Fischer, W. Moersig, S. S. Prince, V. Levitsky, V. Karanikas, D. Lardinois, and A. Zippelius. 2015. Progression of lung cancer is associated with increased dysfunction of T cells defined by coexpression of multiple inhibitory receptors. *Cancer Immunol Res* 3: 1344–1354.

45. Baitsch, L., A. Legat, L. Barba, S. A. Marraco, J. P. Rivals, P. Baumgaertner, C. Christiansen-Jucht, H. Bouzourene, D. Rimoldi, H. Pircher, N. Rufer, M. Matter, O. Michielin, and D. E. Speiser. 2012. Extended Co-Expression of Inhibitory Receptors by Human CD8 T-Cells Depending on Differentiation, Antigen-Specificity and Anatomical Localization. *PLoS One* 7: e30852.

46. Li, X., R. Wang, P. Fan, X. Yao, L. Qin, Y. Peng, M. Ma, N. Asley, X. Chang, Y. Feng, Y. Hu, Y. Zhang, C. Li, G. Fanning, S. Jones, C. Verrill, D. Maldonado-Perez, P. Sopp, C. Waugh, S. Taylor, S. McGowan, V. Cerundolo, C. Conlon, A. McMichael, S. Lu, X. Wang, N. Li, and T. Dong. 2019. A comprehensive analysis of key immune checkpoint receptors on tumor-infiltrating t cells from multiple types of cancer. *Front Oncol* 9: 472080.

47. Chiang, E. Y., and I. Mellman. 2022. TIGIT-CD226-PVR axis: advancing immune checkpoint blockade for cancer immunotherapy. *J Immunother Cancer* 10: e004711.

48. Boissière-Michot, F., M. C. Chateau, S. Thézenas, S. Guiu, A. Bobrie, and W. Jacot. 2022. Correlation of the TIGIT-PVR immune checkpoint axis with clinicopathological features in triple-negative breast cancer. *Front Immunol* 13: 1058424.

49. Worboys, J. D., K. N. Vowell, R. K. Hare, A. R. Ambrose, M.
Bertuzzi, M. A. Conner, F. P. Patel, W. H. Zammit, J. Gali-Moya, K.
S. Hazime, K. L. Jones, C. Rey, S. Jonjic, T. L. Rovis, G. M.
Tannahill, G. D. S. Cruz De Matos, J. D. Waight, and D. M. Davis.
2023. TIGIT can inhibit T cell activation via ligation-induced
nanoclusters, independent of CD226 co-stimulation. *Nature*
Communications 2023 14:1 14: 1–21.
50. Liu, Z., H. Zeng, K. Jin, Y. Yu, R. You, H. Zhang, C. Liu, X. Su,
S. Yan, Y. Chang, L. Liu, L. Xu, J. Xu, Y. Zhu, and Z. Wang. 2022.
TIGIT and PD-1 expression atlas predicts response to adjuvant
chemotherapy and PD-L1 blockade in muscle-invasive bladder cancer.
British Journal of Cancer 2022 126:9 126: 1310–1317.
51. Wherry, E. J., and M. Kurachi. 2015. Molecular and cellular
insights into T cell exhaustion. *Nature Reviews Immunology 2015*
15:8 15: 486–499.
52. Takamatsu, K., N. Tanaka, K. Hakozaki, R. Takahashi, Y.
Teranishi, T. Murakami, R. Kufukihara, N. Niwa, S. Mikami, T.
Shinojima, T. Sasaki, Y. Sato, H. Kume, S. Ogawa, K. Kakimi, T.
Kamatani, F. Miya, T. Tsunoda, E. Aimono, H. Nishihara, K. Sawada,
T. Imamura, R. Mizuno, and M. Oya. 2021. Profiling the inhibitory
receptors LAG-3, TIM-3, and TIGIT in renal cell carcinoma reveals
malignancy. *Nature Communications 2021 12:1* 12: 1–12.
53. Workman, C. J., and D. A. A. Vignali. 2003. The CD4-related
molecule, LAG-3 (CD223), regulates the expansion of activated T
cells. *Eur J Immunol* 33: 970–979.
54. Grosso, J. F., M. V. Goldberg, D. Getnet, T. C. Bruno, H.-R. Yen,
K. J. Pyle, E. Hipkiss, D. A. A. Vignali, D. M. Pardoll, and C. G.
Drake. 2009. Functionally Distinct LAG-3 and PD-1 Subsets on
Activated and Chronically Stimulated CD8 T Cells. *The Journal of*
Immunology 182: 6659–6669.
55. Kallies, A., D. Zehn, and D. T. Utzschneider. 2019. Precursor
exhausted T cells: key to successful immunotherapy? *Nature Reviews*
Immunology 2019 20:2 20: 128–136.
56. Ngiow, S. F., S. Manne, Y. J. Huang, T. Azar, Z. Chen, D.
Mathew, Q. Chen, O. Khan, J. E. Wu, V. Alcalde, A. J. Flowers, S.
McClain, A. E. Baxter, M. Kurachi, J. Shi, A. C. Huang, J. R. Giles,
A. H. Sharpe, D. A. A. Vignali, and E. J. Wherry. 2024. LAG-3
sustains TOX expression and regulates the CD94/NKG2-Qa-1b axis to
govern exhausted CD8 T cell NK receptor expression and
cytotoxicity. *Cell* 187: 4336-4354.e19.
57. Andrews, L. P., S. C. Butler, J. Cui, A. R. Cillo, C. Cardello, C.
Liu, E. A. Brunazzi, A. Baessler, B. Xie, S. R. Kunning, S. F. Ngiow,
Y. J. Huang, S. Manne, A. H. Sharpe, G. M. Delgoffe, E. J. Wherry, J.
M. Kirkwood, T. C. Bruno, C. J. Workman, and D. A. A. Vignali.
2024. LAG-3 and PD-1 synergize on CD8+ T cells to drive T cell

exhaustion and hinder autocrine IFN-γ-dependent anti-tumor immunity. *Cell* 187: 4355-4372.e22.

58. Baixeras, E., B. Huard, C. Miossec, S. Jitsukawa, M. Martin, T. Hercend, C. Auffray, F. Triebel, and D. Piatier-Tonneau. 1992. Characterization of the lymphocyte activation gene 3-encoded protein. A new ligand for human leukocyte antigen class II antigens. *Journal of Experimental Medicine* 176: 327–337.

59. Williams, C. B., E. S. Yeh, and A. C. Soloff. 2016. Tumor-associated macrophages: unwitting accomplices in breast cancer malignancy. *npj Breast Cancer 2016 2:1* 2: 1–12.

60. Medrek, C., F. Pontén, K. Jirström, and K. Leandersson. 2012. The presence of tumor associated macrophages in tumor stroma as a prognostic marker for breast cancer patients. *BMC Cancer* 12: 1–9.

61. Guy, C., D. M. Mitrea, P. C. Chou, J. Temirov, K. M. Vignali, X. Liu, H. Zhang, R. Kriwacki, M. P. Bruchez, S. C. Watkins, C. J. Workman, and D. A. A. Vignali. 2022. LAG3 associates with TCR–CD3 complexes and suppresses signaling by driving co-receptor–Lck dissociation. *Nature Immunology 2022 23:5* 23: 757–767.

62. Bachmann, M. F., M. Barner, A. Viola, and M. Kopf. Distinct kinetics of cytokine production and cytolysis in effector and memory T cells after viral infection. .

63. Veiga-Fernandes, H., U. Walter, C. Bourgeois, A. McLean, and B. Rocha. 2000. Response of naïve and memory CD8+ T cells to antigen stimulation in vivo. *Nature Immunology 2000 1:1* 1: 47–53.

64. Helmich, B. K., and R. W. Dutton. 2001. The role of adoptively transferred CD8 T cells and host cells in the control of the growth of the EG7 thymoma: factors that determine the relative effectiveness and homing properties of Tc1 and Tc2 effectors. *J Immunol* 166: 6500–6508.

65. Dong, C. 2021. Cytokine Regulation and Function in T Cells. *Annu Rev Immunol* 39: 51–76.

66. Sumida, T. S., S. Dulberg, J. C. Schupp, M. R. Lincoln, H. A. Stillwell, P. P. Axisa, M. Comi, A. Unterman, N. Kaminski, A. Madi, V. K. Kuchroo, and D. A. Hafler. 2022. Type I interferon transcriptional network regulates expression of coinhibitory receptors in human T cells. *Nature Immunology 2022 23:4* 23: 632–642.

67. Spranger, S., R. M. Spaapen, Y. Zha, J. Williams, Y. Meng, T. T. Ha, and T. F. Gajewski. 2013. Up-regulation of PD-L1, IDO, and Tregs in the melanoma tumor microenvironment is driven by CD8+ T cells. *Sci Transl Med* 5.

68. Sato, H., A. Niimi, T. Yasuhara, T. B. M. Permata, Y. Hagiwara, M. Isono, E. Nuryadi, R. Sekine, T. Oike, S. Kakoti, Y. Yoshimoto, K. D. Held, Y. Suzuki, K. Kono, K. Miyagawa, T. Nakano, and A. Shibata. 2017. DNA double-strand break repair pathway regulates PD-L1 expression in cancer cells. *Nature Communications 2017 8:1* 8: 1–11.

69. Strickland, K. C., B. E. Howitt, S. A. Shukla, S. Rodig, L. L.
Ritterhouse, J. F. Liu, J. E. Garber, D. Chowdhury, C. J. Wu, A. D.
D'Andrea, U. A. Matulonis, and P. A. Konstantinopoulos. 2016.
Association and prognostic significance of BRCA1/2-mutation status
with neoantigen load, number of tumor-infiltrating lymphocytes and
expression of PD-1/PD-L1 in high grade serous ovarian cancer.
Oncotarget 7: 13587.
70. Rosenbaum, M. W., J. R. Bledsoe, V. Morales-Oyarvide, T. G.
Huynh, and M. Mino-Kenudson. 2016. PD-L1 expression in colorectal
cancer is associated with microsatellite instability, BRAF mutation,
medullary morphology and cytotoxic tumor-infiltrating lymphocytes.
Modern Pathology 29: 1104–1112.
71. Gorvel, L., and D. Olive. 2020. Targeting the "PVR–TIGIT axis"
with immune checkpoint therapies. *F1000Res* 9.
72. Messenheimer, D. J., S. M. Jensen, M. E. Afentoulis, K. W.
Wegmann, Z. Feng, D. J. Friedman, M. J. Gough, W. J. Urba, and B.
A. Fox. Cancer Therapy: Preclinical Timing of PD-1 Blockade Is
Critical to Effective Combination Immunotherapy with Anti-OX40.
Clin Cancer Res 23.
73. Koyama, S., E. A. Akbay, Y. Y. Li, G. S. Herter-Sprie, K. A.
Buczkowski, W. G. Richards, L. Gandhi, A. J. Redig, S. J. Rodig, H.
Asahina, R. E. Jones, M. M. Kulkarni, M. Kuraguchi, S. Palakurthi, P.
E. Fecci, B. E. Johnson, P. A. Janne, J. A. Engelman, S. P.
Gangadharan, D. B. Costa, G. J. Freeman, R. Bueno, F. S. Hodi, G.
Dranoff, K. K. Wong, and P. S. Hammerman. 2016. Adaptive
resistance to therapeutic PD-1 blockade is associated with
upregulation of alternative immune checkpoints. *Nature
Communications 2016 7:1* 7: 1–9.
74. Cho, E., R. Singh, C. Han, S. H. Kim, K. H. Kim, B. M. Park, D.
H. Shin, S. Han, Y. H. Kim, B. S. Kwon, K. T. Nam, and B. K. Choi.
2023. 4-1BB-4-1BBL cis-interaction contributes to the survival of
self-reactive CD8+ T cell. *Cellular & Molecular Immunology 2023
20:9* 20: 1077–1080.
75. Vaishampayan, U. N., A. Thakur, W. Chen, A. Deol, M. Patel, K.
Dobson, B. Dickow, D. Schalk, A. Schienschang, S. Whitaker, A.
Polend, J. A. Fontana, E. I. Heath, and L. G. Lum. 2023. Phase II Trial
of Pembrolizumab and Anti-CD3 x Anti-HER2 Bispecific Antibody-
Armed Activated T Cells in Metastatic Castration-Resistant Prostate
Cancer. *Clinical Cancer Research* 29: 122–133.
1028
1029

Figure captions

1031

Figure 1:

A: UMAP plot of CITE-seq data labeled by bispecific antibody arming. ATC: unarmed Activated T

Cells, BAT: Bispecific antibody-armed Activated T cells

B: A UMAP plot of T cells clustered by similarity in gene expression. Labels were chosen based

on expression of significantly upregulated marker genes. CD4_Rest = Resting CD4 T cells;

CD8_Rest_1 = Group 1 Resting CD8 T cells; CD4_Th1 = Th1 CD4 T cells; CD8_Eff = Effector

CD8 T cells; Div_1 = Group 1 Dividing T cells; CD8_Act = Activated CD8 T cells; CD8_Rest_2 =

Group 2 Resting CD8 T cells; Div_2 = Group 2 Dividing T cells; CD4_Th17 = Th17 CD4 T cells;

CD4_Th2 = Th2 CD4 T cells; MAIT = Mucosal Associated Invariant T cells; NKT = NK-like T cells

C: Heatmap of most differentially expressed genes as determined by a Wilcoxon ranked sum test.

Genes shown most strongly differentiate each group from all other groups and were used to

determine the labels in 1B.

D: A heat map of average gene expression scaled across groups. Genes displayed are involved

in cytotoxicity (top), T cell inhibition (second top), T cell costimulation (second bottom), and T cell

maturation and differentiation (bottom).

E: A re-clustered UMAP of effector CD8 T cells.

F: Differentially expressed genes between the two CD8_Eff clusters.

1049

Figure 2:

A: Diagram illustrating the setup and data collection for the two sets of flow cytometry experiments

performed. Left Panel - BATs generated from Donor 1 challenged against MCF-7, CAMA1, and

T47D breast cancer cell lines. Samples collected at 0, 4, 24, 48, and 72 hours and designated '3

Cell Line Experiment'. Right Panel - BATs generated from Donors 1, 2, and 3 challenged

separately against MCF-7 breast cancer cell line. Samples collected at 0, 4, 24, 48, and 72 hours

and designated '3 Donor Experiment'.

B: Inhibitory panel receptor expression on CD8 (left panel) and CD4 (right panel) BATs. Data

shown was collected from the 3 Cell Line experiment and is presented as the mean value across

all cell lines. Error bars indicate standard error within each data group. Each line represents the

relative proportion of the cells that express a marker regardless of other marker expression.

C: Relative cell proportion of CD8 cells expressing LAG3, TIGIT, PD-1, or a combination thereof.

Cells were binned specificly by their expression or combination of expression of LAG3, TIGIT,

and PD-1. Unlike Panels B and D, there is no overlap between groups containing a common

ligand. For example, the TIGIT+ CD8 T cell subset does not express PD-1 or LAG3. Samples

collected at 0, 4, 24,48, and 72 hours. Data represents mean value across 3 Cell Line experiment

as collected in the coexpression panel. Error bars represent the range within each sample

timepoint.

D: Coexpression panel receptor expression on CD8 (left panel) and CD4 (right panel) BATs. Data

shown was collected from the 3 Cell Line experiment and is presented as the mean value across

all cell lines. Error bars indicate standard error within each data group. Each line represents the

relative proportion of the cells that express a marker and can have other markers simultaneously

expressed.

E: UMAP of BAT cells from the 3 Cell Line experiment clustered by receptor expression as

measured by flow cytometry in the inhibitory panel.

F: UMAP of BAT cells from the Cell Line experiment clustered by receptor expression as

measured by flow cytometry in the coexpression panel.

1079

1080

Figure 3:

A: OPLS-R model of rate of tumor change predicted by BAT subpopulations from the 3 Cell Line

experiment. BATs were binned by CD4 or CD8 expression, and further binned by expression of

LAG3, TIGIT, and/or PD-1. Cells expressing more than one receptor were binned in separate

categories, with cells expressing all three receptors being designated tri-positive (TP). BATs

lacking receptors were designated tri-negative (TN).

B: VIP scores generated by PLS-R model in Figure 3A. Subpopulation impact is designated by

the magnitude of the VIP score, with scores of magnitude >1 designated as greater-than-average

contribution of that cell subpopulation to tumor reduction. Subpopulations with scores colored tan

are associated with reduced rate of tumor change, while subpopulations with scores colored

brown are negatively associated with reduced rate of tumor change.

C: Scatter plots comparing the viable cell count of BAT subpopulations (x-axis) to rate of tumor

change (y-axis) for subpopulations with VIP scores >1. NCI = normalized cell index.

1094   D: Diagram of BAT developmental progression as assumed by the model. BATs are grouped as

1095   NBATs (LAG3-/TIGIT-), LBATs (LAG3+/TIGIT-), TBATs (LAG3-/TIGIT+), or DBATs

1096   (LAG3+/TIGIT+). PD-1 expression does not factor into grouping and can be present within any

1097   group. These four groups are used to govern tumor reduction in the models presented in Figures

1098   4 and 5.

1099

1100

Figure 4:

A: Diagram representing processes represented in the model. Tumor expansion parameters were

determined by fitting a sigmoidal curve to tumor growth data from control conditions in which BATs

were absent. BAT subgroup expansion was fit to individual splines to capture the count data

determined from flow cytometry. CaliPro was used to fit cytotoxicity rates for each BAT subgroup

to match the experimentally measured BAT and tumor cell counts. Data from the 3 Cancer

experiment was used in parameter determination and is shown in B and C.

B: MCF-7 (left), CAMA1 (center), and T47D (right) cell counts as measured by RTCA (black) and

as predicted by the model (blue). Optimal model prediction in sold blue line, while range of

predictions depicted by dotted blue lines.

C: MCF-7 (left), CAMA1 (center), and T47D (right) cell reduction by BAT phenotype as predicted by the model. LBAT (orange), NBAT (blue), TBAT (yellow), and DBAT (purple) cumulative reduction is shown for optimal conditions (solid line) and for total range of predictions (dotted lines).

Figure 5:

A: MCF-7 cell counts as measured by RTCA (black) and as predicted by the model (blue) when challenged by Donor 1 (left), Donor 2 (middle), and Donor 3 (right) BATs. Optimal model prediction in sold blue line, while range of predictions depicted by dotted blue lines.

B: MCF-7 cell reduction by BAT phenotype as predicted by the model when challenged by Donor 1 (left), Donor 2 (middle), and Donor 3 (right) BATs. LBAT (orange), NBAT (blue), TBAT (yellow), and DBAT (purple) cumulative reduction is shown for optimal conditions (solid line) and for total range of predictions (dotted lines).