GABA Neurons in the Amygdala play a central Role in Urocortin-3–Mediated Stress Suppression of Reproduction

Author Information

Junru Yu,[1,†] Saeed Farjami,[2,3,†] Kateryna Nechyporenko,[2,3,†] Xiao Feng Li,[1] Hafsa Yaseen,[1,4] Yanyan Lin,[1] Jinbin Ye,[1] Owen Hollings,[1] Ross de Burgh,[1] Baban Singh,[1] Kevin T. O'Byrne,[1,*] Krasimira Tsaneva-Atanasova,[2,3,5,*] Margaritis Voliotis[2,3,*]

Affiliations

[1]**Department of Women and Children's Health, School of Life Course and Population Sciences, King's College London, Guy's Campus, London SE1 1UL, UK, [2]Department of Mathematics and Statistics, [3]Living Systems Institute, [4]Biological Sciences, University of Missouri, Columbia, MO 65211-7400 and [5]EPSRC Hub for Quantitative Modelling in Healthcare, University of Exeter, Stocker Road, Exeter EX4 4PY, UK**

Abstract

Stress can disrupt menstrual cycles, cause infertility, and lead to other reproductive disorders. The posterodorsal medial amygdala (MePD) processes stress signals and regulates the gonadotropin-releasing hormone (GnRH) pulse generator through GABAergic inhibitory projections to the hypothalamus. However, how stress is processed in the MePD - especially involving its dense GABA and Urocortin-3 (UCN3) neurons - remains poorly understood.

In this study, we combine *in vivo* GRadient-INdex (GRIN) lens mini-endoscopic calcium imaging (to track neuronal activity), optogenetics, clustering analysis, and computational modeling to investigate MePD circuitry. Our findings reveal two anti-correlated GABA sub-populations in the MePD that dictate responses to both UCN3 neuron stimulation and restraint stress. Our computational modeling suggests that mutual inhibition between these GABA groups drives this anti-correlated activity and predicts how these interactions shape downstream responses to stimulation of GABA and UCN3 neurons.

We test these predictions using optogenetics and confirm that GABA neurons are critical for the transmission of UCN3 signals to regulate luteinizing hormone (LH) pulse frequency. Our study is the first to show how GABA neurons in the amygdala mediate stress effects on reproductive health, uncovering key neural mechanisms linking emotional and reproductive functions.

Introduction

Successful reproduction requires normal function of a neural oscillator, the gonadotropin-releasing hormone (GnRH) pulse generator, that drives the pulsatile release of gonadotropic hormones, luteinizing hormone (LH) and follicle stimulating hormone (FSH) to control ovulation and spermatogenesis. This oscillator comprises <u>K</u>isspeptin neurons co-expressing <u>N</u>eurokinin-B (NKB) and <u>Dy</u>norphin A (acronym KNDy) in the hypothalamic arcuate nucleus (ARC). The KNDy neural network not only provides the essential episodic stimulatory kisspeptin signal to the GnRH neurons, but receives diverse afferent inputs and hence is a hub for integrating various internal and external cues that regulates reproductive function[1,2]. Despite the marked heterogeneity of the myriad stressors, including acute versus chronic experimental stress paradigms, the common denominator is suppression of GnRH pulse generator frequency, which is associated with stress-related reproductive disorders and infertility [3–8].

The amygdala, a part of the limbic brain typically associated with emotions and anxiety, has strong projections to the KNDy system[1,2]. The medial amygdala, and more specifically its posterodorsal subnucleus (MePD), has been shown to be a robust upstream regulator of GnRH pulse generator frequency[9–11], pubertal timing[12,13], sociosexual behaviour[14,15], anxiety[14] and psychological stress-induced suppression of pulsatile LH secretion[16,17].

In the MePD, the majority of neurons are GABAergic, accounting for a substantial portion of its projections, many of which target important hypothalamic centers involved in reproduction[18–20], including the KNDy neurons and optogenetic stimulation of the latter suppress LH pulsatility[17]. Moreover, chemogenetic inhibition of GABA neurons in the MePD can reverse the inhibitory effect of psychogenic stress on LH pulsatility[17], suggesting an involvement of GABA signaling in the stress-induced suppression of pulsatile LH secretion. Although the precise way in which the MePD GABA neurons influence LH pulsatility is not fully understood, it has been suggested that they interact with

the stress neuropeptide urocortin-3 (UCN3) neurons to modulate or reinforce stress-induced

suppression of LH pulse frequency[16].

UCN3, part of the corticotropin-releasing factor (CRF) superfamily and selective for CRF type 2

receptors (CRFR2)[21], is highly expressed in the MePD[16,22]. Furthermore, UCN3 neurons in the

MePD mediate psychosocial stress-induced suppression of the GnRH pulse generator and

corticosterone secretion, thus play a central role in the interaction between the reproductive and

stress axes[16,23]. Previously we have shown that intra-MePD GABA antagonism can reverse UCN3-

induced LH pulse suppression[23], suggesting that GABA acts downstream of UCN3 signalling within

the MePD. However, the local neural circuitry within the MePD mediating stress related information

to the GnRH pulse generator remains to be established.

Here, we employed GRadient INdex (GRIN) lens mini-endoscopy to monitor and compare the

calcium activity of GABA neurons at single-cell resolution under acute psychological stress or

optogenetic stimulation of UCN3 neurons. Clustering analysis of the observed calcium transients

uncovered two distinct GABA sub-populations exhibiting anti-correlated activity. To explore the

functional role of these two sub-populations and unpick the mechanisms driving their anti-correlated

activity, we build upon our previously developed MePD circuitry mathematical modeling framework[11].

Our modeling reveals that mutual inhibitory interactions between the GABA populations are sufficient

to drive anti-correlated activity and predicts a complementary inhibitory effect of GABA and UCN3

signaling on the frequency of the GnRH pulse generator, which we tested by applying Cre/Flp

recombinase intersectional strategies to selectively target and manipulate heterogeneous MePD

neuronal populations[17]. Hence, by uncovering precise interactions between distinct sub-populations

of MePD GABA neurons and UCN3 neurons in transmitting stress signals to the GnRH pulse

generator, we provide critical and previously unknown insights into the neural circuitry driving stress-

induced reproductive dysfunction. This discovery significantly advances our understanding of how

emotional stress directly impacts reproductive health and identifies potential neural targets for

therapeutic intervention in stress-related reproductive disorders.

Results

Dynamic reconfiguration of MePD GABA neuronal networks during selective optogenetic stimulation of urocortin-3 (UCN3) neurons and restraint stress

To investigate how the posterodorsal subnucleus of the medial amygdala (MePD) processes stress signals and to understand the role of GABA and UCN3 neurons in this process, we employed *in vivo* calcium imaging using GRadient INdex (GRIN) mini-endoscopy. A cohort of six female UCN3-Cre::VGAT-Flpo transgenic mice were injected with a viral mixture containing pAAV-Syn-Flex-rc[Chrimson-tdTomato] and pAAV-EF1a-fDIO-GCaMP6s. This intersectional viral strategy enabled dual functionality: (i) optogenetically stimulating MePD UCN3 neurons via Chrimson; and (ii) detecting calcium activity in MePD GABA neurons using GCaMP6s (Fig. 1**A**). GRIN lenses were surgically implanted at the same location as the viral injection, enabling imaging of calcium activity with single-neuron resolution (Fig. 1**B-G**). The specificity of the AAV-Syn-Flex-rc[Chrimson-tdTomato] expression in the MePD was quantified by overlapping analysis with the UCN3 immunohistochemical staining and revealed that 95.06 ± 2.24% of tdTomato-positive neurons co-localized with the UCN3 immunohistochemical positive neurons (Supplementary Fig. 1). Further, 81.61 ± 6.03% of the total UCN3-immunoreactive neurons co-expressed Chrimson-tdTomato (Supplementary Fig. 1).

Two series of experiments were conducted to assess GABAergic responses under distinct conditions. First, to assess the role of UCN3 neurons in recruiting GABAergic response, baseline calcium activity from MePD GABA neurons was recorded for 2 min, followed by 2 min of recording during optogenetic stimulation of UCN3 neurons (Fig. 1**A**). Optogenetic stimulation was delivered in a discontinuous pattern, 5 seconds on, 5 seconds off, with a light-pulse frequency of 10 Hz during the 'on' periods (10-ms pulse width, 10 mW power, and 532 nm wavelength). GABA neurons with activity that mimics the pattern of optic stimulation, evidence that they are co-expressing UCN3, are removed from further analysis (see Methods and Supplementary Fig. 2). To further characterize the GABAergic response to acute stressors, baseline calcium activity of GABA neurons was recorded for 2 min, followed by 2 min of recording while the animals were exposed to restraint stress (Fig.

1**A**). The dynamic GABAergic response to optogenetic stimulation of UCN3 neurons and restraint

stress was summarized in functional terms using pairwise (neuron-neuron) association matrices,

constructed from Pearson correlation coefficients between neuronal trace pairs (see Methods).

These correlation matrices represent the MePD GABAergic response as a graph of interconnected

nodes (neurons) with edge weights corresponding to the strength of their pairwise functional

interactions (Fig. 1**H&I**)[24]. To quantify the effect of stress or UCN3 activation on functional

connectivity we calculated the normalized Riemannian distance between the correlation matrices

from baseline and stimulation/stress periods. Our analysis revealed significant alterations in the

functional connectivity of the GABAergic network during both UCN3 stimulation and stress relative

to control animals lacking channelrhodopsin expression (Fig. 1**J**). Furthermore, on average, stress

induces a significant increase in the population's time-averaged calcium activity in contrast to UCN3

stimulation; however, this effect diminishes over the 2-minute stress period (Fig. 1K). These results

indicate that the GABA population in the MePD plays a pivotal role in processing stress signals,

such as those induced by restraint stress. Furthermore, our findings suggest that UCN3 neurons

may serve as a key relay, driving functional reorganization within the GABAergic population to

support the transmission of stress-related information.

Figure 1: Functional connectivity of GABA neurons in posterodorsal medial amygdala (MePD) reconfigures in response to optogenetic stimulation of urocortin-3 (UCN3) neurons or acute restraint stress. (A) Schematic diagram of the experimental design for stimulating MePD UCN3 neurons and simultaneously imaging calcium activity from MePD GABA neurons. Heatmaps of calcium activity (ΔF/F) (left) in response to optogenetic stimulation of UCN3 neurons excluding GABA cells co-expressing UCN3 (see

Supplementary Fig. 2) and (right) during restraint-induced stress. **(B-G)** Representative photomicrographs of

the MePD from a UCN3-Cre::VGAT-Flpo mouse injected with pAAV-Syn-Flex-rc+pAAV-EF1a-fDIO-

GCaMP6s. **(B&E)** Red fluorescence (tdTomoato) labels UCN3 neurons expressing Chrimson-tdTomato,

(C&F) green fluorescence labels GABA neurons, and (D&G) red and green signals merged. **(H&I)** Functional

connectivity matrices during baseline and under UCN3 activation and stress were constructed by calculating

the pairwise Pearson's correlation between calcium time-traces. **(J)** Normalized Riemannian distance between

the correlation matrices from baseline and stimulation/stress periods. Each data point represents a single

experiment, with different animals color-coded; control animals are shown as squares, and channelrhodopsin-

expressing animals as circles. (**K**) Difference in time-averaged calcium activity between periods of UCN3

stimulation or stress and the baseline period. Scale bars: **B**-**D**, 200 µm; **E**-**G**, 50 µm.

141 Clustering analysis reveals two functionally distinct GABA sub-populations in the MePD

To better understand the functional organization of the GABA MePD network during optogenetic

stimulation of UCN3 neurons and under restraint stress, we performed clustering analysis on the

functional connectivity atrices we obtained from the calcium recordings of GABA neurons.

Clustering results consistently revealed two functionally distinct cell populations both during

optogenetic stimulation of UCN3 neurons and restraint stress (Fig. 2**A&C**). Under both conditions,

activity between the two clusters appeared highly anti-correlated as illustrated by two representative

examples depicted in Fig. 2**B&D**. Moreover, we find that clusters are significantly different in size

with the smallest cluster accounting on average for 35% of all cells (Fig 2E&F, Supplementary Fig.

3). The robustness of the above finding was also verified using the K-means clustering algorithm

and an alternative measure of association between neuronal traces that accounts for time-lag and

sign in neuronal interactions (see Methods and Supplementary Figs. 4-6).

Varying the frequency of optogenetic stimulation of UCN3 neurons influenced the functional

connectivity within the GABA network (Supplementary Fig. 7). Stimulation at 5 Hz produced no

significant change compared with controls, 10 Hz resulted in a statistically significant increase, while

20 Hz reduced the effect, consistent with a depolarization block. In contrast, stimulation at 5, 10, or

20 Hz did not significantly affect the correlation between the average activity of the two GABA

clusters (Supplementary Fig. 8). Based on the functional role of the MePD in relaying stress signals

to the hypothalamic arcuate nucleus (ARC), our results suggest that the identified GABA sub-

populations (clusters) could correspond to distinct populations of GABA interneurons and GABA

efferent neurons—defined as neurons that transmit signals from the MePD to downstream targets.

Known efferent projections to ARC play a role in regulating the gonadotropin-releasing hormone

(GnRH) pulse generator, a key driver of reproductive function[11,17].

Figure 2: **Clustering identifies two functionally distinct GABA sub-populations in the posterodorsal**

medial amygdala (MePD) both during UCN3 optogenetic stimulation and restraint stress.

Representative example of (**A**) a clustered connectivity matrix and (**B**) associated calcium time-traces of

MePD GABA neurons during optogenetic stimulation of UNC3 neurons. Representative examples of (**C**) a

clustered connectivity matrix and (**D**) associated calcium activity time-traces of MePD GABA neurons during

restraint stress. (**E&G**) Relative sizes of the two GABA sub-populations identified via clustering analysis. (**##**:

p<0.001).

172 Mutually inhibitory interactions between GABA populations drive anti-correlated activity

To explore the network mechanisms driving the anticorrelated activity of the GABA sub-populations,

we build upon our previously developed MePD circuit model[11]. The model allows us to explore

interactions between MePD neuronal populations and understand emergent patterns of dynamical

activity. Specifically, the model describes the calcium dynamics of two interacting populations of

GABA neurons, corresponding to interneurons and efferent projection neurons. Efferent neurons

project their axons outside the MePD to downstream targets such as the KNDy network. Both GABA

populations interact with a glutamatergic population (Fig. 3**A**). Experimental evidence shows that

antagonizing glutamate signaling completely blocks the suppressive effect of MePD UCN3

activation on LH pulsatility[23], highlighting a functional role of glutamatergic neurons in the circuit.

Therefore, the activity of the glutamatergic population is also included in the MePD model alongside

the two GABA populations. Model equations are provided in Supplementary Information, and the

parameter values for the MePD model are in Supplementary Tab. 1.

To incorporate UCN3 input we include a non-autonomous term in the model informed by recordings

of the activity of UCN3 neurons during optogenetic stimulation (see Fig. 3**A** and Methods). The

recorded traces exhibit a periodic pattern driven by the experimental stimulation protocol with 5

seconds on, 5 seconds off (see Methods). In the model, we approximated this UCN3 input using a

sinusoidal function with roughly 5 seconds on, 5 seconds off pattern, driving the populations GABA

interneurons and the glutamatergic neurons. To reproduce the anti-correlated activity between the

two GABA neuronal sub-populations while also reproducing previously published results on the

effects of neuropharmacological perturbations in the MePD on LH pulsatility[23], we find that we need

to include bi-directional (i.e. mutually inhibitory) GABA interaction in the circuit as well as GABA

interneurons self-inhibition (Fig. 3**A**).

Next, we validate our model by simulating the effects of stress on the MePD circuit. UCN3 neurons

in the MePD have also been shown to mediate psychosocial stress-induced suppression of the

GnRH pulse generator[16]. Experimentally recorded calcium activity of UCN3 neurons during stress

intervention is characterized by a fast rise followed by a slow decay (Fig. 3**B**), which we model by

taking the difference between two exponential functions, one with a fast and one with a slow time

constant (see Methods). Using this input, our mathematical model confirms the emergence of anti-

correlated behavior between the GABA sub-population whereby the activity of GABA interneurons

decreases over the stress period, while the activity of the GABA efferent neuron population

increases (Fig. 3**B**), supporting the proposed MePD circuit structure (Fig. 3**A**).

Figure 3: *In silico* simulation of the activity in the GABA neuronal populations in the MePD under the

increased excitatory input from UCN3 neurons and under restraint stress. Schematic on the left

represents the MePD neuronal circuit interacting with UCN3, consisting of two GABA populations and a

glutamatergic population. **(A)** Calcium time-traces of MePD UCN3 neurons during optogenetic stimulation of

UCN3 neurons and corresponding mean activity. Model simulations of the effects of UCN3 optogenetic

stimulation supports the emergence of anti-correlated activity in GABA sub-populations. **(B)** Calcium time-

traces of MePD UCN3 neurons during restraint stress and corresponding mean activity of the UCN3 calcium

traces during restraint stress with corresponding mean activity. Simulations of the MePD circuit under the

effects of stress also support anti-correlated GABA sub-populations' activity.

In silico **modeling reveals complementary relationship between MePD GABA and UCN3**

populations

Having shown that MePD model can reproduce anti-correlated activity of two GABA sub-

populations, we examine the effects of distinct interventions involving GABA and UCN3 in the MePD

on the LH pulses. To do that we couple the model of the MePD circuitry defined in the previous

section to our network model of the KNDy population in the arcuate nucleus (ARC)[11,25]. KNDy model

details are in Supplementary Information and the model parameter values can be found in

Supplementary Tab. 2. Since KNDy neuronal activity serves as a well-established proxy for LH

secretion, we use the dynamics of the KNDy network model to infer changes in LH pulsatility in

response to MePD circuit manipulations. Our previous in silico findings[11] show that MePD signal

integration is predominately mediated by GABA projections. The effects of MePD GABA and

glutamate antagonism alone and together with UCN3 stimulation on LH pulsatility have been

previously investigated[23], and we use these experimental findings to calibrate our coupled model

parameters by reproducing these effects accordingly (see Supplementary Fig. 9).

Next, using the calibrated model, we predict that providing a periodic excitatory input (see

Supplementary Fig. 10(A)), that mimics the effects of optogenetic stimulation of UCN3, leads to the

increase in the KNDy inter-pulse interval (IPI) from 15.18 min to 23.97 min due to increased activity

in the inhibitory GABAergic input to the KNDy network (Fig. 4A).

Since MePD has an overwhelming number of GABAergic outputs and interneurons, we are

interested in their role in information processing of stress signaling. To this end we provide periodic

excitatory input directly to both populations of GABA neurons (see Methods and Supplementary

Information). This is different to the indirect activation through UCN3 stimulation, as it leads to the

increased activity in both GABA subpopulations (Supplementary Fig. 10(B)). The resulting effect on

the KNDy IPI is an increase from 15.18 min to 42.65 min (Fig. 4B).

In the model we hypothesize that UCN3 signaling to the KNDy is operating through the GABAergic

pathway, hence disrupting GABAergic interactions will suppress UCN3's effects on KNDy. To

investigate it we suppress the strength of the GABAergic interactions by 50% and mimic UCN3

stimulation by providing excitatory input to the circuit. The model shows that this intervention does

not have a significant effect on KNDy IPI, apart from slightly increasing it from 15.18 min to 17.31

min (Fig. 4C & Supplementary Fig. 10(C)).

Taken together the above could be summarized as the following set of model predictions along with

suggested ways to test them experimentally:

1. Optogenetic stimulation modelled via excitation of the MePD UCN3 increases the KNDy IPI

15.18 min to 23.97 min. This could be tested experimentally via in vivo optogenetic activation

of the MePD UCN3 neuronal populations and simultaneous measurement of LH pulsatility.

2. Stimulation of the MePD GABA populations increases the KNDy IPI 15.18 min to 42.65 min.

Such a substantial change underscores the critical role of GABAergic signaling in modulating

KNDy pulsatility. Experimental confirmation could involve in vivo optogenetic activation of the

entire MePD GABA population accompanied by LH pulsatility measurement.

3. Stimulation of the MePD UCN3 with simultaneous suppression of the GABAergic interactions

in the MePD does not significantly affect KNDy IPI, changing it from 15.18 min to 17.31 min.

This finding suggests that UCN3 effects on the KNDy network are primarily dependent on its

interactions with GABA populations. A strategy to verify this hypothesis experimentally could

be to stimulate UCN3 non-GABA-expressing neurons in vivo while simultaneously inhibiting

non-UCN3 GABA neurons and measure LH pulsatility.

Finally, we investigate in silico the effects of stress on LH pulse dynamics mediated via the MePD.

In the restraint-stress experiment it was shown that one GABA cluster was active, while the other

one was inactive (Fig. 2D). By simulating these dynamics (Supplementary Fig. 9(D)) in the coupled

MePD - KNDy network model we are able to demonstrate that pulses in the KNDy network ceased

(Fig. 4D). These modeling findings align with published experimental evidence indicating that

restraint stress suppresses LH pulsatility[16].

Figure 4: Coupled MePD - KNDy model predictions for the effect of stress and stimulation of UCN3

neurons on GnRH pulse generator activity and hence LH pulsatility. (A) Simulation of the optogenetic

stimulation of the UCN3 neurons leads to the increase of the KNDy IPI 15.18 min to 23.97 min. (B)

Optogenetic stimulation of the MePD GABA neurons (providing excitatory input to both GABA populations)

increases KNDy IPI from 15.18 min to 42.65 min. (C) Optogenetic stimulation of the UCN3 neurons and

simultaneous inhibition of GABA neuronal populations do not significantly affect KNDy IPI, where it changes

from 15.18 min to 17.32 min. (D) Simulating the effects of stress leads to the cessation of the pulsatile

dynamics.

Selective optogenetic stimulation of UCN3 non-GABA neurons inhibits pulsatile LH secretion

in female mice.

We now test experimentally our modeling predictions detailed in the previous section regarding the

inhibitory effect of MePD UCN3 activation on LH pulsatility in vivo in freely behaving mice. To this

end a cohort of UCN3-Cre::VGAT-Flpo transgenic female mice was injected with an AAV-nEF-

Con/Foff 2.0-ChRmine-oScarlet viral construct. This intersectional approach enabled conditional

expression of channelrhodopsin (ChRmine) in UCN3 neurons not expressing GABA (UCN3 non-

GABA) (Con/Foff group). Following the viral injection, a unilateral fiber-optic cannula was implanted

into the MePD, above the injection site, to facilitate selective activation of the targeted neuronal

population. The specificity of the AAV-nEF-Con/Foff 2.0-ChRmine-oScarlet expression in the MePD

was quantified by overlapping analysis with the UCN3 immunohistochemical staining and revealed

that 98.92 ± 1.08% of the oScarlet-positive neurons co-localized with the UCN3

immunohistochemical positive neurons (Supplementary Fig. 11). Further, 54.18 ± 5.04% of the total

UCN3-positive neurons co-expressed ChRmine-oScarlet (Supplementary Fig. 11).

The stimulation protocol consisted of a 60-min control period, followed by 60 min of optogenetic

stimulation. Stimulation was delivered, as above, in a pulsed pattern (5 seconds on, 5 seconds off)

with a pulse frequency of 10 Hz during the on period (10-ms pulse width, 10 mW power, 532 nm

wavelength). Blood samples (5 μL) were collected every 5 min throughout the protocol and used to

measure LH levels.

Statistical analysis indicated post-stimulation LH inter-pulse intervals differed across the two groups

($F_{(2, 30)}$ = 15.83, P < 0.0001). In the Con/Foff group, the LH inter-pulse interval significantly

increased after optogenetic stimulation compared to the control period (15.47 ± 0.91 min vs. 23.75 ±

1.07 min, mean ± SEM, n = 6, p<0.01; Fig. 5A&D), validating the first of our modeling predictions. In

contrast, the no-stimulation group, which received the Con/Foff viral construct without optogenetic

stimulation, showed no significant changes in LH inter-pulse interval (15.84 ± 0.37 min vs. 16.68 ±

0.86 min, n = 6, p > 0.05; Fig. 5B&D). Similarly, in a separate cohort of mice injected with a control

virus (DIO-EYFP), optogenetic stimulation resulted in no significant differences in LH inter-pulse

intervals before and after stimulation (19.83 ± 1.02 min vs. 17.78 ± 0.93 min, n = 6, p > 0.05; Fig.

5C&D). The LH pulse amplitude and mean LH levels were quantified, and data are provided in

Supplementary Table 3.

304 Selective optogenetic stimulation of UCN3-expressing GABA neurons fails to affect pulsatile

305 LH secretion in female mice.

To further test the functional interactions between the UCN3 and GABA populations, a separate

cohort of UCN3-Cre::VGAT-Flpo transgenic female mice was injected with an AAV-nEF-Con/Fon

2.0-ChRmine-oScarlet viral construct, which allows optogenetic stimulation of UCN3 neurons co-

expressing GABA (UCN3-expressing GABA) (Con/Fon group). The specificity of the AAV-nEF-

Con/Fon 2.0-ChRmine-oScarlet expression in the MePD was quantified by overlapping analysis with

the UCN3 immunohistochemical staining and revealed that 98.77 ± 1.24% of the oScarlet-positive

neurons co-localized with the UCN3 immunohistochemical positive neurons co-expressing GABA

(Supplementary Fig. 12). Further, 32.73± 2.60% of the total UCN3-positive neurons co-expressed

ChRmine-oScarlet (Supplementary Fig. 12).

Statistical analysis showed no significant interaction effect of group and optogenetic stimulation on

LH inter-pulse intervals (two-way ANOVA, $F_{(1, 18)}$ = 0.2709, p = 0.6091). That is, LH inter-pulse

intervals in the Con/Fon group, showed no change following optogenetic stimulation compared to

the control period (17.92 ± 0.80 min vs. 16.60 ± 0.88 min, mean ± SEM, n = 6, p > 0.05; Fig. 5H&J).

Similarly, in the no-stimulation control experiment for this cohort, which did not receive optogenetic

stimulation, there was no significant difference in LH inter-pulse intervals (16.08 ± 1.45 min vs.

15.83 ± 0.95min, n = 5, p > 0.05; Fig. 5I&J). The LH pulse amplitude and mean LH levels were

quantified, and data are provided in Supplementary Table 3.

Taken together, our results suggest that both UNC3 and GABA populations relay inhibitory

information to the hypothalamic GnRH pulse generator but activation of UCN3 neurons that also co-

express GABA is not sufficient.

Figure 5: Effect of optogenetic stimulation of UCN3 non-GABA and UCN3 neurons co-expressing

GABA in the MePD on LH pulse frequency. (A-D) Optogenetic stimulation of UCN3 non-GABA neurons

inhibited LH pulses. Representative examples of LH pulsatility in UCN3-Cre::VGAT-Flpo mice injected with

AAV-Con/Foff-ChRmine in response to (A) 10Hz optic stimulation or (B) no stimulation as control, and (C) in

mice injected with control virus in response to 10 Hz stimulation. Pulses detected by the DynPeak algorithm

are indicated with an asterisk (*). (D) Mean LH inter-pulse interval (IPI) (±SEM) for each group during the

control period and over the subsequent stimulation period. #: p<0.01 vs stimulation period in Con/Foff group

(Two-way ANOVA, Tukey's post-hoc). (E-G) Representative photomicrographs of expression of AAV-nEF-

Con/Foff 2.0-ChRmine-oScarlet (red) in MePD UCN3 neurons. (H-M) Optogenetic stimulation of neurons co-

expressing UCN3 and GABA in the MePD did not impact LH pulse significantly. Representative examples of

LH pulsatility in UCN3-Cre::VGAT-Flpo mice injected with Con/Fon-ChRmine-oScarlet virus in response to (H)

10 Hz optic stimulation or (I) no stimulation as control. (J) Mean LH IPI (±SEM) for each group during the

control period and over the subsequent stimulation period (Two-way ANOVA). (K-M) Representative

photomicrographs of neurons co-expression UCN3 and GABA, tagged with oScarlet (red). Scale bars

represent (E&K) 200 µm, (F&L) 100 µm, (G&M) 50 µm. OT, optic track. The blue vertical bars in E to G and K

to M indicate the position of the fiber optic cannula.

The MePD GABA neurons relay information from UCN3 neurons and inhibit the GnRH pulse

generator.

To test our remaining model predictions and establish the functional role of GABA neurons in the

MePD in transmitting inhibitory signals to the hypothalamic GnRH pulse generator, we injected the

AAV-EF1a-DIO-ChR2-EYFP viral construct into VGAT-Cre transgenic mice, targeting the entire

MePD GABA neuron population for expressing ChR2 and used the optogenetic stimulation protocol

described above (60 min control period followed by 60 min stimulation) to stimulate the entire MePD

GABA population. In the VGAT-Cre-tdtomato mice, we observed that 93.27 ± 2.99% of the EYFP-

positive neurons were tdtomato-positive and 87.01± 5.21% of the tdtomato-positive neurons were

EYFP-positive. This indicates a high degree of co-expression consistency between the viral and

transgenic reporters. Statistical analysis showed a significant interaction effect of our genetic

intervention and stimulation period on LH pulsatility ($F_{(2,24)}$ = 10.55, p = 0.0005; two-way ANOVA).

Stimulation of the GABA population markedly increased the LH inter-pulse interval compared to the

pre-stimulation control period (18.5 ± 2.03 min vs 49.5 ± 8.46 min, mean ± SEM, n = 5, p<0.01;

Tukey's post-hoc; Fig. 6A&D), validating modeling prediction 2. In contrast, mice that did not receive

optogenetic stimulation showed no change in LH inter-pulse interval (21.25 ± 3.79 min vs 19.33 ±

1.52min, n = 5, p > 0.05; Tukey's post-hoc; Fig. 6**B&D**). Similarly, control virus (AAV-EF1a-DIO-

EYFP) injections did not result in any significant difference in LH inter-pulse interval before and

during stimulation (16.50 ± 0.67 min vs 18.00 ± 0.62min, n = 5, p > 0.05; Tukey's post-hoc; Fig.

6**C**&**D**). The LH pulse amplitude and mean LH levels were quantified, and data are provided in

Supplementary Table 4

Next, UCN3-Cre::VGAT-Flpo transgenic females were injected with an intersectional viral mixture

containing AAV-nEF-Con/Foff 2.0-ChRmine-oScarlet and pAAV-nEF-Coff/Fon-NpHR3.3-EYFP ,

enabling simultaneous excitation of UCN3 non-GABA neurons and inhibition of non-UCN3 GABA

neurons. Statistical analysis showed no significant interaction effect of genetic manipulation and

stimulation period on LH pulsatility ($F_{(1, 18)}$ = 0.131, p = 0.7216; two-way ANOVA). Moreover,

optogenetic stimulation resulted in no significant change in the LH inter-pulse interval relative to the

control period (16.67 ± 1.31 min vs 18.61 ± 1.58 min, mean ± SEM, n = 6, p > 0.05; Fig. 6**H**&**J**),

which aligns with our modeling prediction 3. Similarly, in mice injected with the same viral mixture

but without receiving optogenetic stimulation, no significant change in LH inter-pulse interval was

detected (16.33 ± 1.19 min vs 17.33 ± 0.81min, n = 5, p > 0.05; Fig. 6**I**&**J**). The LH pulse amplitude

and mean LH levels were quantified, and data are provided in Supplementary Table 3.

Collectively, these findings suggest that MePD GABA neurons are integral to regulating LH pulse

frequency. The inhibitory effect on LH pulses observed upon UCN3 neuron activation is attenuated

when GABA neuron activity is optogenetically suppressed, implying that GABA signaling acts

downstream of UCN3 neurons, relaying the inhibitory signal to the hypothalamic GnRH pulse

generator.

Figure 6: Selective optogenetic stimulation of GABA neurons in the MePD inhibits LH pulsatility and their activity is necessary for mediating UCN3 signalling downstream. (**A-G**) Optogenetic activation of the entire MePD GABA population in VGAT-Cre mice led to significant LH pulse suppression. Representative examples showing LH pulsatility in VGAT-Cre mice injected with AAV-double floxed-ChR2-EYFP virus in response to (**A**) 5 Hz stimulation or (**B**) no stimulation as control, and (**C**) 5 Hz stimulation in mice injected with control virus. (**D**) Mean LH IPI (±SEM) for each group during the control period and over the subsequent stimulation period. #: $p < 0.01$, †: $p < 0.05$ vs stimulation period in ChR group (Two-way ANOVA, Tukey's post-

hoc). (**E-G**) Representative photomicrographs of MePD GABA neurons, labeled with (**E**) tdTomato (red), (**F**)

ChR2-EYFP (green), and (**G**) merged. Scale bars represent. (**H-P**) The simultaneous inhibition of non-UCN3

GABA neurons in the MePD restored the LH suppression induced by stimulation of non-GABA UCN3

neurons. Representative examples showing LH pulsatility in UCN3-Cre::VGAT-Flpo female mice injected with

Con/Foff-ChRmine-oScarlet and Coff/Fon-NpHR-EYFP virus mixture in response to (**H**) 10 Hz optogenetic

stimulation or (**I**) no stimulation as control. (**J**) Mean LH IPI (±SEM) for each group during the control period

and over the subsequent stimulation period. (Two-way ANOVA). (**K-P**) Representative photomicrographs of

fluorescence in MePD neurons. (**K&N**) non-GABA UCN3 neurons tagged with oScarlet (red), (**L&O**) non-UCN

GABA neurons tagged with EYFP (green) (**M&P**) merged. Scale bars represent (**E-G**) 150 μm, (**K-M**) 200 μm,

(**N-P**) 100 μm. OT, optic track. The blue vertical bars in **E** to **G** and **K** to **P** indicate the position of the fiber

optic cannula.

Discussion

The posterodorsal subnucleus of the Medial Amygdala (MePD), is a critical regulator of reproductive

function. Single-cell transcriptomics has revealed a mosaic of glutamatergic and GABA populations

in this region, with GABA neurons co-expressing receptors for gonadal steroids as well as

neuromodulators and cognate receptors involved in parental and sociosexual behavior (including

kisspeptin, prolactin and oxytocin)[26]. Furthermore, calcium imaging of neuronal ensembles from this

region have started to decipher how this brain region encodes information about social interactions

and stressful stimuli[27], with stress related information being relayed to the reproductive axis via

potentially direct projections to the hypothalamus[1,2], and eliciting well-documented responses on

pubertal timing[12,13], LH secretion[16,17] and the GnRH pulse generator[9–11]. Here, to elucidate the

neuronal circuitry involved in MePD processing of stress signals, we used GRIN lens mini-

endoscopic calcium imaging to study, for the first time, the dynamic activity of MePD GABA neurons

under acute restraint stress as well as the activity of GABA neurons in response to UCN3 activation.

Functional network analysis of the imaging data revealed that acute restraint stress induces a

reconfiguration of the GABA population, highlighting their role in encoding and transmitting stress

information. Optogenetic stimulation of UCN3 neurons produced a comparable network reconfiguration,

despite no significant change in the magnitude of calcium activity compared to baseline, indicating that both

populations are co-involved in stress processing. Further clustering analysis of the calcium

transients identified two distinct GABA sub-populations with anti-correlated activity. This finding of

two distinct populations is consistent with physiological and morphological studies of GABA neurons

in the medial amygdala characterizing them as either projecting cells or local interneurons[18,20].

Building upon our previously developed computational model of the MePD circuit, we explored the

network connectivity to understand the possible mechanisms through which anti-correlated activity

emerges. The model consists of a glutamatergic sub-population and two GABA sub-populations:

one comprising GABA interneurons and the other GABA efferent projection neurons. Our analysis

identified two conditions necessary for the emergence of anti-correlated activity between these

GABA sub-populations, consistent with our experimental observations. First, mutual inhibition

between the GABA sub-populations is required, such that activation of one population suppresses

the other. In our mechanistic model, mutual inhibition arises from the biophysical characteristics of

GABAergic transmission, where the release of GABA inhibits target neurons, thereby producing

inhibitory interactions between the sub-populations. Second, if both sub-populations receive a

shared input (e.g., from the glutamatergic population or any other population of excitatory cells),

their dynamics become entrained to that common stimulus, meaning that distinct input streams to

each sub-population are necessary for the observed anti-correlation. To satisfy this condition in our

model, GABA interneurons receive inputs from UCN3 neurons, whereas the GABA efferent

projection neurons receive inputs from the glutamatergic population. This circuitry implies that stress

information is processed along two key pathways (one glutamatergic and one GABAergic) and is

integrated by the GABA efferent projection neurons.

We note that an alternative connectivity pattern, with the UCN3 population directly projecting to the

GABA efferent population, could also fit our experimental observations, but does not reproduce

previously published results on the effects of neuropharmacological perturbations in the MePD on

LH pulsatility[23]. Specifically, the suppression of glutamatergic neurons in combination with

stimulation of UCN3 neurons. Under this scenario direct input from stimulated UCN3 to GABA

efferent neurons will lead to the increase in its activity, consequently increasing KNDy IPI. This

finding contrasts with experimental observations, where the combined suppression of glutamate and

stimulation of UCN3 failed to produce any significant change in GnRH pulsatility[23].

We note that our MePD model has potential applications beyond understanding MePD dynamics

and their influence on the GnRH pulse generator. The model incorporates two functionally distinct

populations of GABA neurons (interneurons and efferent projection neurons), which reflect known

differences in GABA neurons within the amygdala[18,20]. This diversity likely originates from the medial

ganglionic eminence (MGE), an embryonic structure that gives rise to GABA interneuron

populations across various brain regions, including the MePD, through migration during

embryogenesis[28]. Therefore, the framework of our model could be extended to other neural circuits

where two functionally distinct GABA populations exist, such as the posterior ventral medial

amygdala[20] and its lateral and basal subdivisions[29].

Consistent with model predictions, we showed that optogenetic activation of non-GABA neurons,

expressing UCN3, suppresses LH pulse frequency. This finding corroborates previous work

demonstrating that LH pulses are inhibited by activation of UCN3 neurons[13,23] or administration of

UCN3[16] in the MePD, and that such effects are attenuated by chemogenetic inhibition of UCN3

neurons or by antagonizing corticotropin-releasing factor type 2 receptors (CRFR2)[16]. Thus, our

findings further support the central role of UCN3 signaling in modulating both stress responses and

reproductive function.

GABA is one of the primary inhibitory neurotransmitters in the nervous system[18]. Previous work has

shown that MePD GABA neurons project to the ARC to regulate LH pulses and chemogenetic

inhibition of MePD GABA neurons attenuates stress-induced suppression of LH pulses[17]. In this

study, we found that optogenetic activation of GABA neurons in the MePD leads to robust LH pulse

suppression, reinforcing the pivotal role of amygdala GABAergic signaling in stress-related

reproductive dysfunction. Antagonizing GABA receptors in the MePD can block UCN3-induced LH

suppression[23], suggesting that GABA neurons may function downstream of UCN3 signaling.

However, this pharmacological approach lacks cell-type specificity, leaving open the possibility that

UCN3 neurons themselves co-expressing GABA receptors, potentially contributing to the observed

effect. Here, by using an intersectional genetic strategy, we demonstrated that optogenetically

inhibiting non-UCN3 GABA neurons can abolish the change in LH pulse intervals induced by

optogenetic stimulation of UCN3 neurons. This finding suggests that these GABA neurons may be

acting downstream of UCN3 signaling to mediating LH pulse suppression. We note, that while in our

current experimental protocol we could also be blocking GABAergic projections to the ARC KNDy

population, which by itself could explain the absence of any effect from UCN3 stimulation, our

modeling predicts that GABAergic interactions within the MePD are additionally crucial for relaying

UCN3 signals to the KNDy network as also suggested by pharmacologically antagonizing GABA

receptors[23].

It is not unusual for neurons to co-express neuropeptides and neurotransmitters[20]. Given that

activating either UCN3 or GABA neurons independently suppresses LH pulses, one might predict

that stimulating neurons co-expressing both UCN3 and GABA would similarly inhibit LH. However,

we observed no significant changes in LH pulses upon optogenetic stimulation of neurons co-

expressing UCN3 and GABA. Our results indicate that UCN3 and GABA are both essential for

transmitting inhibitory signals to the hypothalamic GnRH pulse generator; however, this inhibitory

influence relies on their action through separate, functionally distinct neuronal populations. That is

activation of neurons that exclusively express either UCN3 or GABA suppresses the GnRH pulse

generator, whereas activation of UCN3 neurons that also co-express GABA fails to produce this

inhibitory effect. Within our modeling framework, UCN3/GABA co-expressing neurons might be

considered part of the GABA interneuron population, primarily inhibiting GABA projection neurons

rather than activating them. Consequently, they fail to relay the inhibitory UCN3 signal to the ARC.

Further studies are necessary to define more precisely the distinct role of UCN3/GABA co-

expressing neurons within the MePD circuitry.

By combining mini-endoscopic calcium imaging with computational modeling, we identified two

GABA sub-populations with complementary but distinct roles in encoding stress signals and

modulating LH pulsatility. Our experimental and computational results also stressed the functional

orthogonality between UCN3 and GABA signaling in MePD, highlighting the complexity of the

MePD's contribution to stress-induced reproductive suppression. These results offer improved

understanding of the neuronal circuits in the amygdala, a key hub for emotional processing, involved

in regulating hypothalamic GnRH pulse generator frequency[9–11,16,17,23] providing new insight into

psychogenic stress-related disorders of fertility, such as functional hypothalamic

amenorrhoea[6,7,30,31], and the development of novel therapeutic approaches to improve reproductive

health and well-being.

Methods

Animal

VGAT-Cre-tdTomato mice were genotyped using PCR to determine heterozygosity for VGAT-Cre

(primers 5′-3′: common, 12785—CTTCGTCATCGGCGGCATCTG; wild-type reverse12786—

CAGGGCGATGTGGAATAGAAA; mutant reverse oIMR8292—CCAAAAGACGGCAATATGGT) and

tdtomato (primers 5′-3′: wild-type forward oIMR9020—AAGGGAGCTGCAGTGGAGTA; wild-type

reverse oIMR9021—CCGAAAATCTGTGGGAAGTC; mutant reverse WPRE oIMR9103—

GGCATTAAAGCAGCGTATCC; mutant forward tdTomato oIMR9105—

CTGTTCCTGTACGGCATGG). UCN3-Cre heterozygous mice (strain Tg(Ucn3-cre)KF43Gsat/

Mmucd, MMRRC GENSAT) were crossbred with VGAT-Flpo heterozygous mice (Jax stock

#007909,B6.Cg-Slc32a1 2A-Flpo-D:Het) to obtain female double heterozygous UCN3-Cre::VGAT-

Flpo mice (Slc32a1 2A-Flpo-D:Het|Ucn3-Cre:Het|Rosa26<tomato>WT); genotyped using PCR for

the detection of heterozygosity (primers 5′- 3′: UCN3 Forward: CGAAGTCCCTCTCACACCTGGTT;

Cre Reverse: CGGCAAACGGAC-AGAAGCATT; Slc32a1 Mutant Forward: TGC ATC GCA TTG

TCT GAG TAG; Slc32a1 Mutant Reverse: GAC AGC CGT GAA CAG AAG G). Female mice, aged 8

to 10 weeks, weighing between 20 g and 25 g, were individually housed in ventilated cages sealed

with a HEPA-filter at 25 ± 1°C in a 12:12-hour light/dark cycle, lights on at 07:00 h. Vaginal smears

were performed when mice are 6 weeks old between 10:00 - 12:00 h daily. Only mice showing

regular estrous cycles were included. All animal procedures performed were approved by the King's

College London Animal Welfare and Ethical Review Body. Procedures were in accordance with UK

home office regulation.

524 Stereotaxic injection of adeno-associated virus construct and fiber-optic or mini-endoscope

525 GRIN lens implantation

All surgical procedures were carried out under general anesthesia using ketamine (Vetalar, 100

mg/kg, i.p.; Pfizer, Sandwich, UK) and xylazine (Rompun, 10 mg/kg, i.p.; Bayer, Leverkusen,

Germany) under aseptic conditions. Animals were secured in a David Kopf stereotaxic frame (Model

900, Kopf Instruments, CA, USA). Following bilateral ovariectomy, a midline incision was made in

the scalp to reveal the skull. Two small bone screws were inserted into the skull. A small hole was

drilled above the position of the MePD. The MePD coordinate (2.29 mm lateral, 1.60 mm posterior

from bregma, at a depth of 5.33 mm below the skull surface) was determined according to the

mouse brain atlas of Paxinos and Franklin[32]. A unilateral stereotaxic viral injection was performed to

target opsin or fluorescent protein expression in the MePD neurons, using a robot stereotaxic

system (Neurostar, Tubingen, Germany), with a 5-μL Hamilton microsyringe (Esslab, Essex, UK), at

the speed of 25 nl/min for 10 min.

Two transgenic animal models were used for LH blood sampling under optogenetic stimulation:

UCN3-Cre::VGAT-Flpo and VGAT-Cre-tdTomato. The following types of viral constructs were used

in separated cohorts of UCN3-Cre::VGAT-Flpo mice: (i) AAV-nEF-Con/Foff 2.0-ChRmine-oScarlet

(250 nl, $2.1×10^{12}$ gc/mL, Serotype:8, #137161, Addgene, Massachusetts, USA); (ii) AAV-nEF-

Con/Fon 2.0-ChRmine-oScarlet (250 nl, $1.9×10^{12}$ gc/mL, Serotype:8, # 137159, Addgene,

Massachusetts, USA); (iii) a mixture of AAV-nEF-Con/Foff 2.0-ChRmine-oScarlet ($2.1×10^{12}$ gc/mL,

#137161) and pAAV-nEF-Coff/Fon-NpHR3.3-EYFP ($1.9×10^{12}$ gc/mL; #137159, final volume 300 nl);

(iv) control virus AAV-Ef1a-DIO-EYFP (250 nl, $4.4 × 10^{12}$ gc/mL, Serotype:9, #27056, Addgene,

Massachusetts, USA). VGAT-Cre-tdTomato mice received unilateral injections of (i)

channelrhodopsin (ChR2) viral construct, (250 nl, $1.8 × 10^{12}$ gc/mL; Serotype:9; Addgene, #20298,

Massachusetts, USA) or (ii) control virus, AAV-Ef1a-DIO-EYFP (250 nl, 4.4 × 10^{12} gc/mL,

Serotype:9; Addgene, # 27056, Massachusetts, USA) into the MePD. Following injection, the needle

was kept in position for 10 min before going up slowly over 1 min. A unilateral fiber optic cannula

(200 µm, 0.39 NA, 1.25 mm ceramic ferrule; Doric Lenses, Quebec, Canada) was inserted into the

right MePD 0.15 mm above the injection site. After reaching the target, the fiber optic cannula was

secured on the skull using dental cement (Super-Bond Universal Kit, Prestige Dental, UK), bonded

to the screws and skull. Finally, the incision in the scalp was closed with suture. Animals were given

1 week to recover, after which they were handled 20 min daily to acclimatize to experimental

procedures for a further 2 weeks.

For mini-endoscope *in vivo* calcium imaging, only UCN3-Cre::VGAT-Flpo transgenic animals were

used. Separate mice cohorts received the following viral injection into the MePD: (i) a mixture of

pAAV-Syn-Flex-rc[Chrimson-tdTomato] (2.0×10^{12} gc/mL, Serotype:5, #62723, Addgene,

Massachusetts, USA) + pAAV-EF1a-fDIO-GCaMP6s (2.3×10^{12} gc/mL; Serotype:8; #105715,

Addgene, Massachusetts, USA; final volume 400 nl); (ii) a mixture of pAAV-Syn-Flex-rc[Chrimson-

tdTomato] (2.0×10^{12} gc/mL, Serotype:5, #62723, Addgene, Massachusetts, USA) +

pAAV.CAG.Flex.GCaMP6s.WPRE.SV40 (4.3×10^{12} gc/mL; Serotype:9; # 100842, Addgene,

Massachusetts, USA; final volume 400 nl); (iii) a mixture of control virus pAAV-hSyn-DIO-mCherry

(2.3×10^{12} gc/mL, Serotype:9, #50459) + pAAV-EF1a-fDIO-GCaMP6s (2.3×10^{12} gc/mL, Serotype:8,

#105715, final volume 400nl) . One week after administering the virus, a 600-µm diameter, 7.2-mm

long gradient index (GRIN) ProView™ Integrated Lens (Inscopix, PN:100-004554) was implanted at

the same location as the viral injection. The surface of the skull was scored, using the tip of a

surgical scalpel, and four screws were inserted into the skull. This was done to enhance the

adhesion of dental cement to the skull, thereby reducing the interference of movement on *in vivo*

recording. Before insertion of the GRIN lens, a 23 G needle was slowly lowered at a speed of 5.0

mm/min into the brain (2.42 mm lateral, 1.52 mm posterior from bregma, depth of 5.05 mm below

the skull surface) to make a needle track to facilitate the insertion of the GRIN lens without causing

brain distortion along its route. The GRIN lens was then inserted into the brain at a rate of 0.6

mm/min, using the same coordinates as the needle precutting. To secure the GRIN lens with

integrated baseplate in position, we used a light curing adhesive (iBond Universal, Kulzer GmbH,

Leipziger, Germany) along with a cured radiopaque hybrid dental cement (Gradia Direct Flo, GC

corporation, Tokyo, Japan), bonding it to the head screws and skull. A GRIN lens baseplate dust

cover (Inscopix) was rescued in place to protect the lens. After 4 weeks the mice were re-

anaesthetized as described above and underwent bilateral ovariectomy.

Serial blood sampling for LH measurement

The mouse's tail-tip was excised using a sterile scalpel. Then the chronically unilaterally implanted

fiber optic cannula was connected to a multimode fiber optic rotary joint patch cables (Thorlabs Ltd,

Ely, UK) via a ceramic mating sleeve to allow the animal freedom of movement. Mice were left for 1

h to habituate. A Grass SD9B stimulator-controlled DPSS laser (Laserglow Technologies, Toronto,

Canada) was utilized to deliver either green light (532 nm wavelength, 10 Hz, 5 seconds on 5

seconds off, 10 mW, 10-ms pulse width,) or blue light (473 nm wavelength, 5 Hz, 5 seconds on 5

seconds off, 5 mW, 10-ms pulse width,). The light pattern was controlled using software designed in

house. After 1 h of acclimatization, serial blood samples (5-µL) for LH pulse measurement were

collected every 5 min for 2 h between 10:00 - 12:00 h: 0-60 min for control blood sampling period,

and 60-120 min under stimulation or equivalent no-stimulation control period. Blood samples were

diluted in 45 µL of Phosphate-Buffered Saline (PBS) containing 0.2% Bovine Serum Albumin (BSA)

and 0.05% Tween-20 (PBST) and immediately placed on dry ice and stored at −80°C until later

analysis.

LH assay

LH measurement was performed using an LH ELISA assay as reported previously[33]. A mouse LH

standard (mLH; AFP-5306A, NIDDK-NHPP, USA) was used to establish a standard curve for the

quantitative measurement. The following antibodies were used in the assay: (i) coating antibody

(RRID: AB_2665514, monoclonal antibovine LH beta subunit antiserum, 518B7, University of

California, CA, USA); (ii) anti-LH antibody (RRID: AB_2665533; National Hormone & Peptide

Program, CA, USA); (iii) secondary antibody (RRID: AB_772206, GE Healthcare, Chicago, Illinois,

USA). The inter-assay and intra-assay variations were 10.53% and 5.74%, respectively. The assay

sensitivity was 0.023 ng/mL.

LH pulse statistical analysis

LH pulses were verified using the Dynpeak algorithm in Scilab 5.5.2 programme[34]. The Dynpeak

algorithms were classified for OVX mice based on previously established parameters[35]. The average

LH inter-pulse interval was calculated for the 1-h control period and the 1-h stimulation period in both

experimental and control mice. Statistical significance was assessed using a 2-way ANOVA unless

otherwise stated. Data was represented as mean ± SEM and $p<0.05$ was considered significant.

Mini-endoscope-based Calcium imaging

Under optogenetic stimulation of MePD UCN3 neurons. Prior to recording, animals were

handled daily for 5 to 6 weeks to get accustomed to the experimental setup. The Data Acquisition

(DAQ) box (Inscopix, PN:100-005904) was turned on and connected to the computer wirelessly.

The Data Acquisition Software (IDAS; Inscopix, Inc.) was accessed through the Chrome web

browser on the computer communicating with the DAQ box. Adequate DAQ box storage capacity

was checked routinely before Calcium imaging. The GRIN lens baseplate dust cover was gently

removed from the mouse's head, after which the minscope camera was secured to the baseplate

platform. Equipped with the head-mount miniscope (nVoke2, Inscopix, SN: BB-11238601), the

mouse was then placed back into its home cage. The optimal focal plane was chosen by adjusting

the lens focus to achieve clear cell morphology. The exposure time was fixed at 50 ms. The frame

rate was set to 20 Hz. Subsequently, gain, and LED power were optimized based on image

histograms before initiating recording. Two types of LED were used during the recording: (i) EX-LED

(0.2-0.4 mW, 435-460 nm excitation filter) for GCaMP6s signal detection in the GABA or UCN3

neurons; (ii) OG-LED (10 mW, 590-650 nm excitation filter) for Chrimson optogenetic stimulation of

the UCN3 neurons. After the mice were attached to the miniscope, they were given 1-2 h for

adaptation. For stimulation of the UCN3 neurons, the OG-LED was set to 10 Hz, 5 seconds on 5

seconds off, and a 10-ms pulse width. Once the animals had acclimatized, the GCaMP6s recording

of the GABA neurons or UCN3 neurons was started with a protocol of 2 min baseline before

stimulation for 2 min. Optogenetic stimulation was randomized for the 10 Hz, with a 5-min interval

between each to prevent photobleaching of GCaMP6s. Once the experiment was concluded, the

miniscope was gently detached, and the baseplate cover was reattached and secured with its

screw. The recordings were performed between 9:00 h and 15:00 h. Three groups of animals were

included in this experiment: (i) mice with UCN3 neurons expressing opsin and GABA neurons

expressing GCaMP; (ii) mice with UCN3 neurons expressing both opsin and GCaMP, serving as a

positive control; (iii) mice with UCN3 neurons expressing mCherry and GABA neurons expressing

GCaMP, serving as a negative control.

Under restraint stress. The system was set up and the mouse was connected to the miniscope as

described above but with the OG-LED turned off. Once the mouse had settled, the experiment

started with a 3 min baseline recording, followed by 3 min of recording under hand-held restraint.

Finally, the miniscope was detached and the baseplate cover was reattached. The experiments were

performed between 9:00 h and 15:00 h. The animals in this experiment were the same as those

used in the optogenetic stimulation protocol described above but were divided into two groups

depending on whether GCaMP was expressed in the UCN3 or GABA neurons, regardless of

whether opsin or mCherry was expressed in the UCN3 cells.

In vivo **calcium imaging recording processing**

The video data, with a resolution of 1280 pixels x 800 pixels, was exported from the DAQ box

through the file manager to a desktop computer and processed with the Inscopix Data Processing

Software (IDPS). Within the IDPS, there are six essential steps for extracting single-cell calcium

transient data from the recorded video: down-sampling, cropping, spatial filtering, motion correction,

trace normalization ($\Delta F/F$), and cell identification. First, the video underwent both spatial down-

sampling and temporal down-sampling by a factor of 2 and was cropped to enhance processing

speed, focusing on the region of interest where the GCaMP signal was observed. Spatial filtering,

with a threshold ranging between 0.005 (low) and 0.5 (high) was used to distinguish individual pixels

in the video image, creating a high-contrast image. A mean intensity projection image was

generated. Motion correction was subsequently conducted, using the mean intensity projection

image as a reference. Motion correction was applied again using one of the frames affected by

movement while simultaneously drawing a region of interest (ROI) where UCN3 accumulates to

enhance the effectiveness of motion correction. For the recordings with GCaMP signals from GABA

neurons, we did not use the built-in motion correction function of IDPS. Instead, we used the

MATLAB (version R2022a) implementation of the open-source algorithm NoRMCorre[36] with default

parameters for both rigid and non-rigid (patch size 64 pixels and overlap 32) motion correction as

long as a minimum correlation of the frames with the mean frame above 0.8 is achieved.

Occasionally, if the minimum correlation is not met, an extra step of non-rigid motion correction with

coarser patches is applied. This is performed by doubling the patch size and their overlap to 128 and

64, respectively. Corrected videos were saved in TIFF format for further analysis via IDPS. Following

motion correction, pixel intensity, $F(x,y)$, in each frame of the movie was normalised using the

following formula $\dfrac{(F(x,y) - F_{baseline}(x,y))}{F_{baseline}(x,y)}$, where $F_{baseline}(x,y)$ is the baseline pixel intensity calculated

by taking the mean value of the pixel across the entire movie. Finally, regions of interest were

identified automatically using the cell identification algorithm provided by IDPS, based on principal

component and independent component analysis (PCA-ICA algorithm).

Validation of virus injection, fiber optic cannula and GRIN lens implantation

Once the experimental procedures were completed, a lethal dose of ketamine was used to

humanely euthanize the mice. The mice were perfused transcardially with heparinized saline (5

U/ml) for 5 min, followed by ice-cold 4% paraformaldehyde (PFA) for 15 min using a pump

(Minipuls,156 Gilson, Villiers Le Bel, France). The brains were collected and post-fixed sequentially

at 4°C in a solution of 15% sucrose in 4% PFA and later in a solution of 30% sucrose in phosphate

buffer until they sank. They were then snap frozen on dry ice and stored in -80°C until processing.

The brains were coronally sectioned (30 µm) between +0.5 mm and -2.7 mm from the bregma using

a cryostat (Bright Instrument Co., Huntingdon, UK), and sections were mounted on microscope

slides, air-dried, and covered with slips using Prolonged Antifade mounting medium (Molecular

Probes, Inc. OR, USA). The accuracy of injection sites, fiber optic cannula and GRIN lens

implantation were confirmed by Axioskop 2 Plus microscope equipped with AXIOVISION 4.7

(Zeiss). Images were captured using Axioskop 2 Plus microscope (Carl Zeiss). Only the data of

mice with correct viral infection and implantation of cannula and lens in the MePD were included in

the analysis.

Immunohistochemistry

To assess the targeting of viral labeling to UCN3 neurons in the MePD, we used the indirect

immunofluorescence technique to localize UCN3 neurons. Every 4[th] section from each brain was

washed in potassium phosphate-buffered saline (KPBS) and incubated in blocking solution (KPBS

containing 1% BSA, 0.03% Triton X-100 and 5mg/ml heparin) at room temperature for 1 hour

followed by incubated in a rabbit polyclonal anti-UCN3 antibody (1:2000 dilution)[37] in the blocking

solution (Code: PBL 7218, kindly provided by J. Vaughan, Salk Institute, La Jolla, CA, USA) for 48

hours at 4°C. The brain sections were then processed with goat anti-rabbit IgG (H + L), a

biotinylated secondary antibody (1:1000 dilution; Cat. No. BA-1000, Vector Laboratories,

Burlingame, CA, USA) overnight at 4°C. After rinses in KPBS, visualization of UCN3 immunoactivity

was achieved using Streptavidin, Alexa Fluor 405 conjugate (1:200 dilution; Cat. No. S 32351,

Thermo Fisher Scientific, Waltham, MA, USA) for 2 hours at room temperature. The sections were

finally washed, mounted onto slides and cover slipped. Omission of the UCN primary antibody and

its pre-incubation resulted in the absence of specific staining. Brain sections from each experimental

group were processed on the same day to control for inter-batch variability. To preserve the

fluorescence of the virally expressed proteins, all subsequent steps were performed in a dark box.

Semiquantitative analysis of immunostaining data was carried out on AxioVision microscope image

system. UCN3-positive neurons were identified by Alexa Fluor 405 fluorescence, while native

fluorescence (e.g. tdTomato or oScarlet) was used for virus-transduced neurons identification. The

number of virus-labeled, UCN3-immunoreactive cells and co-localized cells in the MePD were

quantified on 4 sections from each mouse brain. Targeting efficiency was defined as the percentage

of virus-labeled cells that were also positive for UCN3.

Data analysis

The data processing protocol we employ in this study was performed in two phases: pre-intervention

and intervention (either via optogenetic stimulation or restraint stress). For our analysis, we use the

smoothed deconvolved traces generated by the OASIS method[38], which improves the signal-to-

noise ratio (SNR) and mitigates noise interference. All traces are rescaled between 0 and 1 (min-

max normalization), and then further normalized to mean zero and unit variance. The analysis

process is divided into three key steps: i) identification of GABA cells co-expressing UCN3 and

removing them from further analysis; ii) estimating and comparing functional connectivity before and

during intervention; iii) clustering analysis based on pairwise association-derived dissimilarity scores.

In step (i), we use power spectral density (PSD) analysis and K-means clustering (using Euclidean

distance) to identify GABA neurons with activity that mimics the pattern of optic stimulation, evidence

that they are co-expressing UCN3.

PSD filtering. For each trace, we compute the PSD during the stimulation phase. Cells with PSD

720 values at 0.1 Hz (the frequency of the stimulus 5 seconds on and 5 seconds off cycle) above 0.37

721 Hz^{-1} are labelled as co-expressing UCN3. This threshold is determined from recordings of UCN3

722 cells under optic stimulation, by calculating the median of the minimum PSD values at 0.1 Hz from

723 each dataset (see Supplementary Fig. 2**B**).

K-means clustering. Using the Euclidean distance metric, we cluster the traces during the

stimulation phase into two groups, aiming to separate neurons into GABA-only neurons and neurons

co-expressing UCN3 and GABA. Clustering separated a small group of neurons with activity

mimicking the optic stimulation oscillatory pattern (see Supplementary Fig. 2**C**). Recordings of MePD

UCN3 neurons shows that under optogenetic stimulation their activity rises and declines in phase

with on and off periods of stimulation (shown in Fig. 3**A**); therefore, this small number of cells was

labelled as co-expressing UCN3.

The union of neurons identified by either method as co-expressing UCN3 and GABA are excluded

from further analysis. By applying both filters, we aim to robustly identify purely GABA-expressing

neurons. We acknowledge the possibility that some cells may pass through the filters or that a few

purely GABA-expressing neurons may be inadvertently excluded. However, our dual approach of

employing power spectral density (PSD) analysis and K-means clustering minimizes this possibility

and is essential to ensure our analysis remains focused exclusively on GABA-expressing neurons.

Nevertheless, we take a further step and visually inspect the calcium recording traces for any

potential anomalies such as regular oscillations in-phase with the stimulation and/or massive

increase in calcium level during the intervention period. Any identified anomalies of this sort are also

dropped from data sets.

Comparing pre-intervention and intervention cell-cell interaction strength

Following step (i), the remaining dataset contains only traces associated with GABA neurons, which

are not directly activated by our optic stimulation protocol. In step (ii), we assess the functional

connectivity between the observed neurons using Pearson's correlation[39] defined as

745
$$corr(x,y) = \frac{R(x,y)}{\sqrt{R(x,x)R(y,y)}},$$

where $R(x,y) = \sum_{i=1}^{N} (x_i - \bar{x})(y_i - \bar{y})$ represents cross-correlation with N as the length of x and y.

Using the above measures, we construct the corresponding cross-correlation matrices (see for an

example Fig. 1**H&I**).

To evaluate changes in pairwise cell interactions between the pre-intervention and intervention

phases, we compute the pairwise Riemannian distance[40] between correlation matrices from pre-

intervention and intervention episodes. The Riemannian distance here is defined as

752
$$d(A, B) = \left(\sum_i ln(\lambda_i)^2 \right)^{1/2}$$

where ln is the natural logarithm and λ_i indicates the joint eigenvalues of symmetric positive definite

(SPD) cross-correlation matrices A and B. The size of the matrices directly affects the riemannian

distance such that the distance among matrices increases as the size increases. With a good

approximation square root describes this relationship. However, we take a more accurate approach

and normalize all the Riemannian distances by the mean pairwise Riemannian distance between

many SPD matrices generated randomly. To make sure that the random generation of correlation

matrices is uniform, we employ the Metropolis-Hasting method introduced in[41]. Normalizing the

Riemannian distances guarantees that the distance is not influenced by matrix size.

Clustering analysis

In step (iii), we apply agglomerative hierarchical clustering (using the average linkage criterion;

scikit-learn package in Python™) to classify cells based on the dissimilarity of their calcium activity,

given as

765
$$D(x, y) = 1 - corr(x,y).$$

To determine the optimal number of clusters and robustness of clustering, we employed a

bootstrapping technique. This involved resampling (without replacement) 75% of the cells and

clustering the resulting dataset with different numbers of clusters (2, 3, and 4) using hierarchical or

K-means clustering with either the covariance or corr as the dissimilarity measure. This resampling

process was repeated 100 times.

For each resample dataset, we evaluate the clustering performance, using the Silhouette Coefficient

(SC)[42] which measures the cohesion within clusters and their separation, with values ranging from -1

(poor performance) to 1 (good performance). Results of the evaluation are given in Supplementary

Figs. 4-6, showing that the Silhouette Coefficient is significantly higher for 2 clusters compared to 3

or 4 clusters in the case of both hierarchical and K-means clustering.

Modeling

777 We build on our previously published modelling framework of MePD circuit[11] by incorporating

778 dynamic stimulation by UCN3 as observed experimentally (as shown in Fig. 3). To this end we

779 define UCN3 input into the populations of glutamatergic neurons and GABA interneurons with a

780 strength given by the parameter γ. The activity of UCN3 during stimulation is approximated via

781 sinusoidal function:

782
$$U_{stim}(t) = \gamma B + A sin sin (2\pi f t) ,$$

783 where B and A are the baseline UCN3 activity and the amplitude, respectively, while the parameter

784 f is the frequency of the sigmoid wave. We approximate UCN3 activity during restraint stress using

785 two exponential functions:

786
$$U_{stress}(t) = \gamma(atexp(-r_2 t) + b(1 - exp(-r_1 t))),$$

where the function initially increases with magnitude a then decreases to a plateau value given by

the parameter b. The parameters r_1 and r_2 are responsible for the rate of increase of the first term

and rate of decrease to a plateau in the second term.

Briefly, the model captures the dynamics of the MePD intracellular calcium activity of three

populations: glutamatergic neuronal population (excitatory) and the populations of GABA

interneurons and GABA efferent neurons (inhibitory). We build on our previous work[11] by assuming

that excitatory (UCN3) input is provided to the populations of GABA interneurons and glutamate (Fig

3). Previous analysis of the model indicated that negative feedback between GABA and

glutamatergic neuronal populations as well as glutamatergic self-excitation are critical mechanisms

for inducing oscillatory dynamics[11]. However, providing glutamatergic inputs to both populations of

GABA neurons result in a null phase difference between the populations. Therefore, we opted to

provide glutamatergic input only to the population of GABA efferent neurons.

To investigate how the amygdala modulates the activity in the GnRH pulse generator, we couple the

above-described MePD model to our KNDy network model, where the oscillatory dynamics of the

KNDy acts as a proxy of LH pulsatility[25]. Here we couple the model via providing only inhibitory

output from MePD GABA efferent neurons population to KNDy, as it was previously shown that

glutamate plays a modulatory role to the MePD output rather than directly affecting the KNDy

network[11]. We then calibrate the coupled by reproducing the experimental results from Ivanova et

al.[23] (Supplementary Fig. 8). Further modeling details, such as equations governing model

dynamics and model parameter values, can be found in Supplementary Information. The simulations

have been done in MATLAB (version R2023b).

Data Availability

809 Source data are provided with this paper. Further information that supports the findings of this study

810 is available from the corresponding authors upon request.

Code Availability

812 The code for reproducing the data analysis (Python™), mathematical modeling (MATLAB), and

813 figure generation for both the main text and Supplementary information, along with a test dataset, is

814 publicly available at: https://github.com/mv-kr/MePD_GABA.

References

1. Yeo, S.-H., Kyle, V., Blouet, C., Jones, S. & Colledge, W. H. Mapping neuronal inputs to Kiss1

neurons in the arcuate nucleus of the mouse. *PLOS ONE* **14**, e0213927 (2019).

2. Moore, A. M., Coolen, L. M. & Lehman, M. N. Kisspeptin/Neurokinin B/Dynorphin (KNDy) cells

as integrators of diverse internal and external cues: evidence from viral-based monosynaptic

tract-tracing in mice. *Sci. Rep.* **9**, 14768 (2019).

3. Li, X. F. & O'Byrne, K. T. Stress and the Reproductive System. in *Knobil and Neill's Physiology*

of Reproduction: Two-Volume Set vol. 2 1637–1660 (ELSEVIER SCIENCE INC, 2014).

4. Yu, J., Li, X.-F., Tsaneva-Atanasova, K., Zavala, E. & O'Byrne, K. T. Chemogenetic activation of

PVN CRH neurons disrupts the estrous cycle and LH dynamics in female mice. *Front.*

Endocrinol. **14**, 1322662 (2023).

5. Kreisman, M. J., Tadrousse, K. S., McCosh, R. B. & Breen, K. M. Neuroendocrine Basis for

Disrupted Ovarian Cyclicity in Female Mice During Chronic Undernutrition. *Endocrinology* **162**,

bqab103 (2021).

6. Bethea, C. L., Centeno, M. L. & Cameron, J. L. Neurobiology of Stress-Induced Reproductive

Dysfunction in Female Macaques. *Mol. Neurobiol.* **38**, 199–230 (2008).

7. Herod, S. M., Pohl, C. R. & Cameron, J. L. Treatment with a CRH-R1 antagonist prevents

stress-induced suppression of the central neural drive to the reproductive axis in female

macaques. *Am. J. Physiol. Endocrinol. Metab.* **300**, E19-27 (2011).

8. Nair, B. B. *et al.* Impact of chronic variable stress on neuroendocrine hypothalamus and pituitary

in male and female C57BL/6J mice. *J. Neuroendocrinol.* **33**, e12972 (2021).

9. Comninos, A. N. *et al.* Kisspeptin signaling in the amygdala modulates reproductive hormone

secretion. *Brain Struct. Funct.* **221**, 2035–2047 (2016).

10. Lass, G. *et al.* Optogenetic stimulation of kisspeptin neurones within the posterodorsal medial

amygdala increases luteinising hormone pulse frequency in female mice. *J. Neuroendocrinol.*

32, e12823 (2020).

11. Nechyporenko, K. *et al.* Neuronal network dynamics in the posterodorsal amygdala: shaping

reproductive hormone pulsatility. *J. R. Soc. Interface* **21**, 20240143 (2024).

12. Adekunbi, D. A. *et al.* Role of amygdala kisspeptin in pubertal timing in female rats. *PLoS ONE*

12, e0183596 (2017).

13. Ivanova, D. *et al.* Role of Posterodorsal Medial Amygdala Urocortin-3 in Pubertal Timing in

Female Mice. *Front. Endocrinol.* **13**, (2022).

14. Adekunbi, D. A. *et al.* Kisspeptin neurones in the posterodorsal medial amygdala modulate

sexual partner preference and anxiety in male mice. *J. Neuroendocrinol.* **30**, e12572 (2018).

15. Wu, Y. E. *et al.* Neural control of affiliative touch in prosocial interaction. *Nature* **599**, 262–267

(2021).

16. Ivanova, D. *et al.* Urocortin3 in the Posterodorsal Medial Amygdala Mediates Stress-induced

Suppression of LH Pulsatility in Female Mice. *Endocrinology* **162**, 1–14 (2021).

17. McIntyre, C. *et al.* GABA Signaling in the Posterodorsal Medial Amygdala Mediates Stress-

induced Suppression of LH Pulsatility in Female Mice. *Endocrinology* **164**, 1–11 (2023).

18. Keshavarzi, S., Sullivan, R. K. P., Ianno, D. J. & Sah, P. Functional Properties and Projections

of Neurons in the Medial Amygdala. *J. Neurosci.* **34**, 8699–8715 (2014).

19. Choi, G. B. *et al.* Lhx6 delineates a pathway mediating innate reproductive behaviors from the

amygdala to the hypothalamus. *Neuron* **46**, 647–660 (2005).

20. Bian, X. Physiological and morphological characterization of GABAergic neurons in the medial

amygdala. *Brain Res.* **1509**, 8–19 (2013).

21. Lewis, K. *et al.* Identification of urocortin III, an additional member of the corticotropin-releasing

factor (CRF) family with high affinity for the CRF2 receptor. *Proc. Natl. Acad. Sci. U. S. A.* **98**,

7570–7575 (2001).

22. Shemesh, Y. *et al.* Ucn3 and CRF-R2 in the medial amygdala regulate complex social

dynamics. *Nat. Neurosci.* **19**, 1489–1496 (2016).

23. Ivanova, D., Li, X.-F., McIntyre, C. & O'Byrne, K. T. Posterodorsal Medial Amygdala Urocortin-3,

GABA, and Glutamate Mediate Suppression of LH Pulsatility in Female Mice. *Endocrinology*

164, 1–13 (2023).

24. Yuste, R., Cossart, R. & Yaksi, E. Neuronal ensembles: Building blocks of neural circuits.

Neuron **112**, 875–892 (2024).

25. Voliotis, M. *et al.* The Origin of GnRH Pulse Generation: An Integrative Mathematical-

Experimental Approach. *J. Neurosci.* **39**, 9738–9747 (2019).

26. Hochgerner, H. *et al.* Neuronal types in the mouse amygdala and their transcriptional response

to fear conditioning. *Nat. Neurosci.* **26**, 2237–2249 (2023).

27. Li, Y. *et al.* Neuronal Representation of Social Information in the Medial Amygdala of Awake

Behaving Mice. *Cell* **171**, 1176-1190.e17 (2017).

28. Pardo-Bellver, C., Cádiz-Moretti, B., Novejarque, A., Martínez-García, F. & Lanuza, E.

Differential efferent projections of the anterior, posteroventral, and posterodorsal subdivisions of

the medial amygdala in mice. *Front. Neuroanat.* **6**, 33 (2012).

29. Polepalli, J. S., Gooch, H. & Sah, P. Diversity of interneurons in the lateral and basal amygdala.

Npj Sci. Learn. **5**, 10 (2020).

30. Patel, A. H. *et al.* Kisspeptin in functional hypothalamic amenorrhea: Pathophysiology and

therapeutic potential. *Ann. N. Y. Acad. Sci.* **1540**, 21–46 (2024).

31. Barbagallo, F. *et al.* Gene-environment interaction in functional hypothalamic amenorrhea.

Front. Endocrinol. **15**, 1423898 (2024).

32. Paxinos, G. & Franklin, K. B. J. *Paxinos and Franklin's the Mouse Brain in Stereotaxic*

Coordinates. (2019).

33. Steyn, F. J. *et al.* Development of a Methodology for and Assessment of Pulsatile Luteinizing

Hormone Secretion in Juvenile and Adult Male Mice. *Endocrinology* **154**, 4939–4945 (2013).

34. Vidal, A., Zhang, Q., Médigue, C., Fabre, S. & Clément, F. DynPeak: An Algorithm for Pulse

Detection and Frequency Analysis in Hormonal Time Series. *PLOS ONE* **7**, e39001 (2012).

35. Yang, J. A. *et al.* Acute Psychosocial Stress Inhibits LH Pulsatility and Kiss1 Neuronal Activation

in Female Mice. *Endocrinology* **158**, 3716–3723 (2017).

36. Pnevmatikakis, E. A. & Giovannucci, A. NoRMCorre: An online algorithm for piecewise rigid

motion correction of calcium imaging data. *J. Neurosci. Methods* **291**, 83–94 (2017).

37. Battagello, D. S. *et al.* Anatomical Organization of Urocortin 3-Synthesizing Neurons and

Immunoreactive Terminals in the Central Nervous System of Non-Human Primates [Sapajus

spp.]. *Front. Neuroanat.* **11**, (2017).

38. Friedrich, J., Zhou, P. & Paninski, L. Fast online deconvolution of calcium imaging data. *PLOS*

Comput. Biol. **13**, (2017).

Acknowledgements

903 We thank Ms Sumi Mathew for her technical support with genotyping mice.

904 S.F., M.V., K.T.A., K.T.O., X.F.L. gratefully acknowledge the financial support of BBSRC via grants

905 BB/W005913/1 (KCL), BB/W005883/1 (Exeter) and BB/S019979/1. KTA gratefully acknowledges

906 the financial support of the EPSRC via grant EP/T017856/.

Contributions

J.Y. contributed data acquisition and data analysis, S.F. contributed data analysis, K.N. contributed

modeling, M.V. contributed managing data analysis, X.F.L. contributed managing data acquisition,

H.Y. contributed data acquisition, Y.L. contributed data acquisition, J.Y. contributed data acquisition,

O.H. contributed data acquisition, R.D.B. contributed data acquisition, B.S. contributed data

acquisition, K.T.O., K.T.A. and M.V. contributed to managing the project. All authors contributed to

writing the manuscript.

Competing interests

915 The authors declare no competing interests.

Corresponding author

917 Correspondence and requests for materials should be addressed to Krasimira Tsaneva-Atanasova

918 (K.Tsaneva-Atanasova@exeter.ac.uk), Kevin T. O'Byrne (kevin.o'byrne@kcl.ac.uk), Margaritis

919 Voliotis (m.voliotis@exeter.ac.uk).

Open Access