GABA Neurons in the Amygdala play a central Role in Urocortin-3–Mediated Stress Suppression of Reproduction

Author Information

Junru Yu[1,2,†], Saeed Farjami[3,4,5†] , Kateryna Nechyporenko[3,4,†], Xiao Feng Li[1], Hafsa Yaseen[1,6], Yanyan Lin[1], Jinbin Ye[1], Owen Hollings[1], Ross de Burgh[1], Baban Singh[1], Kevin T. O'Byrne[1,*], Krasimira Tsaneva-Atanasova[3,4,7,*], Margaritis Voliotis[3,4,*]

Affiliations

[1]Department of Women and Children's Health, School of Life Course and Population Sciences, King's College London, Guy's Campus, London SE1 1UL, UK, [2]Department of Rehabilitation Medicine, The First Affiliated Hospital of Wenzhou Medical University, Wenzhou, Zhejiang 325000, China, [3]Department of Mathematics and Statistics, Stocker Road, Exeter EX4 4PY, UK, [4]Living Systems Institute, Exeter EX4 4QD, UK, [5]The Pirbright Institute, Ash Road, Pirbright, Surrey GU24 0NF, UK, [6]Biological Sciences, University of Missouri, Columbia, MO 65211-7400 and [7]EPSRC Hub for Quantitative Modelling in Healthcare, University of Exeter, Stocker Road, Exeter EX4 4PY, UK

Correspondence to: Krasimira Tsaneva-Atanasova (k.tsaneva-atanasova@exeter.ac.uk), Kevin T. O'Byrne (kevin.o'byrne@kcl.ac.uk), Margaritis Voliotis (m.voliotis@exeter.ac.uk).

Abstract

Stress can disrupt menstrual cycles, cause infertility, and lead to other reproductive disorders. The posterodorsal medial amygdala (MePD) processes stress signals and regulates the gonadotropin-releasing hormone (GnRH) pulse generator through GABAergic inhibitory projections to the hypothalamus. However, how stress is processed in the MePD - especially involving its dense GABA and Urocortin-3 (UCN3) neurons - remains poorly understood.

In this study, we combine *in vivo* GRadient-INdex (GRIN) lens mini-endoscopic calcium imaging (to track neuronal activity), optogenetics, clustering analysis, and computational modeling to investigate the MePD circuitry. Our findings reveal two anti-correlated GABA subpopulations in the MePD that dictate responses to both UCN3 neuron stimulation and restraint stress. Our computational modeling suggests that mutual inhibition between these GABA groups drives the anti-correlated activity and predicts how these interactions shape downstream responses to stimulation of GABA and UCN3 neurons.

We test these predictions using optogenetics and confirm that GABA neurons are critical for the transmission of UCN3 signals to regulate luteinizing hormone (LH) pulse frequency. Our study is the first to show how GABA neurons in the amygdala mediate stress effects on reproductive health, uncovering key neural mechanisms linking emotional and reproductive functions.

Introduction

Successful reproduction requires normal function of a neural oscillator, the gonadotropin-releasing hormone (GnRH) pulse generator, that drives the pulsatile release of gonadotropic hormones, luteinizing hormone (LH) and follicle stimulating hormone (FSH) to control ovulation and spermatogenesis. This oscillator comprises Kisspeptin neurons co-expressing Neurokinin-B (NKB) and Dynorphin A (acronym KNDy) in the hypothalamic arcuate nucleus (ARC). The KNDy neural network not only provides the essential episodic stimulatory kisspeptin signal to the GnRH neurons, but receives diverse afferent inputs and hence is a hub for integrating various internal and external cues that regulate reproductive function[1,2]. Despite the marked heterogeneity of the myriad stressors, including acute versus chronic experimental stress paradigms, the common denominator is suppression of GnRH pulse generator frequency, which is associated with stress-related reproductive disorders and infertility[3–8].

The amygdala, a part of the limbic brain typically associated with emotions and anxiety, has strong projections to the KNDy system[1,2]. The medial amygdala, and more specifically its posterodorsal subnucleus (MePD), has been shown to be a robust upstream regulator of GnRH pulse generator frequency[9–11], pubertal timing[12,13], sociosexual behavior[14,15], anxiety[14] and psychological stress-induced suppression of pulsatile LH secretion[16,17].

In the MePD, the majority of neurons are GABAergic, accounting for a substantial portion of its projections, many of which target important hypothalamic centers involved in reproduction[18–20], including the KNDy neurons and optogenetic stimulation of the latter suppresses LH pulsatility[17]. Moreover, chemogenetic inhibition of GABA neurons in the MePD can reverse the inhibitory effect of psychogenic stress on LH pulsatility[17], suggesting an involvement of GABA signaling in the stress-induced suppression of pulsatile LH secretion. Although the precise way in which the MePD GABA neurons influence LH pulsatility is not fully understood, it has been suggested that they interact with the stress neuropeptide urocortin-3 (UCN3) neurons to modulate or reinforce stress-induced suppression of LH pulse frequency[16].

UCN3, part of the corticotropin-releasing factor (CRF) superfamily and selective for CRF type 2 receptors (CRFR2)[21], is highly expressed in the MePD[16,22]. Furthermore, UCN3 neurons in the MePD mediate psychosocial stress-induced suppression of the GnRH pulse generator and corticosterone secretion, thus playing a central role in the interaction between the reproductive and stress axes[16,23]. Previously we have shown that intra-MePD GABA antagonism can reverse UCN3-induced LH pulse suppression[23], suggesting that GABA acts downstream of UCN3 signaling within the MePD. However, the local neural circuitry within the MePD mediating stress related information to the GnRH pulse generator remains to be established.

Here, we employed GRadient INdex (GRIN) lens mini-endoscopy to monitor and compare the calcium activity of GABA neurons at single-cell resolution under acute psychological stress or optogenetic stimulation of UCN3 neurons. Clustering analysis of the observed calcium transients uncovered two distinct GABA subpopulations exhibiting anti-correlated activity. To explore the functional role of these two subpopulations and unpick the mechanisms driving their anti-correlated activity, we build upon our previously developed MePD circuitry mathematical modeling framework[11]. Our modeling reveals that mutual inhibitory interactions between the GABA populations are sufficient to drive anti-correlated activity and predicts a complementary inhibitory effect of GABA and UCN3 signaling on the frequency of the GnRH pulse generator, which we tested by applying Cre/Flp recombinase intersectional strategies to selectively target and manipulate heterogeneous MePD neuronal populations[17]. Hence, by uncovering precise interactions between distinct subpopulations of MePD GABA neurons and UCN3 neurons in transmitting stress signals to the GnRH pulse generator, we provide critical and previously unknown insights into the neural circuitry driving stress-induced reproductive dysfunction. This discovery significantly advances our understanding of how emotional stress directly impacts reproductive health and identifies potential neural targets for therapeutic intervention in stress-related reproductive disorders.

Results

Dynamic reconfiguration of MePD GABA neuronal networks during selective optogenetic stimulation of urocortin-3 (UCN3) neurons and restraint stress

To investigate how the posterodorsal subnucleus of the medial amygdala (MePD) processes stress signals and to understand the role of GABA and UCN3 neurons in this process, we employed *in vivo* calcium imaging using GRadient INdex (GRIN) mini-endoscopy. A cohort of six female UCN3Cre::VGAT-Flpo transgenic mice were injected with a viral mixture containing pAAV-Syn-Flexrc[Chrimson-tdTomato] and pAAV-EF1a-fDIO-GCaMP6s. This intersectional viral strategy enabled dual functionality: (i) optogenetically stimulating MePD UCN3 neurons via Chrimson; and (ii) detecting calcium activity in MePD GABA neurons using GCaMP6s (Fig. 1**A**). GRIN lenses were surgically implanted at the same location as the viral injection, enabling imaging of calcium activity with single-neuron resolution Fig. 1**B-G**). The specificity of the AAV-Syn-Flex-rc[Chrimson-tdTomato] expression in the MePD was quantified by overlapping analysis with the UCN3 immunohistochemical staining and revealed that 95.06 ± 2.24% of tdTomato-positive neurons co-localized with the UCN3 immunohistochemical positive neurons (Supplementary Fig. 1). Further, 81.61 ± 6.03% of the total UCN3-immunoreactive neurons co-expressed Chrimson-tdTomato (Supplementary Fig. 1).

Two series of experiments were conducted to assess GABAergic responses under distinct conditions. First, to assess the role of UCN3 neurons in recruiting GABAergic response, baseline calcium activity from MePD GABA neurons was recorded for 3 min, followed by 3 min of recording during optogenetic stimulation of UCN3 neurons (Fig. 1**A**). Optogenetic stimulation was delivered in a discontinuous pattern, 5 seconds on, 5 seconds off, with a light-pulse frequency of 10 Hz during the 'on' periods (10ms pulse width, 10 mW power, and 532 nm wavelength). GABA neurons with activity that mimics the pattern of optic stimulation, evidence that they are co-expressing UCN3, are removed from further analysis (see Methods and Supplementary Fig. 2). To further characterize the GABAergic response to acute stressors, baseline calcium activity of GABA neurons was recorded for 2 min, followed by 2 min of recording while the animals were exposed to restraint stress (Fig. 1**A**). The dynamic GABAergic response to optogenetic stimulation of UCN3 neurons and restraint stress was summarized in

functional terms using pairwise (neuron-neuron) association matrices, constructed from Pearson

correlation coefficients between neuronal trace pairs (see Methods). These correlation matrices

represent the MePD GABAergic response as a graph of interconnected nodes (neurons) with edge

weights corresponding to the strength of their pairwise functional interactions (Fig. 1**H&I**)[24]. To quantify

the effect of stress or UCN3 activation on functional connectivity we calculated the normalized

Riemannian distance between the correlation matrices from baseline and stimulation/stress periods.

Our analysis revealed significant alterations in the functional connectivity of the GABAergic network

during both UCN3 stimulation and stress relative to control animals lacking channelrhodopsin

expression (Fig. 1**J**). Furthermore, on average, stress induces a significantly higher population time-

averaged calcium response than UCN3 stimulation; however, this effect diminishes over the 2-minute

stress period (Fig. 1**K**). These results indicate that the GABA population in the MePD plays a pivotal

role in processing stress signals, such as those induced by restraint stress. Furthermore, our findings

suggest that UCN3 neurons may serve as a key relay, driving functional reorganization within the

GABAergic population to support transmission of stress-related information.

Figure 1: Functional connectivity of GABA neurons in posterodorsal medial amygdala (MePD) reconfigures in response to optogenetic stimulation of urocortin-3 (UCN3) neurons or acute restraint stress. (**A**) Schematic diagram of the experimental design for stimulating MePD UCN3 neurons and simultaneously imaging calcium activity from MePD GABA neurons. Representative heatmaps of calcium activity ($\Delta F/F_0$) recorded from MePD GABA neurons (left) in response to optogenetic stimulation of UCN3 neurons, (animal G7 trial 2, 34 neurons, see Supplementary Table 1; GABA neurons meeting our criteria for UCN3 expression excluded see Supplementary Fig. 2) and (right) during restraint-induced stress (animal G28, 35 neurons, see Supplementary Table 2). **(B-G)** Representative photomicrographs of the MePD from a UCN3-Cre::VGAT-Flpo mouse injected with pAAV-Syn-Flex-rc+pAAV-EF1a-fDIO-GCaMP6s. (**B&E**) Red fluorescence

(tdTomoato) labels UCN3 neurons expressing Chrimson-tdTomato, (**C&F**) green fluorescence labels GABA neurons, and (**D&G**) red and green signals merged. (**H&I**) Functional connectivity matrices during baseline and under UCN3 activation and stress were constructed by calculating the pairwise Pearson's correlation between calcium time-traces. (**J**) Normalized Riemannian distance between the correlation matrices from baseline and stimulation/stress periods. Each data point represents a single experiment, with different animals color-coded; control animals are shown as squares, and channelrhodopsin-expressing animals as circles. (**K**) Difference in time-averaged calcium activity between periods of UCN3 stimulation or stress and the baseline period. Scale bars: **B-D**, 200 μm; **E-G**, 50 μm.

Clustering analysis reveals two functionally distinct GABA subpopulations in the MePD

To better understand the functional organization of the GABA MePD network during optogenetic stimulation of UCN3 neurons and under restraint stress, we performed clustering analysis on the functional connectivity matrices we obtained from the calcium recordings of GABA neurons.

Clustering results consistently revealed two functionally distinct cell populations both during optogenetic stimulation of UCN3 neurons and restraint stress (Fig. 2**A&C**). Under both conditions, activity between the two clusters appeared highly anti-correlated as illustrated by two representative examples depicted in Fig. 2**B&D**. Moreover, we find that clusters are significantly different in size, with the smallest cluster accounting on average for 35% of all cells (Fig 2**E&F**, Supplementary Fig. 3). The robustness of the above finding was also verified using the K-means clustering algorithm and an alternative measure of association between neuronal traces that accounts for time-lag and sign in neuronal interactions (see Methods and Supplementary Figs. 4-6).

Varying the frequency of optogenetic stimulation of UCN3 neurons influenced the functional connectivity within the GABA network (Supplementary Fig. 7). Stimulation at 5 Hz produced no significant change compared with controls, 10 Hz resulted in a statistically significant increase, while 20 Hz reduced the effect, which could possibly reflect a differential release of neurotransmitters/neuropeptides at higher frequencies[25]. In contrast, stimulation at 5, 10, or 20 Hz did not significantly affect the correlation between the average activity of the two GABA clusters (Supplementary Fig. 8). Based on the functional role of the MePD in relaying stress signals to the hypothalamic arcuate nucleus (ARC), our results suggest that the identified GABA subpopulations

(clusters) could correspond to distinct populations of GABA interneurons and GABA efferent

neurons—defined as neurons that transmit signals from the MePD to downstream targets. Known

efferent projections to the ARC play a role in regulating the gonadotropin-releasing hormone (GnRH)

pulse generator, a key driver of reproductive function[11,17].

Figure 2: **Clustering identifies two functionally distinct GABA subpopulations in the posterodorsal**

medial amygdala (MePD) both during UCN3 optogenetic stimulation and restraint stress. Representative

example of (**A**) a clustered connectivity matrix and (**B**) associated calcium time-traces of MePD GABA neurons

during optogenetic stimulation of UNC3 neurons and the population average below. Representative examples

of (**C**) a clustered connectivity matrix and (**D**) associated calcium activity time-traces of MePD GABA neurons

during restraint stress and the average population activity below. (**E&G**) Relative sizes of the two GABA

subpopulations identified via clustering analysis. (##: p<0.001).

Mutually inhibitory interactions between GABA populations drive anti-correlated activity

To explore the network mechanisms driving the anti-correlated activity of the GABA subpopulations,

we build upon our previously developed MePD circuit model[11]. The model allows us to explore

interactions between MePD neuronal populations and understand emergent patterns of dynamical

activity. Specifically, the model describes the calcium dynamics of two interacting populations of GABA

neurons, corresponding to interneurons and efferent projection neurons. Efferent neurons project their

axons outside the MePD to downstream targets such as the KNDy network. Both GABA populations

interact with a glutamatergic population (Fig. 3**A**). Experimental evidence shows that antagonizing

glutamate signaling completely blocks the suppressive effect of MePD UCN3 activation on LH

pulsatility[23], highlighting a functional role of glutamatergic neurons in the circuit. Therefore, the activity

of the glutamatergic population is also included in the MePD model alongside the two GABA

populations. Model equations are provided in Supplementary Information, and the parameter values

for the MePD model are in Supplementary Table 3.

To incorporate input from UCN3 neurons we include a non-autonomous term in the model informed

by recordings of the activity of UCN3 neurons during optogenetic stimulation (see Fig. 3**A** and

Methods). The recorded traces exhibit a periodic pattern driven by the experimental stimulation

protocol with 5 seconds on, 5 seconds off (see Methods). In the model, we approximated this UCN3

input using a constant drive along with a sinusoidal function with roughly 5 seconds on, 5 seconds off

pattern, driving the populations GABA interneurons and the glutamatergic neurons. To simulate the

anti-correlated activity between the two GABA neuronal subpopulations while also reproducing

previously published results on the effects of neuropharmacological perturbations in the MePD on LH

pulsatility[23], we find that we need to include bi-directional (i.e. mutually inhibitory) GABA interaction in

the circuit as well as GABA interneurons self-inhibition (Fig. 3**A**).

Next, we validate our model by simulating the effects of stress on the MePD circuit. UCN3 neurons in

the MePD have also been shown to mediate psychosocial stress-induced suppression of the GnRH

pulse generator[16]. Experimentally recorded calcium activity of UCN3 neurons during stress

intervention is characterized by a fast rise followed by a slow decay (Fig. 3**B**), which we model by

taking the difference between two exponential functions, with one fast and one slow time constant

(see Methods). Using this input, our mathematical model confirms the emergence of anti-correlated

behavior between the GABA subpopulation whereby the activity of GABA interneurons decreases

over the stress period, while the activity of the GABA efferent neuron population increases (Fig. 3**B**),

supporting the proposed MePD circuit structure (Fig. 3**A**).

Figure 3: Mathematical model simulating the activity in the GABA neuronal populations in the MePD

under the increased excitatory input from UCN3 neurons and mimicking the conditions under restraint

stress. The schematics on the left provide a visual representation of the assumptions built into our mathematical

model of the MePD neuronal circuit, consisting of two GABA populations and a glutamatergic population, that

interact with UCN3. **(A)** Calcium time-traces of MePD UCN3 neurons during optogenetic stimulation of UCN3

neurons and corresponding mean activity. Model simulations of the effects of UCN3 optogenetic stimulation

supports the emergence of anti-correlated activity in GABA subpopulations. **(B)** Calcium time-traces of MePD

UCN3 neurons during restraint stress and corresponding mean activity of the UCN3 calcium traces during restraint stress with corresponding mean activity. Simulations of the MePD circuit under the effects of stress also support anti-correlated GABA subpopulations' activity.

***In silico* modeling reveals complementary relationship between MePD GABA and UCN3 populations**

Having shown that the model of the MePD circuitry reproduces the observed anti-correlated dynamics, we next couple it to our network model of the KNDy population[11,26]. KNDy model details are in Supplementary Information, and the model parameter values can be found in Supplementary Table 4. Since KNDy neuronal activity serves as a well-established proxy for LH secretion, we use the dynamics of the KNDy network model to infer changes in LH pulsatility in response to MePD circuit manipulations. Our previous in silico findings[11] show that MePD signal integration is predominately mediated by GABA projections. The effects of MePD GABA and glutamate antagonism alone and together with UCN3 stimulation on LH pulsatility have been previously investigated[23], and we use these experimental findings to calibrate our coupled model parameters by reproducing these effects accordingly (see Supplementary Fig. 9).

Next, using the calibrated model, we predict that providing a periodic excitatory input (see Supplementary Fig. 10**A**), that mimics the effects of optogenetic stimulation of UCN3, leads to the increase in the KNDy inter-pulse interval (IPI) from 15.18 min to 23.97 min due to increased activity in the inhibitory GABAergic input to the KNDy network (Fig. 4**A**).

Since MePD has an overwhelming number of GABAergic outputs and interneurons, we are interested in their role in information processing of stress signaling. To this end we provide periodic excitatory input directly to both populations of GABA neurons (see Methods and Supplementary Information). This is different to the indirect activation through UCN3 stimulation, as it leads to the increased activity in both GABA subpopulations (Supplementary Fig. 10**B**). The resulting effect on the KNDy IPI is an increase from 15.18 min to 42.65 min (Fig. 4**B**).

In the model we hypothesize that UCN3 signaling to the KNDy is operating through the GABAergic pathway, hence disrupting GABAergic interactions will suppress UCN3's effects on KNDy. To

investigate it we suppress the strength of the GABAergic interactions by 50% and mimic UCN3

stimulation by providing excitatory input to the circuit. The model shows that this intervention does not

have a significant effect on KNDy IPI, apart from slightly increasing it from 15.18 min to 17.31 min

(Fig. 4**C** & Supplementary Fig. 10**C**).

Taken together the above could be summarized as the following set of model predictions along with

suggested ways to test them experimentally:

1. Optogenetic stimulation modelled via excitation of the MePD UCN3 increases the KNDy IPI

15.18 min to 23.97 min. This could be tested experimentally via in vivo optogenetic activation

of the MePD UCN3 neuronal population and simultaneous measurement of LH pulsatility.

2. Stimulation of the MePD GABA populations increases the KNDy IPI 15.18 min to 42.65 min.

Such a substantial change underscores the critical role of GABAergic signaling in modulating

KNDy pulsatility. Experimental confirmation could involve in vivo optogenetic activation of the

entire MePD GABA population accompanied by LH pulsatility measurement.

3. Stimulation of the MePD UCN3 with simultaneous suppression of the GABAergic interactions

in the MePD does not significantly affect KNDy IPI, changing it from 15.18 min to 17.31 min.

This finding suggests that UCN3 effects on the KNDy network are primarily dependent on its

interactions with GABA populations. A strategy to verify this hypothesis experimentally could

be to stimulate UCN3 non-GABA-expressing neurons in vivo while simultaneously inhibiting

non-UCN3 GABA neurons and measure LH pulsatility.

Finally, we investigate in silico the effects of stress on LH pulse dynamics mediated via the MePD. In

the restraint-stress experiment it was shown that one GABA cluster was active, while the other one

was inactive (Fig. 2**D**). By simulating these dynamics (Supplementary Fig. 9**D**) in the coupled MePD

- KNDy network model we are able to demonstrate that pulses in the KNDy network ceased (Fig. 4**D**).

These modeling findings align with published experimental evidence indicating that restraint stress

suppresses LH pulsatility[16].

Figure 4: Coupled MePD - KNDy model predictions for the effect of stress and stimulation of UCN3 neurons on GnRH pulse generator activity and hence LH pulsatility. (**A**) Simulation of the optogenetic stimulation of the UCN3 neurons leads to the increase of the KNDy IPI 15.18 min to 23.97 min. (**B**) Mimicking the effects of the optogenetic stimulation of the MePD GABA neurons *in silico* (providing excitatory input to both GABA populations) increases KNDy IPI from 15.18 min to 42.65 min. (**C**) Simulation of the effects of the optogenetic stimulation of the UCN3 neurons and simultaneous inhibition of GABA neuronal populations do not significantly affect KNDy IPI, where it changes from 15.18 min to 17.32 min. (**D**) Simulating the effects of stress leads to the cessation of the pulsatile dynamics.

Selective optogenetic stimulation of UCN3 non-GABA neurons inhibits pulsatile LH secretion in female mice.

We now test experimentally our modeling predictions detailed in the previous section regarding the inhibitory effect of MePD UCN3 activation on LH pulsatility in vivo in freely behaving mice. To this end a cohort of UCN3-Cre::VGAT-Flpo transgenic female mice was injected with an AAV-nEF-Con/Foff 2.0-ChRmine-oScarlet viral construct. This intersectional approach enabled conditional expression of channelrhodopsin (ChRmine) in UCN3 neurons not expressing GABA (UCN3 non-GABA) (Con/Foff group). Following the viral injection, a unilateral fiber-optic cannula was implanted into the MePD, above the injection site, to facilitate selective activation of the targeted neuronal population. The specificity of the AAV-nEF-Con/Foff 2.0-ChRmine-oScarlet expression in the MePD was quantified by overlapping analysis with the UCN3 immunohistochemical staining and revealed that 98.92 ± 1.08% of the oScarlet-positive neurons co-localized with the UCN3 immunohistochemical positive neurons (Supplementary Fig. 11). Further, 54.18 ± 5.04% of the total UCN3-positive neurons coexpressed ChRmine-oScarlet (Supplementary Fig. 11).

The stimulation protocol consisted of a 60-min control period, followed by 60 min of optogenetic stimulation. Stimulation was delivered, as above, in a pulsed pattern (5 seconds on, 5 seconds off) with a pulse frequency of 10 Hz during the on period (10-ms pulse width, 10 mW power, 532 nm wavelength). Blood samples (5 µL) were collected every 5 min throughout the protocol and used to measure LH levels.

Statistical analysis indicated post-stimulation LH inter-pulse intervals differed across the two groups ($F_{(2, 30)} = 15.83$, $P < 0.0001$; two-way ANOVA). In the Con/Foff group, the LH inter-pulse interval significantly increased after optogenetic stimulation compared to the control period (15.47 ± 0.91 min vs. 23.75 ± 1.07 min, mean ± SEM, n = 6, p<0.01; Fig. 5**A&D)**, validating the first of our modeling predictions. In contrast, the no-stimulation group, which received the Con/Foff viral construct without optogenetic stimulation, showed no significant changes in LH inter-pulse interval (15.84 ± 0.37 min vs. 16.68 ± 0.86 min, n = 6, p > 0.05; Fig. 5**B&D**). Similarly, in a separate cohort of mice injected with a control virus (DIO-EYFP), optogenetic stimulation resulted in no significant differences in LH inter-

pulse intervals before and after stimulation (19.83 ± 1.02 min vs. 17.78 ± 0.93 min, n = 6, p > 0.05;

Fig. 5**C&D**). The LH pulse amplitude and mean LH levels were quantified, and data are provided in

Supplementary Table 5.

Selective optogenetic stimulation of UCN3-expressing GABA neurons fails to affect pulsatile

LH secretion in female mice.

To further test the functional interactions between the UCN3 and GABA populations, a separate cohort

of UCN3-Cre::VGAT-Flpo transgenic female mice was injected with an AAV-nEF-Con/Fon

2.0ChRmine-oScarlet viral construct, which allows optogenetic stimulation of UCN3 neurons

coexpressing GABA (UCN3-expressing GABA) (Con/Fon group). The specificity of the AAV-

nEFCon/Fon 2.0-ChRmine-oScarlet expression in the MePD was quantified by overlapping analysis

with the UCN3 immunohistochemical staining and revealed that 98.77 ± 1.24% of the oScarlet-positive

neurons co-localized with the UCN3 immunohistochemical positive neurons co-expressing GABA

(Supplementary Fig. 12). Further, 32.73± 2.60% of the total UCN3-positive neurons co-expressed

ChRmine-oScarlet (Supplementary Fig. 12).

Statistical analysis showed no significant interaction effect of group and optogenetic stimulation on LH

inter-pulse intervals (F (1, 18) = 0.2709, p = 0.6091; two-way ANOVA). That is, LH inter-pulse intervals

in the Con/Fon group, showed no change following optogenetic stimulation compared to the control

period (17.92 ± 0.80 min vs. 16.60 ± 0.88 min, mean ± SEM, n = 6, p > 0.05; Fig. 5**H&J**). Similarly, in

the no-stimulation control experiment for this cohort, which did not receive optogenetic stimulation,

there was no significant difference in LH inter-pulse intervals (16.08 ± 1.45 min vs. 15.83 ± 0.95 min,

n = 5, p > 0.05; Fig. 5**I&J**). The LH pulse amplitude and mean LH levels were quantified, and data are

provided in Supplementary Table 5.

Taken together, our results suggest that although UCN3 and GABAergic neuronal populations overlap,

they serve distinct functional roles within the MePD circuit, with both being critical for relaying inhibitory

signals to the hypothalamic GnRH pulse generator. While independent activation of either population

inhibits the GnRH pulse generator, activation of UCN3 neurons that co-express GABA is insufficient

to elicit this effect.

Figure 5: Effect of optogenetic stimulation of UCN3 non-GABA and UCN3 neurons co-expressing GABA in the MePD on LH pulse frequency. (**A-D**) Optogenetic stimulation of UCN3 non-GABA neurons inhibited LH pulses. Representative examples of LH pulsatility in UCN3-Cre::VGAT-Flpo mice injected with AAV-Con/FoffChRmine in response to (**A**) 10Hz optic stimulation or (**B**) no stimulation as control, and (**C**) in mice injected with control virus in response to 10 Hz stimulation. Pulses detected by the DynPeak algorithm are indicated with an asterisk (*). (**D**) Mean LH inter-pulse interval (IPI) (±SEM) for each group during the control period and over the subsequent stimulation period. #: p<0.01 vs stimulation period in Con/Foff group (two-way ANOVA, Tukey's post-hoc). (**E-G**) Representative photomicrographs of expression of AAV-nEF-Con/Foff 2.0-ChRmine-oScarlet (red) in MePD UCN3 neurons. (**H-M**) Optogenetic stimulation of neurons co-expressing UCN3 and GABA in the MePD did not impact LH pulse significantly. Representative examples of LH pulsatility in UCN3-Cre::VGATFlpo mice injected with Con/Fon-ChRmine-oScarlet virus in response to (**H**) 10 Hz optic

stimulation or (**I**) no stimulation as control. (J) Mean LH IPI (±SEM) for each group during the control period and over the subsequent stimulation period (two-way ANOVA). (**K-M**) Representative photomicrographs of neurons co-expression UCN3 and GABA, tagged with oScarlet (red). Scale bars represent (**E&K**) 200 μm, (**F&L**) 100 μm, (**G&M**) 50 μm. OT, optic track. The blue vertical bars in E to G and K to M indicate the position of the fiber optic cannula.

The MePD GABA neurons relay information from UCN3 neurons and inhibit the GnRH pulse generator.

To test our remaining model predictions and establish the functional role of GABA neurons in the MePD in transmitting inhibitory signals to the hypothalamic GnRH pulse generator, we injected the AAV-EF1a-DIO-ChR2-EYFP viral construct into VGAT-Cre transgenic mice, targeting the entire MePD GABA neuron population for expressing ChR2 and used the optogenetic stimulation protocol described above (60 min control period followed by 60 min stimulation) to stimulate the entire MePD GABA population. In the VGAT-Cre-tdtomato mice, we observed that 93.27 ± 2.99% of the EYFP positive neurons were tdtomato-positive and 87.01± 5.21% of the tdtomato-positive neurons were EYFP-positive. This indicates a high degree of co-expression consistency between the viral and transgenic reporters. Statistical analysis showed a significant interaction effect of our genetic intervention and stimulation period on LH pulsatility ($F_{(2, 24)}$ = 10.55, p = 0.0005; two-way ANOVA). Stimulation of the GABA population markedly increased the LH inter-pulse interval compared to the pre-stimulation control period (18.5 ± 2.03 min vs 49.5 ± 8.46 min, mean ± SEM, n = 5, p<0.01; Tukey's post-hoc; Fig. 6**A&D**), validating modeling prediction 2. In contrast, mice that did not receive optogenetic stimulation showed no change in LH inter-pulse interval (21.25 ± 3.79 min vs 19.33 ± 1.52 min, n = 5, p > 0.05; Tukey's post-hoc; Fig. 6**B&D**). Similarly, control virus (AAV-EF1a-DIO-EYFP) injections did not result in any significant difference in LH inter-pulse interval before and during stimulation (16.50 ± 0.67 min vs 18.00 ± 0.62 min, n = 5, p > 0.05; Tukey's post-hoc; Fig. 6**C&D**). The LH pulse amplitude and mean LH levels were quantified, and data are provided in Supplementary Table 6.

Next, UCN3-Cre::VGAT-Flpo transgenic females were injected with an intersectional viral mixture

containing AAV-nEF-Con/Foff 2.0-ChRmine-oScarlet and pAAV-nEF-Coff/Fon-NpHR3.3-EYFP ,

enabling simultaneous excitation of UCN3 non-GABA neurons and inhibition of non-UCN3 GABA

neurons. Statistical analysis showed no significant interaction effect of genetic manipulation and

stimulation period on LH pulsatility ($F_{(1,18)} = 0.131$, $p = 0.7216$; way ANOVA). Moreover, optogenetic

stimulation resulted in no significant change in the LH inter-pulse interval relative to the control period

(16.67 ± 1.31 min vs 18.61 ± 1.58 min, mean ± SEM, n = 6, $p > 0.05$; Fig. 6**H&J**), which aligns with

our modeling prediction 3. Similarly, in mice injected with the same viral mixture but without receiving

optogenetic stimulation, no significant change in LH inter-pulse interval was detected (16.33 ± 1.19

min vs 17.33 ± 0.81 min, n = 5, $p > 0.05$; Fig. 6**I&J**). The LH pulse amplitude and mean LH levels were

quantified, and data are provided in Supplementary Table 5.

Collectively, these findings suggest that MePD GABA neurons are integral to regulating LH pulse

frequency. The inhibitory effect on LH pulses observed upon UCN3 neuron activation is attenuated

when GABA neuron activity is optogenetically suppressed, implying that GABA signaling acts

downstream of UCN3 neurons, relaying the inhibitory signal to the hypothalamic GnRH pulse

generator.

Figure 6: Selective optogenetic stimulation of GABA neurons in the MePD inhibits LH pulsatility and their activity is necessary for mediating UCN3 signaling downstream. (**A-G**) Optogenetic activation of the entire MePD GABA population in VGAT-Cre mice led to significant LH pulse suppression. Representative examples showing LH pulsatility in VGAT-Cre mice injected with AAV-double floxed-ChR2-EYFP virus in response to (**A**) 5 Hz stimulation or (**B**) no stimulation as control, and (**C**) 5 Hz stimulation in mice injected with control virus. (**D**) Mean LH IPI (±SEM) for each group during the control period and over the subsequent stimulation period. #: p<0.01 vs stimulation period in ChR2 group (two-way ANOVA, Tukey's post-hoc). (**E-G**) Representative photomicrographs of MePD GABA neurons, labeled with (**E**) tdTomato (red), (**F**) ChR2-EYFP

(green), and (**G**) merged. Scale bars represent. (**H-P**) The simultaneous inhibition of non-UCN3 GABA neurons

in the MePD restored the LH suppression induced by stimulation of non-GABA UCN3 neurons. Representative

examples showing LH pulsatility in UCN3-Cre::VGAT-Flpo female mice injected with Con/FoffChRmine-oScarlet

and Coff/Fon-NpHR-EYFP virus mixture in response to (**H**) 10 Hz optogenetic stimulation or (**I**) no stimulation as

control. (**J**) Mean LH IPI (±SEM) for each group during the control period and over the subsequent stimulation

period. (two-way ANOVA). (**K-P**) Representative photomicrographs of fluorescence in MePD neurons. (**K&N**)

non-GABA UCN3 neurons tagged with oScarlet (red), (**L&O**) non-UCN GABA neurons tagged with EYFP (green)

(**M&P**) merged. Scale bars represent (**E-G**) 150 μm, (**K-M**) 200 μm, (**N-P**) 100 μm. OT, optic track. The blue

vertical bars in **E** to **G** and **K** to **P** indicate the position of the fiber optic cannula.

Discussion

The posterodorsal subnucleus of the Medial Amygdala (MePD) is a critical regulator of reproductive

function. Single-cell transcriptomics has revealed a mosaic of glutamatergic and GABAergic

populations in this region, with GABA neurons co-expressing receptors for gonadal steroids as well

as neuromodulators and cognate receptors involved in parental and sociosexual behavior (including

kisspeptin, prolactin and oxytocin)[27]. Furthermore, calcium imaging of neuronal ensembles from this

region has started to decipher how this brain region encodes information about social interactions and

stressful stimuli[28], with stress-related information being relayed to the reproductive axis via potentially

direct projections to the hypothalamus[1,2], and eliciting well-documented responses in pubertal

timing[12,13], LH secretion[16,17] and the GnRH pulse generator[9–11]. Here, to elucidate the neuronal circuitry

involved in MePD processing of stress signals, we used GRIN lens mini-endoscopic calcium imaging

to study, for the first time, the dynamic activity of MePD GABA neurons under acute restraint stress

as well as the activity of GABA neurons in response to UCN3 activation. Functional network analysis

of the imaging data revealed that acute restraint stress induces a reconfiguration of the GABA

population, highlighting their role in encoding and transmitting stress information. Optogenetic

stimulation of UCN3 neurons produced a comparable network reconfiguration indicating that both

populations are co-involved in stress processing. The overall magnitude of calcium activity ($\Delta F/F_0$)

differed between the two conditions, with stress generally inducing a stronger response than

optogenetic stimulation of UCN3 neurons. This discrepancy may reflect the engagement of additional

stress-related pathways or neuromodulatory factors that are not fully replicated by isolated UCN3

stimulation, such as stress-induced GABA transmission that has been well-documented in the

amygdala[17,23]. Clustering analysis of the calcium transients identified two distinct GABA

subpopulations with anti-correlated activity. This finding of two distinct populations is consistent with

physiological and morphological studies of GABA neurons in the medial amygdala characterizing them

as either projecting cells or local[18,20].

Building upon our previously developed computational model of the MePD circuit, we explored the

network connectivity to understand the possible mechanisms through which anti-correlated activity

emerges. The model consists of a glutamatergic subpopulation and two GABA subpopulations: one

comprising GABA interneurons and the other GABA efferent projection neurons. Our analysis

identified two conditions necessary for the emergence of anti-correlated activity between these GABA

subpopulations, consistent with our experimental observations. First, mutual inhibition between the

GABA subpopulations is required, such that activation of one population suppresses the other. In our

mechanistic model, mutual inhibition arises from the biophysical characteristics of GABAergic

transmission, where the release of GABA inhibits target neurons, thereby producing inhibitory

interactions between the subpopulations. Second, if both subpopulations receive a shared input (e.g.,

from the glutamatergic population or any other population of excitatory cells), their dynamics become

entrained to that common stimulus, meaning that distinct input streams to each subpopulation are

necessary for the observed anti-correlation. To satisfy this condition in our model, GABA interneurons

receive inputs from UCN3 neurons, whereas the GABA efferent projection neurons receive inputs

from the glutamatergic population. This circuitry implies that stress information is processed along two

key pathways (one glutamatergic and one GABAergic) and is integrated by the GABA efferent

projection neurons to the arcuate KNDy system.

We note that an alternative connectivity pattern, with the UCN3 population directly projecting to the

GABA efferent population, could also fit our experimental observations, but would not reproduce

previously published results on the effects of neuropharmacological perturbations in the MePD on LH

pulsatility[23]. Specifically, the suppression of glutamatergic neurons in combination with stimulation of

UCN3 neurons. Under this scenario, direct input from stimulated UCN3 to GABA efferent neurons will

lead to the increase in its activity, consequently increasing KNDy inter-pulse interval (IPI). This finding

contrasts with experimental observations, where the combined suppression of glutamate and

stimulation of UCN3 failed to produce any significant change in GnRH pulsatility[23].

Additionally, UCN3 neurons may co-release other fast neurotransmitters, such as glutamate[27]. The

result of our UCN3 neurons stimulations at low frequencies (5 Hz; Supplementary Fig. 7), which

primarily elicits the fast neurotransmitters release[25], suggests that their isolated effects may be limited.

We note that in our modeling framework, we include the constant drive, reflecting the effects of slow

neuropeptide action, and a periodic drive corresponding to the action of fast neurotransmitters aligned

with the stimulation frequency. The constant drive governs the overall dynamical regime, while the

fast periodic term functions primarily as a rhythmic modulator, that is when the periodic drive

dominates over the constant drive both the glutamatergic and GABAergic populations become

entrained to the stimulation frequency. Nonetheless the specific contribution of co-transmitters

remains experimentally unclear, and that may influence how the MePD integrates this tonic input (Fig.

7).

We note that our MePD model has potential applications beyond understanding MePD dynamics and

their influence on the GnRH pulse generator. The model incorporates two functionally distinct

populations of GABA neurons (interneurons and efferent projection neurons), which reflect known

differences in GABA neurons within the amygdala[18,20]. This diversity likely originates from the medial

ganglionic eminence an embryonic structure that gives rise to GABA interneuron populations across

various brain regions, including the MePD, through migration during embryogenesis[29]. Therefore, our

modelling framework could be extended to other neural circuits where two functionally distinct GABA

populations exist, such as the posterior ventral medial amygdala[20] and its lateral and basal

subdivisions[30].

Consistent with model predictions, we showed that optogenetic activation of non-GABA neurons,

expressing UCN3, suppresses LH pulse frequency. This finding corroborates previous work

demonstrating that LH pulses are inhibited by activation of UCN3 neurons[13,23] or administration of

UCN3[16] in the MePD, and that such effects are attenuated by chemogenetic inhibition of UCN3

neurons or by antagonizing corticotropin-releasing factor type 2 receptors (CRFR2)[16]. Thus, our

findings further support the central role of UCN3 signaling in modulating both stress responses and

reproductive function. Future studies could investigate the contribution of UCN3 neurons to stress

integration more directly, for example by silencing UCN3 neurons during stress exposure to determine

whether GABAergic activation and the associated network clustering patterns are abolished.

GABA is one of the primary inhibitory neurotransmitters in the nervous system. Previous work has

shown that MePD GABA neurons project to the ARC to regulate LH pulses and chemogenetic

inhibition of MePD GABA neurons attenuates stress-induced suppression of LH pulses[17]. Here, we

found that optogenetic activation of GABA neurons in the MePD leads to robust LH pulse suppression,

reinforcing the pivotal role of amygdala GABAergic signaling in stress-related reproductive

dysfunction. Antagonizing GABA receptors in the MePD can block UCN3-induced LH suppression[23],

suggesting that GABA neurons may function downstream of UCN3 signaling.

However, this pharmacological approach lacks cell-type specificity, leaving open the possibility that

UCN3 neurons themselves co-expressing GABA receptors, potentially contributing to the observed

effect. Here, by using an intersectional genetic strategy, we demonstrated that optogenetically

inhibiting non-UCN3 GABA neurons can abolish the change in LH pulse intervals induced by

optogenetic stimulation of UCN3 neurons. This finding supports the hypothesis that these GABA

neurons are acting downstream of UCN3 signaling, mediating LH pulse suppression. We note that

while in our current experimental protocol we could also be blocking GABAergic projections to the

ARC KNDy population, which by itself could explain the absence of any effect from UCN3 stimulation,

our modeling predicts that GABAergic interactions within the MePD are additionally crucial for relaying

UCN3 signals to the KNDy network as also suggested by pharmacologically antagonizing

GABAreceptors[23].

It is not unusual for neurons to co-express and co-release neuropeptides and neurotransmitters[20].

Here, our statistical analysis (Supplementary Fig. 2 & Supplementary Table 1) of GABA calcium traces

indicate that certain GABA neurons exhibit activity entrained to the stimulation frequency of UCN3

neurons, suggesting a possibility that UCN3 is co-released with GABA. Given that activating either

UCN3 or GABA neurons independently suppresses LH pulses, one might predict that stimulating

neurons co-expressing both UCN3 and GABA would similarly inhibit LH. However, we observed no

significant changes in LH pulses upon optogenetic stimulation of neurons co-expressing UCN3 and

GABA. Our results indicate that UCN3 and GABA are both essential for transmitting inhibitory signals

to the hypothalamic GnRH pulse generator; however, this inhibitory influence relies on their action

through separate, functionally distinct neuronal populations. That is activation of neurons that

exclusively express either UCN3 or GABA suppresses the GnRH pulse generator, whereas activation

of UCN3 neurons that also co-express GABA fails to produce this inhibitory effect. Within our modeling

framework, UCN3/GABA co-expressing neurons might be considered part of the GABA interneuron

population, primarily inhibiting GABA projection neurons rather than activating them. Consequently,

they fail to relay the inhibitory UCN3 signal to the ARC. Further studies are necessary to define more

precisely the distinct role of UCN3/GABA co-expressing neurons within the MePD circuitry.

Figure 7: Schematic illustration of the proposed functional MePD neurocircuit. UCN3 neurons activate inhibitory GABA interneurons and excitatory glutamate neurons through the release of UCN3 and potentially other fast neurotransmitters. The glutamate and GABA interneuron populations then signal to the GABA efferent projection neurons, that convey the MePD output to the KNDy neurons in the arcuate nucleus.

By combining mini-endoscopic calcium imaging with computational modeling, we identified two GABA subpopulations with complementary but distinct roles in encoding stress signals and modulating LH pulsatility. Our experimental and computational results also stressed the functional orthogonality between UCN3 and GABA signaling in MePD, highlighting the complexity of the MePD's contribution to stress-induced reproductive suppression. These results offer improved understanding of the neuronal circuits in the amygdala, a key hub for emotional processing, involved in regulating hypothalamic GnRH pulse generator frequency[9–11,16,17,23] providing new insight into psychogenic stress-related disorders of fertility, such as functional hypothalamic amenorrhoea[6,7,31,32], and the development of novel therapeutic approaches to improve reproductive health and well-being.

Methods

Animal

VGAT-Cre-tdTomato mice were genotyped using PCR to determine heterozygosity for VGAT-Cre (primers 5′-3′: common, 12785—CTTCGTCATCGGCGGCATCTG; wild-type reverse 12786—CAGGGCGATGTGGAATAGAAA; mutant reverse oIMR8292—CCAAAAGACGGCAATATGGT) and tdtomato (primers 5′-3′: wild-type forward oIMR9020—AAGGGAGCTGCAGTGGAGTA; wild-type reverse oIMR9021—CCGAAAATCTGTGGGAAGTC; mutant reverse WPRE oIMR9103—GGCATTAAAGCAGCGTATCC; mutant forward tdTomato oIMR9105—CTGTTCCTGTACGGCATGG). UCN3-Cre heterozygous mice (strain Tg(Ucn3-cre)KF43Gsat/Mmucd, MMRRC GENSAT) were crossbred with VGAT-Flpo heterozygous mice (Jax stock #007909,B6.Cg-Slc32a1 2A-Flpo-D:Het) to obtain female double heterozygous UCN3-Cre::VGATFlpo mice (Slc32a1 2A-Flpo-D:Het|Ucn3-Cre:Het|Rosa26<tomato>WT); genotyped using PCR for the detection of heterozygosity (primers 5′- 3′: UCN3 Forward: CGAAGTCCCTCTCACACCTGGTT; Cre Reverse: CGGCAAACGGAC-AGAAGCATT; Slc32a1 Mutant Forward: TGC ATC GCA TTG TCT GAG TAG; Slc32a1 Mutant Reverse: GAC AGC CGT GAA CAG AAG G). Female mice, aged 8 to 10 weeks, weighing between 20 g and 25 g, were individually housed in ventilated cages sealed with a HEPA-filter at 25 ± 1°C in a 12:12-hour light/dark cycle, lights on at 07:00 h. Vaginal smears were performed when mice are 6 weeks old between 10:00 - 12:00 h daily. Only mice showing regular estrous cycles were included. All animal procedures performed were approved by the King's College London Animal Welfare and Ethical Review Body. Procedures were in accordance with UK Home Office regulations.

Stereotaxic injection of adeno-associated virus construct and fiber-optic or mini-endoscope GRIN lens implantation

All surgical procedures were carried out under general anesthesia using ketamine (Vetalar, 100 mg/kg, i.p.; Pfizer, Sandwich, UK) and xylazine (Rompun, 10 mg/kg, i.p.; Bayer, Leverkusen, Germany) under aseptic conditions. Animals were secured in a David Kopf stereotaxic frame (Model 900, Kopf

Instruments, CA, USA). Following bilateral ovariectomy, a midline incision was made in the scalp to

reveal the skull. Two small bone screws were inserted into the skull. A small hole was drilled above

the position of the MePD. The MePD coordinate (2.29 mm lateral, 1.60 mm posterior from bregma, at

a depth of 5.33 mm below the skull surface) was determined according to the mouse brain atlas of

Paxinos and Franklin[33]. A unilateral stereotaxic viral injection was performed to target opsin or

fluorescent protein expression in the MePD neurons, using a robot stereotaxic system (Neurostar,

Tubingen, Germany), with a 5-µL Hamilton microsyringe (Esslab, Essex, UK), at the speed of 25 nl/min

for 10 min.

Two transgenic animal models were used for LH blood sampling under optogenetic stimulation:

UCN3Cre::VGAT-Flpo and VGAT-Cre-tdTomato. The following types of viral constructs were used in

separated cohorts of UCN3-Cre::VGAT-Flpo mice: (i) AAV-nEF-Con/Foff 2.0-ChRmine-oScarlet (250

nl, 2.1×10^{12} gc/mL, Serotype:8, #137161, Addgene, Massachusetts, USA); (ii) AAV-nEF-Con/Fon

2.0ChRmine-oScarlet (250 nl, 1.9×10^{12} gc/mL, Serotype:8, # 137159, Addgene, Massachusetts,

USA); (iii) a mixture of AAV-nEF-Con/Foff 2.0-ChRmine-oScarlet (2.1×10^{12} gc/mL, #137161) and

pAAV-nEF-Coff/Fon-NpHR3.3-EYFP (1.9×10^{12} gc/mL; #137159, final volume 300 nl); (iv) control virus

AAV-Ef1a-DIO-EYFP (250 nl, 4.4×10^{12} gc/mL, Serotype:9, #27056, Addgene, Massachusetts, USA).

VGATCre-tdTomato mice received unilateral injections of (i) channelrhodopsin (ChR2) viral construct,

(250 nl, 1.8×10^{12} gc/mL; Serotype:9; Addgene, #20298, Massachusetts, USA) or (ii) control virus,

AAVEf1a-DIO-EYFP (250 nl, 4.4×10^{12} gc/mL, Serotype:9; Addgene, # 27056, Massachusetts, USA)

into the MePD. Following injection, the needle was kept in position for 10 min before going up slowly

over 1 min. A unilateral fiber optic cannula (200 µm, 0.39 NA, 1.25 mm ceramic ferrule; Doric Lenses,

Quebec, Canada) was inserted into the right MePD 0.15 mm above the injection site. After reaching

the target, the fiber optic cannula was secured on the skull using dental cement (Super-Bond Universal

Kit, Prestige Dental, UK), bonded to the screws and skull. Finally, the incision in the scalp was closed

with suture. Animals were given 1 week to recover, after which they were handled 20 min daily to

acclimate to experimental procedures for a further 2 weeks.

For mini-endoscope *in vivo* calcium imaging, only UCN3-Cre::VGAT-Flpo transgenic animals were

used. Separate mice cohorts received the following viral injection into the MePD: (i) a mixture of pAAV-

Syn-Flex-rc[Chrimson-tdTomato] (2.0×10^{12} gc/mL, Serotype:5, #62723, Addgene, Massachusetts,

USA) + pAAV-EF1a-fDIO-GCaMP6s (2.3×10^{12} gc/mL; Serotype:8; #105715, Addgene,

Massachusetts, USA; final volume 400 nl); (ii) a mixture of pAAV-Syn-Flex-rc[ChrimsontdTomato]

(2.0×10^{12} gc/mL, Serotype:5, #62723, Addgene, Massachusetts, USA) +

pAAV.CAG.Flex.GCaMP6s.WPRE.SV40 (4.3×10^{12} gc/mL; Serotype:9; # 100842, Addgene,

Massachusetts, USA; final volume 400 nL); (iii) a mixture of control virus pAAV-hSyn-DIO-mCherry

(2.3×10^{12} gc/mL, Serotype:9, #50459) + pAAV-EF1a-fDIO-GCaMP6s (2.3×10^{12} gc/mL, Serotype:8,

#105715, final volume 400nl) . One week after administering the virus, a 600-μm diameter, 7.2-mm

long gradient index (GRIN) ProView™ Integrated Lens (Inscopix, PN:100-004554) was implanted at

the same location as the viral injection. The surface of the skull was scored, using the tip of a surgical

scalpel, and four screws were inserted into the skull. This was done to enhance the adhesion of dental

cement to the skull, thereby reducing the interference of movement on *in vivo* recording. Before

insertion of the GRIN lens, a 23 G needle was slowly lowered at a speed of 5.0 mm/min into the brain

(2.42 mm lateral, 1.52 mm posterior from bregma, depth of 5.05 mm below the skull surface) to make

a needle track to facilitate the insertion of the GRIN lens without causing brain distortion along its

route. The GRIN lens was then inserted into the brain at a rate of 0.6 mm/min, using the same

coordinates as the needle precutting. To secure the GRIN lens with integrated baseplate in position,

we used a light curing adhesive (iBond Universal, Kulzer GmbH, Leipziger, Germany) along with a

cured radiopaque hybrid dental cement (Gradia Direct Flo, GC corporation, Tokyo, Japan), bonding it

to the head screws and skull. A GRIN lens baseplate dust cover (Inscopix) was rescued in place to

protect the lens. After 4 weeks the mice were re-anaesthetized as described above and underwent

bilateral ovariectomy.

Serial blood sampling for LH measurement

The mouse's tail-tip was excised using a sterile scalpel. Then the chronically unilaterally implanted

fiber optic cannula was connected to a multimode fiber optic rotary joint patch cables (Thorlabs Ltd,

Ely, UK) via a ceramic mating sleeve to allow the animal freedom of movement. Mice were left for 1 h

to habituate. A Grass SD9B stimulator-controlled DPSS laser (Laserglow Technologies, Toronto,

Canada) was utilized to deliver either green light (532 nm wavelength, 10 Hz, 5 seconds on 5 seconds

off, 10 mW, 10-ms pulse width,) or blue light (473 nm wavelength, 5 Hz, 5 seconds on 5 seconds off,

5 mW, 10-ms pulse width,). The light pattern was controlled using software designed in house. After

1 h of acclimatization, serial blood samples (5 μL) for LH pulse measurement were collected every 5

min for 2 h between 10:00 - 12:00 h: 0-60 min for control blood sampling period, and 60-120 min under

stimulation or equivalent no-stimulation control period. Blood samples were diluted in 45 μL of

Phosphate-Buffered Saline (PBS) containing 0.2% Bovine Serum Albumin (BSA) and 0.05% Tween20

(PBST) and immediately placed on dry ice and stored at −80°C until later analysis.

LH assay

LH measurement was performed using an LH ELISA assay as reported previously[34]. A mouse LH

standard (mLH; AFP-5306A, NIDDK-NHPP, USA) was used to establish a standard curve for the

quantitative measurement. The following antibodies were used in the assay: (i) coating antibody

(RRID: AB_2665514, monoclonal antibovine LH beta subunit antiserum, 518B7, University of

California, CA, USA); (ii) anti-LH antibody (RRID: AB_2665533; National Hormone & Peptide

Program, CA, USA); (iii) secondary antibody (RRID: AB_772206, GE Healthcare, Chicago, Illinois,

USA). The inter-assay and intra-assay variations were 10.53% and 5.74%, respectively. The assay

sensitivity was 0.023 ng/mL.

LH pulse statistical analysis

LH pulses were verified using the Dynpeak algorithm in Scilab 5.5.2 programme[35]. The Dynpeak

algorithms were classified for OVX mice based on previously established parameters[36]. The average

LH inter-pulse interval was calculated for the 1-h control period and the 1-h stimulation period in both

experimental and control mice. Statistical significance was assessed using a 2-way ANOVA unless

otherwise stated. Data were represented as mean ± SEM and $p < 0.05$ was considered significant.

Mini-endoscope-based Calcium imaging

Optogenetic stimulation of MePD UCN3 neurons. Prior to recording, animals were handled daily

for 5 to 6 weeks to get accustomed to the experimental setup. The Data Acquisition (DAQ) box

(Inscopix, PN:100-005904) was turned on and connected to the computer wirelessly. The Data

Acquisition Software (IDAS; Inscopix, Inc.) was accessed through the Chrome web browser on the

computer communicating with the DAQ box. Adequate DAQ box storage capacity was checked

routinely before Calcium imaging. The GRIN lens baseplate dust cover was gently removed from the

mouse's head, after which the minscope camera was secured to the baseplate platform. Equipped

with the head-mount miniscope (nVoke2, Inscopix, SN: BB-11238601), the mouse was then placed

back into its home cage. The optimal focal plane was chosen by adjusting the lens focus to achieve

clear cell morphology. The exposure time was fixed at 50 ms. The frame rate was set to 20 Hz.

Subsequently, gain, and LED power were optimized based on image histograms before initiating

recording. Two types of LED were used during the recording: (i) EX-LED (0.2-0.4 mW, 435-460 nm

excitation filter) for GCaMP6s signal detection in the GABA or UCN3 neurons; (ii) OG-LED (10 mW,

590-650 nm excitation filter) for Chrimson optogenetic stimulation of the UCN3 neurons. After the mice

were attached to the miniscope, they were given 1-2 h for adaptation. For stimulation of the UCN3

neurons, the OG-LED was set to 10 Hz, 5 sec on 5 sec off, and a 10-ms pulse width. Once the animals

had acclimatized, the GCaMP6s recording of the GABA neurons or UCN3 neurons was started with

a protocol of 2 min baseline before stimulation for 2 min. Optogenetic stimulation was randomized for

the 10 Hz, with a 5-min interval between each to prevent photobleaching of GCaMP6s. Once the

experiment was concluded, the miniscope was gently detached, and the baseplate cover was

reattached and secured with its screw. The recordings were performed between 9:00 h and 15:00 h.

Three groups of animals were included in this experiment: (i) mice with UCN3 neurons expressing

opsin and GABA neurons expressing GCaMP; (ii) mice with UCN3 neurons expressing both opsin

and GCaMP, serving as a positive control; (iii) mice with UCN3 neurons expressing mCherry and

GABA neurons expressing GCaMP, serving as a negative control. To quantify intra-animal variability

at least two independent stimulation trials were conducted with each animal, with a maximum of four

trials per animal.

Restraint stress. The system was set up and the mouse was connected to the miniscope as

described above but with the OG-LED turned off. Once the mouse had settled, the experiment started

with a 3 min baseline recording, followed by 3 min of recording under hand-held restraint. Finally, the

miniscope was detached and the baseplate cover was reattached. The experiments were performed

between 9:00 h and 15:00 h. The animals in this experiment were the same as those used in the

optogenetic stimulation protocol described above but were divided into two groups depending on

whether GCaMP was expressed in the UCN3 or GABA neurons, regardless of whether opsin or

mCherry was expressed in the UCN3 cells. A single trial per animal was performed to minimize

unnecessary distress, in accordance with the 3Rs principle, apart from one animal in which the

restraint stress experiment was performed twice to ensure this does not result in habituation.

In vivo calcium imaging recording processing

The video data, with a resolution of 1280 pixels x 800 pixels, was exported from the DAQ box through

the file manager to a desktop computer and processed with the Inscopix Data Processing Software

(IDPS). Within the IDPS, there are six essential steps for extracting single-cell calcium transient data

from the recorded video: down-sampling, cropping, spatial filtering, motion correction, trace

normalization ($\Delta F/F_0$), and cell identification. First, the video underwent both spatial down-sampling

and temporal down-sampling by a factor of 2 and was cropped to enhance processing speed, focusing

on the region of interest where the GCaMP signal was observed. Spatial filtering, with a threshold

ranging between 0.005 (low) and 0.5 (high) was used to distinguish individual pixels in the video

image, creating a high-contrast image. A mean intensity projection image was generated. Motion

correction was subsequently conducted, using the mean intensity projection image as a reference.

Motion correction was applied again using one of the frames affected by movement while

simultaneously drawing a region of interest (ROI) where UCN3 accumulates to enhance the

effectiveness of motion correction. For the recordings with GCaMP signals from GABA neurons, we

did not use the built-in motion correction function of IDPS. Instead, we used the MATLAB (version

R2022a) implementation of the open-source algorithm NoRMCorre[37] with default parameters for both

rigid and non-rigid (patch size 64 pixels and overlap 32) motion correction as long as a minimum

correlation of the frames with the mean frame above 0.8 is achieved. This was performed by doubling

the patch size and their overlap to 128 and 64, respectively. Corrected videos were saved in TIFF

format for further analysis via IDPS. Following motion correction, pixel intensity, $F(x, y)$, in each frame

of the movie was normalized using the following formula $\frac{(F(x,y) - F_{baseline}(x,y))}{F_{baseline}(x,y)}$, where $F_{baseline}(x, y)$ is

the baseline pixel intensity calculated by taking the mean value of the pixel across the entire movie.

Finally, regions of interest were identified automatically using the cell identification algorithm provided

by IDPS, based on principal component and independent component analysis (PCA-ICA algorithm).

Validation of virus injection, fiber optic cannula and GRIN lens implantation

Once the experimental procedures were completed, a lethal dose of ketamine was used to humanely

euthanize the mice. The mice were perfused transcardially with heparinized saline (5 U/ml) for 5 min,

followed by ice-cold 4% paraformaldehyde (PFA) for 15 min using a pump (Minipuls,156 Gilson,

Villiers Le Bel, France). The brains were collected and post-fixed sequentially at 4°C in a solution of

15% sucrose in 4% PFA and later in a solution of 30% sucrose in phosphate buffer until they sank.

They were then snap frozen on dry ice and stored in -80°C until processing. The brains were coronally

sectioned (30 µm) between +0.5 mm and -2.7 mm from the bregma using a cryostat (Bright Instrument

Co., Huntingdon, UK), and sections were mounted on microscope slides, air-dried, and covered with

slips using Prolonged Antifade mounting medium (Molecular Probes, Inc. OR, USA). The accuracy of

injection sites, fiber optic cannula and GRIN lens implantation were confirmed by Axioskop 2 Plus

microscope equipped with AXIOVISION 4.7 (Zeiss). Images were captured using Axioskop 2 Plus

microscope (Carl Zeiss). Only the data of mice with correct viral infection and implantation of cannula

and lens in the MePD were included in the analysis.

Immunohistochemistry

To assess the targeting of viral labeling to UCN3 neurons in the MePD, we used the indirect

immunofluorescence technique to localize UCN3 neurons. Every 4[th] section from each brain was

washed in potassium phosphate-buffered saline (KPBS) and incubated in blocking solution (KPBS

containing 1% BSA, 0.03% Triton X-100 and 5 mg/mL heparin) at room temperature for 1 hour

followed by incubated in a rabbit polyclonal anti-UCN3 antibody (1:2000 dilution)[38] in the blocking

solution (Code: PBL 7218, kindly provided by J. Vaughan, Salk Institute, La Jolla, CA, USA) for 48

hours at 4°C. The brain sections were then processed with goat anti-rabbit IgG (H + L), a biotinylated

secondary antibody (1:1000 dilution; Cat. No. BA-1000, Vector Laboratories, Burlingame, CA, USA) overnight at 4°C. After rinses in KPBS, visualization of UCN3 immunoactivity was achieved using Streptavidin, Alexa Fluor 405 conjugate (1:200 dilution; Cat. No. S 32351, Thermo Fisher Scientific, Waltham, MA, USA) for 2 hours at room temperature. The sections were finally washed, mounted onto slides and cover slipped. Omission of the UCN primary antibody and its pre-incubation resulted in the absence of specific staining. Brain sections from each experimental group were processed on the same day to control for inter-batch variability. To preserve the fluorescence of the virally expressed proteins, all subsequent steps were performed in a dark box. Semiquantitative analysis of immunostaining data was carried out on AxioVision microscope image system. UCN3-positive neurons were identified by Alexa Fluor 405 fluorescence, while native fluorescence (e.g. tdTomato or oScarlet) was used for virus-transduced neurons identification. The number of virus-labeled, UCN3-immunoreactive cells and co-localized cells in the MePD were quantified on 4 sections from each mouse brain. Targeting efficiency was defined as the percentage of virus-labeled cells that were also positive for UCN3.

Data analysis

The data processing protocol we employ in this study was performed in two phases: pre-intervention and intervention (either via optogenetic stimulation or restraint stress). For our analysis, we use the smoothed deconvolved traces generated by the OASIS method[39], which improves the signal-to-noise ratio (SNR) and mitigates noise interference. All traces are rescaled between 0 and 1 (min-max normalization), and then further normalized to mean zero and unit variance. The analysis process is divided into three key steps: i) identification of GABA cells co-expressing UCN3 and removing them from further analysis; ii) estimating and comparing functional connectivity before and during intervention; iii) clustering analysis based on pairwise association-derived dissimilarity scores. In step (i), we use power spectral density (PSD) analysis and K-means clustering (using Euclidean distance) to identify GABA neurons with activity that mimics the pattern of optic stimulation, evidence that they are co-expressing UCN3.

PSD filtering. For each trace, we compute the PSD during the stimulation phase. Cells with PSD

values at 0.1 Hz (the frequency of the stimulus 5 sec and 5 sec off cycle) above 0.37 Hz^{-1} are labelled

as co-expressing UCN3. This threshold is determined from recordings of UCN3 cells under optic

stimulation, by calculating the median of the minimum PSD values at 0.1 Hz from each dataset (see

Supplementary Fig. 2**B**).

K-means clustering. Using the Euclidean distance metric, we cluster the traces during the stimulation

phase into two groups, aiming to separate neurons into GABA-only neurons and neurons co-

expressing UCN3 and GABA. Clustering separated a small group of neurons with activity mimicking

the optic stimulation oscillatory pattern (see Supplementary Fig. 2**C**). Recordings of MePD UCN3

neurons shows that under optogenetic stimulation their activity rises and declines in phase with on

and off periods of stimulation (shown in Fig. 3**A**); therefore, this small number of cells was labeled as

co-expressing UCN3.

The union of neurons identified by either method as co-expressing UCN3 and GABA are excluded

from further analysis. By applying both filters, we aim to robustly identify purely GABA-expressing

neurons. We acknowledge the possibility that some cells may pass through the filters or that a few

purely GABA-expressing neurons may be inadvertently excluded. However, our dual approach of

employing PSD analysis and K-means clustering minimizes this possibility and is essential to ensure

our analysis remains focused exclusively on GABA-expressing neurons. Nevertheless, we take a

further step and visually inspect the calcium recording traces for any potential anomalies such as

regular oscillations in-phase with the stimulation and/or massive increase in calcium level during the

intervention period. Any identified anomalies of this sort are also dropped from datasets.

Comparing pre-intervention and intervention cell-cell interaction strength

Following step (i), the remaining dataset contains only traces associated with GABA neurons, which

are not directly activated by our optic stimulation protocol. In step (ii), we assess the functional

connectivity between the observed neurons using Pearson's correlation[40] defined as

773
$$corr(x,y) = \frac{R(x,y)}{\sqrt{R(x,x)R(y,y)}},$$

where $R(x, y) = \sum_{i=1}^{N} (x_i - \bar{x})(y_i - \bar{y})$ represents cross-correlation with N as the length of x and y.

Using the above measures, we construct the corresponding cross-correlation matrices (see for an

example Fig. 1**H&I**).

To evaluate changes in pairwise cell interactions between the pre-intervention and intervention

phases, we compute the pairwise Riemannian distance[41] between correlation matrices from pre-

intervention and intervention episodes. The Riemannian distance here is defined as

$$d(A, B) = \left(\sum_i \ln (\lambda_i)^2 \right)^{1/2},$$

where ln is the natural logarithm and λ indicates the joint eigenvalues of symmetric positive definite

(SPD) cross-correlation matrices A and B. The size of the matrices directly affects the Riemannian

distance, therefore, we normalize each Riemannian distance by the average Riemannian distance

computed from 100 pairs of randomly generated SPD matrices of the same size. To make sure that

random generation is uniform in the space of SPD matrices, we employ the Metropolis-Hasting

method introduced in[42]. Normalizing the Riemannian distances guarantees that the distance is not

influenced by matrix size. To assess the effect of experimental conditions on Riemannian distances

while accounting for the hierarchical structure of the data, we employed a linear mixed-effects model[43].

Specifically, we modeled repeated measurements nested within individual animals, as follows:

$$y \sim condition + (1\,|Animal) + (1\,|\,Animal\!:\!Repeat),$$

where y denotes the computed Riemannian distance, while condition represents whether the

experiment was control, stimulation or restraint stress. The model was fitted using maximum likelihood

estimation.

Clustering analysis

In step (iii), we apply agglomerative hierarchical clustering (using the average linkage criterion;

package in scikit-learn Python™) to classify cells based on the dissimilarity of their calcium activity,

given as

$$D(x,y) = 1 - corr(x,y).$$

To determine the optimal number of clusters and robustness of clustering, we employed a bootstrapping technique. This involved resampling (without replacement) 75% of the cells and clustering the resulting dataset with different numbers of clusters (2, 3, and 4) using hierarchical or K-means clustering with either the covariance or correlation as the dissimilarity measure. This resampling process was repeated 100 times.

For each resample dataset, we evaluate the clustering performance, using the Silhouette Coefficient (SC)[42] which measures the cohesion within clusters and their separation, with values ranging from -1 (poor performance) to 1 (good performance). Results of the evaluation are given in Supplementary Figs. 4-6, showing that the SC is significantly higher for 2 clusters compared to 3 or 4 clusters in the case of both hierarchical and K-means clustering.

Modeling

We build on our previously published modeling framework of MePD circuit[11] by incorporating dynamic stimulation by UCN3 neurons as observed experimentally (as shown in Fig. 3). To this end we define input from UCN3 neurons into the populations of glutamatergic neurons and GABA interneurons with a strength given by the parameter γ. The activity of UCN3 neurons during stimulation is approximated via sinusoidal function:

$$U_{stim}(t) = \gamma B + A sin(2\pi f t),$$

where the parameter f is the frequency of the wave. Parameter B represents the constant excitatory drive, potentially mediated by slow neuropeptide action, while the term $A sin(2\pi f t)$ describes faster neurotransmitter-driven-driven input aligned with optogenetic stimulation frequency. We approximate UCN3 neuron activity during restraint stress using two exponential functions:

$$U_{stress}(t) = \gamma \big(at \exp(-r_2 t) + b(1 - \exp(-r_1 t))\big),$$

where the function initially increases with magnitude a then decreases to a plateau value

given by the parameter b. The parameters r_2 and r_1 are responsible for the rate of increase

of the first term and rate of decrease to a plateau in the second term.

Briefly, the model captures the dynamics of the MePD intracellular calcium activity of three

populations: glutamatergic neuronal population (excitatory) and the populations of GABA interneurons

and GABA efferent neurons (inhibitory). We build on our previous work[11] by assuming that excitatory

(UCN3) input is provided to the populations of GABA interneurons and glutamate (Fig 3). Previous

analysis of the model indicated that negative feedback between GABA and glutamatergic neuronal

populations as well as glutamatergic self-excitation are critical mechanisms for inducing oscillatory

dynamics[11]. However, providing glutamatergic inputs to both populations of GABA neurons results in

a null phase difference between the populations. Therefore, we opted to provide glutamatergic input

only to the population of GABA efferent neurons.

To investigate how the amygdala modulates the activity in the GnRH pulse generator, we couple the

above-described MePD model to our KNDy network model, where the oscillatory dynamics of the

KNDy acts as a proxy of LH pulsatility[26]. Here we couple the model via providing only inhibitory output

from MePD GABA efferent neurons population to KNDy, as it was previously shown that glutamate

plays a modulatory role to the MePD output rather than directly affecting the KNDy network[11]. We then

calibrate the coupled model by reproducing the experimental results from Ivanova et al.[23]

(Supplementary Fig. 8). Further modeling details, such as equations governing model dynamics and

model parameter values, can be found in Supplementary Information. The simulations have been

done in MATLAB (version R2023b).

Data Availability

843 Source data are provided with this paper. Further information that supports the findings of this study

844 is available from the corresponding authors upon request.

Code Availability

The code for reproducing the data analysis (Python™), mathematical modeling (MATLAB), and figure generation for both the main text and Supplementary information, along with the datasets used in the analysis, is publicly available at: https://github.com/mv-kr/MePD_GABA.

References

1. Yeo, S.-H., Kyle, V., Blouet, C., Jones, S. & Colledge, W. H. Mapping neuronal inputs to Kiss1 neurons in the arcuate nucleus of the mouse. *PLOS ONE* **14**, e0213927 (2019).

2. Moore, A. M., Coolen, L. M. & Lehman, M. N. Kisspeptin/Neurokinin B/Dynorphin (KNDy) cells as integrators of diverse internal and external cues: evidence from viral-based monosynaptic tract-tracing in mice. *Sci. Rep.* **9**, 14768 (2019).

3. Li, X. F. & O'Byrne, K. T. Stress and the Reproductive System. in *Knobil and Neill's Physiology of Reproduction: Two-Volume Set* vol. 2 1637–1660 (ELSEVIER SCIENCE INC, 2014).

4. Yu, J., Li, X.-F., Tsaneva-Atanasova, K., Zavala, E. & O'Byrne, K. T. Chemogenetic activation of PVN CRH neurons disrupts the estrous cycle and LH dynamics in female mice. *Front. Endocrinol.* **14**, 1322662 (2023).

5. Kreisman, M. J., Tadrousse, K. S., McCosh, R. B. & Breen, K. M. Neuroendocrine Basis for Disrupted Ovarian Cyclicity in Female Mice During Chronic Undernutrition. *Endocrinology* **162**, bqab103 (2021).

6. Bethea, C. L., Centeno, M. L. & Cameron, J. L. Neurobiology of Stress-Induced Reproductive Dysfunction in Female Macaques. *Mol. Neurobiol.* **38**, 199–230 (2008).

7. Herod, S. M., Pohl, C. R. & Cameron, J. L. Treatment with a CRH-R1 antagonist prevents stress-induced suppression of the central neural drive to the reproductive axis in female macaques. *Am. J. Physiol. Endocrinol. Metab.* **300**, E19-27 (2011).

8. Nair, B. B. *et al.* Impact of chronic variable stress on neuroendocrine hypothalamus and pituitary in male and female C57BL/6J mice. *J. Neuroendocrinol.* **33**, e12972 (2021).

9. Comninos, A. N. *et al.* Kisspeptin signaling in the amygdala modulates reproductive hormone secretion. *Brain Struct. Funct.* **221**, 2035–2047 (2016).

10. Lass, G. *et al.* Optogenetic stimulation of kisspeptin neurones within the posterodorsal medial

amygdala increases luteinising hormone pulse frequency in female mice. *J. Neuroendocrinol.* **32**,

e12823 (2020).

11. Nechyporenko, K. *et al.* Neuronal network dynamics in the posterodorsal amygdala: shaping

reproductive hormone pulsatility. *J. R. Soc. Interface* **21**, 20240143 (2024).

12. Adekunbi, D. A. *et al.* Role of amygdala kisspeptin in pubertal timing in female rats. *PLoS ONE*

12, e0183596 (2017).

13. Ivanova, D. *et al.* Role of Posterodorsal Medial Amygdala Urocortin-3 in Pubertal Timing in Female

Mice. *Front. Endocrinol.* **13**, (2022).

14. Adekunbi, D. A. *et al.* Kisspeptin neurones in the posterodorsal medial amygdala modulate sexual

partner preference and anxiety in male mice. *J. Neuroendocrinol.* **30**, e12572 (2018).

15. Wu, Y. E. *et al.* Neural control of affiliative touch in prosocial interaction. *Nature* **599**, 262–267

(2021).

16. Ivanova, D. *et al.* Urocortin3 in the Posterodorsal Medial Amygdala Mediates Stress-induced

Suppression of LH Pulsatility in Female Mice. *Endocrinology* **162**, 1–14 (2021).

17. McIntyre, C. *et al.* GABA Signaling in the Posterodorsal Medial Amygdala Mediates Stress-

induced Suppression of LH Pulsatility in Female Mice. *Endocrinology* **164**, 1–11 (2023).

18. Keshavarzi, S., Sullivan, R. K. P., Ianno, D. J. & Sah, P. Functional Properties and Projections of

Neurons in the Medial Amygdala. *J. Neurosci.* **34**, 8699–8715 (2014).

19. Choi, G. B. *et al.* Lhx6 delineates a pathway mediating innate reproductive behaviors from the

amygdala to the hypothalamus. *Neuron* **46**, 647–660 (2005).

20. Bian, X. Physiological and morphological characterization of GABAergic neurons in the medial

amygdala. *Brain Res.* **1509**, 8–19 (2013).

21. Lewis, K. *et al.* Identification of urocortin III, an additional member of the corticotropin-releasing

factor (CRF) family with high affinity for the CRF2 receptor. *Proc. Natl. Acad. Sci. U. S. A.* **98**,

7570–7575 (2001).

22. Shemesh, Y. *et al.* Ucn3 and CRF-R2 in the medial amygdala regulate complex social dynamics.

Nat. Neurosci. **19**, 1489–1496 (2016).

23. Ivanova, D., Li, X.-F., McIntyre, C. & O'Byrne, K. T. Posterodorsal Medial Amygdala Urocortin-3,
GABA, and Glutamate Mediate Suppression of LH Pulsatility in Female Mice. *Endocrinology* **164**,
1–13 (2023).

24. Yuste, R., Cossart, R. & Yaksi, E. Neuronal ensembles: Building blocks of neural circuits. *Neuron*
112, 875–892 (2024).

25. Arrigoni, E. & Saper, C. B. What optogenetic stimulation is telling us (and failing to tell us) about
fast neurotransmitters and neuromodulators in brain circuits for wake-sleep regulation. *Curr. Opin.*
Neurobiol. **29**, 165–171 (2014).

26. Voliotis, M. *et al.* The Origin of GnRH Pulse Generation: An Integrative Mathematical-Experimental
Approach. *J. Neurosci.* **39**, 9738–9747 (2019).

27. Hochgerner, H. *et al.* Neuronal types in the mouse amygdala and their transcriptional response to
fear conditioning. *Nat. Neurosci.* **26**, 2237–2249 (2023).

28. Li, Y. *et al.* Neuronal Representation of Social Information in the Medial Amygdala of Awake
Behaving Mice. *Cell* **171**, 1176-1190.e17 (2017).

29. Pardo-Bellver, C., Cádiz-Moretti, B., Novejarque, A., Martínez-García, F. & Lanuza, E. Differential
efferent projections of the anterior, posteroventral, and posterodorsal subdivisions of the medial
amygdala in mice. *Front. Neuroanat.* **6**, 33 (2012).

30. Polepalli, J. S., Gooch, H. & Sah, P. Diversity of interneurons in the lateral and basal amygdala.
Npj Sci. Learn. **5**, 10 (2020).

31. Patel, A. H. *et al.* Kisspeptin in functional hypothalamic amenorrhea: Pathophysiology and
therapeutic potential. *Ann. N. Y. Acad. Sci.* **1540**, 21–46 (2024).

32. Barbagallo, F. *et al.* Gene-environment interaction in functional hypothalamic amenorrhea. *Front.*
Endocrinol. **15**, 1423898 (2024).

33. Paxinos, G. & Franklin, K. B. J. *Paxinos and Franklin's the Mouse Brain in Stereotaxic*
Coordinates. (2019).

34. Steyn, F. J. *et al.* Development of a Methodology for and Assessment of Pulsatile Luteinizing
Hormone Secretion in Juvenile and Adult Male Mice. *Endocrinology* **154**, 4939–4945 (2013).

35. Vidal, A., Zhang, Q., Médigue, C., Fabre, S. & Clément, F. DynPeak: An Algorithm for Pulse
Detection and Frequency Analysis in Hormonal Time Series. *PLOS ONE* **7**, e39001 (2012).

36. Yang, J. A. *et al.* Acute Psychosocial Stress Inhibits LH Pulsatility and Kiss1 Neuronal Activation
in Female Mice. *Endocrinology* **158**, 3716–3723 (2017).

37. Pnevmatikakis, E. A. & Giovannucci, A. NoRMCorre: An online algorithm for piecewise rigid motion
correction of calcium imaging data. *J. Neurosci. Methods* **291**, 83–94 (2017).

38. Battagello, D. S. *et al.* Anatomical Organization of Urocortin 3-Synthesizing Neurons and
Immunoreactive Terminals in the Central Nervous System of Non-Human Primates [Sapajus spp.].
Front. Neuroanat. **11**, (2017).

39. Friedrich, J., Zhou, P. & Paninski, L. Fast online deconvolution of calcium imaging data. *PLOS*
Comput. Biol. **13**, (2017).

40. Rice, J. A. *Mathematical Statistics and Data Analysis*. (Thomson/Brooks/Cole, Belmont, CA,
2007).

41. Moakher, M. A Differential Geometric Approach to the Geometric Mean of Symmetric Positive-
Definite Matrices. *SIAM J. Matrix Anal. Appl.* **26**, 735–747 (2005).

42. Córdoba, I., Varando, G., Bielza, C. & Larrañaga, P. A Fast Metropolis-Hastings Method for
Generating Random Correlation Matrices. in *Intelligent Data Engineering and Automated Learning*
– IDEAL 2018 (eds. Yin, H., Camacho, D., Novais, P. & Tallón-Ballesteros, A. J.) 117–124 (Springer
International Publishing, Cham, 2018). doi:10.1007/978-3-030-03493-1_13.

43. Harrison, X. A. *et al.* A brief introduction to mixed effects modelling and multi-model inference in
ecology. *PeerJ* **6**, e4794 (2018).

Acknowledgements

We thank Ms Sumi Mathew for her technical support with genotyping mice.

S.F., M.V., K.T.A., K.T.O., X.F.L. gratefully acknowledge the financial support of BBSRC via grants
BB/W005913/1 (KCL), BB/W005883/1 (Exeter) and BB/S019979/1. KTA gratefully acknowledges the
financial support of the EPSRC via grant EP/T017856/.

Contributions

J.Y. contributed data acquisition and data analysis, S.F. contributed data analysis, K.N. contributed modeling, M.V. contributed managing data analysis, X.F.L. contributed managing data acquisition and analysis, H.Y. contributed data acquisition, Y.L. contributed data acquisition, J.Y. contributed data acquisition, O.H. contributed data acquisition, R.D.B. contributed data acquisition, B.S. contributed data acquisition, K.T.O., K.T.A., X.F.L. and M.V. contributed to managing the project. All authors contributed to writing the manuscript.

Competing interests

The authors declare no competing interests.

Corresponding author

Correspondence and requests for materials should be addressed to Krasimira Tsaneva-Atanasova (K.Tsaneva-Atanasova@exeter.ac.uk), Kevin T. O'Byrne (kevin.o'byrne@kcl.ac.uk), Margaritis Voliotis (m.voliotis@exeter.ac.uk).

Open Access

42