Amygdala GABA Neurons: Gatekeepers of Stress and Reproduction

Author Information

Junru Yu,[1,†] Saeed Farjami,[2,3,†] Kateryna Nechyporenko,[2,3,†] Xiao Feng Li,[1] Hafsa Yaseen,[1,4] Yanyan Lin,[1] Jinbin Ye,[1] Owen Hollings,[1] Ross de Burgh,[1] Kevin T. O'Byrne,[1,*] Krasimira Tsaneva-Atanasova,[2,3,5,*] Margaritis Voliotis,[2,3,*]

Affiliations

[1]Department of Women and Children's Health, School of Life Course and Population Sciences, King's College London, Guy's Campus, London SE1 1UL, UK, [2]Department of Mathematics and Statistics, [3]Living Systems Institute, [4]Biological Sciences, University of Missouri, Columbia, MO 65211-7400 and [5]EPSRC Hub for Quantitative Modelling in Healthcare, University of Exeter, Stocker Road, Exeter EX4 4PY, UK

Abstract

Stress can disrupt menstrual cycles, cause infertility, and lead to other reproductive disorders. The posterodorsal medial amygdala (MePD) processes stress signals and regulates the gonadotropin-releasing hormone (GnRH) pulse generator through GABAergic inhibitory projections to the hypothalamus. However, how stress is processed in the MePD - especially involving its dense GABAergic and Urocortin-3 (UCN3) neurons - remains poorly understood.

In this study, we combine *in vivo* GRadient-INdex (GRIN) lens mini-endoscopic calcium imaging (to track neuronal activity), optogenetics, clustering analysis, and computational modeling to investigate MePD circuitry. Our findings reveal two anti-correlated GABAergic sub-populations in the MePD that dictate responses to both UCN3 neuron stimulation and restraint stress. Our computational modeling suggests that mutual inhibition between these GABAergic groups drives this anti-correlated activity and predicts how these interactions shape downstream responses to stimulation of GABAergic and UCN3 neurons.

We test these predictions using optogenetics and confirm that GABAergic neurons operate downstream of the UCN3 population, playing a crucial role in transmitting UCN3 signals to regulate luteinizing hormone (LH) pulse frequency. Our study is the first to show how GABAergic neurons in the amygdala mediate stress effects on reproductive health, uncovering key neural mechanisms linking emotional and reproductive functions.

Introduction

Successful reproduction requires normal function of a neural oscillator, the gonadotropin-releasing hormone (GnRH) pulse generator, that drives the pulsatile release of gonadotropic hormones, luteinizing hormone (LH) and follicle stimulating hormone (FSH) to control ovulation and spermatogenesis. This oscillator comprises Kisspeptin neurons co-expressing Neurokinin-B (NKB) and Dynorphin A (acronym KNDy) in the hypothalamic arcuate nucleus (ARC). The KNDy neural network not only provides the essential episodic stimulatory kisspeptin signal to the GnRH neurons, but receives diverse afferent inputs and hence is a hub for integrating various internal and external cues that regulates reproductive function[1,2]. The most common cause of insufficient GnRH drive is functional, that is, due to neuroendocrine effects of psychological and behavioral variables.

The amygdala, a part of the limbic brain typically associated with emotions and anxiety, has strong projections to the KNDy system[1,2]. The medial amygdala, and more specifically its posterodorsal subnucleus (MePD), has been shown to be a robust upstream regulator of GnRH pulse generator frequency[3–5], pubertal timing[6,7], sociosexual behaviour[8,9], anxiety[8] and psychological stress-induced suppression of pulsatile LH secretion[10,11].

In the MePD, the majority of neurons are GABAergic, accounting for a substantial portion of its projections, many of which target important hypothalamic centers involved in reproduction[12–14], including the KNDy neurons and optogenetic stimulation of the latter suppress LH pulsatility[11]. Moreover, chemogenetic inhibition of GABA neurons in the MePD can reverse the inhibitory effect of psychogenic stress on LH pulsatility[11], suggesting an involvement of GABAergic signaling in the stress-induced suppression of pulsatile LH secretion. Although the precise way in which the MePD GABAergic neurons influence LH pulsatility is not fully understood, it has been suggested that they interact with the stress neuropeptide urocortin-3 (UCN3) neurons to modulate or reinforce stress-induced suppression of LH pulse frequency[10].

UCN3, part of the corticotropin-releasing factor (CRF) superfamily and selective for CRF type 2 receptors (CRFR2)[15], is highly expressed in the MePD[10,16]. Furthermore, UCN3 neurons in the MePD mediate psychosocial stress-induced suppression of the GnRH pulse generator and corticosterone secretion, thus play a central role in the interaction between the reproductive and stress axes[10,17]. Previously we have shown that intra-MePD GABA antagonism can reverse UCN3-induced LH pulse suppression[17], suggesting that GABA acts downstream of UCN3 signalling within the MePD. However, the local neural circuitry within the MePD mediating stress related information to the GnRH pulse generator remains to be established.

Here, we employed GRadient INdex (GRIN) lens mini-endoscopy to monitor and compare the calcium activity of GABA neurons at single-cell resolution under acute psychological stress or optogenetic stimulation of UCN3 neurons. Clustering analysis of the observed calcium transients uncovered two distinct GABAergic sub-populations exhibiting anti-correlated activity. To explore the functional role of these two sub-populations and unpick the mechanisms driving their anti-correlated activity, we build upon our previously developed MePD circuitry mathematical modeling framework[5]. Our modeling reveals that mutual inhibitory interactions between the GABAergic populations are sufficient to drive anti-correlated activity and predicts a complementary inhibitory effect of GABA and UCN3 signaling on the frequency of the GnRH pulse generator, which we tested by applying Cre/Flp recombinase intersectional strategies to selectively target and manipulate heterogeneous MePD neuronal populations[11] . Hence, by uncovering precise interactions between distinct sub-populations of MePD GABAergic neurons and UCN3 neurons in transmitting stress signals to the GnRH pulse generator, we provide critical and previously unknown insights into the neural circuitry driving stress-induced reproductive dysfunction. This discovery significantly advances our understanding of how emotional stress directly impacts reproductive health and identifies potential neural targets for therapeutic intervention in stress-related reproductive disorders.

Results

Dynamic reconfiguration of MePD GABA neuronal networks during selective optogenetic stimulation of urocortin-3 (UCN3) neurons and restraint stress

To investigate how the posterodorsal subnucleus of the medial amygdala (MePD) processes stress signals and to understand the role of GABAergic and UCN3 neurons in this process, we employed *in vivo* calcium imaging using GRadient INdex (GRIN) mini-endoscopy. A cohort of six female UCN3-Cre::VGAT-Flpo transgenic mice were injected with a viral mixture containing pAAV-Syn-Flex-rc[Chrimson-tdTomato] and pAAV-EF1a-fDIO-GCaMP6s. This intersectional viral strategy enabled dual functionality: (i) optogenetically stimulating MePD UCN3 neurons via Chrimson; and (ii) detecting calcium activity in MePD GABAergic neurons using GCaMP6s (Fig. 1**A**). GRIN lenses were surgically implanted at the same location as the viral injection, enabling imaging of calcium activity with single-neuron resolution.

Two series of experiments were conducted to assess GABAergic responses under distinct conditions. First, to assess the role of UCN3 neurons in recruiting GABAergic response, baseline calcium activity from MePD GABAergic neurons was recorded for 2 min, followed by 2 min of recording during optogenetic stimulation of UCN3 neurons (Fig. 1**A**). Optogenetic stimulation was delivered in a discontinuous pattern, 5 seconds on, 5 seconds off, with a light-pulse frequency of 10 Hz during the 'on' periods (10-ms pulse width, 10 mW power, and 532 nm wavelength). To further characterize the GABAergic response to acute stressors, baseline calcium activity of GABAergic neurons was recorded for 2 min, followed by 2 min of recording while the animals were exposed to restraint stress (Fig. 1**A**).

The dynamic GABAergic response to optogenetic stimulation of UCN3 neurons and restraint stress was summarized in functional terms using pairwise (neuron-neuron) association matrices, constructed using the signed lagged cross-correlation (SLxCorr) metric. SLxCorr captures the correlation between the activity of two neurons while accounting for both temporal lags and the sign (positive or negative) of their association (see Methods). Hence, SLxCorr matrices represent the

MePD GABAergic response as a set of interconnected nodes (neurons) with connections weighted

by their functional interactions (Fig. 1**H&I**). Our analysis reveals significant differences in the

functional connectivity of the GABAergic network in both experimental conditions (Fig. 1**J**) that we

also confirm using an alternative measure of pairwise association, namely covariance (see Methods

and Supplementary Fig. 1). These results robustly indicate that the GABAergic population in the

MePD plays a pivotal role in processing stress signals, such as those induced by restraint stress.

Furthermore, our findings suggest that UCN3 neurons may serve as a critical relay, transmitting

stress-related information to the GABAergic population.

Figure 1: **Functional connectivity of GABAergic neurons in posterodorsal medial amygdala (MePD)**

reconfigures in response to optogenetic stimulation of urocortin-3 (UCN3) neurons or acute restraint

stress. (**A**) Schematic diagram of the experimental design for stimulating MePD UCN3 neurons and

simultaneously imaging calcium activity from MePD GABAergic neurons. Heatmaps of calcium activity (ΔF/F)

in response to (left) optogenetic stimulation of UCN3 neurons and (right) during restraint-induced stress. (**B**-**G**)

Representative photomicrographs of the MePD from a UCN3-Cre::VGAT-Flpo mouse injected with pAAV-Syn-

Flex-rc+pAAV-EF1a-fDIO-GCaMP6s. (**B&E**) Red fluorescence (tdTomoato) labels UCN3 neurons expressing

Chrimson-tdTomato, (**C&F**) green fluorescence labels GABAergic neurons, and (**D&G**) red and green signals

merged. (**H&I**) Functional connectivity in the GABA populations was quantified using signed lagged cross-

correlation (SLxCorr). (**J**) Functional connectivity weights show a statistically significant difference between

the baseline period and subsequent UCN3 stimulation or restraint stress across animals, assessed via

Kolmogorov-Smirnov tests for each animal trial and using Fisher's method for combining p-values across data

sets recorder in different animals (see Methods). Distribution of combined p-values obtained from different

combinations of trials (one from each animal) is consistently below the 0.05 confidence level. Scale bars: **B**-

D, 200 µm; **E**-**G**, 50 µm.

Clustering analysis reveals two functionally distinct GABAegic sub-populations in the MePD

To better understand the functional organization of the GABAergic MePD network during

optogenetic stimulation of UCN3 neurons and under restraint stress, we performed clustering

analysis of the calcium recordings (time series) from GABAergic neurons.

The Signed Lagged Cross-Correlation (SLxCorr) was transform into a dissimilarity score, based on

which we performed hierarchical clustering. Clustering results consistently revealed two functionally

distinct cell populations both during optogenetic stimulation of UCN3 neurons and restraint stress

(Fig. 2**A&C**). Under both conditions, activity between the two clusters appeared highly anti-

correlated as illustrated by two representative examples depicted in Fig. 2**B&D**. Moreover, we find

that clusters are significantly different in size with the smallest cluster accounting on average for

35% of all cells (Fig 2**D&E**, Supplementary Fig. 2). The robustness of the above finding was also

verified using the K-means clustering algorithm and pairwise covariance as an alternative

dissimilarity score (see Methods and Supplementary Figs. 3-5).

Based on the functional role of the MePD in relaying stress signals to the hypothalamic arcuate

nucleus (ARC), our results suggest that the identified GABAergic sub-populations (clusters) could

correspond to distinct populations of GABA interneurons and GABA efferent neurons projecting to the ARC. These projections play a role in regulating the gonadotropin-releasing hormone (GnRH) pulse generator, a key driver of reproductive function[5,11].

Figure 2: **Clustering identifies two functionally distinct GABAergic sub-populations in the posterodorsal medial amygdala (MePD) both during UCN3 optogenetic stimulation and restraint stress**. Representative example of (**A**) a clustered connectivity matrix and (**B**) associated calcium time-traces of MePD GABA neurons during optogenetic stimulation of UNC3 neurons. Representative examples of (**C**) a clustered connectivity matrix and (**D**) associated calcium activity time-traces of MePD GABA neurons during restraint stress. (**E&G)** Relative sizes of the two GABAergic sub-populations identified via clustering analysis. (##: p<0.001).

Mutually inhibitory interactions between GABA populations drive anti-correlated activity

To explore the functional role of the two GABAergic sub-populations identified through clustering analysis and unpick the mechanisms driving their anti-correlated activity, we build upon our previously developed MePD circuitry mathematical modeling framework[5]. The model captures the dynamics of intracellular calcium activity in sub-populations of GABA neurons (we refer to them as interneurons and efferent neurons) that interact with a glutamatergic neuronal population (Fig. 3**A**). Glutamate antagonism has been shown to completely block the suppressive effect of MePD UCN3 activation on LH pulsatility[17], suggesting a functional importance of glutamatergic neurons in the circuit, hence we keep this neuronal population in the model. To incorporate UCN3 input we include a non-autonomous term in the model informed by recordings of the activity of UCN3 neurons during optogenetic stimulation (see Fig. 3**A** and Methods). The recorded traces exhibit a periodic pattern driven by the experimental stimulation protocol with 5 seconds on, 5 seconds off (see Methods). In the model, we approximated this UCN3 input using a sinusoidal function with roughly 5 seconds on, 5 seconds off pattern, driving the populations GABAergic interneurons and the glutamatergic neurons. We parametrize our MePD model to reproduce anti-correlated activity between the two GABA neuronal sub-populations. To achieve this we need to include bi-directional (i.e. mutually inhibitory) GABA interaction in the circuit as well as GABA interneurons self-inhibition (Fig. 3**A**).

Next we simulate the effects of stress on the MePD circuit in our parametrized model. UCN3 neurons in the MePD have also been shown to mediate psychosocial stress-induced suppression of the GnRH pulse generator[10]. Using our mathematical model, we begin by reproducing experimental observations of GABA activity under an acute (120 seconds) stress intervention (Fig. 3**B**). Experimentally recorded calcium activity of UCN3 neurons during stress intervention is characterized by a fast rise followed by a slow decay (Fig. 3**B**), which we model by taking the difference between two exponential functions, one with a fast and one with a slow time constant (see Methods). The model confirms emergence of anti-correlated behavior between the GABAergic sub-population whereby the activity of GABA interneurons decreases over the stress period, while

the activity of the GABA efferent neuron population increases (Fig. 3**B**), supporting the proposed

MePD circuit structure (Fig. 3**A**).

Figure 3: *In silico* **simulation of the activity in the GABAergic neuronal populations in the MePD under**

the increased excitatory input from UCN3 neurons and under restraint stress. Schematic on the left

represents the MePD neuronal circuit interacting with UCN3, consisting of two GABA populations and a

glutamatergic population. (**A**) Calcium time-traces of MePD UCN3 neurons during optogenetic stimulation of

UCN3 neurons and corresponding mean activity. Model simulations of the effects of UCN3 optogenetic

stimulation supports the emergence of anti-correlated activity in GABAergic sub-populations. (**B**) Calcium

time-traces of MePD UCN3 neurons during restraint stress and corresponding mean activity of the UCN3

calcium traces during restraint stress with corresponding mean activity. Simulations of the MePD circuit under

the effects of stress also support anti-correlated GABAergic sub-populations' activity.

193 ***In silico* modeling reveals complementary relationship between MePD GABA and UCN3**

194 **populations**

195 Having shown that MePD model can reproduce anti-correlated activity of two GABA sub-

196 populations, we examine the effects of distinct interventions involving GABA and UCN3 in the MePD

197 on the LH pulses via coupling the model of the MePD circuitry defined in the previous section to our

198 network model of the KNDy population in the arcuate nucleus (ARC)[5,18]. Building on our previous *in*

199 *silico* findings, which suggest that MePD signal integration is mediated predominately by GABAergic

200 projections[5], here we consider only the input from GABA efferent neuron to the KNDy network in

201 order to complete our coupled MePD – KNDy network model. The effects of MePD GABA and

202 glutamate antagonism alone and together with UCN3 stimulation on LH pulsatility have been

203 previously investigated[17]. We calibrate our coupled model parameters by reproducing these effects

204 accordingly (see Supplementary Fig. 6).

205 Next using the calibrated model, we predict that providing a periodic excitatory input (see

206 Supplementary Fig. 7(**A**)), that mimics the effects of optogenetic stimulation of UCN3, leads to the

207 increase in the KNDy inter-pulse interval (IPI) from 15.18 min to 23.97 min due to increased activity

208 in the inhibitory GABAergic input to the KNDy network (Fig. 4**A**).

209 Since MePD has an overwhelming number of GABAergic outputs and interneurons, we are

210 interested in their role in information processing of stress signaling. To this end we provide periodic

211 excitatory input directly to both populations of GABA neurons (see Supplementary Information). This

212 is different to the indirect activation through UCN3 stimulation, as it leads to the increased activity in

213 both GABA subpopulations (Supplementary Fig. 7(**B**)). The resulting effect on the KNDy IPI is an

214 increase from 15.18 min to 42.65 min (Fig. 4**B**).

215 In the model we hypothesize that UCN3 signaling to the KNDy is operating through the GABAergic

216 pathway, hence disrupting GABAergic interactions will suppress UCN3's effects on KNDy. To

217 investigate it we suppress the strength of the GABAergic interactions by 50% and mimic UCN3

218 stimulation by providing excitatory input to the circuit. The model shows that this intervention does

219   not have a significant effect on KNDy IPI, apart from slightly increasing it from 15.18 min to 17.31

220   min (Fig. 4**C** & Supplementary Fig. 7(**C**)).

221   Taken together the above could be summarized as the following set of model predictions along with

222   suggested ways to test them experimentally:

1. Optogenetic stimulation modelled via excitation of the MePD UCN3 increases the KNDy IPI 15.18 min to 23.97 min. This could be tested experimentally via in vivo optogenetic activation of the MePD UCN3 neuronal populations and simultaneous measurement of LH pulsatility.

2. Stimulation of the MePD GABA populations increases the KNDy IPI 15.18 min to 42.65 min. Such a substantial change underscores the critical role of GABAergic signaling in modulating KNDy pulsatility. Experimental confirmation could involve *in vivo* optogenetic activation of the entire MePD GABAergic population accompanied by LH pulsatility measurement.

3. Stimulation of the MePD UCN3 with simultaneous suppression of the GABAergic interactions in the MePD does not significantly affect KNDy IPI, changing it from 15.18 min to 17.31 min. This finding suggests that UCN3 effects on the KNDy network are primarily dependent on its interactions with GABA populations. A strategy to verify this hypothesis experimentally could be to stimulate UCN3 non-GABA-expressing neurons in vivo while simultaneously inhibiting non-UCN3 GABAergic neurons and measure LH pulsatility.

Finally, we investigate *in silico* the effects of stress on LH pulse dynamics mediated via the MePD.

In the restraint-stress experiment it was shown that one GABAergic cluster was active, while the

other one was inactive (Fig. 2**D**). By simulating these dynamics (Supplementary Fig. 7(**D**)) in the

coupled MePD - KNDy network model we are able to demonstrate that pulses in the KNDy network

ceased (Fig. 4**D**). These modeling findings align with published experimental evidence indicating

that restraint stress suppresses LH pulsatility[10].

Figure 4: Coupled MePD - KNDy model predictions for the effect of stress and stimulation of UCN3 neurons on GnRH pulse generator activity and hence LH pulsatility. (**A**) Simulation of the optogenetic stimulation of the UCN3 neurons leads to the increase of the KNDy IPI 15.18 min to 23.97 min. (**B**) Optogenetic stimulation of the MePD GABA neurons (providing excitatory input to both GABAergic populations) increases KNDy IPI from 15.18 min to 42.65 min. (**C**) Optogenetic stimulation of the UCN3 neurons and simultaneous inhibition of GABAergic neuronal populations do not significantly affect KNDy IPI, where it changes from 15.18 min to 17.32 min. (**D**) Simulating the effects of stress leads to the cessation of the pulsatile dynamics.

Selective optogenetic stimulation of UCN3 non-GABAergic neurons inhibits pulsatile LH secretion in female mice.

We now test experimentally our modeling predictions detailed in the previous section regarding the inhibitory effect of MePD UCN3 activation on LH pulsatility *in vivo* in freely behaving mice. To this end a cohort of UCN3-Cre::VGAT-Flpo transgenic female mice was injected with an AAV-nEF-Con/Foff 2.0-ChRmine-oScarlet viral construct. This intersectional approach enabled conditional expression of channelrhodopsin (ChRmine) in UCN3 neurons not expressing GABA (UCN3 non-GABAergic) (Con/Foff group). Following the viral injection, a unilateral fiber-optic cannula was implanted into the MePD, above the injection site, to facilitate selective activation of the targeted neuronal population.

The stimulation protocol consisted of a 60-min control period, followed by 60 min of optogenetic stimulation. Stimulation was delivered, as above, in a pulsed pattern (5 seconds on, 5 seconds off) with a pulse frequency of 10 Hz during the on period (10-ms pulse width, 10 mW power, 532 nm wavelength). Blood samples (5 µL) were collected every 5 min throughout the protocol and used to measure LH levels.

Statistical analysis indicated post-stimulation LH inter-pulse intervals differed across the two groups. In the Con/Foff group, the LH inter-pulse interval significantly increased after optogenetic stimulation compared to the control period (15.33 ± 0.83 min vs. 23.75 ± 1.07 min, mean ± SEM, n = 6, p<0.01; Fig. 5**A&D**), validating the first of our modeling predictions. In contrast, the no-stimulation group, which received the Con/Foff viral construct without optogenetic stimulation, showed no significant changes in LH inter-pulse interval (16.11 ± 0.56 min vs. 16.70 ± 0.96 min, n = 3, p > 0.05; Fig. 5**B&D**). Similarly, in a separate cohort of mice injected with a control virus (DIO-EYFP), optogenetic stimulation resulted in no significant differences in LH inter-pulse intervals before and after stimulation (18.33 ± 1.18 min vs. 17.08 ± 1.25 min, n = 4, p > 0.05; Fig. 5**C&D**).

Selective optogenetic stimulation of UCN3-expressing GABA neurons fails to affect pulsatile LH secretion in female mice.

To further test the functional interactions between the UCN3 and GABAergic populations, a

separate cohort of UCN3-Cre::VGAT-Flpo transgenic female mice was injected with an AAV-nEF-

Con/Fon 2.0-ChRmine-oScarlet viral construct, which allows optogenetic stimulation of UCN3

neurons co-expressing GABA (UCN3-expressing GABA) (Con/Fon group). Statistical analysis

showed no significant interaction effect of group and optogenetic stimulation on LH inter-pulse

intervals (two-way ANOVA, $F_{(1, 14)} = 0.4306$, $p = 0.5223$). That is, LH inter-pulse intervals in the

Con/Fon group, showed no change following optogenetic stimulation compared to the control period

(17.92 ± 0.80 min vs. 16.60 ± 0.88 min, mean $\pm$ SEM, n = 6, $p > 0.05$; Fig. 5**H&J**). Similarly, in the

no-stimulation control experiment for this cohort, which did not receive optogenetic stimulation, there

was no significant difference in LH inter-pulse intervals (15.42 ± 2.53 min vs. 15.83 ± 1.67 min, n =

3, $p > 0.05$; Fig. 5**I&J**).

Taken together, our results suggest that UNC3 and GABAergic populations are functionally

orthogonal in the MePD circuit, i.e., activation of UCN3 neurons relay inhibitory information to the

hypothalamic GnRH pulse generator but activation of the UCN3 neurons that co-expressing GABA

are not sufficient.

Figure 5: Effect of optogenetic stimulation of UCN3 non-GABAergic and UCN3 neurons co-expressing GABA in the MePD on LH pulse frequency. (**A**-**D**) Optogenetic stimulation of UCN3 non-GABAergic neurons inhibited LH pulses. Representative examples of LH pulsatility in UCN3-Cre::VGAT-Flpo mice injected with AAV-Con/Foff-ChRmine in response to (**A**) 10Hz optic stimulation or (**B**) no stimulation as control, and (**C**) in mice injected with control virus in response to 10 Hz stimulation. Pulses detected by the DynPeak algorithm are indicated with an asterisk (*). (**D**) Mean LH inter-pulse interval (IPI) (±SEM) for each group during the control period and over the subsequent stimulation period. #: $p<0.01$ vs stimulation period in Con/Foff group (Two-way ANOVA, Tukey's post-hoc). (**E**-**G**) Representative photomicrographs of expression of AAV-nEF-Con/Foff 2.0-ChRmine-oScarlet (red) in MePD UCN3 neurons. (**H**-**M**) Optogenetic stimulation of neurons co-expressing UCN3 and GABA in the MePD did not impact LH pulse significantly. Representative

examples of LH pulsatility in UCN3-Cre::VGAT-Flpo mice injected with Con/Fon-ChRmine-oScarlet virus in response to (**H**) 10 Hz optic stimulation or (**I**) no stimulation as control. (**J**) Mean LH IPI (±SEM) for each group during the control period and over the subsequent stimulation period (Two-way ANOVA). (**K**-**M**) Representative photomicrographs of neurons co-expression UCN3 and GABA, tagged with oScarlet (red). Scale bars represent (**E&K**) 200 μm, (**F&L**) 100 μm, (**G&M**) 50 μm. OT, optic track.

The MePD GABAergic neurons relay information from UCN3 neurons and inhibit the GnRH pulse generator.

To test our remaining model predictions and establish the functional role of GABAergic neurons in the MePD in transmitting inhibitory signals to the hypothalamic GnRH pulse generator, we injected the AAV-EF1a-ChR2 viral construct into VGAT-Cre transgenic mice, targeting the entire MePD GABA neuron population for expressing ChR2 and used the optogenetic stimulation protocol described above (60 min control period followed by 60 min stimulation) to stimulate the entire MePD GABAergic population. Statistical analysis showed a significant interaction effect of our genetic intervention and stimulation period on LH pulsatility ($F_{(2,18)}$ = 7.567, p = 0.0041; two-way ANOVA). Stimulation of the GABAergic population markedly increased the LH inter-pulse interval compared to the pre-stimulation control period (18.5 ± 2.03 min vs 49.5 ± 8.46 min, mean ± SEM, n = 5, p<0.01; Tukey's post-hoc; Fig. 6**A&D**), validating modeling prediction 2. In contrast, mice that did not receive optogenetic stimulation showed no change in LH inter-pulse interval (23.13 ± 4.25 min vs 20.00 ± 1.77 min, n = 4, p > 0.05; Tukey's post-hoc; Fig. 6**B&D**). Similarly, control virus (AAV-EF1a-DIO-EYFP) injections did not result in any significant difference in LH inter-pulse interval before and during stimulation (15.83 ± 0.83 min vs 18.33 ± 0.96 min, n = 3, p > 0.05; Tukey's post-hoc; Fig. 6**C&D**).

Next, UCN3-Cre::VGAT-Flpo transgenic females were injected with an intersectional viral mixture containing AAV-nEF-Con/Foff 2.0-ChRmine-oScarlet and pAAV-nEF-Coff/Fon-NpHR3.3-EYFP , enabling simultaneous excitation of UCN3 non-GABA neurons and inhibition of non-UCN3 GABAergic neurons. Statistical analysis showed no significant interaction effect of genetic manipulation and stimulation period on LH pulsatility ($F_{(1, 14)}$ = 0.2338, p = 0.6362; two-way ANOVA).

Moreover, optogenetic stimulation resulted in no significant change in the LH inter-pulse interval

relative to the control period (16.67 ± 1.31 min vs 18.61 ± 1.58 min, mean ± SEM, n = 6, p > 0.05;

Fig. 6**H&J**), which aligns with our modeling prediction 3. Similarly, in mice injected with the same

viral mixture but without receiving optogenetic stimulation, no significant change in LH inter-pulse

interval was detected (16.39 ± 2.47 min vs 16.67 ± 0.83 min, n = 3, p > 0.05; Fig. 6**I&J**).

Collectively, these findings suggest that MePD GABAergic neurons are integral to regulating LH

pulse frequency. The inhibitory effect on LH pulses observed upon UCN3 neuron activation is

attenuated when GABAergic neuron activity is optogenetically suppressed, implying that GABAergic

signaling acts downstream of UCN3 neurons, relaying the inhibitory signal to the hypothalamic

GnRH pulse generator.

Figure 6: Selective optogenetic stimulation of GABA neurons in the MePD inhibits LH pulsatility and their activity is necessary for mediating UCN3 signalling downstream. (**A-G**) Optogenetic activation of the entire MePD GABA population in VGAT-Cre mice led to significant LH pulse suppression. Representative examples showing LH pulsatility in VGAT-Cre mice injected with AAV-double floxed-ChR2-EYFP virus in response to (**A**) 5 Hz stimulation or (**B**) no stimulation as control, and (**C**) 5 Hz stimulation in mice injected with control virus. (**D**) Mean LH IPI (±SEM) for each group during the control period and over the subsequent

stimulation period. #: p<0.01, †: p<0.05 vs stimulation period in ChR group (Two-way ANOVA, Tukey's post-

hoc). (**E-G**) Representative photomicrographs of MePD GABA neurons, labeled with (**E**) tdTomato (red), (**F**)

ChR2-EYFP (green), and (**G**) merged. Scale bars represent. (**H-P**) The simultaneous inhibition of non-UCN3

GABA neurons in the MePD restored the LH suppression induced by stimulation of non-GABAergic UCN3

neurons. Representative examples showing LH pulsatility in UCN3-Cre::VGAT-Flpo female mice injected with

Con/Foff-ChRmine-oScarlet and Coff/Fon-NpHR-EYFP virus mixture in response to (**H**) 10 Hz optogenetic

stimulation or (**I**) no stimulation as control. (**J**) Mean LH IPI (±SEM) for each group during the control period

and over the subsequent stimulation period. (Two-way ANOVA). (**K-P**) Representative photomicrographs of

fluorescence in MePD neurons. (**K&N**) non-GABAergic UCN3 neurons tagged with oScarlet (red), (**L&O**) non-

UCN GABA neurons tagged with EYFP (green) (**M&P**) merged. Scale bars represent (**E-G**) 150 µm, (**K-M**)

200 µm, (**N-P**) 100 µm.

Discussion

The posterodorsal subnucleus of the Medial Amygdala (MePD), is a critical regulator of reproductive

function. Single-cell transcriptomics has revealed a mosaic of glutamatergic and GABAergic

populations in this region, with GABA neurons co-expressing receptors for gonadal steroids as well

as neuromodulators and cognate receptors involved in parental and sociosexual behavior (including

kisppeptin, prolactin and oxytocin)[19]. Furthermore, calcium imaging of neuronal ensembles from this

region have started to decipher how this brain region encodes information about social interactions

and stressful stimuli[20], with stress related information being relayed to the reproductive axis via

potentially direct projections to the hypothalamus[1,2], and eliciting well-documented responses on

pubertal timing[6,7], LH secretion[10,11] and the GnRH pulse generator[3–5]. Here, to elucidate the

neuronal circuitry involved in MePD processing of stress signals, we used GRIN lens mini-

endoscopic calcium imaging to study, for the first time, the dynamic activity of MePD GABAergic and

UCN3 neurons under acute restraint stress as well as the activity of GABAergic neurons in response

to UCN3 activation. Functional network analysis of the imaging data revealed that acute restraint

stress induces a functional reconfiguration of the GABAergic population, emphasizing the pivotal

role of GABA neurons in encoding and transmitting stress information. A comparable reconfiguration

was observed following optogenetic stimulation of UCN3 neurons, suggesting that these two

populations are co-involved in stress processing. Further clustering analysis of the calcium

transients identified two distinct GABAergic sub-populations with anti-correlated activity. This finding

of two distinct populations is consistent with physiological and morphological studies of GABAergic

neurons in the medial amygdala characterizing them as either projecting cells or interneurons[12,14].

Building upon our previously developed computational model of the MePD circuit, we explored the

neuronal pathways underlying stress processing. The model consists of a glutamatergic sub-

population and two GABAergic sub-populations: one comprising GABAergic interneurons and the

other GABAergic efferent neurons. Our analysis identified two conditions necessary for the anti-

correlated activity between these GABAergic sub-populations, consistent with our experimental

observations. First, mutual inhibition between the GABA sub-populations is required, where

activation of one sub-population suppresses the activity of the other. Second, if both sub-

populations receive a shared input (e.g., from the glutamatergic population or in fact any other

population of excitatory cells), their dynamics become entrained to that common stimulus, meaning

that distinct input streams to each sub-population are necessary for the observed anti-correlation.

To satisfy this condition in our model, GABAergic interneurons receive inputs from UCN3 neurons,

whereas the GABAergic efferent sub-population receives inputs from glutamatergic neurons. This

circuitry implies that stress information is processed along two key pathways (one glutamatergic and

one GABAergic) and is integrated by the GABAergic efferent neurons.

We note that an alternative connectivity pattern, with the UCN3 population directly projecting to the

GABA efferent population, could also fit our experimental observations, but does not reproduce

previously published results on the effects of neuropharmacological perturbations in the MePD on

LH pulsatility[17]. Specifically, the suppression of glutamatergic neurons in combination with

stimulation of UCN3 neurons. Under this scenario direct input from stimulated UCN3 to GABA

efferent neurons will lead to the increase in its activity, consequently increasing KNDy IPI. This

finding contrasts with experimental observations, where the combined suppression of glutamate and

stimulation of UCN3 failed to produce any significant change in GnRH pulsatility[17].

Consistent with model predictions, we showed that optogenetic activation of UCN3-expressing non-

GABAergic neurons suppresses LH pulse frequency. This finding corroborates previous work

demonstrating that LH pulses are inhibited by activation of UCN3 neurons[7,17] or administration of

UCN3[10] in the MePD, and that such effects are attenuated by chemogenetic inhibition of UCN3

neurons or by antagonizing corticotropin-releasing factor type 2 receptors (CRFR2)[10]. Thus, our

findings further support the central role of UCN3 signaling in modulating both stress responses and

reproductive function.

GABA is one of the primary inhibitory neurotransmitters in the nervous system[12]. Previous work has

shown that GABAergic neurons in the MePD project to the ARC to regulate LH pulses and

chemogenetic inhibition of MePD GABA neurons attenuates stress-induced suppression of LH

pulses[11]. In this study, we found that optogenetic activation of GABAergic neurons in the MePD

leads to robust LH pulse suppression, reinforcing the pivotal role of GABAergic signaling in stress-

related reproductive dysfunction. Antagonizing GABA receptors in the MePD can block UCN3-

induced LH suppression[17], suggesting that GABAergic neurons function downstream of UCN3

signaling. However, this pharmacological approach lacks cell-type specificity, leaving open the

possibility that UCN3 neurons themselves co-expressing GABA receptors, potentially contributing to

the observed effect. Here, by using an intersectional genetic strategy, we demonstrated that

optogenetically inhibiting non-UCN3 GABAergic neurons can abolish the change in LH pulse

intervals induced by optogenetic stimulation of UCN3 neurons. This finding provides evidence that

these GABAergic neurons act downstream of UCN3 signaling to mediating LH pulse suppression.

We note, that while in our current experimental protocol we could also be blocking GABAergic

projections to the ARC KNDy population, which by itself could explain the absence of any effect

from UCN3 stimulation, our modeling predicts that GABAergic interactions within the MePD are

additionally crucial for relaying UCN3 signals to the KNDy network as also suggested by

pharmacologically antagonizing GABA receptors[17].

It is not unusual for neurons to co-express neuropeptides and neurotransmitters[14]. Given that

activating either UCN3 or GABAergic neurons independently suppresses LH pulses, one might

predict that stimulating neurons co-expressing both UCN3 and GABA would similarly inhibit LH.

However, we observed no significant changes in LH pulses upon optogenetic stimulation of neurons

co-expressing UCN3 and GABA. This suggests the two populations are functionally orthogonal:

although both UCN3 and GABA neurons are essential for LH pulse inhibition, their overlap is not.

Within our modeling framework, UCN3/GABA co-expressing neurons might be considered part of

the GABAergic interneuron population, primarily inhibiting GABA projection neurons rather than

activating them. Consequently, they fail to relay the inhibitory UCN3 signal to the ARC. Further

studies are necessary to define more precisely the distinct role of UCN3/GABA neurons within the

MePD circuitry.

By combining mini-endoscopic calcium imaging with computational modeling, we identified two

GABAergic sub-populations with complementary but distinct roles in encoding stress signals and

modulating LH pulsatility. Our experimental and computational results also stressed the functional

orthogonality between UCN3 and GABA signaling in MePD, highlighting the complexity of the

MePD's contribution to stress-induced reproductive suppression. These results offer valuable

insights into the neuronal circuits potentially involved in processing stress-related effects in

reproduction, paving the way for future research aimed at improving reproductive health and well-

being.

Methods

Animal

VGAT-Cre-tdTomato mice were genotyped using PCR to determine heterozygosity for VGAT-Cre

(primers 5′-3′: common, 12785—CTTCGTCATCGGCGGCATCTG; wild-type reverse12786—

CAGGGCGATGTGGAATAGAAA; mutant reverse oIMR8292—CCAAAAGACGGCAATATGGT) and

tdtomato (primers 5′-3′: wild-type forward oIMR9020—AAGGGAGCTGCAGTGGAGTA; wild-type

reverse oIMR9021—CCGAAAATCTGTGGGAAGTC; mutant reverse WPRE oIMR9103—

GGCATTAAAGCAGCGTATCC; mutant forward tdTomato oIMR9105—

CTGTTCCTGTACGGCATGG). UCN3-Cre heterozygous mice (strain Tg(Ucn3-cre)KF43Gsat/

Mmucd, MMRRC GENSAT) were crossbred with VGAT-Flpo heterozygous mice (Jax stock

#007909,B6.Cg-Slc32a1 2A-Flpo-D:Het) to obtain female double heterozygous UCN3-Cre::VGAT-

Flpo mice (Slc32a1 2A-Flpo-D:Het|Ucn3-Cre:Het|Rosa26<tomato>WT); genotyped using PCR for

the detection of heterozygosity (primers 5′- 3′: UCN3 Forward: CGAAGTCCCTCTCACACCTGGTT;

Cre Reverse: CGGCAAACGGAC-AGAAGCATT; Slc32a1 Mutant Forward: TGC ATC GCA TTG

TCT GAG TAG; Slc32a1 Mutant Reverse: GAC AGC CGT GAA CAG AAG G). Female mice, aged 8

to 10 weeks, weighing between 20 g and 25 g, were individually housed in ventilated cages sealed

with a HEPA-filter at 25 ± 1°C in a 12:12-hour light/dark cycle, lights on at 07:00 h. Vaginal smears

were performed when mice are 6 weeks old between 10:00 - 12:00 h daily. Only mice showing

regular estrous cycles were included. All animal procedures performed were approved by the King's

College London Animal Welfare and Ethical Review Body. Procedures were in accordance with UK

home office regulation.

Stereotaxic injection of adeno-associated virus construct and fiber-optic or mini-endoscope

GRIN lens implantation

All surgical procedures were carried out under general anaesthesia using ketamine (Vetalar, 100

mg/kg, i.p.; Pfizer, Sandwich, UK) and xylazine (Rompun, 10 mg/kg, i.p.; Bayer, Leverkusen,

Germany) under aseptic conditions. Animals were secured in a David Kopf stereotaxic frame (Model

900, Kopf Instruments, CA, USA). Following bilateral ovariectomy, a midline incision was made in

the scalp to reveal the skull. Two small bone screws were inserted into the skull. A small hole was

drilled above the position of the MePD. The MePD coordinate (2.29 mm lateral, 1.60 mm posterior

from bregma, at a depth of 5.33 mm below the skull surface) was determined according to the

mouse brain atlas of Paxinos and Franklin[21]. A unilateral stereotaxic viral injection was performed to

target opsin or fluorescent protein expression in the MePD neurons, using a robot stereotaxic

system (Neurostar, Tubingen, Germany), with a 5-µL Hamilton microsyringe (Esslab, Essex, UK), at

the speed of 25 nl/min for 10 min.

Two transgenic animal models were used for LH blood sampling under optogenetic stimulation:

UCN3-Cre::VGAT-Flpo and VGAT-Cre-tdTomato. The following types of viral constructs were used

in separated cohorts of UCN3-Cre::VGAT-Flpo mice: (i) AAV-nEF-Con/Foff 2.0-ChRmine-oScarlet

(250 nl, $2.1×10^{12}$ gc/mL, Serotype:8, #137161, Addgene, Massachusetts, USA); (ii) AAV-nEF-

Con/Fon 2.0-ChRmine-oScarlet (250 nl, $1.9×10^{12}$ gc/mL, Serotype:8, # 137159, Addgene,

Massachusetts, USA); (iii) a mixture of AAV-nEF-Con/Foff 2.0-ChRmine-oScarlet ($2.1×10^{12}$ gc/mL,

#137161) and pAAV-nEF-Coff/Fon-NpHR3.3-EYFP ($1.9×10^{12}$ gc/mL; #137159, final volume 300 nl);

(iv) control virus AAV-Ef1a-DIO-EYFP (250 nl, $4.4 × 10^{12}$ gc/mL, Serotype:9, #27056, Addgene,

Massachusetts, USA). VGAT-Cre-tdTomato mice received unilateral injections of (i)

channelrhodopsin (ChR2) viral construct, (250 nl, $1.8 × 10^{12}$ gc/mL; Serotype:9; Addgene, #20298,

Massachusetts, USA) or (ii) control virus, AAV-Ef1a-DIO-EYFP (250 nl, $4.4 × 10^{12}$ gc/mL,

Serotype:9; Addgene, # 27056, Massachusetts, USA) into the MePD. Following injection, the needle

was kept in position for 10 min before going up slowly over 1 min. A unilateral fibre optic cannula

(200 μm, 0.39 NA, 1.25 mm ceramic ferrule; Doric Lenses, Quebec, Canada) was inserted into the

right MePD 0.15 mm above the injection site. After reaching the target, the fibre optic cannula was

secured on the skull using dental cement (Super-Bond Universal Kit, Prestige Dental, UK), bonded

to the screws and skull. Finally, the incision in the scalp was closed with suture. Animals were given

1 week to recover, after which they were handled 20 min daily to acclimatize to experimental

procedures for a further 2 weeks.

For mini-endoscope *in vivo* calcium imaging, only UCN3-Cre::VGAT-Flpo transgenic animals were

used. Separate mice cohorts received the following viral injection into the MePD: (i) a mixture of

pAAV-Syn-Flex-rc[Chrimson-tdTomato] ($2.0×10^{12}$ gc/mL, Serotype:5, #62723, Addgene,

Massachusetts, USA) + pAAV-EF1a-fDIO-GCaMP6s ($2.3×10^{12}$ gc/mL; Serotype:8; #105715,

Addgene, Massachusetts, USA; final volume 400 nl); (ii) a mixture of pAAV-Syn-Flex-rc[Chrimson-

tdTomato] ($2.0×10^{12}$ gc/mL, Serotype:5, #62723, Addgene, Massachusetts, USA) +

pAAV.CAG.Flex.GCaMP6s.WPRE.SV40 ($4.3×10^{12}$ gc/mL; Serotype:9; # 100842, Addgene,

Massachusetts, USA; final volume 400 nl); (iii) a mixture of control virus AAV-Ef1a-DIO-EYFP (250

nl, 4.4×10^{12} gc/mL, Serotype:9, #27056, Addgene, Massachusetts, USA) + pAAV-EF1a-fDIO-

GCaMP6s (2.3×10^{12} gc/mL; Serotype:8; #105715, Addgene, Massachusetts, USA; final volume 250

nl). One week after administering the virus, a 600-µm diameter, 7.2-mm long gradient index (GRIN)

ProView™ Integrated Lens (Inscopix, PN:100-004554) was implanted at the same location as the

viral injection. The surface of the skull was scored, using the tip of a surgical scalpel, and four

screws were inserted into the skull. This was done to enhance the adhesion of dental cement to the

skull, thereby reducing the interference of movement on *in vivo* recording. Before insertion of the

GRIN lens, a 23 G needle was slowly lowered at a speed of 5.0 mm/min into the brain (2.42 mm

lateral, 1.52 mm posterior from bregma, depth of 5.05 mm below the skull surface) to make a needle

track to facilitate the insertion of the GRIN lens without causing brain distortion along its route. The

GRIN lens was then inserted into the brain at a rate of 0.6 mm/min, using the same coordinates as

the needle precutting. To secure the GRIN lens with integrated baseplate in position, we used a light

curing adhesive (iBond Universal, Kulzer GmbH, Leipziger, Germany) along with a cured

radiopaque hybrid dental cement (Gradia Direct Flo, GC corporation, Tokyo, Japan), bonding it to

the head screws and skull. A GRIN lens baseplate dust cover (Inscopix) was rescued in place to

protect the lens. After 4 weeks the mice were re-anaesthetized as described above and underwent

bilateral ovariectomy.

Serial blood sampling for LH measurement

The mouse's tail-tip was excised using a sterile scalpel. Then the chronically unilaterally implanted

fibre optic cannula was connected to a multimode fibre optic rotary joint patch cables (Thorlabs Ltd,

Ely, UK) via a ceramic mating sleeve to allow the animal freedom of movement. Mice were left for 1

h to habituate. A Grass SD9B stimulator-controlled DPSS laser (Laserglow Technologies, Toronto,

Canada) was utilised to deliver either green light (532 nm wavelength, 10 Hz, 5 seconds on 5

seconds off, 10 mW, 10-ms pulse width,) or blue light (473 nm wavelength, 5 Hz, 5 seconds on 5

seconds off, 5 mW, 10-ms pulse width,). The light pattern was controlled using software designed in

house. After 1 h of acclimatization, serial blood samples (5-µL) for LH pulse measurement were

collected every 5 min for 2 h between 10:00 - 12:00 h: 0-60 min for control blood sampling period,

and 60-120 min under stimulation or equivalent no-stimulation control period. Blood samples were

diluted in 45 μL of 0.2% BSA-0.05% PBS with Tween-20 (0.2% BSA and 0.05% PBST) and

immediately placed on dry ice and stored at −80°C until later analysis.

LH assay

LH measurement was performed using an LH ELISA assay as reported previously[22]. A mouse LH

standard (mLH; AFP-5306A, NIDDK-NHPP, USA) was used to establish a standard curve for the

quantitative measurement. The following antibodies were used in the assay: (i) coating antibody

(RRID: AB_2665514, monoclonal antibovine LH beta subunit antiserum, 518B7, University of

California, CA, USA); (ii) anti-LH antibody (RRID: AB_2665533; National Hormone & Peptide

Program, CA, USA); (iii) secondary antibody (RRID: AB_772206, GE Healthcare, Chicago, Illinois,

USA). The inter-assay and intra-assay variations were 10.53% and 5.74%, respectively. The assay

sensitivity was 0.023 ng/mL.

LH pulse statistical analysis

546   LH pulses were verified using the Dynpeak algorithm in Scilab 5.5.2 programme[23]. The Dynpeak

547   algorithms were classified for OVX mice based on previously established parameters[24]. The average

548   LH inter-pulse interval was calculated for the 1-h control period and the 1-h stimulation period in both

549   experimental and control mice. Statistical significance was assessed using a 2-way ANOVA unless

550   otherwise stated. Data was represented as mean ± SEM and $p < 0.05$ was considered significant.

Mini-endoscope-based Calcium imaging

Under optogenetic stimulation of MePD UCN3 neurons. Prior to recording, animals were

handled daily for 5 to 6 weeks to get accustomed to the experimental setup. The Data Acquisition

(DAQ) box (Inscopix, PN:100-005904) was turned on and connected to the computer wirelessly.

The Data Acquisition Software (IDAS; Inscopix, Inc.) was accessed through the Chrome web

browser on the computer communicating with the DAQ box. Adequate DAQ box storage capacity

was checked routinely before Calcium imaging. The GRIN lens baseplate dust cover was gently

removed from the mouse's head, after which the minscope camera was secured to the baseplate

platform. Equipped with the head-mount miniscope (nVoke2, Inscopix, SN: BB-11238601), the

mouse was then placed back into its home cage. The optimal focal plane was chosen by adjusting

the lens focus to achieve clear cell morphology. The exposure time was fixed at 50 ms. The frame

rate was set to 20 Hz. Subsequently, gain, and LED power were optimized based on image

histograms before initiating recording. Two types of LED were used during the recording: (i) EX-LED

(0.2-0.4 mW, 435-460 nm excitation filter) for GCaMP6s signal detection in the GABAergic or UCN3

neurons; (ii) OG-LED (10 mW, 590-650 nm excitation filter) for Chrimson optogenetic stimulation of

the UCN3 neurons. After the mice were attached to the miniscope, they were given 1-2 h for

adaptation. For stimulation of the UCN3 neurons, the OG-LED was set to 10 Hz, 5 seconds on 5

seconds off, and a 10-ms pulse width. Once the animals had acclimatized, the GCaMP6s recording

of the GABAergic neurons or UCN3 neurons was started with a protocol of 2 min baseline before

stimulation for 2 min. Optogenetic stimulation was randomized for the 10 Hz, with a 5-min interval

between each to prevent photobleaching of GCaMP6s. Once the experiment was concluded, the

miniscope was gently detached, and the baseplate cover was reattached and secured with its

screw. The recordings were performed between 9:00 h and 15:00 h. Three groups of animals were

included in this experiment: (i) mice with UCN3 neurons expressing opsin and GABAergic neurons

expressing GCaMP; (ii) mice with UCN3 neurons expressing both opsin and GCaMP, serving as a

positive control; (iii) mice with UCN3 neurons expressing mCherry and GABAergic neurons

expressing GCaMP, serving as a negative control.

Under restraint stress. The system was set up and the mouse was connected to the miniscope as

described above but with the OG-LED turned off. Once the mouse had settled, the experiment

started with a 3 min baseline recording, followed by 3 min of recording under hand-held restraint.

Finally, the miniscope was detached and the baseplate cover was reattached. The experiments were

performed between 9:00 h and 15:00 h. The animals in this experiment were the same as those

used in the optogenetic stimulation protocol described above, but were divided into two groups

depending on whether GCaMP was expressed in the UCN3 or GABAergic neurons, regardless of

whether opsin or mCherry was expressed in the UCN3 cells.

In vivo **calcium imaging recording processing**

The video data, with a resolution of 1280 pixels x 800 pixels, was exported from the DAQ box

through the file manager to a desktop computer and processed with the Inscopix Data Processing

Software (IDPS). Within the IDPS, there are six essential steps for extracting single-cell calcium

transient data from the recorded video: down-sampling, cropping, spatial filtering, motion correction,

trace normalization (ΔF/F), and cell identification. First, the video underwent both spatial down-

sampling and temporal down-sampling by a factor of 2 and was cropped to enhance processing

speed, focusing on the region of interest where the GCaMP signal was observed. Spatial filtering,

with a threshold ranging between 0.005 (low) and 0.5 (high) was used to distinguish individual pixels

in the video image, creating a high-contrast image. A mean intensity projection image was

generated. Motion correction was subsequently conducted, using the mean intensity projection

image as a reference. Motion correction was applied again using one of the frames affected by

movement while simultaneously drawing a region of interest (ROI) where UCN3 accumulates to

enhance the effectiveness of motion correction. For the recordings with GCaMP signals from

GABAergic neurons, we did not use the built-in motion correction function of IDPS. Instead, we used

the MATLAB (version R2022a) implementation of the open-source algorithm NoRMCorre[25] with

default parameters for both rigid and non-rigid (patch size 64 pixels and overlap 32) motion

correction as long as a minimum correlation of the frames with the mean frame above 0.8 is

achieved. Occasionally, if the minimum correlation is not met, an extra step of non-rigid motion

correction with coarser patches is applied. This is performed by doubling the patch size and their

overlap to 128 and 64, respectively. Corrected videos were saved in TIFF format for further analysis

via IDPS. Following motion correction, pixel intensity, $F(x,y)$, in each frame of the movie was

normalised using the following formula $\dfrac{(F(x,y) - F_{baseline}(x,y))}{F_{baseline}(x,y)}$, where $F_{baseline}(x,y)$ is the baseline

pixel intensity calculated by taking the mean value of the pixel across the entire movie. Finally,

regions of interest were identified automatically using the cell identification algorithm provided by

IDPS, based on principal component and independent component analysis (PCA-ICA algorithm).

Validation of virus injection, fibre optic cannula and GRIN lens implantation

Once the experimental procedures were completed, a lethal dose of ketamine was used to

humanely euthanize the mice. The mice were perfused transcardially with heparinised saline (5

U/ml) for 5 min, followed by ice-cold 4% paraformaldehyde (PFA) for 15 min using a pump

(Minipuls,156 Gilson, Villiers Le Bel, France). The brains were collected and post-fixed sequentially

at 4°C in a solution of 15% sucrose in 4% PFA and later in a solution of 30% sucrose in phosphate

buffer until they sank. They were then snap frozen on dry ice and stored in -80°C until processing.

The brains were coronally sectioned (30 μm) between +0.5 mm and -2.7 mm from the bregma using

a cryostat (Bright Instrument Co., Huntingdon, UK), and sections were mounted on microscope

slides, air-dried, and covered with slips using Prolonged Antifade mounting medium (Molecular

Probes, Inc. OR, USA). The accuracy of injection sites, fibre optic cannula and GRIN lens

implantation were confirmed by Axioskop 2 Plus microscope equipped with AXIOVISION 4.7

(Zeiss). Images were captured using Axioskop 2 Plus microscope (Carl Zeiss). Only the data of

mice with correct viral infection and implantation of cannula and lens in the MePD were included in

the analysis.

Data analysis

The data processing protocol we employ in this study was performed in two phases: pre-intervention

and intervention (either via optogenetic stimulation or restraint stress). For our analysis, we use the

smoothed deconvolved traces generated by the OASIS method[26], which improves the signal-to-

noise ratio (SNR) and mitigates noise interference. All traces are rescaled between 0 and 1 (min-

max normalization), and then further normalized to mean zero and unit variance. The analysis

process is divided into three key steps: i) identification of GABA cells co-expressing UCN3 and

removing them from further analysis; ii) estimating and comparing functional connectivity before and

during intervention; iii) clustering analysis based on pairwise association-derived dissimilarity scores.

In step (i), we use power spectral density (PSD) analysis and K-means clustering (using Euclidean

distance) to identify GABA neurons with activity that mimics the pattern of optic stimulation, evidence

that they are co-expressing UCN3.

PSD filtering. For each trace, we compute the PSD during the stimulation phase. Cells with PSD

values at 0.1 Hz (the frequency of the stimulus 5 seconds on and 5 seconds off cycle) above 0.37

Hz^{-1} are labelled as co-expressing UCN3. This threshold is determined from recordings of UCN3

cells under optic stimulation, by calculating the median of the minimum PSD values at 0.1 Hz from

each dataset (see Supplementary Fig. 8 **B**).

K-means clustering. Using the Euclidean distance metric, we cluster the traces during the

stimulation phase into two groups, aiming to separate neurons into GABA-only neurons and neurons

co-expressing UCN3 and GABA. Clustering separated a small group of neurons with activity

mimicking the optic stimulation oscillatory pattern (see Supplementary Fig. 8 **C**). Recordings of

MePD UCN3 neurons shows that under optogenetic stimulation their activity rises and declines in

phase with on and off periods of stimulation (shown in Fig. 3**A**); therefore, this small number of cells

was labelled as co-expressing UCN3.

The union of neurons identified by either method as co-expressing UCN3 and GABA are excluded

from further analysis. By applying both filters, we aim to robustly identify purely GABA-expressing

neurons. We acknowledge the possibility that some cells may pass through the filters or that a few

purely GABA-expressing neurons may be inadvertently excluded. However, our dual approach of

employing power spectral density (PSD) analysis and K-means clustering minimizes this possibility

and is essential to ensure our analysis remains focused exclusively on GABA-expressing neurons.

Comparing pre-intervention and intervention cell-cell interaction strength

Following step (i), the remaining dataset contains only traces associated with GABA neurons. In step

(ii), we assess the functional connectivity between the observed neurons using two measures of

pairwise association between calcium activity traces, namely covariance and a modified version of

cross-correlation. The covariance between two traces $x = \{x_i\}$ and $y = \{y_i\}$, $i = 1,..,N$ is defined as

$$cov(x, y) = \frac{1}{N-1}\sum_i (x_i - \underline{x})(y_i - \underline{y}),$$

663   where $\underline{x} = \frac{1}{N}\sum_i x_i$ and $\underline{y} = \frac{1}{N}\sum_i y_i$, quantifying the strength of their linear association[27].

664   The modified cross-correlation, also accounting for lag and sign in cell-cell interactions, is termed

665   *signed lagged cross-correlation* (SLxCorr) and is adapted from the works of Tsuyuzaki et al.[28] and

666   Paparrizos & Gravano[29]. It is defined as

$$SLXCorr(x,y) = \frac{R_{\tau_{max}-N}(x,y)}{\sqrt{R_0(x,x)R_0(y,y)}},$$

668   in which

$$\tau_{max} = \underset{\tau}{\mathrm{argmax}}\left(\left\|\frac{R_{\tau-N}(x,y)}{\sqrt{R_0(x,x)R_0(y,y)}}\right\|\right),$$

where $\quad R_\tau = \begin{cases} \sum\limits_{i=1}^{N-\tau-1} x_{i+\tau}\cdot y_i & \text{for} \quad \tau \geq 0 \\ R_{-\tau}(y,x) & \text{for} \quad \tau < 0 \end{cases}$ represents   cross-correlation   with   lag

$\tau \in \{-N+1, ..., -1, 0, 1, ..., N-1\}$ with N as the length of x and y.

Using the above measures, we construct the corresponding covariance and SLxCorr matrices (see

for an example Fig. 1**H&I**).

To evaluate changes in pairwise cell interactions between the pre-intervention and intervention

phases, we compare the distributions of the covariance - and SLxCorr – derived connectivity weights

using the Kolmogorov-Smirnov nonparametric statistical test[30,31]. This test is applied to each

experimental trial where only the values of weights form the upper triangular entries (excluding the

diagonal) of the covariance and SLxCorr matrices are used to avoid duplication caused by

symmetry.

To account for variability across experiments, we combine the p-values obtained from individual

trials into a single p-value for each animal as shown in Fig. 1**J**. This is done using Fisher's Combined

Probability Test[32], providing an overall measure of the difference between the pre-intervention and

intervention experimental stages for each animal.

Clustering analysis

In step (iii), we apply agglomerative hierarchical clustering (using the average linkage criterion;

scikit-learn package in Python™) to classify cells based on the dissimilarity of their calcium activity,

given as

688
$$D(x, y) = 1 - \mathrm{SLXCorr}(x,y).$$

To determine the optimal number of clusters and robustness of clustering, we employed a

bootstrapping technique. This involved resampling (without replacement) 75% of the cells and

clustering the resulting dataset with different numbers of clusters (2, 3, and 4) using hierarchical or

K-means clustering with either the covariance or SLxCorr as the dissimilarity measure. This

resampling process was repeated 100 times.

For each resample dataset, we evaluate the clustering performance, using the Silhouette Coefficient

(SC)[33] which measures the cohesion within clusters and their separation, with values ranging from -1

(poor performance) to 1 (good performance). Results of the evaluation are given in Supplementary

Figs. 3-5, showing that the Silhouette Coefficient is significantly higher for 2 clusters compared to 3

or 4 clusters in the case of both hierarchical and K-means clustering.

Modeling

We build on our previously published modelling framework of MePD circuit[5] by incorporating

dynamic stimulation by UCN3 as observed experimentally (as shown in Fig. 3). To this end we

define UCN3 input into the populations of glutamatergic neurons and GABA interneurons with a

strength given by the parameter γ. The activity of UCN3 during stimulation is approximated via

sinusoidal function:

$$U_{stim}(t) = \gamma B + A \sin(2\pi f t) ,$$

706 where B and A are the baseline UCN3 activity and the amplitude, respectively, while the parameter

707 f is the frequency of the sigmoid wave. We approximate UCN3 activity during restraint stress using

708 two exponential functions:

$$U_{stress}(t) = \gamma(at \exp(-r_2 t) + b(1 - \exp(-r_1 t))) ,$$

where the function initially increases with magnitude a then decreases to a plateau value given by

the parameter b. The parameters r_1 and r_2 are responsible for the rate of increase of the first term

and rate of decrease to a plateau in the second term.

Briefly, the model captures the dynamics of the MePD intracellular calcium activity of three

populations: glutamatergic neuronal population (excitatory) and the populations of GABA

interneurons and GABA efferent neurons (inhibitory). We build on our previous work[5] by assuming

that excitatory (UCN3) input is provided to the populations of GABA interneurons and glutamate (Fig

7). Previous analysis of the model indicated that negative feedback between GABAergic and

glutamatergic neuronal populations as well as glutamatergic self-excitation are critical mechanisms

for inducing oscillatory dynamics[5]. However, providing glutamatergic inputs to both populations of

GABA neurons result in a null phase difference between the populations. Therefore, we opted to

provide glutamatergic input only to the population of GABA efferent neurons.

To investigate how the amygdala modulates the activity in the GnRH pulse generator, we couple the

above-described MePD model to our KNDy network model, where the oscillatory dynamics of the

KNDy acts as a proxy of LH pulsatility[18]. Here we couple the model via providing only inhibitory

output from MePD GABA efferent neurons population to KNDy, as it was previously shown that

glutamate plays a modulatory role to the MePD output rather than directly affecting the KNDy

network[5]. We then calibrate the coupled by reproducing the experimental results from Ivanova et

al.[17] (Supplementary Fig. 6). Further modeling details, such as equations governing model dynamics

and model parameter values, can be found in Supplementary Information. The simulations have

been done in MATLAB (version R2023b).

Data Availability

732 Source data are provided with this paper. Further information that supports the findings of this study

733 is available from the corresponding authors upon request.

Code Availability

735 The code for reproducing the data analysis (Python™), mathematical modeling (MATLAB), and

736 figure generation for both the main text and Supplementary information, along with a test dataset, is

737 publicly available at: https://github.com/mv-kr/MePD_GABA.

References

1. Yeo, S.-H., Kyle, V., Blouet, C., Jones, S. & Colledge, W. H. Mapping neuronal inputs to Kiss1

neurons in the arcuate nucleus of the mouse. *PLOS ONE* **14**, e0213927 (2019).

2. Moore, A. M., Coolen, L. M. & Lehman, M. N. Kisspeptin/Neurokinin B/Dynorphin (KNDy) cells

as integrators of diverse internal and external cues: evidence from viral-based monosynaptic

tract-tracing in mice. *Sci. Rep.* **9**, 14768 (2019).

3. Comninos, A. N. *et al.* Kisspeptin signaling in the amygdala modulates reproductive hormone

secretion. *Brain Struct. Funct.* **221**, 2035–2047 (2016).

4. Lass, G. *et al.* Optogenetic stimulation of kisspeptin neurones within the posterodorsal medial

amygdala increases luteinising hormone pulse frequency in female mice. *J. Neuroendocrinol.*

32, e12823 (2020).

5. Nechyporenko, K. *et al.* Neuronal network dynamics in the posterodorsal amygdala: shaping

reproductive hormone pulsatility. *J. R. Soc. Interface* **21**, 20240143 (2024).

6. Adekunbi, D. A. *et al.* Role of amygdala kisspeptin in pubertal timing in female rats. *PLOS ONE*

12, e0183596 (2017).

7. Ivanova, D. *et al.* Role of Posterodorsal Medial Amygdala Urocortin-3 in Pubertal Timing in

Female Mice. *Front. Endocrinol.* **13**, (2022).

8. Adekunbi, D. A. *et al.* Kisspeptin neurones in the posterodorsal medial amygdala modulate

sexual partner preference and anxiety in male mice. *J. Neuroendocrinol.* **30**, e12572 (2018).

9. Wu, Y. E. *et al.* Neural control of affiliative touch in prosocial interaction. *Nature* **599**, 262–267

(2021).

10. Ivanova, D. *et al.* Urocortin3 in the Posterodorsal Medial Amygdala Mediates Stress-induced

Suppression of LH Pulsatility in Female Mice. *Endocrinology* **162**, 1–14 (2021).

11. McIntyre, C. *et al.* GABA Signaling in the Posterodorsal Medial Amygdala Mediates Stress-

induced Suppression of LH Pulsatility in Female Mice. *Endocrinology* **164**, 1–11 (2023).

12. Keshavarzi, S., Sullivan, R. K. P., Ianno, D. J. & Sah, P. Functional Properties and Projections

of Neurons in the Medial Amygdala. *J. Neurosci.* **34**, 8699–8715 (2014).

13. Choi, G. B. *et al.* Lhx6 delineates a pathway mediating innate reproductive behaviors from the

amygdala to the hypothalamus. *Neuron* **46**, 647–660 (2005).

14. Bian, X. Physiological and morphological characterization of GABAergic neurons in the medial

amygdala. *Brain Res.* **1509**, 8–19 (2013).

15. Lewis, K. *et al.* Identification of urocortin III, an additional member of the corticotropin-releasing

factor (CRF) family with high affinity for the CRF2 receptor. *Proc. Natl. Acad. Sci. U. S. A.* **98**,

7570–7575 (2001).

16. Shemesh, Y. *et al.* Ucn3 and CRF-R2 in the medial amygdala regulate complex social

dynamics. *Nat. Neurosci.* **19**, 1489–1496 (2016).

17. Ivanova, D., Li, X.-F., McIntyre, C. & O'Byrne, K. T. Posterodorsal Medial Amygdala Urocortin-3,

GABA, and Glutamate Mediate Suppression of LH Pulsatility in Female Mice. *Endocrinology*

164, 1–13 (2023).

18. Voliotis, M. *et al.* The Origin of GnRH Pulse Generation: An Integrative Mathematical-

Experimental Approach. *J. Neurosci.* **39**, 9738–9747 (2019).

19. Hochgerner, H. *et al.* Neuronal types in the mouse amygdala and their transcriptional response

780   to fear conditioning. *Nat. Neurosci.* **26**, 2237–2249 (2023).

781   20. Li, Y. *et al.* Neuronal Representation of Social Information in the Medial Amygdala of Awake

782   Behaving Mice. *Cell* **171**, 1176-1190.e17 (2017).

783   21. Paxinos, G. & Franklin, K. B. J. *Paxinos and Franklin's the Mouse Brain in Stereotaxic*

784   *Coordinates*. (2019).

785   22. Steyn, F. J. *et al.* Development of a Methodology for and Assessment of Pulsatile Luteinizing

786   Hormone Secretion in Juvenile and Adult Male Mice. *Endocrinology* **154**, 4939–4945 (2013).

787   23. Vidal, A., Zhang, Q., Médigue, C., Fabre, S. & Clément, F. DynPeak: An Algorithm for Pulse

788   Detection and Frequency Analysis in Hormonal Time Series. *PLOS ONE* **7**, e39001 (2012).

789   24. Yang, J. A. *et al.* Acute Psychosocial Stress Inhibits LH Pulsatility and Kiss1 Neuronal Activation

790   in Female Mice. *Endocrinology* **158**, 3716–3723 (2017).

791   25. Pnevmatikakis, E. A. & Giovannucci, A. NoRMCorre: An online algorithm for piecewise rigid

792   motion correction of calcium imaging data. *J. Neurosci. Methods* **291**, 83–94 (2017).

793   26. Friedrich, J., Zhou, P. & Paninski, L. Fast online deconvolution of calcium imaging data. *PLOS*

794   *Comput. Biol.* **13**, (2017).

795   27. Rice, J. A. *Mathematical Statistics and Data Analysis*. (Thomson/Brooks/Cole, Belmont, CA,

796   2007).

797   28. Tsuyuzaki, K. *et al.* WormTensor: a clustering method for time-series whole-brain activity data

798   from C. elegans. *BMC Bioinformatics* **24**, 254 (2023).

799   29. Paparrizos, J. & Gravano, L. k-Shape: Efficient and Accurate Clustering of Time Series.

800   *SIGMOD Rec* **45**, 69–76 (2016).

801   30. Kolmogorov, A. Sulla determinazione empirica di una legge didistribuzione. *Giorn Dellinst Ital*

802   *Degli Att* **4**, 89–91 (1933).

803   31. Smirnoff, N. On the estimation of the discrepancy between empirical curves of distribution for

804   two independent samples. *Bull Math Univ Moscou* **2**, 16 (1939).

805   32. Fisher, S. R. A. *Statistical Methods for Research Workers*. (Oliver & Boyd, 1970).

806   33. Rousseeuw, P. J. Silhouettes: A graphical aid to the interpretation and validation of cluster

807   analysis. *J. Comput. Appl. Math.* **20**, 53–65 (1987).

Acknowledgements

809   We thank Ms Sumi Mathew for her technical support with genotyping mice.

810   S.F., M.V., K.T.A., K.T.O., X.F.L. gratefully acknowledge the financial support of BBSRC via grants

811   BB/W005913/1 (KCL), BB/W005883/1 (Exeter) and BB/S019979/1. KTA gratefully acknowledges

812   the financial support of the EPSRC via grant EP/T017856/.

Contributions

814   J.Y. contributed data acquisition and data analysis, S.F. contributed data analysis, K.N. contributed

815   modeling, M.V. contributed managing data analysis, X.F.L. contributed managing data acquisition,

816   H.Y. contributed data acquisition, Y.L. contributed data acquisition, J.Y. contributed data acquisition,

817   O.H. contributed data acquisition, R.D.B. contributed data acquisition, K.T.O., K.T.A. and M.V.

818   contributed to managing the project. All authors contributed to writing the manuscript.

Competing interests

820   The authors declare no competing interests.

Corresponding author

822   Correspondence and requests for materials should be addressed to Krasimira Tsaneva-Atanasova

823   (K.Tsaneva-Atanasova@exeter.ac.uk), Kevin T. O'Byrne (kevin.o'byrne@kcl.ac.uk), Margaritis

824   Voliotis (m.voliotis@exeter.ac.uk).

Open Access