Polymerase mutations underlie early adaptation of H5N1 influenza virus to dairy cattle and other mammals.

Vidhi Dholakia[1#], Jessica L. Quantrill[2,3#], Samuel Richardson[3#], Nunticha Pankaew[4#], Maryn D. Brown[2#], Jiayun Yang[3#], Fernando Capelastegui[1], Tereza Masonou[5], Katie-Marie Case[5], Jila Ajeian[5], Maximillian N.J. Woodall[5], Callum Magill[6], Graham Freimanis[3], Amy McCarron[3], Ecco Staller[7], Carol M. Sheppard[2], Ian H. Brown[3], Pablo R. Murcia[6], Claire M. Smith[5], Munir Iqbal[3], Paul Digard[4], Wendy S. Barclay[2], Rute M. Pinto[4*], Thomas P. Peacock[2,3*] and Daniel H. Goldhill[1*]

[1]Department of Pathobiology and Population Sciences, Royal Veterinary College, London, UK
[2]Department of Infectious Disease, Imperial College London, London, UK
[3]The Pirbright Institute, Woking, UK
[4]The Roslin Institute, University of Edinburgh, UK
[5]Great Ormond Street UCL Institute of Child Health, London, UK
[6]MRC—University of Glasgow Centre for Virus Research, Glasgow, Scotland, UK
[7]Sir William Dunn School of Pathology, The University of Oxford, Oxford, UK

[#]Equal author contributions
*co-corresponding authors: rpinto2@exseed.ed.ac.uk, tom.peacock@pirbright.ac.uk, dgoldhill@rvc.ac.uk

Abstract

In early 2024, an unprecedented outbreak of H5N1 high pathogenicity avian influenza was detected in dairy cattle in the USA[1]. As of mid-2025 the epidemic is ongoing, resulting in spillbacks into poultry, wild birds and other mammals including humans[2]. Here, we present molecular and virological evidence that the cattle B3.13 genotype H5N1 viruses rapidly accumulated adaptations in polymerase genes that enabled better replication in bovine cells and tissues, as well as cells of other mammalian species including humans and pigs. We find evidence of several mammalian adaptations gained early in the evolution of these viruses in cattle including PB2 M631L, which is found in all cattle sequences, and PA K497R, which is found in the majority. Structurally, PB2 M631L maps to the polymerase-ANP32 interface, an essential host factor for viral genome replication. We show that this mutation adapts the polymerase to better interact with bovine ANP32 proteins, particularly ANP32A, and

thereby enhances virus replication in bovine mammary systems and primary human

airway cultures. Importantly, we show that ongoing evolution during 2024 and 2025 in

the PB2 gene, including E627K and a convergently arising D740N substitution, further

increase polymerase activity and virus replication in a range of mammalian cells. Thus,

the continued circulation of H5N1 in dairy cattle not only allows virus adaption improving

replicative ability in cattle, but also increases the risk of zoonotic spillover.

Introduction

Introduction

The natural host reservoirs of influenza A viruses (IAV) are wild waterfowl and

shorebirds. However, IAV can spillover and adapt to infect mammalian species, such as

pigs, humans, horses and dogs[3]. In 2020, a novel genotype of H5N1 clade 2.3.4.4b

emerged in Europe and spread around the world causing a panzootic in birds[4]. This

panzootic genotype and its progeny are notable for their frequent spillover into

mammals, including confirmed or potential mammal-to-mammal spread[5] in farmed mink

in Europe[6-8] (2022-2023), semi-aquatic marine mammals in South America and

Antarctica[9-13] (2023-), and recently, dairy cattle in the USA[1,14] (2024-).

In early 2024, an unexplained drop in milk production in cattle in the USA was

attributed to infection with H5N1 clade 2.3.4.4b, of the B3.13 genotype[1,14]. Infected

cows were initially observed in Texas but as of June 2025, over 1000 infected herds

have been recorded across 17 US states[1,14]. This outbreak was most likely caused by a

single spillover from wild birds[1,14,15], with continuous transmission between dairy cattle

(hereafter referred to as 'cattle') mediated by human activities such as contamination of

milking machinery within farms[16], and movement of infected animals and/or

contaminated equipment between farms and states[1]. The virus does not cause high

mortality in cattle, instead mostly causing mastitis and modest respiratory distress[14,17].

In addition to cattle, several other species have been infected following direct or indirect

exposure to infected cattle including chickens, turkeys, peridomestic birds, farmed

alpaca, domestic cats[1], and humans[2,18]. Infections have been repeatedly fatal in cats

that ingested infected milk, but in humans the B3.13 virus has been associated with

conjunctivitis and/or mild respiratory disease[2,18].

Avian influenza viruses (AIVs) usually have restricted replication in mammalian

hosts[3]. However, they can rapidly accumulate mammalian adaptations that overcome

this block, often in the viral RNA-dependent RNA polymerase[5,19,20]. In particular, ANP32

proteins are essential host-encoded factors co-opted by the virus to support genome

replication[21,22]. All birds excluding palaeognaths (e.g. ostriches and related birds), have

an ANP32A gene with an additional exon resulting in splice variants encoding proteins

with an extra 33 amino acids relative to mammalian orthologues[23]. Avian-origin

influenza viruses require polymerase mutations to utilise the shorter mammalian

ANP32A/B proteins[24]. The most widespread, and best characterised adaptive mutations

are those in the PB2 subunit of the polymerase including PB2 E627K[23], Q591R/K[25,26],

and D701N[26,27], all of which have been detected during other mammalian H5N1

outbreaks[7-13,28] but are absent in the initial dairy cattle viruses. However, alternative

adaptive mutations in the influenza polymerase that enable the virus to co-opt

suboptimal ANP32 proteins are possible[29-31]. Here, we investigated potential

mammalian adaptations in the polymerase of cattle-adapted H5N1 viruses and identify

two key mutations that allow the B3.13 virus to replicate in cells and tissues from cattle

and other mammalian species, including humans.

Results

Phylogenetic analysis of the cattle H5N1 polymerase identified potential adaptive

mutations.

To understand how an avian influenza virus had adapted to efficiently replicate in

cells of a novel mammalian species, we constructed a phylogeny using concatenated

genomes of all influenza A viruses isolated from cattle, virus from the first reported

human case and wild avian viruses from September 2023-March 2024 (Supplementary

Figure S1). Viruses from cattle formed a monophyletic group suggesting a single

spillover event from wild birds[1,15] (Figure 1A). The sequence of the first case of human

infection with a B3.13 H5N1 virus[2] was an outlier relative to all subsequent cattle B3.13

viruses (Figure 1A, Supplementary Figure S1) and contained the known mammalian

adaptations PB2 E627K[32] as well as PA K142E[33] relative to the closest avian relatives

(Supplementary Figure S1). However, these adaptive mutations were not found in any

cattle virus sequences, which instead all contained PB2 M631L as well as PB2 E362G

and PA L219I. PB2 M631L has previously been reported to adapt avian H1N1, H5N1

and H10N7 viruses to mammals[32,34,35], as well as H9N2 viruses to gene-edited chickens

with a non-functional ANP32A[31]. Early cattle sequences grouped into three clades with

additional mutations in the polymerase segments including i) PB2 E677G[36,37], ii) PA

I13V and E613K, and iii) PA K497R, which was found in the largest cattle clade (Figure

1A, Supplementary Figure S1), consisting of ~95% of sequences (as of December

2024) and has previously been associated with influenza adaptation to mammals[38].

Minor clade i (PB2 E677G) was found solely in Texas and appeared only at the

beginning of the outbreak whereas minor clade ii (PA I13V) was detected across

multiple states, including, most recently, in Colorado.

Activity of bovine H5N1 polymerase in mammalian cells maps to PB2 M631L and PA

K497R

117   To test whether the mutations identified from the phylogenetic analysis were

adaptive in cattle, we first constructed a 'minireplicon' system suitable for testing AIV

polymerase activity in bovine cells. This consisted of co-transfecting bovine cells with a

custom-made minigenome reporter that uses a bovine RNA polymerase I (Pol I)

promoter to express a viral-like RNA encoding firefly luciferase flanked with non-coding

regions from influenza A segment 8/NS[39] along with plasmids expressing influenza virus

ribonucleoprotein components, PB2, PB1, PA and nucleoprotein (NP). This species-

specific minireplicon system showed 10-20-fold higher activity in bovine cells than one

using an equivalent human PolI reporter (Supplementary Figure S2). Using this bovine-

specific minireplicon reporter, we showed that the reconstituted polymerase complex of

A/dairy cattle/Texas/24-008749-001-original/2024 (cattle/Texas) was highly active in

MAC-T bovine mammary epithelial cells[40] (Figure 1B).

129   We next reconstructed an avian-like ancestor to represent the polymerase

complex of the virus that mostly likely entered the cattle population, by reverting

potential adaptive mutations from the PB2 and PA segments of cattle/Texas: PB2

M631L, PB2 E362G, PA K497R and PA L219I (Figure 1A, Supplementary Figure S1).

Whilst still active above background, the avian-like ancestral polymerase was much less

active than its WT cattle/Texas counterpart (Figure 1B). To begin to identify key

changes, we assessed the impact of adding back the mutations that arose between this

reconstructed avian-like ancestor, and the cattle virus shared common ancestor (PB2

M631L, PB2 E362G and PA L219I). Neither PB2 E362G nor PA L219I increased

polymerase activity in bovine cells relative to the avian-like ancestor background, either

individually or in combination, and PB2 E362G even reduced activity (Figure 1B). In

contrast, PB2 M631L increased activity by around 10-fold, suggesting this was the key

polymerase mutation that arose early in the adaptation of H5N1 to dairy cattle.

We also investigated later mutations that arose in the major clade iii of the cattle

H5N1 virus (Figure 1A). We first tested the effect of the PA mutation K497R that is

maintained across all sequences of the major bovine clade. Reverting both PB2 631L to

M and PA 497R to K (generating a 'minimal avian-like precursor' or mAP) resulted in a

~100-fold decrease in polymerase activity in bovine cells (Figure 1C). The reintroduction

of PB2 M631L alone increased the signal by around 50-fold. Reintroducing PA K497R

had a more modest effect (~10-fold increase in signal). Neither mutation alone fully

recapitulated the activity of the WT polymerase, indicating that the PA change was

further adaptive. The same mutations had minimal impacts (<2-fold) in avian cells

(chicken fibroblasts, DF-1; Figure 1D), consistent with an adaptive function in cattle cells

rather than simply altering the intrinsic activity of the viral polymerase.

We next tested the impact of these two mutations on polymerase activity in cells

from two other mammalian hosts relevant to influenza viruses - humans and swine,

again using host species-specific minireplicon reporters (Figure 1E, F). As seen in the

bovine cells, both adaptations individually significantly enhanced polymerase activity,

with PB2 M631L having the greater effect, suggesting it is the major mammalian

adaptation across different mammalian species. Also, as observed in bovine cells, we

noted a cumulative effect of the two substitutions together.

To investigate whether PB2 M631L and PA K497R could be universal

mammalian adaptations for H5 viruses, we introduced them into other H5N1

polymerase backgrounds: a recent 2.3.4.4b virus (AIV07-

A/chicken/England/053052/2021), an EU reference laboratory (EURL) genotype C

common in the UK in 2021/22 and a distantly related H5N1 IAV polymerase

(A/turkey/England/50-92/1991; 50-92), both of which originated from low pathogenicity

avian influenza virus donors[41]. Relative to cattle/Texas, AIV07 and 50-92 have amino

acid identities for PB2 of 98% and for PA of 98% and 97%, respectively. In these

additional genetic backgrounds, both PB2 M631L and PA K497R significantly enhanced

polymerase activity in bovine cells and human cells (Supplementary Figure S3A-F).

When compared to the well-characterised mammalian adaptation PB2 E627K, we saw

that neither individual mutation increased polymerase activity to the level of E627K, but

when combined, there was no statistical difference seen between PB2 M631L/PA

K497R and E627K, suggesting the combined mutations are just as potent. There was,

however, a trend that E627K alone had a larger effect relative to PB2 M631L + PA

K497R in the human cells compared to the cattle cells. Overall, this suggests both PB2

M631L and PA K497R are *bona fide* mammalian adaptations, relevant across diverse

AIVs.

PB2 M631L, and to a lesser extent PA K497R, enable efficient replication of bovine

H5N1 in mammalian cells and explants.

To further test the importance of PB2 M631L and PA K497R in adapting virus to

replicate in mammalian cells, we generated a reverse genetics system for cattle/Texas

and made viruses bearing mutations at these sites. To work safely at BSL2, we rescued

viruses as 2:6 reassortants containing the HA and NA of the attenuated vaccine strain,

A/Puerto Rico/8/1934 (PR8)[42] and the remaining genes from cattle/Texas. We also

conducted work with full reverse genetics derived viruses at ACDP3/SAPO4 (BSL3+).

We compared multicycle replication of the WT cattle/Texas virus to either the individual

or double PB2/PA reversions in bovine *ex vivo* mammary gland explants (Figure 2A).

Compared to the minimal avian-like precursor (WT cattle/Texas virus with PB2 M631L

and PA K497R reverted), the viruses with PB2 M631L alone or in combination with PA

K497R showed a significant, >10-fold increase in infectious virus titres at 24 and 48

hours. However, PB2 M631L alone grew to significantly lower titres than the WT virus

that contained K497R as well at 48 and 72 hours post-infection (Figure 2A). This was

reflected both in released virus from the explants, as well as tissue associated virus

harvested at 72 hours post-infection (Figure 2B).

197   Consistent results were also seen in MAC-T cells and bovine alveolar cells (BAT

II; Figure 2C-E) where PB2 M631L, but not PA K497R alone significantly enhanced

replication across multiple tested time points. Again, PB2 M631L replicated to

significantly lower titres than the wild type virus also containing PA K497R. Overall,

these results in the context of infectious virus are consistent with the minireplicon data

that show PB2 M631L, and to a lesser degree PA K497R, are responsible for this virus's

ability to replicate in bovine cells.

204   We next tested the replication of these viruses in several human lung cell lines

(MRC5, A549 and Calu3; Figure 2F-H). Consistent with the minireplicon assays in

human cells (Figure 1E) and virus replication phenotypes in bovine cells (Figure 2A-E),

the PB2 M631L mutation alone, or combined with PA K497R, significantly enhanced

virus titres compared to the minimal avian-like precursor virus across multiple time

points in the human cell lines. While PA K497R alone showed only minor effects, PB2

M631L alone replicated to significantly lower titres than the WT virus (cattle/Texas),

again showing PA K497R, in particular, boosted replication when paired with PB2

M631L. We also tested the replication of the WT and mAP virus in primary human nasal

epithelial cultures (hNECs) from 3 adult donors and maintained at air liquid interface

(Figure 2I). The mAP virus replicated to significantly lower titres than WT in all 3 donors

at 72 hours post-infection. Overall, these data further establish that PB2 M631L is a

major mammalian adaptation that enhances virus replication in both bovine and human

cells.

218   Next, we tested the growth of these viruses in a pair of avian cell lines (chicken

219   CLEC213 and duck CCL-141; Figure 2J,K). All mutants showed similar growth kinetics

220   to the wild type virus, suggesting that these mutations have a minimal impact on virus

221   fitness/replication in avian cells.

222   Across the growth experiments in Figure 2, we also compared the relative

replication of the mAP and mAP containing PB2 E627K. Consistently, across all bovine

cell lines and explants (Figure 2A-D), the combination of PB2 M631L and PA K497R

resulted in significantly higher titres than mAP + E627K. Conversely, in the human cells

(Figure 2F,G), the mAP + E627K virus replicated to higher titres (albeit not significantly)

than the WT cattle virus. Overall, this might suggest why PB2 M631L and PA K497R

were favoured in the cattle virus, whereas PB2 E627K is more common in human

infections.

All available H5N1 sequences from this cattle outbreak contain PB2 M631L but

PA K497R is only found in the major clade of cattle sequences (clade iii). Minor clade i

and ii viruses contain alternative PB2 or PA mutations (Figure 1A, supplementary Figure

1). To investigate whether these minor clade mutations represent alternative bovine

adaptations, we recapitulated these genotypes by individually introducing PA I13V and

E613K from clade i or PB2 E677G from clade ii into the cattle/Texas polymerase

background with PA R497 reverted to the avian consensus 497K. PA I13V and E613K

together exerted a small but significant increase in polymerase activity over PB2 M631L

alone showing a ~3-fold increased minireplicon signal in bovine cells (Supplementary

Figure S4A) and human cells (Supplementary Figure S4B). The introduction of PB2

E677G had no impact in bovine cells but resulted in a significant ~4-fold increase in

polymerase activity in human cells (Supplementary Figure S4B). The polymerase

mutations found in the minor clade i and ii cattle viruses had at most a very minor

impact (<1.5-fold) in polymerase activity in avian cells (Supplementary Figure S4C).

Overall, this suggests that PB2 M631L was the initial potent mammalian adaptation, and

a range of other polymerase mutations then 'fine-tuned' polymerase activity in bovine

cells.

More recent mutations in cattle sublineages further enhance B3.13 H5N1 polymerase

activity in mammalian but not avian cells.

By continually monitoring sequences from the ongoing dairy cattle outbreak, we

detected incidences of further, known mammalian adaptations arising within polymerase

genes of the cattle viruses[26]. This included a single emergence of a cluster of

sequences containing PB2 Q591R, at least three independent emergences of PB2

D740N across the tree (Supplementary Figure S1), as well as a more recent cluster

from 2025 containing PB2 E627K. To test the impact of these changes we introduced

them into the WT cattle/Texas polymerase. PB2 D740N significantly boosted

polymerase activity in bovine (~4-fold increased signal), human (~2-fold increase signal)

and swine cells (1.2-fold increase; Figure 3A-C). PB2 Q591R led to a smaller impact,

reaching statistical significance in the human cells (Figure 3A, B). The introduction of

PB2 E627K on top of M631L consistently led to the highest polymerase activity (Figure

3A-C). However, neither PB2 E627K nor D740N increased activity in avian cells,

relative to the wild-type cattle/Texas polymerase (Figure 3D). We conclude that these

mutations indicate ongoing adaptation of H5N1 viruses to further optimise polymerase

function in cattle.

To further investigate the impact of these ongoing mammalian adaptations, we

rescued WT cattle/Texas + PB2 D740N. For work at CL2, we also rescued PR8:H5N1

2:6 reassortants of cattle/Texas with PB2 D740N or PB2 E627K. We initially tested the

ability of the viruses to replicate in bovine mammary explants, and found that while PB2

D740N closely tracked the wild type virus, PB2 E627K reached significantly higher titres

at early time points (Figure 4A). Similar results were seen in MAC-T and BAT-II cells

where both D740N and E627K significantly enhanced replication in the WT across

multiple time points (Figure 4B,C).

We also tested 2:6 virus in MRC5 and A549 cells (Figure 4D,E) and found both

D740N and E627K significantly increased virus titres across multiple time points. In a

human lung cell line (Calu-3), using the full H5N1 virus, D740N trended towards higher,

albeit but non-significantly higher titres than WT virus (Figure 4F). Finally, we performed

multicycle replication curves with whole H5N1 virus in primary differentiated human

nasal epithelial cultures (hNECs), maintained at air liquid interface, taken from three

independent adult donors (Figure 4G-I). PB2 D740N behaved variably between donors

and did not reach titres significantly differently to the WT virus. We performed whole

genome sequencing on the H5N1 viruses from two of the donors at 72 hours post-

infection and found no evidence of adaptation or selection at the consensus level

following virus replication.

Finally, we tested virus replication kinetics (in the 2:6 background) in chicken and

duck cells (Figure 4J, K). We saw no significant difference in virus replication kinetics.

The PB2 E627K virus showed consistently higher titres than the wild type
suggesting this mutation likely increases the ability to replicate in both human and
bovine cells, while the PB2 D740N containing viruses showed a variable phenotype,
suggesting that if this mutation does impact the virus, the effect is subtle and not
consistently captured by these assays. Both E627K and D740N also appear to have a
neutral impact on the ability to replicate in avian cells, suggesting these mutations could
be maintained if viruses spill back into avian species.
294
295   *Mutations in the polymerase of the cattle H5N1 viruses adapt the polymerase to use*
296   *bovine ANP32 proteins.*
297
PB2 residue 631 is located close to the interface between influenza polymerase
and the host factor ANP32 (Figure 5A, Supplementary Figures S5), nearby several
other mammalian ANP32-adapting mutations such as PB2 E627K and Q591R[26,43]. We
therefore hypothesised that the B3.13 polymerase adapted to bovine cells by increasing
its ability to co-opt bovine ANP32 proteins. To test this, we performed minigenome
assays in human cells which lack ANP32A, ANP32B and ANP32E (eHAP triple
knockout cells[30]) complemented with ANP32 proteins from other relevant species
including cattle, human, pigs, and chicken. Only chicken ANP32A robustly supported
viral polymerases from avian (AIV07, an early European 2.3.4.4b panzootic genotype C
virus; Figure 5B) or avian-like viruses (the minimal avian-like precursor; Figure 5C).
Neither human or bovine ANP32A or B supported any significant activity of avian virus
polymerases. In contrast, both human and bovine ANP32A/B supported mammalian-
adapted polymerases: cattle/Texas (Figure 5D) and a pandemic 2009 H1N1 virus
(Figure 5E), A/England/195/2009 (Eng195). However, bovine ANP32A was consistently
and significantly better able than bovine ANP32B to support polymerase activity of all
constellations tested. We also assessed the effect of the individual cattle/Texas
mammalian adaptations, PB2 M631L and PA K497R on complementation by the
different ANP32 homologues (Figure 5F). Either mutation alone significantly enhanced
polymerase activity supported by bovine ANP32A or ANP32B, with PB2 M631L showing
a stronger effect with all ANP32 orthologues than PA K497R. Thus, we conclude that

the major adaptive mutation PB2 M631L detected in the bovine influenza outbreak has

adapted IAV for replication in cattle by increasing its capacity to co-opt bovine ANP32A.

Finally, to test the mechanism by which the bovine H5N1 virus was able to better

use bovine ANP32 we performed a split-luciferase assay to test protein-protein

interactions (Figure 5G). We tagged the cattle/Texas PB1 with one half of a split

Gaussia luciferase and tagged different ANP32 proteins with the other half of the split

luciferase. We tested the ability of the minimal Avian Precursor or WT cattle/Texas

polymerases to interact with bovine or human ANP32A or B. We found that the

polymerase mutations in cattle/Texas significantly enhanced the ability of the

polymerase to interact with both bovine and human ANP32A/B. A stronger interaction

was seen with bovine ANP32A compared to ANP32B suggesting that during adaptation

to cattle, PB2 M631L and PA K497R have specifically arisen to promote effective

binding and use of bovine ANP32A protein.

Discussion

Our data using reconstructed viral polymerases in host-specific minireplicon and

infection assays in cells and explants from different relevant influenza host species

show that PB2 M631L is the key adaptive mutation that allowed B3.13 H5N1 high

pathogenicity avian influenza virus to efficiently replicate in cattle. PB2 M631L enables

the polymerase complex to co-opt bovine ANP32 proteins to support its activity, with a

strong preference for bovine ANP32A over ANP32B.This is likely due to bovine

ANP32B containing an arginine at position 153, which we have previously shown is

responsible for the reduced ability of canine ANP32B to support influenza polymerase[26].

Structurally, in the complex between the influenza polymerase and host ANP32, PB2

residue 631 is located near amino acids 627 and 591 which are well characterised sites

that mutate to allow adaptation of AIV polymerase to mammalian ANP32 proteins[26,43].

However, the exact mechanism of how PB2 M631L allows the use of mammalian

ANP32s remains unresolved, although our data suggests directly enhanced interaction.

Structural studies show that ANP32 bridges the asymmetric dimer of influenza

polymerase comprising a replicating subunit and a recruited subunit that will be used to

encapsidate the nascent RNA strand. PB2 E627K results in a switch from a negative to

a positive charge enhancing the interaction between the encapsidating subunit of the

polymerase replication dimer and ANP32 residue E151, while also avoiding the clash

between the negatively charged LCAR region of mammalian ANP32s and the

replicating polymerase subunit[43,44]. This mechanism also likely applies to adaptive

charge switch mutations at residue 591 from Q to R/K[43]. PB2 M631L does not result in

an alteration of charge but presumably changes the polymerase structure in this region

to better accommodate the interaction between either or both polymerase subunits and

ANP32. Recent work from Gu et al. using the cattle virus polymerase confirmed that

PB2 M631L increases activity in human cells at both 33°C and 37°C[32]. Previous work

has shown that PB2 M631L adapts avian H10N7 viruses to mice[34] and enables an

avian H9N2 virus to use otherwise suboptimal ANP32 proteins[31,45]. These reports,

combined with our results in the context of a second 2.3.4.4b genotype and an older

non-goose/Guangdong H5N1 suggest that M631L is a general mammalian adaptive

mutation for influenza polymerase that is not specific to a particular virus genetic

background. However, why this mutation emerged in cattle rather than the more potent

PB2 E627K remains unclear. We consistently found when using the minimal Avian

Precursor (cattle WT with PB2 M631L + PA K497R removed), the addition of E627K

showed lower replication in bovine, but not human systems, than the combined addition

of M631L and PA K497R. There was also similar or lower polymerase activity observed

in the corresponding minigenome assays. Our previous work has shown that E627K is

biased towards emerging in hosts or systems where the virus preferentially uses

ANP32B[26], however in cattle we found that ANP32A better supported the polymerase.

E627K has arisen experimentally in cattle infected with a contemporary European H5N1

2.3.4.4b virus[17].

Following fixation of PB2 M631L in cattle, additional mutations in the polymerase

arose in three separate clades. Most cattle sequences contain PA K497R, which

increased polymerase activity in the absence of M631L and virus replication in bovine

mammary cells. Structurally, PA K497R sits in a loop close to the PB2 M631L and

ANP32 interface, though may not be directly interacting with them (Figure 5A).

Additionally, PA K497R sits at the FluPol symmetric dimer interface and therefore may

modulate the dynamics of dimer formation in the same manner as the mammalian
adaptation PA Q556R[46]. Of the mutations defining the other two early cattle virus
clades, the combination of PA I13V + E613K or PB2 E677G also showed modest
increases in activity, although neither group spread particularly widely. Neither PA I13V
or E613K have previously been associated with mammalian adaptation but E613K does
fall on the polymerase asymmetric dimer interface (Supplementary Figure S5B) and
could be altering polymerase regulation[43].

Cattle/Texas virus has extremely robust replication in the bovine mammary
gland, with very high titres of infectious virus being reported from milk[1,14,17]. However,
further mutations that increase polymerase activity in bovine cells are still being
reported as the outbreak continues. There have been three separate emergences of
PB2 D740N which we show here boosted polymerase activity above the wild-type virus.
PB2 E627K has also arisen once in conjunction with M631L in a cluster of cattle and
appears to be continuing to expand as of July 2025, consistant with this combination
giving a robust enhancement in polymerase activity in combination with M631L in our
minireplicon assay, and a replicative advantage in cattle mammary explants. It will be
vital to continue to monitor evolution of the polymerase (and other viral genes), and how
mutations may impact infectivity and transmissibility of these viruses in cattle and other
mammals.

In 2025, USDA reported two further spillovers of H5N1 into cattle in Nevada and
Arizona from the D1.1 genotype[20,28], which was the predominant genotype in wild birds
at the time of spillover. Interestingly, neither cluster has PB2 E627K nor PB2 M631L,
despite D1.1 PB2 having >98.5% amino acid identity with the B3.13 PB2 (and similarly
arising from low pathogenicity avian influenza virus of North American waterfowl). The
Nevada cluster has the known mammalian ANP32 adaptive mutation, PB2 D701N[26],
whereas the Arizona cluster has several mutations in the polymerase relative to the
closest avian sequences, which have yet to be tested for mammalian activity, such as
PB2 R379K and D678N. These independent emergences provide further evidence that
E627K does not have a strong selective advantage to the influenza polymerase in
cattle, and that there are many different viable pathways for H5N1 polymerase adaption
in mammals.

What is the current risk to humans posed by the cattle virus? Our results show

that the cattle virus containing mammalian adaptations can replicate better in

mammalian cells than the minimal avian-like precursor virus, and is therefore a clear

risk to mammals which have consumed infected milk such as cats and raccoons[1]. Thus

far, these cattle viruses show inefficient airborne transmissibility in ferret

experiments[32,47] and have not adapted to transmit via respiratory route or to efficiently

use human receptors (α-2,6-linked sialic acids) which would be essential for human-to-

human transmission[48-51]. In addition, there are also other potential blocks to emergence

posed by antiviral host factors such as MxA to which the cattle virus remains

susceptible[52]. Nonetheless, frequent exposure of an antigenically novel, high

pathogenicity influenza virus that can replicate well in human cells is concerning. Our

data suggest that the emerged cattle influenza virus is also capable of efficiently

infecting avian and swine cells, further increasing the risk of spillover from cattle to other

species. In fact, the mammalian-adapted virus appeared to have no fitness cost in avian

cells, explaining the high propensity of this virus to spillover into poultry causing multiple

outbreaks, and suggesting such a virus could maintain its mammalian adaptations if it

spilt back into wild waterfowl. With each human infection and increased polymerase

activity leading to higher levels of replication, there is a danger of further evolution

changing viral receptor properties. Additionally, a reassortment event with human

seasonal influenza viruses could lead to a novel virus, particularly during the Northern

hemisphere winter influenza season.

Recent experimental infections in cows suggest that the ability of the North

American B3.13 cattle influenza virus to spread via milk is not unique and that other

mammalian viruses are capable of transmitting via this route[17]. Phylogenetic data

suggests that this originated from a single spillover event from wild birds to cattle and,

despite recent spillovers from D1.1, such spillovers are likely unusual as there is scant

evidence of IAV in cattle prior to 2024. Nonetheless, in the absence of an effective

control strategy, the high pathogenicity H5N1 virus may now become endemic in US

dairy cattle, requiring continuous monitoring even in the absence of overt disease.

Moreover, many other clades and strains of H5N1 virus continue to emerge through

reassortment, causing zoonotic infections. Urgent development and testing of broadly

reactive H5 influenza vaccines for both animals and humans is a priority.

Methods

Ethics and biosafety

Virus work was undertaken at either containment level 3 (CL3), SAPO4 (for

whole H5N1 reverse genetics-derived viruses) or CL2 (for reverse genetics-derived

viruses with HA and NA from the attenuated vaccine strain A/Puerto Rico/8/1934 (PR8)

and the remaining internal genes from H5N1 viruses). Viruses carrying H5 HA with a

multibasic cleavage site are categorised as specified animal pathogens order (SAPO) 4

and Advisory Committee on Dangerous Pathogens (ACDP) hazard group 3 by United

Kingdom regulations. Work with these viruses was undertaken in a licensed

CL3/SAPO4 facility of The Pirbright Institute under GMRA (BAG-RA-226). CL2 work

with ACDP Hazard Group 2 recombinant influenza viruses was performed at the Roslin

Institute under biological risk assessments BARA 1011 and GMRA 1811. All virus and

GM risk assessments were approved by the appropriate internal committees, as well as

the UK Health and Safety Executive (HSE) and, where necessary, the UK scientific

advisory committee for genetic modification (SACGM).

Ethics for the use of the primary human airway epithelial cultures (hNECs) were

as described previously[53]. Briefly, donors provided written consent and Ethics approval

was given through the Living Airway Biobank, administered through the UCL Great

Ormond Street Institute of Child Health (REC reference: 19/NW/0171, IRAS project ID:

261511, Northwest Liverpool East Research Ethics Committee). Nasal brushings were

obtained by trained clinicians from adult (30–50 years) donors who reported no

respiratory symptoms in the preceding 7 weeks. Brushings were taken from the inferior

nasal concha zone using cytological brushes (Scientific Laboratory Supplies, CYT1050).

All methods were performed following the relevant guidelines and regulations.

Generation of bovine explants came under approval from the University of

Glasgow School of Biodiversity, One Health and Veterinary Medicine (EA26/25).

Cells

Human Embryonic Kidney 293T (293T), Madin-Darby Canine Kidney (MDCK),
MDCK overexpressing chicken ANP32A (MDCK-ggANP32A), chicken immortalised
fibroblast cells (DF-1), human adenocarcinoma cells (A549), duck immortalised
fibroblasts (CCL-141), human immortalised lung fibroblasts (MRC5), and bovine
mammary epithelial cells (MAC-T) were maintained in Dulbecco's modified Eagle
medium (DMEM) supplemented with 10% fetal bovine serum (FBS), and 1% Pen-Strep.
MAC-T at Roslin additionally had 2 mM L-glutamine, 5 µg/ml insulin and 1 µg/ml of
hydrocortisone. Human lung adenocarcinoma (Calu-3) cells were maintained in DMEM
with 10% FCS, 1% Pen-Strep, 2 mM L-glutamine. Swine foetal testes (ST) cells were
maintained in Advanced DMEM, 5% FCS, 1% Pen-Strep, 2 mM L-glutamine. Chicken
lung epithelial cell (CLEC213[54], kindly gifted by Dr Sascha Trapp) were cultured in
DMEM F12 supplemented with 8% FBS, 2 mM glutamine, 1% Pen-Strep. Triple
knockout cells (tKO- Engineered human haploid (eHAP) cells with ANP32A, ANP32B
and ANP32E knocked out) were generated as previously described[46] and maintained in
Iscove's modified Dulbecco's medium (IMDM), 10% FBS, 1% NEAA, 1% Pen-Strep. All
cells were maintained at 37 °C and 5% CO_2.

Primary human nasal epithelial cultures (hNECs) were generated as described
previously[53]. Briefly, basal epithelial cells from nasal brushings were expanded on
mitotically inactivated 3T3-J2 fibroblasts to passage 1 and cryopreserved. Cells were
thawed as required, expanded with 3T3-J2 fibroblasts to passage 2, and seeded onto
collagen I–coated, semi-permeable membrane supports (3 × 10^5 cells/6.5 mm, 0.4 µm
pore size, Transwell; Corning). Differentiation was performed under air–liquid interface
conditions in PneumaCult™-ALI medium for 4 weeks.

Bovine mammary tissue was collected at Sandyford Abattoir from health lactating
cattle. Excess tissue was removed, teat and gland cisterns were dissected and
transferred to transport media comprising of chilled DMEM supplemented with 100
µg/ml Penicillin/ 100 µg/ml streptomycin and 5 µg/ml fungizone. Every 45 minutes,
tissues were transferred to 500ml of fresh transport media. Inside class II biological

safety cabinet, explants were made using a 5mm biopsy punch. Biopsies were added to

24 well plates with 500ul of transport media supplemented with 10% FBS and incubated

for 24h at 37ºC, 5% CO_2 before infection.

Plasmids

Reverse genetics plasmids of the viruses used in this study were produced as

previously described (Table 1)[51,55,56]. Mutants were generated by site directed

mutagenesis. pCAGGS expression plasmids of the polymerase subunits were

subcloned from reverse genetics plasmids.

Construction of the minireplicon luciferase reporter plasmid for use in bovine cells was

performed using a previous strategy used for an equivalent swine reporter, pSPOM2-

Firefly[39] with swine polI promoter replaced with 428bp bovine polI promoter from the

bovine genome, bosTau8 chrUn_GJ059828v1 (1165-1592)[57]. Plasmid map available at

github.com/Flu1/bovinePolI. *Bos taurus Anp32A* (NCBI reference sequence

NM_001195019.1) and *Anp32B (NM_001035074.1)* were ordered as synthetic GeneArt

constructs from ThermoFisher Scientific and subcloned into pCAGGS as described

previously[58].

Virus strain name	Alias	GISAID ID	Whole virus, 2:6 or minireplicon
A/Puerto Rico/8/1934	PR8		2:6 HA and NA only
A/dairy cattle/Texas/24-008749-001-original/2024 (H5N1)	cattle/Texas	EPI_ISL_19094762	All
A/chicken/England/053052/2021(H5N1)	AIV07	EPI_ISL_9012457	minireplicon only
A/turkey/England/50-92/1991(H5N1)	50-92	EPI_ISL_5888	minireplicon only
A/England/195/2009	Eng195	EPI_ISL_29994	Whole virus and minireplicon only

Table 1. Virus strains, aliases, accessions and uses in this study.

Viruses

Viruses were rescued using 8 plasmid pHW2000 (or derivates) sets containing

each genome segment from the named virus. HEK 293T cells were transfected with a

mix containing 250 ng of each plasmid using lipofectamine 2000 according to the

manufacturer's instructions. Media was replaced 24 h post-transfection with serum-free

DMEM with 0.14% bovine serum albumin and 1 µg/ml of L-(tosylamido-2-phenyl) ethyl

chloromethyl ketone (TPCK)-treated trypsin. Virus-containing supernatants were

collected after 48 h, clarified by centrifugation and used to infect MDCK-ggANP32A

cells or inoculate 10-day-old embryonated hen's eggs. Infected MDCK-*Gg*ANP32A were

monitored twice daily and supernatant was harvested, clarified, aliquoted and stored

when clear cytopathic effect was observed (generally between 48 and 72 hpi).Eggs

were incubated for 48 h before humane euthanization at 4°C, after which virus-

containing allantoic fluid was collected, clarified and stored at -80°C before titration by

plaque assay.

Phylogenetics

Dairy cattle sequences submitted prior to 1st December 2024 , avian H5N1

sequences from the USA submitted between 1st September 2023-31st March 2024 and

A/Texas/37/2024 were downloaded from GISAID. Sequences from isolates which

contained PB2, PB1, PA, NP, HA, NA, NS1, NS2/NEP, M1 and M2 segments were

concatenated. Sequences were aligned using MAFFT v7.490 trimmed using pytrimal

with a gap threshold of 0.85 to ensure correct residue numbering (total length = 4466

amino acids). Duplicate sequences and sequences with > 30 ambiguous residues or

gaps were removed. A phylogenetic tree from the cleaned alignment (n=902) was made

in IQTree2 running automatic ModelFinder and the FLAVI+F+I+R2 model[59,60]. Data

manipulation and visualisation was carried out using Python 3.12.2 and R 4.4.1. Code

and list of GISAID accessions can be found at

https://github.com/fernandocapelastegui/Cattle-H5N1-polymerase-adaptation.

Minireplicon assay

293T, MAC-T, or eHAP tKO cells were transfected in 24-well plates with
Lipofectamine 3000 transfection reagent (Thermo Fisher) with the following amounts of
pCAGGS expression plasmids: 40 ng PB2, 40 ng PB1, 20 ng PA, 80 ng NP, 40 ng
Renilla luciferase, 80 ng species-specific (human, bovine, swine or chicken) polI vRNA
Firefly luciferase and 40 ng ANP32 or empty pCAGGS (where appropriate). DF-1 and
ST cells were transfected in 12 well plates using quadruple the plasmid/transfection
reagent mixes to those used for the other cells in 24-well plates. Cells were lysed 24h
post-transfection using passive lysis buffer (Promega) and polymerase activity
measured using the Dual-luciferase Reporter assay system (Promega) and a FLUOstar
Omega plate reader (BMG Labtech). Assays in MAC-T cells were read using Dual-Glo®
Luciferase Assay System (Promega) on an Infinite MPlex plate reader (Tecan). Firefly
luciferase signal has been normalised to Renilla luciferase signal to give relative
luminescence units (RLU).

Virus replication kinetics

Confluent monolayers of CLEC213, CCL-141, A549, MRC5, MAC-T, BAT II, and
Calu-3 cells were washed once with PBS and infected at multiplicity of infection (MOI)
0.01 with virus diluted in serum-free medium for 1 h at 37 °C. Inoculum was replaced
with serum-free medium supplemented with 0.14% BSA and 1 µg/ml TPCK-treated
bovine pancreas trypsin. Calu-3 and MAC-T experiments performed with whole H5N1
virus excluded TPCK trypsin. Supernatants were collected at various times post
infection and stored at -80°C until infectious titres were determined by plaque assay.
Infection of primary human airway cltures was performed as follows. Before
infection cells were washed with Dulbecco's phosphate-buffered saline supplemented
with calcium and magnesium (DPBS++) to remove mucus and debris. Cells were
infected with 200 µl of virus in DPBS++ at an MOI of 0.01 and incubated at 37°C for 1 h.
Inoculum was removed, and cells were washed twice with DPBS++. Time points were

taken by adding 200 µl of DPBS++ and incubating for 10 mins and 37°C before removal
and titration. The time course was performed at 37 °C, 5% CO_2. Time points were taken
at 24-, 48-, 72- and 96-hours post-infection. Infectious titres were determined by plaque
assay on MDCKs.

Infection of explants was performed as follows: individual explants were
transferred to 96-well plates and infected with 5000 PFU of virus in 100 µl of serum free
DMEM for 2h at 37 °C. Explants were transferred to clean 24-well plates, washed 3x
with warm PBS and overlayed with 1 ml of DMEM supplemented with 0.14% fraction V
BSA and 1 µg/ml TPCK. Time points were collected at the indicated times post
infection.

Split Luciferase assays

Split luciferase assays were performed in 293T cells seeded into 48-well plates.
40 ng of PB2, PA, and PB1 tagged with the N-terminus of *Gaussia* luciferase (Gluc1) on
its C-terminus (after a GGSGG linker), were co-transfected with ANP32 tagged with the
C-terminus of *Gaussia* luciferase (Gluc2) on its C-terminus (after a GGSGG linker),
using lipofectamine 3000 as per the manufacturer's instructions. 24 hours post-
transfection, cells were lysed in 60 µl of *Renilla* lysis buffer (Promega), and luciferase
activity was measured using a *Renilla* luciferase kit (Promega) on a VANTAstar plate
reader (BMG Labtech). Normalised luminescence ratios (NLRs) were calculated by
diving the values of the tagged PB1 and ANP32 wells by the sum of the mean of the
control wells, which contained i) untagged PB1 and free Gluc1, and ii) untagged ANP32
and free Gluc2, as described elsewhere[24].

Immunoblotting

Transfected cells were lysed in radioimmunoprecipitation (RIPA) buffer (150mM
NaCl, 50mM TRIS, 0.1% SDS, 1% NP-40, 0.5% sodium deoxycholate, pH 7.4)
supplemented with an EDTA-free protease inhibitor mini tablet (Thermo Fisher),
incubated on ice for 10 minutes, then centrifuged for 10 minutes to remove precipitate.

Lysates were then diluted in 4 x Laemmli buffer (Bio-rad) and loaded onto 4-20%
gradient gels to perform SDS-PAGE. Proteins were transferred onto a PVDF membrane
by semi-dry transfer. Membranes were blocked in TBS/5% milk for 1 hour at room
temperature and incubated overnight at 4°C with the following primary antibodies:
mouse anti-FLAG (Sigma, F1804; 1:250 dilution), rabbit anti-PB2 (GeneTex,
GTX125926; 1:500 dilution), rabbit anti-PB1 (Genetex, GTX125923, 1:250 dilution),
rabbit anti-PA (Genetex, GTX118991; 1:500 dilution), mouse anti-α-Tubulin (Abcam,
ab7291; 1:1250 dilution) rabbit anti-*Gaussia* luciferase (Invitrogen, PIPA1181; 1:1000).
Following four 10-minute TBS/1% Tween20 (TBS-T) washes, membranes were
incubated with the following near-infrared fluorescent secondary antibodies: goat anti-
mouse IgG Alexa FluorRTM 680 (abcam; ab175775; 1:10,000 dilution), goat anti-rabbit
IRDye 800CW (LI-COR, 926-32211; 1:10,000 dilution) for 45 minutes at room
temperature. After a further four TBST washes, and one TBS wash, membranes were
imaged using an Odyssey DLx (Li-Cor Biosciences) using the Image Studio Lite
software.
631
632   *Virus whole genome sequencing*
633
Viral RNA was extracted from the virus inoculums used in hNECs, and the 72-
hour time point using the QIAamp viral RNA extraction kit (Qiagen) according to the
manufacturer's instructions. Viral cDNA was synthesised using the Veso cDNA
synthesis kit (Thermo) with a mix of UTR-specific primers
(TATTCGTCTCACCCAGCAAAAGCAGG and
TATTCGTCTCACCCAGCGAAAGCAGG) with cycling conditions of 42^{o}C for 75 min,
95^{o}C for 2 min. The Optil-F1 (GTTACGCGCCAGCAAAAGCAGG), Optil-F2
(GTTACGCGCCAGCGAAAGCAGG) and Optil-R1
(GTTACGCGCCAGTAGAAACAAGG) primers were used in a PCR from cDNA with
PFU Ultra II DNA polymerase (Agilent). Cycling conditions were 55^{o}C for 2 minutes,
98^{o}C for 2 minutes. 5 x cycles of 94^{o}C for 30s, 44^{o}C 30s, 68^{o}C 3m30s, 35x cycles of
94^{o}C for 30s, 57^{o}C for 30s, 68^{o}C 3m30s, a final extension step of 68^{o}C for 10m followed
by a holding step of 4^{o}C. The PCR products were then purified using the Agencourt

AMPure XP kit per manufacturer's instructions. Purified amplicons were quantified by

Qubit (Thermo) and diluted in nuclease-free water to a final concentration of 0.2ng/µl. 1

ng total of input nucleic acid was processed using the Illumina Nextera XT kit on a Hamilton

NGStar. The resulting libraries were confirmed using the Tapestation 4200 *Agilent) and bead-

normalised according to the manufacturer's specifications before being pooled and diluted to

12.5 pM final concentration with 1% 12.5 pM PhiX (Illumina). The resulting pool was then run on

an Illumina MiSeq using a 2 x 250 v2 MiSeq reagent cartridge and flow cell.

Raw fastq sequences were then imported into Geneious Prime software and trimmed to

remove primers and aligned to the whole genome reference sequence (A/dairy cattle/Texas/24-

008749-001-original/2024 (H5N1)).

Acknowledgements

MRC-5 cells were a kind gift from Dr Finn Grey of the Roslin Institute. Chicken

lung cells (CLEC213[54]) were kindly gifted by Dr Sascha Trapp of INRA Centre Val de

Loire. CCL-141 cells were a kind gift from Dr Leah Golding of University of Nottingham.

MDCK-ggANP32 cells were a kind gift from Professor Massimo Palmarini of the Centre

for Virological Research, Glasgow. The authors would like to thank Dr Martha Nelson

for feedback on the manuscript and Dr Giulia Gallo for useful suggestions on data

visualisation. The authors thank all researchers who have shared genome data openly

via the Global Initiative on Sharing All Influenza Data (GISAID).

V.D. and F.C. were supported by Academy of Medical Sciences Springboard

Grant 1049 and Royal Society Grant 231225 to D.H.G., J.L.Q., T.M, M.N.J.W., C.M.S.,

N.P., R.M.P., M.D.B., J.Y., S.R., P.M., I.H.B, M.I., P.D., W.B. T.P.P. and D.H.G are

supported by the UK Medical Research Council/Department for Environment, Food and

Rural Affairs (Defra, UK) FluTrailMap-One Health consortium [MR/Y03368X/1]. J.L.Q.,

R.M.P., M.D.B., I.H.B., M.I., P.D., W.S.B., and T.P.P. are also supported by the

Biotechnology and Biological Sciences Research Council (BBSRC)/DEFRA

'FluTrailMap' consortium [BB/Y007298/1]. J.L.Q., J.Y., I.H.B., M. I., P.D., W.S.B.,

R.M.P, and T.P P. were supported by the BBSRC/Defra "FluMAP" consortium

(BB/X006123/1). S.R., G.F., A.M., I.H.B., M.I. and T.P.P. are funded by the BBSRC via

the Pirbright Institute's Strategic Programme Grants (ISPGs) [BBS/E/PI/230002A;

BBS/E/PI/230002B], BBSRC National Bioscience Research Infrastructure: High

Containment and Low Containment Services and Science Platforms grants

[BBS/E/PI/23NB0004, BBS/E/PI/23NB0003]. ES was supported by Medical Research

Council (MRC) programme grant MR/X008312/1. P.D. and R.M.P. acknowledge

BBSRC Strategic Programme Grant funding to the Roslin Institute

(BBS/E/RL/230002C). P.R.M. is supported by BBSRC [BB/V004697/1]. R.M.P is

supported by a Chancellor's Fellowship from the University of Edinburgh. P.D. also

acknowledges BBSRC support from the Evolution & Ecology of Infectious Diseases

programme (BB/V011286/1), MRC support from a ONE Health epidemic preparedness

award (MR/Y015045/1) and Defra award FluSwitch (SE2223).

Credit Contributions:
VD – investigation, resources, visualisation
JLQ – investigation, resources
SR – investigation, Writing – review & editing
NP – investigation, Writing – review & editing
MDB – investigation, resources, Writing – review & editing
JY – investigation, resources, Writing – review & editing
FC – data curation, investigation, software, Writing – review & editing
TM - resources
K-MC - resources
JA – resources
MNJW – resources
CM - resources
GF – methodology, investigation, supervision
AM – methodology, investigation
ES – resources, Writing – review & editing
CMSheppard – investigation, supervision, Writing – review & editing
IHB – funding acquisition, supervision, Writing – review & editing
PRM - funding acquisition, resources, supervision
CMSmith - funding acquisition, resources, supervision
MI - funding acquisition, resources, supervision
PD - funding acquisition, methodology, supervision, Writing – review & editing
WSB – conceptualisation, funding acquisition, methodology, supervision, Writing – review &
editing
RMP – investigation, methodology, project administration, supervision, validation
TPP – conceptualisation, formal analysis, investigation, methodology, project
administration, supervision, validation, visualisation, Writing – original draft

DHG – conceptualisation, data curation, formal analysis, funding acquisition, investigation,
methodology, project administration, software, supervision, visualisation, Writing –
original draft
719
720 **Figures**

721
Figure 1. Identification of key mammalian polymerase adaptations in the major clade of bovine H5N1 that enhance polymerase activity in bovine cells. A)

Simplified schematic of cattle polymerase phylogeny and polymerase mutations that
arose in early branches. B-F) Minigenome in (B, C) bovine MAC-T cells (D) chicken DF-
1 (E) human HEK293T, (F) swine ST cells with cattle Texas (WT). B) Reconstructed
avian-like ancestor has a reversion of PB2 E362G, M631L and PA L219I, K497R. C-F)
Minimal avian-like precursor has the reversions of PB2 M631L and PA K497R. Data
normalised to minimal avian-like precursor. Data throughout plotted as the mean of N =
3 independent repeats. (B, E) Expression of PB2 and PA was measured by western
blot, tubulin was used as a loading control. Data throughout plotted as mean + SD.
Statistics throughout performed by one-way ANOVA with multiple comparisons
comparing all data using log-transformed data. B) significance only plotted if increased
over avian-like ancestor. Log-normality determined by Shapiro-Wilk test and QQ plot.
Significance shown by asterisks indicating: *, $0.05 \geq P > 0.01$; **,
$0.01 \geq P > 0.001$; ***, $0.001 \geq P > 0.0001$; ****, $P \leq 0.0001$.
737

738
739
Figure 2. PB2 M631L, and to a lesser extent PA K497R, enhance virus replication
in mammalian, but not avian culture systems. Reverse-genetics derived viruses,
containing the HA and NA genes of the attenuated laboratory strain A/Puerto Rico/8/34,
and the remaining genes from Cattle Texas (2:6 viruses; panels A-D, F,G,J,K) or

reverse genetics derived full H5N1 viruses (whole virus, panels E, H and I) were used to
infect A, B) bovine ex vivo mammary explants, C, E) bovine mammary cells (MAC-T),
D) bovine respiratory cells (BAT II) cells, F) human lung cells (MRC5), G) human lung
adenocarcinoma cells (A549), H) human lung cells (Calu-3), I) primary human nasal
epithelial cultures (hNECs) maintained at air liquid interface. J) chicken lung cells
(CLEC213) or K) duck fibroblasts (CCL-441). Explants were infected with 5000
pfu/explant. Cell lines were infected at an MOI of 0.01. Infectious virus titres were
determined by plaque assay on MDCKs. Data from bovine explants represents N = 34
repeats from 3 donor cattle from 3 different independent experiments. Data from cell
lines plotted as (C), 2 x N = 4, (D,F,G,J, K) 2 x N = 3 technical repeats, (E,H) N = 3
repeats, from a representative repeat of N = 3 independent repeats and I) 3
independent donors with virus replication tested in triplicate. Data throughout plotted as
mean ± SD. Statistics throughout performed by two-way ANOVA on log-transformed
data with multiple comparisons between all data points within a single time point.
Significance only plotted when differences were significant to the minimal avian
precursor (shown as matched colour asterisks), or between the WT cattle/Texas and
mutants (shown as coloured asterisks next to the purple line). Significance shown by
asterisks indicating: *, 0.05 ≥ P > 0.01; **, 0.01 ≥ P > 0.001; ***,
0.001 ≥ P > 0.0001; ****, P ≤ 0.0001.

Figure 3. Ongoing adaptations found in cattle H5N1 viruses further enhance polymerase activity in mammalian cells. Minigenome in (A) bovine MAC-T cells, (B) human HEK293T, (C) swine ST cells or (D) chicken DF-1 with mutant versions of cattle Texas representing potential ongoing adaptation. Data normalised to WT cattle Texas. Data throughout plotted as mean of N = 3 independent biological repeats. B) Matched western blot showing PB2 expression. Data throughout plotted as mean + SD. Statistics throughout performed by one-way ANOVA with multiple comparisons against WT polymerase using log-transformed data. Log-normality determined by Shapiro-Wilk test and QQ plot. Significance shown by asterisks indicating: *, $0.05 \geq P > 0.01$; **, $0.01 \geq P > 0.001$; ***, $0.001 \geq P > 0.0001$; ****, $P \leq 0.0001$.

774
Figure 4. Further emerging mammalian adaptations in the cattle H5N1 PB2 enhance its replication in human and bovine, but not avian culture systems.
Reverse-genetics derived viruses, containing the HA and NA genes of the attenuated laboratory strain A/Puerto Rico/8/34, and the remaining genes from Cattle Texas (2:6 viruses; panels, A-E, J, K) or reverse genetics derived full H5N1 viruses (whole virus,

panels F-I) were used to infect) A bovine *ex vivo* mammary explants, B) bovine
mammary cells (MAC-T), C) bovine respiratory cells (BAT II) cells, D) human lung cells
(MRC5), E) human lung adenocarcinoma cells (A549), F) human lung cells (Calu-3), G-
I) primary differentiated human nasal epithelial cultures (hNECs) maintained at air liquid
interface (hNECs), J) chicken lung cells, or K) duck fibroblasts (CCL-141). Cells were
infected with an MOI of 0.01, explants were infected with 5000 PFU/explant. Infectious
virus titres were determined by plaque assay on MDCKs. A) Data from bovine explants
represents total N = 34 repeats from 3 donor cattle from 3 different independent
experiments. Data from cell lines plotted as (B), 2 x N = 4, (C-E,J,F) 2 x N = 3 technical
repeats, (F) N = 3 repeats, from a representative repeat of N = 3 independent repeats
and G-I) 3 independent donors with virus replication tested in triplicate. WT and avian
precursor data through the figure is the same shown through Figure 2. Data throughout
plotted as mean ± SD. Statistics throughout performed by two-way ANOVA on log-
transformed data with multiple comparisons against WT cattle/Texas. Significance
shown by asterisks indicating: *, $0.05 \geq P > 0.01$; **, $0.01 \geq P > 0.001$; ***,
$0.001 \geq P > 0.0001$; ****, $P \leq 0.0001$.
796

797
798 **Figure 5. Although bovine ANP32A and B proteins are both able to support**
799 **mammalian-adapted polymerases, bovine ANP32A is dominant which correlates**
800 **with a stronger interaction.** A) Key mammalian adaptations identified in this study
801 mapped to the structure of H5N1 influenza A replicase dimer complex with human
802 ANP32B (PDB: 8R1J). ANP32B is shown in cyan, PB1 in purple, PA in violet and PB2

as a salmon ribbon representation. H5N1 cattle PB2 substitutions (Q591R, E627K, M631L and D740N) shown in red and PA (K497R) shown in blue. B-F) Minigenomes in human engineered haploid cells (eHAP) with endogenous ANP32A, ANP32B and ANP32E knocked out were supplemented with ANP32 proteins from different species (chANP32A = chicken ANP32A, swANP32A = swine ANP32A, swANP32B = swine ANP32B, bovANP32A = bovine ANP32A, bovANP32B = bovine ANP32B, huANP32A = human ANP32A, huANP32B = human ANP32B, -ANP = no ANP32 control). Avianised cattle = PB2 M631L and PA K497R reverted. Data normalised throughout to chANP32A. G) Split luciferase assays in 293T cells showing the relative binding of different ANP32 proteins to the trimeric polymerase from cattle/Texas or the avianised version lacking PB2 M631L and PA K497R. PB1 was tagged with the N-terminal section of *Gaussia* luciferase, while the ANP32 proteins were tagged with the C-terminal portion. NLR, normalised luminescence ratios, were calculated from the ratio between signals between tagged and untagged ANP32/PB1 pairs. Dotted line equals a false positive cutoff of <2.5% as used in Mistry et al[61]. B, G) Matched expression of polymerase and ANP32 constructs shown by western blot. Data throughout plotted the mean of N = 3 (B-F) or N = 4 (G) independent biological repeats. Data throughout plotted as mean + SD. B-E) Statistics performed by one-way ANOVA with multiple comparisons - against -ANP32 control, or between species-matched ANP32A and ANP32B proteins, using log-transformed data. F) Statistics performed by two-way ANOVA with multiple comparisons comparing the different polymerase mutants within the same species ANP32, on log-transformed data. G) Statistics performed by multiple unpaired T tests between cattle/Texas WT and the minimal avian precursor. Log-normality determined by Shapiro-Wilk test and QQ plot. Significance shown by asterisks indicating: *, $0.05 \geq P > 0.01$; **, $0.01 \geq P > 0.001$; ***, $0.001 \geq P > 0.0001$; ****, $P \leq 0.0001$.

References

1   Nguyen, T.-Q. *et al.* Emergence and interstate spread of highly pathogenic avian influenza A(H5N1) in dairy cattle in the United States. *Science* **388**, eadq0900 (2025). https://doi.org/doi:10.1126/science.adq0900

2   Uyeki, T. M. *et al.* Highly Pathogenic Avian Influenza A(H5N1) Virus Infection in a Dairy Farm Worker. *N Engl J Med* **390**, 2028-2029 (2024). https://doi.org/10.1056/NEJMc2405371

3   Long, J. S., Mistry, B., Haslam, S. M. & Barclay, W. S. Host and viral determinants of influenza A virus species specificity. *Nature Reviews Microbiology* **17**, 67-81 (2019). https://doi.org/10.1038/s41579-018-0115-z

4   Xie, R. *et al.* The episodic resurgence of highly pathogenic avian influenza H5 virus. *Nature* **622**, 810-817 (2023). https://doi.org/10.1038/s41586-023-06631-2

5　Peacock, T. *et al.* The global H5N1 influenza panzootic in mammals. *Nature* (2024).
https://doi.org/10.1038/s41586-024-08054-z
6　Agüero, M. *et al.* Highly pathogenic avian influenza A(H5N1) virus infection in farmed
minks, Spain, October 2022. *Eurosurveillance* **28**, 2300001 (2023).
https://doi.org/doi:https://doi.org/10.2807/1560-7917.ES.2023.28.3.2300001
7　Kareinen, L. *et al.* Highly pathogenic avian influenza A(H5N1) virus infections on fur
farms connected to mass mortalities of black-headed gulls, Finland, July to October
2023. *Eurosurveillance* **29**, 2400063 (2024).
https://doi.org/doi:https://doi.org/10.2807/1560-7917.ES.2024.29.25.2400063
8　Lindh, E. *et al.* Highly pathogenic avian influenza A(H5N1) virus infection on multiple fur
farms in the South and Central Ostrobothnia regions of Finland, July 2023.
Eurosurveillance **28**, 2300400 (2023). https://doi.org/doi:https://doi.org/10.2807/1560-
7917.ES.2023.28.31.2300400
9　Uhart, M. M. *et al.* Epidemiological data of an influenza A/H5N1 outbreak in elephant
seals in Argentina indicates mammal-to-mammal transmission. *Nature Communications*
15, 9516 (2024). https://doi.org/10.1038/s41467-024-53766-5
10　de Carvalho Araujo, A. *et al.* Mortality in sea lions is associated with the introduction of
the H5N1 clade 2.3.4.4b virus in Brazil October 2023: whole genome sequencing and
phylogenetic analysis. *BMC Veterinary Research* **20**, 285 (2024).
https://doi.org/10.1186/s12917-024-04137-1
11　Leguia, M. *et al.* Highly pathogenic avian influenza A (H5N1) in marine mammals and
seabirds in Peru. *Nat Commun* **14**, 5489 (2023). https://doi.org/10.1038/s41467-023-
41182-0
12　Tomás, G. *et al.* Highly pathogenic avian influenza H5N1 virus infections in pinnipeds
and seabirds in Uruguay: Implications for bird-mammal transmission in South America.
Virus Evol **10**, veae031 (2024). https://doi.org/10.1093/ve/veae031
13　Gamarra-Toledo, V. *et al.* Mass Mortality of Sea Lions Caused by Highly Pathogenic
Avian Influenza A(H5N1) Virus. *Emerging Infectious Disease journal* **29**, 2553 (2023).
https://doi.org/10.3201/eid2912.230192
14　Caserta, L. C. *et al.* Spillover of highly pathogenic avian influenza H5N1 virus to dairy
cattle. *Nature* (2024). https://doi.org/10.1038/s41586-024-07849-4
15　Worobey, M. *et al.* Preliminary Report on Genomic Epidemiology of the 2024 H5N1
Influenza A Virus Outbreak in US Cattle. *Virological.org* (2024).
16　Le Sage, V., Campbell, A. J., Reed, D. S., Duprex, W. P. & Lakdawala, S. S.
Persistence of Influenza H5N1 and H1N1 Viruses in Unpasteurized Milk on Milking Unit
Surfaces. *Emerg Infect Dis* **30**, 1721-1723 (2024).
https://doi.org/10.3201/eid3008.240775
17　Halwe, N. J. *et al.* H5N1 clade 2.3.4.4b dynamics in experimentally infected calves and
cows. *Nature* (2024). https://doi.org/10.1038/s41586-024-08063-y
18　Garg, S. *et al.* Highly Pathogenic Avian Influenza A(H5N1) Virus Infections in Humans. *N
Engl J Med* (2024). https://doi.org/10.1056/NEJMoa2414610
19　Peacock, T. P., Sheppard, C. M., Staller, E. & Barclay, W. S. Host Determinants of
Influenza RNA Synthesis. *Annual review of virology* **6**, 215-233 (2019).
20　Capelastegui, F. & Goldhill, D. H. H5N1 2.3.4.4b: a review of mammalian adaptations
and risk of pandemic emergence. *Journal of General Virology* **106** (2025).
https://doi.org/https://doi.org/10.1099/jgv.0.002109
21　Staller, E. *et al.* ANP32 Proteins Are Essential for Influenza Virus Replication in Human
Cells. *J Virol* **93** (2019). https://doi.org/10.1128/jvi.00217-19
22　Zhang, H. *et al.* Fundamental Contribution and Host Range Determination of ANP32A
and ANP32B in Influenza A Virus Polymerase Activity. *J Virol* **93** (2019).
https://doi.org/10.1128/jvi.00174-19

23   Long, J. S. *et al.* Species difference in ANP32A underlies influenza A virus polymerase
host restriction. *Nature* **529**, 101-104 (2016). https://doi.org/10.1038/nature16474
24   Long, J. S. *et al.* Species specific differences in use of ANP32 proteins by influenza A
virus. *eLife* **8**, e45066 (2019). https://doi.org/10.7554/eLife.45066
25   Yamada, S. *et al.* Biological and Structural Characterization of a Host-Adapting Amino
Acid in Influenza Virus. *PLoS Pathog* **6**, e1001034 (2010).
https://doi.org/10.1371/journal.ppat.1001034
26   Peacock, T. P. *et al.* Mammalian ANP32A and ANP32B Proteins Drive Differential
Polymerase Adaptations in Avian Influenza Virus. *Journal of Virology*, e00213-00223
(2023).
27   Mehle, A. & Doudna, J. A. An Inhibitory Activity in Human Cells Restricts the Function of
an Avian-like Influenza Virus Polymerase. *Cell Host & Microbe* **4**, 111-122 (2008).
https://doi.org/https://doi.org/10.1016/j.chom.2008.06.007
28   Pekar, J. *et al.* Timing and molecular characterisation of the transmission to cattle of
H5N1 influenza A virus genotype D1.1, clade 2.3.4.4b. *Virological.org* (2025).
29   Gilbertson, B., Duncan, M. & Subbarao, K. Role of the viral polymerase during
adaptation of influenza A viruses to new hosts. *Curr Opin Virol* **62**, 101363 (2023).
https://doi.org/10.1016/j.coviro.2023.101363
30   Sheppard, C. M. *et al.* An Influenza A virus can evolve to use human ANP32E through
altering polymerase dimerization. *Nat Commun* **14**, 6135 (2023).
https://doi.org/10.1038/s41467-023-41308-4
31   Idoko-Akoh, A. *et al.* Creating resistance to avian influenza infection through genome
editing of the ANP32 gene family. *Nature Communications* **14**, 6136 (2023).
https://doi.org/10.1038/s41467-023-41476-3
32   Gu, C. *et al.* A human isolate of bovine H5N1 is transmissible and lethal in animal
models. *Nature* (2024). https://doi.org/10.1038/s41586-024-08254-7
33   Mänz, B., Brunotte, L., Reuther, P. & Schwemmle, M. Adaptive mutations in NEP
compensate for defective H5N1 RNA replication in cultured human cells. *Nature
Communications* **3**, 802 (2012). https://doi.org/10.1038/ncomms1804
34   Zhang, X. *et al.* Enhanced pathogenicity and neurotropism of mouse-adapted H10N7
influenza virus are mediated by novel PB2 and NA mutations. *J Gen Virol* **98**, 1185-1195
(2017). https://doi.org/10.1099/jgv.0.000770
35   Patrono, L. V. *et al.* Archival influenza virus genomes from Europe reveal genomic
variability during the 1918 pandemic. *Nature Communications* **13**, 2314 (2022).
https://doi.org/10.1038/s41467-022-29614-9
36   Mänz, B. *et al.* Multiple Natural Substitutions in Avian Influenza A Virus PB2 Facilitate
Efficient Replication in Human Cells. *J Virol* **90**, 5928-5938 (2016).
https://doi.org/10.1128/jvi.00130-16
37   Welkers, M. R. A. *et al.* Genetic diversity and host adaptation of avian H5N1 influenza
viruses during human infection. *Emerg Microbes Infect* **8**, 262-271 (2019).
https://doi.org/10.1080/22221751.2019.1575700
38   Yamayoshi, S. *et al.* Enhanced Replication of Highly Pathogenic Influenza A(H7N9)
Virus in Humans. *Emerg Infect Dis* **24**, 746-750 (2018).
https://doi.org/10.3201/eid2404.171509
39   Moncorgé, O. *et al.* Investigation of influenza virus polymerase activity in pig cells. *J Virol*
87, 384-394 (2013). https://doi.org/10.1128/jvi.01633-12
40   Huynh, H. T., Robitaille, G. & Turner, J. D. Establishment of bovine mammary epithelial
cells (MAC-T): An in vitro model for bovine lactation. *Experimental Cell Research* **197**,
191-199 (1991). https://doi.org/https://doi.org/10.1016/0014-4827(91)90422-Q

41 Byrne, A. M. P. *et al.* Investigating the Genetic Diversity of H5 Avian Influenza Viruses in
the United Kingdom from 2020-2022. *Microbiol Spectr*, e0477622 (2023).
https://doi.org/10.1128/spectrum.04776-22
42 Beare, A. S., Schild, G. C. & Craig, J. W. TRIALS IN MAN WITH LIVE RECOMBINANTS
MADE FROM A/PR/8/34 (H0 N1) AND WILD H3 N2 INFLUENZA VIRUSES. *The Lancet*
306, 729-732 (1975). https://doi.org/10.1016/S0140-6736(75)90720-5
43 Staller, E. *et al.* Structures of H5N1 influenza polymerase with ANP32B reveal
mechanisms of genome replication and host adaptation. *Nature Communications* **15**,
4123 (2024). https://doi.org/10.1038/s41467-024-48470-3
44 Carrique, L. *et al.* Host ANP32A mediates the assembly of the influenza virus replicase.
Nature **587**, 638-643 (2020). https://doi.org/10.1038/s41586-020-2927-z
45 Camacho-Zarco, A. R. *et al.* Molecular basis of host-adaptation interactions between
influenza virus polymerase PB2 subunit and ANP32A. *Nature Communications* **11**, 3656
(2020). https://doi.org/10.1038/s41467-020-17407-x
46 Sheppard, C. M. *et al.* An Influenza A virus can evolve to use human ANP32E through
altering polymerase dimerization. *Nature Communications* **14**, 6135 (2023).
47 Eisfeld, A. J. *et al.* Pathogenicity and transmissibility of bovine H5N1 influenza virus.
Nature **633**, 426-432 (2024). https://doi.org/10.1038/s41586-024-07766-6
48 Chopra, P. *et al.* Receptor-binding specificity of a bovine influenza A virus. *Nature* **640**,
E21-e27 (2025). https://doi.org/10.1038/s41586-025-08822-5
49 Santos, J. J. S. *et al.* Bovine H5N1 binds poorly to human-type sialic acid receptors.
Nature **640**, E18-e20 (2025). https://doi.org/10.1038/s41586-025-08821-6
50 Good, M. R. *et al.* A single mutation in dairy cow-associated H5N1 viruses increases
receptor binding breadth. *Nature Communications* **15**, 10768 (2024).
https://doi.org/10.1038/s41467-024-54934-3
51 Yang, J. *et al.* The haemagglutinin gene of bovine-origin H5N1 influenza viruses
currently retains receptor-binding and pH-fusion characteristics of avian host phenotype.
Emerg Microbes Infect **14**, 2451052 (2025).
https://doi.org/10.1080/22221751.2025.2451052
52 Ankerhold, J., Kessler, S., Beer, M., Schwemmle, M. & Ciminski, K. Replication
Restriction of Influenza A(H5N1) Clade 2.3.4.4b Viruses by Human Immune Factor,
2023-2024. *Emerg Infect Dis* **31**, 199-202 (2025).
https://doi.org/10.3201/eid3101.241236
53 Woodall, M. N. J. *et al.* Age-specific nasal epithelial responses to SARS-CoV-2 infection.
Nature Microbiology **9**, 1293-1311 (2024). https://doi.org/10.1038/s41564-024-01658-1
54 Esnault, E. *et al.* A novel chicken lung epithelial cell line: characterization and response
to low pathogenicity avian influenza virus. *Virus Res* **159**, 32-42 (2011).
https://doi.org/10.1016/j.virusres.2011.04.022
55 Yang, J. *et al.* The Haemagglutinin Genes of the UK Clade 2.3.4.4b H5N1 Avian
Influenza Viruses from 2020 to 2022 Retain Strong Avian Phenotype. *bioRxiv*,
2024.2007.2009.602706 (2024). https://doi.org/10.1101/2024.07.09.602706
56 Schafers, J. *et al.* Pasteurisation temperatures effectively inactivate influenza A viruses
in milk. *Nature Communications* **16**, 1173 (2025). https://doi.org/10.1038/s41467-025-
56406-8
57 Perez, G. *et al.* The UCSC Genome Browser database: 2025 update. *Nucleic Acids Res*
(2024). https://doi.org/10.1093/nar/gkae974
58 Peacock, T. P. *et al.* Swine ANP32A supports avian influenza virus polymerase. *Journal*
of virology **94**, e00132-00120 (2020).
59 Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods*
14, 587-589 (2017). https://doi.org/10.1038/nmeth.4285

60   Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. *Mol Biol Evol* **37**, 1530-1534 (2020).
https://doi.org/10.1093/molbev/msaa015
61   Mistry, B. *et al.* Elucidating the Interactions between Influenza Virus Polymerase and
Host Factor ANP32A. *J Virol* **94** (2020). https://doi.org/10.1128/jvi.01353-19
1002