High-throughput protein binder discovery by rapid in vivo selection

Matthew J. Styles,[1] Joshua A. Pixley,[1] Tongyao Wei,[1] Christopher Basile,[1] Shannon S. Lu,[1] and Bryan C. Dickinson[1,2]*

[1]*Department of Chemistry, University of Chicago, 5735 S. Ellis Ave., Chicago, IL 60637*
[2]*Chan Zuckerberg Biohub, Chicago, IL 60642*
*dickinson@uchicago.edu

Abstract

Proteins that selectively bind to a target of interest are foundational components of research pipelines[1,2], diagnostics[3], and therapeutics[4]. Current immunization-based[5,6], display-based[7-14], and computational approaches[15-17,18] for discovering binders are laborious and time-consuming – taking months or more, suffer from high false positives – necessitating extensive secondary screening, and have a high failure rate, especially for disordered proteins and other challenging target classes. Here we establish Phage-Assisted Non-Continuous Selection of Protein Binders (PANCS-binders), an *in vivo* selection platform that links the life cycle of M13 phage to target protein binding though customized proximity-dependent split RNA polymerase biosensors, allowing for complete and comprehensive high-throughput screening of billion-plus member protein variant libraries with high signal-to-noise. We showcase the utility of PANCS-Binders by screening multiple protein libraries each against a panel of 95 separate therapeutically relevant targets, thereby individually assessing over 10^{11} protein-protein interaction pairs, completed in two days. These selections yielded large, high-quality datasets and hundreds of novel binders, which we showed can be affinity matured or directly used in mammalian cells to inhibit or degrade targets. PANCS-Binders dramatically accelerates and simplifies the binder discovery process, the democratization of which will help unlock new creative potential in proteome-targeting with engineered binder-based biotechnologies.

Affinity reagents - molecules that bind to a target protein of interest – are critical as basic
research tools for measuring or tracking biomolecules[1], as probes for studying biological
regulation through induced proximity[2], as core elements of diagnostics[3], and as therapeutics,
such as neutralizing antibodies and antibody-drug conjugates[4]. However, even for very well-
studied organisms, including *homo sapiens,* antibody-based binders do not exist for many
proteins of interest and, when available, are notorious for heterogenous quality control, function,
and specificity[5].
Binders can be developed by generating antibodies to a target of interest through animal
immunization, molecular display methods, or by computational design[6]. Immunization-mediated
binder generation typically costs thousands of dollars, takes many months, and is limited to
creating antibody-based reagents[6]. *In vitro* selection approaches, such as display-based
methods (e.g. phage display, mRNA display), cell sorting methods (e.g. FACS, MACS), and
growth-based methods (e.g. bacterial-2-hybrid selections) can be used to mine high diversity
libraries of protein variants to identify binders to a target of interest[7-11]. However, these
selection-based methods take several months to complete, primarily due to high false positive
rates necessitating time-consuming secondary screening[12-14]. Finally, while rapidly improving,
computational approaches require significant computing capacities, expertise, and subsequent
display-based selections, affinity maturation, and/or screening[15,16].
Creating a novel binder to a protein generally requires months of highly specialized
work, thousands of dollars, and often results in failure. Collectively, the costs and time
associated with protein binder generation restrict production to labs with a significant focus and
expertise in these techniques, prohibiting exploratory work. In comparison, creating selective
binders to DNA or RNA is as simple as designing complementary oligos, which can be
synthesized and delivered in a matter of days for ~$10. The programmable nature of nucleic
acid binders has led to the rapid explosion of diverse CRISPR technologies and other genetic
tools. To address this critical bottleneck, we sought to develop a platform for protein binder
discovery that can find novel binders to protein targets of interest in a matter of days with high
fidelity, that has the capacity to perform multiplexed screens, and that is easy enough any lab
can do it. Such a platform would both accelerate and democratize the binder discovery process.
In this work, we establish Phage-Assisted Non-Continuous Selection of protein
Binders (PANCS-Binders; **Fig. 1a**), a viral life cycle-based selection platform that can
comprehensively screen high diversity (10^{10+}) libraries of M13 phage-encoded protein variants
and identify binders to panels of dozens or more proteins of interest in a matter of days.
PANCS-Binders uses replication-deficient phage that encode protein variant libraries tagged
with one half of a proximity-dependent split RNA polymerase ($RNAP_N$) biosensor (**Fig. 1b**)[19]. *E.*
coli host cells are engineered to express a target protein of interest tagged with the other half of
the split RNA polymerase ($RNAP_C$). Protein-protein interaction (PPI) between a phage encoded
variant and the target reconstitutes the RNA polymerase (RNAP) and triggers expression of a
required phage gene, allowing phage encoding that variant to replicate, in line with the basic
principles of PACE[20,21]. After optimization and trial selections, we demonstrated the versatility of

PANCS-Binders by performing selections on 95 different protein targets with two *de novo*

phage-encoded protein variant libraries, each encoding ~10^8 unique protein variants, thereby

completing 190 independent selections in 2 days. The hit rate of this screen was 55%, resulting

in new binders for 52 diverse targets. We scaled up our library size 100-fold (~10^{10}), which

expanded the hit rate to 72% and dramatically improved the affinity of hits from PANCS – a 40-

2000x improvement with affinities as low as 206 pM. Additionally, we showcased how hits can

be quickly affinity matured though PACE, resulting in >20x improvement in affinity (to 8.4 nM).

Finally, we demonstrated that the binders for two targets, Mdm2 and KRAS, engage their

targets in mammalian cells: our Mdm2 binders inhibit the Mdm2-p53 interaction and fusion of

our KRAS binder with an LIR motif leads to LC3B mediated degradation of endogenous

KRAS[22]. The ease-of-use, speed, and reliability of PANCS-Binders will facilitate a transition of

binder generation from an expensive specialty requiring months of work with high failure to a

laboratory tool requiring less than 2 weeks (**Fig. 1c**) and available to any researcher.

Fig. 1: <u>P</u>hage-<u>A</u>ssisted <u>N</u>on-<u>C</u>ontinuous <u>S</u>election of protein <u>B</u>inders (PANCS-Binders)

a, Schematic of PANCS-Binders process. Serial passaging of *de novo*, high diversity libraries of protein

variants encoded in phage for the rapid discovery of binding variants via conditional phage replication on

an *E. coli* selection strain. Selections include the extinction of inactive variants and enrichment of the

active variants. **b,** Schematic of PANCS-binder molecular biology. A split RNAP biosensor is used for *in*

vivo selection of binder variants. *gIII* is removed from the M13 phage and placed on the positive selection

plasmid (+AP). Phage encode the N-terminal portion of the split RNAP (RNAP$_N$) fused to a protein variant

(potential binder). The +AP encodes the target fused to the C-terminal portion of RNAP (RNAP$_C$). If the

binder variant interacts with the target, then RNAP$_N$ and RNAP$_C$ recombine and transcribe *gIII*. A

simultaneous counterselection is performed using a negative selection plasmid (-AP). The -AP encodes

an off-target protein fused to an orthogonal RNAP$_C$. If the binder variant interacts with the off-target or

RNAP$_C$, then the RNAP$_N$ and RNAP$_C$ recombine and transcribe *gIII*$_{neg}$ – a dominant negative variant of

gIII that poisons phage amplification by preventing release of phage from the *E. coli* host. **c,** Timeline for

2-week binder discovery using PANCS-Binders: clone the desired target(s) into a +AP(s) and construct

the selection strain, 4-6 passages over 2-3 day of PANCS, a plaque assay or qPCR to assess endpoint

titer, and then subcloning and sequencing to validate specific enriched variants.

Optimizing Binder-PANCS

Recently, we established a split RNAP-based PPI-PACE platform for reprogramming the

binding specificity of proteins[23] (**Fig. 1b**), which we demonstrated could swap the binding

specificity of BCL2 and MCL1 using continuous evolution. In general, PACE has been shown to

be powerful for altering or tuning existing functions of molecules[24-27], primarily from initial

variants with minimal or closely related function, rather than *de novo* discovery of function. We

aimed to adapt the components of our PPI-PACE platform for the use of mining high diversity

libraries for *de novo* discovery of binders. To accomplish this, we cloned a phage-encoded,

$RNAP_N$-tagged 10^8 unique variant affibody library (**Supplementary Fig. 1**)[28]. We then

performed PACE with this library on two targets, the RAS binding domain of RAF (RAF) and

IFNG (see **Table S3** for target details). Both evolutions went extinct (**Supplementary Fig. 2**).

Prior efforts have established that PACE can enrich active phage from pools of inactive phage

(1:1000 active-to-inactive ratio)[21,27]; however, *de novo* libraries are likely to have an active-to-

inactive ratio closer to $1:10^{7+}$. To assess if the PACE evolution process itself led to phage

extinction (as opposed to no binders being present in our library), we constructed a mock library

selection system that included known, active variants. We performed PACS (Phage-Assisted

Continuous Selection[21]), PACE without the mutagenesis plasmid, using KRAS as a protein

target (+AP) and a mock library of containing a mixture of active phage encoding RAF that binds

KRAS and inactive phage encoding an affitin, evolved to bind SasA, that does not bind KRAS

(**Fig. 2a**)[29]. From these mock selections, we found that PPI-PACS could successfully enrich

active phage from mock libraries of $1:10^5$ (active:inactive phage; **Supplementary Figure 3**), but

failed to do so at any lower ratio ($1:10^{6-9}$). This indicates that continuous selection does not

sample every variant in the mock library in the initial infection step. PANCE, non-continuous

passaging, has frequently been used as a less stringent version of PACE, and we suspected

that part of this lower stringency could be due to a higher percentage of phage that infect cells

prior to being washed away in the continuous flow versus in passaging[30,31]. We hypothesized

that by extending the incubation time of phage with selection cells, we could more completely

sample every variant in our *de novo* library, and therefore succeed in *de novo* selections.

To test this hypothesis, we optimized a non-continuous selection procedure. First, we

established 6 hours as a minimum time for incubating phage and cells to obtain nearly complete

infection of our phage sample by monitoring the rate at which phage infect our cells

(**Supplementary Fig. 4**); 12 hour incubations were chosen for convenience. To determine how

quickly active phage would enrich and how quickly inactive phage would de-enrich, we

measured the rate of amplification for active (RAF variants with known affinities) and inactive

phage (Affitin (SasA)) in a KRAS selection strain (**Fig. 2a, b**). The amplification rates spanned 8

orders of magnitude: 10^6 for high affinity WT RAF, 10^{1-2} for low affinity RAF mutants, and 10^{-1-2}

for non-binders (**Fig. 2c**).

Based on the replication rates, we predicted that serial passaging with 5% of phage

transferred between passages would result in selective enrichment of the high affinity WT RAF

from 10 phage to >10^9 phage in just 2 passages and the complete de-enrichment of the inactive

phage from 10^9 to 0 in just 4 passages (**Supplementary Fig. 5**). We tested this prediction by

passaging mock libraries of 10 phage of each RAF variant spiked into 10^{10} inactive Affitin

(SasA) phage (**Supplementary Fig. 2d**). As expected, over 2 days, the high affinity WT RAF

variant enriched (a >10^{15}-fold relative enrichment) during the four-passage selection; all weaker

binders and inactive phage went extinct (**Fig. 2e**). We performed additional mock PANCS to

understand the effects of several variables on this relative enrichment rate: +AP selection

stringency (**Supplementary Fig. 6),** -AP selection stringency (**Supplementary Fig. 7**), transfer

rate (**Supplementary Fig. 8**), and initial cell-to-phage ratio (**Supplementary Fig. 9**). Finally, we

tested mock selections using several published binder-target pairs using our optimized AP

strengths, transfer rate, and initial cell-to-phage ratios (**Supplementary Fig. 10**). Collectively,

these mock selections indicate that this new system, which we named P̲hage-A̲ssisted N̲on-

C̲ontinuous S̲election of Protein B̲inders (PANCS-Binders), can perform *de novo* selections of

up to 10^{10+} variant libraries (above the typical 10^{9-10} *E. coli* transformation limit) in 2 days, using

simple serial phage out-growths in culture tubes or even 96-well plates. Therefore, we next

performed pilot selections with a *de novo* phage-encoded binder library to demonstrate that

PANCS-Binders can be used to discover novel binders.

Fig. 2: Development of PANCS-Binders for selection from *de novo*-like mock libraries.

a, The selection system for mock libraries consists of an *E. coli* selection strain with KRAS4b (WT)-

$RNAP_{C,CGG}$ as the positive selection target (+AP) and ZB_{neg}-$RNAP_{C,T7}$ as the counter selection target (-

AP). The mock library consisted of a mixture of two selection plasmids (SP): an active phage with

$RNAP_N$-RAF(RBD) and an inactive phage with $RNAP_N$-Affitin (SasA). **b,** Phage amplification assay: 1000

PFU of each phage are incubated with 1 mL of a selection strain for 12 hours and then the titer is

determined: amplification rate is output titer/input titer. **c,** Amplification rate for RAF variants and the non-

binding affitin (SasA) phage on the KRAS selection strain. Published affinities (K_d) are listed above each

RAF variant amplification rate[32]. Each amplification rate was obtained in duplicate and error bars indicate

SD. The green line indicates an amplification rate of 20 (no enrichment if passaging at 5%) and the red

line indicates an amplification rate of 1 (no amplification). **d,** Four passage PANCS starting from a mock

library (10 PFU active phage with 10^{10} PFU inactive phage (affitin (SasA)) in 5 mL KRAS4b (WT)

selection strain (**Fig. 2a**) with a 12 hr passage outgrowth and 5% transfer of supernatant phage into fresh

cells to seed each passage. **e,** Titers at the end of each passage (passage 0 indicates the initial titer to

start PANCS). The limit of detection (LOD) in our plaque assays is $5*10^2$ PFU/mL (1 PFU); if a titer had 0

PFU, it was set to 0.2 PFU for calculation purposes (Affitin (SasA) in C). Each phage was passaged in

triplicate and error bars indicate SD.

Pilot *de novo* library PANCS-Binders

We selected six protein targets to attempt to discover novel binders for, each with

varying architectures and degrees of structural order: KRAS4b(G12D), RAF (RBD), Mdm2 (1-

188), IFNG, Myc DNA binding domain (DBD), and Sos1 disordered domain (**Fig. 3a**,

Supplementary Table 3). We simply cloned each target into the +AP as a RNAP$_C$ fusion and

transformed into *E. coli* host cells with the ZB$_{neg}$ -AP used in our final mock selections

(**Supplementary Fig. 10**) to prepare the selection materials. We passaged the 10^8 affibody

library, which had gone extinct in PACE-based selections, on each *E. coli* selection strain in

culture tubes for 12-hour outgrowths and 5% transfer between passages (**Fig. 3a** and

Supplementary Fig. 1). After 4 rounds of passaging (48 hours total), we measured the titer in

each condition. KRAS(G12D), RAF (RBD), IFNG, and Mdm2 had high titers ($>10^8$), indicating

successful selections, while Sos1 and Myc (DBD) selections had titers near the limit of detection

($<10^5$), indicating failure to enrich binders (**Supplementary Table 4**).

We performed next generation sequencing (NGS) on the four selections with a high titer

(**Fig. 3b**), which revealed that the selections on KRAS(G12D), RAF (RBD), and IFNG each

converged onto a single sequence (i.e., >80% of reads belonged to that sequence). The most

dominant Mdm2 variant comprised only 2.6% of the population; in retrospect, this lack of

convergence is unsurprising as Mdm2 binds an FXXXWF/Y motif common to ~3% of library

variants, and therefore, many variants were enriched. For KRAS G12D, RAF (RBD), and IFNG

selections, we also sequenced the library, passage 2, and passage 3, which revealed that the

relative ratio between active variants is set by passage 2 (**Supplementary Fig. 11**), in line with

our Mock PANCS results. Finally, we used AlphaFold3 to predict the binding interface for each

hit, which, as expected, showed the randomized region of the affibody at the predicted

interaction interface (**Fig. 3a** and **Supplementary Fig. 12**).

Fig. 3: PANCS-Binders to discover novel binders from *de novo* libraries.

a, Cloning a target +AP panel, transformation of selection strain, parallel 4-passage PANCS of each

selection strain with a 10^8 variant affibody library. Titer assessed after the fourth passage indicated which

selections enriched binders (indicated as AlphaFold predictions of the binder-target pair (if titer was high)

or target alone (if titer was low), see **Supplementary Table 4** for titers). The affibody sequence from

passage 4 phage was then PCR amplified for NGS, subcloned into a luciferase assay system (shown

bottom right), and subcloned into pET vectors for protein purification for SPR. Binding validation

luciferase assay (bottom right): the target is fused to the $RNAP_C$ on an expression plasmid, the binder is

fused to the $RNAP_N$ on a separate expression plasmid, and a reporter plasmid where LuxAB expression

is determined by PPI dependent recombination of the spRNAP. **b,** Top variant amino acid sequences

identified from the NGS (top) and the percentage of reads for each unique variant in the NGS of passage

4 (bottom left) for each successful selection. Variants shown in color with the associated sequence were

examined further in the luciferase and SPR assays. The second S-WYS variant for Mdm2 isolated for

testing in lux (pink) did not have a single read in the NGS but is indicated as having a single read for

plotting on a log scale. Fold-change in luciferase signal (bottom right) for select variants from the *de novo*

screens over the template affibody used in cloning the library (a non-binder), and when available,

previously published binders to these targets (NS1 Monobody (70 nM K_d[33]), Z(RAF322) Affibody (190 nM

K_d[34]), Cl2-based binder (K_d not determined[35]). Each condition has four independent data points, and error

bars indicate SD. The *in vitro* binding affinity, K_d, is reported as text for select variants (as measured by

SPR, see **Supplementary Fig. 15**). **c,** Percentage of reads for each unique variant in the original PANCS

compared to the percentage of reads for each unique variant in a biological replicate of the PANCS at

passage 4 (comparison of parallel replicates in **Supplementary Fig. 17**). If a variant was present in one

NGS sample but not in another, it was coded as 0.01%. of reads.

We subcloned the top variants from each selection (those >1% of reads in KRAS G12D,

RAF (RBD), and IFNG selections and 4 random variants from the Mdm2 selection) into an

expression plasmid (Lux-N) for measuring binding in a previously established *E. coli* luciferase

assay (**Fig. 3a**)[19,21,23]. Reconstitution of the proximity dependent split RNAP, measured by the

production of luminescence, was observed to be induced following co-expression of each

affibody variant and its respective selection target. This indicated variant:target binding in *E. coli*

(**Fig. 3b**). Positive binding controls, previously published binders discovered by ribosome

display (Z(RAF322)[34], phage display (NS1 Monobody)[33], and rational engineering (Cl2 12.1A)[35],

produced comparable signal to the newly selected affibodies for RAF (RBD), KRAS G12D, and

Mdm2, respectively. We confirmed specificity of binding by assaying for binding between one

binder from each successful selection with the four targets that gave hits (**Supplementary Fig.**

13), which revealed high specificity of each binder (only S-VVD had off-target binding to Mdm2).

We purified each of the top binders (**Supplementary Fig. 14**) and performed surface-plasmon

resonance (SPR) binding assays, revealing binding between top variants and their target of

interest with *in vitro* affinities between 176 and 635 nM (**Fig. 3b** and **Supplementary Fig. 15**).

These results confirmed that the selections successfully enriched binder variants.

To assess reproducibility of the selections, we repeated the entire 6-target selection four

additional times in parallel several months later. This yielded highly consistent results in terms of

the extinction events and endpoint phage titers (**Supplementary Fig. 16**). We performed NGS

on each of the replicate selections and observed high reproducibility (r = 0.72; average of each

pairwise Pearson's Correlation for variants >0.1% of NGS reads; variants enriched >5% were

identical) between biological replicates (**Fig. 3c)** and within parallel replicates **(Supplementary**

Fig. 17; r = 0.95). These results demonstrate that PANCS-Binders can rapidly - in just 48 hours,

comprehensively, and reproducibly screen and isolate binder variants from *de novo* libraries

without the need for replicates or additional screening.

High-Throughput PANCS

We next sought to challenge the PANCS-Binders technology in a multiplexed high-

throughput selection by attempting to simultaneously identify binders for a large panel of diverse

protein targets in 96-well plate format (**Fig. 4a**). In addition to scaling down the selection

volumes to 1 mL for plate compatibility, we made three additional adjustments for this selection:

we extended the linker length between the target and $RNAP_C$ to ensure that the position and

orientation of the binder was not constrained (**Supplementary Fig. 18**), we reduced the

selection stringency from 5% to 10% transfer, and we created a second ~10^8 phage library

based on an affitin scaffold to have two different scaffold libraries to compare to one another

(**Supplementary Fig. 19**). We simply cloned each protein of interest (a total of 95 targets) into

+APs without additional optimization, as well as a negative control no-fusion +AP consisting of a

start codon followed by the 60-amino acid GS linker and RNAPc, to establish a 96-well plate of

target selection strains. The 95 targets (**Supplementary Table 3**) vary in origin (mammalian,

bacterial, or viral), localization (secreted, extracellular domains of membrane proteins,

membrane proteins, cytosolic, or nuclear), function, and structure (fully ordered, significant

disordered regions, fully disordered). We reasoned that such a diverse target panel should

assess the capability for performing PANCS-Binders in a high-throughput manner and provide a

realistic estimate of the expected hit rate for 10^8 libraries. To confirm that the selection +APs

were functional, we performed an amplification assay (**Supplementary Fig. 20**) with phage

encoding $RNAP_N$ WT, which is not proximity dependent (recombines with $RNAP_C$ independent

of target and binder interacting), with the -AP (demonstrating sufficient counterselection to

prevent non-selective replication) and without the -AP (indicating sufficient target expression for

binding induced replication; only four targets did not amplify >10-fold). Notably, this panel has

many targets that are difficult to purify from *E. coli*, which highlights the superior properties of

targets expression plasmids over *in vitro* selection methods.

Fig. 4: High-throughput PANCS-Binders for 95 targets.

a, Cloning a 96-panel set of target selection strains, 96-deep well plate-based parallel PANCS-binder

selection of two 10^8 libraries (affibody (**Supplementary Fig. 1**) and affitin (**Supplementary Fig. 18**) using

binding assays and qPCR to measure the endpoint titer in a high-throughput manner. **b,** *E. coli* spRNAP

complementation luciferase assay heatmap (**Fig. 3a**) on all preliminary hit variants (titer > 10^7 PFU/mL

when top variant is full-length protein (see Note 1 in supporting information and **Supplementary Fig. 23**

for individual plots). Fold/change in binding >20 is set equal to 20; selections which were not preliminary

hits are indicated in grey. **c,** Fold-change in luciferase signal for selection of 15 variants from the *de novo*

screens assessed across 15 targets and additional controls to assess selectivity. Within each target (y-

axis) each binder is normalized to the signal for two non-binders (set to 1): an affibody that was evolved

to bind PD-L1[36] and an affitin that was evolved to bind SasA[29]. Only one off-diagonal had a fold-change

>2.

After preparing the selection cells, we performed the 4 passage PANCS-Binders over

the course of 48 hours and collected endpoint titers using qPCR to identify preliminary hits –

titers >10^7 PFU/mL (**Supplementary Fig. 21** and **Supplementary Table 5;** see

Supplementary Note 1 for a detailed discussion of how we selected this threshold including

Supplementary Figs. 22-26). For all preliminary hit wells, we collected NGS, performed

AlphaFold2 multimer predictions for the top variants (NGS and AlphaFold compiled in

Supplementary Fig. 22), and subcloned variants from passage 4 phage and collected

luciferase binding assay for the top variant(s) cloned (**Supplementary Fig. 23**). Overall, we

validated 79 new binders to 52 targets (**Fig. 4b**). These results further demonstrate the high

correlation between endpoint titer and binder enrichment. We used 16 of these binders to

investigate the specificity of our selected binders (**Fig. 4c**), identifying only one off-target

interaction between GABARAP and our LC3B binder which is not surprising given the high

sequence similarity between GABARAP and LC3B (both of which bind LIR domains). With this

dataset of 288 pairwise binding measurements, we sought to evaluate the ability of

AlphaFold3[37] to identify binding vs non-binding pairs; iPTM values were not well correlated with

binding (**Supplementary Fig. 27**).

Improving affinity and hit rate in PANCS-Binders

While the initial 55% hit rate and 100s of nM K_d showcase the viability of PANCS-

Binders, we wanted to assess whether we could improve the hit rate and identify higher affinity

binders by simply using larger libraries in PANCS-Binders. We cloned 10^{10} variant libraries for

both the affibody and affitin scaffolds using parallel transformations (**Fig. 5a**) – a 100-fold

improvement of our initial library size (**Supplementary Table 6**). We selected 8 targets that did

not give hits in our initial screens, including VHL, TRIM, and PNCA, and 4 targets that

previously generated hits, including KRAS G12D and RAF, and performed a large library

PANCS-Binders screen (**Fig. 5a**) with 6 rounds of passaging over 72 hours (see Methods for

additional details and protocol differences). In addition to getting new hits for the targets that

also initially gave hits with smaller libraries, 3 out of the 8 previously failed targets now also

yielded hits, as confirmed by the *E. coli* luciferase assay (38%; **Fig. 5a**, **Supplementary Table**

7). We analyzed these hits by NGS (**Supplementary Fig. 28**) and compared the new hits for

KRAS G12D both using the *E. coli* luciferase assay and *in vitro* (**Fig. 5b,** binders for other

targets shown in **Supplementary Figs. 29** and **31**): the affinity of the best binder obtained from

this selection improved from 384 nM to 0.2 nM, representing a ~2000x improvement in affinity.

These results confirm that PANCS-Binders is capable of quickly mining 10^{10} variant libraries for

novel binders, improving our hit rate (predicted from 55% to 72%), and identifying higher affinity

binders, all within a 72-hour selection.

One advantage of the PANCS-Binders technology is that the same selection strains and
phage can quickly be adapted to PACE by transforming the selection strain with a mutagenesis
plasmid to allow mutations to accrue during a directed evolution campaign. To test this, we
performed PACE-based directed evolution on the passage 4 phage from our initial Mdm2-
affibody selection (**Fig. 3b**) using adaptations of our previous method[23]. First, we identified
higher stringency positive APs with reduced propagation of passage 4 phage (as measured by
activity dependent plaque assays). Then we used these two strains to perform PACE over the
course of 60 hours of evolution, after which the phage populations converged (6/8 subclones
sequenced) on an affibody variant Y-WCT with an additional A46T mutation and then an
additional L51F mutation (in 1/8 with A46T; **Fig. 5c**). Assessment using the *E. coli* luciferase
binding assays showed the evolved variants have improved affinity for their targets (**Fig. 5d**).
We confirmed this *in vitro*: K_d = 8.4 nM for A46T, and 26 nM for A46T/L51F (**Fig. 5d** and
Supplementary Fig. 31) compared with 176 nM for the highest affinity Mdm2 binder identified
from the initial PANCS (a >20x improvement). Critically, the mutations arising from the
supplemental PACE are not in the initial randomization sites, nor are they predicted to make
direct contacts with the target protein (**Fig. 5c**). This is not altogether surprising; the power of
unbiased directed evolution via PACE for optimizing existing function though non-intuitive
mutations is well-established.
19

Fig. 5. Larger libraries, affinity maturation, and mammalian applications of PANCS-binder hits.

a, Large, 10^{10}, libraries were prepared using parallel transformations (**Supplementary Table 6**) that were

pooled to initiate large library PANCS-binder (**Supplementary Table 7**), which required starting at an

initial volume of 500 mL and decreasing the volume of each passage slowly as the selection progressed

(**Supplementary Fig. 28** for NGS results). *E. coli* spRNAP complementation luciferase assay heatmap

shown for successful selections, with slashes indicating selections that also gave hits in the 95 panel

selections (**Fig. 4**). **b,** Luciferase assay results for large KRas G12D selection (**Supplementary Fig. 29**

for others) with *in vitro* affinity listed above each bar, comparing literature reference binder in grey,

original binder from smaller scale selection in red, and new binder in purple (**Supplementary Fig. 30** for

SPR). **c,** Affinity maturation of Mdm2 Affibody hit by PACE. Passage 4 from the Mdm2 selection with the

10^8 affibody library (**Fig. 3b**) was used to seed PACE. PACE was initiated on a +AP selection strain with

lower Mdm2 expression than the original PANCS. After 24 hours, there was a 12-hour mixing step with an

even lower expression level strain for an additional 36 hours (total of 60 hours). Titers were monitored by

activity independent plaque assay, and variants were subcloned at 60 hours (isolated variants shown

here). **d,** Variants were tested in the luciferase binding assay alongside T-WDN and the non-binding

Affibody (PDL1) as in **Fig. 3n**. Affinities measured by SPR (**Supplementary Fig. 28**) are shown above

each luciferase bar graph. **e,** Initial KRAS G12D (**Fig. 3b**) and high affinity KRAS G12D (**Fig. 5b**) binders

were tested for binding in mammalian cells using a split-Nano-luciferase assay in HEK293T cells.

Transfections were performed in triplicate; error bars indicate SD. **f,** Schematic showing LC3B recruiting

element (LIR motif) fused to binders for targeted degradation by autophagy.[38] **g,** Representative WB of

endogenous KRAS degradation by Ab N-LHY – LIR (**Supplementary Fig. 33** for full blots and replicates).

h, Quantification of replicate degradation results as shown in **g** (**Supplementary Fig. 33**). **i,** Mdm2/Mdm2

binder co-localization: mCherry-tagged Mdm2 and GFP-tagged Mdm2 binder co-localize in the nucleus,

while a control GFP does not (replicates in **Figure S34**). **j,** Representative WB of Mdm2-p53 PPI

inhibition; inhibition of this PPI activates p53 transcription resulting in increased expression of p21 and

Mdm2 (**Supplementary Fig. 35** for full blots and replicates). Quantification of replicate WBs for Mdm2-

p53 PPI inhibition in U2OS cells: **k,** Mdm2 expression and **l,** p21 expression (**Supplementary Fig. 35**).

For Western blot quantification, statistical analyses were performed using one-way ANOVA with Dunnett's

multiple comparison test binder vs. control. *$P < 0.05$; **$P < 0.01$.

Binders Function in Mammalian Cells

Lastly, we sought to test whether the novel binders could be taken directly from PANCS-

Binders into mammalian cells and bind their targets in a functionally relevant manner. We

cloned the KRAS G12D and KRAS binders Ab-N-VCD (**Fig. 3b**) and Ab-N-LHY (**Fig. 5b**) into a

split nano-luciferase complementation assay[39] and demonstrated robust binding in HEK293T

cells (**Fig. 5e**). Next, we sought to convert Ab-N-LHY into a KRAS degrader by fusing it to an

LIR (LC3B interacting region) domain for targeted protein degradation through authophagy[22]

(**Fig. 5f**). We observed robust degradation of endogenous KRAS in U2OS as assessed by WB

in a binder0dependent manner (**Fig. 5g** and **Supplementary Fig. 33**). We then demonstrate

that the high affinity Mdm2 binder (Ab-Y-WCT A46T) co-localizes with Mdm2 in the nucleus

(**Fig. h** and **Supplementary Fig. 34**), confirming binding in mammalian cells. We then

overexpressed AB-Y-WCT A46T and several other Mdm2 binders in U2OS cells to see if the

binders could inhibit the MDM2-p53 interaction, by monitoring expression levels of Mdm2 and

p21, which are both transcriptionally regulated by p53. We observed a strong induction of both

targets upon MDM2 binder expression, which indicates robust inhibition of Mdm2 and activation

of p53 (**Fig. 5i** and **Supplementary Fig. 35**). These results demonstrate that PANCS-Binders

can produce binder variants with functional binding activity in mammalian cells.

Discussion

PANCS-Binders is a rapid, reproducible, and reliable method for discovering protein

binders. The entire process of cloning a target into the +AP/RNAP$_C$ expression plasmids,

selection, assessment, and secondary validation assays can routinely be performed in 2 weeks

without the requirement for highly specialized expertise or equipment. The speed of PANCS-

Binders comes from its strong de-enrichment of weak and non-binders and high enrichment of

binders, which abrogates the need for secondary screening campaigns common in display-

based selection techniques. We propose that two fundamental aspects of such techniques limit

the relative enrichment of binding variants over non-binding variants commonly achieved in

display methods: threshold selection (bound or not bound) and activity-independent

amplification. PANCS-Binders utilizes a split RNAP-based biosensor with a large dynamic range

that links the degree of variant function to the degree of phage replication for that variant, both

creating a gradient rather than threshold selection and removing the activity-independent

amplification step. Notably, neither Alphafold2 Multimer (**Supplementary Fig. 24**) nor

AlphaFold3 (**Supplementary Fig. 27** and **32**) could predict binding (false negatives) or non-

binding (false positives) from our screen, illustrating the enduring importance and value of real

experimentation and limitations still inherent in computational modeling. Furthermore, the high-

quality binding data generated through PANCS can be used in improving computational

modeling and AI-based design techniques.

The ability to utilize high diversity libraries in phage-assisted selections and evolutions is

a powerful tool in the directed evolution arsenal and should expand the range of evolutions

possible. PACE and PANCE have been applied to alter or tune the specificity of a variety of

protein functions[8,19-21,23-25,27,31,40]; however, because mutations accumulate incrementally, these

campaigns require a nearly continuous evolutionary pathway starting from low or non-functional

initial variants. In a recent tour de force, the evolution of a protein that binds a small molecule-

protein complex, an elaborate pathway was needed to access the 5 mutations needed for

minimal function and the 8 mutations eventually reached for high function. This included

repeated high mutagenesis drift periods, a 3-position randomized library, steppingstone states

(a panel of 16 small molecules), and testing of a wide range of selection stringencies (PACE

and PANCE across 26 different stringency selection plasmids). High diversity libraries, like

those used here in PANCS-Binders, are prepared using *in vitro* diversification techniques

capable of making tens of targeted mutations in the initial variant. PANCS is a powerful

approach to jumpstart more difficult evolutionary campaigns by increasing the navigable

distance between functional states.

PANCS-Binders can screen multiplexed libraries of 10^{10} phage-encoded variants across

dozens of targets in 2-3 days, yielding high affinity, selective binders with sufficient fidelity such

that hits can be directly used in secondary assays, such as mammalian cell experiments. In our

96-well based high-throughput PANCS-Binders with 10^8 variant libraries, we achieved a 55% hit

rate across a range of targets with a high correlation between endpoint titer and validated

binding (79/92). Repeating a subset of failed selections with a 100-fold larger library produced

hits for 38% of those initial failures, showcasing how simply scaling up library size can yield hits

for otherwise challenging targets. Additionally, either with large 10^{10+} libraries or through rapid

affinity maturation via PACE, high affinity binders (<10 nM) can be obtained quickly.

PANCS-Binders generated hits for disordered protein targets and proteins that are

challenging or impossible to purify, showcasing the potential of this screening and selection

platform to discover binders for proteins that lack structural data or are incompatible with *in vitro*

selection strategies. PANCS-Binders did not require significant optimization of selection

conditions, as is commonly the case for 2-hybrid selections; while target solubility does impact

the target-RNAP$_C$ expression level, the differences were small enough to abrogate the need for

target dependent tuning of expression. In other words, with PANCS-Binders, one can "plug-and-

play" targets into the system.

Methods

Cloning and Bacterial Strain Handling

All plasmids and phage (**Supplementary Table 1)** were cloned by Gibson Assembly (GA) of PCR

fragments generated using Q5 DNA polymerase (NEB). All primers (**Supplementary Table 2**) were

ordered from IDT. For plasmids, GA mixtures were transformed into chemically competent DH10β *E. coli*

and after a 1 h outgrowth in 2xYT media, were plated on antibiotic selective agar plates to isolate

individual clones. For phage, GA mixtures were transformed into chemically competent S1030-1059 *E.*

coli, and after a 2 h outgrowth, a plaque assay was performed to isolate individual phage clones. All

plasmids and phage were confirmed by Sanger sequencing. All plasmid maps with annotations of key

features are available in **Supplementary Table 1**. For constructing selection (+AP/-AP) and *E. coli*

luciferase (2-22/N-lux/C-lux) strains, S1030 *E. coli* was made chemically competent and then single or

double transformations were used (and then repeated as needed until all plasmids were incorporated). *E.*

coli strains was grown on agar plates static at 37 °C or in solution at 37 °C with 200 rpm shaking with

Luria Broth (LB) supplemented with the appropriate antibiotic unless otherwise indicated. Antibiotics were

used at standard concentrations: kanamycin (40 ug/mL), chloramphenicol (33 ug/mL), and carbenicillin

(100 ug/mL).

Plaque Assays

Activity independent plaque assays can be used to determine the phage titer via plaque counting. Activity

dependent plaque assays can be used to check for robust phage replication on a given strain. For activity

independent plaque assays, an S1030-1059 *E. coli* culture (1059 plasmid encodes *gIII* expressed from

the phage shock promoter to produce *gIII* after phage infection), is grown to stationary phase in LB with

carbenicillin, subcultured 1/10 in fresh LB with antibiotic to an OD600 of 0.4-0.6, and then used as the

selection strain in the plaque assay. Similarly, for activity dependent strains, S1030 with a +AP (and -AP)

were grown similarly for use in the plaque assay. For the plaque assay, an initial dilution of the stock can

be added based on the expected titer, but generally, 2 μL of a phage stock (or diluted stock) is added to

100 μL of subculture, mixed, and then serially diluted (2 μL into 100 μL) to create 4 dilutions. 750 μL of 50

°C top agar (7 g/L agar, 25 g/L LB) was added to each dilution and then transferred in its entirety to one

quadrant of a bottom agar plate (15 g/L agar, 15 g/L LB). After 10-16 h of incubation at 37 °C, plaques

become visible and were counted in the quadrant with 10-200 plaque forming units (PFU).

Phage Amplification Rates

To determine phage amplification rate, the titer of a phage stock is determined using an activity

independent plaque assay. Based on this titer, a diluted stock is made that should be 500 PFU/μL (the

titer of this diluted stock is confirmed using an activity independent plaque assay). The activity dependent

strain (+AP/-AP) is grown to stationary phase in LB with carbenicillin and kanamycin, subcultured 1/10 in

fresh LB with antibiotic to an OD600 of 0.4-0.6. 2 μL, 1000 PFU, are added to 1 mL of this subculture and

then incubated at 37 °C with shaking for 12 h. The cells are then pelleted and the cell-free supernatant

collected for use in an activity independent plaque assay to determine the titer at the end of the

amplification assay. The endpoint titer is divided by the starting titer (1000 PFU/mL) to determine the

amplification rate.

PACS/PACE

General procedures for continuous flow experiments. PACS[21] and PACE[19] were performed as previously

described. All tubing, chemostat bottles, and lagoon flasks were bleached thoroughly, rinsed with DI

water, and then autoclaved to ensure sterility. 10 L carboys of Davis Rich media were prepared as

described previously[19]. Inlet lines consist of short needles unable to reach the culture and outlet lines

consist of long needles able to reach the culture. Each chemostat had an inlet line for fresh media, an

inlet line with a sterile filter for airflow, an outlet line for waste, and an outlet line for each lagoon. Each

lagoon had an inlet line from the chemostat, an inlet line with a sterile filter for airflow, and an outlet line

for waste. For PACE lagoons, each lagoon also has an inlet line for arabinose. Each chemostat and

lagoon has a magnetic stir bar. Colonies of the selection strain were used to inoculate a 5 mL culture in

the relevant media (see below) and grown to stationary phase. This culture was then used to inoculate a

200 mL chemostat (250 mL bottle). This culture was stirred in a 37 °C cabinet until an OD600 of ~0.5 and

then fresh Davis Rich media was flowed into the chemostat at ~1 vol/h. The chemostat was monitored for

4 h to ensure that the flow rate maintained a stable OD600 of ~0.5. Phage was then added to each

lagoon and then culture was flowed into the lagoon to a volume of 20-25 mL and let incubate for 1 h prior

to beginning flow of 1 vol/h. Samples from the lagoons were collected from the waste lines at various

timepoints.

PACE with libraries and with PANCS output: PACE was performed as describe previously[19] in line with

standard PACE protocols[40]. For PACE with RAF and IFNG with the affibody library (Figure S2), two

selection strength strains were used for 36 h each with a 12 h mixing step (60 h total). +APs (ori,

gIII/RNAP$_C$ RBS strength): p15 SD8/SD8 to pSC101 SD8/SD8 for RAF and from pSC101 SD8/SD8 to

p15 SD8/sd5 for IFNG. In addition to the +AP, each strain had a ZB$_{neg}$ -AP (20-1) and MP6 (see Table

S1). The initial selection strain supported activity dependent plaques of the affibody binders isolated from

PANCS of the affibody library, confirming that binders capable of propagating on the initial selection strain

exist in the library. One lagoon was used for each chemostat, initially seeded with 10^{10} PFU of library

phage and arabinose began flowing during the 1 h of incubation of phage with culture in the lagoon prior

to beginning flow at 1 vol/h. Samples were collected prior to beginning the mixed strain phase (24 h) and

after completion of the second strain (60 h) and titers were assessed by activity independent plaque

assay. For the PACE with Mdm2 (Figure 5A) starting from the final passage of PANCS with the Affibody

library (Figure 3A), the same protocol was followed with the selection strengths of the +AP being pSC101

SD8/SD8 to p15 SD8/sd5 (both strains produced small activity dependent plaques using the passage 4

phage and large activity dependent plaques at the 60 h timepoint of PACE).

PACS with mock libraries. PACS was performed as describe previously[21]. The selection strain (S1030/31-
69/20-6) was prepared by double transformation into chemically competent *E. coli*. Mock libraries were
composed of 10^{10} PFU inactive phage (Affitin (SasA)) and varying amounts (0 (negative control), 10, 10^2,
10^3, 10^4, or 10^5 PFU) of active phage (RAF WT). In addition, a lagoon was seeded with only 10^3 active
phage as a positive control. Each of these was done in duplicate lagoons/chemostats. Samples were
taken at 12, 24, and 48 h and the titer was determined by activity independent plaque assay (Figure S3).

PANCS

General protocol for PANCS. For each passage, a selection strain (+AP/-AP) is grown to stationary phase
in LB with carbenicillin and kanamycin, subcultured 1/10 in fresh LB with carbenicillin and kanamycin to
an OD600 of 0.4-0.6 prior to adding phage. For passage 1, stock phage are added to the subculture and
incubated at 37 °C with shaking (200 rpm) for 12 h, then centrifuged to pellet the cells and collect the cell-
free supernatant (referred to as passage 1 phage). For subsequent passages, some fraction of the cell-
free supernatant from the prior passage is added to the subculture and incubated at 37 °C with shaking
(200 rpm) for 12 h, then centrifuged to pellet the cells and collect the cell-free supernatant (referred to as
passage # phage). Titers of each passage or just the final passage were then determined using activity
independent plaque assays, or for the 96-target panel, using qPCR (see below).

Details for specific PANCS. For PANCS development, a variety of culture volumes, transfer rates, and
number of passages were used. For the Mock PANCS (**Fig. 2e**) and the 6-target panel PANCS (**Fig. 3a,**
Supplementary Table 4), we performed 4-passage PANCS with 5 mL cultures, initially seeded with 10^{10}
PFU for passage 1, and seeded with 250 µL of prior passage for passages 2-4 (5% transfer). For the 96-
target panel PANCS (**Fig. 4a, Supplementary Table 5**), we performed 4-passage PANCS with 1 mL
cultures (2 mL deep 96-well plates), initially seeded with $2*10^9$ PFU for passage 1, and seeded with 100
µL of prior passage for passages 2-4 (10% transfer). For additional passaging of this PANCS, we did 2%
transfer for two passages (**Supplementary Fig. 26**). For the 10^{10} library PANCS (**Fig. 5a**), 6 passage
PANCS was performed using either a 2% (RAF, IFNG, KRAS G12D, and HRAS) or 5% (PCNA, ALKBH5,
PINK, PIK3CA, TRIM21, VHL, NIX, and BAX) transfer rate between passages and passage 1 was
seeded with $5*10^{10}$ PFU; however, unlike previous PANCS, the volume of each passage was changed as
well: 500 mL for passage 1, 125 mL for passage 2, 25 mL for passage 3, and 5 mL for passage 4-6.

Split-RNAP *E. coli* Luciferase Assays

We followed a slightly modified version of our previously reported assay[19]. For each target, a two-plasmid
strain (S1030/2-22/C-lux) was made chemically competent and each binder and non-binder N-Lux
plasmid was transformed to make the three-plasmid luciferase strain. Colonies were picked for each
binder and non-binder for each strain to inoculate 1 mL of LB with kanamycin, chloramphenicol, and
carbenicillin and grown at 37 °C with shaking (200 rpm) for 12-16 h. Strains were then subcultured 7.5 µL
into 143 µL of LB with kanamycin, chloramphenicol, carbenicillin, and L-arabinose (2 mg/mL final

concentration) in white side, clear bottom 96-well assay plates (Corning 3610) and incubated at 37 °C

with shaking (200 rpm) for 3.5 h prior to reading the OD600 and luminescence signal on a BioTek

Synergy Neo2 plate reader. Luminescence signal is first divided by the OD600 to normalize luminescence

to cell growth. Then the luminescence/OD600 is normalized for all strains with the same C-Lux plasmid

(target-RNAP$_C$ expression plasmid) are divided by the non-binder signal (non-binders set equal to 1). Due

to differences in expression levels of each target and differences in how binding impacts expression level

of a target, we do not believe direct comparisons in fold change over non-binder can be made across

different targets, and therefore, we plot all binders for an RNAP$_C$ expression plasmid separately from

other RNAP$_C$ expression plasmids.

NGS of PANCS Hits

We used the Amplicon EZ service provided by Genewiz (Azenta) for Illumina sequencing of each of our

hits ([GENEWIZ from Azenta | Amplicon-EZ](#)) which provides 50,000+ paired end reads per sample. We

used primers to install Illumina partial adaptors (red/purple) and barcodes (blue) to PCR products

extending from the linker to after the stop codon of our scaffold in phage (primed with green regions) –

see **Supplementary Table 2**. PCR was performed with Q5 DNAP polymerase (NEB) directly from phage

(1 µL) in a 25 µL PCR reaction; the initial denaturation step was 10 min at 98 °C to release the ssDNA

from the phage particle, a 63 °C Ta, and a 40 second extension time were used with 30 cycles. PCR

products are confirmed by gel (5 µL), and the remaining 20 µL of PCR product is pooled with other

barcoded PCR products and column purified (Zymo DCC5). Qubit dsDNA High Sensitivity kit is used to

determine an accurate concentration of the sample prior to dilution and submission for sequencing. For

the NGS data in **Fig. 3, Supplementary Figs. 1, 11** and **17**, we used 7 barcodes/sequencing sample

yielding >15,000 reads per condition (library or PANCS passage). For the NGS data in **Supplementary**

Figs. 22, 25, and **28**, we used 24 barcodes/sequencing sample yielding >1000 reads for nearly all

PANCS samples (reads listed in tables for each library-target pairing in each figure). BB Merge was used

to merge each paired end reads ([https://jgi.doe.gov/data-and-tools/software-tools/bbtools/bb-tools-user-](#)

[guide/bbmerge-guide/](#)) and then MatLab was used to separate reads by barcode and to translate reads

using modified scripts as described previously (MatLab scripts provided as a supplement)[41].

Alpha fold predictions

As a preliminary estimate of how our binders interact with their target, we used AlphaFold2 multimer

collab ([AlphaFold2.ipynb - Colab (google.com)](#))[42,43] for predicting the interaction between binder and

target for the top 4 variants above 1% of reads (**Supplementary Fig. 22**). AlphaFold3 was released after

this analysis, and we subsequently used AlphaFold3 ([https://golgi.sandbox.google.com/](#))[37] to predict

binding interactions for each top variants from our 6-target panel PANCS (**Supplementary Fig. 12**), our

10[10] library PANCS (**Supplementary Fig. 28**), and for all of the binder-target pairs examined in **Fig. 4c**

(**Supplementary Fig. 27**). We implore readers to utilize these predictions only for hypothesis generation
rather than as data indicative of an actual interaction.

qPCR to estimate phage titers

qPCR was tested across several primers that prime to M13 phage genes for linear response of a phage
serial dilution. Primers VC-525 and VC-526 were chosen (**Supplementary Table 2**). Power Up SYBR mix
was used with the following PCR protocol: 10 minutes at 95 °C (to denature phage particle and release
ssDNA); 40 cycles of 20 seconds at 95 °C, 20 seconds at 60 °C, and 20 seconds at 72 °C; then 10
seconds at 95 °C and 60 seconds at 65 °C. qPCR was run on a QuantStudio6Pro. Each run includes a
standard curve for which the titer is assessed using activity independent plaque assay.

Library Construction

General Protocol: Libraries were designed based on previously published randomizations[28,44].
Randomization was installed into a template phage using primers with degenerate codons (IDT; see
Supplementary Table 2). We optimized each step of this protocol to maximize the number of clones
obtained. PCR conditions were optimized for each library to produce robust PCR product at 25 cycles and
then tested for production at lower cycles to reduce amplification bias (18 or fewer cycles were used for
each library reported here). PCR was then scaled up to produce 20-100 μg of PCR product. PCR
products were concentrated using the Wizard Kit (Promega) and then digested using DpnI and NheI-HF
(NEB) using a multidose cycle: for ~20-50 μg of PCR product in 300-400 μL, and then digested with DpnI
and NheI, purified using a Zymo Gel Extraction kit, and then ligated with T4 DNA Ligase (NEB). Ligated
products were then electroporated into 1059 *E. coli* cells. The cells were then recovered in 50 mL of 37
°C SOC media and incubated for 2 h at 37 °C with shaking – samples were collected throughout this time
for determining the titer by plaque assay. At 2 h, the cells were pelleted and the cell-free supernatant was
collected. 1030-1059 (activity independent replication strain) was grown overnight and then subcultured
1:10 to and OD600 of 0.6 at 37 °C with shaking (200 rpm). The phage (cell-free supernatant) was then
amplified by adding the phage to this subculture for 8-10 h. At the conclusion of this outgrowth, cells were
pelleted and the cell-free supernatant was sterile filtered to create the final library stock (titer determined
by activity independent plaque assay).

Affibody Library: The affibody library (**Supplementary Fig. 1**) was cloned from the Affibody (PDL1)
phage (**Supplementary Table 1**) using MS-783 and MS-618 (**Supplementary Table 2**) by Q5 DNAP
(NEB) with a T_a of 68 °C. For generating the 10^8 size library, the *E. coli* strain used for the electroporation
was SS320 (a highly electrocompetent strain that is capable of phage replication) rather than 10β. The
10^8 library size was generated with a single transformation of 3 μg ligation product. For the 10^{10} library
size, eight transformations of 10 μg ligation product were performed (**Supplementary Table 6**).

Affitin Library: The affitin library (**Supplementary Fig. 19**) was cloned from either Affitin (SasA), 10^{10}

library, or a version of the Affitin (SasA) phage with three stop codons inserted into a region randomized

by the primers (**Supplementary Table 1**), 10^8 library, using MS-624 and MS-799 primers

(**Supplementary Table 2**) by Q5 DNAP with GC enhancer with a T_a of 68 °C. Both the 10^8 and 10^{10}

libraries were generated following the general protocol with a single 4 μg and eight 8 μg transformations,

respectively, (**Supplementary Table 6**).

Protein Purification

General Protocol for Target Proteins: Each target protein (KRAS G12D (1-169), RAF, IFNG, and Mdm2

was cloned into a pET28 vector with a C-terminal 6xHis tag and transformed into BL21 *E. coli*

(**Supplementary Table 1**). Cells were grown to an OD600 of 0.8 (37 °C with shaking), chilled on ice,

induced with 1 mM IPTG, and then incubated with shaking at 16 °C overnight. Cells were pelleted and

resuspended in a lysis buffer (25 mM Tris (pH 7.8), 10% glycerol, 200 mM NaCl). Prior to lysing by

sonication, cells were treated with PMSF. The soluble fraction of the lysate was incubated with Ni^{2+} resin,

washed with lysis buffer containing 50 mM imidazole, then eluted in lysis buffer containing 250 mM

imidazole, and finally buffer exchanged into lysis buffer and concentrated.

General Protocol for Binder Proteins: Each binder variant was cloned into a pET30 vector with an N-

terminal 3xFLAG and GST tag and transformed into BL21 *E. coli* (**Supplementary Table 1**). Cells were

grown to an OD600 of 0.8 (37 °C with shaking), chilled on ice, induced with 1 mM IPTG, and then

incubated with shaking at 16 °C overnight. Cells were pelleted and resuspended in a lysis buffer (25 mM

Tris (pH 7.8), 10% glycerol, 100 mM NaCl). The soluble fraction of the lysate was incubated with GST

resin, washed with lysis buffer, then eluted in lysis buffer containing 10 mM L-glutathione, and finally

buffer exchanged into lysis buffer and concentrated. Purified binders shown in **Supplementary Fig. 14**.

Surface Plasmon Resonance

Surface Plasmon Resonance was performed on a Biacore 8000 using a NTA chip for immobilizing the

His-tagged target proteins. Target concentrations were optimized to elicit a response of ~50-100 RU (180

s of 5 uL/s) and then a range of binder concentrations were tested to identify concentrations that

produced robust binding (90 s of 30 uL/s). All SPR conducted at 10 °C to maintain slow dissociation of the

His-tagged immobilized protein. All dose-responses were fit to a kinetic model for 1:1 binding using the

Biacore evaluation software – all fits passed the quality checks in this software (**Supplementary Table 8**).

Split Nano-Luciferase Assay

62.5 ng of the N-terminus of Nano-Luciferase-binder fusion plasmid and 62.5 ng of the KRas(G12D)-C-

terminus of Nano-Luciferase fusion plasmid were co-transfected into HEK293T cells using 500 ng of PEI

in 96-well glass bottom plate (Cellvis, P96-1-N). Transfection was performed in triplicate. After 36 hours,

the Nano-luciferase activity was measured using Nano-Glo® Live Cell Assay System (Promega, N2011).

Endogenous KRAS Degradation Assay

5   1000 ng of binder-LIR fusion plasmids were transfected into U2OS cells by 0.3 uL of Lipofectamin 3000 in

6   a 24-well plate. After 4 h, the media was replaced. After 48 h, the cells were collected and subjected to

7   western blot analysis with the appropriate antibodies.

Mdm2 binder-Mdm2 Co-Localization Assay

10   125 ng of the GFP-binder fusion plasmid and 125 ng of the mCherry-Mdm2 fusion plasmid were co-

11   transfected into HEK293T cells using 0.075 µL of Xfect™ Transfection Reagent (Takara Bio, 631317) in

12   96-well glass bottom plate (Cellvis, P96-1-N). After 4 h, the media was replaced. After 36 h, cells were

13   imaged with a Leica fluorescence microscope.

Mdm2-p53 Inhibition Assay

16   1000 ng of Mdm2 binder plasmids were transfected into U2OS cells by 0.3 uL of Xfect™ Transfection

17   Reagent (Takara Bio, 631317) in a 24-well plate. After 4 h, the media was replaced. After 48 h, the cells

18   were collected and subjected to western blot analysis with the appropriate antibodies.

Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary

linked to this article.

Data Availability

Links to electronic vector maps are included in Supplementary Information. Key vectors will be

deposited with Addgene and all physical vectors will be made available upon reasonable request.

Source data are provided with paper.

References

Bandrowski, A., Pairish, M., Eckmann, P., Grethe, J. & Martone, M. E. The Antibody Registry: ten years of registering antibodies. *Nucleic Acids Res* **51**, D358-D367 (2023). https://doi.org:10.1093/nar/gkac927

Stanton, B. Z., Chory, E. J. & Crabtree, G. R. Chemically induced proximity in biology and medicine. *Science* **359** (2018). https://doi.org:10.1126/science.aao5902

Park, M. Surface Display Technology for Biosensor Applications: A Review. *Sensors (Basel)* **20** (2020). https://doi.org:10.3390/s20102775

Carter, P. J. & Lazar, G. A. Next generation antibody drugs: pursuit of the 'high-hanging fruit'. *Nat Rev Drug Discov* **17**, 197-223 (2018). https://doi.org:10.1038/nrd.2017.227

Ayoubi, R. *et al.* Scaling of an antibody validation procedure enables quantification of antibody performance in major research applications. *Elife* **12** (2023). https://doi.org:10.7554/eLife.91645

Laustsen, A. H., Greiff, V., Karatt-Vellatt, A., Muyldermans, S. & Jenkins, T. P. Animal Immunization, in Vitro Display Technologies, and Machine Learning for Antibody Discovery. *Trends Biotechnol* **39**, 1263-1273 (2021). https://doi.org:10.1016/j.tibtech.2021.03.003

Sidhu, S. S., Lowman, H. B., Cunningham, B. C. & Wells, J. A. Phage display for selection of novel binding peptides. *Methods Enzymol* **328**, 333-363 (2000). https://doi.org:10.1016/s0076-6879(00)28406-1

Xie, V. C., Styles, M. J. & Dickinson, B. C. Methods for the directed evolution of biomolecular interactions. *Trends Biochem Sci* **47**, 403-416 (2022). https://doi.org:10.1016/j.tibs.2022.01.001

Wellner, A. *et al.* Rapid generation of potent antibodies by autonomous hypermutation in yeast. *Nat Chem Biol* **17**, 1057-1064 (2021). https://doi.org:10.1038/s41589-021-00832-4

Philpott, D. N. *et al.* Rapid On-Cell Selection of High-Performance Human Antibodies. *ACS Cent Sci* **8**, 102-109 (2022). https://doi.org:10.1021/acscentsci.1c01205

Lopez-Morales, J. *et al.* Protein Engineering and High-Throughput Screening by Yeast Surface Display: Survey of Current Methods. *Small Science* **3** (2023). https://doi.org:10.1002/smsc.202300095

Porebski, B. T. *et al.* Rapid discovery of high-affinity antibodies via massively parallel sequencing, ribosome display and affinity screening. *Nat Biomed Eng* **8**, 214-232 (2024). https://doi.org:10.1038/s41551-023-01093-3

McConnell, A., Batten, S. L. & Hackel, B. J. Determinants of Developability and Evolvability of Synthetic Miniproteins as Ligand Scaffolds. *J Mol Biol* **435**, 168339 (2023). https://doi.org:10.1016/j.jmb.2023.168339

Kordon, S. P. *et al.* Isoform- and ligand-specific modulation of the adhesion GPCR ADGRL3/Latrophilin3 by a synthetic binder. *Nat Commun* **14**, 635 (2023). https://doi.org:10.1038/s41467-023-36312-7

Cao, L. *et al.* Design of protein-binding proteins from the target structure alone. *Nature* **605**, 551-560 (2022). https://doi.org:10.1038/s41586-022-04654-9

Bennett, N. R. *et al.* Atomically accurate de novo design of single-domain antibodies. *bioRxiv* (2024). https://doi.org:10.1101/2024.03.14.585103

Sappington, I. *et al.* Improved protein binder design using beta-pairing targeted RFdiffusion. *bioRxiv* (2024). https://doi.org:10.1101/2024.10.11.617496

Huang, B. *et al.* Designed endocytosis-inducing proteins degrade targets and amplify signals. *Nature* (2024). https://doi.org:10.1038/s41586-024-07948-2

Pu, J., Zinkus-Boltz, J. & Dickinson, B. C. Evolution of a split RNA polymerase as a versatile biosensor platform. *Nat Chem Biol* **13**, 432-438 (2017). https://doi.org:10.1038/nchembio.2299

Esvelt, K. M., Carlson, J. C. & Liu, D. R. A system for the continuous directed evolution of biomolecules. *Nature* **472**, 499-503 (2011). https://doi.org:10.1038/nature09929

Zinkus-Boltz, J., DeValk, C. & Dickinson, B. C. A Phage-Assisted Continuous Selection Approach for Deep Mutational Scanning of Protein-Protein Interactions. *ACS Chem Biol* **14**, 2757-2767 (2019). https://doi.org:10.1021/acschembio.9b00669

He, H., Zhou, C. & Chen, X. ATNC: Versatile Nanobody Chimeras for Autophagic Degradation of Intracellular Unligandable and Undruggable Proteins. *J Am Chem Soc* **145**, 24785-24795 (2023). https://doi.org:10.1021/jacs.3c08843

Xie, V. C., Pu, J., Metzger, B. P., Thornton, J. W. & Dickinson, B. C. Contingency and chance erase necessity in the experimental evolution of ancestral proteins. *Elife* **10** (2021). https://doi.org:10.7554/eLife.67336

Miller, S. M. *et al.* Continuous evolution of SpCas9 variants compatible with non-G PAMs. *Nat Biotechnol* **38**, 471-481 (2020). https://doi.org:10.1038/s41587-020-0412-8

Hubbard, B. P. *et al.* Continuous directed evolution of DNA-binding proteins to improve TALEN specificity. *Nat Methods* **12**, 939-942 (2015). https://doi.org:10.1038/nmeth.3515

Packer, M. S., Rees, H. A. & Liu, D. R. Phage-assisted continuous evolution of proteases with altered substrate specificity. *Nat Commun* **8**, 956 (2017). https://doi.org:10.1038/s41467-017-01055-9

Thuronyi, B. W. *et al.* Continuous evolution of base editors with expanded target compatibility and improved activity. *Nat Biotechnol* **37**, 1070-1079 (2019). https://doi.org:10.1038/s41587-019-0193-0

Woldring, D. R., Holec, P. V., Stern, L. A., Du, Y. & Hackel, B. J. A Gradient of Sitewise Diversity Promotes Evolutionary Fitness for Binder Discovery in a Three-Helix Bundle Protein Scaffold. *Biochemistry* **56**, 1656-1671 (2017). https://doi.org:10.1021/acs.biochem.6b01142

Behar, G. *et al.* Whole-bacterium ribosome display selection for isolation of highly specific anti-Staphyloccocus aureus Affitins for detection- and capture-based biomedical applications. *Biotechnol Bioeng* **116**, 1844-1855 (2019). https://doi.org:10.1002/bit.26989

Hu, J. H. *et al.* Evolved Cas9 variants with broad PAM compatibility and high DNA specificity. *Nature* **556**, 57-63 (2018). https://doi.org:10.1038/nature26155

Dewey, J. A., Azizi, S. A., Lu, V. & Dickinson, B. C. A System for the Evolution of Protein-Protein Interaction Inducers. *ACS Synth Biol* **10**, 2096-2110 (2021). https://doi.org:10.1021/acssynbio.1c00276

Block, C., Janknecht, R., Herrmann, C., Nassar, N. & Wittinghofer, A. Quantitative structure-activity analysis correlating Ras/Raf interaction in vitro to Raf activation in vivo. *Nat Struct Biol* **3**, 244-251 (1996). https://doi.org:10.1038/nsb0396-244

Spencer-Smith, R. *et al.* Inhibition of RAS function through targeting an allosteric regulatory site. *Nat Chem Biol* **13**, 62-68 (2017). https://doi.org:10.1038/nchembio.2231

Grimm, S., Salahshour, S. & Nygren, P. A. Monitored whole gene in vitro evolution of an anti-hRaf-1 affibody molecule towards increased binding affinity. *N Biotechnol* **29**, 534-542 (2012). https://doi.org:10.1016/j.nbt.2011.10.008

Karlsson, G. B. *et al.* Activation of p53 by scaffold-stabilised expression of Mdm2-binding peptides: visualisation of reporter gene induction at the single-cell level. *Br J Cancer* **91**, 1488-1494 (2004). https://doi.org:10.1038/sj.bjc.6602143

Jing, L. *et al.* Screening and production of an affibody inhibiting the interaction of the PD-1/PD-L1 immune checkpoint. *Protein Expr Purif* **166**, 105520 (2020). https://doi.org:10.1016/j.pep.2019.105520

Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **630**, 493-500 (2024). https://doi.org:10.1038/s41586-024-07487-w

He, H., Zhou, C. & Chen, X. ATNC: Versatile Nanobody Chimeras for Autophagic Degradation of Intracellular Unligandable and Undruggable Proteins. *Journal of the American Chemical Society* **145**, 24785-24795 (2023). https://doi.org:10.1021/jacs.3c08843

Dixon, A. S. *et al.* NanoLuc Complementation Reporter Optimized for Accurate Measurement of Protein Interactions in Cells. *ACS Chem Biol* **11**, 400-408 (2016). https://doi.org:10.1021/acschembio.5b00753

Miller, S. M., Wang, T. & Liu, D. R. Phage-assisted continuous and non-continuous evolution. *Nat Protoc* **15**, 4101-4127 (2020). https://doi.org:10.1038/s41596-020-00410-3

Rentero Rebollo, I., Sabisz, M., Baeriswyl, V. & Heinis, C. Identification of target-binding peptide motifs by high-throughput sequencing of phage-selected peptides. *Nucleic Acids Res* **42**, e169 (2014). https://doi.org:10.1093/nar/gku940

Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679-682 (2022). https://doi.org:10.1038/s41592-022-01488-1

Evans, R. *et al.* (2022). https://doi.org:10.1101/2021.10.04.463034

Mouratou, B. *et al.* Remodeling a DNA-binding protein as a specific in vivo inhibitor of bacterial secretin PulD. *Proc Natl Acad Sci U S A* **104**, 17983-17988 (2007). https://doi.org:10.1073/pnas.0702963104

Acknowledgements

This work was supported by the National Institute of General Medical Sciences (GM119840 to B.C.D and F32GM147968 to M.J.S) and then National Cancer Institute (P30CA014599) of the National Institutes of Health, and by the Camille and Henry Dreyfus Foundation Teacher Scholar Award (B.C.D.). We thank S. Ahmadiantehrani for assistance with preparing this paper.

Contributions: Conceptualization: M.J.S. and B.C.D. Methodology: M.J.S. Investigation: M.S., J.A.P.; T.W.; C.B.; and S.L. Writing – original draft: M.J.S., J.A.P., and B.C.D. Writing – reviewing and editing: T.W. Supervision: B.C.D.

Ethics Declaration

Competing interests

B.C.D. is an inventor on the patent describing the split RNAP biosensors. The University of Chicago has filed a provisional patent on the PANCS-Binders technology with M.J.S and B.C.D. listed as inventors.

SUPPORTING INFORMATION

Details of bacterial strains, plasmids, primers, additional data figures and tables.