Title: Maximizing Nature-based Solutions using Artificial Intelligence to align global biodiversity, climate, and water targets.

Camilo Alejo[1,2,3,*], Amy Luers[2,3,4], Andréa Ventimiglia[2,3], H. Damon Matthews[1,2,3,**]

[1] Department of Geography, Planning and Environment, Concordia University, Montreal, QC, Canada.

[2] Sustainability in the Digital Age, Concordia University, Montreal, QC, Canada.

[3] Future Earth Canada Hub, Montreal, QC, Canada.

[4] Microsoft Corporation, Seattle, WA, USA

*Correspondence: camilo.alejo.monroy@futureearth.org

**Correspondence: damon.matthews@concordia.ca

Summary

Nature-based Solutions (NbS) can represent holistic pathways to reach sustainable biodiversity, climate, and water outcomes through conservation and restoration. Yet, existing prioritization frameworks rarely identify holistic NbS that maximize these outcomes. Here, we present an AI-driven framework that prioritizes NbS locations, maximizing biodiversity protection with ecological intactness, carbon storage, and water surface stability. We implement this framework through scenarios combining specific biodiversity and co-benefits outcomes to achieve Canada's 30x30 conservation and restoration targets. Our results reveal inevitable environmental trade-offs and critical implications for equitable NbS and extractive activities. To achieve conservation targets, protecting threatened biodiversity and irrecoverable carbon storage in forests would effectively address trade-offs and enhance Protected Areas outcomes. However, minimizing trade-offs to achieve restoration targets will require targeted interventions in existing forested and agricultural lands. These findings demonstrate that frameworks integrating AI and biodiversity-co-benefit scenarios reveal strategic land use planning, policies, and investments for holistic NbS.

Keywords

Biodiversity, carbon stocks, surface water, conservation, restoration, nature-based solutions, spatial prioritization, reinforcement learning, artificial intelligence, Canada

Introduction

Global biodiversity loss is closely intertwined with climate change, water scarcity, and social inequality. Given these interconnections, many global targets emphasize the

need for holistic pathways that simultaneously address environmental and social
challenges. Centered around biodiversity, the Kunming-Montreal Framework aims to
restore 30% of degraded land and establish 30% of area-based conservation by 2030
(i.e., 30x30 targets), ensuring enhanced biodiversity, ecosystems' integrity, and related
co-benefits [1]. Since the establishment of the Paris Agreement [2], reducing land use
emissions while conserving and restoring ecosystems' carbon sinks have become
relevant components of Nationally Determined Contributions (NDCs) to reach net-zero
emissions by 2050. Similarly, the Sustainable Development Goals (SDGs) set targets to
protect and restore ecosystems, including water-related ecosystems such as forests,
wetlands, rivers, and lakes [3].
48
Nature-based Solutions (NbS), understood as actions to protect, restore, or sustainably
manage ecosystems (natural or modified) to address societal challenges [4], may play a
critical role in reaching global targets. NbS encompass a wide set of interventions [5,6],
many of which provide holistic pathways for countries to reach the 30x30 conservation
and restoration targets [1] in synergy with climate change mitigation [7], water management
[8], and Sustainable Development Goals [9]. Reaching these global targets through NbS
implies strategically aligning biodiversity outcomes with related co-benefits emerging
from ecological integrity, carbon storage, and water. This alignment will necessarily
invoke trade-offs and NbS monitoring frameworks that can identify optimal conservation
and restoration priorities.
59
Over the past decade, analytical tools for NbS implementation have become
increasingly available. State-of-the-art monitoring tools typically leverage field
observations, remote sensing, and machine learning to predict spatial-temporal patterns
of biodiversity [10], carbon stocks [11,12], and water quality [13]. In parallel with this increased
monitoring capacity, other approaches identify optimal NbS priorities for conservation
and restoration actions. Usually, conservation priority frameworks have weighted the
importance of biodiversity aspects such as irreplaceability (e.g., species and habitats
rarity) and vulnerability through a reactive (e.g., threatened species, habitat loss) or
proactive (low vulnerability and low cost) approach [14,15]. From an ecosystem services
perspective, Mitchell et al.[16] instead prioritized conservation co-benefits emerging from
carbon storage, water supply, and tourism. Others have aimed to inform conservation
targets by simultaneously prioritizing biodiversity outcomes with climate suitability [17–19],
carbon storage [20], or multiple co-benefits simultaneously [21,22].
73
The identification of optimal NbS has not been restricted to conservation targets. For
instance, biodiversity and carbon outcomes have been ranked and prioritized on lands
that have been converted to pastures and croplands [23]. Vettorazzi & Valente[24] proposed
land suitability assessments for prioritizing restoration actions that could improve water
supply. A common approach between conservation [25] and restoration [26] has resulted
from prioritizing vulnerability (e.g., habitat conversion, human footprint) on global
ecoregions. However, AI's potential to find optimal solutions has only recently begun to
be explored in conservation [27] and restoration [28] priority frameworks. Furthermore, few
studies leverage AI to develop common frameworks for conservation and restoration
targets that maximize outcomes related to biodiversity as well as related co-benefits.

The complexity of optimizing biodiversity and co-benefit outcomes for conservation and
restoration targets is further challenged by the interplay of local and economic priorities.
Governments and decision-makers have dominated conservation and restoration
planning and decision-making spaces, relegating local priorities and capacities [29]. This
is the case of Indigenous Peoples, who have been historically marginalized from these
spaces [30] despite being increasingly acknowledged in global environmental targets [1–3].
Indigenous Peoples' presence in 40% of global protected areas [31] and 36% of intact
forests [32] suggest that any feasible and equitable pathway for conservation and
restoration targets requires the inclusion of Indigenous land governance.

Economic priorities also challenge the feasibility of conservation and restoration targets.
For example, some priority frameworks highlight the restoration potential of converted
lands for pasture and agriculture to reach environmental targets [23,28]. However, realizing
this potential may be unfeasible due to economic barriers [33] and may require
considering other degraded lands to reach restoration targets. Similarly, prioritizing
energy development will limit governments' pathways to achieve environmental targets
while diminishing current biodiversity and co-benefits outcomes [34–36]. Tackling this
complexity requires prioritization frameworks that explore diverse conservation and
restoration pathways and that address their local and economic impacts.

Here, we present a framework that prioritizes NbS locations maximizing biodiversity and
co-benefits outcomes to reach 30x30 conservation and restoration targets. Leveraging
Reinforcement Learning, this framework trains an agent to learn from the state of a
system (i.e., exploration) and cost-effectively select areas (i.e., exploitation) that
maximize a reward (Figure 1). A cost-effective selection follows monitoring (e.g.,
species occurrence) and prioritization (e.g., minimize species loss) policies while
avoiding a predefined cost (e.g., low ecological intactness). We test this framework in
Canada under a series of conservation and restoration scenarios that maximize the
occurrence of threatened species in synergy with co-benefits from ecological intactness,
irrecoverable carbon storage, and water surface stability. Additionally, we examine
some local and economic implications of these scenarios. Overall, aligning biodiversity
outcomes with other co-benefits across conservation and restoration scenarios results
in varied environmental trade-offs and spatial alignments with Indigenous-led NbS and
natural resource projects. To conserve 30% of lands, prioritizing the protection of
threatened biodiversity and carbon stocks in forests can effectively address these trade-
offs. Yet, minimizing trade-offs to restore 30% of degraded lands will require a
combination of scenarios and interventions in existing forested and agricultural lands.
Taken together, this prioritization framework demonstrates that integrating AI, multiple
co-benefits, and scenario analysis can reveal strategic land use planning, policies, and
investments for holistic NbS.

Results

Scenarios to conserve 30% of land by aligning biodiversity, climate, and water outcomes.

We defined conservation priority scenarios (Figure 1) based on the spatial distribution of existing Protected Areas (Figure S1), threatened species (Figure S2), ecological intactness, carbon storage, and water surface stability (Table S1). The conservation scenarios identified an additional ~17.75% of Canadian land areas to be protected. These scenarios achieve Canada's 30X30 targets by maximizing the protection of threatened biodiversity and guaranteeing at least 10% of each threatened species range, while also prioritizing co-benefits of ecological intactness (Conservation Scenario 1), irrecoverable carbon stocks (Conservation Scenario 2), and water surface stability (Conservation Scenario 3) (Figure 2).

When only ecological intactness was prioritized (Conservation Scenario 1; Figure 2), more than 60% of conservation priorities were concentrated in the Southern and Northern Artic (Quebec, Nunavut, Northwest Territories). Relative to existing Protected Areas, this scenario would result in an expected increase of ecological intactness from 0.80 to 0.91 on a normalized scale. However, it would reduce the mean number of threatened species protected (2.55 to 1.52 species/100 km^2), species richness (76.49 to 40.58 species/100 km2), carbon stocks (441.42 to 145.43 tC/ha), and absolute water surface change (2.30%/100 km^2 in 2000-2015 vs 1984-1999) (Figure 3).

Conservation Scenario 2 (prioritizing irrecoverable carbon stocks; Figure 2) led to conservation priorities that are concentrated in the forest-dominated Boreal ecozones (> 60%) and the Montane Cordillera (~14%) throughout southern and western provinces, respectively. In contrast to Scenario 1, aligning biodiversity and climate change mitigation targets led to lower ecological intactness (0.74) but consistently higher number of protected threatened species (3.42 species/100km^2), species richness (95 species/100km^2), water surface change (0.67%), and carbon stocks (631.34 tC/ha), compared to Protected Areas (Figure 3).

Conservation Scenario 3 that prioritized water surface stability (Figure 2), had a dispersed geographic distribution, partially overlapping with Conservation Scenarios 1 and 2. However, some priorities were particularly aggregated in the eastern Taiga Shield (Northern Quebec and Newfoundland and Labrador), Hudson Plains (Quebec and Ontario), and Taiga Plains (British Columbia), known to be partially dominated by peatlands. Except for an overall reduction of water surface change, Scenario 3 exhibited similar outcomes to existing Protected Areas.

After overlapping the results from all three conservation scenarios (Figure 2), we were able to identify the areas of land that were prioritized in two or more of the individual scenarios. Of the target 17.75% of land area prioritized by the three scenarios, 11.4% appeared in two scenarios and 0.89% in all three scenarios, suggesting significant potential for achieving multiple co-benefits. The conservation scenarios overlaps also reveal a reduced portion of priorities in strategic ecozones with substantial outcomes on biodiversity, ecological intactness, carbon stocks, and water stability co-benefits. For

example, compared to existing Protected Areas, the overlapped priorities in the
Mixewood Plains (0.01%), Pacific Maritime (0.1%), and Hudson Plains (13.2%) (Figure
S3) would simultaneously increase biodiversity outcomes, ecological intactness, and
carbon stocks (Table S2). The overlaps also highlight three conservation priorities
clusters that could potentially realize biodiversity and co-benefits outcomes. These
clusters are the ecozone transition of the Montane Cordillera, Boreal Plains, and Taiga
Plains (British Columbia); the southern Hudson Bay and James Bay Lowlands (i.e.,
Hudson Plains, Québec, Ontario); and Lac Temiscamingue Lowlands (i.e., Boreal
Shield, Québec, Ontario). Overall, these Conservation Scenarios imply that aligning
biodiversity outcomes with other co-benefits results in varied spatial patterns of
conservation priorities and trade-offs between co-benefits. Despite these trade-offs,
maximizing the protection of threatened biodiversity and irrecoverable carbon stocks
(Conservation Scenario 2) represents a comprehensive pathway to improve
biodiversity, climate change mitigation, and water stability outcomes in existing
Protected Areas.

188
189 **Figure 1. Flowchart summarizing the steps to identify Conservation and**
190 **Restoration priorities to reach 30X30 targets aligned with co-benefits, using**
191 **Reinforcement Learning.**

192
193

194
195 **Figure 2. The spatial distribution of Conservation Priority Scenarios to reach**
196 **30X30 targets obtained through Reinforcement Learning**. All Scenarios assume a
197 baseline extent of Protected Areas of 12.25% and a 17.75% remaining area budget to
198 maximize the occurrence of threatened species with co-benefits emerging from
199 Ecological Intactness (A), Irrecoverable Carbon Stocks (B), Water Surface Stability (C).
200 Priority Areas had a 10km resolution. The Scenarios Overlaps (D) suggest areas that
201 simultaneously maximize the occurrence of threatened species with multiple co-
202 benefits.
203
204

205
Figure 3. The distribution of threatened species, species richness, ecological
intactness, carbon stocks, and absolute surface water change in existing
Protected Areas and Conservation Priority Scenarios. The conservation scenarios
aim to maximize the occurrence of threatened species with co-benefits emerging from
Ecological Intactness, Irrecoverable Carbon Stocks, and Water Surface Stability. The
Scenarios Overlaps represent the co-occurrence of conservation priorities in at least
two scenarios. The points and bars refer to the mean and standard deviation.
213

214 Scenarios to restore 30% of degraded land by aligning biodiversity, climate, and
215 water outcomes.
The restoration scenarios suggested priorities in 30% of degraded land (i.e., ~3.5% total
surface) that would maximize the occurrence of threatened biodiversity in areas
experiencing reduced ecological intactness (Restoration Scenario 1), carbon stocks
losses (Restoration Scenario 2), and water surface instability (Restoration Scenario 3)
(Figure 4). Restoration Scenario 1 priorities were mostly concentrated in the Prairies
(~72.8%) and Mixed Wood Plains (~12.4%) that are dominated by agriculture. Among
all Restoration Scenarios, this scenario was particularly effective for protecting
threatened biodiversity (~4) and species richness in general (108) and the least
effective for carbon storage (147.48 tC/ha) (Figure 5). When threatened biodiversity was
prioritized with irrecoverable carbon stocks losses, a considerable part of the priorities
occurred in the Boreal Shield (31.8%), Montane Cordillera (22.2%), and Boreal Plains
(21.3%), largely dominated by forests (Figure 4). Although Restoration Scenario 2
would result in lower biodiversity outcomes than Restoration Scenario 1, prioritizing the
recovery of carbon stock losses exhibit the most effective outcomes for ecological
intactness (0.70), carbon stocks (513.94 tC/ha), and water surface stability (1.2%)
(Figure 5). The restoration scenario 3 displayed a wide geographical range of priorities
across contrasting ecozones (Figure 4, Figure S3). However, we did not identify wide-
scale improvements in multiple outcomes after prioritizing actions for threatened
biodiversity and water surface instability.
235
The overlaps among Restoration Scenarios represented approximately a third (1.28%)
of the desired land target (3.5%). Specifically, 0.05% simultaneously aligned reduced
ecological intactness, carbon storage, and water co-benefits, and 1.23% aligned two of
these co-benefits. Consequently, we did not identify wide potential or spatial clusters
(Figure 4), requiring coordinated actions to improve biodiversity and all co-benefits
(Figure 5). Nevertheless, some overlapped restoration priorities that moderately
clustered across the western Boreal Shield and eastern Montane Cordillera (British
Columbia) (Figure 4) suggest moderate effects on threatened species (~3 species/100
km^2) and species richness (90-110 species/100 km^2) if aligned with the restoration of
ecological intactness and carbon stocks (Figure 5, Table S3). Restoring these co-
benefits would potentially result in larger biodiversity outcomes in the Mixedwood Plains
from Southern Ontario, which also displayed a cluster of overlapped priorities. In the
Prairies (Alberta, Saskatchewan, Manitoba), a cluster of overlapped priorities suggested
that aligning restoration actions that manage and adapt to water surface instability and
improve ecological intactness could positively impact threatened species (~3) and
several species in general (91). These spatial patterns imply that each restoration
scenario would potentially improve conditions for threatened biodiversity in tandem with
one or multiple co-benefits in limited spatial extents, suggesting diverse and context-
dependent restoration pathways.
255

Figure 4. The spatial distribution of Restoration Priority Scenarios to reach 30X30 targets obtained through Reinforcement Learning. All Scenarios assume a baseline extent of degraded land of 11.7% and an area budget of 3.5% to maximize the occurrence of threatened species while prioritizing the occurrence of low ecological intactness (A), irrecoverable carbon stocks losses (B), and water surface instability (C). Priority Areas had a 10km resolution. The Restoration Scenarios overlaps (D) suggest areas that maximize the occurrence of threatened species and are currently experiencing reduced ecological intactness, carbon stocks losses, and water surface instability.

267
Figure 5. The distribution of threatened species, species richness, ecological intactness, carbon stocks, and absolute surface water change intensity in restoration priority scenarios to reach 30X30 targets. All restoration scenarios aim to maximize the occurrence of threatened species while prioritizing the occurrence of low ecological intactness, irrecoverable carbon stocks losses, and water surface instability. The Scenarios Overlaps represent the co-occurrence of conservation or restoration priorities in at least two scenarios. The points and bars refer to the mean and standard deviation.

276

Conservation and restoration priorities are near Indigenous-led NbS and major natural resource projects.

279   To identify the local implications of the priority scenarios, we quantified the overlap and
280   distance of Indigenous-led NbS programs and other Indigenous Lands to conservation
281   and restoration priorities (Figure 6, Table S4). Indigenous-led NbS included Indigenous
282   Protected Conserved Areas and the Indigenous Guardians Federal Program. These
283   federally supported initiatives displayed the highest overlap (47%) and closest proximity
284   (130 km) to conservation priorities maximizing Irrecoverable Carbon stocks (Scenario
285   2). Conversely, priorities maximizing Ecological Intactness (Scenario 1) showed minimal
286   overlap (8%) and the greatest distance (~180 km). Similarly, Restoration priorities
287   focusing on Carbon Stocks recovery (Restoration Scenario 2) demonstrated the highest
288   overlap (17%) and shortest distance (~150 km) to Indigenous-led NbS. Notably, these
289   carbon stocks focused Conservation and Restoration priorities, displaying more
290   proximity to these government-funded initiatives, mainly occurred in the Pacific and
291   Atlantic Maritimes, the Montane Cordillera and Hudson Plains.

292

293   Compared to Indigenous-led NbS, other Indigenous Lands without federal government
294   funding were considerably closer to Conservation and Restoration Priorities. The
295   overlap of Indigenous Lands with Conservation priorities ranged between ~34%
296   (Conservation Scenario 2: Irrecoverable Carbon Stocks) and ~3% (Conservation
297   Scenario 1: Ecological Intactness) and an average distance range of 48 km to 160 km
298   (Figure 5). For Restoration priorities, other Indigenous Lands showed 12-23% overlap
299   and averaged ~80 km in distance. These spatial patterns imply that a significant fraction
300   of government-funded Indigenous-led NbS overlap or are near conservation and
301   restoration priorities, especially those that align with climate change mitigation.
302   Additionally, both conservation and restoration priorities, which displayed disparate
303   spatial patterns across ecozones, share a wide alignment with Indigenous Lands that
304   lack government funding.

305

Figure 6. The distribution of conservation and restoration priorities distance to Indigenous-led NbS and other Indigenous Lands. Indigenous-led NbS corresponded to government-recognized Indigenous Protected Conserved Areas and the Indigenous Guardians Federal Program. Other Indigenous Lands corresponded to areas that are not federally funded in Indigenous-led NbS programs. The points and bars refer to the mean and standard deviation.

Finally, we assessed the spatial distribution of conservation and restoration priority scenarios in relation to major forestry, mining, and energy sector projects under construction or planned in the next ten years (Figure 7). The average proximity of forestry projects to restoration priorities (~370-380 km) tended to be higher than conservation priorities (~340-880 km). Compared to forestry, conservation priorities were, on average, closer (140-450 km) to energy projects (i.e., solar, hydro, and oil), particularly when ecological intactness, carbon stocks, and water stability co-benefits were overlapped (i.e., Scenarios Overlaps). Notably, conservation priorities were very close to energy projects in the Boreal Plains, Montane Cordillera and Pacific Maritimes. Moreover, the average distance of all restoration priorities to energy projects was the lowest (83-120 km) among all scenarios and sector comparisons. In fact, 25% of the restoration scenarios that prioritized low ecological intactness (Restoration Scenario 1), carbon stock losses (Restoration Scenario 2), and scenarios overlaps were in the 30 km of an energy project. Relative to energy projects, restoration priorities shared ecozone locations with conservation priorities but were additionally identified in the Mixewood Plains and Prairies. Furthermore, energy linear infrastructure, in particular, displayed the most significant overlap (1-4%) with both conservation and restoration priorities (Table S5).

The distance of mining projects to conservation and restoration priorities tended to be higher than energy projects but lower than those corresponding to forestry (Figure 7). For example, the average conservation priority that maximized the protection of threatened biodiversity and ecological intactness (Conservation Scenario 1) was ~130

km from mining projects, and 25% of these priorities were in a ~80 km range.
Restoration priorities across the different scenarios were in a similar average range to
mining projects (~160 km). Moreover, the spatial distribution of both conservation and
restoration priorities to mining was relatively similar to energy projects across Boreal
ecozones and the Prairies. According to these spatial distributions, restoration priorities
across all scenarios are more likely to be close to major natural resource projects than
conservation priorities, especially in the energy sector. Nevertheless, some
conservation priorities that maximize ecological intactness (Conservation Scenario 1),
irrecoverable carbon stocks (Conservation Scenario 2), and overlapped co-benefits
exhibit a marked immediacy to major energy and mining projects.
347

348
349   **Figure 7. The distribution of conservation and restoration priorities distance to**
350   **major natural resource projects in the forestry, energy, and mining sectors.** The
351   points and bars refer to the mean and standard deviation.
352

Discussion

Our study strengthens a growing body of research aiming to inform the impactful
implementation of Nature-based Solutions. Leveraging Reinforcement Learning, we
provide a prioritization framework assessing conservation and restoration scenarios that
maximize threatened biodiversity protection in synergy with co-benefits from ecological
intactness, carbon storage, and water (Figure 1). Moreover, we analyze some local and
economic implications of these scenarios. We test this holistic framework in the context
of Canada's 30x30 conservation and restoration targets stated in the Kunming-Montreal
Global Biodiversity Framework. The results exhibit complex environmental trade-offs
along with potential synergies, demonstrating how Artificial Intelligence can help
optimize conservation and restoration decisions. By testing the framework at the
national level, we show its versatility in incorporating various environmental co-benefits
and scenario analyses to inform pathways to reach environmental targets in different

land use contexts and scales. Particularly, this framework versatility reveals
combinations of conservation and restoration scenarios that maximize both biodiversity
protection and environmental co-benefits.
369

Holistic pathways to reach 30x30 conservation targets
Our results highlight opportunities to reach 30x30 conservation targets that improve
biodiversity while maximizing co-benefits. Currently, our results confirm that the existing
Protected Areas network in Canada does not effectively protect threatened and non-
threatened species [19] and only partially captures the potential co-benefits from carbon
storage and water [16]. These results are not exclusive to Canada and reflect a known
global trend where Protected Areas are biased to locations with reduced land-use
pressures that do not necessarily prioritize biodiversity and other co-benefits outcomes
[37].
379

To improve Protected Areas outcomes, our results show that prioritizing the protection
of threatened biodiversity and irrecoverable carbon stocks yields the highest
improvements in biodiversity and co-benefits. Expanding the Protected Areas network
with this prioritization approach could maintain ecological intactness (i.e., species
abundance, habitat quality) and lead to improved carbon storage and surface water
stability. Most of these priorities are located in southern provinces, extending across
forest-dominated lands in the boreal ecozones to the western Montane Cordillera.
Without accounting for biodiversity, Mitchell et al.[16] have already suggested the potential
of these ecozones for carbon storage, water provision, and nature-based recreation.
Although not assessed in our analysis, these forest lands serve as climate refugia for
multiple taxonomic groups, including vertebrates, plants, and invertebrates [17,18]. Our
results, therefore, demonstrate that conservation targets can be effectively maximized
with avoided land use emissions targets in forested lands while also achieving co-
benefits from ecological intactness and water.
394

Our prioritization framework also identified locations that simultaneously maximize
biodiversity, irrecoverable carbon storage, ecological intactness, and water outcomes.
These win-win conservation priorities could moderately contribute to 30x30
conservation targets. Some are dispersed in the Mixewood Plains and Pacific Maritimes
and surpass biodiversity outcomes in the Boreal ecozones and Montane Cordillera.
These ecozones may guarantee the occurrence of climate-resilient species [19] but
contain regions under high anthropogenic pressures (e.g., St. Lawrence Lowlands,
Manitoulin Lake Simcoe, Lake Erie Lowland) [38]. Other priority locations were clustered:
the southern Hudson Bay and James Bay Lowlands (Quebec and Ontario Hudson
Plains); the Lac Temiscamingue Lowlands (Ontario Boreal Shield); and the ecozones
transition of the Montane Cordillera, Taiga, and Boreal Plains (British Columbia). The
ecozones transition, in particular, may strategically increase the connectivity of
Protected Areas in the Yellowstone to Yukon corridor [39]. Moreover, these biodiversity
and co-benefits clusters have been identified as either climate corridors or refugia [17,18].
Although strategically important, these win-win conservation priorities remain vulnerable
to pervasive climate pressures. This challenge urge exploring suitable adaptive actions,

such as the reduction of land use pressures [40,41], fire prescription [42,43], translocating or
slowing dispersal of species, along with other forms of ecosystem engineering [41,44].
413

Holistic pathways to reach 30x30 restoration targets
Compared to conservation, the restoration priority scenarios suggest diverse and
context-dependent pathways to reach the 30x30 targets. We identify that biodiversity
outcomes are potentially maximized by restoring areas experiencing reduced ecological
intactness in the Prairies and Mixewood Plains that are currently dedicated to
agriculture. These ecozones have already been highlighted for their potential
biodiversity outcomes for both conservation [19] and restoration [28] targets. More aligned
with the latter, our results indicate that these heavily disturbed landscapes are suitable
for diverse interventions. For instance, avoiding further grassland conversion into
cropland, maintaining shelterbelts, and increasing tree intercropping and silvopasture
are essential actions to reduce anthropogenic pressures on biodiversity and the delivery
of co-benefits [45,46]. Additional interventions include restoring wetlands [47] and riparian
corridors across grasslands and croplands [48], which are expected to improve
biodiversity outcomes and water co-benefits (e.g., water provision and quality). These
restoration actions are of particular importance in the Prairies, where we identified that
aligning restoration actions to manage water surface instability and improve ecological
intactness could positively impact local biodiversity. Evidently, small to medium-scale
interventions alone may not achieve restoration targets. Nevertheless, moderately
improving biodiversity and co-benefits by combining management and restoration
interventions can lead to direct benefits for agricultural production [49].
434

In addition to small and medium-scale interventions, our results show the potential for
large-scale interventions in the Prairies and Mixewood Plains to achieve restoration
targets. The significant losses of carbon stocks in the Prairies and Mixewood plains due
to agriculture represent one of the largest opportunities to simultaneously restore
carbon and biodiversity in Canada [28]. However, this restoration potential carries some
caveats. The prairies are not extensive climate refugia or corridors [17–19], and the
Mixewood Plains are expected to experience increased climate instability [50].
Furthermore, given the need to maintain food production, shifting land use in most of
these areas might be unfeasible [6,45], limiting the scale of restoration actions. Addressing
climate pressures and safeguarding food production will require further prioritization
scenarios at higher resolutions that account for opportunity costs [33]. Consequently,
while agricultural lands are fundamental priorities for biodiversity, there are also inherent
tradeoffs that would need to be addressed to realize any substantial contribution to
Canada's 30x30 restoration targets.
449

In contrast to frameworks focusing on converted lands for pasture and agriculture [23], we
identify complementary restoration pathways in forested lands. These restoration
pathways emerge when biodiversity outcomes were prioritized with the loss of
irrecoverable carbon stocks. Most priorities are dispersed across Boreal ecozones and
the Montane Cordillera, mirroring previous findings on restoration scenarios for
biodiversity and carbon [28]. Restoring carbon storage in forested lands produced slightly
lower biodiversity outcomes than restoration scenarios prioritizing agricultural lands.

However, considerable levels of ecological intactness would be maintained by restoring
carbon storage in forested lands. If ecological intactness is maintained, combining
interventions within and around remaining forests, such as improved forest
management and restoring forest cover, could result in short-term benefits for
biodiversity [51]. Additionally, restoring and mitigating further carbon stocks losses in
these forest lands would result in longer-term climate benefits than restoration actions in
agricultural lands [52]. Considering that most of these restoration priorities are near
conservation priorities, coordinating efforts at different government levels [53,54] and
investments from the public and private sectors [33] will be essential.
466
Local and economic implications of conservation and restoration pathways
Any pathway to achieve conservation and restoration targets will be intrinsically related
to Indigenous land governance. According to our priority scenarios, Indigenous-led NbS
currently funded by the federal government (e.g., Indigenous Guardians, Indigenous
Protected Conserved Areas) are already playing a key role in protecting biodiversity,
irrecoverable carbon storage, and other co-benefits in the Pacific and Atlantic
Maritimes, the Montane Cordillera and Hudson Plains. This is the case of the Guardian
Watchmen program in the Pacific, which has enhanced local capacity for monitoring
environmental change and facilitated co-management arrangements [55]. Interestingly,
both conservation and restoration priorities in forested and agricultural lands are closer
to currently unfunded Indigenous Lands than government-funded Indigenous-led NbS.
478
It is worth emphasizing that our analysis likely underestimates the influence of
Indigenous Land governance in conservation and restoration. First, we use government-
recognized Indigenous Lands locations that do not entirely represent the extent of
Indigenous Land governance [30]. Second, several Indigenous Guardians and Indigenous
Protected Areas are not recognized or funded by the Federal Government [56].
Additionally, the federal government Initiative Canada Target 1 Challenge [57], which
explores the establishment of Indigenous Protected Conserved Areas, is not included in
our study and could increase the influence of Indigenous-led NbS on conservation and
restoration priorities. Nevertheless, it remains the case that the proximity of
conservation and restoration priorities to Indigenous Lands exhibits an opportunity for
equitable NbS implementation. To realize this opportunity within the 30x30 targets,
governments at different levels need to coordinate efforts that assert Indigenous
governance [58–61].
492
Along with navigating environmental trade-offs and land governance coordination,
achieving effective 30x30 targets may also result in land use conflicts with major natural
resource projects. Forestry is well known to have impacted forested lands, especially in
the Boreal ecozone [62]. However, only a fraction of planned forestry concessions were
close to conservation and restoration priorities that maximize biodiversity and carbon
outcomes. Instead, energy and mining are the most immediate pressures for these
priorities in forested lands. The Peace, Trans Mountain, and other pipelines planned
across Alberta and British Columbia ecozones transition represent a challenge for
numerous conservation and restoration priorities, including those maximizing multiple
co-benefits. This linear infrastructure causes landscape fragmentation, species

mortality, and changes in species structure and composition [63]. Further challenges and
reduced mitigation capacity emerge in Boreal forested lands across northeastern
Alberta due to oil sand extraction and mining in the Ontario-Québec border. Besides
complete habitat removal, these activities result in landscape-scale disturbances due to
linear infrastructure, light pollution, and chemical pollution in soils and water bodies,
altering biodiversity through complex pathways [64]. While aligning conservation and
restoration priorities across forested lands represent a complementary and effective
pathway to achieve 30x30 targets, major natural resource projects will likely alter
current biodiversity and co-benefits outcomes.
512
Mining and energy project development are also expected to impact restoration
priorities in agricultural lands. In the Prairies, restoration priorities might be impacted by
potash mining in Saskatchewan, wind power in Alberta, and electric facilities and
transmission lines in Manitoba. Similarly, electric power will represent an immediate
pressure for restoration priorities in Ontario's and Quebec's Mixewood Plains. Although
mining methods exert varying effects on biodiversity, they generally exacerbate existing
pressures in areas that have already lost ecological intactness, such as agricultural
lands [65]. Particularly, potash mining's high water use intensity, in combination with
climate change, will further impact water availability for an increasing population and
growing agricultural production [66]. Energy projects effects are expected to extend
beyond habitat removal from in-situ and surrounding linear infrastructure [67]. Wind power
can increase bird and bat species mortality [68], resulting in cascading effects on other
trophic levels [69]. Run-of-river hydropower alters flow regimes, physical habitats,
connectivity, sediment and nutrient dynamics, affecting biodiversity locally as well as
upstream and downstream [70]. In some cases (e.g., wind power, hydropower), project
impacts in agricultural lands would diminish by coordinating mitigation and restoration
actions [71]. Overall, major natural resource projects will increase the baseline of
degraded land, leading to an increased challenge in achieving restoration targets.
531

Conclusions
NbS have the potential to address the intertwined biodiversity, climate, water, and social
equality crises. Using an AI-driven framework, we assess this potential by exploring
conservation and restoration scenarios to reach Canada's 30x30 biodiversity targets in
synergy with climate and water co-benefits. Our results indicate that prioritizing the
protection of threatened species and irrecoverable carbon storage, mostly in forested
lands, represents a key pathway to conserve 30% of land and improve biodiversity and
co-benefits outcomes relative to existing Protected Areas. Restoring 30% of degraded
land for effective biodiversity outcomes will require a combination of scenarios: restoring
carbon stocks in forest lands and mitigating the effects of reduced ecological intactness
and water surface instability in agricultural lands. These findings offer complementary
insights for environmental decision-making. On the one hand, implementing holistic
frameworks with diverse priority scenarios may reveal strategic conservation and
restoration actions that minimize trade-offs and maximize outcomes. On the other hand,
targeted land use planning, policies, and investments can strategically maximize
biodiversity, climate, and water outcomes.
548

To accelerate the strategic implementation and investments of holistic NbS, our results
outline four key governance recommendations that emphasize multi-dimensional
coordination. First, NbS for conservation and restoration actions must be interconnected
to guarantee long-term outcomes [72]. Second, conservation and restoration actions
should be coordinated across different government levels, asserting Indigenous land
governance. Part of this administrative coordination will imply balancing extractive
activities against conservation and restoration priorities. The third dimension of
coordination involves the public and private sectors. Traditionally, the former relies on
policy interventions for conservation, and the latter invests in restoration [33].
Nevertheless, these traditional roles do not have to be static. Finally, decision-making in
the digital age requires coordinating scientific, collective, and artificial intelligence [73]
through ethical and human-centered frameworks [74]. These transdisciplinary frameworks
will require co-developing with regional stakeholders and local rightsholders on which
targets, species, co-benefits, interventions, scale, data, and model training are most
relevant in their specific contexts.
564

Study limitations

Despite developing a holistic framework to prioritize conservation and restoration
targets, our study has some limitations. First, the complexity of conservation and
restoration planning underscores the need for further research at higher resolutions and
smaller spatial scales. For instance, high-resolution scenarios could display low-scale
conservation priorities in agricultural landscapes that are ruled out in our framework.
Aligning species area of habitat maps and land cover changes with habitat quality and
connectivity assessments at high resolutions could identify strategic areas for targeted
actions[75]. Second, our framework does not directly account for the demand or
immediate benefits from biodiversity and related co-benefits. This limitation is illustrated
by water surface, which represent the most accessible water source for people[76] but
could be complemented by assessing water provision and demand[16]. Similarly, carbon
storage co-benefits from NbS in forested or agricultural lands would not necessarily
result in interchangeable climate benefits. To address disparate climate benefits,
Matthews et al.[52] propose tracking the amount of carbon storage achieved over time
and estimating the near-term temperature benefits from that storage. Lastly, while we
compare our results with previous work on climate scenarios and assess potential land
use conflicts, our prioritization framework does not directly incorporate these
assessments. Future prioritization frameworks would benefit from integrating climate
and land use scenarios to envision NbS in different shared socio-economic pathways.
585

Methods

Our study aims to leverage geospatial analysis and Reinforcement Learning to prioritize
NbS locations and reach Canada's 30x30 targets while identifying synergies with other
co-benefits. To this end, we explored two sets of NbS priority scenarios that maximize
the occurrence of threatened vertebrate and plant species within selected land areas.
The first set of scenarios focuses on conservation priorities to maximize threatened

biodiversity occurrence and reach 30X30 conservation targets (i.e., conserve 30% of
land by 2030) in areas with high ecological intactness (Conservation Scenario 1), high
and irrecoverable carbon stocks (Conservation Scenario 2), and stable surface water
(Conservation Scenario 3). The other set of priorities is based on maximizing threatened
biodiversity occurrence to reach 30X30 restoration targets (i.e., restore 30% of
degraded land by 2030) in areas with low ecological intactness (Restoration Scenario
1), carbon stocks losses (Restoration Scenario 2), and significant historical changes in
surface water (Restoration Scenario 3). Based on the resulting priority scenarios, we
analyze the spatial patterns of conservation and restoration priorities across ecozones
and provinces. After, we compare them to current Indigenous-led NbS projects and
ongoing and planned major natural resource projects.
603
Study Area
We use Canada as a case study, which has the second largest country land mass with
more than 9.9 million square kilometers [77]. It is estimated that Canada is home to
around 80000 species, including mammals, birds, fishes, amphibians, reptiles, plants,
and invertebrates [78]. These species are found across diverse ecozones that contain
approximately ~25% of global wetlands, boreal forests, and temperate forests [77].
Summing aboveground biomass, belowground biomass, and soil organic carbon,
Canada's carbon stocks are estimated at 33.7 PgC, representing 13.7% of global
carbon storage, surpassed only by Brazil (18.2%) and Russia (20.9%) [12]. Moreover,
Canada has a surface water area of 1.1 million square kilometers, accounting for 26%
of the global surface water distribution [76]. Given these ecological and geographic
features, Canada presents significant potential to implement NbS initiatives aimed at
achieving combined biodiversity, carbon storage, and surface water outcomes.
617
Canada is also considered to have one of the most extensive areas of global intact
ecosystems [79,80]. The current network of Protected Areas covers 12.7% of terrestrial
land, partially securing these ecosystems' biodiversity, carbon stocks, and surface water
[81]. However, 11.7% of Canada's land area is considered degraded (Department of
Economic and Social Affairs - UN, 2019), and 10% of vertebrate and plant species
registered by the IUCN red list are classified under some category of extinction risk [83].
Additionally, significant portions of Canada's carbon stocks are at risk of being lost and
not recovering by 2050 due to land use and climate change, particularly in ecozones
along the Pacific (e.g., Pacific Maritimes), Southern Artic, and Hudson Plains [84]. Surface
water faces similar risks, which have resulted in transitions from permanent water cover
to seasonal or complete loss [76].
629
Data processing
To determine NbS priority scenarios of conservation and restoration through
Reinforcement Learning, we used a range of datasets on the spatial patterns of
biodiversity, carbon stocks, and surface water occurrence in Canada (Table S6). We
used the International Union for Conservation of Nature species range polygons to
establish the spatial distribution of mammals, fishes, amphibians, reptiles, and plant
species in terrestrial ecosystems [83]. Bird species range polygons were obtained from
BirdLife International [85]. Among these range maps, we selected species experiencing

some risk of extinction among the IUCN Red List categories: Near
Threatened, Vulnerable, Endangered, and Critically Endangered. These selected
categories resulted in freshwater fish (n = 24), migratory birds (n = 21), and plants (n =
17) representing 80% of threatened species (n = 77), while the remaining 20%
corresponded to mammals (n = 8), reptiles (n = 4), and amphibians (n = 1).
643
In addition to the distribution of threatened species, we determined ecological intactness
at different spatial scales following Soto-Navarro et al.[20] (Table S6). Local ecological
intactness was determined through the GLOBIO 4-Mean Species Abundance index that
represents the mean abundance of species in a grid cell (~300m) given a particular
anthropogenic pressure (e.g., land use, climate change, fragmentation, hunting) and
relative to undisturbed areas [86]. This measure of local ecological intactness can be
understood as a β diversity measure because it reflects the compositional turnover
between disturbed and undisturbed lands [20]. Regional ecological intactness derived
from the CSIRO-Biodiversity Habitat Index and provided a measure of habitat
transformation impacts on the retention of terrestrial biodiversity [87]. This ~1km
resolution index relies on estimating habitat conditions based on land cover and spatial
turnover of vascular plant species and reflects the proportion of effective habitat
remaining among grid cells with similar species composition before habitat
transformation. After resampling to the same resolution (~1km), we estimated the
average of both indices to obtain a composite assessment of ecological intactness. The
spatial patterns of threatened species and ecological intactness were resampled to a
10km resolution, which corresponded to the size of Planning Units for potential
conservation and restoration priority scenarios.
662
Total carbon stocks were estimated using publicly available biomass and soil organic
carbon maps. Biomass was obtained from a 2010 global harmonized map of
aboveground and belowground carbon with ~300 m resolution that leverages different
vegetation-specific maps (e.g., forest, tundra, grassland, agriculture, tundra) and land
cover [88]. Soil organic carbon was obtained from Sothe et al., [11], which produced ~250 m
resolution maps at different depths (0-2m) of Canadian terrestrial ecosystems based on
soil ground measurements, covariates (e.g., elevation, slope, precipitation, vegetation
indices, remote sensing spectral bands) and a random forest algorithm. While this map
does not directly correspond to 2010, we assumed the soil carbon estimates
approximate that temporal range, considering that the covariates involved represent an
average of 20 years (2000-2019). Following Noon et al.[84] and using global peatlands
maps [89], we accounted for carbon in peatlands at 1 m depth, which is potentially
vulnerable to human activities, and 0.3 m depth elsewhere. After resampling the soil
organic carbon maps to the resolution of aboveground and belowground biomass
carbon, these estimates were added to determine the stocks of total carbon. Hence, the
total carbon estimates we have used here represent the distribution of stocks in 2010
that might be released by subsequent land cover changes.
680
To determine current total carbon (ca. 2020) and land use emissions, we tracked land
cover transitions since 2010. The transitions were determined through annual land
cover maps for 2010 and 2020 from the ESA CCI products [90]. We opted for the ESA-

CCI maps due to their involvement in the harmonization process of biomass carbon
maps [88]. Using the approach of Hu et al., [91], ESA-CCI land cover classes were
translated to 9 IPCC classes to track transitions in land cover and estimate carbon
stocks losses or gains for the period 2010-2020. Based on the land cover transitions,
the Canadian terrestrial ecozones, and the occurrence of peatlands, carbon stocks
changes were estimated separately for biomass carbon and soil organic carbon (Table
S7).
691
Regarding carbon stock losses, ~60% of land cover transitions for the period 2010-2020
involved forests. The transitions from vegetation classes (i.e., forest, wetland,
shrubland, grassland, sparse vegetation) to non-vegetation (i.e., urban, bare land)
resulted in complete losses of biomass carbon stocks, while partial losses were
estimated for transitions involving vegetation classes and agriculture (Table S7). For
instance, biomass losses from forest-agriculture transitions were estimated by the
difference between the initial biomass carbon in each forest pixel and the terrestrial
ecozone average of biomass carbon stocks in agricultural lands. We followed the same
approach for similar transitions involving high (e.g., forest) to low-carbon vegetation land
cover classes (e.g., shrubland). In the case of soil organic carbon, we assumed that
stocks remained stable for transitions involving forests, shrubland, grassland, and
sparse vegetation [84,92]. Transitions from vegetation classes to agriculture, urban, and
bare land involved a carbon loss function in non-peat soils and complete carbon loss in
peat soils [84,93].
706
Most of the carbon stocks gains involved transitions to forest land cover (~20%).
Biomass gains were estimated according to sequestration rates reported in the boreal,
temperate lowland, temperate mountain [84], and tundra regions [94]. Gains in soil organic
carbon involved transitions from low carbon land covers (i.e., urban, bare land, or
agriculture) to vegetation land cover classes (i.e., shrubland, grassland, sparse-
vegetation, wetland, forest). Low-high carbon transitions were estimated according to
carbon recovery functions reported for temperate and boreal regions, boreal peatlands,
temperate peatlands [93], and tundra [95]. We did not assume changes in carbon stocks for
land cover transitions involving wetlands and forests, as previous studies suggest that
these transitions may not be permanent and are frequently associated with classification
errors [96,97]. Based on the changes in biomass and soil organic carbon for the period
2010-2020, both sources of carbon stocks were added to create a total carbon map for
2020. At the same time, the total carbon maps for 2010 and 2020 were used to estimate
total carbon stocks changes. All resulting total carbon stocks and total carbon change
maps were resampled to a 10km resolution.
722
In addition to the spatial patterns of threatened species, ecological intactness, and total
carbon stocks, we quantified water surface changes. We used the Landsat-derived
estimates of changes in global surface water from 1984 to 2015 [76]. This dataset exhibits
the spatial and temporal changes of any stretch of open, fresh, and salt water, larger
than 30m by 30m. Specifically, we focused on water occurrence change intensity that
results from comparing the frequency of monthly water detection between two periods:
1984-1999 and 2000-2015. Thus, the water occurrence change intensity provides a

measure of gains, losses, and persistence. After masking other land covers, we
estimated the mean water occurrence change intensity in a 10km resolution across
Canada. Capturing these long-term changes in water occurrence may reveal the
impacts of climate change and land use while highlighting adaptation actions affecting
one of the most accessible sources of water for human populations [76]. Water surface
changes were estimated in Google Earth Engine, and the spatial patterns of threatened
species, ecological intactness, and total carbon stocks were established using the R
package Terra [98].
738
Reinforcement learning to determine conservation and restoration priorities
We used the Reinforcement Learning algorithm CAPTAIN [27] to identify conservation
and restoration priorities to reach 30x30 targets in Canada. The process of identifying
priorities in this algorithm can be summarized in four steps: (1) simulate hypothetical
spatial training data; (2) train an optimized model of prioritization through Reinforcement
Learning; (3) build an empirical environment using observed real-world data; and (4)
apply the optimized model within the empirical environment to identify priority
conservation and restoration areas (Figure 1).
747
In the first step, hypothetical natural environments are simulated with a resolution,
spatial extent, and several hypothetical species that resemble the empirical
environment. Each hypothetical species displays unique spatial distributions,
demographic traits, dispersal capacity, and sensitivity to climate and anthropogenic
disturbances. Based on the hypothetical environments, we implemented a monitoring
policy, a budget (target), and a priority policy to be used for model training. In our case,
the monitoring policy gathers information about species (including their presence) and
anthropogenic pressures in each planning unit. The budget defines the maximum
fraction of planning units that can be prioritized and is a function of the anthropogenic
costs. Here, the budget was arbitrarily defined as 10%, that is, a tenth of the total cost of
all planning units in the simulated natural environment. Constrained by the budget, the
prioritization policy was set to reward the selection of planning units that maximize the
occurrence of species and guarantee at least 10% of the spatial distribution of all
species while avoiding high costs (e.g., high anthropogenic disturbance).
762
In the second step (Figure 1), an optimization model was trained to prioritize planning
units (i.e., spatial 10X10 km grids) using Reinforcement Learning according to the
predetermined monitoring policy, budget, and prioritization policy. To this end, CAPTAIN
optimizes the parameters of a neural network to cost-effectively maximize the rewards
resulting from prioritizing planning units. As explained in detail by Silvestro et al.[27], this
training process relies on a two-phase parallelized evolution strategy that repeats in
several epochs. During the first phase, changes in policy parameters are randomly
perturbed and tested in a simulated change in the environment (i.e., species
distributions and anthropogenic pressures), resulting in priority actions and a reward.
After simulating these changes in different iterations that occur in parallel, the second
phase aggregates the results of these iterations to identify improvements in the rewards
and update the neural network parameters. The two-phase process repeats across
different epochs (here, 100), and the rewards from a current epoch are compared with

the weighted average results of previous epochs in a process defined as an advantage
function. Thus, based on the simulated natural environments, we produced a trained
optimization model capable of cost-effectively identifying priority planning units that
maximize the occurrence of species.
780
In the third step, we integrated observed spatial datasets for Canada as an empirical
environment. This environment was created by dividing the country's land surface into
planning units of 10X10 km that included information regarding the presence of
threatened species, the occurrence of Protected Areas, composite ecological
intactness, total carbon stocks, total carbon stocks changes, and water occurrence
change intensity. Using the Canadian Protected and Conserved Areas database [81], we
excluded planning units with a Protected Area surface equal to or higher than 50 km2,
resulting in 12.25% of the area assumed to be already protected and therefore excluded
from priority scenarios. Estimates regarding ecological intactness, carbon stocks and
water occurrence change intensity across planning units were normalized to create a
scale of costs for conservation and restoration priority scenarios.
792
In the final step, the priority scenarios for conservation and restoration aimed to reach
30X30 targets that would maximize the occurrence of threatened biodiversity,
guaranteeing at least 10% of all species ranges while avoiding specific costs. We aimed
to reach 30x30 targets for the conservation scenarios by identifying at least ~16200
priority planning units (~17.75% of the total area). In Conservation Scenario 1, the
normalized ecological intactness values were recalculated so that the highest value (~
1), and therefore the highest cost, corresponded to the lowest ecological intactness.
This cost scale resulted in rewarding the selection of planning units displaying low cost-
high ecological intactness (~0). Similarly, the Conservation Scenario 2 was based on a
cost scale that rewarded the selection of low cost-high total carbon. To prioritize carbon
stocks that, if lost, would not recover by 2050 [84], the original cost of planning units found
in areas with irrecoverable carbon stocks was further reduced to a tenth of the original
value. Using a cost scale displaying the largest values for the most intense water
occurrence changes, the Conservation Scenario 3 rewarded the prioritization of
planning units displaying low cost-low water surface change. Consequently, the
conservation scenarios prioritized ecological intactness, climate change mitigation
potential, and surface water stability, suggesting proactive NbS with low to moderate
anthropogenic pressures.
811
Following the removal of conservation priorities from the empirical environment, we
performed the restoration scenarios. The three scenarios aimed to identify priorities
areas to restore 30% of degraded land in 2019, which corresponded to 3.5% of the total
land area (~3300 planning units) [82]. While the restoration scenarios had the same
themes as the conservation scenarios, we used a different scale of costs and rewards
for the prioritization process. Restoration Scenario 1 and Scenario 3 were based on
scales of costs that rewarded the prioritization of planning units with low intactness and
high-water occurrence change intensity, respectively. Using a normalized scale where
values closer to zero represented the largest carbon stock losses, the Restoration
Scenario 2 rewarded the prioritization of areas that have experienced carbon stocks

losses over the period 2010-2020. Similar to the Conservation Scenario 2, the cost of
carbon stocks losses in planning units with irrecoverable carbon stocks (i.e.,
irrecoverable by 2050 via natural processes) was reduced to a tenth of the original
value. By prioritizing reduced ecological intactness, carbon stock losses, and surface
water instability, the restoration scenarios produced priority areas for reactive NbS
requiring active management and adaptation actions under high anthropogenic
pressures.
829
Conservation and Restoration priorities relative to Indigenous-led NbS
We identified spatial alignments between conservation and restoration priorities with
Indigenous Lands, including Indigenous-led NbS funded by the federal government of
Canada. We focused on two indigenous-led NbS: government-recognized Indigenous
Protected Conserved Areas (IPCAs) and the Indigenous Guardians Federal Program.
IPCAs are part of Indigenous-Led Area-Based Conservation programs [99]. Based on the
Canadian Protected and Conserved Areas Database [81], three IPCAs were included:
Edéhzhíe Protected Area, Thaidene Nëné Territorial Protected Area, and Ts'udé Niļiné
Tuyeta Territorial Protected Area. The Indigenous Guardians Federal program funded
projects from 2018-2022 (Government of Canada, 2023), aiming to support monitoring,
stewardship, and conservation actions across traditional lands and waters [101]. Using the
public database of Indigenous Guardians funded initiatives [100], we related proponents,
such as Indigenous governments, to Indigenous Lands' legislative boundaries [102]. When
these boundaries did not relate to an Indigenous Guardians project, we defined them as
Indigenous Lands. After, for each conservation and restoration priority planning unit, we
identified the nearest distance to an Indigenous-led NbS or other Indigenous Land
centroid location, using the function nearest in the R package Terra [98]. The conservation
and restoration priorities' distance to Indigenous-led NbS and Indigenous Lands allowed
us to identify local opportunities for equitable NbS.
849
Conservation and Restoration priorities relative to major natural resource
projects
Finally, we compared the spatial distribution of resulting conservation and restoration
priorities with major natural resource projects in Canada. The Major Projects Inventory
provided the location of forestry, mining, and energy sector projects currently under
construction or planned within the next ten years [103]. This inventory included high-
impact projects with a minimum investment between 20 (forestry and electricity) and 50
million (mining and energy) CAD. In addition to extractive activities, energy sector
projects (i.e., solar, hydro, and oil), displayed diverse infrastructure facilities for
processing and transportation. Similar to Indigenous-led NbS and other Indigenous
Lands, we identified the nearest distance of each conservation and restoration priority to
major natural resource projects in each sector. By analyzing this spatial distribution, we
aimed to identify potential land conflicts of conservation and restoration actions with the
extraction of natural resources.
864
865

Resource availability

Lead contact
Requests for further information and resources should be directed to and will be fulfilled by the lead contact, Camilo Alejo (camilo.alejo.monroy@futureearth.org).

Materials availability
This study did not generate new unique materials.

Data and code availability
The data and code that support the findings of this study are openly available in the Open Science Framework at https://osf.io/k7wjq/

Acknowledgements

This work has been supported by a Natural Sciences and Engineering Research Council of Canada (NSERC) Alliance grant with Microsoft (ALLPR 571271-21). C.A. was supported by the Horizon Postdoctoral fellowship at Concordia University.

Author Contributions

Conceptualization, C.A., A.L., H.D.M.; Methodology, C.A., H.D.M., Formal Analysis, Investigation, Data Curation, Writing – Original Draft, Visualization, Project Administration C.A.; Writing – Review & Editing, C.A., H.D.M., A.L., A.V.; Supervision, H.D.M.; Funding Acquisition H.D.M., A.L.

Declaration of interests

The authors declare no competing interests.

Supplemental Information

Document S1. **Figure S1.** The land surface extent of Protected Areas in Canada. A. Area distribution across Canadian Terrestrial Ecozones. **Figure S2.** The distribution of threatened vertebrate and plant species in Canada. **Figure S3.** The distribution of Conservation and Restoration Priority Scenarios across Canadian Terrestrial Ecozones. **Table S1.** Summary statistics of ecological intactness, carbon storage, and water occurrence change intensity in Canadian Terrestrial Ecozones. **Table S2.** Summary statistics of threatened species, species richness, ecological intactness, carbon storage, and absolute water occurrence change intensity in Protected Areas and Conservation Priority Scenarios across Canadian Terrestrial Ecozones. **Table S3.** Summary statistics

of threatened species, species richness, ecological intactness, carbon storage, and
absolute water occurrence change intensity in Protected Areas and Conservation
Priority Scenarios across Canadian Terrestrial Ecozones. **Table S4.** Conservation and
restoration priorities overlaps with Indigenous-led NbS and Indigenous Lands. **Table**
S5. Conservation and restoration priorities overlaps with major development projects.
Table S6. Datasets used to determine and analyze Conservation and Restoration
priority scenarios. **Table S7.** Biomass and soil organic carbon change estimates from
land cover transitions.
911
912
913

References

1. CBD (2022). Kunming-Montreal global biodiversity framework - Draft decision submitted by the President.

2. UNFCCC (2015). Paris Agreement.

3. United Nations (2015). Goal 6: Ensure availability and sustainable management of water and sanitation for all. Sustainable Development Goals. https://sdgs.un.org/goals/goal6.

4. Cohen-Shacham, E., Walters, G., Janzen, C., and Maginnis, S. (2016). Nature-based solutions to address global societal challenges E. Cohen-Shacham, G. Walters, C. Janzen, and S. Maginnis, eds. (IUCN International Union for Conservation of Nature) https://doi.org/10.2305/IUCN.CH.2016.13.en.

5. Seddon, N., Chausson, A., Berry, P., Girardin, C.A.J., Smith, A., and Turner, B. (2020). Understanding the value and limits of nature-based solutions to climate change and other global challenges. Preprint at Royal Society Publishing, https://doi.org/10.1098/rstb.2019.0120 https://doi.org/10.1098/rstb.2019.0120.

6. Griscom, B.W., Adams, J., Ellis, P.W., Houghton, R.A., Lomax, G., Miteva, D.A., Schlesinger, W.H., Shoch, D., Siikamäki, J. V., Smith, P., et al. (2017). Natural climate solutions. Proceedings of the National Academy of Sciences 114, 11645–11650. https://doi.org/10.1073/pnas.1710465114.

7. Matthews, H.D., Zickfeld, K., Dickau, M., MacIsaac, A.J., Mathesius, S., Nzotungicimpaye, C.M., and Luers, A. (2022). Temporary nature-based carbon removal can lower peak warming in a well-below 2 °C scenario. Commun Earth Environ 3. https://doi.org/10.1038/s43247-022-00391-z.

8. Ferreira, C.S.S., Kašanin-Grubin, M., Solomun, M.K., Sushkova, S., Minkina, T., Zhao, W., and Kalantari, Z. (2023). Wetlands as nature-based solutions for water management in different environments. Preprint at Elsevier B.V., https://doi.org/10.1016/j.coesh.2023.100476 https://doi.org/10.1016/j.coesh.2023.100476.

945   9.   *Gómez Martín, E., Giordano, R., Pagano, A., van der Keur, P., and Máñez*
946   *Costa, M. (2020). Using a system thinking approach to assess the*
947   *contribution of nature based solutions to sustainable development goals.*
948   *Science of the Total Environment 738.*
949   *https://doi.org/10.1016/j.scitotenv.2020.139693.*

950   10.   *Chollet Ramampiandra, E., Scheidegger, A., Wydler, J., and Schuwirth, N.*
951   *(2023). A comparison of machine learning and statistical species*
952   *distribution models: Quantifying overfitting supports model interpretation.*
953   *Ecol Modell 481. https://doi.org/10.1016/j.ecolmodel.2023.110353.*

954   11.   *Sothe, C., Gonsamo, A., Arabian, J., Kurz, W.A., Finkelstein, S.A., and*
955   *Snider, J. (2022). Large Soil Carbon Storage in Terrestrial Ecosystems of*
956   *Canada. Global Biogeochem Cycles 36.*
957   *https://doi.org/10.1029/2021GB007213.*

958   12.   *Walker, W.S., Gorelik, S.R., Cook-Patton, S.C., Baccini, A., Farina, M.K.,*
959   *Solvik, K.K., Ellis, P.W., Sanderman, J., Houghton, R.A., Leavitt, S.M., et al.*
960   *(2022). The global potential for increased storage of carbon on land.*
961   *Proceedings of the National Academy of Sciences 119.*
962   *https://doi.org/10.1073/pnas.2111312119.*

963   13.   *Chen, J., Chen, S., Fu, R., Li, D., Jiang, H., Wang, C., Peng, Y., Jia, K., and*
964   *Hicks, B.J. (2022). Remote Sensing Big Data for Water Environment*
965   *Monitoring: Current Status, Challenges, and Future Prospects. Preprint at*
966   *John Wiley and Sons Inc, https://doi.org/10.1029/2021EF002289*
967   *https://doi.org/10.1029/2021EF002289.*

968   14.   *Brooks, T.M., Mittermeier, R.A., Da Fonseca, G.A.B., Gerlach, J.,*
969   *Hoffmann, M., Lamoreux, J.F., Mittermeier, C.G., Pilgrim, J.D., and*
970   *Rodrigues, A.S.L. (2006). Global Biodiversity Conservation Priorities.*
971   *Science (1979) 313.*

972   15.   *Funk, S.M., and Fa, J.E. (2010). Ecoregion prioritization suggests an*
973   *armoury not a silver bullet for conservation planning. PLoS One 5.*
974   *https://doi.org/10.1371/journal.pone.0008923.*

975   16.   *Mitchell, M.G.E., Schuster, R., Jacob, A.L., Hanna, D.E.L., Dallaire, C.O.,*
976   *Raudsepp-Hearne, C., Bennett, E.M., Lehner, B., and Chan, K.M.A. (2021).*
977   *Identifying key ecosystem service providing areas to inform national-scale*
978   *conservation planning. Environmental Research Letters 16.*
979   *https://doi.org/10.1088/1748-9326/abc121.*

980   17.   *Saunders, S.P., Grand, J., Bateman, B.L., Meek, M., Wilsey, C.B.,*
981   *Forstenhaeusler, N., Graham, E., Warren, R., and Price, J. (2023).*
982   *Integrating climate-change refugia into 30 by 30 conservation planning in*
983   *North America. Front Ecol Environ 21, 77–84.*
984   *https://doi.org/10.1002/fee.2592.*

985   18.   *Stralberg, D., Carroll, C., and Nielsen, S.E. (2020). Toward a climate-*
986   *informed North American protected areas network: Incorporating climate-*
987   *change refugia and corridors in conservation planning. Preprint at Wiley-*
988   *Blackwell, https://doi.org/10.1111/conl.12712*
989   *https://doi.org/10.1111/conl.12712.*

990   19.   *Eckert, I., Brown, A., Caron, D., Riva, F., and Pollock, L.J. (2023). 30×30*
991   *biodiversity gains rely on national coordination. Nat Commun 14.*
992   *https://doi.org/10.1038/s41467-023-42737-x.*

993   20.   *Soto-Navarro, C., Ravilious, C., Arnell, A., De Lamo, X., Harfoot, M., Hill,*
994   *S.L.L., Wearn, O.R., Santoro, M., Bouvet, A., Mermoz, S., et al. (2020).*
995   *Mapping co-benefits for carbon storage and biodiversity to inform*
996   *conservation policy and action. Philosophical Transactions of the Royal*
997   *Society B: Biological Sciences 375. https://doi.org/10.1098/rstb.2019.0128.*

998   21.   *Jung, M., Arnell, A., de Lamo, X., García-Rangel, S., Lewis, M., Mark, J.,*
999   *Merow, C., Miles, L., Ondo, I., Pironon, S., et al. (2021). Areas of global*
importance for conserving terrestrial biodiversity, carbon and water. Nat
Ecol Evol 5, 1499–1509. https://doi.org/10.1038/s41559-021-01528-7.

22.   *O'connor, L.M.J., Pollock, L.J., Renaud, J., Verhagen, W., Verburg, P.H.,*
Lavorel, S., Maiorano, L., and Thuiller, W. (2021). Balancing conservation
priorities for nature and for people in Europe.

23.   *Strassburg, B.B.N., Iribarrem, A., Beyer, H.L., Cordeiro, C.L., Crouzeilles,*

1006   *R., Jakovac, C.C., Braga Junqueira, A., Lacerda, E., Latawiec, A.E.,*

1007   *Balmford, A., et al. (2020). Global priority areas for ecosystem restoration.*

1008   *Nature 586, 724–729. https://doi.org/10.1038/s41586-020-2784-9.*

1009   24.   *Vettorazzi, C.A., and Valente, R.A. (2016). Priority areas for forest*

1010   *restoration aiming at the conservation of water resources. Ecol Eng 94,*

1011   *255–267. https://doi.org/10.1016/j.ecoleng.2016.05.069.*

1012   25.   *Chauvenet, A.L.M., Watson, J.E.M., Adams, V.M., Di Marco, M., Venter, O.,*

1013   *Davis, K.J., Mappin, B., Klein, C.J., Kuempel, C.D., and Possingham, H.P.*

1014   *(2020). To Achieve Big Wins for Terrestrial Conservation, Prioritize*

1015   *Protection of Ecoregions Closest to Meeting Targets. In One Earth (Cell*

1016   *Press), pp. 479–486. https://doi.org/10.1016/j.oneear.2020.04.013.*

1017   26.   *Mappin, B., Chauvenet, A.L.M., Adams, V.M., Di Marco, M., Beyer, H.L.,*

1018   *Venter, O., Halpern, B.S., Possingham, H.P., and Watson, J.E.M. (2019).*

1019   *Restoration priorities to achieve the global protected area target. Preprint at*

1020   *Wiley-Blackwell, https://doi.org/10.1111/conl.12646*

1021   *https://doi.org/10.1111/conl.12646.*

1022   27.   *Silvestro, D., Goria, S., Sterner, T., and Antonelli, A. (2022). Improving*

1023   *biodiversity protection through artificial intelligence. Nat Sustain 5, 415–424.*

1024   *https://doi.org/10.1038/s41893-022-00851-6.*

1025   28.   *Currie, J., Merritt, W., Liang, C., Sothe, C., Beatty, C.R., Shackelford, N.,*

1026   *Hirsh-Pearson, K., Gonsamo, A., and Snider, J. (2023). Prioritizing*

1027   *ecological restoration of converted lands in Canada by spatially integrating*

1028   *organic carbon storage and biodiversity benefits. Conserv Sci Pract 5.*

1029   *https://doi.org/10.1111/csp2.12924.*

1030   29.   *Tauli-Corpuz, V., Alcorn, J., Molnar, A., Healy, C., and Barrow, E. (2020).*

1031   *Cornered by PAs: Adopting rights-based approaches to enable cost-*

1032   *effective conservation and climate action. World Dev 130.*

1033   *https://doi.org/10.1016/j.worlddev.2020.104923.*

1034   30.   *Zurba, M., Beazley, K.F., English, E., and Buchmann-Duck, J. (2019).*

1035   *Indigenous Protected and Conserved Areas (IPCAs), Aichi target 11 and*

1036   *Canada's pathway to target 1: Focusing conservation on reconciliation.*

1037   *Land (Basel) 8. https://doi.org/10.3390/land8010010.*

1038   31.   *Garnett, S.T., Austin, B., Collier, N.F., Duncan, T., Geyle, H., Jackson,*

1039   *M.V., McGowan, B., Sivongxay, A., Leiper, I., Burgess, N.D., et al. (2018). A*

1040   *spatial overview of the global importance of Indigenous lands for*

1041   *conservation. Nat Sustain.*

1042   32.   *Fa, J.E., Watson, J.E.M., Leiper, I., Potapov, P., Evans, T.D., Burgess,*

1043   *N.D., Molnár, Z., Fernández-Llamazares, Á., Duncan, T., Wang, S., et al.*

1044   *(2020). Importance of Indigenous Peoples' lands for the conservation of*

1045   *Intact Forest Landscapes. Front Ecol Environ 18, 135–140.*

1046   *https://doi.org/10.1002/fee.2148.*

1047   33.   *Cook-Patton, S.C., Drever, C.R., Griscom, B.W., Hamrick, K., Hardman, H.,*

1048   *Kroeger, T., Pacheco, P., Raghav, S., Stevenson, M., Webb, C., et al.*

1049   *(2021). Protect, manage and then restore lands for climate mitigation.*

1050   *Preprint at Nature Research, https://doi.org/10.1038/s41558-021-01198-0*

1051   *https://doi.org/10.1038/s41558-021-01198-0.*

1052   34.   *Carr, J.A., Petrokofsky, G., Spracklen, D. V., Lewis, S.L., Roe, D., Trull, N.,*

1053   *Vidal, A., Wicander, S., Worthington-Hill, J., and Sallu, S.M. (2021).*

1054   *Anticipated impacts of achieving SDG targets on forests - a review. Preprint*

1055   *at Elsevier B.V., https://doi.org/10.1016/j.forpol.2021.102423*

1056   *https://doi.org/10.1016/j.forpol.2021.102423.*

1057   35.   *Lawley, C.J.M., Mitchell, M.G.E., Stralberg, D., Schuster, R., McIntire, E.,*

1058   *and Bennett, J.R. (2022). Mapping Canada's Green Economic Pathways for*

1059   *Battery Minerals: Balancing Prospectivity Modelling With Conservation and*

1060   *Biodiversity Values. Earth Science, Systems and Society 2.*

1061   *https://doi.org/10.3389/esss.2022.10064.*

1062   36.   *Rehbein, J.A., Watson, J.E.M., Lane, J.L., Sonter, L.J., Venter, O.,*

1063   *Atkinson, S.C., and Allan, J.R. (2020). Renewable energy development*

1064   *threatens many globally important biodiversity areas. Glob Chang Biol 26,*

1065   *3040–3051. https://doi.org/10.1111/gcb.15067.*

1066   37.   *Joppa, L.N., and Pfaff, A. (2009). High and far: Biases in the location of*
1067   *protected areas. PLoS One 4.*
1068   *https://doi.org/10.1371/journal.pone.0008273.*

1069   38.   *Kraus, D., and Hebb, A. (2020). Southern Canada's crisis ecoregions:*
1070   *identifying the most significant and threatened places for biodiversity*
1071   *conservation. Biodivers Conserv 29, 3573–3590.*
1072   *https://doi.org/10.1007/s10531-020-02038-x.*

1073   39.   *Chester, C.C. (2015). Yellowstone to Yukon: Transborder conservation*
1074   *across a vast international landscape. Environ Sci Policy 49, 75–84.*
1075   *https://doi.org/10.1016/j.envsci.2014.08.009.*

1076   40.   *Welch, D. (2005). What Should Protected Areas Managers Do in the Face*
1077   *of Climate Change?*

1078   41.   *Rannow, S., Macgregor, N.A., Albrecht, J., Crick, H.Q.P., Förster, M.,*
1079   *Heiland, S., Janauer, G., Morecroft, M.D., Neubert, M., Sarbu, A., et al.*
1080   *(2014). Managing Protected Areas Under Climate Change: Challenges and*
1081   *Priorities. Environ Manage 54, 732–743. https://doi.org/10.1007/s00267-*
1082   *014-0271-5.*

1083   42.   *Pereira, P., Mierauskas, P., Ubeda, X., Mataix-Solera, J., and Cerda, A.*
1084   *(2012). Fire in Protected Areas - the Effect of Protection and Importance of*
1085   *Fire Management. Environmental Research, Engineering and Management*
1086   *59. https://doi.org/10.5755/j01.erem.59.1.856.*

1087   43.   *Wang, W., Wu, W., Guo, F., and Wang, G. (2022). Fire regime and*
1088   *management in Canada's protected areas. International Journal of*
1089   *Geoheritage and Parks 10, 240–251.*
1090   *https://doi.org/10.1016/j.ijgeop.2022.04.003.*

1091   44.   *Lemieux, C.J., and Scott, D.J. (2011). Changing climate, challenging*
1092   *choices: Identifying and evaluating climate change adaptation options for*
1093   *protected areas management in Ontario, Canada. Environ Manage 48,*
1094   *675–690. https://doi.org/10.1007/s00267-011-9700-x.*

1095   45.   *Drever, C.R., Cook-Patton, S.C., Akhter, F., Badiou, P.H., Chmura, G.L.,*
1096   *Davidson, S.J., Desjardins, R.L., Dyk, A., Fargione, J.E., Fellows, M., et al.*

(2021). Natural climate solutions for Canada. Sci Adv 7.

https://doi.org/10.1126/sciadv.abd6034.

46. *Keesstra, S., Nunes, J., Novara, A., Finger, D., Avelar, D., Kalantari, Z., and*

Cerdà, A. (2018). The superior effect of nature based solutions in land

management for enhancing ecosystem services. Preprint at Elsevier B.V.,

https://doi.org/10.1016/j.scitotenv.2017.08.077

https://doi.org/10.1016/j.scitotenv.2017.08.077.

47. *Salaria, S., Howard, R., Clare, S., and Creed, I.F. (2019). Incomplete*

recovery of plant diversity in restored prairie wetlands on agricultural

landscapes. Restor Ecol 27, 520–530. https://doi.org/10.1111/rec.12890.

48. *Scherr, S.J., and McNeely, J.A. (2008). Biodiversity conservation and*

agricultural sustainability: Towards a new paradigm of "ecoagriculture"

landscapes. Philosophical Transactions of the Royal Society B: Biological

Sciences 363, 477–494. https://doi.org/10.1098/rstb.2007.2165.

49. *Rey Benayas, J.M., and Bullock, J.M. (2012). Restoration of Biodiversity*

and Ecosystem Services on Agricultural Land. Ecosystems 15, 883–899.

https://doi.org/10.1007/s10021-012-9552-0.

50. *Herrando-Moraira, S., Nualart, N., Galbany-Casals, M., Garcia-Jacas, N.,*

Ohashi, H., Matsui, T., Susanna, A., Tang, C.Q., and López-Pujol, J.

(2022). Climate Stability Index maps, a global high resolution cartography of

climate stability from Pliocene to 2100. Sci Data 9.

https://doi.org/10.1038/s41597-022-01144-5.

51. *Rayden, T., Jones, K.R., Austin, K., and Radachowsky, J. (2023). Improving*

climate and biodiversity outcomes through restoration of forest integrity.

Conservation Biology 37. https://doi.org/10.1111/cobi.14163.

52. *Matthews, H.D., Zickfeld, K., Koch, A., and Luers, A. (2023). Accounting for*

the climate benefit of temporary carbon storage in nature. Nat Commun 14.

https://doi.org/10.1038/s41467-023-41242-5.

53. *Cardona Santos, E.M., Kinniburgh, F., Schmid, S., Büttner, N., Pröbstl, F.,*

Liswanti, N., Komarudin, H., Borasino, E., Ntawuhiganayo, E.B., and

Zinngrebe, Y. (2023). Mainstreaming revisited: Experiences from eight

countries on the role of National Biodiversity Strategies in practice. Preprint
at Elsevier B.V., https://doi.org/10.1016/j.esg.2023.100177
https://doi.org/10.1016/j.esg.2023.100177.

54.   Milner-Gulland, E.J., Addison, P., Arlidge, W.N.S., Baker, J., Booth, H.,
Brooks, T., Bull, J.W., Burgass, M.J., Ekstrom, J., zu Ermgassen, S.O.S.E.,
et al. (2021). Four steps for the Earth: mainstreaming the post-2020 global
biodiversity framework. Preprint at Cell Press,
https://doi.org/10.1016/j.oneear.2020.12.011
https://doi.org/10.1016/j.oneear.2020.12.011.

55.   Whitney, C.K., and Ban, N.C. (2019). Barriers and opportunities for social-
ecological adaptation to climate change in coastal British Columbia. Ocean
Coast Manag 179. https://doi.org/10.1016/j.ocecoaman.2019.05.010.

56.   Mansuy, N., Staley, D., Alook, S., Parlee, B., Thomson, A., Littlechild, D.B.,
Munson, M., and Didzena, F. (2023). Indigenous protected and conserved
areas (IPCAs): Canada's new path forward for biological and cultural
conservation and Indigenous well-being. Facets 8, 1–16.
https://doi.org/10.1139/facets-2022-0118.

57.   Government of Canada (2021). Canada Target 1 Challenge. Canada's
Nature Legacy: protecting our nature.
https://www.canada.ca/en/environment-climate-change/services/nature-
legacy/canada-target-one-challenge.html.

58.   Artelle, K.A., Zurba, M., Bhattacharrya, J., Chan, D.E., Brown, K., Housty,
J., and Moola, F. (2019). Supporting resurgent Indigenous-led governance:
A nascent mechanism for just and effective conservation. Biol Conserv 240.
https://doi.org/10.1016/j.biocon.2019.108284.

59.   Nitah, S. (2021). Indigenous peoples proven to sustain biodiversity and
address climate change: Now it's time to recognize and support this
leadership. One Earth 4, 907–909.
https://doi.org/10.1016/j.oneear.2021.06.015.

60.   Reed, G., Brunet, N.D., Longboat, S., and Natcher, D.C. (2021). Indigenous
guardians as an emerging approach to indigenous environmental

governance. Preprint at Blackwell Publishing Inc.,

https://doi.org/10.1111/cobi.13532 https://doi.org/10.1111/cobi.13532.

61.   Vogel, B., Yumagulova, L., McBean, G., and Charles Norris, K.A. (2022).

1162   Indigenous-Led Nature-Based Solutions for the Climate Crisis: Insights from

1163   Canada. Sustainability (Switzerland) 14.

1164   https://doi.org/10.3390/su14116725.

1165   62.   Wells, J. V., Dawson, N., Culver, N., Reid, F.A., and Morgan Siegers, S.

1166   (2020). The State of Conservation in North America's Boreal Forest: Issues

1167   and Opportunities. Frontiers in Forests and Global Change 3.

1168   https://doi.org/10.3389/ffgc.2020.00090.

1169   63.   Richardson, M.L., Wilson, B.A., Aiuto, D.A.S., Crosby, J.E., Alonso, A.,

1170   Dallmeier, F., and Golinski, G.K. (2017). A review of the impact of pipelines

1171   and power lines on biodiversity and strategies for mitigation. Preprint at

1172   Springer Netherlands, https://doi.org/10.1007/s10531-017-1341-9

1173   https://doi.org/10.1007/s10531-017-1341-9.

1174   64.   Roberts, D.R., Bayne, E.M., Beausoleil, D., Dennett, J., Fisher, J.T.,

1175   Hazewinkel, R.O., Sayanda, D., Wyatt, F., and Dubé, M.G. (2022). A

1176   synthetic review of terrestrial biological research from the Alberta oil sands

1177   region: 10 years of published literature. Integr Environ Assess Manag 18,

1178   388–406. https://doi.org/10.1002/ieam.4519.

1179   65.   Sonter, L.J., Ali, S.H., and Watson, J.E.M. (2018). Mining and biodiversity:

1180   Key issues and research needs in conservation science. Preprint at Royal

1181   Society Publishing, https://doi.org/10.1098/rspb.2018.1926

1182   https://doi.org/10.1098/rspb.2018.1926.

1183   66.   Clark, R., Andreichuk, I., Sauchyn, D., and McMartin, D.W. (2017).

1184   Incorporating climate change scenarios and water-balance approach to

1185   cumulative assessment models of solution potash Mining in the Canadian

1186   Prairies. Clim Change 145, 321–334. https://doi.org/10.1007/s10584-017-

1187   2099-5.

1188   67.   Niebuhr, B.B., Sant'Ana, D., Panzacchi, M., van Moorter, B., Sandström, P.,

1189   Morato, R.G., and Skarin, A. (2022). Renewable energy infrastructure

1190   *impacts biodiversity beyond the area it occupies. Preprint at National*

1191   *Academy of Sciences, https://doi.org/10.1073/pnas.2208815119*

1192   *https://doi.org/10.1073/pnas.2208815119.*

1193   68.   *Schuster, E., Bulling, L., and Köppel, J. (2015). Consolidating the State of*

1194   *Knowledge: A Synoptical Review of Wind Energy's Wildlife Effects. Environ*

1195   *Manage 56, 300–331. https://doi.org/10.1007/s00267-015-0501-5.*

1196   69.   *Thaker, M., Zambre, A., and Bhosale, H. (2018). Wind farms have*

1197   *cascading impacts on ecosystems across trophic levels. Nat Ecol Evol 2,*

1198   *1854–1858. https://doi.org/10.1038/s41559-018-0707-z.*

1199   70.   *Anderson, D., Moggridge, H., Warren, P., and Shucksmith, J. (2015). The*

1200   *impacts of "run-of-river" hydropower on the physical and ecological*

1201   *condition of rivers. Water and Environment Journal 29, 268–276.*

1202   *https://doi.org/10.1111/wej.12101.*

1203   71.   *Gorman, C.E., Torsney, A., Gaughran, A., McKeon, C.M., Farrell, C.A.,*

1204   *White, C., Donohue, I., Stout, J.C., and Buckley, Y.M. (2023). Reconciling*

1205   *climate action with the need for biodiversity protection, restoration and*

1206   *rehabilitation. Preprint at Elsevier B.V.,*

1207   *https://doi.org/10.1016/j.scitotenv.2022.159316*

1208   *https://doi.org/10.1016/j.scitotenv.2022.159316.*

1209   72.   *Wiens, J.A., and Hobbs, R.J. (2015). Integrating conservation and*

1210   *restoration in a changing world. Preprint at Oxford University Press,*

1211   *https://doi.org/10.1093/biosci/biu235 https://doi.org/10.1093/biosci/biu235.*

1212   73.   *Luers, A.L. (2021). Planetary intelligence for sustainability in the digital age:*

1213   *Five priorities. One Earth 4, 772–775.*

1214   *https://doi.org/10.1016/j.oneear.2021.05.013.*

1215   74.   *Shneiderman, B. (2021). Human-Centered AI. Issues Sci Technol, 56–61.*

1216   75.   *Arce-Plata, M.I., Burbano Giron, J., Norden, N., Diaz Corzo, M.C.,*

1217   *Larocque, G., Vasquez Peinado, A.J., and Poisot, T. (2023). Evaluation of*

1218   *the state and trends of habitat change on the Colombian Tropical Dry*

1219   *Forest. In GEO BON Global Conference: Monitoring Biodiversity for Action.*

76. Pekel, J.F., Cottam, A., Gorelick, N., and Belward, A.S. (2016). High-
resolution mapping of global surface water and its long-term changes.
Nature 540, 418–422. https://doi.org/10.1038/nature20584.

77. Statistics Canada (2011). Geography. In Canada Year Book.

78. Canadian Endangered Species Conservation Council (2016). Wild Species
2015: The General Status of Species in Canada.

79. Williams, B.A., Venter, O., Allan, J.R., Atkinson, S.C., Rehbein, J.A., Ward,
M., Di Marco, M., Grantham, H.S., Ervin, J., Goetz, S.J., et al. (2020).
Change in Terrestrial Human Footprint Drives Continued Loss of Intact
Ecosystems. One Earth 3, 371–382.
https://doi.org/10.1016/j.oneear.2020.08.009.

80. Watson, J.E.M., Evans, T., Venter, O., Williams, B., Tulloch, A., Stewart, C.,
Thompson, I., Ray, J.C., Murray, K., Salazar, A., et al. (2018). The
exceptional value of intact forest ecosystems. Preprint at Nature Publishing
Group, https://doi.org/10.1038/s41559-018-0490-x
https://doi.org/10.1038/s41559-018-0490-x.

81. Environment and Climate Change Canada (2023). Canadian Protected and
Conserved Areas Database (CPCAD). ECCC Data Catalogue. https://data-
donnees.az.ec.gc.ca/data/species/protectrestore/canadian-protected-
conserved-areas-database/?lang=en.

82. Department of Economic and Social Affairs - UN (2019). SDG Indicators
Database. Indicator 15.3.1, Series : Proportion of land that is degraded over
total land area (%) AG_LND_DGRD.
https://unstats.un.org/sdgs/dataportal/database.

83. IUCN (2023). IUCN Red List of Threatened Species. Version 2023-1.
https://www.iucnredlist.org/.

84. Noon, M.L., Goldstein, A., Ledezma, J.C., Roehrdanz, P.R., Cook-Patton,
S.C., Spawn-Lee, S.A., Wright, T.M., Gonzalez-Roglich, M., Hole, D.G.,
Rockström, J., et al. (2022). Mapping the irrecoverable carbon in Earth's
ecosystems. Nat Sustain 5, 37–46. https://doi.org/10.1038/s41893-021-
00803-6.

85. *BirdLife International, and Handbook of the Birds of the World (2022).*
Species Distribution data. Bird species distribution maps of the world -
Version 7.0. https://datazone.birdlife.org/species/requestdis.

86. *Kim, H., Rosa, I.M.D., Alkemade, R., Leadley, P., Hurtt, G., Popp, A., Van*
Vuuren, D.P., Anthoni, P., Arneth, A., Baisero, D., et al. (2018). A protocol
for an intercomparison of biodiversity and ecosystem services models using
harmonized land-use and climate scenarios. Geosci Model Dev 11, 4537–
4562. https://doi.org/10.5194/gmd-11-4537-2018.

87. *Hoskins, A.J., Harwood, T.D., Ware, C., Williams, K.J., Perry, J.J., Ota, N.,*
Croft, J.R., Yeates, D.K., Jetz, W., Golebiewski, M., et al. (2020). BILBI:
Supporting global biodiversity assessment through high-resolution
macroecological modelling. Environmental Modelling and Software 132.
https://doi.org/10.1016/j.envsoft.2020.104806.

88. *Spawn, S.A., Sullivan, C.C., Lark, T.J., and Gibbs, H.K. (2020). Harmonized*
global maps of above and belowground biomass carbon density in the year
2010. Sci Data 7. https://doi.org/10.1038/s41597-020-0444-4.

89. *Xu, J., Morris, P.J., Liu, J., and Holden, J. (2018). PEATMAP: Refining*
estimates of global peatland distribution based on a meta-analysis. Catena
(Amst) 160, 134–140. https://doi.org/10.1016/j.catena.2017.09.010.

90. *Defourny, P., Lamarche, C., Brockmann, C., Boettcher, M., Bontemps, S.,*
De Maet, T., Duveiller, G.L., Harper, K., Hartley A., Kirches, G., et al.
(2023). Observed annual global land-use change from 1992 to 2020 three
times more dynamic than reported by inventory-based statistics.

91. *Hu, X., Næss, J.S., Iordan, C.M., Huang, B., Zhao, W., and Cherubini, F.*
(2021). Recent global land cover dynamics and implications for soil erosion
and carbon losses from deforestation. Anthropocene 34.
https://doi.org/10.1016/j.ancene.2021.100291.

92. *Li, W., Ciais, P., Guenet, B., Peng, S., Chang, J., Chaplot, V., Khudyaev,*
S., Peregon, A., Piao, S., Wang, Y., et al. (2018). Temporal response of soil
organic carbon after grassland-related land-use change. Glob Chang Biol
24, 4731–4746. https://doi.org/10.1111/gcb.14328.

93. Poeplau, C., Don, A., Vesterdal, L., Leifeld, J., Van Wesemael, B., Schumacher, J., and Gensior, A. (2011). Temporal dynamics of soil organic carbon after land-use change in the temperate zone - carbon response functions as a model approach. Glob Chang Biol 17, 2415–2427. https://doi.org/10.1111/j.1365-2486.2011.02408.x.

94. Rocha, A. V., and Shaver, G.R. (2011). Burn severity influences postfire CO_2 exchange in arctic tundra. Ecological Applications 21, 477–489. https://doi.org/10.1890/10-0255.1.

95. Loranty, M.M., Natali, S.M., Berner, L.T., Goetz, S.J., Holmes, R.M., Davydov, S.P., Zimov, N.S., and Zimov, S.A. (2014). Siberian tundra ecosystem vegetation and carbon stocks four decades after wildfire. J Geophys Res Biogeosci 119, 2144–2154. https://doi.org/10.1002/2014JG002730.

96. Amani, M., Mahdavi, S., Kakooei, M., Ghorbanian, A., Brisco, B., Delancey, E., Toure, S., and Reyes, E.L. (2021). Wetland Change Analysis in Alberta, Canada Using Four Decades of Landsat Imagery. IEEE J Sel Top Appl Earth Obs Remote Sens 14, 10314–10335. https://doi.org/10.1109/JSTARS.2021.3110460.

97. Li, N., Zhou, S., and Margenot, A.J. (2023). From prairie to crop: Spatiotemporal dynamics of surface soil organic carbon stocks over 167 years in Illinois, U.S.A. Science of the Total Environment 857, 159038. https://doi.org/10.1016/j.scitotenv.2022.159038.

98. Hijmans, Robert., Bivand, R., Pebesma, E., and Summer, Michael.D. (2024). Package "terra." Preprint.

99. Government of Canada (2022). Indigenous-Led Area-Based Conservation. https://www.canada.ca/en/environment-climate-change/services/nature-legacy/indigenous-led-area-based-conservation.html#toc6.

100. Government of Canada (2023). Indigenous Guardians - Canada.ca. https://www.canada.ca/en/environment-climate-change/services/environmental-funding/indigenous-guardians.html.

1312   101. *Artelle, K.A., Zurba, M., Bhattacharrya, J., Chan, D.E., Brown, K., Housty,*

1313   *J., and Moola, F. (2019). Supporting resurgent Indigenous-led governance:*

1314   *A nascent mechanism for just and effective conservation. Biol Conserv 240,*

1315   *108284. https://doi.org/10.1016/j.biocon.2019.108284.*

1316   102. *Government of Canada (2022). Aboriginal Lands of Canada Legislative*

1317   *Boundaries - Open Government Portal.*

1318   *https://open.canada.ca/data/en/dataset/522b07b9-78e2-4819-b736-*

1319   *ad9208eb1067.*

1320   103. *Natural Resources Canada (2024). Major Projects Inventory. Open*

1321   *Government Portal. https://open.canada.ca/data/en/dataset/f5f2db55-31e4-*

1322   *42fb-8c73-23e1c44de9b2.*

1339