Signatures of genomic adaptation to Cuckoo brood parasitism in Reed Warbler hosts across Europe

William J. Smith[1,2,3*†], Katja Rönkä[1,2*], Staffan Bensch[4], Nora Bergman[1,2], Rachael Y. Dudaniec[5], Fabrice Eroukhmanoff[6], Frode Fossøy[7], Bengt Hansson[4], Michał T. Jezierski[1,2,8], Edward Kluen[1,2], Petr Procházka[9], Camilla Lo Cascio Sætre[6], Bård G. Stokke[7], & Rose Thorogood[1,2]

1. Helsinki Institute of Life Science, University of Helsinki, Finland.
2. Research Programme in Organismal and Evolutionary Biology, Faculty of Biological and Environmental Sciences, University of Helsinki, Finland.
3. Department of Biology, University of Oxford, United Kingdom.
4. Department of Biology, Lund University, Sweden.
5. School of Natural Sciences, Macquarie University, Sydney, NSW, Australia.
6. Department of Biology, University of Oslo, Norway.
7. Norwegian Institute for Nature Research (NINA), Trondheim, Norway.
8. School Of Geography, Earth and Environmental Sciences, University of Birmingham, United Kingdom.
9. Institute of Vertebrate Biology, Czech Academy of Sciences, Czech Republic.

*Joint first authors

† Correspondence should be addressed to william.smith@helsinki.fi

Abstract

Predicting coevolutionary outcomes is critical for understanding the consequences of climate

change and biodiversity loss. The geographic mosaic theory of coevolution posits that

coevolutionary interactions vary among populations, facilitating the persistence of such

interactions over evolutionary timescales. Characterisation of geographic mosaics of

coevolution is difficult, with mobile organisms and behavioural adaptations posing temporal

and spatial challenges. The Reed Warbler (*Acrocephalus scirpaceus*) and its brood parasite,

the Cuckoo (*Cuculus canorus*), have been used to study coevolution for decades. By leveraging

RAD-sequencing and brood parasitism data for 188 Reed Warblers across 22 European sites,

we first demonstrate that the magnitude of gene flow varies across the species' distribution,

allowing for variable selection which is a fundamental component of the geographic mosaic

theory. We then use genotype-environment association analysis to reveal genes that are

putatively involved in the Reed Warbler's response to brood parasitism, providing the first

evidence for a genetic basis to this geographic mosaic of coevolution. Together, these results

help to characterise this important model of host–parasite coevolution. Further work is needed

to confirm a causal relationship between parasitism and genomic variation, and to identify the

exact selection mechanisms underpinning this relationship.

Keywords: coevolution, brood parasitism, gene flow, genotype-environment interactions, reed

warbler, selection

Introduction

Antagonistic species interactions are major drivers of evolution. Predators and prey, as well as parasites or pathogens and their hosts, directly impact one another's fitness, exerting strong selective pressure. This can result in antagonistic coevolution[1], where adaptation and counter-adaptation lead to reciprocal evolution in interacting species. Such 'arms races' are of fundamental importance, impacting both contemporary population dynamics[2] and evolutionary diversification[3]. Despite its central role in ecology and evolution, the genetic basis of many aspects of antagonistic coevolution remains underexplored, hindering our ability to understand this phenomenon[4]. Recent advances in genomic approaches offer unprecedented opportunities to address these gaps. In ecology, we are now able to identify the functional genomic underpinnings of physiological changes required to colonise new environments[5]. Genomic approaches hold similar promise in helping us to deepen our understanding of antagonistic coevolution and its impact on interacting species, including in case studies where behavioural traits are fundamental components of these interactions[6].

The geographic mosaic theory of coevolution (GMTC) is the prevailing theory explaining how antagonistic interactions can be maintained over time[7]. Without spatio-temporal variation in species interactions, such arms-races are not predicted to last for long, since sooner or later one entity 'wins': either host defences render parasitism impossible, or the parasites overcome host defences[8]. However, if this happens only locally and there is enough gene flow from other populations, interactions can continue at the landscape level. Host shifts by parasites might also play an important role. Landscape-level interactions enable antagonistic evolutionary relationships to persist over time at a metapopulation level (in many bird species, for example,

brood parasitism relationships may operate at a continental scale)[7]. The study of GMTC-related

questions remains particularly challenging when host and parasite adaptations manifest as

behavioural traits, given their inherent plasticity. The use of genotype-environment analyses

could provide a potent tool with which to characterise geographic mosaics of coevolution in

such systems, particularly in non-model and mobile species where common garden

experiments would be incredibly challenging to design and execute.

Despite plenty of attention on this topic, evidence for geographic mosaics has largely remained

at a descriptive level, demonstrating hot and cold spots of coevolution[9] without a detailed

understanding of the underlying mechanisms, such as the strength and direction of selection

within either the parasite or host, or levels of gene flow within the host and parasite (although

see Kaur *et al.* 2019 for work carried out in an invertebrate system[10]). This is a particular issue

in systems where coevolutionary adaptations are largely behavioural in nature – because (1)

genomic data resolution has only recently advanced to the point where we can scan genomes

for such genetic mechanisms, and (2) behavioural ecologists studying these systems have

traditionally ignored or simplified the complex genetic basis of behavioural traits (i.e. the

"phenotypic gambit"[11]). It is also debated whether plasticity explains observed variation in host

responses to parasitism in particular systems[12]. Genetic signals in response to parasitism are

important to identify if we are to strengthen the evidence for variable responses not being due

to such (e.g. behavioural) plasticity. The successes of genomic approaches in identifying the

genetic underpinnings of behavioural variation more generally e.g.[13,14], combined with an

awareness that local adaptation can persist despite significant gene flow among

populations[15,16], suggests that such an assessment is worthwhile.

Avian brood parasitism is a textbook example of antagonistic coevolution, where reciprocal

arms-race selection has led to the development of complex behavioural adaptations in brood

parasites and their hosts[17-19]. For example, the Reed Warbler (*Acrocephalus scirpaceus*), one

of the preferred European hosts of the brood parasitic Cuckoo (*Cuculus canorus*)[20], defends by

mobbing the Cuckoo at the nest site[21], or by recognising and ejecting foreign eggs[22]. For their

part, Cuckoos evolved host-specific female races (gentes), which mimic their preferred host's

eggs with increasing accuracy (see Stokke *et al.* for discussion of the role of crypsis as an

alternative to egg mimicry[23]), and so Cuckoo parasitism could also impact host egg colour, as

well as nest building and incubation behaviour. The existence of gentes perhaps explains why

genetic research has tended to focus on Cuckoos, perhaps at the expense of their hosts[26,27].

Despite over 40 years of research into this classic case study for behavioural coevolution[22,24,25],

our understanding of the way brood parasitism shapes host defences has been almost

completely limited to the phenotypic level[28], meaning that the underlying heritable mechanisms

remain unknown. This is a major drawback in terms of characterising a geographic mosaic of

coevolution[9], as it is widely recognised that knowledge of the genetic basis of adaptive traits is

needed to understand and ultimately predict coevolutionary trajectories[29]. The few existing

studies which have attempted to delineate the relationship between brood parasitism and

genetics have provided mixed evidence. The first such attempt, by Martín-Gálvez *et al.*[30,31],

detected an apparent association between an allele at microsatellite locus Ase64 and egg

rejection behaviour in European Magpies (*Pica pica*), parasitised by Great Spotted Cuckoos

(*Clamator glandarius*), and this has been used since to argue that anti-parasite behaviours have

a genetic basis[32]. Corroborating this, European Magpie populations across Europe vary in the

rate at which they reject model eggs depending on both geography and genetics[33]. However, a

more recent study, also using microsatellite markers, found no such association in Great Reed

Warblers (*Acrocephalus arundinaceus*) parasitised by Cuckoos[34]. The use of higher resolution

genetic data is therefore necessary to explore this question further, in light of prior mixed

evidence and the need for modern genomic approaches. The Reed Warbler/Cuckoo system is

a particularly valuable case study with which to do this, given the generally good understanding

of the traits involved in brood parasitism thanks to years of research. Although some studies of

parasite–host interactions more generally have found genomic regions putatively associated

with non-brood parasitism (e.g. Atlantic Salmon *Salmo salar* and ectoparasites[35]), others have

not (e.g. Red Grouse (*Lagopus scotica*) and the nematode *Trichostrongylus tenuis*[36]).

Reed Warbler and Cuckoo natural history[37,38], breeding behaviour[39,40], egg colouration[41,42],

predators[43,44], migration[45,46], taxonomy[47,48], and diet[49,50] are well understood and have been

subject to increasingly sophisticated experimental and observational research. The GMTC has

been proposed to explain the persistence of brood parasite / host coevolutionary dynamics[7],

and much of the existing body of work on these two species provides supporting evidence for

this[28,51]. On the other hand, certain aspects vital to the GMTC have so far not been assessed in

this system. These include fundamental components such as characterisation of gene flow

patterns at a continental scale, and assessment of genomic selection imposed by brood

parasitism. The putative geographic mosaic of Reed Warbler/Cuckoo interactions provides a

valuable opportunity to evaluate the genomic consequences of variation in brood parasitism

risk across a host's breeding range, where breeding habitats are scattered with varying levels

of isolation from one another, and varying levels of allopatry/sympatry with Cuckoos of the

host-specific gens (i.e. the Reed Warbler-specific race of the Cuckoo). Sites at which Reed

Warblers and Cuckoos regularly interact, or coevolutionary 'hot spots', have been monitored

for long periods of time e.g.[28]. On the other hand, there are also 'cold spots' where the two

species very rarely or never interact (e.g. regions in Finland colonised by Reed Warblers within

the past century[52]). The GMTC predicts that this could lead to relaxation or even reversal of

anti-parasite defences in such 'cold spots'. Although behavioural defences (such as egg

rejection) can rescue hosts that are in sympatry with parasitic Cuckoos, accidental rejection of

one's own egg is costly and decreases host fitness[53]. Thus, where they are in allopatry with

Cuckoos, Reed Warblers could potentially lose resistance from fitness costs or drift[54]. Re-

encounters are predicted to promote cyclic trajectories[7]. Cyclic changes across the Reed

Warbler/Cuckoo range mosaic could retain the genetic variation needed for adaptation (as in

sticklebacks *Gasterosteus aculeatus*, where metapopulation structuring maintains enough

genetic variation to fuel novel colonisations[55]).

Our work, combining behavioural and genomic information in a well-studied natural system,

represents an exciting application of this approach. Using a restriction site associated (RAD)

sequencing approach, we characterise gene flow among host populations to assess whether we

see patterns of variable gene flow that are a fundamental component of the GMTC.

Furthermore, using genotype-environment association analyses, we test for signals of selection

on biotic interactions by identifying SNP (single nucleotide polymorphism) loci that associate

with Cuckoo parasitism rates as sites across continental Europe, controlling for environmental

(i.e. climatic) factors. Overall, this study provides further evidence for the existence of a

geographic mosaic of coevolution in an avian-cuckoo system, and highlights where future

efforts could focus to provide a more complete characterisation of the GMTC.

Material and methods

Data collection

Reed Warblers' anti-Cuckoo defences vary in extent according to the level of Cuckoo parasitism the population is exposed to[54]. The species' range is also expanding to the north[52] and, at least in certain regions, to the south[56]. Our sampling sites form a latitudinal and longitudinal gradient across much of Europe, and include locations with known range expansion histories. Birds were caught and blood sampled as part of prior projects[57-59]. The inclusion of Reed Warblers from sites across much of their European breeding range enables comparisons among sites which vary in parasitism level. It also reduces the impact of other genetic or demographic effects (e.g. allele surfing[60,61]) as well as from selection imposed by other forms of environmental variation that commonly impact genomic evolution (e.g. climate[62]), on our inferences.

We collated spatial data on brood parasitism rate, making use of a combination of existing literature[63] and information provided by academic and citizen scientist collaborators. For some sites, we also had access to the rate at which Reed Warblers at that site reject a 'foreign' egg from their nest (Supplementary Table S1). All 'parasitised' sites had a parasitism rate (i.e. percentage of parasitised nests) of > 5%. All 'unparasitised' sites had an observed parasitism rate of 0% during the period in which the samples were collected (although parasitism may occur very rarely – e.g. in Helsinki, an 'unparasitised' Reed Warbler population, two nests were found to be parasitised by a Cuckoo in 2023, out of 667 nests monitored over six years (pers. obs. RT, KR, EK, WJS)). For two sampling sites, parasitism status was unknown (Figure 1A).

Fig. 1: (a) We used DNA samples from Reed Warblers at 22 sites. Red points represent sites where Reed Warblers are not parasitised by Cuckoos. Blue points represent locations where they are. Grey points represent sites where brood parasitism rate is not known. Location codes are indicated (see Supplementary Table S1). **(b)** Principal Component Analysis reveals genetic differences across the Reed Warbler range. PC1 shows a north-south gradient of genetic differences, and PC2 shows differentiation amongst southern European birds. The variance explained by PC1 and PC2 is shown in parentheses. Colours of points depict sampling regions. Sampling regions are ordered latitudinally in the figure legend. Location codes correspond to sites depicted in (A). **(c)** Effective migration surfaces inferred by EEMS. Orange regions have reduced gene flow, and blue regions have higher gene flow. The scale represents effective

migration rate on log10 scale, relative to overall migration. A value of '1' denotes effective

migration that is 10 times faster than average. Gene flow is higher in central Europe, with lower

gene flow occurring in the Mediterranean and Baltic regions. Sampling locations are indicated.

Genomic data generation and pre-processing

Genomic DNA extraction and RAD-sequencing was carried out as detailed in prior work[57,58].

We generated and filtered two genotype files, one for use in population genetic structure and

gene flow analyses, and another for use in genotype-environment analyses (GEA). Both files

included both autosomes and sex chromosomes. For both datasets, we scanned for individuals

with high relatedness (see[64]) (none of the pairs of individuals had relatedness of > 0.1, and so

no individuals were excluded). To calculate this information, we used VCFtools v0.1.17[65] --

relatedness. We split Finnish samples into three groups called 'West', 'Middle', and 'East'

Finland. The subsequent 'Dataset 1' included 188 individuals from 22 sampling sites. For the

genotype file used in GEA pipelines, 'Dataset 2', we excluded sampling sites for which there

was no brood parasitism data (Sweden and one of the Czech sites) and split the Finnish samples

in the same way as in Dataset 1. Dataset 2 included 177 individuals from 20 sampling sites.

Full details on the genomic datasets are available in Supplementary Table S1.

For both Dataset 1 and Dataset 2, we used STACKS v2.65[66] to sort, filter, and demultiplex

reads with the *process_radtags* script. We aligned reads to the bAcrSci1 reference genome[67]

with bwa v0.7.17[68]. We used SAMTOOLS v1.18 to sort files[69]. Following this, we used the

ref_map.pl pipeline to build the initial catalogue to be used as an input for *populations*. Using

VCFtools, we filtered by depth, retaining sites with mean depth of more than or equal to 10,

and less than or equal to 100. We then assessed sites for Hardy-Weinberg Equilibrium (using

an exact test (--hwe), see[70]) and filtered to exclude SNPs out of HWE in all populations with

more than ten individuals. Using *populations*, we selected sites where all individuals were

represented (-R 1), where there was a minimum minor allele count of 3 (--min-mac 3), and

where there was a maximum observed heterozygosity of 0.75 (--max-obs-het 0.75). We

restricted analysis to the first SNP of each locus (--write-single-snp). For Dataset 1, we used

PLINK v2.00[71] to detect SNP pairs with linkage disequilibrium of 0.5, and then exclude one

of them (--indep-pairwise 50 5 0.5), retaining 13,208 SNPs ('Dataset 1') for our analyses. For

Dataset 2, we did not filter for linkage disequilibrium as this would risk removing parts of the

genome under strong selection. Hence, we retained 13,346 SNPs.

Population genetic structure and gene flow analyses

We first carried out PCA with PLINK v2.00[71], as well as ADMIXTURE v1.3.0[72], ranging the

number of assumed populations between 1 and 22. We then examined gene flow among

sampling sites. To do so, we used EEMS v0.0.0.9, which makes use of geo-referenced genetic

samples to detect regions of higher- and lower-than-average gene flow and has been simulated

to accurately recover effective migration surfaces (visualised gene flow) in scenarios with

complicated histories of gene flow[73]. We first converted the Dataset 1 VCF into PLINK input

files using the –make-bed command. We then used the bed2diffs_script to convert these into a

matrix of average pairwise genetic dissimilarities, which was used as the input file for EEMS.

Following this, we carried out 8 runs each with an MCMC chain of 10,000,000 iterations, with

a 20,000-iteration burn-in and 1800 thinning iterations. The number of geographic demes was

set to 100, 300, 500, 700, 900, 1100, 1300 and 1500, and runs were combined using

rEEMSplots in R. Data is presented on a $\log_{10}$ scale, where an increase of one represents a

tenfold increase in migration rate relative to average levels. We also used VCFtools to output

mean pairwise F_{ST} values for each population pair.

Genotype-environment analysis

To explore genotype-environment associations we used three different GEA analytical

frameworks: univariate latent factor mixed modelling (LFMM)[74], multivariate redundancy

analysis (RDA) and partial redundancy analysis (pRDA)[75], and the F_{ST}-based method

BayeScEnv[76]. We used the vcf2geno command from the R package *LEA*[77] to convert the

relevant VCF files into suitable input files. Finally, we estimated additive polygenic scores to

determine the cumulative signal of selection across candidate loci identified across all GEA

methods as in[78,79]. Parasitism rate tends to be highest in the centre of Europe, and lowest in its

northern and southern parts[80]. To summarise climatic variation, we downloaded the 19 BioClim

climate variables[81]. To avoid high correlations amongst the variables, we calculated Pearson's

correlation coefficient in R and selected four variables which had correlation coefficients of <

0.7 across our sites. These were temperature seasonality (BIO4), mean temperature of the

wettest quarter (BIO8), precipitation of the wettest month (BIO13), and precipitation

seasonality (BIO15). The latter was particularly important to include as it was identified as an

important contributor to genetic variation in European populations of the Reed Warbler in prior

work[57]. Measures of precipitation are important to include given the significance of rainfall in

determining water level and therefore reedbed growth (i.e. Reed Warbler habitat availability

and quality[82]). We also included latitude in our models, to account for neutral geographical

variation. Both LFMM and RDA/pRDA are run with parasitism rate, climatic variables, and

latitude together. BayeScEnv is run individually for each variable.

We carried out LFMM with the *lfmm* v2 package in R[83]. This uses an MCMC algorithm to

explore associations between allele frequency and environmental variables. We included

parasitism status, our four selected BioClim variables, and latitude as our environmental

variables. We then ran LFMM using *lfmm_ridge* to estimate the model, and *lfmm_test* to assess

the significance of the SNPs. The number of latent factors was chosen based on our population

genetic structure analyses (allowing us to control for neutral genomic processes e.g. due to

colonisation history). We selected K = 1 as we inferred limited differentiation across Europe

from the PCA, ADMIXTURE, and F_{ST} results. We then calculated the genomic inflation factor

(GIF)[84,85]. We confirmed that GIF values were close to 1, demonstrating that population

structure was adequately controlled for. LFMM then adjusts the p-values by the GIF values,

and we applied the Benjamini-Hochberg procedure[86] to tolerate a false discovery rate of 10%,

to return significant candidate loci.

Following LFMM, we carried out RDA, which models relationships between environmental

predictors and genomic variation (i.e. the SNPs). RDA is a constrained ordination and has been

demonstrated to outperform other GEA analyses when selection acts over multiple loci[75]. We

carried out RDA using the *vegan* R package's *rda* function. We calculated adjusted R^2 (based

on the number of predictors), before assessing variance explained by each eigenvalue. We

assessed the RDA model for significance using F-statistics via an analysis of variance

(ANOVA) with 999 permutations. We assessed both the full model and each constrained axis.

We also assessed the Variance Inflation Factors of each predictor variable, confirming that they

are below ten, and therefore that collinearity amongst predictors is not a significant problem.

We identified candidate SNPs on significant constrained axes based on their loadings in

ordination space, using a cut-off of ±3 SD (see[87]), and removed duplicate candidates. We

repeated this in a 'partial RDA' context, which accounts for population genomic structure.

Applying both RDA approaches provides a more holistic assessment of SNPs under selection

according to variation in levels of brood parasitism.

Finally, we used BayeScEnv v1.1[76]. BayeScEnv does not directly assess associations between

allele frequencies and environmental variables, but rather environmental and genetic

differentiation (i.e. F_{ST}), providing an additional GEA test. We used VCFR[88] to convert the

VCF file to a genind object (of the adegenet R package[89]), and then used hierfstat[90] to convert

this into an input file. We divided parasitism rate at each site by the standard deviation, as

recommended by the BayeScEnv authors[76]. We then ran BayeScEnv with twenty pilot runs of

length 5000, a thinning interval size of ten, 5000 outputted iterations, a burn-in length of

50,000, prior probability against the neutral model of 0.5, and prior preference for the locus-

specific effect model of 0.5. We identified SNPs with a posterior error probability (see[91,92]) of

< 0.01 as candidate loci. We repeated BayeScEnv using the four Bioclim variables and latitude

as the focal parasitism variable, to identify any common candidate loci.

We filtered the annotation file for the bAcrSci1 genome to identify candidate genes from our

outlier SNPs (i.e. SNPs that fall within a gene in the annotation file). Then, we (1) performed

Gene Ontology Enrichment Analysis (geneontology.org, Panther v18.0), with *Gallus gallus*

(the most complete avian genome annotation) as the reference set to identify whether genes

associated with specific biological processes were overrepresented, and (2) delineated the

cumulative impact of selection at the individual level, by calculating per-individual additive

polygenic scores (estimates of the cumulative signal of selection across candidate loci; referred

to henceforth as 'polygenic scores') (see[78,79]). We assessed the correlation between the

polygenic scores and parasitism rate to determine the extent to which the candidate alleles vary

with parasitism. We tested linear and quadratic models and selected the model with the lowest

Bayesian information criterion (BIC) value. We also assessed the correlation of polygenic

scores with the bioclimatic variables and latitude, as well as the rate at which Reed Warblers

at each site reject a foreign egg, again testing both linear and quadratic models.

Results

Population structure and gene flow analyses

PCA reveals genetic differentiation with geography, albeit without discrete groupings (Figure

1B) and overall low differences between populations (corroborating F_{ST} results; Supplementary

Table S2). PC1 reflects differentiation along a north-south gradient, whereas PC2 shows

differentiation within the southern European individuals. In particular, Turkish and Maltese

birds differ from those elsewhere in southern Europe. ADMIXTURE highlights similar

patterns. Although K = 1 is best supported, higher K values highlight some population

structure, with K = 2 splitting birds from Nordic regions from those from southern Europe, and

characterising birds from within the range core as genetically admixed individuals (Figure S1).

Higher K values identify genetic structure amongst the southern European birds. Gene flow

analysis depicts a varying degree of gene flow throughout Europe, with higher gene flow across

the core of the continent flanked by areas of lower gene flow in both the north and south (Figure

1C).

Genotype-environment associations

LFMM identified 11 candidate SNPs associated with parasitism rate (Supplementary Table S3)

and 129 SNPs associated with the other environmental variables (i.e. latitude or one of the four

bioclimatic variables). Whilst there was some SNP overlap amongst the five environmental

variables, the candidate parasitism SNPs were not identified as associated with any of these

other variables. The RDA model, as a global model, was significant to p = 0.001, and explained

0.9% of variance in the dataset (n.b. variance explained is typically low in RDA models

assessing the genetic impacts of environmental variation). The variance inflation factor (VIF)

between predictor variables was < 5, suggesting limited impact of collinearity on the RDA.

The first five axes were significant at p = 0.001 and together explained 86.03% of variation.

Using these five significant axes for outlier detection, we identified 41 SNPs as being

associated with parasitism, 86 with latitude, 30 with BIO4, 14 with BIO8, and 17 with BIO13

(Figure 2A). Partial RDA was also significant as a global model (p = 0.001) and explained

0.05% of the variance. The first four axes were significant to p = 0.001 and together explained

69.91% of variation. From this method, we identified 43 SNPs associated with parasitism, 78

with latitude, 29 with BIO4, 10 with BIO8, and 9 with BIO13 (Figure 2B). A total of 37

parasitism candidate SNPs were identified by both RDA and pRDA. Four parasitism candidate

SNPs were unique to RDA, and six to pRDA. Whilst the RDA/pRDA may be somewhat

confounded by BIO8 and BIO13 (Figure 2), all of the 11 candidate SNPs identified by LFMM

(which tested each variable in a univariate manner and did not identify overlap with the

parasitism SNPs) were also identified by both RDA and pRDA. Finally, BayeScEnv detected

2 parasitism candidate SNPs, and 26 SNPs associated with latitude or a bioclimatic variable.

One of the two parasitism candidate SNPs was also a candidate associating with BIO8 and

BIO15 (and there was extensive overlap amongst SNPs associating with the five environmental

variables, with 10 of the 26 being candidates for more than one environmental variable).

Neither of the BayeScEnv parasitism loci were identified by either LFMM or RDA/pRDA.

Together, LFMM, RDA/pRDA, and BayeScEnv detected 49 candidate SNPs (Figure 2C). We

identified a known gene associated with 24 of these candidate SNPs.

Fig. 2: Genotype–environment association with brood parasitism. **(a)** Redundancy analysis (RDA) using 13,346 SNPs, depicting the first two axes and using the following variables: parasitism, latitude, BIO4 (temperature seasonality), BIO8 (mean temperature of the wettest quarter), BIO13 (precipitation of the wettest month), and BIO15 (precipitation seasonality). SNP/individual scores are scaled by the square root of the eigenvalues to give appropriate weight to the axes based on their contribution to the model. SNP genotypes are represented as grey crosses, and the direction of variation in each constraining variable is indicated with black arrows (arrow length represents the strength of the correlation of that variable). Individuals are represented by different colours according to location. Percentages on the axes indicate the proportion of total variance explained. **(b)** Partial redundancy analysis (pRDA), accounting for population genetic structure. **(c)** Manhattan plot showing genotype–parasitism associations,

estimated with LFMM, across the genome. The red dot indicates the candidate SNPs identified

by LFMM itself (as well as RDA and pRDA). Blue dots represent the candidate SNPs identified

by RDA and/or pRDA. Pink dots represent candidate SNPs identified by BayeScEnv. Amongst

the candidate SNPs, those with the top 11 LFMM p-values are circled, and the top ten candidate

SNPs which are associated with known genes are annotated. Full gene names are presented in

Supplementary Table S3.

Gene ontology analysis of the SNPs associated with parasitism rate highlighted genes with a

diverse array of functions and associations, with overrepresentation of genes related to cell

communication, luteinising hormone secretion, tissue morphogenesis, and the cell surface

receptor protein tyrosine kinase signalling pathway. Specific genes (Supplementary Table S3)

included those with neurological (e.g. *nrp2*, *ctnna2,* rtn4rl1), metabolic (e.g. *cit*), and cell-cycle

related (e.g. *babam2, znf830*) associations. The additive polygenic scores for the 49 candidate

loci increased with parasitism rate at the location at which the individual was sampled ($n = 177$

individuals). Both the linear model ($F_{(1,175)} = 178.9$, adjusted $R^2 = 0.5027$, p < 0.05) and the

quadratic model were significant ($F_{(2,174)} = 188$, adjusted $R^2 = 0.68$, p < 0.05). The quadratic

model explained more variation than the linear model and had a lower BIC score (linear model

= 868.0, quadratic model = 794.2, ΔBIC = 73.86) (Figure 3).

Fig. 3: Correlation of brood parasitism rate and individual polygenic scores (an estimate of the

cumulative signal of selection across all the candidate loci). Each dot represents the polygenic

score for an individual Reed Warbler and the parasitism rate at its location. The solid line

depicts the regression line from the quadratic model. Points are jittered for clarity. Sampling

location is indicated by point colour (see Figure 1 for more information on sampling sites).

For all other models (looking at the variables other than parasitism, including rejection rate)

(Figure S2), the linear model was chosen as the BIC difference could not separate the quadratic

and the linear models, so the simpler was chosen. R-squared values for all of them were an

order of magnitude or more lower than for the relationship for parasitism.

Discussion

Our analyses provide two novel pieces of information which contribute to our understanding of the Reed Warbler–Cuckoo mosaic of coevolution. Firstly, we show that the level of gene flow varies within the European breeding range of the Reed Warbler. Generally, gene flow is higher within the range core and lower in the north and south of our sampling area – which is probably attributable to the Baltic and Adriatic seas acting as barriers to gene flow. This variation in gene flow also corresponds to general patterns in the intensity of brood parasitism, wherein parasitism rate is generally higher in the host's range core when compared to the range edge. Principal component analysis suggests some isolation by distance between core and edge populations, however the edge populations cluster as three connected groups in the PCA, largely corresponding to their geographical location. Differentiation of Turkish individuals could be related to subspecies-level admixture in this region[93]. Prior work, using microsatellite data, also identified restricted gene flow in southern Europe[94], corroborating our results. Low gene flow to the west of Romania might be attributable to the known migratory divide in this species[95]. Whilst Reed Warblers lack significant population genetic structure, they are known to have some geographic differentiation[47,57,96] (in comparison to many Palearctic bird species, which are panmictic[97]). This may facilitate the loss, gain or maintenance of genetic variation involved in responses to Cuckoo brood parasitism, allowing adaptation to parasitism to exhibit spatial variation at a continental scale. This would allow for a geographic mosaic of coevolution to be maintained.

Our second novel piece of evidence for a geographic mosaic of coevolution is the identification of genetic associations with Cuckoo parasitism. Using three different types of genotype-environment association analysis, we have identified candidate genes under selection that

highly correlate with parasitism. We have therefore described, for the first time, a continental

signature of genomic adaptation to brood parasitism pressure. Four of the identified genes (and

11 of the identified SNPs) were detected by multiple method types. The candidate genes

represent a diverse suite of physiological functions, as assessed by gene ontology analysis.

Among those previously studied in birds, several genes stand out as particularly interesting.

Notably, *nrp2*, which was identified as correlated with parasitism rate by RDA, partial RDA,

and LFMM, is associated with neurodevelopmental functions (potentially relating to vocal

learning in birds [98] and also to memory and learning in laboratory mice[99]). More generally,

ctnna2 is known to have a major role in the function and growth of neurons[100]. These findings

suggest a potential role for these genes in learning or communication, or at least some cognitive

response, in the defence against Cuckoo parasitism (perhaps relating to mobbing for example,

which is a key defence against Cuckoos at Reed Warbler nest sites[21]). Other candidate genes

have less obvious links to parasite responses. For example, *ppil2* is associated with immunity

(and was identified as involved in immune system changes during chicken moult[101]). Note that

there were no SNPs near the Ase64 microsatellite locus (found to be associated with egg

rejection capability in magpies[30]), precluding us from testing whether this region is associated

with parasitism rate in our study species. The closest SNP in our dataset was 7670 bp away,

and was not among the parasitism candidate SNPs.

Despite the associations found here, there are several caveats which are important to highlight.

Firstly, correlation does not necessarily mean causation. There will undoubtedly be various

abiotic and biotic correlates of parasitism in the Reed Warbler–Cuckoo system. Whilst we have

explored the potential confounding role of climatic variation (e.g. none of focal environmental

variables correlate with the parasitism polygenic scores), a well-known biotic correlate of

parasitism in this system is the increase in parasitism rate with host population density[63]. Our

candidate SNPs might therefore be impacted by selection according to local conspecific

density, or some correlate of this (e.g. disease, intraspecific competition). It is also known that

habitat structure plays a role in determining host defences and parasite adaptations[102,103]. The

identification of genes which are putatively involved in immunity and growth provides support

for this. Disentangling such factors represents a major challenge. Common garden experiments,

which have proven useful in studying the geographic mosaic theory of coevolution in plants

(e.g.[104,105]) are impractical for wild bird populations. A feasible alternative might involve

characterising the responses of individual Reed Warblers to Cuckoos (e.g. extent of mobbing

behaviour or egg rejection), and then using genomic data for these individuals to assess

differentiation at our candidate regions and across the genome. This would also allow for

characterisation of differential selection on male and female hosts. If carried out both within

and between different sites that differ in parasitism rates across the Reed Warbler's European

range, this could provide additional evidence for the role of these candidate genes in mediating

Reed Warbler–Cuckoo interactions. A second caveat is that, given that RAD sequencing data

covers only a small fraction of the genome, many of the candidate genes are likely to be in

linkage with genes that are adaptively responding to parasitism rate, rather than having direct

mechanistic roles. A cautionary example is provided by the gene responsible for the black coat

of some North American Grey Wolves (*Canis lupus*), which does not determine immunological

differences as was previously thought, but is linked to immunological genes[106]).

Whilst there is limited value in speculating about the exact functions of our candidate genes

and their potential links to parasitism, it is worth outlining potential pathways or adaptations in

which we would expect parasitism to underpin genetic differences. For example, genetic

differences could relate to the visual systems of the Reed Warbler and therefore contribute to

egg-rejection capacity, which is known to be a major component of the Reed Warbler's

antiparasitic defences[107]. There might also be a genetic underpinning to the willingness of Reed

Warblers to mob Cuckoos (perhaps relating to varying aggression amongst individuals). The

extent of mobbing is known to vary across Europe[21]. Nesting behaviour (i.e. how well-hidden

nests are) and phenology (i.e. timing of laying) are also likely to have a genetic component and

would have an impact on parasitism likelihood. Our parasitism polygenic scores do not

correlate with known rates of rejection of foreign eggs across sampling sites (Figure S2), so it

seems unlikely that these genes impact this particular anti-cuckoo behaviour. Overall, it is very

probable that all of these axes of phenotypic variation and others would be highly polygenic

and therefore will require further research to parse the contributions of different genes, and,

more broadly, the respective contributions of genetics and environment to parasitism response.

Whilst our results provide support for a genomic association with brood parasitism in the Reed

Warbler–Cuckoo system, a more complete characterisation of the geographic mosaic of

coevolution will require further research. Although our reduced-representation based

assessment provides a significant advance on attempts using microsatellite markers, analysis

of whole-genome sequence data in the Reed Warbler and other hosts would be invaluable in

providing a more fine-grain assessment of the architecture of genomic selection. This is

because it is well-known that using RAD-sequence data to infer local adaptation may lead to

candidate loci being missed[108]. As well as this, we have highlighted a need for better

understanding of parasitism status within the range of this host and others, given that some

locations do not have a known parasitism rate. Furthermore, the impact of human activity has,

in recent centuries, changed landscapes across Europe[109,110], likely affecting distributions and

abundance of the Cuckoo and its hosts. It would be fascinating to delineate the genomic impacts

of such changes, perhaps using museum specimens. Furthermore, attempting to extend this

type of study to the Cuckoo itself would be worthwhile. The existence of gentes, each

potentially subject to differing selection pressures, adds complexity to the Reed Warbler–

Cuckoo coevolutionary mosaic. Might we expect different genomic regions to be impacted by

selection in different Cuckoo lineages (gentes), due to variation in parasitism pressures across

host populations? Sampling of Cuckoos and their hosts across varying habitats and regions, at

continental scale, will be needed to resolve this.

Our characterisation of this antagonistic coevolutionary interaction represents the most in-

depth assessment of the genetic underpinnings of brood parasitism carried out to date. A similar

approach – using estimates of interaction strength – could hold promise in elucidating biotic

interactions more generally. For example, in tight predator–prey relationships, predation rate

could be used in much the same way as we have used brood parasitism. Whilst studies of

parasite–host interactions have made use of in-depth population genomic assessments since the

advent of widely available whole-genome data[35,36], population genomic research that has

involved the study of predation (and competition) has historically focused on understanding

changes in overall levels of genetic variation[111]. Genotype–environment analysis would,

however, allow for the identification of candidate genes involved in this ecologically

fundamental biotic interaction, if combined with a recognition that genes involved in

behavioural expression are likely to work in a polygenic fashion, meaning that their individual

effects are small. Such approaches are being increasingly used as part of wider population

genetic studies (e.g.[112]) (as well as gene expression-based assessments[113] or experimental

approaches[114] which have been deployed in laboratory settings). More broadly, the application

of genomic research to study behaviour in well-studied systems such as humans[115] and mice[116]

is developing rapidly. Nevertheless, genotype–environment analyses deserve specific attention

in non-model systems in the wild, especially in tightly linked predator–prey species pairs which

are known to experience cyclical demographic changes. Overall, our work highlights that, even

with traits which are likely to be highly polygenic or involve significant plastic elements,
genotype–environment analyses offer opportunities to significantly improve our understanding
of the genetic underpinnings of biotic interactions, especially when deployed alongside a
thorough grounding in the natural history and field biology of the species in question.

In conclusion, this investigation represents a significant step forward in understanding the
geographic mosaic of coevolution in the Reed Warbler–Cuckoo system. We not only provide
support for variable gene flow across the European range of the Reed Warbler, but also, for the
first time, identify putative signature of genomic selection exerted by brood parasitism. This
work highlights the need for additional characterisation of the genetic and environmental
factors responsible for this antagonistic coevolutionary interaction.

574
Acknowledgements: We are grateful to those who supplied information on egg rejection and
brood parasitism rates in prior works. These include Geir Rudolfsen, Inge Hafstad, Bruno
Bargain, Josef Beier, David Bigas Campàs, Bernd Leisler, Péter L. Pap, Ričardas Patapavičius,
Karl Schulze-Hagen, Johannes Bang, Bernd Leisler, Jesper J. Madsen, Anders Pape Møller,
Andrzej Dyrcz, Marcel Honza, Vytautas Jusys, Eivin Røskaft, and Arne Moksnes. We
acknowledge the use of the CSC – IT Center for Science, Finland, for computational resources.
BGS was funded by the Research Council of Norway, grant no. 151641/432. The work was
funded by Academy of Finland project grant number 333801 and HiLIFE Fellows grant to RT.

Ethics statement: All blood sampling was conducted under permission from animal ethics
experimentation boards and with permission to derogate from wildlife protection laws in
respective countries. In Finland these were granted by the Project Authorization Board of the
Regional State Administrative Agency (ESAVI/7857/2018) and the Centre for Economic

Development, Transport and the Environment (VARELY/758/2018). See references 57-59 for

a list of permissions for other locations.

Data availability statement:

Data and code will be made available on publication.

Author contributions:

Conceptualisation: Rose Thorogood, Katja Rönkä, William J. Smith. **Data curation**: William

J Smith, Katja Rönkä, Rose Thorogood, Nora Bergman. **Formal analysis**: William J. Smith,

Katja Rönkä. **Funding acquisition**: Rose Thorogood. **Investigation**: William J. Smith, Katja

Rönkä, Michał T. Jezierski. **Methodology**: William J. Smith, Kata Rönkä, Michał T. Jezierski.

Project administration: William J. Smith, Katja Rönkä, Rose Thorogood. **Resources**: Rose

Thorogood, Katja Rönkä, Nora Bergman, Edward Kluen, Petr Procházka, Bård Stokke, Bengt

Hansson. **Supervision**: Rose Thorogood. **Validation**: William J. Smith, Katja Rönkä, Michał

T. Jezierski. **Visualisation**: William J. Smith, Michał T. Jezierski, Katja Rönkä, Rose

Thorogood. **Writing – original draft**: William J. Smith, Katja Rönkä, Rose Thorogood.

Writing – review & editing: William J. Smith, Katja Rönkä, Staffan Bensch, Nora Bergman,

Rachael Y. Dudaniec, Fabrice Eroukhmanoff, Frode Fossøy, Bengt Hansson, Michał T.

Jezierski, Edward Kluen, Petr Procházka, Camilla Lo Cascio Sætre, Bård G. Stokke, & Rose

Thorogood.

611
612
613

References

Paterson, S. *et al.* Antagonistic coevolution accelerates molecular evolution. *Nature* **464**, 275-278, doi:10.1038/nature08798 (2010).

Yamamichi, M. & Ellner, S. P. Antagonistic coevolution between quantitative and Mendelian traits. *Proc Biol Sci* **283**, 20152926, doi:10.1098/rspb.2015.2926 (2016).

Laine, A-L. Role of coevolution in generating biological diversity: spatially divergent selection trajectories. *Journal of Experimental Botany* **60**, 2957-2970, doi:10.1093/jxb/erp168 (2009).

Ebert, D. Open questions: what are the genes underlying antagonistic coevolution? *BMC Biol* **16**, 114, doi:10.1186/s12915-018-0583-7 (2018).

Thomas, M. A. & Klaper, R. Genomics for the ecological toolbox. *Trends in Ecology & Evolution* **19**, 439-445, doi:10.1016/j.tree.2004.06.010 (2004).

Živković, D., John, S., Verin, M., Stephan, W. & Tellier, A. Neutral genomic signatures of host-parasite coevolution. *BMC Evolutionary Biology* **19**, 230, doi:10.1186/s12862-019-1556-3 (2019).

Thompson, J. N. *The Geographic Mosaic of Coevolution*. (University of Chicago Press, 2005).

Gomulkiewicz, R., Thompson, J., Holt, R., Nuismer, S. & Hochberg, M. Hot Spots, Cold Spots, and the Geographic Mosaic Theory of Coevolution. *The American naturalist* **156**, 156-174, doi:10.1086/303382 (2000).

Gomulkiewicz, R. *et al.* Dos and don'ts of testing the geographic mosaic theory of coevolution. *Heredity (Edinb)* **98**, 249-258, doi:10.1038/sj.hdy.6800949 (2007).

Kaur, R. *et al.* Ant behaviour and brain gene expression of defending hosts depend on the ecological success of the intruding social parasite. *Philosophical Transactions of the Royal Society B: Biological Sciences* **374**, 20180192, doi:10.1098/rstb.2018.0192 (2019).

Springer, S. A., Crespi, B. J. & Swanson, W. J. Beyond the phenotypic gambit: molecular behavioural ecology and the evolution of genetic architecture. *Molecular Ecology* **20**, 2240-2257 (2011).

Lindholm, A. K. & Thomas, R. J. Differences between Populations of Reed Warblers in Defences against Brood Parasitism. *Behaviour* **137**, 25-42 (2000).

Lank, D. B., Smith, C. M., Hanotte, O., Burke, T. & Cooke, F. Genetic polymorphism for alternative mating behaviour in lekking male ruff *Philomachus pugnax*. *Nature* **378**, 59-62, doi:10.1038/378059a0 (1995).

Saino, N. *et al.* Polymorphism at the Clock gene predicts phenology of long-distance migration in birds. *Molecular Ecology* **24**, 1758-1773 (2015).

Recuerda, M. *et al.* Adaptive phenotypic and genomic divergence in the common chaffinch (*Fringilla coelebs*) following niche expansion within a small oceanic island. *Journal of Evolutionary Biology* **36**, 1226-1241, doi:10.1111/jeb.14200 (2023).

Clark, J. D., Benham, P. M., Maldonado, J. E., Luther, D. A. & Lim, H. C. Maintenance of local adaptation despite gene flow in a coastal songbird. *Evolution* **76**, 1481-1494, doi:10.1111/evo.14538 (2022).

Rothstein, S. I. A Model System for Coevolution: Avian Brood Parasitism. *Annual Review of Ecology and Systematics* **21**, 481-508 (1990).

Thorogood, R., Spottiswoode, C. N., Portugal, S. J. & Gloag, R. The coevolutionary biology of brood parasitism: a call for integration. *Philosophical Transactions of the Royal Society B: Biological Sciences* **374**, 20180190, doi:10.1098/rstb.2018.0190 (2019).

Langmore, N. E. *et al.* Coevolution with hosts underpins speciation in brood-parasitic cuckoos. *Science* **384**, 1030-1036, doi:10.1126/science.adj3210 (2024).

Rothstein, S. I. A Model System for Coevolution: Avian Brood Parasitism. *Annual Review of Ecology and Systematics* **21**, 481-508, doi:10.1146/annurev.es.21.110190.002405 (1990).

Tolman, D., Campobello, D., Rönkä, K., Kluen, E. & Thorogood, R. Reed Warbler Hosts Do Not Fine-Tune Mobbing Defenses During the Breeding Season, Even When Cuckoos Are Rare. *Frontiers in Ecology and Evolution* **9**, doi:10.3389/fevo.2021.725467 (2021).

Soler, M. *Avian Brood Parasitism: Behaviour, Ecology, Evolution and Coevolution*. (Springer International Publishing, 2018).

Stokke, B. G., Fossøy, F., Røskaft, E. & Moksnes, A. in *Avian Brood Parasitism: Behaviour, Ecology, Evolution and Coevolution* (ed. Manuel Soler) 363-384 (Springer International Publishing, 2017).

Davies, N. B. & Quinn, D. *Cuckoos, Cowbirds and Other Cheats*. (Bloomsbury Publishing Plc, 2011).

Gibbs, H. L. *et al.* Genetic evidence for female host-specific races of the common cuckoo. *Nature* **407**, 183-186, doi:10.1038/35025058 (2000).

Fossøy, F. *et al.* Ancient origin and maternal inheritance of blue cuckoo eggs. *Nature Communications* **7**, 10272, doi:10.1038/ncomms10272 (2016).

Merondun, J. *et al.* Evolution and genetic architecture of sex-limited polymorphism in cuckoos. *Science Advances* **10**, eadl5255, doi:10.1126/sciadv.adl5255.

Stoddard, M. C. & Kilner, R. M. The past, present and future of 'cuckoos versus reed warblers'. *Animal Behaviour* **85**, 693-699 (2013).

715   29   Bay, R. A. *et al.* Predicting Responses to Contemporary Environmental Change Using
716   Evolutionary Response Architectures. *The American Naturalist* **189**, 463-473,
717   doi:10.1086/691233 (2017).
718

719   30   MartÍN-GÁLvez, D. *et al.* A quantitative trait locus for recognition of foreign eggs in
720   the host of a brood parasite. *Journal of Evolutionary Biology* **19**, 543-550 (2006).
721

722   31   Martín-Gálvez, D. *et al.* Cuckoo parasitism and productivity in different magpie
723   subpopulations predict frequencies of the 457bp allele: a mosaic of coevolution at a
724   small geographic scale. *Evolution* **61**, 2340-2348, doi:10.1111/j.1558-
725   5646.2007.00194.x (2007).
726

727   32   Ruiz-Raya, F. Ecophysiology of egg rejection in hosts of avian brood parasites: new
728   insights and perspectives. *Current Zoology* **67**, 631-638, doi:10.1093/cz/zoab042
729   (2021).
730

731   33   Soler, J. J., Martinez, J. G., Soler, M. & Møller, A. P. Genetic and geographic
732   variation in rejection behavior of cuckoo eggs by European Magpie populations: an
733   experimental test of rejector-gene flow. *Evolution* **53**, 947-956, doi:10.1111/j.1558-
734   5646.1999.tb05388.x (1999).
735

736   34   Procházka, P. *et al.* Host genotype and age have no effect on rejection of parasitic
737   eggs. *Naturwissenschaften* **101**, 417-426, doi:10.1007/s00114-014-1171-5 (2014).
738

739   35   Zueva, K. J. *et al.* Footprints of directional selection in wild Atlantic salmon
740   populations: evidence for parasite-driven evolution? *PLoS One* **9**, e91672,
741   doi:10.1371/journal.pone.0091672 (2014).
742

743   36   Wenzel, M. A., Douglas, A., James, M. C., Redpath, S. M. & Piertney, S. B. The role
744   of parasite-driven selection in shaping landscape genomic structure in red grouse
745   (*Lagopus lagopus scotica*). *Molecular Ecology* **25**, 324-341 (2016).
746

747   37   Moksnes, A., Fossøy, F., Røskaft, E. & Stokke, B. Reviewing 30 years of studies on
748   the Common Cuckoo - accumulated knowledge and future perspectives. *Chinese
749   Birds* **4**, 3-14, doi:10.5122/cbirds.2013.0001 (2013).
750

751   38   Pearson, D. J., Small, B. J. & Kennerley, P. R. Eurasian Reed Warbler: The characters
752   and variation associated with the Asian form fuscus. **95**, 42-61 (2002).
753

754   39   Moksnes, A. *et al.* Behavioural Responses of Potential Hosts Towards Artificial
755   Cuckoo Eggs and Dummies. *Behaviour* **116**, 64-89 (1991).
756

757   40   Wang, L., Yang, C., He, G., Liang, W. & Møller, A. P. Cuckoos use host egg number
758   to choose host nests for parasitism. *Proceedings of the Royal Society B: Biological
759   Sciences* **287**, 20200343, doi:doi:10.1098/rspb.2020.0343 (2020).
760

761   41   Avilés, J. M., Stokke, B. G., Moksnes, A., Røskaft, E. & Møller, A. P. Environmental
762   conditions influence egg color of reed warblers *Acrocephalus scirpaceus* and their

763   parasite, the common cuckoo *Cuculus canorus*. *Behavioral Ecology and Sociobiology*
764   **61**, 475-485, doi:10.1007/s00265-006-0275-0 (2007).
765

766 42   Stoddard, M. C. & Stevens, M. Avian vision and the evolution of egg color mimicry
767   in the common cuckoo. *Evolution* **65**, 2004-2013 (2011).
768

769 43   Jelínek, V., Karasová, T., Weidinger, K., Procházka, P. & Honza, M. Do common
770   cuckoo chicks suffer nest predation more than host nestlings? *Behavioral Ecology and*
771   *Sociobiology* **70**, 1975-1987, doi:10.1007/s00265-016-2203-2 (2016).
772

773 44   Kameníková, M., Navrátil, J. & Rajchard, J. Dependence of clutch predation rate of
774   Eurasian reed warbler *Acrocephalus scirpaceus* on nesting site selection: a model
775   study. *Biologia* **71**, 452-456, doi:10.1515/biolog-2016-0053 (2016).
776

777 45   Buchan, C. *et al.* Combining remote sensing and tracking data to quantify species'
778   cumulative exposure to anthropogenic change. *Global Change Biology* **29**, 6679-6692
779   (2023).
780

781 46   Procházka, P. *et al.* Delineating large-scale migratory connectivity of reed warblers
782   using integrated multistate models. *Diversity and Distributions* **23**, 27-40 (2017).
783

784 47   Olsson, U. *et al.* Mitochondrial phylogeny of the Eurasian/African reed warbler
785   complex (*Acrocephalus*, Aves). Disagreement between morphological and molecular
786   evidence and cryptic divergence: A case for resurrecting *Calamoherpe ambigua*
787   Brehm 1857. *Molecular Phylogenetics and Evolution* **102**, 30-44 (2016).
788

789 48   Wei, C. *et al.* Geographic variation in the calls of the Common Cuckoo (*Cuculus*
790   *canorus*): isolation by distance and divergence among subspecies. *Journal of*
791   *Ornithology* **156**, 533 - 542 (2015).
792

793 49   Ientile, R. *et al.* Analysis of Prey Composition in Eurasian Reed Warblers
794   *Acrocephalus scirpaceus* Droppings at Four Breeding Sites in Italy. *Diversity* **14**,
795   1134 (2022).
796

797 50   Mills, L. J. *et al.* Using molecular and crowd-sourcing methods to assess breeding
798   ground diet of a migratory brood parasite of conservation concern. *Journal of Avian*
799   *Biology* **51** (2020).
800

801 51   Lindholm, A. K. Brood parasitism by the cuckoo on patchy reed warbler populations
802   in Britain. *Journal of Animal Ecology* **68**, 293-309 (1999).
803

804 52   Fransson, T. & Stolt, B.-O. Migration routes of North European Reed Warblers
805   *Acrocephalus scirpaceus*. *Ornis Svecica* **15**, 153-160, doi:10.34080/os.v15.22735
806   (2005).
807

808 53   Stokke, B. G., Takasu, F., Moksnes, A. & Røskaft, E. The Importance of Clutch
809   Characteristics and Learning for Antiparasite Adaptations in Hosts of Avian Brood
810   Parasies. *Evolution* **61**, 2212-2228 (2007).
811

812 54 Thorogood, R. & Davies, N. B. Reed warbler hosts fine-tune their defenses to track
813   three decades of cuckoo decline. *Evolution* **67**, 3545-3555, doi:10.1111/evo.12213
814   (2013).
815

816 55 Lescak, E. A. *et al.* Evolution of stickleback in 50 years on earthquake-uplifted
817   islands. *Proceedings of the National Academy of Sciences* **112**, E7204-E7212,
818   doi:10.1073/pnas.1512020112 (2015).
819

820 56 Lo Cascio Sætre, C. *et al.* Rapid adaptive phenotypic change following colonization
821   of a newly restored habitat. *Nature Communications* **8**, 14159,
822   doi:10.1038/ncomms14159 (2017).
823

824 57 Sætre, C. L. C. *et al.* Genomic consequences of range expansion and colonisation in
825   the reed warbler (*Acrocephalus scirpaceus*). *bioRxiv*, 2022.2011.2028.518135,
826   doi:10.1101/2022.11.28.518135 (2022).
827

828 58 Bergman, N. *et al.* Conserved genetic diversity and no persisting founder effects
829   following rapid range expansion of a long-distance migrant (*Acrocephalus
830   scirpaceus*). *bioRxiv* (2025)
831

832 59 Procházka, P. *et al.* Low genetic differentiation among reed warbler *Acrocephalus
833   scirpaceus* populations across Europe. *Journal of Avian Biology* **42**, 103-113 (2011).
834

835 60 Travis, J. M. J. *et al.* Deleterious Mutations Can Surf to High Densities on the Wave
836   Front of an Expanding Population. *Molecular Biology and Evolution* **24**, 2334-2343,
837   doi:10.1093/molbev/msm167 (2007).
838

839 61 Rougemont, Q., Leroy, T., Rondeau, E. B., Koop, B. & Bernatchez, L. Allele surfing
840   causes maladaptation in a Pacific salmon of conservation concern. *PLOS Genetics* **19**,
841   e1010918, doi:10.1371/journal.pgen.1010918 (2023).
842

843 62 Lancaster, L. T. *et al.* Understanding climate change response in the age of genomics.
844   *Journal of Animal Ecology* **91**, 1056-1063 (2022).
845

846 63 Stokke, B. G. *et al.* Host density predicts presence of cuckoo parasitism in reed
847   warblers. *Oikos* **116**, 913-922 (2007).
848

849 64 Yang, J. *et al.* Common SNPs explain a large proportion of the heritability for human
850   height. *Nature Genetics* **42**, 565-569, doi:10.1038/ng.608 (2010).
851

852 65 Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156-
853   2158, doi:10.1093/bioinformatics/btr330 (2011).
854

855 66 Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A. & Cresko, W. A. Stacks: an
856   analysis tool set for population genomics. *Molecular ecology* **22**, 3124-3140,
857   doi:10.1111/mec.12354 (2013).
858

859 67 Sætre, C. L. C. *et al.* A Chromosome-Level Genome Assembly of the Reed Warbler
860   (*Acrocephalus scirpaceus*). *Genome Biology and Evolution* **13**, evab212,
861   doi:10.1093/gbe/evab212 (2021).

863 68 Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler
864   transform. *Bioinformatics* **25**, 1754-1760, doi:10.1093/bioinformatics/btp324 (2009).
865

866 69 Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
867   2078-2079, doi:10.1093/bioinformatics/btp352 (2009).
868

869 70 Wigginton, J. E., Cutler, D. J. & Abecasis, G. R. A note on exact tests of Hardy-
870   Weinberg equilibrium. *Am J Hum Genet* **76**, 887-893, doi:10.1086/429864 (2005).
871

872 71 Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-
873   Based Linkage Analyses. *The American Journal of Human Genetics* **81**, 559-575,
874   doi:10.1086/519795 (2007).
875

876 72 Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry
877   in unrelated individuals. *Genome Res* **19**, 1655-1664, doi:10.1101/gr.094052.109
878   (2009).
879

880 73 Petkova, D., Novembre, J. & Stephens, M. Visualizing spatial population structure
881   with estimated effective migration surfaces. *Nat Genet* **48**, 94-100,
882   doi:10.1038/ng.3464 (2016).
883

884 74 Caye, K., Jumentier, B., Lepeule, J. & François, O. LFMM 2: Fast and Accurate
885   Inference of Gene-Environment Associations in Genome-Wide Studies. *Mol Biol Evol*
886   **36**, 852-860, doi:10.1093/molbev/msz008 (2019).
887

888 75 Capblancq, T. & Forester, B. R. Redundancy analysis: A Swiss Army Knife for
889   landscape genomics. *Methods in Ecology and Evolution* **12**, 2298-2309 (2021).
890

891 76 de Villemereuil, P. & Gaggiotti, O. E. A new FST-based method to uncover local
892   adaptation using environmental variables. *Methods in Ecology and Evolution* **6**, 1248-
893   1258 (2015).
894

895 77 Frichot, E. & François, O. LEA: An R package for landscape and ecological
896   association studies. *Methods in Ecology and Evolution* **6**, 925-929 (2015).
897

898 78 Babin, C., Gagnaire, P-A., Pavey, S. A. & Bernatchez, L. RAD-Seq Reveals Patterns
899   of Additive Polygenic Variation Caused by Spatially-Varying Selection in the
900   American Eel (*Anguilla rostrata*). *Genome Biology and Evolution* **9**, 2974-2986,
901   doi:10.1093/gbe/evx226 (2017).
902

903 79 Xuereb, A., Kimber, C. M., Curtis, J. M. R., Bernatchez, L. & Fortin, M-J. Putatively
904   adaptive genetic variation in the giant California sea cucumber (*Parastichopus
905   californicus*) as revealed by environmental association analysis of restriction-site
906   associated DNA sequencing data. *Molecular Ecology* **27**, 5035-5048 (2018).
907

908 80 Stokke, B. G. *et al.* Predictors of resistance to brood parasitism within and among
909   reed warbler populations. *Behavioral Ecology* **19**, 612-620,
910   doi:10.1093/beheco/arn007 (2008).
911

Busby, J. R. BIOCLIM - a bioclimate analysis and prediction system. (1991).

Higgisson, W., Cobb, A., Tschierschke, A. & Dyer, F. The Role of Environmental Water and Reedbed Condition on the Response of *Phragmites australis* Reedbeds to Flooding. *Remote Sensing* **14**, 1868 (2022).

Caye, K., Jumentier, B., Lepeule, J. & François, O. LFMM 2: Fast and Accurate Inference of Gene-Environment Associations in Genome-Wide Studies. *Molecular Biology and Evolution* **36**, 852-860, doi:10.1093/molbev/msz008 (2019).

Yang, J. *et al.* Genomic inflation factors under polygenic inheritance. *European Journal of Human Genetics* **19**, 807-812 (2011)

Devlin, B. & Roeder, K. Genomic control for association studies. *Biometrics* **55**, 997-1004, doi:10.1111/j.0006-341x.1999.00997.x (1999).

Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B (Methodological)* **57**, 289-300 (1995).

Forester, B. R., Lasky, J. R., Wagner, H. H. & Urban, D. L. Comparing methods for detecting multilocus adaptation with multivariate genotype–environment associations. *Molecular Ecology* **27**, 2215-2233 (2018).

Knaus, B. J. & Grünwald, N. J. vcfr: a package to manipulate and visualize variant call format data in R. *Molecular Ecology Resources* **17**, 44-53 (2017).

Jombart, T. adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics* **24**, 1403-1405, doi:10.1093/bioinformatics/btn129 (2008).

Goudet, J. hierfstat, a package for r to compute and test hierarchical F-statistics. *Molecular Ecology Notes* **5**, 184-186 (2005).

Storey, J. D. A direct approach to false discovery rates. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **64**, 479-498 (2002).

Käll, L., Storey, J. D., MacCoss, M. J. & Noble, W. S. Posterior Error Probabilities and False Discovery Rates: Two Sides of the Same Coin. *Journal of Proteome Research* **7**, 40-44, doi:10.1021/pr700739d (2008).

Roselaar, C. Songbirds of Turkey. An atlas of biodiversity of Turkish passerine birds. A & C Black (1995).

Procházka, P. *et al.* Low genetic differentiation among Reed Warbler *Acrocephalus scirpaceus* populations across Europe. *Journal of Avian Biology* **42**, 103-113, doi:10.2307/23018103 (2011).

Procházka, P., Van Wilgenburg, S. L., Neto, J. M., Yosef, R. & Hobson, K. A. Using stable hydrogen isotopes ($\delta 2H$) and ring recoveries to trace natal origins in a Eurasian passerine with a migratory divide. *Journal of Avian Biology* **44**, 541-550 (2013).

Arbabi, T., Gonzalez, J., Witt, H.-H., Klein, R. & Wink, M. Mitochondrial phylogeography of the Eurasian Reed Warbler *Acrocephalus scirpaceus* and the first genetic record of *A. s. fuscus* in Central Europe. *Ibis* **156**, 799-811 (2014).

Pârâu, L. G. & Wink, M. Common patterns in the molecular phylogeography of western palearctic birds: a comprehensive review. *Journal of Ornithology* **162**, 937-959, doi:10.1007/s10336-021-01893-x (2021).

Cahill, J. A. *et al.* Positive selection in noncoding genomic regions of vocal learning birds is associated with genes implicated in vocal learning and speech functions in humans. *Genome Res* **31**, 2035-2049, doi:10.1101/gr.275989.121 (2021).

Shiflett, M. W., Gavin, M. & Tran, T. S. Altered hippocampal-dependent memory and motor function in neuropilin 2-deficient mice. *Transl Psychiatry* **5**, e521, doi:10.1038/tp.2015.17 (2015).

Smith, J., Lipkin, E., Soller, M., Fulton, J. E. & Burt, D. W. Mapping QTL Associated with Resistance to Avian Oncogenic Marek's Disease Virus (MDV) Reveals Major Candidate Genes and Variants. *Genes (Basel)* **11**, doi:10.3390/genes11091019 (2020).

Charton, C. *et al.* The transcriptomic blueprint of molt in rooster using various tissues from Ginkkoridak (Korean long-tailed chicken). *BMC Genomics* **22**, 594, doi:10.1186/s12864-021-07903-9 (2021).

Røskaft, E., Moksnes, A., Stokke, B. G., Moskát, C. & Honza, M. The spatial habitat structure of host populations explains the pattern of rejection behavior in hosts and parasitic adaptations in cuckoos. *Behavioral Ecology* **13**, 163-168 (2002).

Røskaft, E., Takasu, F., Moksnes, A. & Stokke, B. G. Importance of spatial habitat structure on establishment of host defenses against brood parasitism. *Behavioral Ecology* **17**, 700-708, doi:10.1093/beheco/ark019 (2006).

Nogueira, A., Rey, P. J., Alcántara, J. M., Feitosa, R. M. & Lohmann, L. G. Geographic mosaic of plant evolution: extrafloral nectary variation mediated by ant and herbivore assemblages. *PLoS One* **10**, e0123806, doi:10.1371/journal.pone.0123806 (2015).

Craig, T. P. & Itami, J. K. A geographic mosaic of coevolution between Eurosta solidaginis (Fitch) and its host plant tall goldenrod *Solidago altissima* (L.). *Evolution* **75**, 3056-3070 (2021).

Cubaynes, S. *et al.* Disease outbreaks select for mate choice and coat color in wolves. *Science* **378**, 300-303, doi:10.1126/science.abi8745 (2022).

Davies, N. B. & Brooke, M. d. L. Cuckoos versus reed warblers: Adaptations and counteradaptations. *Animal Behaviour* **36**, 262-284 (1988).

1011   108   Lowry, D. B. *et al.* Breaking RAD: an evaluation of the utility of restriction site-
1012   associated DNA sequencing for genome scans of adaptation. *Mol Ecol Resour* **17**,
1013   142-152, doi:10.1111/1755-0998.12635 (2017).
1014
1015   109   Palmero-Iniesta, M., Espelta, J. M., Gordillo, J. & Pino, J. Changes in forest
1016   landscape patterns resulting from recent afforestation in Europe (1990–2012):
1017   defragmentation of pre-existing forest versus new patch proliferation. *Annals of*
1018   *Forest Science* **77**, 43, doi:10.1007/s13595-020-00946-0 (2020).
1019
1020   110   Mitchell, F. J. G. How open were European primeval forests? Hypothesis testing
1021   using palaeoecological data. *Journal of Ecology* **93**, 168-177 (2005).
1022
1023   111   Hoy, S. R. *et al.* The far-reaching effects of genetic process in a keystone predator
1024   species, grey wolves. *Sci Adv* **9**, eadc8724, doi:10.1126/sciadv.adc8724 (2023).
1025
1026   112   Bassitta, M. *et al.* Genomic signatures of drift and selection driven by predation and
1027   human pressure in an insular lizard. *Sci Rep* **11**, 6136, doi:10.1038/s41598-021-
1028   85591-x (2021).
1029
1030   113   Tams, V., Nickel, J. H., Ehring, A. & Cordellier, M. Insights into the genetic basis of
1031   predator-induced response in *Daphnia galeata*. *Ecol Evol* **10**, 13095-13108,
1032   doi:10.1002/ece3.6899 (2020).
1033
1034   114   Stewart, B. *et al.* The genetic architecture underlying prey-dependent performance in
1035   a microbial predator. *Nature Communications* **13**, 319, doi:10.1038/s41467-021-
1036   27844-x (2022).
1037
1038   115   Abdellaoui, A. & Verweij, K. J. H. Dissecting polygenic signals from genome-wide
1039   association studies on human behaviour. *Nature Human Behaviour* **5**, 686-694,
1040   doi:10.1038/s41562-021-01110-y (2021).
1041
1042   116   Bendesky, A. *et al.* The genetic basis of parental care evolution in monogamous mice.
1043   *Nature* **544**, 434-439, doi:10.1038/nature22074 (2017).