Structural insights into heterohexameric assembly of epilepsy-related ligand–receptor complex LGI1–ADAM22

Takayuki Yamaguchi[1,*], Kei Okatsu[1,*], Masato Kubota[1], Ayuka Mitsumori[1], Atsushi Yamagata[2], Yuko Fukata[3], Masaki Fukata[4], Mikihiro Shibata[5,6] & Shuya Fukai[1]

[1]Department of Chemistry, Graduate School of Science, Kyoto University, Kyoto 606-8502, Japan.

[2]Laboratory for Protein Functional and Structural Biology, RIKEN Center for Biosystems Dynamics Research, Kanagawa 230-0045, Japan

[3]Division of Molecular and Cellular Pharmacology, Nagoya University Graduate School of Medicine, Nagoya 466-8550, Japan

[4]Division of Neuropharmacology, Nagoya University Graduate School of Medicine, Nagoya 466-8550, Japan

[5]Institute for Frontier Science Initiative, Kanazawa University, Kanazawa, Ishikawa 920-1192, Japan.

[6]WPI Nano Life Science Institute (WPI-NanoLSI), Kanazawa University, Kanazawa, Ishikawa 920-1192, Japan.

*These authors contributed equally to this work.

Correspondence should be addressed to S.F. (fukai@iam.u-tokyo.ac.jp) or M.S. (msshibata@staff.kanazawa-u.ac.jp)

Significance

The neuronal secretory protein Leucine-rich glioma-inactivated 1 (LGI1) and its receptor protein ADAM22 play a critical role in maintaining normal synaptic signal transmission. Genetic mutations in LGI1 are linked to epilepsy. Structural analysis of the LGI1–ADAM22 complex is crucial for understanding the molecular mechanisms of epileptogenesis and for developing targeted treatments. In this study, we determined the cryo-EM structure of the heterohexameric LGI1–ADAM22 complex and visualized the dynamics of this complex by high-speed atomic force microscopy.

Abstract

Leucine-rich glioma-inactivated 1 protein (LGI1) is a secreted neuronal protein consisting of the N-terminal leucine-rich repeat (LRR) and C-terminal epitempin repeat (EPTP) domains. LGI1 is

linked to epilepsy, a neurological disorder caused by genetic mutations. ADAM22 is a membrane receptor that binds to LGI1 extracellularly and interacts with AMPA-type glutamate receptors via PSD-95 intracellularly to maintain normal synaptic signal transmission. Structural analysis of the LGI1–ADAM22 complex is important for understanding the molecular mechanism of epileptogenesis and developing new therapies against epilepsy. We previously reported the crystal structure of a 2:2 complex consisting of two molecules of LGI1 and two molecules of the ADAM22 ectodomain (ECD), which is suggested to bridge neurons across the synaptic cleft. On the other hand, multiangle light scattering, small-angle X-ray scattering, and cryo-EM analyses have suggested the existence of a 3:3 complex consisting of three molecules of LGI1 and three molecules of ADAM22. In the previous cryo-EM analysis, many observed particles were in a dissociated state, making it difficult to determine the three-dimensional (3D) structure of the 3:3 complex. In this study, we stabilized the 3:3 LGI1–ADAM22$_{ECD}$ complex using chemical crosslinking and determined the cryo-EM structures of the LGI1$_{LRR}$–LGI1$_{EPTP}$–ADAM22$_{ECD}$ and 3:3 LGI1–ADAM22$_{ECD}$ complexes at 2.78 Å and 3.79 Å resolutions, respectively. Furthermore, high-speed atomic force microscopy (HS-AFM) visualized the structural features and flexibility of the 3:3 LGI1–ADAM22$_{ECD}$ complex in solution. We discuss new insights into the interaction modes of the LGI1–ADAM22 higher-order complex and the structural properties of the 3:3 LGI1–ADAM22 complex.

Introduction

Epilepsy is a prevalent neurological disorder, affecting approximately 1% of the population. Epilepsy is characterized by recurrent, unprovoked seizures, resulting from an imbalance between excitation and inhibition within neural circuits. Mutations associated with epilepsy frequently occur in genes that regulate neuronal excitability through ion channels such as voltage-gated ion channels (*e.g.*, K^+, Na^+, and Ca^{2+} channels) and ligand-gated ion channels (*e.g.*, nicotinic acetylcholine and GABAA receptors) (1-3). Additionally, some epilepsy-related mutations have been identified in genes encoding non-ion channel proteins such as *LGI1*.

LGI1 is a 60-kDa secreted neuronal protein that consists of the N-terminal leucine-rich repeat (LRR) domain and the C-terminal epitempin-repeat (EPTP) domain (4) (Fig. 1a). Mutations in the LGI1 gene, resulting in incorrect folding and post-translational modifications, cause autosomal dominant lateral temporal lobe epilepsy (ADLTE) (5-7). For example, the E383A mutant of LGI1 is eliminated by the endoplasmic reticulum and becomes deficient in secretion (8). Meanwhile, the S473L mutation of LGI1 causes epileptiform seizures due to reduced binding to ADAM22, which is a member of the a disintegrin and metalloproteinase (ADAM) family (9, 10) and acts as a receptor for LGI1 in neurons without protease activity (11,

12). ADAM22 is a single-pass transmembrane protein. The ectodomain (ECD) of ADAM22 consists of a metalloprotease-like domain, a disintegrin domain, a cysteine-rich domain, and an EGF-like domain (13). The metalloprotease-like domain interacts with the EPTP domain of LGI1 in the extracellular space. In the cytoplasm, the PDZ-binding motif (PBM)-containing C-terminal tail of ADAM22 binds to a synaptic scaffolding protein PSD-95, which regulates the cellular dynamics of α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptors through binding to Stargazin, an auxiliary subunit of AMPA receptors, in the postsynapse (14, 15). Furthermore, LGI1 forms a complex with the voltage-gated potassium channel (VGKC) through ADAM22/23. As such, the LGI1-ADAM22 complex maintains normal nerve signal transduction through the interaction network including synaptic ion channels (10, 11, 16). Structural analysis of the LGI1-ADAM22 complex is important for understanding the molecular basis of epileptogenesis and for developing therapeutic strategies based on this understanding.

We previously reported the crystal structure of a 2:2 LGI1–ADAM22$_{ECD}$ complex consisting of two molecules of LGI1 and two molecules of ADAM22$_{ECD}$ (17). The long-axis length of the 2:2 complex is approximately 190 Å, which is comparable to the width of the synaptic cleft. This 2:2 complex structure and the structure-guided study on a mouse model for familial epilepsy suggested that the formation of the 2:2 complex bridges neurons in the synaptic cleft. The results revealed the structural basis of the interaction between the EPTP domain of one LGI1 and the LRR domain of the other LGI1, as well as the interaction between the EPTP domain of LGI1 and the metalloproteinase-like domain of ADAM22. On the other hand, size-exclusion chromatography-multiangle light scattering (SEC-MALS), size-exclusion chromatography-small-angle X-ray scattering (SEC-SAXS), and cryo-electron microscopy (cryo-EM) analyses suggested that three molecules of LGI1 and three molecules of ADAM22$_{ECD}$ bind to each other to form a 3:3 complex at near physiological salt concentrations (17). Regarding the function of the 3:3 complex, it has been suggested that the 3:3 complex regulates the density of synaptic molecules such as scaffolding proteins and synaptic vesicles. The 3:3 complex might serve as an extracellular scaffold to stabilize Kv1 channels or AMPA receptors. Otherwise, the 3:3 assembly on the same membrane might compete with the transsynaptic 2:2 assembly. It is also possible that the 3:3 complex regulates the accumulation of VGKC at axon terminals rather than at synapses, thereby suppressing neuronal hyperexcitability. However, no clear evidence has been provided, and the three-dimensional structure of the 3:3 complex has not yet been determined.

In this study, the three-dimensional (3D) structure of the 3:3 LGI1–ADAM22$_{ECD}$ complex was determined at a nominal resolution of 3.79 Å by cryo-EM single-particle analysis. We also determined the 3D structure of the LGI1$_{LRR}$–LGI1$_{EPTP}$–ADAM22$_{ECD}$ complex at a nominal resolution of 2.78 Å. These higher-resolution 3D structures provide more detailed interaction mechanisms for the LGI1–ADAM22 higher-order assembly. We also performed high-speed

atomic force microscopy (HS-AFM) (18, 19) to directly visualize the molecular dynamics of the LGI1–ADAM22$_{ECD}$ assembly and characterize its structural property. The cryo-EM and HS-AFM results suggest that the 3:3 LGI1–ADAM22$_{ECD}$ assembly does not form a rigid 3-fold symmetric structure but rather a flexible triangular structure accompanying relative motions between each protomer.

Results

Cryo-EM single particle analysis of the LGI1–ADAM22$_{ECD}$ complex

As we reported previously, the molar mass of the complex between the full-length LGI1 and ADAM22$_{ECD}$ determined by SEC-MALS at 150 mM NaCl was 356 kDa, corresponding to the 3:3 hexameric assembly of LGI1–ADAM22$_{ECD}$ in solution (17). This is consistent with our initial cryo-EM analysis, where 5% of the reference-free 2D class averaged images showed particles with pseudo-$C3$ symmetry suggestive of the 3:3 assembly. Since the 3:3 complex class was clearly seen and appeared structurally stable, we tried to determine the 3D structure of the 3:3 LGI1-ADAM22$_{ECD}$ complex by single particle analysis but failed due to insufficient numbers of particles of the 3:3 complex. On this basis, we decided to perform a structural analysis of the complex stabilized by chemical crosslinking. We co-expressed His$_6$-tagged LGI1 and non-tagged ADAM22$_{ECD}$ in Expi293F cells and purified the complex using Ni-NTA affinity chromatography and gel filtration chromatography at 50 mM NaCl (Fig. S1a,b). Then, fractions containing the LGI1–ADAM22$_{ECD}$ complex were collected and treated with glutaraldehyde to chemically crosslink the 3:3 complex. The sample was purified again by gel filtration chromatography. The chromatogram showed a peak likely corresponding to the 3:3 LGI1–ADAM22$_{ECD}$ complex (Fig S1c,d). One of the peak fractions was collected and subjected to single-particle analysis by cryo-EM. Among 2,766,936 particles picked from 7,625 movies without templates, 1,403,037 particle images were extracted with a box size of 544 pixels (0.752 Å/pixel) and downsampled to a size of 136 pixels (3.008 Å/pixel) by Fourier cropping. The downsampled images were subjected to reference-free 2D classification, which generated images of triangle-shaped particles considered to be the 3:3 LGI1–ADAM22$_{ECD}$ complex (Fig. S2). Other images looked like two particles stacked on top of each other (Fig. S2) or single particles clearly smaller than the 3:3 complex. The stacked particle images might represent the 2:2 complex viewed along the long axis or the 3:3 complex viewed from the side, which were difficult to distinguish. Six classes with 176,443 particles of the putative 3:3 LGI1-ADAM22$_{ECD}$ complex were used as templates for picking particles again. 2,530,790 particle images were extracted, downsampled, and subjected to the second run of reference-free 2D classification, resulting in a set of images similar to that generated in the first run of 2D classification (Fig. S3). Eighty 2D classes with 2,006,398 particles were selected for *ab initio* 3D reconstruction. Six classes of *ab initio* 3D models were generated and

then refined by heterogeneous refinement. The resultant 3D map in one class corresponded to a complex comprising LGI1$_{LRR}$, LGI1*$_{EPTP}$, and ADAM22$_{ECD}$ (* indicates a distinct molecule hereafter). Non-uniform refinement using the particle images with the original pixel size yielded a density map at a nominal resolution of 2.78 Å (Figs. 1b and S4 and Table S1). This map corresponds to a part of the 3:3 complex, where two-thirds of the complex may adopt various conformations. The 3D map of one other class corresponded to the 3:3 LGI1–ADAM22$_{ECD}$ complex. Non-uniform refinement using the particle images with the original pixel size yielded a density map at a nominal resolution of 3.79 Å (Figs. 1c,d and S4 and Table S1). Using these two maps, we constructed atomic models of the LGI1$_{LRR}$–LGI1*$_{EPTP}$–ADAM22$_{ECD}$ complex and the 3:3 LGI1-ADAM22$_{ECD}$ complex (Table S1). The resolutions of the present cryo-EM analysis (2.78 Å and 3.79 Å) are better than that of the crystal structure of the 2:2 complex (7.125 Å).

Interaction of the LGI1$_{EPTP}$–ADAM22$_{ECD}$ complex

In the cryo-EM structure of the LGI1$_{LRR}$–LGI1*$_{EPTP}$–ADAM22$_{ECD}$ complex, the LGI1$_{EPTP}$–ADAM22$_{ECD}$ structure is essentially identical to that determined by X-ray crystallography (Fig. 2a-c). At the interface, Trp398, Tyr408, and Tyr409 of ADAM22 are stacked in a layer and project into the hydrophobic inner rim of the central channel of LGI1$_{EPTP}$, which consists of Leu237, Phe256, Val284, Leu302, Tyr433, Met477, and Phe541 of LGI1 (Fig. 2b, c). In addition, several hydrogen bonds are formed between LGI1 and ADAM22: Arg330 and Lys353 of LGI1 form hydrogen bonds with Asp405 and Glu359 of ADAM22, respectively, whereas Arg378 of LGI1 forms hydrogen bonds with Ser340 and Thr406 of ADAM22 (Fig. 2d-g). The hydrogen bond between Asp431 of LGI1 and Lys362 of ADAM22 was also found in the cryo-EM structure (Fig. S5). These hydrogen bonds differ slightly from those in the previous crystal structure: In the previous crystal structure, Lys331 of LGI1 formed a hydrogen bond with the main-chain carbonyl of ADAM22 Asp405. On the other hand, in the present cryo-EM structure, Lys331 of LGI1 is reoriented and does not form the hydrogen bond. Arg378 of LGI1 hydrogen bonds with Ser340 and Glu359 in the crystal structure, while it does with Ser340 and Thr406 in the cryo-EM structure (Fig. 2f,g). In the crystal structure, Asp431 of LGI1 faces outward and does not form a hydrogen bond with Lys362 of ADAM22. These differences in hydrogen bonding may reflect the varying contribution of each interacting residue to the LGI1$_{EPTP}$–ADAM22$_{ECD}$ affinity. As shown in our previous pull-down experiments, mutations of the hydrophobic residues at the interface of ADAM22 (*i.e.*, Trp398, Tyr408, and Tyr409) almost or completely abolished binding to LGI1, whereas the E359A or D405A mutation of ADAM22 decreased but did not abolish its binding to LGI1 (17). The hydrogen bonds involving the LGI1$_{EPTP}$–ADAM22$_{ECD}$ interaction might be so weak that the orientation of the hydrogen bonding residues could alter. In the cryo-EM structure, we also found hydrogen bonds between Ser282 of LGI1 and the main-chain carbonyl of Thr397

of ADAM22, between the main-chain carbonyl of Trp376 of LGI1 and Gln334 of ADAM22, and between Lys353 of LGI1 and the main-chain carbonyl of Phe335 of ADAM22 (Fig. S5). These three hydrogen bonds were also formed in the crystal structure.

Intermolecular interactions between $LGI1_{LRR}$ and $LGI1*_{EPTP}$

The present cryo-EM map of the $LGI1_{LRR}$–$LGI1*_{EPTP}$–$ADAM22_{ECD}$ complex provides a more detailed view of the interaction between $LGI1_{LRR}$ and $LGI1*_{EPTP}$ (Fig. 3a,b) than the previous crystal structure of the 2:2 complex at a moderate resolution of 7.125 Å. Specifically, Glu123 and Arg76 in $LGI1_{LRR}$ form hydrogen bonds with Arg474 and Glu516 in $LGI1*_{EPTP}$, respectively. The hydrogen bond between Arg76 and Glu516* is reinforced by stacking with Phe121 of the LRR domain. Additionally, Asn52 and Ser73 of the LRR domain form hydrogen bonds with the main chains of Tyr496 and Asp495 of the EPTP domain, respectively. Leu54, Val75, Leu97, and Phe121 of the LRR domain also form extensive hydrophobic interactions (Fig. 3a,b). The R474A mutation in $LGI1_{EPTP}$ is a missense ADLTE mutation in LGI1, known to cause epileptic symptoms by inhibiting the assembly of the LGI1–ADAM22 higher-order complex (20, 21). The present higher-resolution structure demonstrates that Arg474 in the EPTP domain of one LGI1 molecule forms a hydrogen bond with Glu123 in the LRR domain of another LGI1 molecule, which was ambiguous in the previous moderate-resolution crystal structure. Regarding interactions for the higher-order assembly, the previous crystal structure suggested a possible interaction between His116 of LRR and Glu446 of ADAM22 (17). However, in the present cryo-EM structure, the distance between His116 and Glu446 is 6.6 Å (Fig. S6), indicating that they do not interact with each other. The present cryo-EM structure indicates no interaction between $LGI1_{LRR}$ and $ADAM22_{ECD}$.

Cryo-EM structure of the 3:3 LGI1–$ADAM22_{ECD}$ complex

The cryo-EM structure of the $LGI1_{LRR}$–$LGI1*_{EPTP}$–$ADAM22_{ECD}$ complex could fit a density map of the 3:3 LGI1–$ADAM22_{ECD}$ complex well. Three $LGI1_{LRR}$–$LGI1*_{EPTP}$–$ADAM22_{ECD}$ complex structures were put into the 3.79-Å-resolution map of the 3:3 complex. The density of LGI1 and the metalloprotease domain of $ADAM22_{ECD}$ appears relatively strong. The density of one interface between $LGI1_{EPTP}$ and $ADAM22_{ECD}$ is especially resolved well (Fig. S7a). Although the density of amino-acid side chains in $LGI1_{LRR}$ is not well separated, the main-chain structure of $LGI1_{LRR}$ could be fitted to the density map (Fig. S7b). On the other hand, the density of the disintegrin, cysteine-rich, and EGF-like domains after Pro445 of ADAM22 is relatively weak in all three molecules. Correspondingly, in the local resolution map, the resolution of these ADAM22 domains after Pro445 is low (Fig. S7c). Like the crystal structure of the 2:2 LGI1–$ADAM22_{ECD}$ complex, the LRR and EPTP domains of LGI1 are linked by a two-residue linker

(Ile222-Ile223) in an extended conformation. LGI1$_{EPTP}$ interacts with the metalloprotease-like domain of ADAM22 to form the LGI1$_{EPTP}$–ADAM22$_{ECD}$ complex, while the LRR domain of one LGI1 molecule interacts with the EPTP domain of the neighboring LGI1, thereby bridging three distant ADAM22 molecules in the 3:3 complex (Fig. 1c,d). The C-termini of the three ADAM22 molecules point in opposite directions. When the three assembled LGI1–ADAM22$_{ECD}$ complexes were superimposed with the LGI1$_{EPTP}$–ADAM22$_{ECD}$ structure as the reference, all three LGI1 LRR domains were oriented differently (Fig. 3c). Although the triangular shape observed in the 2D class averaged image suggested (pseudo-)$C3$ symmetry of the 3:3 complex, the determined structure of the 3:3 complex was not symmetric. We then analyzed the interdomain motion of the three LGI1 domains (assigned to chains B, D, and F in the deposited PDB file；Fig. 3d) in the 3:3 LGI1–ADAM22$_{ECD}$ complex by the DynDom server (22) (Fig. S8 and Table S2). Analysis of the motion between the three pairs of chains suggested that the static domain corresponds to the EPTP domain, while the mobile domain corresponds to the LRR domain, predicting a hinge region for interdomain bending. The LRR domain of chain D was rotated 69.8° around the hinge axis compared to chain B, and the LRR domain of chain F was rotated 69.0° around the hinge axis compared to chain B. The LRR domains of chain D and chain F were also rotated 70.0° around the hinge axis relative to each other, suggesting that in the 3:3 LGI1–ADAM22$_{ECD}$ complex, each LRR domain rotates approximately 70° about the hinge axis (Table S2). Furthermore, the motion of each LRR domain varied widely: The movement of the LRR domain of chain D with respect to chain B follows a standard closure motion of 14 %, whereas the movement of the LRR domain of chain F with respect to chain D follows a standard closure motion of 99 % (Table S2).

Dynamics of the LGI1−ADAM22 higher-order complex observed by HS-AFM

To directly visualize the molecular dynamics of the LGI1–ADAM22$_{ECD}$ complex and characterize its structural properties in solution, we performed HS-AFM (18, 19). HS-AFM images of gel-filtration chromatography fractions containing the 3:3 LGI1–ADAM22$_{ECD}$ complex predominantly revealed triangular-shaped molecules, which appeared to exist stably with no drastic structural changes (Fig. 4a-c and Movie S1). A comparison with the simulated AFM image suggests that the protrusion on the exterior of the triangle is likely ADAM22 (Fig. 4b). This site frequently appeared to dissociate during HS-AFM scanning (Fig. 4c and Movie S1), indicating that the interaction between LGI1 and ADAM22 is weaker than the interactions among LGI1 molecules within the 3:3 LGI1–ADAM22$_{ECD}$ assembly. Additionally, HS-AFM also captured the dynamics of the 2:2 LGI1-ADAM22$_{ECD}$ complex (Fig. 4d-e and Movie S2). A comparison of the experimental HS-AFM image with the simulated AFM image based on the

crystal structure of the 2:2 complex indicates that the outer site corresponds to ADAM22, suggesting that LGI1 and ADAM22 are facing each other. The 2:2 complex also exhibited fragility during HS-AFM imaging, similar to the 3:3 complex (Fig. 4e and Movie S2). In individual LGI1 molecules dissociated from the complex, the LRR domain moved freely relative to the EPTP domain (Fig. 4f and after 10.2 s in Movie S2), likely reflecting the conformational difference in the relative orientation between LGI1$_{LRR}$ and LGI1$_{EPTP}$ observed within the cryo-EM structure of the 3:3 complex. Thus, single-molecule observations using HS-AFM demonstrated that the 3:3 LGI1–ADAM22$_{ECD}$ complex is present in solution, that the binding between ADAM22 and LGI1 is relatively weak within the 3:3 LGI1–ADAM22$_{ECD}$ assembly, and that the LRR domain of LGI1 exhibits flexibility.

Discussion

In this study, the ligand-receptor complex between LGI1, a secreted protein of neurons, and its receptor protein, ADAM22, was investigated by cryo-EM and HS-AFM. By chemically crosslinking with glutaraldehyde, we successfully captured much larger numbers of particle images of the 3:3 LGI1–ADAM22$_{ECD}$ complex than in our initial, preliminary cryo-EM study (17). We could determine not only the overall 3D structure of the 3:3 LGI1–ADAM22$_{ECD}$ complex but also the 3D structure of the LGI1$_{LRR}$–LGI1*$_{EPTP}$–ADAM22$_{ECD}$ complex at a nominal resolution of 2.78 Å, revealing the interaction mode of the LGI1–ADAM22 higher-order complex at a higher resolution than before. HS-AFM successfully imaged the LGI1–ADAM22$_{ECD}$ higher-order complex and confirmed that LGI1–ADAM22$_{ECD}$ forms a 3:3 complex in solution.

The present 3.79-Å-resolution cryo-EM map of the 3:3 LGI1–ADAM22$_{ECD}$ complex was calculated from 120,708 particle images selected after two rounds of heterogeneous refinement. On the other hand, a few classes of other particle images display triangular shapes with missing parts, suggesting domain motions or conformational heterogeneity in the 3:3 complex (Fig. S4). This raised a possibility that motion-based refinement might improve resolution in flexible regions. Therefore, we first performed 3D flexible refinement (3D Flex) using CryoSPARC (Movie S3) (23). However, even with the 3D Flex refinement, the density of the disintegrin, cysteine-rich, and EGF-like domains in the ADAM22 molecule remained poorly resolved. Then, we next hypothesized that the 3:3 LGI1–ADAM22$_{ECD}$ complex undergoes discrete, non-rigid motions and performed 3D Variability Analysis (3D VA) in CryoSPARC (24). Using three orthogonal principal modes, the 3D VA analysis indicated two twisting motions and one stretching motion of the triangular-shaped 3:3 LGI1–ADAM22$_{ECD}$ complex (Movie S4). This analysis also visualized relatively large motions of the disintegrin, cysteine-rich, and EGF-like domains of ADAM22 (Movie S4), suggestive of intrinsic conformational flexibilities of these domains, likely resulting in lower local resolution in the periphery of the complex.

Previous studies have shown that LGI1 is enriched at the axon initial segment and colocalized with ADAM22/23 and the voltage-gated potassium (Kv1) channels (25, 26) besides at the synapse. The Kv1-associated cell-adhesion molecules, TAG-1 and Caspr2, likely mediate this colocalization by binding to ADAM22/23. LGI1 knock-out mice show a reduced density of Kv1 channels, which is associated with increased neuronal excitability of hippocampal CA3 neurons, indicating that LGI1 regulates Kv1 channel function. In addition, the LGI1 R474Q mutation has been found to interfere with the colocalization of ADAM22/23 and Kv1 channels. Given that the LGI1 R474Q mutation inhibits the higher-order assembly of LGI1–ADAM22 with little impact on LGI1 secretion and binding to ADAM22, it is likely that the LGI1–ADAM22 higher-order complex regulates the axonal Kv1 channel function. The 3:3 LGI1–ADAM22 complex observed *in vitro* might facilitate efficient clustering of the axonal Kv1 channels to control their density and inhibit epilepsy. A recent study has shown that LGI3, a member of the LGI family, selectively co-assembles with Kv1 channels by using ADAM23 as the receptor in axons (27). LGI3 is secreted by oligodendrocytes in the brain and enriched at juxtaparanodes of myelinated axons to form subclusters. Both LGI1 and LGI3, along with ADAM22 and ADAM23, belong to the same subfamily and share a similar domain organization. This suggests that the LGI3–ADAM23 complex, like the LGI1–ADAM22 complex, may form a higher-order assembly (especially the 3:3 complex), which could be related to the clustering mechanism of Kv1 channels in axons. Finally, this study proposes that the LGI1–ADAM22 complex is an interesting therapeutic target for epilepsy and other neurological disorders. The 3:3 LGI1–ADAM22 complex structure revealed in this study could serve as a useful platform for structure-based drug design and facilitate the development of antiepileptic drugs.

Materials and methods

Protein preparation

For preparation of the LGI1–ADAM22$_{ECD}$ complex, the C-terminally His$_6$-tagged LGI1 was co-expressed with the non-tagged ADAM22$_{ECD}$ (R470A) in Expi293F cells (Thermo Fisher Scientific). As reported previously, the R470A mutation of ADAM22 increases the yield of the LGI1–ADAM22$_{ECD}$ complex without affecting the higher-order assembly of LGI1–ADAM22$_{ECD}$. The vectors for the co-expression were the pEBMulti-Neo vector (Wako chemicals) harboring the gene encoding human LRR (full length, residues 37–557) with the N-terminal Igκ signal sequence and that harboring ADAM22$_{ECD}$ (R470A) including the N-terminal prosequence (residues 1–729), both of which were reported previously. The culture media were loaded onto a Ni-NTA (Qiagen) column pre-equilibrated with 20 mM Tris-HCl (pH 8.0) containing 300 mM NaCl. After the column was washed with 20 mM Tris-HCl (pH 8.0) containing 300 mM NaCl and 25 mM imidazole, the proteins were eluted with 20 mM Tris-HCl (pH 8.0) containing 300

mM NaCl and 250 mM imidazole. The eluted proteins were further purified by size-exclusion chromatography using Superdex200 (GE healthcare) with 20 mM Tris-HCl (pH 7.5) buffer containing 50 mM NaCl. The purified proteins were concentrated to 0.25 g/L in Amicon Ultra-15 50,000 MWCO filter (Millipore), and glutaraldehyde was added at a final concentration of 0.1% (v/v). The sample was concentrated to 500 µL in Amicon Ultra-4 50,000 MWCO filter (Millipore) and purified by gel filtration chromatography using a Superose6 10/300 GL (GE Healthcare) column with 20 mM Tris-HCl buffer (pH 7.5) containing 50 mM NaCl. One of the peak fraction was concentrated to 0.5 g/L by Amicon Ultra-0.5 50,000 MWCO filter (Millipore).

Cryo-EM single particle analysis

Cu grids (R1.2/1.3, 300 mesh, Quantifoil) with holey carbon films were hydrophilized using a JEC-3000FC Auto Fine Coater (JEOL) at 7 Pa, 10 mA, and 10 sec. A 3 µL aliquot of 0.5 g/L protein solution was added to the grids using a Vitrobot Mark IV (Thermo Fisher Scientific) at a temperature of 8°C and 100% humidity. The grids were then immersed in liquid ethane and rapidly frozen under the following conditions: waiting time of 0 sec, blotting time of 3 sec, and blotting force of 10. Data collection were carried out on a CRYO ARM 300 transmission electron microscope (JEOL Ltd., Japan) operating at 300 kV, equipped with an Omega-type in-column energy filter (slit width 20 eV) and a Gatan K3 electron detector (operated in correlated doubling sample mode) at SPring-8 (Hyogo, Japan). A total of 7,625 movies were automatically collected using SerialEM (28). Hole centering was performed using yoneoLocr (29) integrated as a SerialEM macro. Movies were collected using the beam-image shift method (5×5×1 matrices), at a target defocus range of −1.4 to −1.7 µm and a nominal magnification of ×60,000, corresponding to a calibrated pixel size of 0.752 Å/pixel. Each movie was recorded with an exposure time of 2.79627 s, subdivided into 60 frames with a total electron dose of 60.8046 e^{-1} $Å^{-2}$.

All processing was performed in CryoSPARC v.4.1.2 or higher (30). The collected micrographs were processed using patch motion correction and patch CTF correction. Particles were picked from 7,635 particle images with using a template, yielding 3,336,559 particles. After extracting the particles from the micrographs and classifying them into 2D classes, about 60% of the particles based on the average 2D image of the protein were selected. An initial 3D reconstruction and heterogeneous refinement were performed on six classes, and one of these classes was further refined by non-uniform refinement to obtain a 2.78 Å resolution 3D map of the LGI1$_{LRR}$–LGI1*$_{EPTP}$–ADAM22$_{ECD}$ complex. Additionally, by performing *ab initio* reconstruction and heterogeneous refinement of one of the six classes into three subclasses and refining one of these subclasses further with non-uniform refinement (31), a 3D map of the 3:3 LGI1–ADAM22$_{ECD}$ complex with a resolution of 3.79 Å was obtained. Overall resolution estimates correspond to an Fourier shell correlation of 0.143 using an optimized mask that is

automatically determined after refinement. Local resolution maps were obtained using local resolution estimation. Movies were created using 3D VA (24) and 3D Flex (23).

Model building

Model building was performed using the programs Coot (32) and UCSF ChimeraX (33). The initial model of the $LGI1_{LRR}$–$LGI1*_{EPTP}$–$ADAM22_{ECD}$ complex was built using a part of the crystal structure of the 2:2 $LGI1$–$ADAM22_{ECD}$ complex (PDB 5Y31) (17) and fitted into the cryo-EM map. The initial model of the 3:3 $LGI1$–$ADAM22_{ECD}$ complex was built using the cryo-EM structure of the $LGI1_{LRR}$–$LGI1*_{EPTP}$–$ADAM22_{ECD}$ complex. Three $LGI1_{LRR}$–$LGI1*_{EPTP}$–$ADAM22_{ECD}$ complexes were fitted into the cryo-EM map and $LGI1_{LRR}$ and $LGI1_{EPTP}$ were connected by modeling the linker region. The structure refinement was performed using the Phenix software package (34). All figures and movies were created using the programs PyMOL (35) and UCSF ChimeraX (33).

HS-AFM observations

HS-AFM experiments were conducted using a custom-built high-speed atomic force microscope operating in tapping mode (19). Briefly, an optical beam deflection detector monitored the cantilever's (BL-AC10DS-A2, Olympus, Japan) deflection using a 780 nm, 0.8 mW infrared (IR) laser. The cantilever exhibited a spring constant of approximately 100 pN/nm, a resonant frequency near 400 kHz, and a quality factor around 2 in a liquid environment. The IR beam was directed onto the cantilever's back surface through a 60× objective lens (CFI S Plan Fluor ELWD 60X, Nikon, Japan), and its reflection was detected by a two-segmented PIN photodiode (MPR-1, Graviton, Japan). The initial AFM tip had a triangular section resembling a bird's beak. To enhance spatial resolution, an amorphous carbon tip was grown on the bird beak tip via electron beam deposition using a scanning electron microscope (FE-SEM; Verious 5UC, Thermo Fisher Scientific, USA). The additional AFM tip was roughly 500 nm long, with an apex radius of approximately 1 nm after plasma etching with a plasma cleaner (Tergeo, PIC Scientific, USA). All HS-AFM images were obtained using the cantilever with the additional AFM tip. The cantilever's free oscillation amplitude was less than 1 nm, and during HS-AFM scanning, the set-point amplitude was adjusted to roughly 90% of the free amplitude. To minimize interaction forces between the sample and the AFM tip, "only trace imaging" (OTI) mode (36) was employed for HS-AFM. Data acquisition and operation were managed using custom software based on Visual Basic. NET (Microsoft).

HS-AFM observations of the $LGI1$–$ADAM22_{ECD}$ complex were conducted on AP-mica, prepared by treating the mica surface with 0.00005% (3-aminopropyl)triethoxysilane (APTES) (Shin-Etsu Chemical, Japan) in MilliQ water for 3 min. Samples at a concentration of 10 nM were

added to the AP-mica surface after 3 min incubation of 3 μL. HS-AFM observations of the LGI1–ADAM22$_{ECD}$ complex were performed in 20 mM Tris-HCl (pH 7.4) buffer containing 150 mM NaCl. All HS-AFM experiments were performed at room temperature (24–26°C) and was independently repeated at least three times, consistently yielding similar results. For image processing, the HS-AFM images were processed using Fiji (ImageJ) software (NIH, USA) (37). A mean filter with a radius of 0.5 pixels was utilized to lower noise levels in each image. The Template Matching and Slice Alignment plugin for ImageJ was used to correct for drift between sequential images.

Simulation of AFM Images

The BioAFMviewer software (38) was utilized to validate the captured topographies of the LGI1–ADAM22$_{ECD}$ complex. The simulated scanning was based on the nonelastic collisions between a rigid cone-shaped tip model and the rigid van der Waals atomic model of the protein structure . Automatized fitting (39) was employed to generate a simulated image that closely matched the HS-AFM target image (image correlation coefficients reported in each figure). In the simulation presented in Fig. 4, the tip shape parameters were set to $R = 0.4$ nm for the tip probe sphere radius and $α = 5.0°$ for the cone half angle.

Data, materials, and software availability

The coordinates and maps of the LGI1$_{LRR}$–LGI1*$_{EPTP}$–ADAM22$_{ECD}$ complex and the 3:3 LGI1–ADAM22$_{ECD}$ complex have been deposited in the Protein Data Bank/Electron Microscopy Data Bank under the accession codes of 9KZC/EMD-62659 and 9KZT/EMD-62668, respectively. Other data are available from the corresponding authors upon reasonable request.

References

1.   Y. Fukata, M. Fukata, Epilepsy and synaptic proteins. *Curr Opin Neurobiol* **45**, 1-8 (2017).

2.   J. Noebels, Pathway-driven discovery of epilepsy genes. *Nat Neurosci* **18**, 344-350 (2015).

3.   O. K. Steinlein, Genetic mechanisms that underlie epilepsy. *Nat Rev Neurosci* **5**, 400-408 (2004).

4.   E. Staub *et al.*, The novel EPTP repeat defines a superfamily of proteins implicated in epileptic disorders. *Trends Biochem Sci* **27**, 441-444 (2002).

5.   W. Gu, E. Brodtkorb, O. K. Steinlein, LGI1 is mutated in familial temporal lobe epilepsy characterized by aphasic seizures. *Ann Neurol* **52**, 364-367 (2002).

6. S. Kalachikov *et al.*, Mutations in LGI1 cause autosomal-dominant partial epilepsy with auditory features. *Nat Genet* **30**, 335-341 (2002).

7. J. M. Morante-Redolat *et al.*, Mutations in the LGI1/Epitempin gene on 10q24 cause autosomal dominant lateral temporal epilepsy. *Hum Mol Genet* **11**, 1119-1128 (2002).

8. N. Yokoi *et al.*, Chemical corrector treatment ameliorates increased seizure susceptibility in a mouse model of familial epilepsy. *Nat Med* **21**, 19-26 (2015).

9. U. Schulte *et al.*, The epilepsy-linked Lgi1 protein assembles into presynaptic Kv1 channels and inhibits inactivation by Kvβ1. *Neuron* **49**, 697-706 (2006).

10. Y. Fukata *et al.*, Disruption of LGI1-linked synaptic complex causes abnormal synaptic transmission and epilepsy. *Proc Natl Acad Sci U S A* **107**, 3799-3804 (2010).

11. Y. Fukata *et al.*, Epilepsy-related ligand/receptor complex LGI1 and ADAM22 regulate synaptic transmission. *Science* **313**, 1792-1795 (2006).

12. K. Sagane, K. Yamazaki, Y. Mizui, I. Tanaka, Cloning and chromosomal mapping of mouse ADAM11, ADAM22 and ADAM23. *Gene* **236**, 79-86 (1999).

13. H. Liu, A. H. Shim, X. He, Structural characterization of the ectodomain of a disintegrin and metalloproteinase-22 (ADAM22), a neural adhesion receptor instead of metalloproteinase: insights on ADAM function. *J Biol Chem* **284**, 29077-29086 (2009).

14. L. Chen *et al.*, Stargazin regulates synaptic targeting of AMPA receptors by two distinct mechanisms. *Nature* **408**, 936-943 (2000).

15. R. A. Nicoll, S. Tomita, D. S. Bredt, Auxiliary subunits assist AMPA-type glutamate receptors. *Science* **311**, 1253-1256 (2006).

16. K. L. Lovero, Y. Fukata, A. J. Granger, M. Fukata, R. A. Nicoll, The LGI1-ADAM22 protein complex directs synapse maturation through regulation of PSD-95 function. *Proc Natl Acad Sci U S A* **112**, E4129-4137 (2015).

17. A. Yamagata *et al.*, Structural basis of epilepsy-related ligand-receptor complex LGI1-ADAM22. *Nat Commun* **9**, 1546 (2018).

18. T. Ando *et al.*, A high-speed atomic force microscope for studying biological macromolecules. *Proc Natl Acad Sci U S A* **98**, 12468-12472 (2001).

19. A. Sumino *et al.*, High-speed atomic force microscopy reveals fluctuations and dimer splitting of the N-terminal domain of GluA2 ionotropic glutamate receptor-auxiliary subunit complex. *ACS Nano* **18**, 25018-25035 (2024).

20. J. Kawamata *et al.*, Mutations in LGI1 gene in Japanese families with autosomal dominant lateral temporal lobe epilepsy: the first report from Asian families. *Epilepsia* **51**, 690-693 (2010).

21. E. Dazzo *et al.*, Secretion-positive LGI1 mutations linked to lateral temporal epilepsy impair binding to ADAM22 and ADAM23 receptors. *PLoS Genet* **12**, e1006376 (2016).

22. R. Veevers, S. Hayward, Methodological improvements for the analysis of domain movements in large biomolecular complexes. *Biophys Physicobiol* **16**, 328-336 (2019).

23. A. Punjani, D. J. Fleet, 3DFlex: determining structure and motion of flexible proteins from cryo-EM. *Nat Methods* **20**, 860-870 (2023).

24. A. Punjani, D. J. Fleet, 3D variability analysis: Resolving continuous flexibility and discrete heterogeneity from single particle cryo-EM. *J Struct Biol* **213**, 107702 (2021).

25. M. Seagar *et al.*, LGI1 tunes intrinsic excitability by regulating the density of axonal Kv1 channels. *Proc Natl Acad Sci U S A* **114**, 7719-7724 (2017).

26. B. Hivert, L. Marien, K. N. Agbam, C. Faivre-Sarrailh, ADAM22 and ADAM23 modulate the targeting of the Kv1 channel-associated protein LGI1 to the axon initial segment. *J Cell Sci* **132** (2019).

27. Y. Miyazaki *et al.*, Oligodendrocyte-derived LGI3 and its receptor ADAM23 organize juxtaparanodal Kv1 channel clustering for short-term synaptic plasticity. *Cell Rep* **43**, 113634 (2024).

28. D. N. Mastronarde, Automated electron microscope tomography using robust prediction of specimen movements. *J Struct Biol* **152**, 36-51 (2005).

29. K. Yonekura, S. Maki-Yonekura, H. Naitow, T. Hamaguchi, K. Takaba, Machine learning-based real-time object locator/evaluator for cryo-EM data collection. *Commun Biol* **4**, 1044 (2021).

30. A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296 (2017).

31. A. Punjani, H. Zhang, D. J. Fleet, Non-uniform refinement: adaptive regularization improves single-particle cryo-EM reconstruction. *Nat Methods* **17**, 1214-1221 (2020).

32. P. Emsley, K. Cowtan, Coot: model-building tools for molecular graphics. *Acta Crystallogr D Biol Crystallogr* **60**, 2126-2132 (2004).

33. E. C. Meng *et al.*, UCSF ChimeraX: Tools for structure building and analysis. *Protein Sci* **32**, e4792 (2023).

34. P. D. Adams *et al.*, The Phenix software for automated determination of macromolecular structures. *Methods* **55**, 94-106 (2011).

35. L. Schrödinger, W. DeLano (2020) PyMOL, Available at: http://www.pymol.org/pymol.

36. S. Fukuda, T. Ando, Faster high-speed atomic force microscopy for imaging of biomolecular processes. *Rev Sci Instrum* **92**, 033705 (2021).

37.   J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676-682 (2012).

38.   R. Amyot, H. Flechsig, BioAFMviewer: An interactive interface for simulated AFM scanning of biomolecular structures and dynamics. *PLoS Comput Biol* **16**, e1008444 (2020).

39.   R. Amyot, A. Marchesi, C. M. Franz, I. Casuso, H. Flechsig, Simulation atomic force microscopy for atomic reconstruction of biomolecular structures from resolution-limited experimental images. *PLoS Comput Biol* **18**, e1009970 (2022).

Acknowledgements

This research was partially supported by Research Support Project for Life Science and Drug Discovery (Basis for Supporting Innovative Drug Discovery and Life Science Research (BINDS)) from AMED under Grant Number JP24ama121001. Preliminary cryo-EM experiments were performed at EM01CT/EM02CT with the approval of the Japan Synchrotron Radiation Research Institute (JASRI) (Proposal No. 2021B2542, 2022A2542, 2022B2543, and 2023A2543). This work was supported by JSPS/MEXT KAKENHI (19H03162 to A.Y., 23H00374 to M.F., 18H03983 and 23H04070 to S.F.), Japan Agency for Medical Research and Development (JP24wm0625319 to Y.F. and JP24ek0109649 to M.F.), Takeda Science Foundation (to S.F.), the World Premier International Research Center Initiative (WPI), MEXT, Japan (to M.S.).

Author contributions

T.Y. prepared the sample for cryo-EM. T.Y., M.K., and A.M. performed cryo-EM analysis. K.O., A.Y., and S.F. assisted with cryo-EM analysis. A.M. prepared the sample for HS-AFM. M.S. performed HS-AFM experiments and analysis. Y.F., M.F., M.S., and S.F. designed the experiments. A.M., M.S., and S.F. wrote the paper. All authors discussed the results and commented on the manuscript.

Competing interests

The authors declare no competing interest.

Figure legends
Figure 1. Structure of the LGI1–ADAM22$_{ECD}$ complex

a Domain organizations of LGI1 and ADAM22. LGI1 consists of the LRR (blue) and EPTP (purple) domains. The N-terminal secretion signal peptide (SP, enclosed by dotted lines) is

removed in the secreted LGI1. The shaded blue boxes represent the N- and C-terminal caps, whereas the filled blue boxes represent the LRRs. The premature form of ADAM22 contains the N-terminal prosequence (enclosed by dotted lines). The mature ADAM22 consists of the metalloprotease-like (magenta), disintegrin (salmon pink), cysteine-rich (brown), EGF-like (orange), transmembrane (white), and cytoplasmic domains. The major ADAM22 isoform has a PDZ-binding motif in the C-terminal region of the cytoplasmic domain **b** Cryo-EM map and structure of the $LGI1_{LRR}$–$LGI1^*_{EPTP}$–$ADAM22_{ECD}$ complex at 2.78 Å resolution. **c** Cryo-EM map and structure of the 3:3 LGI1–$ADAM22_{ECD}$ complex at 3.79 Å resolution. **d** Schematic diagram of the 3:3 LGI1–$ADAM22_{ECD}$ complex. Black circles indicate the $LGI1_{LRR}$–$LGI1^*_{EPTP}$–$ADAM22_{ECD}$ complex.

Figure 2. Interactions between $LGI1_{EPTP}$ and $ADAM22_{ECD}$

a Superposition of the cryo-EM structure of the present $LGI1_{LRR}$–$LGI1^*_{EPTP}$–$ADAM22_{ECD}$ complex and the previously reported crystal structure of the $LGI1_{EPTP}$–$ADAM22_{ECD}$ complex. **b, c** Overall view of the interface between $LGI1_{EPTP}$ and $ADAM22_{ECD}$. The structure (**b**) and corresponding map (**c**) are shown. **d, e** Close-up view of the interaction between Arg330 of LGI1 and Asp405 of ADAM22. The structure (**d**) and corresponding map (**e**) are shown. **f, g** Close-up view of the interaction around Arg378 of LGI1. The structure (**f**) and corresponding map (**g**) are shown.

Figure 3. Structure and interaction of the higher-order LGI1–$ADAM22_{ECD}$ complex

a, b Overall view of the interface between $LGI1_{LRR}$ and $LGI1^*_{EPTP}$. The structure (**a**) and corresponding map (**b**) are shown. **c** Cryo-EM structure of the 3:3 LGI1–$ADAM22_{ECD}$ complex. **c** Superposition of the three LGI1-$ADAM22_{ECD}$ complexes in the cryo-EM structure of the 3:3 LGI1-$ADAM22_{ECD}$ complex. **d** Chain IDs of the individual LGI1 or $ADAM22_{ECD}$ molecules in the 3:3 LGI1-$ADAM22_{ECD}$ complex, assigned in this study.

Figure 4. HS-AFM observations of LGI1–$ADAM22_{ECD}$ complexes

a A representative HS-AFM image of the 3:3 LGI1–$ADAM22_{ECD}$ complex. The color bars on the right indicate height in nanometers. The frame rate was 1.0 frames/s.

b, d Magnified HS-AFM images of the 3:3 (left in b) and 2:2 (left in d) LGI1–$ADAM22_{ECD}$ complexes. The simulated AFM images (middle) were derived from fitting to the experimental HS-AFM image (left). The well-fitting simulated AFM images and the coefficient of correlation (CC) are indicated. The tertiary structures of the 3:3 (b) and 2:2 (d) LGI1–$ADAM22_{ECD}$ complexes are shown in the same orientation as the simulated AFM images.

c, e, f Sequential HS-AFM images of the 3:3 (c; see also Movie S1) and 2:2 (e, f; see also Movie S2) LGI1–ADAM22$_{ECD}$ complexes. A schematic illustration of the interpretation of HS-AFM images is shown at the bottom. Imaging parameters: scanning area = 120 × 96 nm^2 (240 × 192 pixels); frame rate = 3.3 frames/s. HS-AFM experiments were repeated independently at least 3 times with consistent results.

a

LGI1

SP N-CAP LRR C-CAP EPTP

ADAM22

Pro Metalloprotease Disintegrin Cys-rich EGF TM PDZ binding

ECD

b

LGI1*EPTP

LGI1LRR

ADAM22

LGI1LRR–LGI1*EPTP–ADAM22ECD complex
(Resolution 2.78 Å)

c

ADAM22

LGI1

ADAM22

LGI1

ADAM22

LGI1

3:3 LGI1–ADAM22ECD complex
(Resolution 3.79 Å)

d

ADAM22

LGI1

LGI1

ADAM22

LGI1

ADAM22

Fig. 1

Fig. 2

Fig. 3

a

12.0

0.0
(nm)

50 nm

b

HS-AFM image

Simulated AFM

Cryo-EM [XXXX.pdb]

3:3

LGI1

ADAM22

10 nm

CC 0.80

LGI1

ADAM22

LGI1

LGI1$_{EPTP}$

ADAM22

c

Height (nm) 0.0 ▬ 7.9

0 s

3:3

1:1

20 nm

0.3 s

12.0 s

LGI1 collapse

19.2 s

ADAM22 release

34.5 s

LGI1-ADAM22 release

ADAM22

LGI1$_{EPTP}$

LGI1$_{LRR}$

d

HS-AFM image

Simulated AFM

5Y31.pdb

2:2

LGI1

ADAM22

10 nm

CC 0.80

LGI1

ADAM22

LGI1

ADAM22

e

Height (nm) 0.0 ▬ 7.0

0 s

2:2

2:2

20 nm

6.0 s

ADAM22 release

6.3 s

8.1 s

LGI1 release

10.5 s

ADAM22 release

f

16.5 s

LGI1

LGI1$_{EPTP}$

LGI1$_{LRR}$

18.0 s

18.9 s

21.3 s

27.6 s

Fig. 4