1

Main Manuscript for

Improved synapsis dynamics accompany meiotic stability in
Arabidopsis arenosa autotetraploids

Authors: Adrián Gonzalo[1]*, Aditya Nayak[1], Kirsten Bomblies[1]*

1. Institute of Molecular Plant Biology, Department of Biology, ETH Zürich, Zürich, Switzerland.

*Correspondence to Kirsten Bomblies and Adrian Gonzalo

Email: kirsten.bomblies@biol.ethz.ch and adrian.gonzalo@biol.ethz.ch

Author Contributions: A.G. and K.B. conceptualized the research, A.G. and A.N. obtained the
materials and reagents, A.G. performed the research. A.G. analyzed the data, A.G. and K.B.
wrote and edited the manuscript.

Competing Interest Statement: The authors declare no competing interests.

Classification: Biological Sciences. Genetics.

Keywords: Meiosis, polyploidy, synaptonemal complex, crossovers, Arabidopsis.

This PDF file includes:
        Main Text
        Figures 1 to 6

Abstract

During meiosis, the correct pairing, synapsis, and recombination of homologous chromosome pairs is critical for fertility of sexual eukaryotes. These processes are challenged in polyploids, which possess additional copies of each chromosome. Polyploidy thus provides a unique context to study how evolution can modify meiotic programs in response to challenges. We previously observed that in *Arabidopsis arenosa* newly formed (neo-)polyploids, synapsis defects precede chromosomes associating in aberrant multivalent and univalent configurations. Here we study synapsis dynamics in genotypes with varying levels of meiotic stability to ask whether efficient synaptic progression is a key component of evolving stable tetraploid meiosis. We develop a method to quantify synapsis dynamics using the progression of foci of the pro-crossover factor HEI10 as a reference. HEI10 initially appears at many small loci before accumulating only at crossover sites. In diploids, this transition begins while there is still significant asynapsis, which quickly declines as HEI10 accumulation to fewer foci progresses. In neo-tetraploids, suboptimal elongation of synaptic initiation sites, and perhaps defective pairing, precedes synapsis stalling before the onset of HEI10 accumulation. In established tetraploids, HEI10 accumulation begins only when asynapsis is minimal, suggesting an enhanced HEI10/synapsis co-dynamic (even compared to diploids). Hybrids generated by crossing neo- and established tetraploids exhibit intermediate phenotypes. We find the extent of asynapsis correlates positively with crossover numbers, as well as a higher frequency of multivalents and univalent, which can disturb chromosome segregation. Our work supports the hypothesis that improving the efficiency of synapsis is important for evolving polyploid meiotic stability.

Significance Statement

During meiosis, homologous chromosomes pair and subsequently form the synaptonemal complex, which provides a context for maturation of DNA recombination events. How synapsis and crossover formation are coordinated in polyploids, where multiple chromosome sets complicate pair-wise interactions, remains unclear. Leveraging the dynamics of the pro-crossover factor HEI10 as a 'developmental clock,' we quantified synapsis progression in new and established tetraploids of *Arabidopsis arenosa*. Synapsis is severely compromised in neo-tetraploids, while in established ones it is more efficient even than diploids. Notably, the extent of synaptic defects correlated with excess crossovers, which compromises meiotic stability in tetraploids by connecting connect multiple homologs in multivalents. These findings highlight that improving synapsis dynamics are important for the evolution of meiotic stability in polyploids.

Main Text

Introduction

Meiosis is a specialized cell division that halves chromosome numbers through two rounds of segregation, producing haploid gametes. Crossovers generate new trait combinations and also hold homologous chromosomes in tight apposition, which ensures proper chromosome segregation – and thus fertility – in meiosis I (1). In polyploids, whole genome duplication leads to a genome with more than the usual two copies of each chromosomes, which can disrupt key meiotic processes, leading to abnormal chromosome configurations and impaired segregation (2). Understanding which processes are perturbed by genome duplication, and how they are adjusted in evolved polyploids, provides novel insights into meiosis and its evolution(3), as well as helping us understand how polyploids re-establish meiotic stability.

In diploids, proper pairing and segregation of homologs requires the correct number and pattern of crossovers, which is regulated by a meiotic program that ensures pair-wise interactions between homologs (forming bivalents) during prophase I of meiosis. Recombination begins with the

formation of DNA double-strand breaks (4), which are processed in the context of linear
proteinaceous axes that form along the length of each homolog (5). DSBs are typically required for
subsequent homolog co-alignment, and polymerization of the synaptonemal complex during
zygotene (6). The synaptonemal complex holds homologs in close alignment as recombination
events mature, and its formation coincides with the partial removal of some axis proteins, such as
ASY1 in plants (7, 8). The synaptonemal complex acts as a platform for recombination proteins,
including the dosage-sensitive HEI10, which transitions from dispersed weak foci in early
pachytene to bright foci at crossover sites in late pachytene (9–13). This process of HEI10
accumulation has been modeled based on data from *Arabidopsis thaliana* and *Caenorhabditis*
elegans to follow the biophysics of a "coarsening" process that has been proposed to explain
crossover patterning (14, 15). However, this model is debated, as it implies that crossover
designation can occur later than proposed in well-supported models from budding yeast and
Sordaria (16, 17). Regardless of whether HEI10 moves along it, defective synaptonemal complex
formation is known to cause abnormal crossover numbers and patterns in some species, including
Arabidopsis (18–21).

The processes of both synapsis and HEI10 accumulation are conserved across eukaryotic
kingdoms (10, 11, 16, 22–24). However, it is not known how they might be affected by whole
genome duplication or how they might evolve in response. In our previous work on natural auto-
tetraploid *A. arenosa* (which arose from a single within-species whole genome duplication about
30,000 generations ago (25)) we found that defects in neo-polyploids in crossover patterning were
accompanied by a defect in synapsis (26). Meiotically stable established tetraploids, in contrast,
not only evolved improved crossover patterning, they also recovered full synapsis. It remained
unknown at what stage synapsis becomes defective in neo-polyploids, whether it is stalled or
merely slow, to what extent established tetraploid synapsis is enhanced, or whether the synaptic
defects in neo-polyploids relate directly to defects in crossover patterning.

To clarify when and how synapsis becomes defective in neo-tetraploids and whether evolved
tetraploids have more efficient synapsis, we studied the co-dynamics of synapsis and HEI10
progression. Quantifying these dynamics would ideally require live imaging (27), but this is limited
to a few model species. Therefore, we developed a quantitative framework to assess synapsis
dynamics in male meiocytes of four *A. arenosa* genotypes with varying meiotic stability. We found
that synapsis is fully stalled, not just slow, in neo-tetraploids, which is associated with defects in
chromosome co-alignment. Though synaptic initiation appears normal in the neo-tetraploids, they
cannot efficiently elongate the synaptonemal complex from the initial sites. These defects correlate
with increased crossover rates, suggesting crossover patterning issues, which in turn may cause
the observed extensive multivalent and univalent formation. In contrast, established tetraploids
exhibit more efficient synapsis, even surpassing diploids, suggesting that synapsis improvements
contributed to meiotic stabilization and improved crossover regulation.

Results

Crossover number and meiotic stabilities of all genotypes at metaphase I

To explore the link between prophase I defects and meiotic instability in metaphase I, we quantified
meiotic features in natural diploids (2X), neo-tetraploid lines, created by colchicine treatment in the
lab (NEO-4X, see Material and Methods), evolved tetraploids (EST-4X), and hybrids (HYB-4X) from
crosses between EST-4X and NEO-4X (Fig. S1). We analyzed metaphase I and diakinesis spreads
(Fig. 1A, B). As previously reported, (26, 28), 2X showed full meiotic stability with all chromosome
pairs forming bivalents. EST-4X formed on average less than 1 quadrivalent per cell (Fig. 1C) and
very rarely univalents (0.1 per cell; Fig. 1D). HYB-4X had significantly more quadrivalents than
EST-4X (Fig. 1C), and a similarly low number of univalents (Fig. 1D). In contrast, NEO-4X formed
significantly more quadrivalents (3.0 ± 1.9 per cell, $p < 0.0041$, Fig. 1C) and also more univalents
(0.9 ± 1.4 per cell, $p > 0.0031$, Fig. 1D).

153
Because the frequency of quadrivalents should be positively (and univalents negatively) correlated
with crossover frequency (29–31), we also measured crossover number in 2X, NEO-4X, HYB-4X
and EST-4X by immunostaining for HEI10, which in late prophase I stages of diplotene and
diakinesis forms discrete foci at Class I crossover sites (32) (Fig. 1B). Consistent with previous
reports (26) we scored ~10 HEI10 foci per cell in 2X, about 22 in NEO-4X and 17 in EST-4X. HYB-
4X had an intermediate number (about 18 foci) between its two parent genotypes (Fig. 1E),
suggesting crossover number, like multivalent frequency, has a semidominant genetic basis.
161
162 **Quantification of HEI10 accumulation enables modeling of synapsis progression**
163
To obtain a detailed picture of prophase I dynamics in the different tetraploid materials, we
developed a quantitative framework by exploiting observable changes in HEI10 localization during
pachytene as a "developmental clock." HEI10 signal typically transitions from a very dispersed
pattern of numerous weak foci in early pachytene, to a few bright and enlarged foci that mark the
position of crossovers in late pachytene (10, 14, 22–24). We will refer to this dynamic process as
"HEI10 accumulation." We used HEI10 accumulation as a quantitative proxy for the developmental
progression of meiosis to study the dynamics of other concomitant events, mainly synapsis.
171
We imaged male meiocytes in the prophase I stages of zygotene and pachytene in a set of 17
plants (Fig. S1), including 2X (4 plants), NEO-4X (6 plants), HYB-4X (3 plants), and EST-4X (4
plants) using Structured Illumination Microscopy (SIM). We immunolocalized three meiotic proteins
(Fig. 2A): ZYP1 (the transverse filament of the synaptonemal complex, which marks synapsed
axes), ASY1 (an axis component largely removed upon synapsis, thus serving as a marker for
unsynapsed chromosomes), and HEI10 (to mark intermediates at different stages of the
recombination process). Prior to image analysis, we verified that these three antibodies specifically
stain meiotic cells; as expected, only a dim background signal was detected upon overexposure in
somatic cells (Fig. S2A).
181
Using a Fiji macro (see Material and Methods; Fig. 2A, S2B), we quantified HEI10 signal distribution
while preventing bias from human decisions through particle analysis of HEI10 foci. The "HEI10
accumulation level" was calculated on each cell as the percentage of HEI10 signal intensity in the
most relatively prominent foci compared to the total HEI10 signal intensity in that cell (Fig. 2A, Fig.
S2B, C). We defined "prominent foci" vs. "total signal" using two different automated detection
thresholds (Fig. 2A, S2B, C, see also Material and Methods). Each image serves as a "snapshot"
of a dynamic process, showing diverse HEI10 accumulation levels across different cells (Fig. 2A,
Fig. 3A), from low (where prominent foci account for a small share of total HEI10 signal intensity;
Fig. 2A) to high (where most total HEI10 signal is concentrated in prominent foci, presumably
marking crossover sites; Fig. 2A).
192
To assess the potential effect of background signal on HEI10 accumulation, we measured somatic
and meiotic leptotene cells, where HEI10 accumulation should be zero (Fig. S2D). These cells
registered an average of 7% accumulation level, with a maximum of 12% (Fig. S2E). Therefore, we
excluded cells with HEI10 accumulation levels below 12% from further analyses, as these values
might be indistinguishable from background. This resulted in a final dataset of 523 imaged
meiocytes from all genotypes (2X: 125 cells from 4 plants; NEO-4X: 172 cells from 6 plants; HYB-
4X: 134 cells from 3 plants; EST-4X: 94 cells from 4 plants). To quantify synapsis levels, we used
the ASY1 3D length (in μm) as a negative proxy for synapsis completion. Unfortunately, the
punctate pattern of ZYP1 immunostaining precluded reliable automated measurement of its 3D
length (Fig. S2F).
203
204 Next, we fitted a Gamma-Generalized Linear Mixed Model (Gamma-GLMM, see Material and
205 Methods and SI Appendix) using HEI10 accumulation level and genotype as predictors
206 (explanatory variables) for the extent of asynapsis as the response variable (Fig. 2B, Fig. S3A).

GLMMs are especially recommended when data from different individuals (with different sample sizes each) are pooled to deal with pseudoreplication. This is particularly important in heterozygous outcrossing species like *A. arenosa*, which have extensive genetic and phenotypic variation among individuals (33). The individual plant is specified as a random effect in the model formula (see SI Appendix). This model suggests a negative correlation between HEI10 accumulation and extent of asynapsis, described by an exponential decay curve with substantial explanatory power (R^2 = 0.660; see Material and Methods and SI Appendix). This fits prior descriptions that large HEI10 foci marking crossovers are associated with full synapsis (10, 14, 22–24, 34). Overall, these analyses illustrate the utility of HEI10 accumulation level as a quantitative marker for developmental progression during mid-prophase I, and as a reference to quantify synapsis dynamics.

Synapsis stalls in NEO-4X but is optimized in EST-4X

Despite the overall negative association between HEI10 accumulation level and the extent of asynapsis, plotting asynapsis (μm of ASY1 signal) against HEI10 accumulation revealed strong differences among genotypes, particularly for NEO-4X (Fig. 2B). Notably, NEO-4X exhibited higher levels of asynapsis at the onset of HEI10 accumulation (which we consider to be our above-calculated threshold of 12%) and little or no progress afterward, indicated by a flat trend line. To quantitatively analyze the differences in synapsis among genotypes, we examined the predictions and coefficients from our fitted model (Fig. 3B, C, D). We doubled the values of ASY1 length in 2X so they can be directly compared with those of tetraploids. At the onset of HEI10 accumulation (defined as our 12% cutoff), the extent of asynapsis was notably higher in NEO-4X (nearly 200 μm of ASY1; Fig. 3A, B) compared to 2X (about 50 μm of ASY1). In contrast, EST-4X showed minor asynapsis at this point (less than 10 μm of ASY1), significantly different from both NEO-4X and 2X (Fig. 3A, B). HYB-4X presented intermediate values (almost 30 μm ASY1; Fig. 3A, B), not significantly different from 2X or EST-4X but differing from NEO-4X. Overall, this analysis demonstrates that asynapsis at the onset of HEI10 accumulation varies significantly across genotypes, with NEO-4X exhibiting the highest levels. Interestingly, EST-4X displayed lower asynapsis than 2X, indicating it has more efficient early synapsis relative to HEI10 accumulation.

The analyses above raised the question of whether synapsis in NEO-4X is terminally stalled or merely very slow. Our Gamma-GLMM allowed us to compare the pace of synapsis progression across all genotypes by testing for differences in decay exponent coefficients of ASY1 length relative to HEI10 accumulation levels (Fig. 3C). We found statistically comparable decay exponent coefficients in 2X, HYB-4X, and EST-4X (Fig. 3D), which are all significantly different from 0 ($p < 0.0001$ for all). These means that ASY1 length changes exponentially as HEI10 accumulation increases. In contrast, the decay exponent of NEO-4X was not significantly different from 0 ($p = 0.8$, Fig. 3A and 3D) and differed significantly from all other genotypes (Fig. 3D), —hence the flat trendline observed in Fig. 2B. This result suggests that NEO-4X does not significantly reduce asynapsis as HEI10 accumulation advances, indicating that the extensive asynapsis present at the beginning persists through to the end of pachytene. These findings support the hypothesis that NEO-4X synapsis stalls before HEI10 accumulation begins. Although GLMM analysis excels in dealing with differences between individuals, plotting each plant individually hinds some differences in behavior (Fig S3), so it should not be ruled out that some individual plants may differ from the overall trend.

We also examined the final levels of asynapsis by determining when asynapsis ceases to decline. To do so, we identified the HEI10 accumulation level at which decay exponent coefficients for all genotypes were not significantly different from zero. We found that models including only cells with HEI10 accumulation levels greater than 49% had decay exponent coefficients for all genotypes not significantly different from zero (at least, $p = 0.12$, Gamma-GLMM, $R^2 = 0.849$). This model suggests, that after this cutoff, cells have already reached their asynapsis minimum. At this HEI10 accumulation level, both 2X and EST-4X had reached full synapsis (less than 2 μm of ASY1, Fig. 3D, S4). NEO-4X in contrast, retained very high levels of asynapsis (Fig. 3D, S4) and interestinlgy

HYB-4X showed intermediate levels (almost 10 of µm of ASY1) significantly different from both all the other genotypes.

Neo-tetraploids have less co-alignment of late asynaptic regions

Having observed that synapsis stalls in NEO-4X already before HEI10 starts accumulating, we hypothesized asynapsis in NEO-4X may be due to even earlier problems, e.g., during pairing, which leads to co-alignment. Thus, we examined co-alignment in cells with 30-150 µm of linear ASY1 signals, which allows assessment relative axis position (this becomes difficult in cells with greater ASY1 lengths). In this sub-sampled set we observed two types of geometry: "irregular", where pairs of asynaptic axes are not near one another, and "parallel", where asynaptic axes could be easily observed as coaligned ASY1 signals (Fig. 4A,B). These parallel asynaptic structures were typically 0.5 to 10 µm long, and could be seen as either terminal asynaptic regions at chromosome ends, or interstitial bubbles where asynaptic regions lie between two synapsed regions (Fig. 4A,B). We believe these parallel asynapsis regions are regions that did not have time to synapse yet (or in NEO-4X had stalled) as opposed to unresolved interlocks, as we did not observe interlocks when these structures were examined in 3D. We scored the number of parallel and irregular structures per cell and used it as a response variable in a Negative-binomial-GLMM (Fig. S5A) where ASY1 length was the explanatory variable. We indeed observed (with a substantial explanatory power; $R^2 = 0.7$) that there was a positive correlation between the length of ASY1 and the number of parallel axes. We interrogated this model to ask whether there are differences in the number of parallel axes between genotypes at comparable levels of asynapsis (ASY1 length = 100 microns), and found that NEO-4X has fewer parallel asynaptic axes than the other three genotypes (at most p < 0.0052, Negative-binomial-GLMM, Fig. 4C).

EST-4X and NEO-4X show different early synapsis dynamics.

Issues with pairing might also manifest as problems with synaptic initiation and/or elongation (19, 35), so we sought to get more insights into the initiation of synapsis by examining ZYP1 signals in early to mid-zygotene cells with little synapsis. We observed two kinds of ZYP1 signals in these cells: punctate signals, likely corresponding to synapsis initiation sites (SIS), and more elongated signals that we interpret as events where synapsis has begun to elongate from a previous synapsis initiation site (synapsis elongation sites; SES). To better understand the dynamics of ZYP1 signals, we first modeled their number (response variable) as a function of total ZYP1 length (explanatory variable). Although we cannot rule out that some synaptic stretches are reverted, we used the total length of ZYP1 as a proxy of developmental time (the longer the total ZYP1 length, the later the progress). Here, we used Generalized Linear Models (GLM, which differs from GLMM in that it does not control for individual variation or any other random effect) as they showed a better fit to the data than GLMMs (see SI Appendix).

Although we observed no differences in the behavior of punctate signals (Fig S5B, C and D), when we modeled the dynamics of the appearance of elongated ones, we fitted a negative-binomial-GLM with very strong explanatory power ($R^2 = 0.98$, Fig. 4E), which suggests a linear correlation between the number of SES and the total length of ZYP1. Our GLM analysis showed a greater initial number of SES (the model's intercept) in NEO-4X than EST-4X (Fig. 4F). Interestingly, we also observed that the rate at which the number of elongated signals increases as the total ZYP1 length grows (the model's slope) is significantly greater in EST-4X than in NEO-4X. Since we already saw NEO-4X has fewer co-aligned regions and does not complete synapsis, we hypothesize that these differences in slope mean that synapsis initiation sites elongate more readily in EST-4X, whereas in NEO-4X synapsis initiates normally, but a lower proportion of the initiation sites are proficient to elongate normally.

Preferential localization of prominent HEI10 foci to synapsed regions is stage-dependent

We next asked whether or how HEI10 focus development is affected by the synaptic defects in
NEO-4X by examining HEI10 localization in more detail. It has been shown in other species that
HEI10 forms foci at synaptic initiation sites (11, 17, 36–38), thus, we began by analyzing cells in
early zygotene, when synapsis is just starting. Irrespective of genotype, in nuclei with incipient
synapsis, the few prominent HEI10 foci localize with no apparent preference to either asynaptic
axis (which accounts for the vast majority of axis at this stage) or occasionally on the few small
ZYP1 signals where synapsis has initiated (early zygotene cells in Fig 4D). This observation is
consistent with reports in *Arabidopsis thaliana* at the earliest stages of synapsis (10, 39), but not
other species, where HEI10 orthologs are specifically associated with synapsis initiation sites (11,
17, 36, 37). Although, non-prominent foci are more difficult to distinguish from background signal,
the lack of preference of prominent foci for synapsed stretches, suggests that in *A. arenosa*, HEI10
only marks (at most) a subset of synapsis initiation sites, and is not a reliable marker for them. In
all genotypes, HEI10 is mostly found in discrete punctate foci, though on rare occasions, we
observed linear HEI10 signal on unsynapsed chromosomes overlapping with ASY1, resembling a
pattern previously described in wheat (24) (Fig S6). Importantly, we did not consider these cells in
the previously described analyses as these linear signals could skew the results.
331
In more advanced stages, where HEI10 condensation to fewer foci has begun (HEI10 accumulation
levels of 12-24%), we observed that prominent HEI10 foci are more commonly located on synapsed
regions in all genotypes (early-mid zygotene cells in Fig. 4D, 5A, S6A,B), though they can
occasionally also occur on unsynapsed regions at this stage. Later, at 45-100% HEI10
accumulation levels, enlarged discrete foci were found almost exclusively in synapsed regions.
This was true for 2X, HYB-4X, EST-4X, that are fully or nearly fully synapsed, but also in NEO-4X,
that still retains extensive asynapsis where only 3 of 919 prominent foci were localized on
unsynapsed axis (Fig 5A). Remarkably, we observed that in these late asynaptic NEO-4X cells,
even small tracts of synapsis (< 5 µm) can contain a prominent HEI10 focus (Fig 5B,C), resembling
observations in mutants with patchy synapsis in *C. elegans* (18, 20) and *A. thaliana* (21, 39).
342
343 **Extensive asynapsis accompanies elevated crossover number**
344
Since synaptic alterations have been shown to alter crossover number and/or patterning in other
species (18–21), we asked whether defective synapsis in NEO-4X is related to their elevated
crossover number. Since the three tetraploid genotypes (NEO-4X, HYB-4X, EST-4X) have different
levels of asynapsis (Fig. 3) and different crossover numbers (Fig. 1), we tested if there is a
correlation among cells within a cytotype between degree of asynapsis and crossover number. We
filtered for cells where HEI10 foci likely specifically mark crossovers (late pachytene), by
determining after what HEI10 accumulation level the number of detected prominent foci resembled
the most the crossover count at diakinesis for that genotype. This corresponded to HEI10
accumulation levels greater than 44, 59, 48, and 65% for 2X, NEO-4X, HYB-4X, and EST-4X,
respectively (Fig. S7B). Once we calculated the number of HEI10 foci in each late pachytene cell,
we used it as a response variable in a Poisson-GLM with the extent of asynapsis (µm of ASY1) as
an explanatory variable. Our Poisson-GLM suggested, with substantial explanatory power
(R^2=0.608) that there is a significant (p < 0.0001) logarithmic relationship between the extent of
asynapsis and crossover number (Fig. 5D). This non-linear relationship, along with the fact that we
found no correlation within genotypes (R^2=0.141, at most), suggests that the potential relationship
between asynapsis and mis-regulation of crossovers is likely complex.
361
362 **Discussion**
363
We previously showed that appropriate crossover number and patterning can prevent meiotic
instability in autopolyploids (26, 29). We previously also reported that the synaptonemal complex
is compromised in neo-tetraploids of *Arabidopsis arenosa* (26). Synaptic defects have also been
reported in neo-autotetraploid *C. elegans* (40), as well as neo-allopolyploid *A. suecica* (41).
Interestingly, established tetraploids of both *A. suecica* and *A. arenosa* exhibit complete synapsis

(26, 42), suggesting its re-establishment is important for polyploids. In this study, we thus explored
the dynamics of synapsis in both neo- and established tetraploid *A. arenosa* to better understand
the origin of the problem itself, and more importantly, to ask if improving the dynamics of this
process might be an important adaptation for meiotic stability in established tetraploids.
373
Prophase I, the stage where crossover patterning and synapsis take place (6), is a highly dynamic
stage of meiosis. Quantitative analysis of a dynamic process like synapsis ideally would involve
live imaging experiments (27), but this limits image resolution, and is not tenable in non-model
systems where appropriate fluorescently tagged lines are not available. Moreover, tagging HEI10
at least in the *C. elegans* ortholog ZHP-3, has been reported to compromise its wild type function
(43). An appealing alternative is to measure quantifiable dynamics of one process relative to
quantifiable dynamics of another process in images of fixed meiocytes, which also allows analysis
using super-resolution microscopy. Such an approach has been used to study e.g., the dynamics
of DSB repair relative to synapsis progression in *A. thaliana* (44). Here we developed this approach
to analyze synapsis dynamics using HEI10 progression as a reference (Fig. 6). While informative,
this approach does carry the caveat that it provides only relative information and does not measure
how long events take in real time.
386
Recent work in several species has focused on the role that the synaptonemal complex (which
links the axes of homologous chromosomes and provides a platform for the maturation and
regulation of recombination (45–48)), plays in the final number and pattern of crossovers on
chromosomes (47, 49–51). Comparing synaptic progression to the progression of HEI10
accumulation, we found that there are differences among genotypes in synapsis dynamics already
from the very beginning of the process (Fig. 6). Our data suggest that synapsis initiation occurs
seemingly normally in neo-polyploids but elongates less efficiently in NEO-4X than EST-4X. This
may result from defects in homologous axis pairing or synaptonemal complex polymerization. The
former is supported based on asynaptic axes being more commonly parallel in 2X, HYB-4X, and
EST-4X than in NEO-4X. In late zygotene in NEO-4X, many regions remain unaligned, likely
unpaired, which may directly hinder synapsis completion. This diverges somewhat from the
conclusions in autotetraploid *C. elegans*, where synaptic defects were observed even when axes
were co-aligned in groups of four, leading to the idea that homologous chromosomes may "compete
for establishing synapsis connections" (40). The same may happen locally in *A. arenosa* as well,
where regional co-alignment of all four axes has also been observed, but these four-way alignments
occur in neo- as well as established polyploids, and thus they seem in themselves to not block
synapsis (26). It is nevertheless certainly possible that there are pairing problems that arise from
competition among the four homologs in neo-polyploids that could hinder synaptonemal complex
elongation, while the established polyploids evolved a solution to this challenge. Another
explanation for inefficient synapsis in neo-polyploids, could be that extra chromosome sets
complicate chromosome movements during prophase I, which are important to promote pairing
and synapsis (19, 35, 52).
409
How might evolved tetraploids have solved the synaptic issues their neo-tetraploid counterparts
face? Previously, we identified genes that show evidence of having been targets of natural selection
in the established tetraploid *A. arenosa* lineage, including numerous potentially relevant meiosis
genes (28, 53–55). If pairing itself is the ultimate problem, a possible candidate for the evolved
solution in established tetraploids is *PRD3* (56), which shows strong evidence of having been
targeted by selection in tetraploid *A. arenosa* (28). *PRD3* encodes a homolog of yeast *MER2* (57),
and is essential for the formation of the DSBs required for pairing and synapsis (56, 58). Evolved
variants of *PRD3* could thus enhance pairing, but whether or how they might do so remains to be
tested. DNA strand invasion is a requisite in plants for pairing and synapsis (59, 60) which is
regulated antagonistically by SDS (56) and FLIP(61) (and other factors), and the genes encoding
them also show strong evidence of selection in the tetraploid lineage (55). *RMF1*, another gene
under with evidence of having been targeted by selection in *A. arenosa*, was also recently shown
to negatively regulate strand invasion (62). We also previously suggested that changes to the

chromosome axis (e.g., stiffening), could lead to more efficient chromosome pairing and synapsis
in the established tetraploids (26, 33). This which would fit with the strong evidence for selection
having targeted the axis proteins ASY1 and ASY3 in the tetraploid lineage (28, 55), and where
tetraploid alleles have already been shown to reduce multivalent frequency. Moreover, recently
ASY3 dosage has been shown to affect synapsis and crossover number in allotetraploid *Brassica*
napus (63). However, genetically generated established tetraploid plants homozygous for diploid
alleles of ASY1 and ASY3 do not have defects in synapsis, suggesting that while they may
contribute, they are not solely responsible for the difference in synapsis between diploids and
tetraploid genotypes (33). Prominently, selection also targeted the synaptonemal complex protein
ZYP1 (28, 55), suggesting the derived allele found in tetraploids may have a very direct role of
modifying the synaptonemal complex itself, perhaps by enhancing the efficiency of polymerization,
but this, too, remains to be tested.
435
An important question regarding the synaptic defects observed in neo-polyploids is whether they
relate directly to the increased recombination rate observed in neo-tetraploids of *A. arenosa* (26)
and other species (64, 65). We first found a broad correlation across genotypes with neo-tetraploids
having both defective synapsis as well as higher crossover numbers relative to both established
tetraploids and neotetraploid/established tetraploid hybrids. We found additional support for this
correlation from the finding that asynapsis and crossover number correlate positively across cells,
which showed a non-linear but logarithmic correlation. However, this non-linear relationship could
indicate that crossover number is not proportional to the extent of asynapsis, but maybe to other
factors like the number of synaptic patches (as observed in A. thaliana *pch2* mutants (21)).
Alternatively, a threshold-like effect could disrupt crossover regulation at certain level of asynapsis,
but not less. For example, DSBs are known to be increased in situations where homologs fail to
engage through synapsis or crossovers in budding yeast and *C. elegans* (46, 66, 67).
448
We hypothesize this link may be caused at least in part by the fact that neo-tetraploids have HEI10
dynamics taking place in the context of abnormal patchy synapsis. This might allow recombination
intermediates that would otherwise not develop into crossovers to accumulate sufficient HEI10 to
become crossover-fated. Similar alterations of crossover patterning are reported for the *axr1* (39),
pss1 (19) and *pch2* (21) mutants in *A. thaliana* and *syp-1* knock down mutants in C. *elegans* (18,
20) which have patchy synapsis. Because of its preference for associating with synapsed regions,
HEI10 accumulates locally to higher concentrations per length of synaptonemal complex when
synapsis is patchy, resulting in nearly every synaptic "patch" developing at least one prominent
HEI10 focus (19). Importantly, this would imply that local concentration of HEI10 in synapsed
regions can alter the fate of at least some recombination intermediates, implying they retain some
plasticity and remain sensitive to HEI10 levels.
460
We suggest that patchy synapsis in neo-tetraploids might similarly locally concentrate HEI10,
assuming about the same amount of protein per unit DNA now accumulates on a smaller total
amount of synaptonemal complex length. Conversely, in the context of full synapsis (as in the
established tetraploids), HEI10 could be siphoned off recombination intermediates that are not
irreversibly crossover-fated; in the context of patchy synapsis this may not be possible, allowing
HEI10 to be retained on these plastic sites. The HEI10 present on a synaptic patch would
accumulate where it can – a crossover-designated site when one is available, a plastic site when
not. When this happens in *A. thaliana* patchy synapsis mutants, it does not result in a crossover
increase, but this is because the number of synaptic patches is lower than the wildtype crossover
number. However, in *C. elegans* where in wild type meiosis there is only 1 crossover per bivalent,
mutants with patchy synapsis do show a crossover increase (18, 20). In *A. arenosa* crossover rates
per chromosome pair are also low (1.06 and 1.25 crossovers per chromosome pair in EST-4X and
2X, respectively), and the number of synaptic patches in neo-tetraploids is greater than this (26).
Overall, the increased efficiency of synapsis we see in the evolved polyploids of *A. arenosa* likely
helps prevent excess recombination intermediates taking on crossover fates, and may also help
solve other patterning issues (26).

477
Materials and Methods

479
Plant material. We used lab propagated descendants of natural populations of diploid (2X) and
autotetraploid (EST-4X) *A. arenosa* from Strečno, Slovakia (SNO, 2X) and TBG Triberg, Germany
(TBG, EST-4X) (25, 68). To produce neo-tetraploids (NEO-4X), we treated 2X 14-days-old
seedlings in the apical meristem with 0.05% colchicine (Sigma) diluted in sterile water with 0.05%
Silwet- 77 (Anawa). Neo-tetraploid branches of treated plants were identified in colchicine-derived
chimeric plants using flow cytometry. Confirmed tetraploid branches were used both for crosses (to
obtain a second-generation non-chimeric neo-tetraploids). For the experiments we used both
colchicine-treated and second-generation neo-tetraploids (as indicated in Figure S1), since our
analysis showed no significant differences between them (see SI Appendix). To generate tetraploid
hybrids (HYB-4X), we crossed EST-4X as male with NEO-4X as a female (as described in Figure
S1). We verified the karyotype of second-generation NEO-4X, HYB-4X and EST-4X plants in
meiotic and mitotic spreads and only proceed to further analyses with confirmed euploids.

Metaphase I and late prophase I spreads. We used chromosomes spreads of flower buds fixed
in ethanol:acetic acid 3:1 at diplotene, diakinesis (immunostaining of HEI10 for crossover
quantification), and metaphase I (univalent and quadrivalent counts). We performed spreads
following the protocol in ref. (32), with the minor modifications explained in ref. (29). Spreads were
visualized and imaged using Leica Thunder Imager 3D Tissue epifluorescence microscope.

Immunostaining. For HEI10 foci count for crossover quantification in late prophase I cells
(diplotene and diakinesis) we followed the protocol in ref. (32), with the same minor modifications
explained in ref. (29). We used a primary monoclonal antibody against *A. thaliana* HEI10 (29) and
a secondary goat anti-guinea pig Alexa-488 antibody (both at 1:200 dilution). We visualized and
imaged immunostained late prophase I cells with a Leica Thunder Imager 3D Tissue
epifluorescence microscope. To visualize HEI10 accumulation and the extent of asynapsis in
zygotene/pachytene cells, we used 3D Structured Illumination Microscopy (SIM) following the
protocol described in ref. (69). For this immunostaining protocol, we used the following primary
antibodies: rabbit polyclonal anti-HEI10, guinea pig monoclonal anti-ZYP1, guinea pig polyclonal
anti-ZYP1, guinea pig polyclonal anti-ASY1 (all with 1:500 dilution) and rat polyclonal anti-ASY1
(with 1:1000 dilution). We used the following secondary antibodies with a 1:200 dilution: goat anti-
rabbit Alexa-647, goat anti-rat Alexa-555, goat anti-guinea pig Alexa-488. Cells were imaged using
a Deltavision OMX SIM microscope in 3D stacks of 0.125 microns optical section spacing.

Data Generation and Analysis.

Image analysis and scoring. Multivalent and univalent scoring was performed blindly in
randomized metaphase I images. Each metaphase I image was assigned to a "scorability" class
(A, B, C, D, or E), depending on the quality of the spread and the confidence of the count, with A
the best and E the least reliable (examples shown in Fig. S8). Scorability class was considered for
statistical analyses. 2-chanel multi-image files for HEI10 focus count in late prophase I spreads did
not allow image randomization. To analyzed HEI10 accumulation and the extent of asynapsis in
zygotene/pachytene cells, we processed SIM images in 3D stacks using three different ImageJ Fiji
macros applied in batch to the full image set. These macros apply different thresholding methods
to detect fluorescence signals (70, 71) of HEI10 or ASY1. Macro 1 analyzes HEI10 signal to
estimate the number of prominent foci, and the intensity of both prominent foci and total signal
(which we used to calculate the HEI10 accumulation level). Macro 2, analyzes ASY1 signal to
measure its 3D length (in µm) to estimate the extent of asynapsis. Macro 3 analyzes the signal
from both ASY1 and ZYP1 for quality control purposes. Details about the macros are further
explained in the SI Appendix.
529

Statistical analysis. All statistical analyses were performed using Rstudio with the version 4.4.0 of R, and plots were made using ggplot2 (72). To analyze late prophase I HEI10 focus count data, since only one plant per individual was included per genotype, we used the classical statistical tests. Since there was unequal variance, we used Kruskal-Wallis and Dunnett tests (for multiple group comparisons) or t-test/Mann-Whitney U for simple comparisons (depending on whether equal variance criteria is met or not, respectively; details in Supplementary Dataset S8). For other datasets, since there were always genotypes with more than one individual, we used Generalized Linear Mixed Models, to account for random variation between individuals and prevent pseudoreplication. Occasionally we used Generalized Linear Models (which do not control for variability between individuals), when they showed a better fit than GLMMs (see SI Appendix). We used gaussian-GLM(M) or Gamma-GLM(M) (more appropriate for skewed data) for continuous data (such as ASY1 length) and Poisson-GLMM or negative binomial-GLMM for discrete data (such as multivalent/univalent or foci counts). We used the glmmTMB R package (73) to fit models. For each analysis we fitted several models and chose the best fit using the Performance (74) and DHARMa (75) R packages. Details on criteria and procedure to select the best fit are explained in the SI Appendix and Supporting Dataset S8 and S9. For the best fit GLMM, we estimated means, confidence intervals, medians, standard errors and p-values using the emmeans R package (76) with Holm-Bonferroni correction, based on Wald's tests performed by glmmTMB.

Data, Materials, and Software Availability. All image data used in this study are available freely from the ETH Research Collection. For SIM immunostaining images from 2x, DOI: 10.3929/ethz-b-000696798. For SIM immunostaining images from NEO-4X, DOI: 10.3929/ethz-b-000696907. For SIM immunostaining images from HYB-4X, DOI: 10.3929/ethz-b-000696850. For SIM immunostaining images from EST-4X, DOI: 10.3929/ethz-b-000696797. For epifluorescence microscopy images in metaphase, diakinesis and diplotene spreads, DOI; 10.3929/ethz-b-000696798. The ImageJ Fiji macros, R scripts and raw data files to reproduce the statistical analyses are available in https://github.com/adgon/MeioScope.

Acknowledgments

We thank the entire Bomblies group for scientific discussions, particularly Marinela Dukic for constructive feedback and Tadeas Priklopil for critical reading and guidance on statistics. We are thankful to Sophie Thüring for technical support in preparing metaphase I spreads. We also thank Joiselle B. Fernandes for scientific discussions, critical reading of the manuscript and support in editing. We also thank Tobias Schwarz and ETH ScopeM for technical support with SIM. We thank Chris Franklin for providing some of the antibodies used in this work. This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie (MSC) grant agreement No 101029732 to AG, as well as ETH core funds (KB).

References

1. G. S. Roeder, Meiotic chromosomes: it takes two to tango. *Genes Dev.* **11**, 2600–21 (1997).
2. L. Grandont, E. Jenczewski, A. Lloyd, Meiosis and Its Deviations in Polyploid Plants. *Cytogenet. Genome Res.* **14**, 171–184 (2013).
3. A. Gonzalo, All Ways Lead to Rome—Meiotic Stabilization Can Take Many Routes in Nascent Polyploid Plants. *Genes (Basel).* **13** (2022).
4. J. Szostak, T. Orr-Weaver, R. Rothstein, F. Stahl, The double-strand-break repair model for recombination. *Cell* **33**, 25–35 (1983).
5. C. Morgan, A. Nayak, N. Hosoya, G. R. Smith, C. Lambing, "Meiotic chromosome organization and its role in recombination and cancer" in *Current Topics in Developmental Biology*, 1st Ed., (Elsevier Inc., 2023), pp. 91–126.
6. D. Zickler, N. Kleckner, Meiotic chromosomes: integrating structure and function. *Annu.*

Rev. Genet. **33**, 603–754 (1999).

7.   A. Caryl, S. Armstrong, G. Jones, F. Franklin, A homologue of the yeast HOP1 gene is
inactivated in the Arabidopsis meiotic mutant asy1. *Chromosoma* **109**, 62–71 (2000).

8.   C. Lambing, P. C. Kuo, A. J. Tock, S. D. Topp, I. R. Henderson, ASY1 acts as a dosage-
dependent antagonist of telomere-led recombination and mediates crossover interference
in Arabidopsis. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 13647–13658 (2020).

9.   J. C. Fung, B. Rockmill, M. Odell, G. S. Roeder, Imposition of crossover interference
through the nonrandom distribution of synapsis initiation complexes. *Cell* **116**, 795–802
(2004).

10.   L. Chelysheva, *et al.*, The Arabidopsis HEI10 is a new ZMM protein related to Zip3. *PLoS*
Genet. **8** (2012).

11.   A. Reynolds, *et al.*, RNF212 is a dosage-sensitive regulator of crossing-over during
mammalian meiosis. *Nat. Genet.* **45**, 269–78 (2013).

12.   P. A. Ziolkowski, *et al.*, Natural variation and dosage of the HEI10 meiotic E3 ligase
control Arabidopsis crossover recombination. *Genes Dev.* **31**, 306–317 (2017).

13.   A. Gonzalo, *et al.*, Reducing MSH4 copy number prevents meiotic crossovers between
non-homologous chromosomes in Brassica napus. *Nat. Commun.* **10**, 2354 (2019).

14.   C. Morgan, *et al.*, Diffusion-mediated HEI10 coarsening can explain meiotic crossover
positioning in Arabidopsis. *Nat. Commun.* **12**, 4674 (2021).

15.   L. Zhang, W. Stauffer, D. Zwicker, A. F. Dernburg, @ Affiliations, Crossover patterning
through kinase-regulated condensation and coarsening of recombination nodules. *bioRxiv*,
2021.08.26.457865 (2021).

16.   C. M. Anderson, A. Oke, P. Yam, T. Zhuge, J. C. Fung, Reduced Crossover Interference
and Increased ZMM-Independent Recombination in the Absence of Tel1/ATM. *PLoS*
Genet. **11**, 1–27 (2015).

17.   L. Zhang, E. Espagne, A. De Muyt, D. Zickler, N. E. Kleckner, Interference-mediated
synaptonemal complex formation with embedded crossover designation. *Proc. Natl. Acad.*
Sci. U. S. A. **111**, E5059–E5068 (2014).

18.   D. E. Libuda, S. Uzawa, B. J. Meyer, A. M. Villeneuve, Meiotic chromosome structures
constrain and respond to designation of crossover sites. *Nature* **502**, 703–706 (2013).

19.   Y. Duroc, *et al.*, The Kinesin AtPSS1 Promotes Synapsis and is Required for Proper
Crossover Distribution in Meiosis. *PLoS Genet* **10** (2014).

20.   C. K. Cahoon, J. M. Helm, D. E. Libuda, Synaptonemal Complex Central Region Proteins
Promote Localization of Pro-crossover Factors to Recombination Events During
Caenorhabditis elegans Meiosis. *Genetics* **213**, 395–409 (2019).

21.   J. A. Fozard, C. Morgan, M. Howard, Coarsening dynamics can explain meiotic crossover
patterning in both the presence and absence of the synaptonemal complex. *Elife* **12**, 1–25
(2023).

22.   K. Wang, *et al.*, The role of rice HEI10 in the formation of meiotic crossovers. *PLoS Genet.*
8 (2012).

23.   L. Grandont, *et al.*, Homoeologous Chromosome Sorting and Progression of Meiotic
Recombination in Brassica napus: Ploidy Does Matter! *Plant Cell* **26**, 1448–1463 (2014).

24.   S. D. Desjardins, *et al.*, MutS homologue 4 and MutS homologue 5 Maintain the Obligate
Crossover in Wheat Despite Stepwise Gene Loss following Polyploidization. *Plant Physiol.*
183, 1545–1558 (2020).

25.   B. Arnold, S.-T. Kim, K. Bomblies, Single Geographic Origin of a Widespread
Autotetraploid Arabidopsis arenosa Lineage Followed by Interploidy Admixture. *Mol. Biol.*
Evol. **32**, 1382–95 (2015).

26.   C. Morgan, *et al.*, Evolution of crossover interference enables stable autopolyploidy by
ensuring pairwise partner connections in Arabidopsis arenosa. *Curr. Biol.*, 1–14 (2021).

27.   M. A. Prusicki, *et al.*, Live cell imaging of meiosis in arabidopsis thaliana. *Elife* **8**, 1–31
(2019).

28.   L. Yant, *et al.*, Meiotic Adaptation to Genome Duplication in Arabidopsis arenosa. *Curr.*
Biol., 1–6 (2013).

29.   A. Gonzalo, P. Parra-Nunez, A. L. Bachmann, E. Sanchez-Moran, K. Bomblies, Partial
639   cytological diploidization of neoautotetraploid meiosis by induced cross-over rate
640   reduction. *Proc. Natl. Acad. Sci.* **120**, 2017 (2023).
641   30.   K. Bomblies, G. Jones, C. Franklin, D. Zickler, N. Kleckner, The challenge of evolving
642   stable polyploidy: could an increase in "crossover interference distance" play a central
643   role? *Chromosoma* **125**, 287–300 (2016).
644   31.   D. Kostoff, Fertility and chromosome length. *J. Hered.* **31**, 33–34 (1940).
645   32.   L. Chelysheva, L. Grandont, M. Grelon, Immunolocalization of meiotic proteins in
646   Brassicaceae: method 1. *Methods Mol Biol*, 990:93 (2013).
647   33.   C. Morgan, H. Zhang, C. E. Henry, C. F. H. Franklin, K. Bomblies, Derived alleles of two
648   axis proteins affect meiotic traits in autotetraploid Arabidopsis arenosa. *Proc. Natl. Acad.*
649   *Sci. U. S. A.* **117**, 8980–8988 (2020).
650   34.   M. Ito, *et al.*, Distinct and interdependent functions of three RING proteins regulate
651   recombination during mammalian meiosis. *bioRxiv*, 2023.11.07.566091 (2023).
652   35.   J. Varas, *et al.*, Absence of SUN1 and SUN2 proteins in Arabidopsis thaliana leads to a
653   delay in meiotic progression and defects in synapsis and recombination. *Plant J.* **81**, 329–
654   46 (2015).
655   36.   M. Castellani, *et al.*, Meiotic recombination dynamics in plants with repeat-based
656   holocentromeres shed light on the primary drivers of crossover patterning. *Nat. plants* **10**,
657   423–438 (2024).
658   37.   S. Agarwal, G. S. Roeder, Zip3 Provides a Link between Recombination Enzymes and
659   Synaptonemal Complex Proteins. **102**, 245–255 (2000).
660   38.   N. Bhalla, D. J. Wynne, V. Jantsch, A. F. Dernburg, ZHP-3 acts at crossovers to couple
661   meiotic recombination with synaptonemal complex disassembly and bivalent formation in
662   C. elegans. *PLoS Genet.* **4** (2008).
663   39.   M. T. Jahns, *et al.*, Crossover Localisation Is Regulated by the Neddylation
664   Posttranslational Regulatory Pathway. *PLoS Biol* **12** (2014).
665   40.   S. Mlynarczyk-Evans, B. Roelens, A. M. Villeneuve, Evidence That Masking of Synapsis
666   Imperfections Counterbalances Quality Control to Promote Efficient Meiosis. *PLoS Genet.*
667   **9**, 14–16 (2013).
668   41.   C. Nibau, A. Evans, H. King, D. W. Phillips, A. Lloyd, Homoeologous crossovers are
669   distally biased and underlie genomic instability in first-generation neo-allopolyploid
670   Arabidopsis suecica. *New Phytol.* (2024) https:/doi.org/10.1111/nph.20095.
671   42.   C. Nibau, *et al.*, Meiosis in allopolyploid Arabidopsis suecica. *Plant J.* **111**, 1110–1122
672   (2022).
673   43.   L. von Diezmann, C. Bristow, O. Rog, Diffusion within the synaptonemal complex can
674   account for signal transduction along meiotic chromosomes. *bioRxiv* (2024).
675   44.   M. Martinez-Garcia, C. I. White, F. C. H. Franklin, E. Sanchez-Moran, The role of
676   topoisomerase II in DNA repair and recombination in arabidopsis thaliana. *Int. J. Mol. Sci.*
677   **22** (2021).
678   45.   K. Voelkel-meiman, C. Johnston, Y. Thappeta, Separable Crossover-Promoting and
679   Crossover-Constraining Aspects of Zip1 Activity during Budding Yeast Meiosis. **5**, 1–39
680   (2015).
681   46.   X. Mu, H. Murakami, N. Mohibullah, S. Keeney, Chromosome-autonomous feedback
682   down-regulates meiotic DNA break competence upon synaptonemal complex formation.
683   *Genes Dev.* **34**, 1605–1618 (2020).
684   47.   S. G. Gordon, L. E. Kursel, K. Xu, O. Rog, Synaptonemal Complex dimerization regulates
685   chromosome alignment and crossover patterning in meiosis. *PLoS Genet.* **17**, e1009205
686   (2021).
687   48.   L. I. Láscarez-Lagunas, *et al.*, ATM/ATR kinases link the synaptonemal complex and DNA
688   double-strand break repair pathway choice. *Curr. Biol.* **32**, 4719-4726.e4 (2022).
689   49.   L. Capilla-Pérez, *et al.*, The synaptonemal complex imposes crossover interference and
690   heterochiasmy in Arabidopsis. *Proc. Natl. Acad. Sci. U. S. A.* **118**, 1–11 (2021).
50.   M. G. France, *et al.*, ZYP1 is required for obligate cross-over formation and cross-over

interference in Arabidopsis. *Proc. Natl. Acad. Sci. U. S. A.* **118**, 1–11 (2021).

51. N. Vrielynck, *et al.*, SCEP1 and SCEP2 are two new components of the synaptonemal
complex central element. *Nat. plants* **9**, 2016–2030 (2023).

52. L. Cromer, *et al.*, Rapid meiotic prophase chromosome movements in Arabidopsis
thaliana are linked to essential reorganization at the nuclear envelope. *Nat. Commun.* **15**,
5964 (2024).

53. J. D. Hollister, *et al.*, Genetic Adaptation Associated with Genome-Doubling in
Autotetraploid Arabidopsis arenosa. *PLoS Genet* **8** (2012).

54. K. M. Wright, *et al.*, Selection on Meiosis Genes in Diploid and Tetraploid Arabidopsis
arenosa. *Mol Biol Evol* **32**, 944–955 (2014).

55. M. Bohutínská, *et al.*, De Novo Mutation and Rapid Protein (Co-)evolution during Meiotic
Adaptation in Arabidopsis arenosa. *Mol. Biol. Evol.* **38**, 1980–1994 (2021).

56. A. De Muyt, *et al.*, A high throughput genetic screen identifies new early meiotic
recombination functions in Arabidopsis thaliana. *PLoS Genet.* **5** (2009).

57. S. Tessé, *et al.*, Asy2/Mer2: an evolutionarily conserved mediator of meiotic
recombination, pairing, and global chromosome compaction. *Genes Dev.* **31**, 1880–1893
(2017).

58. M. Grelon, D. Vezon, G. Gendrot, G. Pelletier, AtSPO11-1 is necessary for efficient
meiotic recombination in plants. *EMBO J.* **20**, 589–600 (2001).

59. F. Couteau, *et al.*, Random chromosome segregation without meiotic arrest in both male
and female meiocytes of a dmc1 mutant of Arabidopsis. *Plant Cell* **11**, 1623–34 (1999).

60. C. Oliver, J. L. Santos, M. Pradillo, On the role of some ARGONAUTE proteins in meiosis
and DNA repair in Arabidopsis thaliana. *Front Plant Sci* **5**, 177 (2014).

61. J. B. Fernandes, *et al.*, FIGL1 and its novel partner FLIP form a conserved complex that
regulates homologous recombination. *PLoS Genet.* **14**, e1007317 (2018).

62. W. Xu, *et al.*, SCFRMF mediates degradation of the meiosis-specific recombinase DMC1.
Nat. Commun. **14**, 5044 (2023).

63. L. Chu, *et al.*, ASYNAPSIS3 has diverse dosage-dependent effects on meiotic crossover
formation in Brassica napus. *Plant Cell* **36**, 3838–3856 (2024).

64. M. Leflon, *et al.*, Crossovers Get a Boost in Brassica Allotriploid and Allotetraploid
Hybrids. *Plant Cell* **22**, 2253–2264 (2010).

65. A. Pecinka, W. Fang, M. Rehmsmeier, A. a Levy, O. Mittelsten Scheid, Polyploidization
increases meiotic recombination frequency in Arabidopsis. *BMC Biol.* **9**, 24 (2011).

66. E. L. Stamper, *et al.*, Identification of DSB-1, a Protein Required for Initiation of Meiotic
Recombination in Caenorhabditis elegans, Illuminates a Crossover Assurance
Checkpoint. *PLoS Genet.* **9**, 1–18 (2013).

67. D. Thacker, N. Mohibullah, X. Zhu, S. Keeney, Homologue engagement controls meiotic
DNA break number and distribution. *Nature* **510**, 241–6 (2014).

68. S. Marburger, *et al.*, Interspecific introgression mediates adaptation to whole genome
duplication. *Nat. Commun.* **10**, 5218 (2019).

69. C. Morgan, E. Wegel, Cytological Characterization of Arabidopsis arenosa Polyploids by
SIM. *Methods Mol. Biol.* **2061**, 37–46 (2020).

70. J. C. Yen, F. J. Chang, S. Chang, A New Criterion for Automatic Multilevel Thresholding.
IEEE Trans. Image Process. **4**, 370–378 (1995).

71. W. H. Tsai, Moment-preserving thresholding: a new approach. *Comput. Vision, Graph.*
Image Process. **29**, 377–393 (1985).

72. H. Wickham, Elegant Graphics for Data Analysis. *ttps://ggplot2.tidyverse.org* (2016).

73. E. Brooks, Mollie, *et al.*, glmmTMB Balances Speed and Flexibility Among Packages for
Zero-inflated Generalized Linear Mixed Modeling. *R J.* **9**, 378 (2017).

74. D. Lüdecke, M. Ben-Shachar, I. Patil, P. Waggoner, D. Makowski, performance: An R
Package for Assessment, Comparison and Testing of Statistical Models. *J. Open Source*
Softw. **6**, 3139 (2021).

75. F. Hartig, DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)
Regression Models. *http://florianhartig.github.io/DHARMa/* (2022).

76.   R. V. Lenth, emmeans: Estimated Marginal Means, aka Least-Squares Means.
747   *https://rvlenth.github.io/emmeans/* (2024).

748
749 **Figures and Tables**
750

751
752
Figure 1. Crossover number and meiotic stability in different tetraploid populations. Scale Bar, 10
µm. (A) Examples of metaphase I spreads from all genotypes. Red Q, T, and U, mark examples of
quadrivalents, trivalent, and univalent, respectively. (B) Examples of diakinesis cells with HEI10
immunostaining to detect crossovers. (C,D) Box plots of quadrivalent (C) and univalent (D) counts
per cell. (E) HEI10 foci count in late prophase I (diplotene plus pachytene) cells. For (C), (D) and
(E) and (E), population name, mean ± SD, and sample size. P-values are indicated on top of the

plots for each comparison (according to Wald's test on Negative binomial-GLMM, Poisson-GLMM
coefficients for (C) and (D); and Dunnett's test for (E).

761
762
763
Figure 2. Modeling synapsis dynamics. Scale bar 10 microns. (A) Examples of imaged cells where
two different thresholds (highlighted in red) were applied to the HEI10 channel to detect the total
signal and prominent foci. Based on the percentage of signal in prominent foci compared to the
total, the "HEI10 accumulation level" shown at left was calculated (see also Fig. S2). (B) Plots of
all analyzed cells showing HEI10 accumulation level (x axis) by µm of ASY1 (asynapsis, in the Y-
axis). Trendlines and 95% confidence intervals (shaded areas) are based in estimations from
GLMM. Grey-shadowed region marks datapoints that were not included in GLMM analysis because
they were under the HEI10 accumulation cutoff (12%) we estimated as our detection threshold.
772

Figure 3. Differences in synapsis dynamics. Scale bar = 10 μm. (A) Examples of imaged cells of all genotypes with different levels of HEI10 accumulation showing how asynapsis (marked by ASY1, green) generally declines as HEI10 accumulation level increases, except in NEO-4X. (B) Plot of predicted extent of asynapsis by a Gamma-GLMM at the onset of HEI10 accumulation. Error bars represent 95% confidence intervals. (C) Plot of decay exponent coefficients estimated by a Gamma-GLMM for different genotypes. Error bars represent 95% confidence intervals. (D) is a box plot of the final level of asynapsis, at levels of HEI10 accumulation where synapsis does not progress further. P-values are indicated on top of the plots for each comparison (according to Wald's test on Gamma-GLMM coefficients). Significant p-values were highlighted in red. Mean values (Mean ± SE) and sample sizes are indicated in the lower part of the plots.

Figure 4. Early synaptic defects in NEO-4X. Scale bar = 10 µm. (A) Example illustration of two types of geometry of asynaptic regions observed: parallel and irregular. (B) Examples of these structures observed in cells from the four genotypes. Arrowheads indicate regions classed as "parallel". (C) GLMM results showing differences in number of parallels when ASY1 length is 100 µm in all the genotypes, indicating differences in the frequencies of these structures. (D) Examples of early and early-to-mid zygotene meiocytes showing punctate (shaded arrowheads) and elongated (white arrowheads) ZYP1 signals. (E) Plot of data and predicted trendlines for NEO-4X and EST-4X according to our fitted model (Poisson-GLM) for elongated ZYP1 segments by total ZYP1 length. (F,G) Plots showing differences predicted by the fitted model in (F) the intercept for both genotypes (the initial number of SES, when ZYP1 length is zero) and (G) the slope (the increase of the number of SES as ZYP1 length grows). P-values are indicated on top of the plots for each comparison (according to Wald's test on Poisson-GLMM coefficients). Significant p- values were highlighted in red. Mean values (Mean ± SE) and sample sizes are indicated in the lower part of the plots.