Dorsal hippocampus mediates light-tone associations in male mice

Julia S. Pinho[1,2], Carla Ramon-Duaso[1], Irene Manzanares-Sierra[1], Arnau Busquets-

García[1]

[1] Cell-Type Mechanisms in Normal and Pathological Behavior Research Group,

Neuroscience Research Program, Hospital del Mar Research Institute, Barcelona,

Spain

[2] Current address: Gulbenkian Institute for Molecular Medicine, Oeiras, Portugal

Running title: Hippocampal involvement in mouse sensory preconditioning

Corresponding author:

Arnau Busquets-Garcia, PhD

Cell-Type Mechanisms in Normal and Pathological Behavior Research Group

Neuroscience Research Program, Hospital del Mar Research Institute.

PRBB

88 Dr. Aiguader street

Barcelona 08003

SPAIN

abusquets@researchmar.net

Tel: +34 93 316 0390

Abstract

Daily choices are often influenced by environmental cues that are not directly associated with reinforcers. This phenomenon, known as higher-order conditioning, can be studied using sensory preconditioning tasks in rodents. This behavioral paradigm involves the repeated pairing of two innocuous stimuli, such as a light and a tone, followed by a devaluation phase in which one stimulus is associated with an unconditioned stimulus, such as a mild foot-shock. The result is a conditioned response (e.g., freezing) to both the conditioned stimulus (direct learning) and the non-conditioned stimulus (mediated learning). In our study, we successfully established a light-tone sensory preconditioning task specifically in male mice, as sex differences were observed in both control experimental groups and in sensory preconditioning responses. We employed in vivo, freely moving fiber photometry to monitor neural activity in the dorsal and ventral subregions of the hippocampus in male mice during the formation of associations between innocuous stimuli and reinforcers. Additionally, we combined our sensory preconditioning task with chemogenetic approaches to investigate the roles of these hippocampal subregions in sensory preconditioning. Our results indicate that dorsal, but not ventral, CaMKII-positive neurons are involved in encoding innocuous stimuli during the preconditioning phase. Overall, we developed a novel light-tone sensory preconditioning protocol in male mice, enabling the detection of sex differences and furthering our understanding of how specific hippocampal subregions and cell types regulate complex cognitive processes.

1 **Introduction**
2
The brain mechanisms underlying associative learning have traditionally been
elucidated through classical conditioning paradigms involving salient stimuli as
reinforcers—such as pairing an electric foot-shock with innocuous stimuli (e.g., a light,
tone, or odor). However, these paradigms are not sufficient to fully capture the
complexity of animal behavior, as daily choices are not always dictated by stimuli that
have been directly associated with a potent reinforcer. In both humans and animals,
responses often occur to cues that were never explicitly conditioned but had previously
been associated with other stimuli linked to specific aversive or rewarding outcomes[1-5].
11
These higher-order conditioning processes can be studied in laboratory settings using
sensory preconditioning procedures[2,6-11]. These tasks involve incidental associations
between neutral stimuli (e.g., odors, tastes, lights, or tones) during a preconditioning
phase, followed by classical conditioning of one of these stimuli with an aversive or
appetitive unconditioned reinforcer[2,6-11]. As a result, subjects exhibit aversion or
preference towards the stimulus that was never directly paired with the reinforcer,
enabling the assessment of mediated learning[2,6-11]. While the brain circuits underlying
classical associative memory between neutral cues and reinforcers have been
extensively studied, the mechanisms responsible for incidental associations between
innocuous sensory stimuli—and their behavioral consequences—remain much less
understood.
23
The hippocampus and other cortical regions (e.g., the perirhinal, retrosplenial, and
orbitofrontal cortices) have been implicated in higher-order conditioned responses[6,8,10-14].
Sensory preconditioning tasks using various sensory modalities have shown that the
involvement of different brain regions may depend on the behavioral phase being
studied (i.e., preconditioning or testing) or the modality of the sensory cues (visual,
gustatory, olfactory, or auditory). However, it is still unknown whether there are
common brain regions where different types of stimuli are integrated, regardless of the
behavioral phase or the modality. The hippocampus has been proposed to play a
central role in multiple behavioral phases of sensory preconditioning procedures across
species[1,4,11]. This region, which maintains continuous information exchange with other
cortical areas, is thought to act as an integrator of past experiences into broader
cognitive representations through a tightly regulated excitatory-inhibitory balance[15,16].
Previous studies have highlighted a specific role for hippocampal mechanisms during
the encoding of incidental associations[11]. However, the contribution of specific

hippocampal subregions or cell types to sensory preconditioning remains unclear. The dorsoventral axis of the rodent hippocampus is known to be structurally and functionally segregated. The dorsal hippocampus, connected to cortical regions and the thalamus, is primarily involved in cognitive processes such as navigation and exploration[17-21]. In contrast, the ventral hippocampus, which communicates with the amygdala, nucleus accumbens, and hypothalamus, plays a key role in motivated and emotional behaviors[17,18,22]. Based on this, our initial hypothesis was that the dorsal and/or ventral hippocampus may differentially contribute to sensory preconditioning in mice.

To test this, we employed chemogenetic and imaging techniques in combination with an adapted light-tone sensory preconditioning ($_{LT}$SPC) procedure[6]. Our results suggest that the dorsal hippocampus, and in particular CaMKII-positive neurons, plays a crucial role in the formation of incidental associations between visual and auditory stimuli, ultimately driving sensory preconditioning responses. Understanding the differential contribution of the dorsal and ventral hippocampus to sensory preconditioning provides valuable insights into the mechanisms of higher-order conditioning and the broader role of the hippocampus in associative learning.

Material and methods:

Animals

Male and female C57BL/6J mice were purchased from Charles River Laboratories and

were used for behavioral and chemogenetic experiments. Male Parvalbumin (PV)-Cre

mice were obtained by own breedings in our animal facility. All the mice used in this

study were 8 weeks old at the beginning of the experiments and they were grouped-

housed and maintained in a temperature (20–24°C) and humidity (40%–70%)

controlled condition under a 12 h light/dark cycle and had ad libitum access to food and

water. All behavioral studies were performed during the dark cycle (from 9am to 5pm)

by trained researchers who were blind to the different experimental conditions.

All experimental procedures shown in this work have followed European guidelines

(2010/62/EU) and were approved by the Committee on Animal Health and Care of

Barcelona Biomedical Research Park (PRBB) and from the Generalitat de Catalunya.

All experiments were performed in the animal facility of the PRBB, which has the full

accreditation from the Association for Assessment and Accreditation of Laboratory

Animal Care (AAALAC).

Light-tone sensory preconditioning task ($_{LT}$SPC)

The task was performed using automatized conditioning chambers (Imetronic, France)

and was divided in different phases (Figure 1A):

Habituation: Animals were exposed to the chambers with the presence of background

noise (65db) (like white noise but allocated to different speakers to avoid that animals

associate the sound with a location) during one session of 20 minutes.

Preconditioning Phase: This phase was composed by six sessions (2 per day; 3 hours

of intersession interval) of 510 seconds each under background noise. These sessions

started with an OFF period (only background noise) of 3 min and were followed by 5

simultaneous presentations of a light (CS1, white LED light of 1.8cm of diameter

located in one side of the box) and a tone (CS2, 65db,3000Hz) during 30 seconds with

an intertrial interval of 30 seconds and a final OFF period of 1 minute. This protocol

details apply to the Paired and No-Shock experimental groups. For the unpaired

groups, light and tone were never associated together during this phase. Indeed,

animals were exposed to the same amount of time to light and tone but in different

sessions (S1 light, S2 tone, S3 tone, S4 light, S5 light, S6 tone, in a pseudo-

randomized way) separated by 3h (male and female mice) or 6h (female mice).

<u>Conditioning Phase:</u> This phase consisted of two training sessions in males and one

training session in females. Each session lasted 510 seconds under background noise.

In the case of males, there was an inter-session interval of 3 hours. Specifically, the

sessions started with an off period of 3 min and were followed by 5 presentations of a

10-seconds light stimulus (CS1) that co-terminated during 2 seconds with a mild foot-

shock (0.4 mA, see results) with an intertrial interval of 1 min and a final OFF period of

1 min. This protocol details apply to the Paired and Unpaired experimental groups. For

the No-Shock group, animals followed the same conditioning session but without the

exposure to the electric foot-shock. For the control group performed in female mice, no

light was associated with the foot-shock during this conditioning session.

<u>Probe Tests Phase:</u> Mice were subjected on the same day to two Probe Test sessions

where they were exposed to the tone (Mediated Learning, Probe Test 1) and the light

(Direct Learning, Probe Test 2) in a new context (different from the previous phases).

We changed 4 features of the context: floor texture, wall design, smell (from 70%

ethanol to a CR36 disinfectant solution), and absence of background noise to avoid

fear responses elicited by the context itself. The two Probe Tests lasted 6 minutes and

were separated by at least 1 hour. In this session, animals underwent an OFF period of

3 min followed by an ON period of 3 min where the tone (Probe Test 1) or the light

(Probe Test 2) were continuously exposed. All experimental groups (Paired, Unpaired

and No-Shock) perform the Probe Tests in the same manner. The presence of

mediated and direct learning is shown by the percentage of freezing during ON and

OFF periods.

The behavior analysis was automatized using Deeplabcut to track the animal's position

across time and a Python homemade script to compute all the different analyses,

graphs, and statistics performed in the present paper. The main behavior measured

was the freezing response (i.e. immobility), which was defined as an Euclidean

distance lower than 0.02 cm per pair of frames for videos with 25 fps, resulting in a

speed lower than 0.5 cm/s. We validated this automated behavioral counting by

performing correlations with manual counts and another software dedicated to freezing

scoring (EzTrack)[23], with which we found a high correlation (higher than 90%)

(Supplementary Figure 1). The Deeplabcut project and all the scripts will be available in

the GitHub repository of the lab.

***In vivo* Fiber Photometry**

Male C57BL/6J and PV-Cre mice were anesthetized with a mixture of ketamine (75

mg/Kg, Imalgene 500, Merial, Spain) and medetomidine (1 mg/Kg, Domtor, Spain) by

intraperitoneal (i.p.) injection. Then, animals were placed into a stereotaxic apparatus

(World Precision Instruments, FL, USA) with a mouse adaptor and lateral ear bars. For

viral intra-hippocampus delivery, AAV vectors were injected with a Hamilton syringe

coupled with a nanofill attached to a pump (UMP3-1, World Precision Instruments, FL,

USA). PV-cre mice were injected with a mixture of 200 nl of

AAV.Syn.NES.jRCaMP1a.WPRE.SV40 (titer: 1×10^{13}, addgene 100848-AAV9) and

200nl of a Cre-dependent AAV-syn-FLEX-jGCaMP8f-WPRE (titer: 1×10^{13}, addgene

162379-AAV9) in order to monitor both the activity of the synapsin-positive cells in red

and the PV interneurons in green. C57BL/6J mice were injected with 400nl of AAV-

CaMKII-GCaMP6m (titer: 1×10^{12}, Tebu-BIO 189SL-AAV9) to monitor activity of the

CaMKII positive neurons in green. To check DREADD functionality, 200 nl of the AAV-

CaMKII-GCaMP6m and 200 nl of the inhibitory DREADD were infused in a similar

manner. These viral vectors were infused (1 nl/s) directly into the dorsal (in one

hemisphere) or ventral hippocampus (in the other hemisphere) with the following

coordinates in mm: dorsal, AP ±1.5, ML ± 2, DV -1.5 and ventral, AP ±3.5, ML ± 3.3,

DV -3.5, according to Paxinos and Franklin brain atlas (Paxinos and Franklin, 2001).

After the AAV infusions, an optic Fiber (core 400 μm, N.A 0.5, RWD, China) was

implanted using dental cement following the same coordinates except for DV that was

placed 0.25 mm above viral infusions. Four weeks after this surgery, animals were

used for *in vivo* calcium recordings where they undergo a $_{LT}$SPC and *in vivo* recordings

were performed during Light-Tone associations (preconditioning phase) and Light-

Shock associations (conditioning phase). Before this experiment, animals were

habituated for 3 days to connect and disconnect the optic fibers and to be habituated to

the cable in the same chamber where the behavioral experiment was performed.

During these two behavioral phases, *in vivo* recordings were performed using a

commercial Fiber Photometry equipment (RWD, China) where we used 470nM LED to

excite the GCaMP sensor, and 560 nM for the RCaMP signal. In all mice used, after

the recording experiments, the signal was checked to validate the viral infusions.

To analyze the Fiber Photometry experiments, a custom python code was used and

the behavioral videos and photometry recordings were synchronized by TTL signals.

Raw calcium signals were pre-processed by removing the first minute of the recording

to decrease the effect of the initial exponential photobleaching, and by removing point

artifacts. The 470nM signal was fitted to the isosbestic 405nM using a linear fit and for

each time point, ΔF/F was calculated as (F470nm –F405nm(fitted))/F405nm(fitted).

This procedure is the same as described in previous works[24]. The codes used for the
analysis will be found on the dedicated github repository.
3

Chemogenetic modulation

Stereotaxic surgeries were performed as described above for fiber photometry. In this
case, C57BL/6J or PV-cre mice were injected with 500 nl AAV-CaMKII-mCherry (titer:
7×10^{12}, 114469-AAV5) and 500 nl AAV-CaMKII-hM4Di (titer:7×10^{12}, 50477-AAV2)
directly into the hippocampus, with the following coordinates: dorsal, AP ±1.5, ML ± 2,
DV -1.5 and ventral, AP ±3.5, ML ± 3.3, DV -3.5, according to Paxinos and Franklin
brain atlas (Paxinos and Franklin, 2001). On the other hand, PV-Cre mice were infused
with AAV-DIO-hM4Di using the same coordinates to target the dorsal hippocampus.
Three control groups were performed for each subregion (dorsal and ventral
hippocampus): animals infused with control virus (pAAV-CaMKIIa-mCherry) and
injected with saline or JHU37160 dihydrochloride (J60, 0.1 mg/Kg, i.p., HelloBio,
HB6261)[25], and animals infused with AAV-CaMKII-mCherry or AAV-DIO-hM4Di
injected with saline. Animals were used for chemogenetics experiments four weeks
after injections to get an optimal expression of the viruses. The injection of J60 was
one hour before each preconditioning session (C57BL/6J and PV-Cre mice), before the
Probe Test 1 (C57BL/6J mice) or the two conditioning sessions. In all mice used in the
behavioral experiments, the signal was checked, and representative images are shown
in Supplementary Figures.
22

Histology

After the chemogenetic and imaging experiments, mice were anesthetized i.p with a
mixture of ketamine (50 mg/Kg) and xylazine (20 mg/Kg) in overdose (3-4x body
weight), transcardially perfused with cold 4% paraformaldehyde to fix tissues. Brains
were removed and sectioned in serial coronal sections of 20 um and collected directly
to the slide for further analysis in the microscope. All sections were counterstained with
4',6-diamidino-2-phenylindole (DAPI, ref. 00-4959-52, Fluoromount-G w/DAPI, Life
Technologies, USA) to visualize cell nuclei in the mounting medium. Slides were
coverslipped and imaged by an epifluorescence Nikon Eclipse Ni-E. All animals were
signal checked to guarantee the target of the hippocampus subregion.
33

Data collection and statistical analysis

<u>Data collection</u>
All mice were randomly assigned to experimental conditions. Researchers performing
the experiments were always blind to the condition of the subject and we used an

automated way to analyze behavioral responses throughout the study. Raw data was

processed and analyzed using homemade python packages.

<u>Statistical analysis</u>

Graphs and statistical analysis were performed through homemade python scripts that

will be openly shared. All data comes from distinct mice and is shown as independent

data points per animal ± standard error of the median (SEM). Normality and

homoscedasticity of the data were assessed with the Kolmogorov-Smirnoff and Levene

tests, respectively. Due to non-parametric properties of the data, a generalized linear

model with gamma distribution was used for multivariable analysis (after a goodness of

fit to select the appropriate distribution), followed by planned comparisons corrected

with false discovery rate. For univariate analysis, the Kruskal-Wallis test was performed

followed by planned comparisons corrected with false discovery rate. For simple

comparisons the Mann-Whitney test was performed. Detailed statistical data for each

experiment can be found in Extended Table 1 (for Main Figures) and Extended Table 2

(for Supplementary Figures).

Results

Simultaneous light-tone associations during preconditioning are required for

sensory preconditioning responding in male mice.

Sensory preconditioning tasks using olfactory and gustatory stimuli have already been

demonstrated in mice in several previous studies[11,14]. However, other sensory

modalities such as visual and auditory stimuli are also highly relevant for animals' daily

decision-making. While the use of these cues to assess aversive sensory

preconditioning has been shown in rats[6,8,13], their use in mice—particularly for aversive

paradigms—remains limited[11]. The first aim of the present work was to establish a

sensory preconditioning task in mice using auditory and visual cues, enabling the

assessment of both mediated and direct learning. To this end, we developed a sensory

preconditioning protocol (see Methods, Figure 1A), alongside a novel automated tool to

analyze the freezing response (i.e., immobility) in mice. This tool combines

DeepLabCut pose estimation with a newly developed Python script and its accuracy

was validated showing a high correlation with manual scoring and ezTrack[23] freezing

assessments (Supplementary Figure 1). Although for freezing scoring our tool is similar

to ezTrack measurements, there are several key added benefits offered by this

approach such as an enhanced flexibility to combine freezing measurements with other

customized behavioral assessments, which can be particularly beneficial for labs

conducting large-scale or longitudinal behavioral studies in rodents.

Using both male and female mice, we conducted a long-term sensory preconditioning

($_{LT}$SPC) experiment with three experimental groups in male (Figure 1A) and female

mice (Supplementary Figure 2A): Paired group (protocol described in methods

section); Unpaired group (same protocol, but the light and tone were never

simultaneous); No-Shock group (same as the Paired group but without electric foot-

shock exposure) (Figure 1A and Supplementary Figure 2A). According to this design,

male (Figures 1B and 1C) and female mice (Supplementary Figure 2B and 2C) in the

Paired group exhibited an enhanced freezing response at the onset of the tone or light,

respectively. Specifically, exposure to the tone (Probe Test 1, ON period) led to

increased freezing behavior (Figure 1B) compared to the OFF period, indicating

sensory preconditioning responding in male mice. When the light was presented

(Probe Test 2), animals showed greater freezing during the ON period compared to the

OFF period (Figure 1C), revealing direct learning.

When the light and tone were temporally separated (Unpaired group), male mice failed

to exhibit sensory preconditioning (Figure 1B), while their direct learning response to

the light remained intact (Figure 1C). In contrast, female mice still displayed a behavioral response to the tone despite the separation of stimuli, which could be attributed to fear generalization (Supplementary Figure 2B), along with preserved direct learning (Supplementary Figure 2C). These results reveal sex differences in sensory preconditioning performance: male mice clearly demonstrated both mediated and direct learning, whereas female mice did not. To further explore this, we conducted an independent experiment to test whether the fear generalization observed in the female Unpaired group could be reduced by either extending the interval between cue presentations during preconditioning or by removing the light during the conditioning phase (Supplementary Figure 2D). Results showed that under both conditions, female mice still exhibited cue-induced fear generalization, which appears to mask their capacity for mediated learning responding.

Finally, in the No-Shock group, neither male (Figure 1B) nor female (Supplementary Figure 2B) mice showed a stimulus-mediated response, suggesting that tone or light exposure alone during the Probe Tests does not elicit behavioral changes. Altogether, these results confirm the successful establishment of an $_{LT}$SPC protocol in male mice, which can now be used to further investigate the brain circuits involved in higher-order conditioning. However, additional research will be required to optimize an $_{LT}$SPC protocol for female mice that avoids the confounding effect of fear generalization. Indeed, supporting these sex differences in this sensory preconditioning task, in male mice we observed a clear correlation between direct learning and mediated learning responding (Supplementary Figure 3A), whereas this is not shown in female mice (Supplementary Figure 3B).

Hippocampal cells are engaged during $_{LT}$SPC

Previous studies using sensory preconditioning paradigms with other sensory modalities have suggested a key role for the hippocampus in processing incidental associations between innocuous stimuli during the preconditioning phase[1,11]. Specifically, hippocampal GABAergic neurons expressing the type-1 cannabinoid receptors, but not the activity of PV interneurons, have been implicated in odor–taste associations in a mouse odor-taste sensory preconditioning task[26]. However, whether similar cellular mechanisms are involved in sensory preconditioning using different sensory modalities remains unknown. To better characterize the involvement of the dorsal and ventral hippocampus, as well as their specific cell types, during our $_{LT}$SPC task, we performed simultaneous in vivo fiber photometry recordings by implanting an optic fiber in each hippocampal subregion of PV-Cre mice and infusing a mixture of

AAV-Syn-RCaMP and AAV-DIO-GCaMP viruses (see *Methods*) in both hippocampal

subregions. This allowed us to monitor general neuronal activity (via RCaMP) and the

activity of PV-positive interneurons (via GCaMP) during both the preconditioning (light-

tone pairings) and the conditioning phase (light-foot-shock pairings). Four weeks after

surgery, animals were subjected to the $_{LT}$SPC task (Figure 2A). Neuronal activity in the

dorsal and ventral hippocampus was quantified as ΔF/F following stimulus pairings

(dHPC: Figure 2B on left, vHPC: Figure 2C on left). Peri-event time histograms

(PETHs) of Z-scored ΔF/F revealed the average dynamic response in each subregion

during light–tone associations (dHPC: Figure 2B on right, vHPC: Figure 2C on right).

These PETHs highlight a lower trial-by-trial variability in dHPC (Figure 2B) and a more

sustained response over time compared to vHPC (Figure 2C). To quantify this activity,

we measured the peak Z-scored ΔF/F within a 1-second window following stimulus

onset and compared it to baseline values (Figure 2D). Notably, this increase in activity

was consistent across all preconditioning sessions, with no significant differences

observed when data were analyzed per session (Supplementary Figure 4A).

Additionally, both neuronal activity markers (RCaMP and GCaMP) showed clear

increases during the conditioning phase, both at the onset of the light stimulus

(conditioned stimulus; Supplementary Figure 5) and at the time of foot-shock delivery

(unconditioned stimulus; Supplementary Figure 6). These increases were consistent

across sessions (Supplementary Figure 4B and C), with no significant session-

dependent variation. Together, these findings demonstrate the engagement of

hippocampal neurons, including PV-positive interneurons, in both the dHPC and vHPC

during associative learning in our $_{LT}$SPC task. This supports the role of the

hippocampus in processing sensory associations involving non-olfactory modalities.

Although our data showed a clear engagement of hippocampal activity at different

phases of our $_{LT}$SPC task, the use of a synapsin promoter is too broad to selectively

point to a specific cell-type mediating the behavioral responses observed. Thus, we

decided to use a similar approach and specifically monitor CaMKII+ neurons during the

preconditioning sessions by infusing an AAV- CaMKII -GCAMP6 (see Methods, Figure

3A). Interestingly, and matching what was observed with a pan promoter (*i.e.*

synapsin), PETHs of Z-scored ΔF/F (Figure 3B and C) and the subsequent histograms

representing the peak Z-scored ΔF/F within a 1-second window following stimulus

onset (Figure 3D) revealed a clear engagement of CaMKII-positive neurons from both

hippocampal subregions during light–tone associations, which is constant across

preconditioning sessions (Figure 3E).

CaMKII-positive neurons in dorsal hippocampus mediate $_{LT}$SPC

The in vivo photometry results suggest a prominent engagement of the dorsal

hippocampal (dHPC) subregion, characterized by increased activity of various

hippocampal cell types, including CaMKII-positive neurons, during light–tone

presentations in the preconditioning phase. However, no study to date has explored a

potential causal dissociation between the roles of the dHPC and vHPC in mouse

sensory preconditioning. To investigate this, adeno-associated viral vectors expressing

an inhibitory DREADD (hM4DGi, hereafter referred to as DREADD-Gi)[27] under the

CaMKII promoter were infused into either the dHPC or vHPC to selectively inhibit

principal excitatory neurons in these hippocampal subregions (Figure 4A). Control

animals, which received either saline or the DREADD agonist J60, were pooled as they

exhibited no significant behavioral differences and showed a reliable mediated and

direct learning in the $_{LT}$SPC task (Figure 4B and, Supplementary Figure 7A-D).

Notably, inhibition of CaMKII-positive neurons in the dHPC, which decrease calcium

activity in CaMKII-positive neurons (Supplementary Figure 7E and F), during the

preconditioning phase (i.e., J60 administration in DREADD-Gi mice) completely

abolished mediated learning (Figure 4B) without affecting direct learning (Figure 4D).

This effect was specific to the preconditioning phase, as inhibition prior to conditioning

sessions (Supplementary Figure 7G) or before Probe Test 1 (Figure 4B) did not impair

mediated learning. In contrast, inhibition of CaMKII-positive neurons in the vHPC,

whether during preconditioning or Probe Test 1, had no effect on either mediated

(Figure 4C) or direct learning (Figure 4E). These findings indicate that the dHPC, but

not the vHPC, plays a specific and causal role in encoding associations between

innocuous stimuli (e.g., light and tone) that underlie sensory preconditioning responses.

Given the additional results observed in our photometry experiments and the crucial

role of dHPC, we next investigated whether PV-positive interneurons, which were also

activated during stimuli presentation in our photometry recordings, played a role in this

behavior. To test this, a Cre-dependent DREADD-Gi virus was infused into the dHPC

of PV-Cre mice to allow for selective inhibition of PV interneurons (Supplementary

Figure 8A-C) during light–tone pairings. Remarkably, silencing PV interneurons during

the preconditioning phase did not affect either mediated or direct learning

(Supplementary Figure 8D).

Taken together, these results suggest that CaMKII-positive principal neurons, but not

PV interneurons, in the dHPC, are critical for encoding light–tone associations that

drive sensory preconditioning responses in mice. Importantly, no statistical behavioral

differences were observed during the conditioning across the different experiments

(Supplementary Figure 9).

Discussion

Our study provides novel insights into the role of the hippocampus in higher-order

conditioning by establishing and validating a light-tone sensory preconditioning ($_{LT}$SPC)

task in male mice. Using in vivo calcium imaging, we linked the activity of specific

hippocampal cells in the dorsal and ventral subregions to distinct behavioral phases of

the task. Furthermore, chemogenetic inhibition experiments allowed us to causally

identify the dorsal hippocampus (dHPC)—but not the ventral hippocampus (vHPC)—as

essential for encoding light–tone pairings, which are critical for the behavioral

outcomes observed in sensory preconditioning.

Sex differences in classical fear conditioning have been reported, with conflicting

findings showing stronger conditioning in males, in females, or no differences at all[28-35].

However, to our knowledge, no previous studies have examined sex differences in

sensory preconditioning paradigms. Our results demonstrate that, using the same

preconditioning and conditioning settings, only male mice show a reliable light-tone

sensory preconditioning responding, whereas female mice show behavioral responses

mostly associated to fear generalization. Indeed, our results could suggest that female

mice are more vulnerable to aversive stimuli responding, which might be consistent

with prior evidence showing sex-dependent differences in conditioned freezing

responses[36-39]. When the light and the tone were presented separately during the

preconditioning, or when the foot-shock was omitted, both mediated and direct learning

were abolished in male mice. Instead, female mice still showed reduced but detectable

cue-induced responses under these conditions, even if we increased the time between

light and tone exposures during the preconditioning phase. Thus, we did not

successfully set up a light-tone sensory preconditioning paradigm for female mice and

future research will be required to optimize and adapt a protocol to reliably observe

mediated learning in females. Given that female mice also tend to exhibit greater fear

generalization[29,40,41], the observed differences in the unpaired group could reflect this

enhanced generalization[33]. In this regard, changes on the number of conditioning

sessions and/or trials, intensity of the electric foot-shock or additional contextual

changes could avoid this fear generalization in future sensory preconditioning

paradigms in female mice. In addition, an interesting unresolved issue is how sex

hormones could alter sensory preconditioning responding or the different performance

of male and female mice in different protocols. Overall, our data clearly revealed the

necessity to adapt behavioral protocols for each sex in sensory preconditioning

paradigms involving auditory and visual stimuli and aversive conditioning.

Prior research has established the hippocampus as critical for encoding spatial

location, context, cues, and memory related to appetitive and aversive behaviors[42-46].

However, few studies have addressed its causal involvement in reinforcement

learning[47], and even fewer have explored its role in forming associations between

innocuous stimuli. Using in vivo fiber photometry, we observed increased neuronal

activity in both the dorsal and ventral hippocampus during light–tone pairings and

during light–shock associations. These findings support the notion that the

hippocampus is actively engaged in encoding both conditioned (CS) and unconditioned

(US) stimuli, in agreement with studies showing hippocampal activation during

reinforcement behavior[47-49]. Still, some prior work has argued that the hippocampus is

not required for forming simple cue–US associations, which may instead rely on the

amygdala[50].

Our photometry results show that the simultaneous presentation of sensory cues is

encoded by the hippocampus, reinforcing its role in associative learning and sensory

integration[51]. This is consistent with previous work in both animals and humans

supporting hippocampal involvement during associative phases of sensory

preconditioning[1,11,13,14,52]. Importantly, however, none of these earlier studies

investigated the specific roles of hippocampal subregions or cell types. Our data

indicate that modulation of hippocampal activity—specifically in the dHPC—affects

encoding during the preconditioning phase but not during conditioning or retrieval

phases. Chemogenetic inhibition of CaMKII-positive neurons in the dHPC during the

preconditioning blocked mediated learning responding, while inhibition in the vHPC had

no such effect. Moreover, silencing PV-positive interneurons in the dHPC during

preconditioning did not disrupt sensory preconditioning, suggesting that principal

excitatory neurons, but not PV interneurons, are essential for forming associations

between innocuous stimuli. Indeed, it is somehow contradictory to observe an activity

enhancement of PV-positive interneurons during light-tone associations and the lack of

effect of its inhibition during this phase. To fully discard any role of the activity of these

interneurons during this phase, future experiments should aim to particularly inhibit

these cells during restricted temporal windows such as some preconditioning sessions

rather than the full preconditioning phase or to specifically inhibit its activity during the

light-tone exposures (e.g. close-loop optogenetic approaches).

Other brain regions, including the orbitofrontal, retrosplenial, and perirhinal cortices,

have also been implicated in processing associations between neutral sensory cues[6-]

[8,10]. These areas maintain reciprocal anatomical connections with the hippocampus[53-55], suggesting a broader, hippocampus-guided network for encoding higher-order associations. Our findings showing that the dHPC—but not the vHPC—is crucial for encoding light–tone pairings may offer new insight into this circuit. A likely candidate for mediating such interactions is the retrosplenial cortex, which receives direct projections from the dHPC[56]. Thus, inhibiting CaMKII-positive neurons in the dHPC, which project to the retrosplenial cortex could disrupt sensory preconditioning, supporting this potential circuit model. Future studies will be necessary to explore this hypothesis in detail.

Despite the robust findings presented by our data, several limitations must be acknowledged. First, while we successfully validated the $_{LT}$SPC protocol in male mice, further optimization is needed for its reliable use in female mice as commented above. Second, the fixed order of Probe Tests may have introduced extinction effects for the tone[57], potentially affecting responses to the light cue. Future studies should counterbalance Probe Tests to address this possibility. Third, our behavioral protocol required context changes between experimental phases, which may have engaged hippocampal circuits[58] more strongly than if a single context had been used. Thus, the impact of context-switching on hippocampal involvement should be addressed in future work. Finally, this work has been focused on the activity of CaMKII-positive neurons and PV-interneurons although we cannot discard the role of other hippocampal cell types such as somatostatin interneurons, which have been recently involved in learning and associative processes[59,60]. Future experiments will be required to fully characterize the role of different hippocampal subtypes in sensory preconditioning. Indeed, other techniques such as the optogenetic modulation of cell activity could give more details on the temporal engagement of these hippocampal circuits in higher-order conditioning.

In summary, our results identify a key role for hippocampal circuits—particularly CaMKII-positive neurons in the dorsal hippocampus—in mediating associations between innocuous stimuli during light-tone sensory preconditioning. This process reflects a fundamental cognitive mechanism underpinning higher-order learning and decision-making in both animals and humans. Understanding the neural substrates of sensory preconditioning is crucial, as this form of learning may contribute to adaptive behavior but also underlie maladaptive processes seen in disorders such as psychosis[61,62].

Acknowledgements

We would like to thank the personnel of the Animal Facility of the Parc de
Recerca Biomedica de Barcelona (PRBB) for mouse care. We thank all the members
of our lab for useful discussions during the development of the project and Remi
Proville (Aquineuro, Bordeaux, France) for the great help in the analysis of behavior
and *in vivo* photometry. Finally, we would like to also thank Dr. Maria Victoria Puig
Velasco for providing the PV-Cre colony. This work was supported by la Generalitat de
Catalunya (SGR (00022) and "Jo Investigo" (2022 INV-1 00005/100005TG3)
programmes) from the Departament d'Economia i Coneixement de la Generalitat de
Catalunya (Spain) and from the European Research Council (ERC) under the
European Union's Horizon 2020 research and innovation programme (Grant
agreement No. [948217]).

Author Contributions

15 A.B-G. and J.P. contributed to the conception of the project and performed and
16 analyzed all the experiments. C.R-D. helped on all surgical procedures and
17 immunohistochemistry   assays.   I.M-S.   helped   with   behavioral   and
18 immunohistochemistry assays. A.B-G. and J.P wrote the manuscript. All authors have
19 corrected and revised the manuscript.
20

21 **Conflict of interest**
22 The authors declare no conflict of interest.
23
24

Figure Legends

Figure 1.   Simultaneous associations between light and tone are required for sensory preconditioning responding. (A) Schematic representation of the $_{LT}$SPC task in males (2 training sessions) with a representation of paired, unpaired and no-shock experimental groups. The temporal dynamics of freezing represented in bins of 10 seconds across time of experiment of the Probe Test 1 (Tone) (B on the left) and Probe Test 2 (light) (C on the left). The percentage of time spent freezing during OFF and ON periods of the Probe Test 1 (Tone) (B on the right) and Probe Test 2 (light) (C on the right). * Significant p-value (<0.05) after false discovery rate (FDR). GLM: generalized linear model fitted to gamma distribution with planned comparisons. KW: Kruskal-Wallis across experimental groups: paired, unpaired, no-shock. See statistical details in Supplementary Table 1.

Figure 2. Hippocampal cell activity during $_{LT}$SPC. A) Schematic representation of $_{LT}$SPC task while fiber photometry recordings of RCaMP (calcium sensor in synapsin-positive neurons) and Cre-dependent GCaMP (calcium sensor in PV-positive interneurons) in dHPC and vHPC of PV^{cre} mice. B) dHPC modulation during preconditioning of $_{LT}$SPC: on the left upper panel z scores of $\frac{\Delta f}{f}$ (where f represents fluorescence) of GCaMP PV-positive interneurons on dHPC (green), left bottom z scores of $\frac{\Delta f}{f}$ of RCaMP in neurons of dHPC (red), right upper number of events with a $\frac{\Delta f}{f} > 2$ of GCaMP PV-positive interneurons on dHPC, and right bottom number of trials with a $\frac{\Delta f}{f} > 2$ of Rcamp in neurons of dHPC. C) vHPC modulation during $_{LT}$SPC: on left upper painel z scores of $\frac{\Delta f}{f}$ of GCaMP PV-positive inter neurons of vHPC (green), left bottom z scores of $\frac{\Delta f}{f}$ of RCaMP in neurons of vHPC (red), right upper number of trials with a $\frac{\Delta f}{f} > 2$ of Gcamp in PV-positive interneurons of vHPC, and right bottom number of trials with a $\frac{\Delta f}{f} > 2$ of RCaMP in neurons of vHPC. D) maximal value of $\frac{\Delta f}{f}$ in the first 1 second window after pairings compared with baseline, on the left (dHPC) and on the right (vHPC). * Significant p-value (<0.05). See statistical details in Supplementary Table 1.

Figure 3. CaMKII neurons during preconditioning on dHPC and vHPC. A) maximal

value of $\frac{\Delta f}{f}$ in the first 1 second window after stimulus compared with baseline, for each

stimulus during preconditioning session, on the left (dHPC) and on the light (vHPC). B)

dHPC modulation during preconditioning of $_{LT}$SPC: on left z scores of $\frac{\Delta f}{f}$ (where f

represents fluorescence) of GCaMP in CaMKII-positive neurons on dHPC (green), on

right number of events with a $\frac{\Delta f}{f} > 2$ of GCaMP CaMKII-positive neurons on dHPC. C)

vHPC modulation during preconditioning of $_{LT}$SPC: on left z scores of $\frac{\Delta f}{f}$ (where f

represents fluorescence) of GCaMP in CaMKII-positive neurons on vHPC (green), on

right number of events with a $\frac{\Delta f}{f} > 2$ of GCaMP CaMKII-positive neurons on vHPC. D)

maximal value of $\frac{\Delta f}{f}$ in the first 1 second window after stimulus compared with baseline,

for the average of the 6 pairings stimuli during preconditioning session, on the left

(dHPC) and on the light (vHPC). *, p-value (<0.05). Statistical details in Supplementary

Table 2.

Figure 4. Chemogenetic modulation of dorsal and ventral hippocampus during

$_{LT}$SPC. (A) Schematic representation of $_{LT}$SPC task combined with chemogenetic

approaches, where an inhibitory DREADD was infused in dHPC (representative image

on the left) and vHPC (representative image on the right). During the preconditioning

(DPC) or the Probe Test (DPT), a DREADD agonist (J60) was injected intraperitoneally

(controls where injected with the agonist in both phases) (diagram of DREADD agonist

administration on the center). (B,C,D,E on the left) The temporal dynamics of freezing

represented in bins of 10 seconds across time of experiment of the Probe Test 1

(Tone) of dHPC (B on the left) and vHPC (C on the left) and Probe Test 2 (light) of

dHPC (D on the left) and vHPC (E on the left). (B, C on the right) The percentage of

time spent in freezing during OFF and ON periods (left) of the Probe Test 1 (Tone) of

animals infused in dHPC (B on the right) and vHPC (C on the right) D, E). (D, E on the

right) The percentage of time spent in freezing during OFF and ON periods (left) of the

Probe Test 2 (Light) of animals infused in dHPC (D on the right) and vHPC (E on the

right). Controls are labeleld as Constrols_d (on dHPC) and Controls_V (on vHPC). The

preconditioning groups are labelled as DPCd (on dHPC) and DPCv (on vHPC). The

Probe Test groups are labelled as DPTd (on dHPC) and DPTv (on vHPC). *Significant

p-value (<0.05) after false discovery rate (FDR). GLM: generalized linear model fitted

to gamma distribution with planned comparisons. See statistical details in

Supplementary Table 1.

References

Wimmer, G. E. & Shohamy, D. Preference by association: how memory mechanisms in the hippocampus bias decisions. *Science* **338**, 270-273 (2012). https://doi.org:10.1126/science.1223252

Gewirtz, J. C. & Davis, M. Using pavlovian higher-order conditioning paradigms to investigate the neural substrates of emotional learning and memory. *Learn Mem* **7**, 257-266 (2000).

Parkes, S. L. & Westbrook, R. F. Role of the basolateral amygdala and NMDA receptors in higher-order conditioned fear. *Rev Neurosci* **22**, 317-333 (2011). https://doi.org:10.1515/RNS.2011.025

Holmes, N. M., Fam, J. P., Clemens, K. J., Laurent, V. & Westbrook, R. F. The neural substrates of higher-order conditioning: A review. *Neurosci Biobehav Rev* **138**, 104687 (2022). https://doi.org:10.1016/j.neubiorev.2022.104687

Ioannidou, C., Busquets-Garcia, A., Ferreira, G. & Marsicano, G. Neural Substrates of Incidental Associations and Mediated Learning: The Role of Cannabinoid Receptors. *Front Behav Neurosci* **15**, 722796 (2021). https://doi.org:10.3389/fnbeh.2021.722796

Holmes, N. M., Parkes, S. L., Killcross, A. S. & Westbrook, R. F. The basolateral amygdala is critical for learning about neutral stimuli in the presence of danger, and the perirhinal cortex is critical in the absence of danger. *J Neurosci* **33**, 13112-13125 (2013). https://doi.org:10.1523/JNEUROSCI.1998-13.2013

Parkes, S. L. & Westbrook, R. F. The basolateral amygdala is critical for the acquisition and extinction of associations between a neutral stimulus and a learned danger signal but not between two neutral stimuli. *J Neurosci* **30**, 12608-12618 (2010). https://doi.org:10.1523/JNEUROSCI.2949-10.2010

Robinson, S. *et al.* Chemogenetic silencing of neurons in retrosplenial cortex disrupts sensory preconditioning. *J Neurosci* **34**, 10982-10988 (2014). https://doi.org:10.1523/JNEUROSCI.1349-14.2014

Gostolupce, D., Lay, B. P. P., Maes, E. J. P. & Iordanova, M. D. Understanding Associative Learning Through Higher-Order Conditioning. *Front Behav Neurosci* **16**, 845616 (2022). https://doi.org:10.3389/fnbeh.2022.845616

Hart, E. E., Sharpe, M. J., Gardner, M. P. & Schoenbaum, G. Responding to preconditioned cues is devaluation sensitive and requires orbitofrontal cortex during cue-cue learning. *Elife* **9** (2020). https://doi.org:10.7554/eLife.59998

Busquets-Garcia, A. *et al.* Hippocampal CB(1) Receptors Control Incidental Associations. *Neuron* **99**, 1247-1259 e1247 (2018). https://doi.org:10.1016/j.neuron.2018.08.014

Kahnt, T. & Schoenbaum, G. Cross-species studies on orbitofrontal control of inference-based behavior. *Behav Neurosci* **135**, 109-119 (2021). https://doi.org:10.1037/bne0000401

Iordanova, M. D., Good, M. & Honey, R. C. Retrieval-mediated learning involving episodes requires synaptic plasticity in the hippocampus. *J Neurosci* **31**, 7156-7162 (2011). https://doi.org:10.1523/JNEUROSCI.0295-11.2011

Wheeler, D. S., Chang, S. E. & Holland, P. C. Odor-mediated taste learning requires dorsal hippocampus, but not basolateral amygdala activity. *Neurobiol Learn Mem* **101**, 1-7 (2013). https://doi.org:10.1016/j.nlm.2012.12.015

Caroni, P. Inhibitory microcircuit modules in hippocampal learning. *Curr Opin Neurobiol* **35**, 66-73 (2015). https://doi.org/10.1016/j.conb.2015.06.010

Chevaleyre, V. & Piskorowski, R. Modulating excitation through plasticity at inhibitory synapses. *Front Cell Neurosci* **8**, 93 (2014). https://doi.org/10.3389/fncel.2014.00093

Fanselow, M. S. & Dong, H. W. Are the dorsal and ventral hippocampus functionally distinct structures? *Neuron* **65**, 7-19 (2010). https://doi.org/10.1016/j.neuron.2009.11.031

Strange, B. A., Witter, M. P., Lein, E. S. & Moser, E. I. Functional organization of the hippocampal longitudinal axis. *Nat Rev Neurosci* **15**, 655-669 (2014). https://doi.org/10.1038/nrn3785

Jones, B. F. & Witter, M. P. Cingulate cortex projections to the parahippocampal region and hippocampal formation in the rat. *Hippocampus* **17**, 957-976 (2007). https://doi.org/10.1002/hipo.20330

Galvez-Marquez, D. K. *et al.* Spatial contextual recognition memory updating is modulated by dopamine release in the dorsal hippocampus from the locus coeruleus. *Proc Natl Acad Sci U S A* **119**, e2208254119 (2022). https://doi.org/10.1073/pnas.2208254119

Moser, E., Moser, M. B. & Andersen, P. Spatial learning impairment parallels the magnitude of dorsal hippocampal lesions, but is hardly present following ventral lesions. *J Neurosci* **13**, 3916-3925 (1993). https://doi.org/10.1523/JNEUROSCI.13-09-03916.1993

Groenewegen, H. J., Vermeulen-Van der Zee, E., te Kortschot, A. & Witter, M. P. Organization of the projections from the subiculum to the ventral striatum in the rat. A study using anterograde transport of Phaseolus vulgaris leucoagglutinin. *Neuroscience* **23**, 103-120 (1987). https://doi.org/10.1016/0306-4522(87)90275-2

Pennington, Z. T. *et al.* ezTrack-A Step-by-Step Guide to Behavior Tracking. *Curr Protoc* **1**, e255 (2021). https://doi.org/10.1002/cpz1.255

Lerner, T. N. *et al.* Intact-Brain Analyses Reveal Distinct Information Carried by SNc Dopamine Subcircuits. *Cell* **162**, 635-647 (2015). https://doi.org/10.1016/j.cell.2015.07.014

Bonaventura, J. *et al.* High-potency ligands for DREADD imaging and activation in rodents and monkeys. *Nat Commun* **10**, 4627 (2019). https://doi.org/10.1038/s41467-019-12236-z

Busquets-Garcia, A. *et al.* Hippocampal CB1 Receptors Control Incidental Associations. *Neuron* **99**, 1247-1259 e1247 (2018). https://doi.org/10.1016/j.neuron.2018.08.014

Robinson, S. & Adelman, J. S. A Method for Remotely Silencing Neural Activity in Rodents During Discrete Phases of Learning. *J Vis Exp*, e52859 (2015). https://doi.org/10.3791/52859

Moore, M. D., Cushman, J., Chandra, D., Homanics, G. E., Olsen, R. W. & Fanselow, M. S. Trace and contextual fear conditioning is enhanced in mice lacking the alpha4 subunit of the GABA(A) receptor. *Neurobiol Learn Mem* **93**, 383-387 (2010). https://doi.org/10.1016/j.nlm.2009.12.004

Keiser, A. A., Turnbull, L. M., Darian, M. A., Feldman, D. E., Song, I. & Tronson, N. C. Sex Differences in Context Fear Generalization and Recruitment of Hippocampus and Amygdala during Retrieval. *Neuropsychopharmacology* **42**, 397-407 (2017). https://doi.org/10.1038/npp.2016.174

Anagnostaras, S. G. *et al.* Testicular hormones do not regulate sexually dimorphic Pavlovian fear conditioning or perforant-path long-term potentiation in adult male rats. *Behav Brain Res* **92**, 1-9 (1998). https://doi.org/10.1016/s0166-4328(97)00115-0

Maren, S., De Oca, B. & Fanselow, M. S. Sex differences in hippocampal long-term potentiation (LTP) and Pavlovian fear conditioning in rats: positive

correlation between LTP and contextual learning. *Brain Res* **661**, 25-34 (1994). https://doi.org/10.1016/0006-8993(94)91176-2

Colon, L., Odynocki, N., Santarelli, A. & Poulos, A. M. Sexual differentiation of contextual fear responses. *Learn Mem* **25**, 230-240 (2018). https://doi.org/10.1101/lm.047159.117

Urien, L., Stein, N., Ryckman, A., Bell, L. & Bauer, E. P. Extended amygdala circuits are differentially activated by context fear conditioning in male and female rats. *Neurobiol Learn Mem* **180**, 107401 (2021). https://doi.org/10.1016/j.nlm.2021.107401

Kosten, T. A., Lee, H. J. & Kim, J. J. Early life stress impairs fear conditioning in adult male and female rats. *Brain Res* **1087**, 142-150 (2006). https://doi.org/10.1016/j.brainres.2006.03.009

Dachtler, J., Fox, K. D. & Good, M. A. Gender specific requirement of GluR1 receptors in contextual conditioning but not spatial learning. *Neurobiol Learn Mem* **96**, 461-467 (2011). https://doi.org/10.1016/j.nlm.2011.07.001

Colom-Lapetina, J., Li, A. J., Pelegrina-Perez, T. C. & Shansky, R. M. Behavioral Diversity Across Classic Rodent Models Is Sex-Dependent. *Front Behav Neurosci* **13**, 45 (2019). https://doi.org/10.3389/fnbeh.2019.00045

Marcinkiewcz, C. A. *et al.* Sex-Dependent Modulation of Anxiety and Fear by 5-HT(1A) Receptors in the Bed Nucleus of the Stria Terminalis. *ACS Chem Neurosci* **10**, 3154-3166 (2019). https://doi.org/10.1021/acschemneuro.8b00594

Shansky, R. M. & Murphy, A. Z. Considering sex as a biological variable will require a global shift in science culture. *Nat Neurosci* **24**, 457-464 (2021). https://doi.org/10.1038/s41593-021-00806-8

Gruene, T. M., Roberts, E., Thomas, V., Ronzio, A. & Shansky, R. M. Sex-specific neuroanatomical correlates of fear expression in prefrontal-amygdala circuits. *Biol Psychiatry* **78**, 186-193 (2015). https://doi.org/10.1016/j.biopsych.2014.11.014

Day, H. L. L., Reed, M. M. & Stevenson, C. W. Sex differences in discriminating between cues predicting threat and safety. *Neurobiol Learn Mem* **133**, 196-203 (2016). https://doi.org/10.1016/j.nlm.2016.07.014

Krueger, J. N. & Sangha, S. On the basis of sex: Differences in safety discrimination vs. conditioned inhibition. *Behav Brain Res* **400**, 113024 (2021). https://doi.org/10.1016/j.bbr.2020.113024

Moser, M. B., Moser, E. I., Forrest, E., Andersen, P. & Morris, R. G. Spatial learning with a minislab in the dorsal hippocampus. *Proc Natl Acad Sci U S A* **92**, 9697-9701 (1995). https://doi.org/10.1073/pnas.92.21.9697

Trouche, S. *et al.* Recoding a cocaine-place memory engram to a neutral engram in the hippocampus. *Nat Neurosci* **19**, 564-567 (2016). https://doi.org/10.1038/nn.4250

Trouche, S. *et al.* A Hippocampus-Accumbens Tripartite Neuronal Motif Guides Appetitive Memory in Space. *Cell* **176**, 1393-1406 e1316 (2019). https://doi.org/10.1016/j.cell.2018.12.037

Dragoi, G., Harris, K. D. & Buzsaki, G. Place representation within hippocampal networks is modified by long-term potentiation. *Neuron* **39**, 843-853 (2003). https://doi.org/10.1016/s0896-6273(03)00465-3

Robinson, N. T. M. *et al.* Targeted Activation of Hippocampal Place Cells Drives Memory-Guided Spatial Behavior. *Cell* **183**, 1586-1599 e1510 (2020). https://doi.org/10.1016/j.cell.2020.09.061

Ibrahim, K. M. *et al.* Dorsal hippocampus to nucleus accumbens projections drive reinforcement via activation of accumbal dynorphin neurons. *Nat Commun* **15**, 750 (2024). https://doi.org/10.1038/s41467-024-44836-9

Ursin, R., Ursin, H. & Olds, J. Self-stimulation of hippocampus in rats. *J Comp Physiol Psychol* **61**, 353-359 (1966). https://doi.org/10.1037/h0023253

van der Kooy, D., Fibiger, H. C. & Phillips, A. G. Monoamine involvement in hippocampal self-stimulation. *Brain Res* **136**, 119-130 (1977). https://doi.org:10.1016/0006-8993(77)90136-6

Kim, J. J. & Fanselow, M. S. Modality-specific retrograde amnesia of fear. *Science* **256**, 675-677 (1992). https://doi.org:10.1126/science.1585183

Voss, J. L. & Cohen, N. J. Hippocampal-cortical contributions to strategic exploration during perceptual discrimination. *Hippocampus* **27**, 642-652 (2017). https://doi.org:10.1002/hipo.22719

Talk, A. C., Gandhi, C. C. & Matzel, L. D. Hippocampal function during behaviorally silent associative learning: dissociation of memory storage and expression. *Hippocampus* **12**, 648-656 (2002). https://doi.org:10.1002/hipo.10098

Agster, K. L. & Burwell, R. D. Hippocampal and subicular efferents and afferents of the perirhinal, postrhinal, and entorhinal cortices of the rat. *Behav Brain Res* **254**, 50-64 (2013). https://doi.org:10.1016/j.bbr.2013.07.005

Witter, M. P., Kleven, H. & Kobro Flatmoen, A. Comparative Contemplations on the Hippocampus. *Brain Behav Evol* **90**, 15-24 (2017). https://doi.org:10.1159/000475703

Ritchey, M., Libby, L. A. & Ranganath, C. Cortico-hippocampal systems involved in memory and cognition: the PMAT framework. *Prog Brain Res* **219**, 45-64 (2015). https://doi.org:10.1016/bs.pbr.2015.04.001

Opalka, A. N. & Wang, D. V. Hippocampal efferents to retrosplenial cortex and lateral septum are required for memory acquisition. *Learn Mem* **27**, 310-318 (2020). https://doi.org:10.1101/lm.051797.120

Polack, C. W., Molet, M., Miguez, G. & Miller, R. R. Associative structure of integrated temporal relationships. *Learn Behav* **41**, 443-454 (2013). https://doi.org:10.3758/s13420-013-0119-5

Keinath, A. T., Mosser, C. A. & Brandon, M. P. The representation of context in mouse hippocampus is preserved despite neural drift. *Nat Commun* **13**, 2415 (2022). https://doi.org:10.1038/s41467-022-30198-7

Abbas, A. I. *et al.* Somatostatin Interneurons Facilitate Hippocampal-Prefrontal Synchrony and Prefrontal Spatial Encoding. *Neuron* **100**, 926-939 e923 (2018). https://doi.org:10.1016/j.neuron.2018.09.029

Lacagnina, A. F. *et al.* Ventral hippocampal interneurons govern extinction and relapse of contextual associations. *Cell Rep* **43**, 114880 (2024). https://doi.org:10.1016/j.celrep.2024.114880

Busquets-Garcia, A. *et al.* Pregnenolone blocks cannabinoid-induced acute psychotic-like states in mice. *Mol Psychiatry* **22**, 1594-1603 (2017). https://doi.org:10.1038/mp.2017.4

Busquets-Garcia, A., Soria-Gomez, E., Ferreira, G. & Marsicano, G. Representation-mediated Aversion as a Model to Study Psychotic-like States in Mice. *Bio Protoc* **7** (2017). https://doi.org:10.21769/BioProtoc.2358

A

AAV-CaMKII-GCaMP6m
AAV-CamKIIa-hM4D(Gi)-mCherry

Dorsal | Ventral

C57BL/6J

Dorsal Hippocampus   Ventral Hippocampus

B   Dorsal Hippocampus

C   Ventral Hippocampus

D

dHPC   vHPC

E

dHPC - GCaMP   vHPC - GCaMP