AAVone: A Cost-Effective, Single-Plasmid Solution for Efficient AAV Production with Reduced

DNA Impurities

Rongze Yang,[1]* Ngoc Tam Tran,[2]* Taylor Chen,[2] Mengtian Cui,[2] Yuyan Wang,[1] Tapan Sharma,[2] Yu

Liu,[1] Jiantao Zhang,[1] Xinxu Yuan,[4] Danmeng Zhang,[4] Cuiping Chen,[1] Zhen Shi,[1] Leming Wang,[1] Yuling

Dai,[1] Haniya Zaidi,[1] Jiarui Liang,[1] May Chen[1], Dabbu Jaijyan,[4] Huan Hu,[1] Bing Wang,[1] Cheng Xu,[1]

Wenhui Hu,[4] Guangping Gao,[2,3] Daozhan Yu,[1] Phillip W.L. Tai,[2,3#] and Qizhao Wang[1#]

[1]AAVnerGene Inc, 9620 Medical Center Dr, Suite 100, Rockville, MD, USA;

[2]Horae Gene Therapy Center, UMass Chan Medical School, Worcester, Massachusetts, USA;

[3]Department of Microbiology, UMass Chan Medical School, Worcester, Massachusetts, USA.

[4]Department of Anatomy and Neurobiology, Virginia Commonwealth University, Richmond, VA 23298,

USA

*Co-first authors

#Correspondences should be addressed to:

P.W.L.T. (Phillip.Tai2@umassmed.edu)   Q.W. (Qizhao.Wang@aavnergene.com)

UMass Chan Medical School   AAVnerGene Inc.

New Education and Research Building (N8-1039) 9620 Medical Center Dr, Suite 100

366 Plantation Street   Rockville, MD 20850

Worcester, MA 01605   (267)-240-3270

Office phone: (774) 455-4559

18 **Short title: AAVone system for efficient AAV vector production**

ABSTRACT

Currently, the most common approach for manufacturing GMP-grade adeno-associated virus (AAV) vectors involves transiently transfecting mammalian cells with three plasmids that carry the essential components for production. The requirement for all three plasmids to be transfected into a single cell and the necessity for high quantities of input plasmid DNA, limits AAV production efficiency, introduces variability between production batches, and increases time and labor costs. Here, we developed an all-in-one, single-plasmid AAV production system, called AAVone. In this system, the adenovirus helper genes (*E2A*, *E4orf6*, and *VA RNA*), packaging genes (*rep* and *cap*), and the vector transgene cassette are consolidated into a single compact plasmid with a 13-kb backbone. The AAVone system achieves a two- to four-fold increase in yields compared to the traditional triple-plasmid system. Furthermore, the AAVone system exhibits low batch-to-batch variation and eliminates the need for fine-tuning the ratios of the three plasmids, simplifying the production process. In terms of vector quality, AAVs generated by the AAVone system show similar *in vitro* and *in vivo* transduction efficiency, but a substantial reduction in sequences attributed to plasmid backbones and a marked reduction in non-functional snap-back genomes. In Summary, the AAVone platform is a straightforward, cost-effective, and highly consistent AAV production system – making it particularly suitable for GMP-grade AAV vectors.

INTRODUCTION

Adeno-associated virus (AAV) vectors have become the preferred gene delivery vehicles because of their efficient transduction, long-term gene expression, low immunogenicity, and favorable safety profile. They have been successfully utilized in numerous clinical trials and have received regulatory approvals as drugs for human use.[1-3] Although AAV production systems based on herpes simplex virus, vaccinia virus, and adenovirus (AdV) have been developed, they have not been widely adopted because of concerns related to potency, quality, safety, and processing complexity.[4-11] The mainstay technology for AAV vector production with human cell lines (HEK293) is still the triple-plasmid (tri-plasmid) transfection method, which involves co-transfecting cells with three plasmids containing the necessary *cis* and *trans* components.[12] The inverted terminal repeats (ITRs) are the only *cis* elements of viral origin and are required for AAV replication and packaging. Within the AAV vector plasmid that carries the transgene cassette, also called the *cis* plasmid, the gene of interest (GOI) is positioned between two ITRs. The AAV genes necessary for production, *rep* and *cap*, are supplied by a separate plasmid called the *trans* plasmid. Among the four non-structural Rep proteins encoded by the wildtype AAV (wtAAV) genome, only Rep78 and Rep52 are necessary for AAV vector production.[6] The capsid proteins VP1, VP2, and VP3 are required for proper capsid assembly. Additionally, assembly-activating protein (*AAP*) and membrane-associated accessory protein (*MAAP*) genes, which are embedded within the *cap* gene in different open reading frames (ORFs), aid in viral capsid assembly and egress.[13-15] To facilitate AAV vector production, a third plasmid is used to provide essential AdV genes. This "helper" plasmid expresses the *E2A*, *E4orf6*, and *VA RNA* genes.[16] AAV vectors produced using the tri-plasmid method have demonstrated safety, convenience, and effectiveness. However, this method has limited scalability,[17,18] and requires co-transfection of all three plasmids into a single cell.[19] Moreover, the requirement of high quantity of plasmid DNA (pDNA) and the need for ratio optimization among the three plasmids not only increases time and labor costs, but also introduces variability between production batches.[20-22]

In addition to packaging intact GOI genomes, AAV can also encapsulate nucleic acid

contaminants, including incomplete AAV vector genomes, producer plasmid genomes, and host cells

DNAs (hcDNAs).[23] The tri-plasmid systems yield a relatively high amount of impurities linked to the

input plasmids. These impurities can encompass sequences related to the bacterial origin of replication

and antibiotic resistance genes.[10,17,23-25] The level of DNA impurities encapsulated in AAV vector

products varies significantly, depending on the vector design and production process. Self-

complementary AAVs (scAAVs) have shown that as much as 26% of the total capsids can contain

backbone plasmid DNA.[26] For standard single-stranded AAVs (ssAAVs), this figure can be as high as

10%.[23,27-29] This concern has become more important, since recent trials have revealed significant

safety concerns at doses exceeding 1E14 vg/kg.[30,31] For instance, even if 95% of the packaged DNA

within a clinical preparation were fully functional vectors, a dose of 1E14 vg/kg would comprise around

5E12 vg/kg of non-therapeutic vector.[23] The use of an "oversized" backbone can significantly reduce

residual plasmid backbone packaging.[32,33] [34]

To address the increasing demand for AAV gene therapies, it is crucial to make improvements

to current production systems. One approach is to develop more efficient upstream processes,

including new cell lines and transfection reagents.[17,35,36] Another approach is to optimize the design and

function of the helper and *cis* plasmids used in production systems.[12,37,38] A third approach is to

streamline the process to optimize the transient-transfection step by reducing the number of plasmids

required. In a dual-plasmid system, one plasmid (pDG) carries the necessary AdV helper genes, as

well as the AAV *rep* and *cap* genes, while the other contains the GOI cassette.[37,39] Although this

approach has been proposed for over 20 years, it has not been widely adopted for AAV vector

production.[40] One possible reason for its limited use is the need for a large plasmid (~22 kb), posing

challenges during manufacturing, particularly in terms of scalability, purification, and stability.

Additionally, large plasmids exhibit lower transfection rates and high toxicities compared to smaller

plasmids.[41-43] Recently, another dual-plasmid system, pOXB, which combines the AAV vector genome

and *rep-cap* sequences together on a single plasmid, and provides AdV helper genes on a separate

plasmid.[44] A third possible dual-plasmid configuration (pLV) is to combine the AAV genome with the

AdV helper sequences, while expressing *rep* and *cap* on a separate plasmid. Interestingly, the pOXB

design outperforms the pDG-like and pLV dual-plasmid systems in terms of productivity.[44]

Here, we have developed an all-in-one single-plasmid AAV production system called AAVone. In the

AAVone system, the AdV helper genes (*E2A*, *E4orf6* and *VA RNA*), AAV helper genes (*rep* and *cap*),

and the AAV vector genome are assembled into one compact pAAVone plasmid with a 13-kb

backbone. High-quality AAV vectors can be consistently, efficiently, and easily produced by transfecting

packaging cells with relatively low amounts of a single plasmid. This makes the AAVone platform a

straightforward, cost-effective, and highly reliable system for AAV production, particularly well-suited for

generating GMP-grade AAV vectors.

RESULTS

Creation of AAVone single-plasmid system for AAV vector production

To create an efficient single-plasmid system for AAV vector production, it is critical to control the

total plasmid size, as it can modulate gene transfer efficiency.[41-43] Among the three plasmids for AAV

vector production, the AdV helper plasmid is the largest. Researchers have continually made efforts to

reduce the size of the AdV helper, such as pBHG10, pXX5, pXX6, and pAdDeltaF6.[12,45] The AdV helper

plasmid used in this study, pHelper, is 11.6 kb in size and contains 9.3 kb of the AdV2 genome. We

minimized the pHelper plasmid by removing the introns present in the *E2A* and *E4orf6* genes **(Figure**

S1). This modification led to a mini-pHelper that retains only 6.1 kb of the AdV2 genome with a total

plasmid size of 8.4 kb **(Figure 1A)**. When co-transfected with the *trans* and *cis* plasmids, the mini-

pHelper based system (mTri-plasmid) was capable of supporting AAV vector production, with higher

productivity compared with parental pHelper (pTri-plasmid) **(Figure 1B)**.[46] We then created two

versions of dual-plasmid systems that are based on the mini-pHelper: a pDG-like version and a

AAVdual (pLV-like) version.[44] In the pDG-like system, *rep-cap* was assembled into mini-pHelper,

resulting in a 13.4-kb plasmid, which is much smaller than the pDG plasmid.[39,40] In the AAVdual

version, the GOI cassette was integrated into the mini-pHelper, resulting in a construct with a 8.4-kb

backbone (pAAVdual) **(Figure 1A)**. The AAVdual system exhibited AAV vector yields that were

comparable to, or surpassed those of the pTri-plasmid system, whereas the pDG-like system displayed

comparable or reduced packaging efficiencies **(Figure 1B)**.

Encouraged by the positive outcomes from the dual-plasmid systems, we consolidated all of the

cis and *trans* essential elements for AAV production into a single-plasmid, pAAVone, with a 13-kb

compact backbone **(Figure 1A)**. In this system, the *rep-cap* expression cassette was positioned

between *E4orf6* and *VA RNA*, while the GOI cassette was situated between the *VA RNA* and the

bacterial origin of replication (*ori*). In this configuration, the *VA RNA* is closer to the left (L)-ITR and

bacterial elements are located after the right (R)-ITR. The compact pAAVone plasmids are easy to

clone and amplify without reducing yields (**Figure S2**).

With the AAVone system, AAV vectors can be produced by transfecting a single-plasmid into

producer cells. For example, AAV2-*CMV-Egfp* vectors were generated by transfecting pAAVone-AAV2-

CMV-Egfp plasmid into adherent HEK-293T cells. As shown in **Figure 1B**, the output from the tri- and

dual-plasmid systems varied greatly with changes in the proportions of the three plasmids, while the

AAVone system produces consistent and higher yields in multiple tests.

When the plasmids were tested by transient transfection **(Figure S3)**, the AAVone system

demonstrated comparable levels of EGFP expression (**Figure S3A,B**) and equivalent percentage of

positively transfected cells (**Figure S3C**) to those achieved by the pTri-plasmid system. It is important

to highlight that in these experiments, the pAAVone-AAV2-*CMV-Egfp* construct (15.8 kb) used in the

AAVone system and pAAV-*CMV-Egfp* construct (5.4 kb) used in pTri-plasmid system carried the same

AAV-*CMV-Egfp* genome design. In addition, the AAVone and the pTri-plasmid system (with molecular

ratio of pHelper:pRep-Cap:pAAV-*CMV-Egfp*=1:1:1) used the same amount of total pDNA masses.

Under this experimental parameter, the molecular ratio of pAAVone-AAV2-*CMV-Egfp* to pAAV-*CMV-*

Egfp plasmid is 1:0.64. When pAAVone-AAV2-*CMV-Egfp* and pAAV-*CMV-Egfp* were transfected at the

same mass (molecular ratio is 1:2.93), EGFP from the pAAV-*CMV-Egfp* plasmid was much stronger

than what was produced by transfection with pAAVone (**Figure S3**). This data suggests that the

increased AAV productivity of the AAVone system is not due to increased transfection efficiency.

The AAVone system achieves high AAV vector yields in suspension-cultured cells

To assess the compatibility of the AAVone platform for large-scale manufacturing, we initially

evaluated its packaging yields in suspension-cultured HEK-293T cells. AAVone consistently resulted in

a two- to four-fold increase in packaging yields compared to the pTri-plasmid system across various

serotypes **(Figure 1C and Table 1)**. We next tested various parameters, including different cell

densities (ranging from 2.5E6 to 3.5E6 cells/mL), PEI/DNA ratios (ranging from 2:1 to 3:1), and pDNA

amounts (ranging from 0.5 to 1.0 µg per 1E6 cells). The AAVone system demonstrated optimal

performance when 0.5 µg of pDNA was used for every 1E6 cells, with a PEI/DNA ratio of 2:1, and

transfecting at a cell density of 3E6 cells/mL **(Figure 1D)**. The most important parameter for vector

production was the pDNA amount. The highest yields were obtained for transfections using 0.5 µg

pDNA per 1E6 cells condition.

For high-yield AAV serotypes such as AAV1, AAV5, AAV8, AAV9, and AAVrh.10, the AAVone

system easily achieved crude lysate titers exceeding 1E15 vg/L in different production scales without

any optimization **(Table S1)**. Among these serotypes, AAV9 demonstrated the highest yields, with

crude titers ranging from 1.96E15 to 3.7E15 vg/L and purified titers ranging from 4.22E14 to 1.03E15

vg/L. AAV9 also exhibited the highest fold changes with AAVone, achieving up to a 4-fold increase in

yield compared to the pTri-plasmid system. Even with low-yield AAV serotypes, like AAV2, the AAVone

system reliably produced between 4.9E14 and 8.9E14 vg/L of crude lysate titers across various

production scales, showcasing its consistent performance. Remarkably, the AAVone system

demonstrated the ability to generate AAV vectors with transgene genome sizes ranging from 2.2 to 4.7

kb, while the sizes of the corresponding plasmids varied between 15.2 and 17.7 kb **(Table S1)**. Despite

a slight decrease in AAV vector yields with increasing transgene sizes, the AAVone system attained

1.94E15 vg/L of crude lysate titer for a 4.7-kb vector packaged with AAV9.

We further evaluated the productivity of the AAVone system in suspension-cultured HEK-293

cells, which are commonly used in GMP-level AAV production. In general, the yields achieved by the

AAVone system in the suspension HEK-293 cells were about half of those obtained with HEK-293T

cells (**Table S1**). The AAVone system showed adaptability with various transfection agents, yielding

high outputs with both PEImax and FectoVIR. Specifically, for AAV9 packaged with a 2.2-kb cassette,

AAVone reached up to 1.53E15 vg/L of crude lysate titers with PEImax. With AAV2, the system

produced over 4E14 vg/L with PEImax, and more than 6E14 vg/L with FectoVIR. These results

collectively indicate that the AAVone system is highly effective across various transgenes, AAV

serotypes and cell types, and shows compatibility with different transfection reagents. Moreover, the

AAVone system can be easily scaled for production using bioreactors **(data not shown)**.

AAV vectors generated by AAVone are similar to those produced by tri-plasmid systems

Assessing the quality of AAV vectors is of paramount importance when appraising a novel

production system. To directly compare the performance of the AAVone system with tri-plasmid

systems, AAV2-*CMV-Egfp* vectors were produced across all platforms using suspension-cultured HEK-

293T cells. A total of 0.75 µg/1E6 cells of pDNA was used for this examination. Vectors were isolated at

specific refractive index (RI) values, ranging from 1.369 to 1.371, following two rounds of CsCl gradient

ultracentrifugation. The composition of the capsid proteins, VP1, VP2, and VP3 showed an approximate

1:1:10 ratio across the three systems, with no observable differences between the vectors **(Figure 2A)**.

Purified vector DNA also showed that the predominant species were the expected 2.7-kb full-length

genomes (**Figure 2B**). All three preparations indicated the presence of truncated or partial genomes

with no visually discernable differences in the sizes and abundances of the partial genomes. It is

notable that the Tapestation software identified additional low-molecular weight species under 500 bp in

size for the pTri- and mTri-plasmid systems, but these species were not identified in the AAVone

system (**Figure 2B**). Mass photometry analyses also did not reveal major differences in particle masses

between vectors produced by the three platforms (**Figure 2C**). We next evaluated whether there were

differences in the transduction profiles between vectors produced by the three systems. Transduction of HeLa cells indicated no clear differences in EGFP expression at 48-hrs post-transduction (**Figure 2D**).

Vectors packaged by the AAVone system and tri-plasmid systems exhibited similar retinal transduction profiles.

Retinal tissues are among the most prominent targets for AAV vector-based gene therapies being tested in preclinical and clinical trials; especially for platforms that are based on AAV2. Vectors packaged with AAV2-*CMV-Egfp* using the AAVone, mTri-, and pTri-plasmid systems were administered to mouse retinas by intravitreal (IVT) injection at 1E9 vg/eye. At four-weeks post-injection, we conducted funduscopy imaging to assess EGFP expression and distribution across the retina. We found that administration of the three vectors led to similar levels of EGFP expression and distribution (**Figure 3A**). At six-weeks post-injection, retinas were collected, cryo-sectioned, and stained with anti-EGFP antibody (green), peanut agglutin (PNA, cone photoreceptor marker, white), and DAPI (blue). We observed comparable levels of EGFP and PNA signal overlap across all three vector groups (**Figure 3B**) and similar percentages of EGFP-positive photoreceptor outer segments per section (56-59%) (**Figure 3C**). This finding suggested that vectors produced by the AAVone system performed similarly to those produced by the tri-plasmid systems. To quantify the transduction of the vectors produced by the three systems, we extracted bulk DNA and RNA from the harvested retinas to measure the presence of vector genomes and transgene transcripts. Using digital droplet PCR (ddPCR) we found that the vector packaging systems did not influence transduction (**Figure 3D,E**).

We next aimed to assess whether vectors produced by the three AAV packaging systems elicit differences in immune responses. It has been well documented that following IVT injection of AAV vectors, microglia are activated and migrate to regions of pathogen-associated molecular pattern (PAMP) activation.[47-50] We therefore stained cryosections of treated retinas to mark microglia (anti-IBA1). Cells positive for IBA1 were found to increase in the retinas treated by AAV vectors (**Figure 3F**). Notably, we found similar counts of Iba1-positive cells per section across all three treatment groups.

We also examined the distribution of Iba1-positive cells infiltrating into the deeper layers of the

transduced retinas (**Figure 3G,H**). All three vectors exhibited equivalent stimulation of IBA1-positive cell

infiltrates across the retinal across the outer plexiform layer (OPL), the inner nuclear layer (INL), the

inner plexiform layer (IPL), and the ganglion cell layer (GCL).

Vectors packaged AAV1, AAV8, AAVrh.10, and AAV-PHP.eB using the AAVone system show

similar *in vivo* transduction profiles as vectors produced with tri-plasmid systems

We next aimed to determine whether the AAVone system was also compatible with other

serotypes. We packaged the same ssAAV-*CMV-Egfp* transgene as above with AAV1, AAV8, AAVrh.10,

and AAV-PHP.eB capsids by the AAVone, pTri-, and mTri-plasmid systems. Since AAV1 and AAV8

are known to be tropic to muscle, we injected the AAV1- and AAV8-packaged vectors into the tibialis

anterior (TA) muscle of adult female mice via intramuscular (IM) injection. Animals were sacrificed at

week 2, for those injected with AAV1 vectors; and at week 8, for those injected with AAV8 vectors. TA

muscles were harvested and cryosectioned to assess differences in transduction efficiency between the

three vectors systems. Fluorescence microscopy of TA cross-sections revealed similar intensities and

distributions of EGFP throughout the TA across the three treatment groups for AAV1 and AAV8 (**Figure**

S4A) vectors. Additionally, we assessed the immunogenic profiles of vectors produced by the AAVone

system compared to vectors produced by pTri- or mTri-plasmid systems. Two-weeks post-injection,

there were no observable differences in cellular infiltrates of the TA muscle across the treatment groups

(**Figure S4B**).

Both AAV-PHP.eB and AAVrh.10 are well known for their neurotropic properties. We therefore

injected adult male C57BL/6J mice at a dose of 1E14 vg/kg by intravenous administration. Two-weeks

post-injection, animals were euthanized and brains were harvested. Immunohistochemical staining of

brains cryosections showed that AAV-PHP.eB and AAVrh.10 exhibited different patterns of EGFP-

positive cells, as expected (**Figure S5**); AAV-PHP.eB primarily transduced neurons, while AAVrh.10

transduced both neurons and glial cells. Importantly, there were no significant differences in EGFP

transduction between vectors produced by the three packaging systems among the AAVrh.10 and

AAV-PHP.eB groups.

The AAVone system produces vectors with slightly reduced genome heterogeneity

We next assessed the heterogeneity of the vector genomes using single molecule, real-time

(SMRT) sequencing and AAV genome population (AAV-GPseq) analyses.[11,28,51] Using this approach,

we were able to analyze individual AAV vector genomes at a single molecule level. The majority of the

reads obtained by SMRT sequencing spanned the entirety of the ssAAV-*CMV-Egfp* reference genome

(**Figure S6**). Truncated genomes with snap-back configurations[52,53] were also observed as reads that

only partially spanned the reference from ITR to ITR. These species appear to be similar across the

three production platforms, suggesting that there were no substantial differences in truncation events

between the vectors analyzed. Instead, the major truncations appear to be centered at the woodchuck

hepatitis virus post-transcriptional regulatory element (*WPRE*) (**Figure S5**). We next aimed to assess

whether the AAVone system produced vectors with different amounts of non-unit length genomes. The

lengths of the reads mapping to the references were tabulated and graphed (**Figure 4A**). The vector

genomes from the two tri-plasmid systems had nearly identical read-length distributions, with a major

peak at 2,770 nt and multiple smaller peaks at 347 nt, 591 nt, and 1,878 nt. The 2,770-nt population

represents the intact AAV genome size, plus two intact 145 nt ITRs, suggesting at ITR repair.[54]

Interestingly, the vector produced by the AAVone system displayed a predominant read length peak at

2,770 nt, with some minor peaks at other lengths. The AAVone system exhibited a slightly higher

percentage of intact genomes (58.77%), compared to the pTri-plasmid (51.34%) and the mTri-plasmid

(51.41%) systems (**Figure 4A**). Analyses of read alignment start and end positions also indicate a clear

reduction in truncation hotspots with the AAVone system when compared to the two tri-plasmid

systems (**Figure 4B**). As noted before, the major truncation hotspot appears to be centered at the

WPRE. Additional truncations are revealed to be centered at the *hGH* polyA sequences (**Figure 4B**).

Intact ITRs in AAV vectors are fundamental for transgene expression, and for the overall

success of the gene therapy. wtAAV2 ITRs are 145-nt in length and can take on two configurations, flip

and flop [55]. Packaged vectors that undergo proper rolling-hairpin replication show equal distribution of

flip and flip at both ITRs. The AAV GOI cassette used by the different production systems are the same.

Within the plasmid construct, the R-ITR is 130 nt long, which is 15 nt shorter than the full-length 145-nt

ITR. The L-ITR is 119 nt in length and differs from the L-ITR by an 11-nt deletion in the C arm **(Figure**

4C). Both ITRs were designed in the flop orientation. We found that the ITRs in all AAV vectors were

145 bp in length at both L- and R-ITRs, **(Figure 4D)**. The majority of reads obtained from vectors

produced by the two systems were fully aligned to the reference ITRs. Flip and flop are the two major

configurations observed in both L- and R-ITRs in AAVone and tri-plasmid systems. They occupy over

90% of the population and display similar flip:flop ratios **(Figure S7)**. Other mutated and truncated

forms were also observed in both L- and R-ITRs but they were in the minority. The presence of ITRs

missing the 11 nt in the C arm constituted only ~1.5% of the total genomes, implying that the deletion

was largely repaired during replication (**Figure 4E**).[56] Taken together, AAV vectors produced by the

AAVone system have nearly identical genomic compositions as compared to those produced by the tri-

plasmid systems.

The AAVone system exhibits alternative plasmid-related impurities in the AAV vectors

Residual input pDNAs are one major concern for AAV production. In the tri-plasmid systems,

contaminants mainly originate from the backbone of the *cis* plasmid, while *trans* plasmid sequences are

packaged at lower levels.[29] One major difference between the AAVone and the tri-plasmid systems is

the oversized 13-kb backbone, which not only includes the bacterial *ori* and the antibiotic selection

gene (*kanR*), but also contains *rep*, *cap*, *E2A*, *E4orf6* and *VA RNA*. In order to examine the impurities

originating from plasmids found in vectors produced by AAVone and tri-plasmid systems, we aligned

SMRT reads to the AAV-GOI plasmid references. We found that the two ITRs were where the majority

of the reads mapped **(Figure 5A-C)**. We also showed that the reads mapping to the plasmid backbone

tend aggregate proximal to the ITRs, with decreasing coverage when sequences are further away from

the ITRs. Within the AAVone system, *VA RNA* constituted 2.35% of the total sequences **(Table S2)**,

making it the most prevalent contaminant. The next most abundant type of contaminant was the

plasmid bacterial backbone (0.87%). Other elements, such as *E2*, *E4*, *rep,* and *cap*, which are more

distant from the ITRs, exhibit considerably lower percentage of mapped reads **(Table S2)**. The

dominant contaminant in vectors produced by the tri-plasmid systems originated from the *cis* plasmid

backbone (3.95% in the pTri-plasmid and 2.69% in the mTri-plasmid system). These observations were

validated through qPCR analysis focusing on various regions of the pAAVone plasmid **(Figure 5D)**.

Thus, the dominant contaminants in AAVone-produced vectors were different from the contaminants

related to vectors produced by the tri-plasmid systems.

hcDNA contaminants originating from the packaging cell line used in vector production

represent another form of genomic contaminant that may impact the safety of gene therapy vectors.

Overall, there was no notable difference in hcDNAs among the AAV vectors produced by the three

production systems, with each showing approximately 3% of the total reads **(Table S2, Figure S8)**. In

all of the reads that mapped to the hg38 genome, the average (~1,300 nt) and mean (~1,140 nt) read

lengths were very similar among the three systems **(Figure S9)**. These read lengths are shorter than

the smallest oncogene size (1,664 bp).[57] The number of unique reads spread across each

chromosome, suggested at an overall random distribution among all three AAV samples **(Figure**

S8).[27,28] Interestingly, reads mapping to mitochondrial DNA (chrM) appeared to be enriched at the D-

loop[27] **(Figure S10).** However, chrM contaminants that were chimeric with the GOI were also rare (data

not shown). Among all of the hcDNA-mapped reads, irrespective of the production system, only 15%

appear to be chimeric genomes, hcDNAs recombined with vector genomic sequences **(Table S2,**

Figure S8). However, when these chimeric reads were mapped back to the GOI reference, and

visualized, the majority of reads mapped imperfectly with the hGH polyA **(Figure S11)**. These results

indicated that the chimeric reads were predominantly false positives.

In summary, our SMRT sequencing results demonstrate that the AAVone system exhibits

significantly lower levels of bacterial backbone sequences, comparable or slightly higher *rep, cap*, *E2*,

E4, and hcDNA contaminations, and a increase in *VA RNA* sequences compared with the tri-plasmid

systems. Despite this shift in the nature of contaminants, the overall total number of genomic

contaminations for the AAVone system were decreased (**Table S2**).

The AAVone system does not pose a risk for generating rcAAV contaminants.

Another concern for the AAVone system is the potential for generating replication-competent

AAV (rcAAV). The FDA requires that AAV products be screened for the presence of possible rcAAVs.

Current vector production systems in use have been optimized to reduce rcAAV to levels ≤1 rcAAV per

1E8 vector genomes.[24] In tri-plasmid systems, rcAAVs arise through the recombination of DNAs

carrying ITRs with *rep* and *cap* sequences originating from the pRep-Cap plasmid.[58] In the AAVone

system, *rep*, *cap,* and the ITRs coexist within the same plasmid. A cell-based rcAAV assay performed

by an external contract research organization indicated that rcAAV levels were below one per 1E9

vector genomes for AAV vectors generated by the AAVone system. After three rounds of amplification

with AdV, all three flasks in the 1E9 vector genome group showed no presence of rcAAV, and only one

out of three samples tested positive for rcAAV in 1E10 vector genomes. This underscores the AAVone

system can effectively produce safe AAV vectors with low rcAAV. Using qPCR primers targeting both

ITR and *rep* to detect ITR-*rep* hybrids.[59] which represent a potential for rcAAV contamination,

comparable levels were observed among the AAVone and tri-plasmid systems (**Figure 5D**). Analyses

of the SMRT reads representing genomes obtained from vectors produced by the AAVone system

showed that the majority of reads mapping to the *rep* or *cap* genes also co-mapped with the GOI

sequence (89 out of 99 (89.9%) and 201 out of 230 (87.39%), respectively). It is noteworthy that in the

tri-plasmid systems, reads that map to either *rep* or *cap* from the *trans* plasmid, which inherently lack

ITRs, also displayed a high degree of mapping to ITRs **(Table S2)**. For a read to represent an rcAAV

genome, it has to contain the ITRs, *rep*, and *cap* in one sequence.[58] Our analysis revealed that among

the vector genomes from the AAVone system, such chimeric reads were present in 90 out of 196,753

reads (0.045%) **(Figure 6)**, which was slightly higher than those of the pTri-plasmid system (27 out of

131,989; 0.021%) and mTri-plasmid system (22 out of 119,029; 0.018%). Nevertheless, among all of

these tri-chimeric sequences, none contained an intact ITR-*rep-cap*-ITR configuration **(Figure 6)**.

The AAVone system enhances production and full-particle ratios while reducing impurities

using less plasmid DNA

In the tri-plasmid systems, a relatively high amount of input plasmids (1 µg/1E6 cells) is typically

needed to achieve high AAV vector yields.[60-63] Surprisingly, the AAVone system allows for a substantial

reduction of input plasmid, while still increasing AAV yield in both HEK-293T and HEK-293 cells

(Figure 7A and 7B). In HEK-293T cells, using plasmids of 15.2 kb or 15.8 kb carrying *CMV-Egfp*

genomes, the AAVone system reached its highest yield at 0.25 µg/1E6 cells, a result that was

independent of the serotype **(Figure S12)**. In HEK-293 cells, the optimal plasmid amounts were

between 0.25 to 0.50 µg/1E6 cells **(Figure 7A)**. The reduced input plasmid also increased transfection

efficiency for AAV2 vectors (**Figure S12A**). The use of reduced plasmids slightly increased cell viability

for both AAV2 and AAV9 vectors **(Figure S12B),** while cell densities reached up to 6E6 cells/mL

(**Figure S12C**). Overall, the optimal plasmid dose required for the AAVone system is approximately one

half to one fourth of the amount typically utilized in tri-plasmid systems.

The use of reduced input plasmid with the AAVone system has additional benefits. It led to

increased full-to-empty capsid ratios in the whole production system (**Figure 7C-E**), which are desirable

for obtaining higher-quality AAV vectors with a larger proportion of functional, transgene-carrying

capsids. When utilizing the same quantity of input pDNAs to produce AAV2 vectors (0.75 µg/1E6 cells),

the AAVone system exhibited a similar full-to-empty ratio as compared to what was achieved by the tri-

plasmid systems (**Figure 7C**). All systems demonstrate two prominent peaks at 66 S and 92 S, which

correspond to empty AAV2 particles and full capsids with 2.8-kb genomes, respectively. Approximately

30% of AAV particles found in crude lysates for each production system were full capsids. When the

plasmid was reduced to 0.25 µg/1E6 cells with the AAVone system, the full particle ratio rose from

30.02% to 38.48%. Similarly, titers for AAV8 and AAV9 exhibited a ~10% increase in full AAV particles

when using a reduced amount of input pDNA (**Figure 7D,E**).

The use of reduced pDNA with the AAVone system can also help minimize plasmid-related

impurities. Reducing the pDNA amount led to a decrease in plasmid impurities on both sides of the

ITRs **(Figure 7F)**. In summary, reducing the input plasmid increases packaging efficiency, leads to

increased full-to-empty particle ratios, and reduced plasmid-related impurities.

DISCUSSION

Advantages of the AAVone system

The AAVone system described here is designed to bring all the essential components for AAV

production into a single-plasmid (**Figure 1**), while retaining all of the essential vector performance

features and qualities expected of those produced by standard tri-plasmid transfection platforms

(**Figures 2 and 4**). These features include the preservation of important characteristics such as capsid

content, genome and ITR integrity, transduction efficiency, ratio of full-to-empty particles. With this

system, users can integrate their existing GOIs into the AAVone manufacturing process without the

need to develop additional upstream and downstream processes, analytical release, or characterization

assays.

A notable benefit of using the AAVone system is the considerable reduction in costs associated

with raw materials, as it only uses a small quantity of a single-plasmid, instead of three plasmids.

Plasmids represent one of the most expensive components in AAV vector production. The large-scale

production of three GMP grade plasmids with over 95% purity and absent of process-related impurities,

continues to be a significant challenge. Although numerous researchers have dedicated efforts to

developing dual-plasmid transfection systems for AAV vector production,[37,39,40,44] this study stands out

as the first to investigate the production of AAV vectors using a single-plasmid. The system enables the

production of AAV vectors using only about one-quarter to one-half of the total pDNA normally required

with conventional approaches **(Figure 7)**. Lowering the input of pDNA not only reduces costs, but also

leads to decreased genomic contaminants, increased full-to-empty capsid ratios, and increased titers

(**Figure 7**). Combined with a two- to four-fold increase in AAV vector productivity, the AAVone system

uses only one-tenth of the total pDNA to achieve the same titers produced from the pTri-plasmid

system. The AAVone system's single-plasmid approach also addresses the challenges faced with

traditional tri-plasmid systems, such as obtaining a consistent vector product **(Figure 1B** and **Table**

S1). Finding the optimal ratio of the three plasmids, maintaining plasmid stability, and ensuring

consistent copy numbers across different production batches is complex and often times challenging

when considering the different cell lines, AAV serotype, and the size and composition of the rAAV

cassette used.[60-63] The single-plasmid system minimizes the potential for introducing plasmid-related

variation; thereby, enhancing the consistency of the AAV vectors produced.

The AAVone system potentially offers a higher effective AAV vector production efficiency

compared to the pTri-plasmid system. The increased productivity of AAVone system is not due to

increased transfection efficiency, but increased production efficiency **(Figure S3)**. Theoretically, within

the AAVone system, there is a linear relationship between AAV vector production efficiency and

transfection efficiency. However, in the tri-plasmid systems, transfection efficiency cannot linearly

convert to AAV vector productivity, as only cells that are co-transfected with all three plasmids are able

to produce AAV vectors **(Figure S13)**. This effect becomes significantly pronounced at lower

transfection efficiencies. It has been reported that at a transfection efficiency of ~60%, only ~7% of

cells produce measurable amounts of assembled AAV capsid with the tri-plasmid transfection method,

while transfection with the dual-plasmid system increased cells positive for assembled AAV capsid by

four- to five-fold.[19] Moreover, it has been reported that with three different dual-plasmid configurations,

pOXB showed significantly superior results compared to the pDG-like and pLV systems.[44] We also

found that the pLV configuration exhibited comparable productivity, and the pDG-like configuration

showed a lower production efficiency than achieved with the tri-plasmid method (**Figure 1B**). These

findings suggest that co-existence of the AAV vector genome and AAV helper genes may play an

important role in the increased vector productivity by AAVone.

Genomic impurities by the AAVone system

Besides intact GOI genomes, AAV vector production also yields contaminants in the final

product, including incomplete AAV vector genomes, and DNAs from the producer plasmids and host

genome.[23] The presence of contaminants in clinical vectors is considered undesirable because it can

lead to increased immune responses against the vector, potential integration into the host genome, and

other safety concerns.

Different types of packaged incomplete vector genome species are present in AAV vectors,[52]

but most of them are snap-back forms (**Figure S4**), which are double-stranded DNAs containing either

5′ or 3′ ITRs.[23] The proportion of incomplete genomes are sequence dependent, and regions with high

secondary structure such as short hairpin RNA (shRNA) result in a greater proportion of snap-back

genomes.[28,52] In our AAV-*CMV-Egfp* constructs, the major truncation hotspot appears to be centered at

the *WPRE*, *hGH* polyA, and *CMV* sequences, which are close to ITRs (**Figure 4A,B** and **Figure S6**). In

our previous studies, byproduct empty rAAV particles were not truly empty, but contained different

types of short vector genomes that were enriched for sequences corresponding to the ITRs.[64] We were

unable to completely remove these aberrant genomes from rAAV preparations, even after extracting

thin bands related to full capsid fractions following CsCl ultracentrifugation (**Figure S2** and **Figure**

4A,B). However, the larger size of the pAAVone plasmid may contribute to the reduction of aberrant

genomes, as the AAVone system yielded slightly more intact genomes and significantly less the snap-

back genomes in the final purified AAV vectors when compared to what is produced by tri-plasmid

systems (**Figure 4** and **Table S2**). Furthermore, decreasing the amount of the pAAVone plasmid during

production significantly reduced the impurities as well as the empty particles (**Figures 5 and 8**). Both

the DNA template switch model and NHEJ mechanism may result in the formation of snap-back

genomes.[52] The use of one large plasmid may overall reduce recombination events between ITR and

other plasmid elements, which in turn leads to decreases in genomic impurities. Previous studies have

concluded that increasing the size of the plasmid backbone with stuffer DNA can reduce the level of

plasmid backbone contaminants, as the two ITRs are not in range of each other to facilitate reverse

packaging. The pAAVone plasmid has 13 kb of non-GOI-cassette sequences existing beyond the two

ITRs. The oversized design did decrease the abundance of sequencing spanning the bacterial *ori* and

the antibiotic resistance gene. All of the high abundance elements were positioned close to ITRs on the

plasmid, which supports the idea that the majority of vector product-related impurities are ITR-

containing fragments. In addition, the abundance of contaminants is mainly determined by the distance

from the ITRs (**Figures 5 and 8D**) and sequences positioned more than 2.5 kb away from the ITRs,

such as *rep*, *E2*, and *E4* genes, are rarely packaged into AAV vectors (**Figure 5).** Thus, incorporating

genes into one single-plasmid did not significantly increase their related impurities. As the *rep* is

located 5.6 kb away from ITR, the pAAVone design did not increase the potential for rcAAV formation.

Overall, we found that an oversized backbone design reduced total plasmid-related impurities (**Table

S2**).

In conclusion, the AAVone packaging system represents a simple, cost-effective, and highly

reproducible AAV production system, particularly suitable for GMP-grade AAV production.

MATERIALS AND METHODS

AAV plasmids

All vector constructs were generated using HiFi assembly (NEB), Gibson assembly, or T4

ligation and were scaled up with midi/maxi preparation kits (Takara Bio). The mini-pHelper was

generated by deleting introns of *E2* and *E4* genes from the pHelper parent construct. pAAVdual

plasmids were cloned by inserting AAV-GOI genomes (including ITRs) into mini-pHelper at the *Pme*I

site, located between pMB1 *ori* and *VA RNA*. mini-pHelper-Rep-Cap plasmids were cloned by inserting

the *rep-cap* cassette into the *Cla*I site of mini-pHelper, which is located between *E4* and *VA RNA*.

pAAVone-AAV2-*CMV-Egfp* plasmids were achieved by insertion AAV2-*CMV-Egfp* genome into the

mini-pHelper-Rep-Cap-AAV2 at the *Pme*I site. pAAVone plasmids with different capsids and

transgenes based one pAAVone-AAV2-*CMV-Egfp*. All the ITR-bearing plasmids were subjected to

restriction enzyme analysis using *Sma*I and/or *Ahe*I, ITR sequencing and/or whole plasmid sequencing

to confirm the ITR structures.

AAV production, purification, and titration.

Adherent HEK-293T cells were cultured in DMEM medium with 10% fetal bovine serum (FBS)

and 1% penicillin/streptomycin (P/S) in various culture vessels including 24-well plates, 15 cm dishes,

or 2 L roller bottles. Suspension HEK-293T and HEK-293 cells were expanded for at least three

passages and inoculated into a shake flask with F17 culture medium (Thermo Fisher, Waltham, MA)

prior to transfection. The transfections were carried out using either PEImax (Polyscience, Warrington,

PA) or FectoVIR (Polyplus, NY), adhering strictly to the provided guidelines. Experimental variations

included different cell densities, amounts of pDNA, and ratios of transfection reagent to DNA. Cells

were collected for analyses 72 hours after transfection. For initial titer estimations, cell samples

underwent a single freeze-thaw cycle and sonication for lysis. Subsequent centrifugation allowed for the

collection of supernatants, from which crude titer productivity was assessed. The AAV vectors extracted

from these samples were then purified through one or two CsCl-gradient ultracentrifugation steps

before undergoing quality evaluation. Prior to titer determination, both crude lysate and purified

samples were treated with DNase I to remove un-encapsidated residual DNA, followed by Proteinase K

treatment. The quantification of AAV vector productivity was achieved using either Droplet Digital PCR

(ddPCR) or qPCR, employing primers and/or probes targeting the vector's transgene. To establish a

standard curve, a plasmid carrying the relevant sequence was linearized. Each experimental batch

included a well-characterized AAV vector as an internal standard, ensuring data normalization against

this control.

Mice

Six- to eight-week-old female C57BL/6J mice were administered AAV2-*CMV-Egfp* or saline IVT

injection to the left and right eyes at 1E9 GC/dose (n=10 per group). All AAV2 vectors were diluted with

saline and fast green to 1.0E9 vg/µL. The IVT injections were performed with glass needles (Clunbury

Scientific LLC; Cat no. B100-58-50) to deliver ~1 µL of fluid into the vitreous using the FemtoJet from

Eppendorf with a constant pressure of 300 psi and injection time of 1.5 s. At four-weeks post-injection,

funduscopy imaging was performed to assess the EGFP expression in the mouse retina. Mice were

euthanized at four- and six-weeks post-injection. Eye cups were collected for immune-staining and

molecular quantification.

Six- to eight-week-old female C57BL/6 mice were administered AAV1-*CMV-Egfp*, AAV8-*CMV-*

Egfp, or saline via intramuscular (IM) injection to the left and right tibialis anterior (TA) muscle. Vectors

were diluted with saline to 1E10 vg/µL and administered at 1E11 GC/dose (n=10 per group). At two-

and 6-weeks post-injection, TA muscles were harvested and either placed in 4% paraformaldehyde

(PFA, for histology) or snap frozen (for RNA and DNA extraction). Following 4% PFA overnight fixation,

tissues were sent to UMass Chan Medical School Morphology Core for H&E staining or placed in 30%

sucrose for 72 hours before embedding in Tissue-Tek® O.C.T. Compound (Sakura Finetek USA Inc

#4583). The embedded tissues were cryosectioned (10 µM slices) and mounted on slides with

VECTASHIELD® HardSet™ Antifade Mounting Medium, with DAPI (Vector Laboratories #H-1500-10)

for microscopy. All images were visualized with a Leica DM6 Thunder microscope with a 16-bit

monochrome camera. Images were processed by LAS X Life Science microscope software.

AAV vectors were intravenously administered to adult male C57BL/6J mice (8–10 weeks old,

n=3) at a dose of 1E14 vg/kg per mouse. Two-weeks post-injection, animals were euthanized and

intracardially perfused with cold 0.1 M PBS followed by 4% PFA in 0.1 M PBS. The brains were

harvested, fixed overnight in 4% PFA, and cryoprotected in 30% sucrose at 4 °C for 2-3 days. Eight

series of 40-µm-thick sections were cut using a frozen sliding microtome, collected in sterile Eppendorf

tubes containing 30% sucrose in 0.1M PBS, and stored at -20°C until use.

Immunohistochemistry

Treated retinas were cross-sectioned as previously described [48]. In brief, eye cups were

dissected in cold 1X PBS and fixed in 4% PFA overnight at 4 °C. Cryosections were cut to 12-micron

thickness. The following primary antibodies and dilutions were used: chicken anti-EGFP antibody

(1:1000; Abcam; Cat no. ab13970), and rabbit anti-IBA1 (1:300; Wako; Cat no. 019-19741). The

chromophore conjugate, fluorescein peanut agglutinin (PNA) (1:1,000; Vector Laboratories; Cat no.

FL1071) was used to stain cone photoreceptors. All antibodies were diluted in PBS with 0.3% Triton X-

100 and 5% bovine serum albumin (CST). Nuclei were counterstained with 4′,6-diamidino-2-

phenylindole (1:2000, Sigma-Aldrich; Cat no. 9542). The following secondary antibodies were used:

Donkey anti-Chicken IgY (H+L) Highly Cross Adsorbed Secondary Antibody, Alexa Fluor™ 488,

A78948 and Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™

594, A-11037. All secondary antibodies were diluted 1:1000 in PBST (PBS with 0.25% tween-20). All

images were visualized with a Leica DM6 Thunder microscope with a 16-bit monochrome camera.

Images were processed by LAS X Life Science Microscope Software.

Floating brain sections were washed three times with 0.5% Triton X-100/Tris-buffered saline

(TBS) and blocked for 30 minutes in a blocking buffer containing 2% BSA in TBS with 0.5% Triton X-

100. Primary anti-GFP antibodies were then added, and the sections were incubated overnight at 4 °C.

After three washes, the sections were treated with Alexa fluorescent-labeled secondary antibodies in

blocking buffer for one hour at room temperature. Following washing and Hoechst counterstaining,

brain sections were mounted onto glass microscope slides and cover slipped with Fluoroshield (Sigma

Aldrich Cat. #F6182). Tiled confocal images of entire brain sections were acquired and analyzed using

the Leica Stellaris-5 confocal microscope system.

RNA and DNA extractions

Snap-frozen TA muscles were homogenized using Bullet Blender Green Bead Lysis Kit

(NextAdvance #GREENE5) in conjunction with a Qiagen TissueLyzer II. Half of the homogenized

tissues were processed for RNA extraction using mirVana™ miRNA Isolation Kit (ThermoScientific

#AM1561) according to the standard protocol. The remaining homogenized tissues were processed

using QIAamp DNA Mini Kit (Qiagen #51306) according to manufacturer's recommended procedures.

Extracted RNAs were reverse transcribed into cDNA using Iscript™ Reverse Transcription Supermix

(BioRad #1708841) and Iscript™ gDNA Clear cDNA Synthesis Kit (BioRad #1725035).

Snap-frozen retinas were homogenized using Qiagen 5mm TissueLyser bead (Qiagen #69965)

in conjunction with Omni Bead Ruptor Elite. Homogenized retina tissue was processed for RNA and

DNA extraction using the Norgen Biotek RNA/DNA/Protein Purification Plus Kit (Norgen Biotek #47700)

according to the standard protocol. Extracted RNA was reverse transcribed into cDNA using Iscript™

Reverse Transcription Supermix (BioRad #1708841).

Digital Droplet PCR

Digital Droplet PCR (ddPCR) was performed on gDNA and cDNA using ddPCR Supermix for

Probes (No dUTP), 2X (BioRad #1863024) with the following probe sets: *Egfp*-FAM (Thermofisher

#4400291, Assay ID Mr00660654_cn), mouse *Gapdh*-VIC (Thermofisher #4448484, Assay ID

Mm99999915_g1), and TaqMan™ Copy Number Reference Assay, mouse, Tfrc-VIC (ThermoFisher

#4458366). Input for AL 479, AL 480, and AL 481 ddPCR reactions were as follows: 40 ng gDNA per

20 uL reaction and 0.5 uL cDNA (diluted 1:15) per 20 uL reaction. Droplet generation was performed

using a BioRad Automated Droplet Generator using the corresponding manufacturer recommended

consumables into a 96-well plate. PCR reactions were conducted on a thermocycler using the following

conditions: 95 °C for 10 minutes, 32 cycles of 95 °C for 30 seconds and 61 °C for 1 minute, and a final

step of 95 °C for 10 minutes. Droplets were read using a BioRad QX200 Droplet Reader and analyzed

using QuantSoft software. Raw reads were exported and *Egfp* values were normalized either to diploid

genomes or to housekeeping gene transcripts.

AUC analysis

AUC was used to quantify empty and full particles. AAV from CsCl index 1.360-1.380 were collected after one round of ultracentrifugation. AUC measurements were performed at 20,000 rpm and 20 °C on a Beckman AUC Optima instrument (Beckman Coulter, Brea, CA, USA) equipped with an An-50 Ti analytical rotor (Beckman Coulter, Brea, CA, USA) using a total number of 150 scans per sample. Analysis of AAV samples by the SEDFIT c(S) model yields a distribution of sedimentation coefficients with each peak in the distribution representing AAV particles with full genomes or empty capsids. Integration of individual peaks yields the sedimentation coefficient (S) and the relative concentration of each species in the distribution.

Mass photometry

Mass photometry measurements were carried out on a SamuxMP instrument (Refeyn Ltd., Oxford, UK). Prior to each measurement, a calibration was conducted using truly-empty AAV particles, which was produced by transfection of one mini-pHelper-Rep-Cap plasmid in HEK-293T cells.

qPCR titration plasmid related impurities contaminant amplicons

Purified AAV vectors were treated by DNase I and proteinase K. Vector genome titers and contaminant amplicon titers were assessed by qPCR using SYBR Green. Viral titers were quantified on a StepOnePlus Real-Time PCR System (Applied Biosystems) using the default settings. The procedure involved denaturation at 95 °C for 5 min, followed by 40 cycles of amplification (95 °C for 10 s, 60 °C for 30 s), and concluding with a melting curve. Primers targeting different regions of the pAAVone plasmid are shown in **Table S1**. A plasmid containing the corresponding target sequences was used to make the standard curve. All data were normalized to *Egfp* transgene ct values.

Characterization of the AAV capsid

The capsid compositions of AAV vectors were determined by Bio-Safe Coomassie G-250 stain.

AAV vectors (~5E11 particles) were resolved by electrophoresis on a 10% SDS-PAGE gel and

standard Coomassie G-250 stain was performed according to the manufacturer's procedures (Bio-

Rad).

AAV vector transduction in vitro*.*

The HeLa cell line used in this study were grown in DMEM with 10% FBS and 1% P/S at 37 °C

in a humidified environment supplied with 5% CO_2. For each transduction experiment, ~50,000 viable

cells were seeded onto a 24-well plate 24 h before transduction. AAV vectors were added directly to

each well at an MOI=10,000. EGFP expression was determined 48 h post-transduction by

epifluorescence microscopy.

Flow cytometry

Cells transfected with the tri-plasmid systems or the AAVone system was collected at

approximately 48 h after transfection and processed using a BD Accuri C6 Plus flow cytometer

equipped with an autosampler (BD Biosciences, San Jose, CA). At least 50,000 events were collected

per sample with the fluidics rate set to slow (14 µL/min) with one agitation/SIP clean cycle between

samples. The acquired data were next analyzed with FlowJo software (v10.8; FlowJo, Ashland, OR)

using standard gating strategies to select for single cells (based on SSC-A vs. FSC-A and FSC-H vs.

FSC-A scatterplots) and to calculate the percentage of GFP-expressing cells (based on FITC-A

histogram).

NGS analysis of AAV vector genomes

NGS of AAV vector genomes was performed by Azenta. The ssAAV viral DNA was extracted

using the Invitrogen PureLink Viral RNA/DNA kit following manufacturer's instructions. DNA samples

were quantified using a Qubit 2.0 Fluorometer (Life Technologies, Carlsbad, CA, USA) and sample

integrity was checked using Agilent TapeStation 4200 (Agilent Technologies, Palo Alto, CA, USA).

Amplicon libraries were prepared using SMRTbell Express Template Prep Kit 3.0. The sequencing

library was validated on the Agilent TapeStation and quantified by using Qubit 2.0 Fluorometer. All

libraries were combined into a single sample and sequenced across two PacBio Sequel IIe SMRT cells

using Sequel II binding kit 3.0 with heteroduplex detection mode on.

SMRT sequencing and AAV-GPseq

The bioinformatic workflows were conducted using the Galaxy platform.[65,66] Resulting

consensus fastq files were mapped to the references as reported in the study using the Burrows–

Wheeler aligner-maximal exact match (BWA-MEM)[67,68] in PacBio mode (-x pacbio). Aligned reads were

visualized with the Integrated Genomics Viewer (IGV) tool version 2.14.0 [69] with soft clipping on.

Statistical Analyses

The results were analyzed by one-way or two-way ANOVA statistical tests were performed in

GraphPad Prism 10 along with Tukey's multiple comparison statistical tests where applicable as

described.

DATA AVAILABILITY STATEMENT

All data are provided in the main text, figures, or the supplemental information, and raw

data files are available upon request. Requests for reagents or cell lines used in this study

can directed to the corresponding author, Qizhao Wang.

Declaration of Generative AI and AI-assisted Technologies in the Writing Process.

During the preparation of this work the author(s) used ChatGPT to assist with revising the

format, grammar, and spelling. After using this tool/service, the author(s) reviewed and edited the

content as needed and take(s) full responsibility for the content of the publication.

ACKNOWLEDGMENTS

G.G. is supported by grants from the UMass Chan Medical School (an internal grant) and by the National Institutes of Health (R01NS076991-01, P01HL131471-05, R01AI121135, UG3HL147367-01, R01HL097088, R01HL152723-02, U19AI149646-01, and UH3HL147367-04). P.W.L.T. is supported by an award from The Bassick Family Foundation, a BRIDGE Fund Award (a UMass Chan Medical School internal grant), and by the National Institutes of Health (1R21AI183080-01A1). W.H. is supported by the National Institutes of Health (R01DA056876)

AUTHOR CONTRIBUTIONS

R.Y., J.Z., Y.L. designed and performed the AAV productivity experiments; Y.W. and Y.L. conducted all AAV-related characterizations; N.T.T. analyzed the NGS data and prepared figures. M.Cu., T.C., and T.S. designed and carried out animal studies involving AAV1, AAV2, and AAV8, and drafted the parts of the manuscript that encompass this work; X.Y., D.Z., and D.J. conducted animal studies for AAV-PHP.eB and AAVrh.10; Y.L., Z.S., Y.D., L.F., H.H., B.W., and C.X. constructed and prepared all plasmids; L.W., H.Z., Y.L., M.Ch, J.L., and C.C. performed AAV production, purification, and quality checks; G.G. revised and organized the manuscript, and supported the AAV1, AAV2, and AAV8 animal studies; W.H. provided support for the AAV-PHP.eB and AAVrh.10 animal studies. D.Y. revised the manuscript, contributed to the design of the experimental strategy, and provided overall support for the study; P.W.L.T. analyzed the data, drafted the manuscript, and prepared the figures; Q.W. designed constructs, drafted the manuscript and figures, and provided overarching support for the study.

DECLARATION OF INTERESTS

KEYWORDS (X/5-10)

REFERENCES

1.   Kuzmin, D.A., Shutova, M.V., Johnston, N.R., Smith, O.P., Fedorin, V.V., Kukushkin, Y.S., van

der Loo, J.C.M., and Johnstone, E.C. (2021). The clinical landscape for AAV gene therapies.

Nat Rev Drug Discov *20*, 173-174. 10.1038/d41573-021-00017-7.

2.   Wang, D., Tai, P.W.L., and Gao, G. (2019). Adeno-associated virus vector as a platform for

gene therapy delivery. Nat Rev Drug Discov *18*, 358-378. 10.1038/s41573-019-0012-9.

3.   Pupo, A., Fernandez, A., Low, S.H., Francois, A., Suarez-Amaran, L., and Samulski, R.J.

(2022). AAV vectors: The Rubik's cube of human gene therapy. Mol Ther *30*, 3515-3541.

10.1016/j.ymthe.2022.09.015.

4.   Zhang, X., De Alwis, M., Hart, S.L., Fitzke, F.W., Inglis, S.C., Boursnell, M.E., Levinsky, R.J.,

Kinnon, C., Ali, R.R., and Thrasher, A.J. (1999). High-titer recombinant adeno-associated virus

production from replicating amplicons and herpes vectors deleted for glycoprotein H. Hum Gene

Ther *10*, 2527-2537. 10.1089/10430349950016861.

5.   Urabe, M., Ding, C., and Kotin, R.M. (2002). Insect cells as a factory to produce adeno-

associated virus type 2 vectors. Hum Gene Ther *13*, 1935-1943. 10.1089/10430340260355347.

6.   Wang, Q., Wu, Z., Zhang, J., Firrman, J., Wei, H., Zhuang, Z., Liu, L., Miao, L., Hu, Y., Li, D.,

702   Diao, Y., et al. (2017). A Robust System for Production of Superabundant VP1 Recombinant

703   AAV Vectors. Mol Ther Methods Clin Dev *7*, 146-156. 10.1016/j.omtm.2017.11.002.

704   7.   Su, W., Patricio, M.I., Duffy, M.R., Krakowiak, J.M., Seymour, L.W., and Cawood, R. (2022).

705   Self-attenuating adenovirus enables production of recombinant adeno-associated virus for high

706   manufacturing yield without contamination. Nat Commun *13*, 1182. 10.1038/s41467-022-28738-

707   2.

708   8.   Penaud-Budloo, M., Francois, A., Clement, N., and Ayuso, E. (2018). Pharmacology of

709   Recombinant Adeno-associated Virus Production. Mol Ther Methods Clin Dev *8*, 166-180.

710   10.1016/j.omtm.2018.01.002.

711   9.   Srivastava, A., Mallela, K.M.G., Deorkar, N., and Brophy, G. (2021). Manufacturing Challenges

712   and Rational Formulation Development for AAV Viral Vectors. J Pharm Sci *110*, 2609-2624.

713   10.1016/j.xphs.2021.03.024.

714   10.   Jiang, Z., and Dalby, P.A. (2023). Challenges in scaling up AAV-based gene therapy

715   manufacturing. Trends Biotechnol *41*, 1268-1281. 10.1016/j.tibtech.2023.04.002.

716   11.   Tran, N.T., Lecomte, E., Saleun, S., Namkung, S., Robin, C., Weber, K., Devine, E., Blouin, V.,

717   Adjali, O., Ayuso, E., Gao, G., et al. (2022). Human and insect cell-produced rAAVs show

718   differences in genome heterogeneity. Hum Gene Ther. 10.1089/hum.2022.050.

719   12.   Xiao, X., Li, J., and Samulski, R.J. (1998). Production of high-titer recombinant adeno-

720   associated virus vectors in the absence of helper adenovirus. J Virol *72*, 2224-2232.

721   13.   Sonntag, F., Schmidt, K., and Kleinschmidt, J.A. (2010). A viral assembly factor promotes AAV2

722   capsid formation in the nucleolus. Proc Natl Acad Sci U S A *107*, 10220-10225.

723   10.1073/pnas.1001673107.

724   14.   Cao, M., You, H., and Hermonat, P.L. (2014). The X gene of adeno-associated virus 2 (AAV2) is

725   involved in viral DNA replication. PLoS One *9*, e104596. 10.1371/journal.pone.0104596.

15.   Elmore, Z.C., Patrick Havlik, L., Oh, D.K., Anderson, L., Daaboul, G., Devlin, G.W., Vincent,

H.A., and Asokan, A. (2021). The membrane associated accessory protein is an adeno-

associated viral egress factor. Nat Commun *12*, 6239. 10.1038/s41467-021-26485-4.

16.   Meier, A.F., Fraefel, C., and Seyffert, M. (2020). The Interplay between Adeno-Associated Virus

and its Helper Viruses. Viruses *12*. 10.3390/v12060662.

17.   Fu, Q., Polanco, A., Lee, Y.S., and Yoon, S. (2023). Critical challenges and advances in

recombinant adeno-associated virus (rAAV) biomanufacturing. Biotechnol Bioeng *120*, 2601-

2621. 10.1002/bit.28412.

18.   Young, P. (2023). Treatment to cure: Advancing AAV gene therapy manufacture. Drug Discov

Today *28*, 103610. 10.1016/j.drudis.2023.103610.

19.   Dash, S., Sharon, D.M., Mullick, A., and Kamen, A.A. (2022). Only a small fraction of cells

produce assembled capsids during transfection-based manufacturing of adeno-associated virus

vectors. Biotechnol Bioeng *119*, 1685-1690. 10.1002/bit.28068.

20.   Lavado-Garcia, J., Perez-Rubio, P., Cervera, L., and Godia, F. (2022). The cell density effect in

animal cell-based bioprocessing: Questions, insights and perspectives. Biotechnol Adv *60*,

108017. 10.1016/j.biotechadv.2022.108017.

21.   Moco, P.D., Xu, X., Silva, C.A.T., and Kamen, A.A. (2023). Production of adeno-associated viral

vector serotype 6 by triple transfection of suspension HEK293 cells at higher cell densities.

Biotechnol J *18*, e2300051. 10.1002/biot.202300051.

22.   Chung, C.H., Murphy, C.M., Wingate, V.P., Pavlicek, J.W., Nakashima, R., Wei, W., McCarty,

D., Rabinowitz, J., and Barton, E. (2023). Production of rAAV by plasmid transfection induces

antiviral and inflammatory responses in suspension HEK293 cells. Mol Ther Methods Clin Dev

28, 272-283. 10.1016/j.omtm.2023.01.002.

23.   Brimble, M.A., Winston, S.M., and Davidoff, A.M. (2023). Stowaways in the cargo:

Contaminating nucleic acids in rAAV preparations for gene therapy. Mol Ther *31*, 2826-2838.

10.1016/j.ymthe.2023.07.025.

24. Wright, J.F. (2014). Product-Related Impurities in Clinical-Grade Recombinant AAV Vectors:

Characterization and Risk Assessment. Biomedicines *2*, 80-97. 10.3390/biomedicines2010080.

25. Vandermeulen, G., Marie, C., Scherman, D., and Preat, V. (2011). New generation of plasmid

backbones devoid of antibiotic resistance marker for gene therapy trials. Mol Ther *19*, 1942-

1949. 10.1038/mt.2011.182.

26. Schnodt, M., Schmeer, M., Kracher, B., Krusemann, C., Espinosa, L.E., Grunert, A.,

Fuchsluger, T., Rischmuller, A., Schleef, M., and Buning, H. (2016). DNA Minicircle Technology

Improves Purity of Adeno-associated Viral Vector Preparations. Mol Ther Nucleic Acids *5*, e355.

10.1038/mtna.2016.60.

27. Lecomte, E., Tournaire, B., Cogne, B., Dupont, J.B., Lindenbaum, P., Martin-Fontaine, M.,

Broucque, F., Robin, C., Hebben, M., Merten, O.W., Blouin, V., et al. (2015). Advanced

Characterization of DNA Molecules in rAAV Vector Preparations by Single-stranded Virus Next-

generation Sequencing. Mol Ther Nucleic Acids *4*, e260. 10.1038/mtna.2015.32.

28. Tai, P.W.L., Xie, J., Fong, K., Seetin, M., Heiner, C., Su, Q., Weiand, M., Wilmot, D., Zapp, M.L.,

and Gao, G. (2018). Adeno-associated Virus Genome Population Sequencing Achieves Full

Vector Genome Resolution and Reveals Human-Vector Chimeras. Mol Ther Methods Clin Dev

9, 130-141. 10.1016/j.omtm.2018.02.002.

29. Chadeuf, G., Ciron, C., Moullier, P., and Salvetti, A. (2005). Evidence for encapsidation of

prokaryotic sequences during recombinant adeno-associated virus production and their in vivo

persistence after vector delivery. Mol Ther *12*, 744-753. 10.1016/j.ymthe.2005.06.003.

30. Samelson-Jones, B.J., and George, L.A. (2023). Adeno-Associated Virus Gene Therapy for

Hemophilia. Annu Rev Med *74*, 231-247. 10.1146/annurev-med-043021-033013.

31. Salabarria, S.M., Corti, M., Coleman, K.E., Wichman, M.B., Berthy, J.A., D'Souza, P., Tifft, C.J.,

Herzog, R.W., Elder, M.E., Shoemaker, L.R., Leon-Astudillo, C., et al. (2024). Thrombotic

microangiopathy following systemic AAV administration is dependent on anti-capsid antibodies.

J Clin Invest *134*. 10.1172/JCI173510.

32.   Hauck, B., Murphy, S.L., Smith, P.H., Qu, G., Liu, X., Zelenaia, O., Mingozzi, F., Sommer, J.M.,

779   High, K.A., and Wright, J.F. (2009). Undetectable transcription of cap in a clinical AAV vector:

780   implications for preformed capsid in immune responses. Mol Ther *17*, 144-152.

781   10.1038/mt.2008.227.

782   33.   Zelenaia, O., Sumaroka, M., Tai, L., Hauck, B., and Wright, J.F. (2012). 365. Characterization of

783   Residual Plasmid DNA Impurities in Recombinant AAV; Correlation with Size of Transgene

784   Expression Cassette. American Society of Gene and Cell Therapy. Molecular Therapy.

785   34.   Brimble, M.A., Zhou, J., Morton, C., Meagher, M., Nathwani, A.C., Gray, J.T., and Davidoff, A.M.

786   (2016). 547. AAV Preparations Contain Contamination from DNA Sequences in Production

787   Plasmids Directly Outside of the ITRs. Molecular Therapy.

788   35.   Jalsic, L., Lytvyn, V., Elahi, S.M., Hrapovic, S., Nassoury, N., Chahal, P.S., Gaillet, B., and

789   Gilbert, R. (2023). Inducible HEK293 AAV packaging cell lines expressing Rep proteins. Mol

790   Ther Methods Clin Dev *30*, 259-275. 10.1016/j.omtm.2023.07.002.

791   36.   Whiteley, Z., Ho, H.M.K., Gan, Y.X., Panariello, L., Gkogkos, G., Gavriilidis, A., and Craig,

792   D.Q.M. (2021). Microfluidic synthesis of protein-loaded nanogels in a coaxial flow reactor using

793   a design of experiments approach. Nanoscale Adv *3*, 2039-2055. 10.1039/d0na01051k.

794   37.   Emmerling, V.V., Pegel, A., Milian, E.G., Venereo-Sanchez, A., Kunz, M., Wegele, J., Kamen,

795   A.A., Kochanek, S., and Hoerer, M. (2016). Rational plasmid design and bioprocess

796   optimization to enhance recombinant adeno-associated virus (AAV) productivity in mammalian

797   cells. Biotechnol J *11*, 290-297. 10.1002/biot.201500176.

798   38.   Asaad, W., Volos, P., Maksimov, D., Khavina, E., Deviatkin, A., Mityaeva, O., and Volchkov, P.

799   (2023). AAV genome modification for efficient AAV production. Heliyon *9*, e15071.

800   10.1016/j.heliyon.2023.e15071.

801   39.   Grimm, D., Kern, A., Rittner, K., and Kleinschmidt, J.A. (1998). Novel tools for production and

802   purification of recombinant adenoassociated virus vectors. Hum Gene Ther *9*, 2745-2760.

803   10.1089/hum.1998.9.18-2745.

40.   Tang, Q., Keeler, A.M., Zhang, S., Su, Q., Lyu, Z., Cheng, Y., Gao, G., and Flotte, T.R. (2020).

Two-Plasmid Packaging System for Recombinant Adeno-Associated Virus. Biores Open Access

9, 219-228. 10.1089/biores.2020.0031.

41.   Kreiss, P., Cameron, B., Rangara, R., Mailhe, P., Aguerre-Charriol, O., Airiau, M., Scherman,

D., Crouzet, J., and Pitard, B. (1999). Plasmid DNA size does not affect the physicochemical

properties of lipoplexes but modulates gene transfer efficiency. Nucleic Acids Res *27*, 3792-

3798. 10.1093/nar/27.19.3792.

42.   Hornstein, B.D., Roman, D., Arevalo-Soliz, L.M., Engevik, M.A., and Zechiedrich, L. (2016).

Effects of Circular DNA Length on Transfection Efficiency by Electroporation into HeLa Cells.

PLoS One *11*, e0167537. 10.1371/journal.pone.0167537.

43.   Lesueur, L.L., Mir, L.M., and Andre, F.M. (2016). Overcoming the Specific Toxicity of Large

Plasmids Electrotransfer in Primary Cells In Vitro. Mol Ther Nucleic Acids *5*, e291.

10.1038/mtna.2016.4.

44.   van Lieshout, L.P., Rubin, M., Costa-Grant, K., Ota, S., Golebiowski, D., Panico, T., Wiberg, E.,

Szymczak, K., Gilmore, R., Stanvick, M., Burnham, B., et al. (2023). A novel dual-plasmid

platform provides scalable transfection yielding improved productivity and packaging across

multiple AAV serotypes and genomes. Mol Ther Methods Clin Dev *29*, 426-436.

10.1016/j.omtm.2023.05.004.

45.   Joshi, P.R.H., Venereo-Sanchez, A., Chahal, P.S., and Kamen, A.A. (2021). Advancements in

molecular design and bioprocessing of recombinant adeno-associated virus gene delivery

vectors using the insect-cell baculovirus expression platform. Biotechnol J *16*, e2000021.

10.1002/biot.202000021.

46.   Philpott, N.J., Gomos, J., Berns, K.I., and Falck-Pedersen, E. (2002). A p5 integration efficiency

element mediates Rep-dependent integration into AAVS1 at chromosome 19. Proc Natl Acad

Sci U S A *99*, 12381-12385. 10.1073/pnas.182430299.

47. Xiong, W., Wu, D.M., Xue, Y., Wang, S.K., Chung, M.J., Ji, X., Rana, P., Zhao, S.R., Mai, S.,

and Cepko, C.L. (2019). AAV cis-regulatory sequences are correlated with ocular toxicity. Proc

Natl Acad Sci U S A *116*, 5785-5794. 10.1073/pnas.1821000116.

48. Cheng, S.Y., Luo, Y., Malachi, A., Ko, J., Su, Q., Xie, J., Tian, B., Lin, H., Ke, X., Zheng, Q., Tai,

P.W.L., et al. (2021). Low-Dose Recombinant Adeno-Associated Virus-Mediated Inhibition of

Vascular Endothelial Growth Factor Can Treat Neovascular Pathologies Without Inducing

Retinal Vasculitis. Hum Gene Ther *32*, 649-666. 10.1089/hum.2021.132.

49. Chan, Y.K., Wang, S.K., Chu, C.J., Copland, D.A., Letizia, A.J., Costa Verdera, H., Chiang, J.J.,

Sethi, M., Wang, M.K., Neidermyer, W.J., Jr., Chan, Y., et al. (2021). Engineering adeno-

associated viral vectors to evade innate immune and inflammatory responses. Sci Transl Med

13. 10.1126/scitranslmed.abd3438.

50. Cui, M., Su, Q., Yip, M., McGowan, J., Punzo, C., Gao, G., and Tai, P.W.L. (2024). The

AAV2.7m8 capsid packages a higher degree of heterogeneous vector genomes than AAV2.

Gene Ther *31*, 489-498. 10.1038/s41434-024-00477-7.

51. Namkung, S., Tran, N.T., Manokaran, S., He, R., Su, Q., Xie, J., Gao, G., and Tai, P.W.L.

(2022). Direct ITR-to-ITR Nanopore Sequencing of AAV Vector Genomes. Hum Gene Ther *33*,

1187-1196. 10.1089/hum.2022.143.

52. Zhang, J., Guo, P., Yu, X., Frabutt, D.A., Lam, A.K., Mulcrone, P.L., Chrzanowski, M., Firrman,

J., Pouchnik, D., Sang, N., Diao, Y., et al. (2022). Subgenomic particles in rAAV vectors result

from DNA lesion/break and non-homologous end joining of vector genomes. Mol Ther Nucleic

Acids *29*, 852-861. 10.1016/j.omtn.2022.08.027.

53. Xie, J., Mao, Q., Tai, P.W.L., He, R., Ai, J., Su, Q., Zhu, Y., Ma, H., Li, J., Gong, S., Wang, D., et

al. (2017). Short DNA Hairpins Compromise Recombinant Adeno-Associated Virus Genome

Homogeneity. Mol Ther *25*, 1363-1374. 10.1016/j.ymthe.2017.03.028.

54.   Shitik, E.M., Shalik, I.K., and Yudkin, D.V. (2023). AAV- based vector improvements unrelated

to capsid protein modification. Front Med (Lausanne) *10*, 1106085.

10.3389/fmed.2023.1106085.

55.   Wilmott, P., Lisowski, L., Alexander, I.E., and Logan, G.J. (2019). A User's Guide to the Inverted

Terminal Repeats of Adeno-Associated Virus. Hum Gene Ther Methods *30*, 206-213.

10.1089/hgtb.2019.276.

56.   Tran, N.T., Heiner, C., Weber, K., Weiand, M., Wilmot, D., Xie, J., Wang, D., Brown, A.,

Manokaran, S., Su, Q., Zapp, M.L., et al. (2020). AAV-Genome Population Sequencing of

Vectors Packaging CRISPR Components Reveals Design-Influenced Heterogeneity. Mol Ther

Methods Clin Dev *18*, 639-651. 10.1016/j.omtm.2020.07.007.

57.   Muralidharan, N. (2023). Quantitative Risk Assessment of Limits for Residual Host-Cell DNA:

Ensuring Patient Safety for In Vitro Gene Therapies Produced Using Human-Derived Cell Lines.

Bioprocess International *21*, 33-37.

58.   Yip, M., Chen, J., Zhi, Y., Tran, N.T., Namkung, S., Pastor, E., Gao, G., and Tai, P.W.L. (2023).

Querying Recombination Junctions of Replication-Competent Adeno-Associated Viruses in

Gene Therapy Vector Preparations with Single Molecule, Real-Time Sequencing. Viruses *15*.

10.3390/v15061228.

59.   Shmidt, A.A., and Egorova, T.V. (2021). PCR-Based Analytical Methods for Quantification and

Quality Control of Recombinant Adeno-Associated Viral Vector Preparations. Pharmaceuticals

(Basel) *15*. 10.3390/ph15010023.

60.   Huang, X., Hartley, A.V., Yin, Y., Herskowitz, J.H., Lah, J.J., and Ressler, K.J. (2013). AAV2

production with optimized N/P ratio and PEI-mediated transfection results in low toxicity and

high titer for in vitro and in vivo applications. J Virol Methods *193*, 270-277.

10.1016/j.jviromet.2013.06.008.

61.   Chahal, P.S., Schulze, E., Tran, R., Montes, J., and Kamen, A.A. (2014). Production of adeno-

878   associated virus (AAV) serotypes by transient transfection of HEK293 cell suspension cultures

879   for gene delivery. J Virol Methods *196*, 163-173. 10.1016/j.jviromet.2013.10.038.

880   62.   Grieger, J.C., Soltys, S.M., and Samulski, R.J. (2016). Production of Recombinant Adeno-

881   associated Virus Vectors Using Suspension HEK293 Cells and Continuous Harvest of Vector

882   From the Culture Media for GMP FIX and FLT1 Clinical Vector. Mol Ther *24*, 287-297.

883   10.1038/mt.2015.187.

884   63.   Zhao, H., Lee, K.J., Daris, M., Lin, Y., Wolfe, T., Sheng, J., Plewa, C., Wang, S., and Meisen,

885   W.H. (2020). Creation of a High-Yield AAV Vector Production Platform in Suspension Cells

886   Using a Design-of-Experiment Approach. Mol Ther Methods Clin Dev *18*, 312-320.

887   10.1016/j.omtm.2020.06.004.

888   64.   Tran, N.T., Lecomte, E., Saleun, S., Namkung, S., Robin, C., Weber, K., Devine, E., Blouin, V.,

889   Adjali, O., Ayuso, E., Gao, G., et al. (2022). Human and Insect Cell-Produced Recombinant

890   Adeno-Associated Viruses Show Differences in Genome Heterogeneity. Hum Gene Ther *33*,

891   371-388. 10.1089/hum.2022.050.

892   65.   Afgan, E., Sloggett, C., Goonasekera, N., Makunin, I., Benson, D., Crowe, M., Gladman, S.,

893   Kowsar, Y., Pheasant, M., Horst, R., and Lonie, A. (2015). Genomics Virtual Laboratory: A

894   Practical Bioinformatics Workbench for the Cloud. PLoS One *10*, e0140829.

895   10.1371/journal.pone.0140829.

896   66.   Blankenberg, D., Von Kuster, G., Coraor, N., Ananda, G., Lazarus, R., Mangan, M., Nekrutenko,

897   A., and Taylor, J. (2010). Galaxy: a web-based genome analysis tool for experimentalists. Curr

898   Protoc Mol Biol *Chapter 19*, Unit 19 10 11-21. 10.1002/0471142727.mb1910s89.

899   67.   Giardine, B., Riemer, C., Hardison, R.C., Burhans, R., Elnitski, L., Shah, P., Zhang, Y.,

900   Blankenberg, D., Albert, I., Taylor, J., Miller, W., et al. (2005). Galaxy: a platform for interactive

901   large-scale genome analysis. Genome Res *15*, 1451-1455. 10.1101/gr.4086505.

902   68.   Goecks, J., Nekrutenko, A., Taylor, J., and Galaxy, T. (2010). Galaxy: a comprehensive

903   approach for supporting accessible, reproducible, and transparent computational research in the

904   life sciences. Genome Biol *11*, R86. 10.1186/gb-2010-11-8-r86.

905   69.   Robinson, J.T., Thorvaldsdottir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and

906   Mesirov, J.P. (2011). Integrative genomics viewer. Nat Biotechnol *29*, 24-26. 10.1038/nbt.1754.

FIGURE LEGENDS

Figure 1. Design and testing of the AAVone production system. (A) Schematic illustrations of the

AAVone, triple-plasmid (standard, pTri-plasmid; and mini-pHelper, mTri-plasmid) systems, dual-plasmid

(pDG-like and AAVdual) systems, and the AAVone system. **(B)** A comparative analysis of AAV

productivity using AAVone and alternative systems in adherent HEK-293T cells. For every system

compared, the total pDNA amount was maintained at 0.5 µg/well (24-well plate), and the PEImax to

DNA ratio was consistently kept at 3:1. The plasmid molecular ratios for each system is displayed

under each data set. **(C)** A comparison of vector yields with different serotypes between the AAVone

and the pTri-plasmid system in suspension HEK-293T cells. The pTri-plasmid set was transfected at

equal plasmid molecular ratios at a total DNA amount of 1 µg per million cells, at a cell density of 3E6

cells/mL. Crude lysate samples were quantified using qPCR with primers targeting *Egfp*, and data were

normalized to the average values of each pTri-plasmid group. **(D)** Productivity optimization of the

AAVone system in suspension cultured HEK-293T cells as assessed by crude lysate titration. Different

cell densities, total pDNA, and transfection reagent (PEImax) were examined.

Figure 2. Characterization of AAV vectors produced by the AAVone and tri-plasmid systems.

Purified AAV vectors produced in suspension HEK-293T under different production platforms. **(A)**

Purified AAV vectors (~5E11 particles) from pTri-plasmid, mTri-plasmid, and AAVone were

electrophoresed and visualized by Coomassie G-250 staining. **(B)** Automated electrophoresis analysis

(TapeStation) of vector DNAs illustrating full-length genomes (arrow) and partial genomes (bracket).

Arrow heads designate specific bands identified by software. **(C)** Mass photometry analysis of AAV

particles. y-axis, particle counts; x-axis, mass of particles. **(D)** Transduction activity of vectors

packaging the AAV2-*CMV-Egfp* reporter produced by AAVone or triple-plasmid systems in HeLa cells

at an MOI of 10,000 vg/cell. and EGFP expression was observed 48 hrs post-transduction.

Figure 3. AAVone-packaged AAV2 vectors exhibit similar *in vivo* efficacies as vectors produce d

by tri-plasmid systems. (A) Representative funduscopy images of murine retinas at 4 weeks post-

injection. **(B)** Representative retina cross-sections stained to visualize EGFP (anti-EGFP, green), nuclei

(DAPI, blue), cone photoreceptors (PNA, white), and microglia (IBA1, red) at 6-weeks post-injection.

Cross sections of representative whole eye cups (top row) and zoom ins to show transduction of retinal

layers (bottom row). Dotted box in top panels show position of zoom-in areas. Scale bars = 0.5 mM.

(C) Quantification of EGFP$^+$ cells in the photoreceptor (PR) layer. **(D,E)** ddPCR quantification of vector

genomes **(D)** and vector transcripts **(E)** in retina samples at 6-weeks post-injection (n=4). **(F)** Counts of

total microglia in retina cross-sections at 6 weeks post-injection. **(G)** Representative cross-section of a

vector-treated retina illustrating microglial (IBA1$^+$) infiltration in retinal layers: photoreceptor segment

layer (PS); outer nuclear layer (ONL); outer plexiform layer (OPL); inner nuclear layer (INL); inner

plexiform layer (IPL); ganglion cell layer (GCL). IBA1$^+$ cells (red) and DAPI (blue). **(H)** Percentage of

IBA1$^+$ cells in each retinal layer. Values represent means ±SD, **$p<0.01$; ****$p<0.0001$, ns = not

significant; by one-way ANOVA and Tukey's multiple comparison

Figure 4. Characterization of AAV vectors produced by the AAVone and tri-plasmid systems. **(A)**

Distribution of SMRT sequencing read lengths from vectors produced by pTri-plasmid system (left),

mTri-plasmid(center), and AAVone (right). Read lengths are binned into 10-nt increments. The

percentage of full-lengh genomes (2,770 ±10 nt) are shown for each graph. The y axes shown are

separated into two segments to highlight the low prevalence of truncated genomes. **(B)** Relative counts

of mapped read start (blue) and end (red) positions, indicating truncation hotspots. The y axes shown

are separated into two segments to highlight the low prevalence of start and end positions beyond the

ITR ends. **(C)** Sequences of the 119-nt left ITR (L-ITR) and the 130-nt right ITR (R-ITR) used in the

plasmid vector designs. **(D)** Length distribution of the L-ITRs and R-ITRs from the three production

systems examined. Read counts are on the x axes, and the read lengths are on the y axes. **(E)** A

summary table of the number of 11-nt C arm deleted ITRs at the L-ITRs and R-ITRs identified among

the SMRT reads. The percentages of the Δ11-nt C arm ITRs are also shown.

Figure 5. **Abundance of producer plasmid contaminants in AAV vectors generated by tri-**

plasmid systems and AAVone. (A-B) Coverage traces of aligned SMRT reads to the pAAVone **(A),**

pAAV-CMV-EGFP **(B)** and AAVone **(C)** plasmid references. Dashed lines indicate the boundaries of

the ITR references. **(D)** ddPCR quantification of *trans* plasmid sequences (*rep* and *cap*), AdV helper

plasmid (*VA RNA*), backbone sequences (pMB1 *ori* and *F1 ori*), and rcAAV in vector preparations.

Values represent means ±SD (n=3). *, $p \leq 0.05$; ****, $p \leq 0.0001$ by 2-way ANOVA. Color of astrisks

correspond to *p* values when compared to pTri-plasmid (red), mTri-plasmid (blue), and AAVone

(purple) data.

Figure 6. Examination of SMRT reads for the presence of rcAAV. IGV display of SMRT reads

representing genomes obtained from vectors produced by pTri-plasmid (top), mTri-plasmid (middle)

and AAVone (bottom) systems. Alignments are shown with soft-clipped bases visible. Read matches

(gray), mismatches (colored), and insertions/deletions (speckles) are shown.

Figure 7. Reduction of AAVone plasmid increases the quality of vectors. (A) and **(B)** Effect of

pDNA amount on AAV productivity of suspension HEK 293T(A) and HEK 293 (B) cells. **(C-E)** Analytical

ultracentrifugation analyses of AAV2 vectors produced by triple-plasmid transfections (pTri-plasmid and

mTri-plasmid) and AAVone produced under different plasmid amounts **(C)**. AUC analyses of AAV8 **(D)**

and AAV9 **(E)** vectors under different plasmid amounts x axis = sedimentation coefficient (S). The y

axis = normalized distribution $c(s)$. **(F)** qPCR analyses of DNase-resistant target regions packaged with

AAVone-produced vectors. x axes represent positions moving away from the ITRs.

.

Figure 1. Design and testing of the AAVone production system. (A) Schematic illustrations of the AAVone, triple-plasmid (standard, pTri-plasmid; and mini-pHelper, mTri-plasmid) systems, dual-plasmid (pDG-like and AAVdual) systems, and the AAVone system. **(B)** A comparative analysis of AAV productivity using AAVone and alternative systems in adherent HEK-293T cells. For every system compared, the total pDNA amount was maintained at 0.5 µg/well (24-well plate), and the PEImax to DNA ratio was consistently kept at 3:1. The plasmid molecular ratios for each system is displayed under each data set. **(C)** A comparison of vector yields with different serotypes between the AAVone and the pTri-plasmid system in suspension HEK-293T cells. The pTri-plasmid set was transfected at equal plasmid molecular ratios at a total DNA amount of 1 µg per million cells, at a cell density of 3E6 cells/mL. Crude lysate samples were quantified using qPCR with primers targeting Egfp, and data were normalized to the average values of each pTri-plasmid group. **(D)** Productivity optimization of the AAVone system in suspension cultured HEK-293T cells as assessed by crude lysate titration. Different cell densities, total pDNA, and transfection reagent (PEImax) were examined.

Figure 2. Characterization of AAV vectors produced by the AAVone and tri-plasmid systems. Purified AAV vectors produced in suspension HEK-293T under different production platforms. **(A)** Purified AAV vectors (~5E11 particles) from pTri-plasmid, mTri-plasmid, and AAVone were electrophoresed and visualized by Coomassie G-250 staining. **(B)** Automated electrophoresis analysis (TapeStation) of vector DNAs illustrating full-length genomes (arrow) and partial genomes (bracket). Arrow heads designate specific bands identified by software. **(C)** Mass photometry analysis of AAV particles. y-axis, particle counts; x-axis, mass of particles. **(D)** Transduction activity of vectors packaging the AAV2-*CMV-Egfp* reporter produced by AAVone or triple-plasmid systems in HeLa cells at an MOI of 10,000 vg/cell. and EGFP expression was observed 48 hrs post-transduction.

Figure 3. AAVone-packaged AAV2 vectors exhibit similar *in vivo* efficacies as vectors produce d by triple-plasmid systems. (A) Representative funduscopy images of murine retinas at 4 weeks post-injection. **(B)** Representative retina cross-sections stained to visualize EGFP (anti-EGFP, green), nuclei (DAPI, blue), cone photoreceptors (PNA, white), and microglia (IBA1, red) at 6-weeks post-injection. Cross sections of representative whole eye cups (top row) and zoom ins to show transduction of retinal layers (bottom row). Dotted box in top panels show position of zoom-in areas. Scale bars = 0.5 mM. **(C)** Quantification of EGFP+ cells in the photoreceptor (PR) layer. **(D,E)** ddPCR quantification of vector genomes **(D)** and vector transcripts **(E)** in retina samples at 6-weeks post-injection (n=4). **(F)** Counts of total microglia in retina cross-sections at 6 weeks post-injection. **(G)** Representative cross-section of a vector-treated retina illustrating microglial (IBA1+) infiltration in retinal layers: photoreceptor segment layer (PS); outer nuclear layer (ONL); outer plexiform layer (OPL); inner nuclear layer (INL); inner plexiform layer (IPL); ganglion cell layer (GCL). IBA1+ cells (red) and DAPI (blue). **(H)** Percentage of IBA1+ cells in each retinal layer. Values represent means ±SD, **$p<0.01$; ****$p<0.0001$, ns = not significant; by one-way ANOVA and Tukey's multiple comparison

Figure 4. Characterization of AAV vectors produced by the AAVone and tri-plasmid systems. **(A)** Distribution of SMRT sequencing read lengths from vectors produced by pTri-plasmid system (left), mTri-plasmid(center), and AAVone (right). Read lengths are binned into 10-nt increments. The percentage of full-lengh genomes (2,770 ±10 nt) are shown for each graph. The y axes shown are separated into two segments to highlight the low prevalence of truncated genomes. **(B)** Relative counts of mapped read start (blue) and end (red) positions, indicating truncation hotspots. The y axes shown are separated into two segments to highlight the low prevalence of start and end positions beyond the ITR ends. **(C)** Sequences of the 119-nt left ITR (L-ITR) and the 130-nt right ITR (R-ITR) used in the plasmid vector designs. **(D)** Length distribution of the L-ITRs and R-ITRs from the three production systems examined. Read counts are on the x axes, and the read lengths are on the y axes. **(E)** A summary table of the number of 11-nt C arm deleted ITRs at the L-ITRs and R-ITRs identified among the SMRT reads. The percentages of the Δ11-nt C arm ITRs are also shown.

Figure 5. **Abundance of producer plasmid contaminants in AAV vectors generated by triple-plasmid systems and AAVone. (A-B)** Coverage traces of aligned SMRT reads to the pAAVone **(A),** pAAV-CMV-EGFP **(B)** and AAVone **(C)** plasmid references. Dashed lines indicate the boundaries of the ITR references. **(D)** ddPCR quantification of *trans* plasmid sequences (*rep* and *cap*), AdV helper plasmid (*VA RNA*), backbone sequences (pMB1 *ori* and *F1 ori*), and rcAAV in vector preparations. Values represent means ±SD (n=3). *, $p \leq 0.05$; ****, $p \leq 0.0001$ by 2-way ANOVA. Color of astrisks correspond to *p* values when compared to pTri-plasmid (red), mTri-plasmid (blue), and AAVone (purple) data.

pTri-plasmid

Count:
27/131,989

mTri-plasmid

Count:
22/119,029

AAVone

Count:
90/196,753

Figure 6. Examination of SMRT reads for the presence of rcAAV. IGV display of SMRT reads representing genomes obtained from vectors produced by pTri-plasmid (top), mTri-plasmid (middle) and AAVone (bottom) systems. Alignments are shown with soft-clipped bases visible. Read matches (gray), mismatches (colored), and insertions/deletions (speckles) are shown.

Figure 7. Reduction of AAVone plasmid increases the quality of vectors. (A) and **(B)** Effect of pDNA amount on AAV productivity of suspension HEK 293T(A) and HEK 293 (B) cells. **(C-E)** Analytical ultracentrifugation analyses of AAV2 vectors produced by triple-plasmid transfections (pTri-plasmid and mTri-plasmid) and AAVone produced under different plasmid amounts **(C)**. AUC analyses of AAV8 **(D)** and AAV9 **(E)** vectors under different plasmid amounts x axis = sedimentation coefficient (S). The y axis = normalized distribution c(s). **(F)** qPCR analyses of DNase-resistant target regions packaged with AAVone-produced vectors. x axes represent positions moving away from the ITRs.

Table S1. AAV productivity of AAVone systems in suspension cultured HEK 293T and HEK 293 cells.

Cell lines	pAAVone plasmids	pAAVone size (kb)	AAV Serotype	AAV size (kb)	Scale (mL)	Cell density (1E6 cells/ml)	DNA Amount (ug/1E6 cells)	Transfection reagents (ratios with DNA)	Crude lysate titer (GC/L)	Purified vector yield (GC/L)
HEK 293T	pAAVone-AAV1-CMV-EGFP-2.7kb	15.7	AAV1	2.7	130	3.0	0.75	PEIMax (2:1)	1.83E+15	2.84E+14
	pAAVone-AAV2-CMV-EGFP-2.7kb	15.7	AAV2	2.7	130	3.0	0.75	PEIMax (2:1)	4.90E+14	1.59E+14
					160	3.0	0.5	PEIMax (2:1)	6.92E+14	1.46E+14
					160	3.0	0.5	PEIMax (3:1)	7.91E+14	1.61E+14
					400	3.0	0.5	PEIMax (2:1)	8.90E+14	2.14E+14
	pAAVone-AAV2-CMV-EGFP-2.2K	15.2	AAV2	2.2	90	3.0	0.5	PEIMax (2:1)	/	1.90E+14
	pAAVone-AAV5-CMV-EGFP-2.7kb	15.7	AAV5	2.7	130	3.0	0.75	PEIMax (2:1)	1.23E+15	1.65E+14
	pAAVone-AAV6-CMV-EGFP-2.7kb	15.7	AAV6	2.7	130	3.0	0.75	PEIMax (2:1)	2.01E+14	5.20E+13
	pAAVone-AAV8-CMV-EGFP-2.7kb	15.7	AAV8	2.7	130	3.0	0.75	PEIMax (2:1)	1.06E+15	1.66E+14
	pAAVone-AAV8-CMV-EGFP-2.2K	15.2	AAV8	2.2	90	3.0	0.75	PEIMax (2:1)	1.56E+15	3.29E+14
	pAAVone-AAV9-CMV-EGFP-2.7kb	15.7	AAV9	2.7	130	3.0	0.75	PEIMax (2:1)	3.70E+15	1.03E+15
					2100	3.0	0.75	PEIMax (2:1)	2.04E+15	5.20E+14
	pAAVone-AAV9-CMV-EGFP-2.2kb	15.2	AAV9	2.2	90	3.0	1.0	PEIMax (2:1)	1.96E+15	4.77E+14
					90	3.0	0.75	PEIMax (2:1)	2.49E+15	5.99E+14
					90	3.0	0.5	PEIMax (2:1)	3.07E+15	6.80E+14
					90	3.5	0.5	PEIMax (2:1)	3.01E+15	7.87E+14
					90	4.0	0.5	PEIMax (2:1)	2.38E+15	4.22E+14
	pAAVone-AAV9-CMV-EGFP-4.4kb	17.4	AAV9	4.4	90	3.0	0.25	PEIMax (2:1)	2.26E+15	/
	pAAVone-AAV9-CMV-EGFP-4.6kb	17.6	AAV9	4.6	90	3.0	0.25	PEIMax (2:1)	1.76E+15	/
	pAAVone-AAV9-CMV-EGFP-4.7kb	17.7	AAV9	4.7	90	3.0	0.25	PEIMax (2:1)	1.94E+15	/
	pAAVone-AAVrh.10-CMV-EGFP-2.7kb	15,719	AAVrh.10	2.7	130	3.0	0.75	PEIMax (2:1)	1.61E+15	3.42E+14
HEK 293	pAAVone-AAV2-CMV-EGFP-2.2K	15.2	AAV2	2.2	30	3.0	0.35	PEIMax (2:1)	3.54E+14	/
	pAAVone-AAV6-CMV-EGFP-2.2K	15.2	AAV6	2.2	30	3.0	0.35	PEIMax (2:1)	2.91E+14	/
	pAAVone-AAV8-CMV-EGFP-2.2K	15.2	AAV8	2.2	30	3.0	0.35	PEIMax (2:1)	8.69E+14	/
	pAAVone-AAV9-CMV-EGFP-2.2K	15.2	AAV9	2.2	30	3.0	0.50	PEIMax (2:1)	1.55E+15	/
	pAAVone-AAV2-CMV-EGFP-2.2K	15.2	AAV2	2.2	30	3.0	0.375	PEIMax (2:1)	4.98E+14	/
					30	3.0	0.25	PEIMax (2:1)	4.66E+14	/
					30	3.0	0.375	FectoVIR (1:1)	4.42E+14	/
					30	3.0	0.25	FectoVIR (1:1)	5.12E+14	/
					30	3.0	0.375	FectoVIR (2:1)	6.48E+14	/
					30	3.0	0.25	PEIMax (2:1)	6.53E+14	/
	pAAVone-AAV9-CMV-EGFP-2.2K	15.2	AAV9	2.2	30	3.0	0.35	PEIMax (2:1)	1.53E+15	/
					30	3.0	0.35	PEIMax (2:1)	1.21E+15	/
					30	3.0	0.50	PEIMax (1:1)	3.19E+15	/
					30	3.0	0.30	PEIMax (1:1)	2.79E+15	/

Table S2. Summary of aligned SMRT reads to references

References	pHelper counts (%)	mini-pHelper counts (%)	AAVone counts (%)
GOI from ITR-to-ITR	126,756 (96.04%)	115,005 (96.62%)	190,658 (96.90%)
GOI Full Reference	127,018 (96.23%)	115,164 (96.75%)	191,168 (97.16%)
GOI from ITR-to-ITR Carrying Backbone	5,209 (3.95%)	3,204 (2.69%)	1,715 (0.87%)
GOI backbone Only	152 (0.12%)	85 (0.07%)	105 (0.05%)
Rep Only	15 (0.01%)	3 (0.00%)	10 (0.01%)
Rep Chimeric with GOI	377 (0.29%)	243 (0.20%)	89 (0.05%)
Cap only	28 (0.02%)	28 (0.02%)	29 (0.01%)
Cap Chimeric with GOI	336 (0.25%)	240 (0.20%)	201 (0.10%)
Rep and Cap overlap	51 (0.04%)	18 (0.02%)	66 (0.03%)
Rep-Cap Carrying Backbone	63 (0.05%)	51 (0.04%)	N/A
Rep-Cap Backbone	224 (0.17%)	14 (0.01%)	N/A
Rep-Cap Carrying with GOI	687 (0.52%)	466 (0.39%)	319 (0.16%)
E2 only	63 (0.05%)	34 (0.03%)	90 (0.05%)
E2 chimeric with GOI	389 (0.29%)	335 (0.28%)	610 (0.31%)
E4 only	12 (0.01%)	11 (0.01%)	34 (0.02%)
E4 chimeric with GOI	147 (0.11%)	150 (0.13%)	260 (0.13%)
VA RNA only	1 (0.00%)	0 (0.00%)	0 (0.00%)
VA RNA chimeric with GOI	131 (0.10%)	123 (0.10%)	4,614 (2.35%)
pHelper/mini-pHelper carrying backbone	278 (0.21%)	181 (0.15%)	N/A
pHelper/mini-pHelper backbone	0 (0.00%)	0 (0.00%)	N/A
Backbone combined(AAVone only)	N/A	N/A	7,950 (4.04%)
hg38	4,453 (3.37%)	3,471 (2.92%)	5,311 (2.70%)
hg38 chimeric with GOI	677 (0.51%)	527 (0.44%)	814 (0.41%)
unmapped reads	921 (0.70%)	715 (0.60%)	990 (0.50%)
Total non-AAV containing reads	4,697 (3.56%)	3,659 (3.07%)	5,487 (2.79%)
Total plasmid impurities	8,303 (6.29%)	5,591 (4.70%)	7,899 (4.01%)
Total reads	131,989	119,029	196,753

Figure S1. Construction of mini-pHelper. The original pHelper contains 9.3 kb of the AdV2 genome with three expression cassettes (gray), including *VA RNA* (731 bp), *E4orf6* (3,192 bp) and *E2A* (5,343 bp). The *E2A* and *E4orf6* components are the main contributors to the plasmid's large size. The *E2A* gene transcripts: early (E2A-E) and late (E2A-L) mRNAs share a small second intron, but have different first intron. The *E4orf6* has a 1,275 bp intron. In the mini-pHelper, intron 1 and 2 of E2A-L, as well as the intron of *E4orf6* were eliminated, resulting in a 8.4-kb plasmid with 6.1 kb of AdV2. Red, promoters; green, ORFs; arrows, introns.

Figure S2. Examination of pAAVone plasmid yields. Comparisons of plasmid concentrations produced by mini-preps **(A)** and midi-preps **(B)** across multiple serotypes.

Figure S3. Transfection efficiency of the pAAV-CMV-Egfp, pTri-plasmid system and the pAAVone in adherent HEK 293T cells. At 48-h post-transfection, EGFP was observed by epifluorescence **(A),** and the mean fluorescence intensity **(B)** and percentage of EGFP-positive cells **(C)** were analysis with flow cytometry.

Figure S4. AAVone-packaged AAV2 and AAV8 vectors exhibit similar muscle transduction efficacies to vectors packaged by triple-plasmid transfection. Representative epifluorescence microscopy images **(A)** and H&E staining **(B)** of TA cross-sections at 2 weeks post-injection with ssAAV1.*CMV.Egfp* **(top row)** and ssAAV8.*CMV.Egfp* **(bottom row)** produced via the three three packaging systems. EGFP (green) and nuclei (blue). Tissues from mice injected by saline solution serve as negative controls.

Figure S5. AAVone-packaged AAV-PHP.eB and AAVrh.10 vectors exhibit similar brain transduction efficacies to vectors packaged by triple-plasmid transfection. Epifluorescence images of mouse brain cryosections treated with AAV-PHP.eB vectors (top row) or AAVrh.10 vectors (bottom row) produced by AAVone or the triple-plasmid systems. Sections were stained with anti-GFP antibodies and Hoechst dye to label nuclei.

Figure S6. Genome integrity analysis of vectors produced by tri-plasmid systems and AAVone.
(**A, B**) IGV display of SMRT reads representing AAV vector genomes aligned to the standard *cis* plasmid
(**A**) from the pTri-plasmid (top) and the mTri-plasmid (bottom) packaging systems, and the AAVone (**B**)
reference genomes. Both constructs carry the *CMV-Egfp* cassette. Reads are shown in squished displays
with soft-clipped bases visible. Read matches (gray), mismatches (colored), and insertions/deletions
(speckles) are shown. Alignment coverages are shown above each display.

Figure S7. Repair of the Δ11-nt C arm ITR. (A) IGV display of left and right ITR regions extracted from SMRT reads from pTri-plasmid-, mTri-plasmid-, and AAVone-produced vectors aligned to the 145-nt flop-oriented ITR references. Alignments are shown in squished displays with soft-clipped bases visible. Read matches (gray), mismatches (colored), and insertions/deletions (speckles) are shown. Alignment coverages are shown above each display. **(B)** Distributions of ITR configurations at the left and right ITRs from the three production platforms. Flip (red); Flop (blue); 11-nt deletion, Δ11-nt in C arm (orange); B arm deletion, ΔB arm (grey); C arm deletion, ΔC arm (yellow); external A region deletion, ΔExt. A region (purple); truncation of the A region, Trunc. Ext. A region (green); and trident-shaped ITR (cyan).

Figure S8. Profiling of hcDNA contaminants associated with triple-plasmid systems and AAVone. The Venn diagrams (left) show mapped reads to vector genome and to human genome hg38, as well as chimeric mapped reads for the pTri-plasmid (top), mTri-plasmid (middle), and AAVone (bottom) packaging systems. The number of reads mapped exclusively to the vector genome (ITR-to-ITR, white) and hg38 (human genome) are shown. The distribution of mapped reads to individual chromosomes can also be observed in the bar graph.

	pTri-plasmid	mTri-plasmid	AAVone
Average hcDNA size	1,327	1,298	1,346
Mean hcDNA size	1,146	1,146	1,142

Figure S9. Length distributions of SMRT reads that map to hg38. Reads from vectors produced by the pTri-plasmid (top), mTri-plasmid (middle), and AAVone (bottom) systems are shown. A table summarizing the average amdn mean sizes are shown.

Figure S10. Alignment of SMRT reads from DNase-resistant DNA isolated from vectors to chrM.
Reads from vectors produced by the pTri-plasmid (top), mTri-plasmid (middle), and AAVone (bottom)
systems are shown mapping to the mitochondrial genome. Tracks of the mitochondrial genes are shown
above.

Figure S11. Alignment of chimeric reads to the GOI reference. Reads from vectors produced by the pTri-plasmid (top), mTri-plasmid (middle), and AAVone (bottom) systems are shown. Alignments are shown in squished displays with soft-clipped bases visible. Read matches (gray), mismatches (colored), and insertions/deletions (speckles) are shown. Multiple **r**eads falsely map to the human GH polyA (bracket).

Figure S12. Reduction of AAVone plasmid used in production increases vector titers. (A-C) Cell metrics following transfection with different amounts of packaging plasmids. Percentage of EGFP-positive cells **(A),** cell viability **(B),** and cell density **(C)** are reported at 48-hours (blue bars) or 72-hours (orange bars) post-transfection.

Figure S13. Theoretical relationship of transfection (TF) efficiency and AAV production efficiency with AAVone and triple-plasmid systems.