Neural encoding of biomechanically (im)possible human movements in occipitotemporal cortex

Giuseppe Marrazzo[1], Federico De Martino[1,2], Albert Mukovskiy[3], Martin A. Giese[3], Beatrice de Gelder [1*]

[1] Department of Cognitive Neuroscience, Faculty of Psychology and Neuroscience, Maastricht University, Limburg 6200 MD, Maastricht, The Netherlands

[2] Center for Magnetic Resonance Research, Department of Radiology, University of Minnesota, Minneapolis, MN 55455, United States and Department of NeuroInformatics

[3] Hertie Institute for Clinical Brain Research and Center for Integrative Neuroscience, University Clinic Tübingen, 72074 Tübingen, Germany

* Corresponding author

E-mail: b.degelder@maastrichtuniversity.nl

Abstract

Understanding how the human brain processes body movements is essential for clarifying the mechanisms underlying social cognition and interaction. This study investigates the encoding of biomechanically possible and impossible body movements in occipitotemporal cortex using ultra-high field 7Tesla fMRI. By predicting the response of single voxels to impossible/possible movements using a computational modelling approach, our findings demonstrate that a combination of postural, biomechanical, and categorical features significantly predicts neural responses in the ventral visual cortex, particularly within the extrastriate body area (EBA), underscoring the brain's sensitivity to biomechanical plausibility. Lastly, these findings highlight the functional heterogeneity of EBA, with specific regions (middle/superior occipital gyri) focusing on detailed biomechanical features and anterior regions (lateral occipital sulcus and inferior temporal gyrus) integrating more abstract, categorical information.

Introduction

Human bodies convey essential information about others' actions, intentions, and emotions and provide critical cues in social communication ([1-4]). previous research using functional magnetic resonance imaging to investigate the neural basis of body perception (fMRI) has primarily focused on localizing high level visual category-specific representations. Specific regions in the occipitotemporal and fusiform cortex are selectively responsive to images of bodies, the extrastriate body area (EBA) and the fusiform body area (FBA) ([5], [6]). Similar findings of distinct body sensitive patches were found in monkeys in the ventral bank superior temporal sulcus (STS), namely the middle STS body patch (MSB) and the anterior STS body patch (ASB), with a putative homology between MSB and EBA, and ASB and FBA ([7]). When dynamic images or functional aspects of body perception like action and

emotional expression are also considered, body sensitivity was reported in other areas (8). This has raised interest in investigating the neural mechanisms underlying body sensitivity, notably in the specific computational mechanisms operating across these different body sensitive areas.

Some studies suggested that EBA is more involved in processing body parts and local features and FBA devoted to holistic processing (9, 10). There is also some evidence that EBA and FBA might process a combination of local and global body features (11-14), depending on semantic attributes such as emotion and action (15-17), and that EBA is sensitive to task demands (18). Additionally, recent findings further suggest that activity in the Default Mode Network (DMN) is sensitive to the contrast between biological and non-biological motion based on the naturalness of kinematic patterns. Specifically, the DMN's stronger response to human-like motion, particularly when it matches expected kinematics, suggests that it may modulate or support EBA and FBA processing by enhancing sensitivity to motion patterns that carry social and biological relevance (19). However, despite these insights, there is no clear understanding of a functional division of labour between different body-sensitive areas remain inconclusive. A better understanding of the computational processes within these body-selective areas should clarify their specific contributions to body perception.

Over the past decade, (linearized) encoding (20, 21) has been used to compare different computational hypotheses of brain function. In these approaches, brain activity (e.g., blood oxygen level-dependent (BOLD) signals in a voxel or brain region during fMRI) is predicted based on stimulus features derived from computational models. The accuracy of these predictions can then be compared to adjudicate between competing models, or to determine the relative contribution (variance explained) of each model (22-28). Encoding models predict neural responses based on specific stimulus features and have been successfully

applied to visual processing, in early visual cortex ([20], [21]) as well as higher visual cortex ([13], [24], [29], [30]). An earlier study using still images did apply encoding models to human body-selective regions ([13]) and shed light on the relevance of joint positions and their spatial configuration for the response in the EBA. Like most prior research in the field, the use of still images, only addressed postural aspects rather than movement, thus limiting our understanding of how the brain processes more complex, dynamic information. We approached the issue of dynamic feature computation in EBA by using biomechanical manipulations of natural movements based on 3D. Creating videos that disrupt the natural spatial configuration of joints allowed us to investigate how EBA processes biomechanical plausibility. This approach is particularly important with moving bodies, as dynamic stimuli capture the temporal and kinematic properties essential for understanding how the brain encodes real-world, biologically relevant movements.

We specifically tested the hypothesis that EBA is sensitive to biomechanical characteristics of body movements, building on some earlier indications in the literature. For instance, participants exhibit automatic imitation effects even for impossible movements, indicating the brain's predisposition to process action dynamics despite biomechanical violations ([31]). Recognition of human bodies is significantly affected by inversion, reflecting specialized perceptual mechanisms for recognizing human form in upright configurations ([32]). More recently, prior knowledge of biomechanical constraints has been shown to bias visual memory, with participants misremembering extreme postures as less extreme, adjusting their perceptions toward more biomechanically plausible positions ([33]). Developmental evidence also points to an early sensitivity to biomechanical constraints on human movement: both 12-month-old infants and adults spend more time looking at the elbows during impossible arm movements compared to possible ones ([34]), and newborns can differentiate between biomechanically possible and impossible hand movements ([35]). Investigating the neural

correlates of humanly impossible movements has further revealed that impossible finger movements elicit distinct neural responses compared to possible ones in EBA (36). The influence of biomechanics to the processing of visual information related to the body may be fundamental to how body representations are formed in the brain, and may involve areas like the EBA.

To investigate the computations underlying the neural responses to body movements in the occipitotemporal cortex, we utilized ultra-high-field 7 Tesla fMRI and linearized encoding models, assessing macroscopic and mesoscopic (layer-specific) responses related to biomechanical sensitivity. We aimed to identify how different cortical layers within the EBA encode biomechanical information and distinguish between possible and impossible movements. We employed three distinct encoding models to probe these computations: the 3D Keypoints (kp3d) model, which represents three-dimensional coordinates of body joints and captures precise postural information; the Similarity Distances (simdist) model, which quantifies biomechanical differences between possible (natural) and morphed (impossible) movements based on motion capture data (37); and the categorical differences model, which provides a higher-level distinction by categorizing movements as biomechanically possible or impossible. By comparing model performance across cortical layers, we aimed to test the hypothesis that superficial cortical layers encode categorical information—indicating sensitivity to global, higher-order features—while deeper layers encode joint-specific and biomechanical information (contained in the kp3d and simdist models).

Results

Behavioural analysis confirms consistent categorization of possible and impossible stimuli

The analysis of the questionnaire responses showed that all stimuli were accurately

categorized. In the "possible" condition, every stimulus received the highest rating,

confirming correct classification. Results for the "impossible" videos showed more variability

while consistently scoring below 4 on the 1-7 Likert scale. Notably, 95% (57 out of 60) of

these stimuli had a median rating between 1 and 2, with the remaining three videos rated

between 2 and 3 (see supplementary material for more information).

Localizer stimuli reveal activation in ventral visual cortex and EBA for voxel selection

In each subject, voxels that significantly responded to the localizer conditions (Body +

Objects) with a false discovery rate (FDR) of less than 0.05 were selected for the encoding

analysis. While selection took place at the individual level, in Figure 2 we report group-level

maps obtained by averaging individual thresholded (q(FDR)<0.05) single-subject maps. All

group maps are projected on group aligned (cortex based aligned - CBA) surface.   The

localizer conditions consistently activated regions in the occipitotemporal cortex, specifically

in the superior, middle, and inferior occipital gyri (SOG/MOG/IOG), fusiform gyrus (FG),

lingual gyrus (LG)

middle temporal gyrus (MTG), inferior temporal sulcus (ITS), lateral occipital sulcus (LOS),

and superior temporal sulcus (STS). These clusters overlap with areas identified in our

previous study (13). By subtracting the responses to object stimuli from the responses to body

stimuli, we defined the extrastriate body area (EBA) in each individual and computed

probabilistic maps of the overlap of EBA across individuals in cortex based aligned space.

The EBA spanned the MOG, MTG, and ITS (Fig. 2) with the probabilistic maps showing an

overlap between 20 (white in the Fig.2) and 100% (Green) of subjects.

Joint model significantly predicts responses to novel stimuli in ventral visual cortex

The main effect of the response in the localizer (objects + bodies) was used to select voxels

for the encoding in the individual subjects' data. In these voxels, the response elicited by

body stimuli in the main experiment, independent of the localizer, was modelled using

banded ridge regression. The group performance of the joint encoding model (kp3d,

categorical, simdist) is shown in Figure 3a. Statistical significance at the group level was

assessed via a permutation test, with correction for multiple comparisons using FDR

(q<0.05). The joint model significantly predicted responses to novel stimuli throughout the

ventral visual cortex (SOG, MOG, IOG, ITG, MTG, FG, LOS)

Spatial differences in model contributions to fMRI responses were assessed using an RGB

map (Figure 3b), where each vertex is color-coded to show the relative contribution of each

model to the joint model's accuracy. The kp3d model (red) and simdist model (blue) showed

varying contributions across regions, with the categorical model (green) also playing a role.

In early visual cortical areas, the response to both possible and impossible bodies was best

captured by a combination of the kp3d and simdist models, as indicated by magenta and

purple hues. The categorical model (green) contributed more to the voxels' response in

ventral occipital regions, either on its own or in combination with the one of other models

(reflected by light-blue or orange colors).

EBA encodes postural, biomechanical and categorical information

Within the EBA, the joint model accounted approximately 10% of the variance of the BOLD

signal (Fig. 4, top panel). When considering the model fit across cortical layers, we did not

observe significant differences in (joint) model fit across layers, despite a trend for the model

fit to increase from inner to superficial layers (Fig. 4 bottom left panels). Although not

statistically significant, the percentage of R^2 explained by each model showed a trend, with

the kp3d model accounting for a larger portion of the variance (approximately 40%)

(permutation test subject wise sign-flipping, 2^N=2048 times with N=11 on the differences

of variance explained: kp3d-simdist, p=0.0698; kp3d-categorical, p=0.083)　in the left

hemisphere　(Fig. 4, bottom right panels). " Moreover, a layer-specific analysis within EBA

revealed that the joint model's performance increased from inner to superficial layers in the

right hemisphere (inner-middle: t(10) = -3.546, p = 0.005; inner-superficial: t(10) = -2.325, p

= 0.042),

An ANOVA testing for the difference in model performance between hemispheres and layers

revealed a significant main effect of models (F=4.408, p=0.014) and a significant interaction

between hemispheres and models (F=3.572, p=0.030).

Discussion

The present study investigated how dynamic body stimuli, specifically biomechanically

possible and impossible movements, are encoded in the occipitotemporal cortex. Specifically,

we compared the predictive performance of encoding models based on 3D keypoints,

similarity distances, and categorical differences (kp3d, simdist, categorical). At the group

level, we observed that a combination of the three models significantly predicted fMRI

BOLD responses in the ventral visual cortex after applying permutation testing and correcting

for multiple comparisons. The variance partitioning across the different models of body

posture in EBA across cortical layers revealed hemispheric differences between models. In

the left hemisphere, the kp3d model appeared to explain a larger portion of the variance

(approximately 40%) compared to the simdist and categorical models (both around 30%),

with this pattern observed consistently across cortical depths (see Fig. 4). In the right

hemisphere, the kp3d model accounted for approximately one-third of the variance, while the

simdist and categorical models showed differing trends across cortical layers. Specifically,

the simdist model tended to explain more variance in the inner layers, whereas the categorical

model appeared to account for more variance in the superficial layers. Although these

differences were not statistically significant.

Our findings reveal that a combination of low and high-level features contribute to the

dynamic perception of body movement in occipitotemporal cortex. In early visual cortical

areas, the kp3d and simdist models alone, or in combination, best predicted neural responses

(red, blue, magenta-purple color patches in Fig. 4b) indicating that postural and

biomechanical features play a significant role in these regions. These results align with the

understanding that early visual areas process low-level features such as orientation, spatial

frequency, and basic shape attributes ([20], [21], [38-40]). As processing advances to higher visual

areas, the categorical model becomes increasingly dominant. This shift aligns with previous

literature showing that higher-order areas integrate lower-level features into more abstract

representations, reflecting a progression toward semantic processing ([29], [41-43]).

Our analysis revealed that a combination of the three encoding models—kp3d, simdist, and

categorical—significantly predicted neural responses, accounting for approximately 10% of

the variance in the BOLD signal in EBA (see Fig. 4). This indicates that in EBA various

types of information, including postural and biomechanical features and categorical

distinctions are combined.

While all models contributed significantly to the response elicited by dynamic bodies in

EBA, the kp3d and simdist models were more prominent (purple-magenta patches - in Fig. 3b

and 4) in the superior part of EBA, covering middle occipital gyrus (MOG) and superior

occipital gyrus (SOG). In contrast, the anterior inferior part of EBA —spanning anterior part

of the inferior temporal gyrus (aITG) and anterior lateral occipital sulcus (aLOS)— tended

towards categorical encoding (cyan-orange-green patches in Fig. 3b and 4) suggesting an integration of postural information with more abstract representations. This may involve linking specific body configurations to semantic concepts such as the type of action being performed or the emotional state conveyed by the body movement (44, 45)

This functional heterogeneity found in EBA aligns with anatomical findings that identify distinct body-selective areas within the occipitotemporal cortex (46). Recent findings by Li et al. (47) using data-driven methods identified four adjacent body-selective nodes within the temporo-occipital cortex further support this notion by identifying. Specifically, the superior subregions' emphasis on kp3d and simdist models may explain their role in detailed sensory processing, as they show stronger connectivity with regions involved in processing fine-grained visual details (47). In contrast, the anterior inferior subregions' reliance on categorical encoding supports their involvement in higher-order interpretation and integration of body-related information, consistent with their broader connectivity profiles (47). Our findings thus reinforce the view of the EBA as functionally heterogeneous, highlighting specialized subregions dedicated to different aspects of body and action perception. (47).

Our results are consistent with previous findings showing that EBA is more functionally and structurally connected to dorsal stream regions compared to other body-related areas, such as FBA and the lateral occipital complex (LOC) (48). This connectivity supports the EBA's role in bridging perceptual and motor functions, particularly in specifying goal-directed postural configurations for motor planning. Notably, the study suggests that EBA's connectivity with parietal regions, such as the superior parietal lobule and postcentral gyrus, may enable it to access somatosensory information, which is essential for planning and executing actions based on body posture.

Our layer-specific analysis within EBA revealed that the joint model's performance increased

from inner to superficial layers in the right hemisphere (inner-middle: $t(10) = -3.546$, $p = 0.005$;

0.005; inner-superficial: $t(10) = -2.325$, $p = 0.042$), which may hint towards a gradient of

sensitivity to postural features in right EBA. Conversely, the joint model performed

uniformly across layers in the left hemisphere. The variance partitioning also hinted at a

potential hemispheric difference, with the kp3d model accounting for a substantial portion of

the variance (approximately 40%) across all cortical depths in the left hemisphere. This trend

may suggest a specialization for encoding detailed three-dimensional postural information.,

which is essential for precise spatial judgments and the accurate interpretation of body

movements ([49], [50]). This left-lateralization aligns with previous findings indicating a

dominance of the left hemisphere in processing detailed aspects of body stimuli ([12], [14]) and

may enhance the ability to recognize and interpret fine-grained body movements, facilitating

action recognition and understanding others' intentions. ([1], [51], [52]).

In the right hemisphere, we observed a varying contributions trend of the simdist and

categorical models across cortical layers—higher simdist influence in inner layers and greater

categorical influence in superficial layers— which may hint at a layer-specific encoding

strategy. Although these differences were not statistically significant, they hint at a

potentially differentiated role of cortical layers in processing biomechanical and categorical

information. This aligns with the notion that deeper cortical layers may handle more input-

driven, sensory information, while superficial layers integrate higher-order, contextual, or

semantic information ([53-57]). Additionally, a similar depth-dependent organization has been

demonstrated in the ventral temporal cortex, where superficial layers predominantly encoded

broader, domain-level distinctions, while deeper layers were more sensitive to specific

category-level information ([58]).

The substantial predictive power of the simdist model from early to high-level visual cortex

underscores the visual system's sensitivity to biomechanical constraints from the initial stages

of processing. This suggests that, beyond simply recognizing body parts, the brain may be

encoding midlevel features (8) that reflect the biomechanical characteristics of human

bodies. Midlevel features, including biomechanical constraints on posture and movement,

could serve as a crucial link between the configurations driven by body joints we previously

identified (see (13)) and more abstract, higher-level representations of the body.

Furthermore, differentiating between possible and impossible movements likely involves

detecting deviations from typical joint configurations and movement patterns, which could

suggest the presence of an internal model of human biomechanics (59, 60). Our results extend

previous findings by demonstrating that encoding models can effectively capture neural

responses to biomechanical plausibility (36). This sensitivity may reflect a mechanism for

detecting errors or anomalies in observed body movement, acting as a filtering mechanism

for upstream processing of actions, which is critical for social cognition (47, 51, 61-63).

The relatively small sample size (n=11), which is common in laminar fMRI studies, may

limit the generalizability of our findings. Replication with larger samples will be necessary to

confirm the observed effects and strengthen the reliability of these results. Moreover, our

scanning parameters focused primarily on occipitotemporal and frontal regions, excluding

areas such as the motor and premotor cortices. These regions are known to play a crucial role

in the recognition of both static and dynamic bodily actions (51, 64), responding to

biomechanically possible and impossible stimuli alike (36) and contributing to the

distinction between actions that can be performed and those that cannot (63). Incorporating

these regions in future studies will help clarify their role in the perception and discrimination

of biomechanical plausibility, offering a more comprehensive view of the neural mechanisms

underlying action recognition. Additionally, our stimulus creation was limited to

manipulations of the elbows and knees to generate impossible movements. While these

manipulations provided a targeted approach, future research could expand to include a

broader range of movements and joint manipulations to evaluate the generality of encoding

mechanisms across different biomechanical contexts. Investigating how the brain processes a

variety of joint configurations could uncover additional insights into neural sensitivity to

biomechanical plausibility. Finally, further exploration of hemispheric differences, along with

the potential influence of attention and task demands on encoding, would enrich our

understanding of the factors shaping these neural processes.

In summary, this study investigated whether occipitotemporal cortex, particularly the first

body sensitive area EBA, encodes biomechanically possible and impossible body

movements. By comparing three encoding models—3D keypoints, similarity distances, and

categorical differences—we found that a combination of these models significantly predicted

neural responses in the ventral visual cortex. In the left hemisphere, 3D keypoints explained a

larger portion of the variance across cortical layers, while in the right hemisphere we saw an

emerging trend for preference for similarity distances in deeper layers, with categorical

differences becoming more prominent in superficial layers. The study underscores the brain's

sensitivity to biomechanical plausibility, with the biomechanical (simdist) model explaining a

significant portion of the variance from early stages of visual processing. Lastly, these

findings highlight the EBA's functional heterogeneity, with superior regions (middle/superior

occipital gyri) focusing on detailed biomechanical features and anterior regions (lateral

occipital sulcus and inferior temporal gyrus) integrating more abstract, categorical

information.

Material and methods

Participants

12 right-handed subjects (5 males, mean age = 27.8 ± 3.8 years) participated in this study. They all had normal (or corrected to normal) vision and reported no history of psychiatric or neurological disorders. All subjects were naïve to the task and the stimuli and received monetary compensation for their participation. Scanning sessions took place at the neuroimaging facility Scannexus at Maastricht University. All experimental procedure conformed to the Declaration of Helsinki and the study was approved by the Ethics Committee of the faculty of Psychology and Neuroscience of Maastricht University. One participant was excluded from the main analysis for excessive head motion across multiple runs.

Main experiment stimuli

The stimulus set consisted of 120 videos of two avatars (1 male). The videos were generated by animating mocap data from the MoVi dataset (37), which includes recordings from 60 female and 30 male actors performing 21 daily actions and sports movements. For this experiment, we animated six specific actions (kicking, pointing, waving, jumping, jumping jacks, and walking sideways) performed by 17 actors (9 males). The movements of these 17 actors were then used to animate the two avatars, ensuring that the presented stimuli maintained diversity in motion while being standardized in appearance. This process resulted in 96 videos depicting natural body movements. Additionally, we modified the joint angles of the limbs to create 96 biomechanically impossible videos. To refine the set for the fMRI experiment, we conducted a behavioral validation, to select stimuli showing the greatest difference between possible and impossible movements. This ultimately reduced the set to 120 videos (60 possible videos created from 17 actors performing 4 actions: kicking, jumping, pointing, waving). More details are provided in the behavioral validation section

333 below. Each video was edited to have a length between 60 and 90 frames, corresponding to 2

334 to 3 seconds at 30 frames per second. Additionally, the avatars in each video were aligned to

335 be centered relative to the fixation cross, ensuring a consistent starting position across all

336 videos. During the experiment, the stimuli spanned a mean width and height of 1.84º x 4.32º

337 of visual angle (Fig. 1a).

Localizer stimuli

340 Stimuli for the localizer experiment consisted of videos depicting two object categories:

341 bodies, objects. Additionally, also a scrambled version of each stimulus was included. (Fig.

342 1b). The size of the stimuli was 3.5 * 7.5 degrees for human bodies and objects. For more

343 details about the localizer stimuli we refer to (65). None of the stimuli from the localizer were

344 used in the main experiment.

Behavioural validation

The stimuli created from the mocap data comprised 96 videos of natural body movements

(possible) and their corresponding modified versions, for a total of 192 stimuli. These

modified versions (impossible) were created by altering the joint angles of the limbs to

produce biomechanically impossible movements. We violated the anatomical constraints of

the elbows and knees, by mirroring those joints orientations for each time point of a

trajectory. Accordingly, we modified the shoulders and wrist joint angles, as well as ankles

and hips, in order to preserve the end-effectors (hands and feet) orientations to be as close as

possible to the original (possible) ones for every time point.

Out of the total 192 videos, we selected 120 (60 possible and their impossible version) for the

fMRI experiment through a process of behavioral validation. This selection was based on

identifying the stimuli that best demonstrated the intended differences between possible and

impossible movements, ensuring the most effective set for the experiment. We asked 136

participants (25 males, mean age = 21.45 ± 2 years) to rate the stimuli using a questionnaire

consisting of two Likert-scale questions and one categorical question. Participants were

presented with half (96) of the total stimuli (192) once. For each participant, the stimuli were

pseudo-randomized (96 stimuli randomly selected for each participant, but evenly distributed

so that each stimulus was rated by approximately the same number of participants: mean

number of responses = 68 ± 2.24). After each presentation, participants were asked to answer

a total of three questions about the plausibility/realism of the body movement, action content

and salience of specific body parts (see Supplementary materials).

MRI acquisition and experimental procedure

Participants viewed the stimuli while lying supine in the scanner. Stimuli were presented on a

screen positioned behind participant's head at the end of the scanner bore (distance

screen/eye = 99 cm) which the participants could see via a mirror attached to the head coil.

The screen had a resolution of 1920x1200 pixels, and its angular size was 16º (horizontal) x

10º (vertical). The experiment was coded in Matlab (v2021b The MathWorks Inc., Natick,

MA, USA) using the Psychophysics Toolbox extensions (66-68).

Each participant underwent two MRI sessions, we collected a total of twelve functional runs

(six runs per session) and one set of anatomical images. Images were acquired in a 7T MR

scanner (Siemens Magnetom) using a 32-channel (NOVA) head coil. Anatomical (T1-

weighted) images were collected using MP2RAGE MP2RAGE: 0.7 mm isotropic, repetition

time (TR) = 5000 ms, echo time (TE) = 2.47 ms, matrix size= 320 x 320, number of slices =

240. The functional dataset (T2*-weighted) covered the occipitotemporal cortex and was

acquired using a Multi-Band accelerated 2D-EPI BOLD sequence, multiband acceleration

factor = 2, voxel size = 0.8 mm isotropic, TR = 2300 ms, TE = 27 ms, number of slices = 58

without gaps; matrix size = 224 x 224; number of volumes = 300, GRAPPA factor =3. In

addition to functional images, phase images were simultaneously acquired along with five

noise volumes appended at the end of each run.

During the main experiment, stimuli were presented on the screen for 2-3 seconds (depending

on the length of each video) with an inter stimulus interval that was pseudo-randomised to be

2, 3 or 4 TRs. Participants were asked to fixate at all times on a white cross at the centre of

the screen (Fig. 1c).

To control for attention, participants were asked to detect a shape change at the fixation cross

(cross to circle) and respond via button press with the index finger of the right hand. Within

each run, 20 stimuli (10 possible and 10 impossible) were presented and repeated 3 times.

Three target trials were added for a total of 63 trials per run. The two sessions were identical

therefore each of the 120 videos was repeated 6 times (3 repetitions x 2 sessions) across the

12 runs. Additionally, three blank trials were added in each run lengthening the baseline

period.

Across sessions, we collected 2 to 3 runs of localizer depending on available scanning time.

Each localizer run contained 10 videos per category presented following a block design. Each

block lasted 25 seconds (10 videos x 1 sec + 1.5 sec intertrial interval) and was followed by a

jittered fixation period of 11 seconds on average. Each category block was repeated 3 times

per run. During the localizer participants performed the same task as in the main experiment.

Preprocessing for the functional images was performed using BrainVoyager software (v22.2,

Brain Innovation B.V., Maastricht, the Netherlands), Matlab (v2021b) and ANTs (69). To

lower thermal noise, we performed NOise reduction with DIstribution Corrected (NORDIC)

using both magnitude and phase images (70). EPI Distortion was corrected using the

Correction based on Opposite Phase Encoding (COPE) plugin in BrainVoyager, where the

amount of distortion is estimated based on volumes acquired with opposite phase-encoding

(PE) with respect to the PE direction of the main experiment volumes (71), after which

subsequent corrections is applied to the functional volumes. Other preprocessing steps

included scan slice time correction using cubic spline, 3D motion correction using

trilinear/sinc interpolation and high-pass filtering (GLM Fourier) cut off 3 cycles per run.

During the 3D motion correction process, all runs were aligned to the first volume of the first

run using the scanner's intersession auto-align function, ensuring consistent spatial alignment

across sessions. Anatomical images were resampled at 0.4mm isotropic resolution using sinc

interpolation. To ensure a correct functional-anatomical and functional-functional alignment,

the first volume of the first run was coregistered to the anatomical data in native space using

boundary based registration (72). Functional images were exported in nifti format for further

processing in ANTs. To reduce non-linear intersession distortions, functional images were

corrected using the antsRegistration command in ANTs using as target image the first volume

of the first run and as moving image the first volume of all the other runs. Volume Time

Courses (VTCs) were created for each run in the normalized space (sinc interpolation). Prior

to the encoding analysis (and following an initial general linear model [GLM] analysis aimed

at identifying regions of interest based on the response to the localizer blocks), we performed

an additional denoising step of the functional time series by regressing out the stimulus onset

(convolved with a canonical hemodynamic response function [HRF]) and the motion

parameters. This step was crucial for minimizing the influence of external confounds, such as

the timing of stimulus presentation and participant head motion, on the neural data. By

removing these factors, we ensured that the model's training focused exclusively on learning

patterns directly associated with the features of the encoding models. However, this

approach, while effective in isolating feature-driven neural responses, can lead to smaller

accuracies as it also removes some of the variance explained by the stimulation paradigm

itself. Despite this trade-off, this method provides a cleaner and more specific evaluation of the encoding models' ability to capture the relevant neural patterns.

Segmentation of white matter (WM) and gray matter (GM) boundary as well as cortical layers estimation was performed using a custom pipeline. First, the UNI image and T1 image obtained from MP2RAGE were exported to nifti. We performed gaussian noise reduction using the DenoiseImage command in ANTs ([73]), and bias field correction in SPM12 as described on layer fMRI blog (https://layerfmri.com/2017/12/21/bias-field-correction/). After preprocessing of anatomical images, Cortical reconstruction and volumetric segmentation was performed using Basic SAMSEG (cross-sectional processing) command of the Freesurfer image analysis suite (http://surfer.nmr.mgh.harvard.edu/), using the UNI images as T1w contrast and the T1 map of the MP2RAGE (which has flipped intensities between white and gray matter, resembling a T2w image) as T2w contrast. Lastly, cortical thickness and layers extraction were performed using surf_laynii.sh script (https://github.com/srikash/surf_laynii/blob/main/surf_laynii) which enables layering in LAYNII ([74]) using the Freesurfer segmentations output. Three layers were then calculated in LAYNII using the equi-volume approach. While all analyses were performed in the individual subject space, for visualization purposes we projected single-subject statistical or encoding maps onto a group cortex-based aligned surface and then averaged the results across subjects ([75]).

Voxels selection for encoding analysis

The functional time series of the localizer runs collected in each participant were analysed using a fixed-effect GLM with 5 predictors (4 conditions in the localizer: Body Objects and

their scrambled version and 1 modelling the catch trials). Motion parameters were included in

the design matrix as nuisance regressors. The estimated regressor coefficients representing

the response to the localizer blocks were used for voxel selection. A voxel was selected for

the encoding analysis if significantly active (q(FDR)<0.05) in response to the Body and

Objects categories. Note that this selection is unbiased to the response to the stimuli

presented in the experimental section of each run.

Functional ROI definition

Using the functional localizer we also defined body selective regions at the single subject

level. Specifically, the EBA was defined using the contrast [Body + Body Scrambled] >

[Objects + Objects Scrambled] ([76]) with a statistical threshold of q(FDR) < 0.05. All

subsequent ROI-level analyses were conducted by identifying the intersection between the

voxels assigned to the EBA and those selected for the encoding analysis.

Encoding models

In order to understand what determines the response to body images we tested several

hypotheses, represented by different computational models, using fMRI encoding ([20], [21], [25],

[77]). We compared the performance (accuracy in predicting left out data) of three encoding

models.

The first represented body stimuli using the position of joints in three dimensions (kp3d)

using 71 keypoints (main skeleton joints like hips, knees, shoulders, elbows, hands and facial

features like eyeballs, neck and jaw) extracted from the MoVi dataset. This   model

represents the stimuli as a collection of points in space forming a human skeleton. To focus

on joints that significantly influence perception while minimizing variability from less

relevant keypoints, we excluded constant (or almost constant) keypoints ending up with a

subset that included 56 keypoints (shoulders, elbows, wrists, hips, knees, and ankles, hands,

fingers and facial features from both sides of the body).

The second model quantifies the similarity distances (simdist) between morphed movements

(impossible) and normal movements (possible) by analyzing motion capture data extracted

during stimulus creation. For each video, both the modified and original motion data were

loaded. Initially, all 71 joints defined in the MoVi skeleton were considered. However, to

focus on joints with meaningful movement and reduce variability from less relevant joints

(such as fingers and toes), joints without rotation data (i.e., joints with empty rotation indices)

were excluded, reducing the original set to 56 keypoints (the same as in the previous

paragraph). For each selected joint at each time frame, we converted the original Euler angles

representing the joint rotation to axis-angle representation. This process yielded a set of three-

dimensional vectors in Euclidian space representing the rotation of each joint over time. To

measure the similarity between test movements (both modified and original) and the manifold

of normal (original) movements, a Gaussian kernel-based approach was employed. This

method quantifies the proximity of motion data in the high-dimensional joint angle space,

allowing for a robust assessment of movement similarity (see supplementary material).

Keypoints for which the computed similarity distances to the normative manifold were not

finite (e.g., containing NaN or Inf values) were identified and excluded to maintain data

quality, reducing the original 56 keypoints to 29. Similarity distances for all joints were then

concatenated to form feature vectors representing each movement's similarity across all

considered joints. This model encoded biomechanical differences because it evaluates the

kinematic properties of human joint movements by measuring their distances to a manifold of

normal actions, thereby allowing for the differentiation between biomechanically plausible

(possible) and implausible (impossible) movements, with the latter exhibiting higher

distances due to their deviation from typical human motion patterns. (for the mathematical

formulation see supplementary materials).

The third model encodes categorical differences between possible and impossible stimuli by

incorporating two features that explicitly indicate the (im)possibility of each stimulus. Unlike

the other models, this approach does not account for variations within each category,

focusing instead on the binary classification of stimuli as either possible or impossible. This

model is considered more abstract (or higher-order) compared to the kp3d and simdist

models, as it goes beyond image computable approaches (like keypoints) and instead

recapitulates a conceptual distinctions.

Banded ridge regression and model estimates'

In the context of fMRI, the linearized encoding framework typically uses L2-regularized

(ridge) regression to extract information from brain activity (78). This method is effective for

improving the performance of models with nearly collinear features and helps minimize

overfitting. When dealing with multiple encoding models, ridge regression can either

estimate parameters for a combined feature space or for each model separately. However,

using a single regularization parameter for all models may not be optimal due to varying

feature space requirements. To address this, banded ridge regression optimizes separate

regularization parameters for each feature space, enhancing model performance by reducing

spurious correlations and ignoring non-predictive features.(23, 24). In the present work we

used banded ridge regression to fit the three encoding models and performed a decomposition

of the variance explained by each of the models following established procedures (13, 23).

Model training and testing were performed in cross-validation (3-folds: training on 8 runs [80

stimuli repeated 6 times] and testing on 4 runs [40 repeated 6 times]). For each fold, the

training data were additionally split in training set and validation set using split-half

crossvalidation. Within the (split-half) training set a combination of random search and

gradient descent (23) was used to optimize the model fit to the data (regularization strength

and model parameters). Ultimately, the best model over the two (split-half) validation folds

was selected to be tested on the independent test data (4 runs). Within each fold, the models'

representations of the training stimuli were normalized (each feature was standardized to zero

mean and unit variance withing the training set). The feature matrices representing the stimuli

were then combined with the information of the stimuli onset during the experimental runs.

This resulted in an experimental design matrix (nrTRs x NrFeatures) in which each stimulus

was described by its representation by each of the models. To account for the hemodynamic

response, we delayed each feature of the experimental design matrix (5 delays spanning 11.5

seconds). The same procedure was applied to the test data, with the only difference that when

standardizing the model matrices, the mean and standard deviation obtained from the training

data were used.

We used banded ridge regression to determine the relationship between the features of the

encoding models (stimulus representations) and the fMRI response at each voxel. The

encoding was limited to voxels that significantly responded to the localizer stimuli

(p(FDR)<0.05) in each individual volunteer's data.

For each cross-validation, we assessed the accuracy of the model in predicting fMRI time

series by computing the correlation between the predicted fMRI response to novel stimuli (4

runs, 40 stimuli) and the actual responses. The accuracy obtained across the three folds were

Z-transformed and then averaged. To obtain the contribution of each of the models to the

overall accuracy we computed the partial correlation between the measured time series and

the prediction obtained when considering each of the models individually (23). Statistical

significance was assessed at the group level via permutation test (subject wise sign-flipping,

2^N=2048 times with N=11), and correction for multiple comparison was performed using

FDR (q<0.05). Additionally, for each subject, we obtained the average coefficient of

determination within EBA (defined in the localizer) and we tested significance with a t-test

against zero.

Acknowledgements

562 This work was supported by the European Research Council (ERC) Synergy grant (Grant

563 agreement 856495; Relevance), by the Horizon 2020 Programme H2020-FETPROACT430

564 2020-2 (grant 101017884 GuestXR).

Data availability statement

567 Data and code are being prepared to be made publicly available.

CRediT authorship contribution statement

Giuseppe Marrazzo: Conceptualization, Investigation, Software, Formal analysis,

Validation, Visualization, Writing – Original Draft Preparation, Writing - review & editing.

Federico De Martino: Conceptualization, Investigation, Supervision, Validation, Writing -

review & editing. **Albert Mukovskiy:** Software, Validation, Writing - review & editing.

Martin A. Giese: Conceptualization, review & editing, Funding Acquisition. **Beatrice de**

Gelder: Conceptualization, Project Administration, Resources, Supervision, Funding

Acquisition, Writing – Original Draft Preparation, Writing - review & editing

References

1.   de Gelder B, Van den Stock J, Meeren HKM, Sinke CBA, Kret ME, Tamietto M.

Standing up for the body. Recent progress in uncovering the networks involved in the

perception of bodies and bodily expressions. Neuroscience & Biobehavioral Reviews.

2010;34(4):513-27.

2.   Peelen MV, Downing PE. The neural basis of visual body perception. Nature Reviews

Neuroscience. 2007;8(8):636-48.

3.   de Gelder B. Towards the neurobiology of emotional body language. Nature Reviews

Neuroscience. 2006;7(3):242-9.

4.   Tipper CM, Signorini G, Grafton ST. Body language in the brain: constructing

meaning from expressive movement. Frontiers in Human Neuroscience. 2015;9.

5.   Downing PE, Jiang Y, Shuman M, Kanwisher N. A cortical area selective for visual

processing of the human body. Science. 2001;293(5539):2470-3.

6.   Peelen MV, Downing PE. Selectivity for the human body in the fusiform gyrus.

Journal of Neurophysiology. 2005;93(1):603-8.

7.   Vogels R. More Than the Face: Representations of Bodies in the Inferior Temporal

Cortex. Annual Review of Vision Science. 2022;8(1):383-405.

8.   de Gelder B, Poyo Solanas M. A computational neuroethology perspective on body

and expression perception. Trends in Cognitive Sciences. 2021;25(9):744-56.

9.   Taylor JC, Downing PE. Division of labor between lateral and ventral extrastriate

representations of faces, bodies, and objects. J Cogn Neurosci. 2011;23(12):4122-37.

10.   Taylor JC, Wiggett AJ, Downing PE. Functional MRI analysis of body and body part

representations in the extrastriate and fusiform body areas. J Neurophysiol. 2007;98(3):1626-

33.

11.   Downing PE, Peelen MV. The role of occipitotemporal body-selective regions in

person perception. Cogn Neurosci. 2011;2(3-4):186-203.

12.   Downing PE, Peelen MV. Body selectivity in occipitotemporal cortex: Causal

evidence. Neuropsychologia. 2016;83:138-48.

13.   Marrazzo G, De Martino F, Lage-Castellanos A, Vaessen MJ, de Gelder B.

Voxelwise encoding models of body stimuli reveal a representational gradient from low-level

visual features to postural features in occipitotemporal cortex. Neuroimage.

2023;277:120240.

14.   Bracci S, Ietswaart M, Peelen MV, Cavina-Pratesi C. Dissociable neural responses to

hands and non-hand body parts in human left extrastriate visual cortex. J Neurophysiol.

2010;103(6):3389-97.

15.   de Gelder B, Snyder J, Greve D, Gerard G, Hadjikhani N. Fear fosters flight: a

mechanism for fear contagion when perceiving emotion expressed by a whole body. Proc

Natl Acad Sci U S A. 2004;101(47):16701-6.

16.   Hadjikhani N, de Gelder B. Seeing fearful body expressions activates the fusiform

cortex and amygdala. Curr Biol. 2003;13(24):2201-5.

17.   Downing PE, Peelen MV, Wiggett AJ, Tew BD. The role of the extrastriate body area

in action perception. Soc Neurosci. 2006;1(1):52-62.

18.   Marrazzo G, Vaessen MJ, de Gelder B. Decoding the difference between explicit and

implicit body expression representation in high level visual, prefrontal and inferior parietal

cortex. NeuroImage. 2021;243:118545.

19.   Dayan E, Sella I, Mukovskiy A, Douek Y, Giese MA, Malach R, et al. The Default

Mode Network Differentiates Biological From Non-Biological Motion. Cerebral Cortex.

2016;26(1):234-45.

20.   Naselaris T, Kay KN, Nishimoto S, Gallant JL. Encoding and decoding in fMRI.

Neuroimage. 2011;56(2):400-10.

21.   Kay KN, Naselaris T, Prenger RJ, Gallant JL. Identifying natural images from human

brain activity. Nature. 2008;452(7185):352-5.

22.   Dumoulin SO, Wandell BA. Population receptive field estimates in human visual

cortex. Neuroimage. 2008;39(2):647-60.

23.   Dupré la Tour T, Eickenberg M, Gallant JL. Feature-space selection with banded

ridge regression. Neuroimage. 2022;264:119728.

24.   Nunez-Elizalde AO, Huth AG, Gallant JL. Voxelwise encoding models with non-

spherical multivariate normal priors. Neuroimage. 2019;197:482-92.

25.   Santoro R, Moerel M, De Martino F, Goebel R, Ugurbil K, Yacoub E, et al. Encoding

of Natural Sounds at Multiple Spectral and Temporal Resolutions in the Human Auditory

Cortex. PLOS Computational Biology. 2014;10(1):e1003412.

26.   Thirion B, Duchesnay E, Hubbard E, Dubois J, Poline JB, Lebihan D, et al. Inverse

retinotopy: inferring the visual content of images from brain activation patterns. Neuroimage.

2006;33(4):1104-16.

27.   Wandell BA, Dumoulin SO, Brewer AA. Visual field maps in human cortex. Neuron.

2007;56(2):366-83.

28.   Moerel M, De Martino F, Formisano E. Processing of natural sounds in human

auditory cortex: tonotopy, spectral tuning, and relation to voice sensitivity. The Journal of

neuroscience : the official journal of the Society for Neuroscience. 2012;32(41):14205-16.

29.   Huth AG, Nishimoto S, Vu AT, Gallant JL. A continuous semantic space describes

the representation of thousands of object and action categories across the human brain.

Neuron. 2012;76(6):1210-24.

30.   Yamins DLK, Hong H, Cadieu CF, Solomon EA, Seibert D, DiCarlo JJ. Performance-

optimized hierarchical models predict neural responses in higher visual cortex. Proceedings

of the National Academy of Sciences. 2014;111(23):8619-24.

31.   Longo MR, Kosobud A, Bertenthal BI. Automatic Imitation of Biomechanically

Possible and Impossible Actions: Effects of Priming Movements Versus Goals. Journal of

Experimental Psychology: Human Perception and Performance. 2008;34(2):489-501.

32.   Reed CL, Stone VE, Bozova S, Tanaka J. The Body-Inversion Effect. Psychological

Science. 2003;14(4):302-8.

33.   Han Q, Gandolfo M, Peelen MV. Prior knowledge biases the visual memory of body

postures. iScience. 2024;27(4):109475.

34.   Morita T, Slaughter V, Katayama N, Kitazaki M, Kakigi R, Itakura S. Infant and adult

perceptions of possible and impossible body movements: an eye-tracking study. J Exp Child

Psychol. 2012;113(3):401-14.

35.   Longhi E, Senna I, Bolognini N, Bulf H, Tagliabue P, Cassia VM, et al.

Discrimination of biomechanically possible and impossible hand movements at birth. Child

Dev. 2015;86(2):632-41.

36.   Costantini M, Galati G, Ferretti A, Caulo M, Tartaro A, Romani GL, et al. Neural

Systems Underlying Observation of Humanly Impossible Movements: An fMRI Study.

Cerebral Cortex. 2005;15(11):1761-7.

37.   Ghorbani S, Mahdaviani K, Thaler A, Kording K, Cook DJ, Blohm G, et al. MoVi: A

large multi-purpose human motion and video dataset. PLoS One. 2021;16(6):e0253157.

38.   Carandini M, Demb JB, Mante V, Tolhurst DJ, Dan Y, Olshausen BA, et al. Do we

know what the early visual system does? J Neurosci. 2005;25(46):10577-97.

39.   Nishimoto S, Gallant JL. A Three-Dimensional Spatiotemporal Receptive Field

Model Explains Responses of Area MT Neurons to Naturalistic Movies. The Journal of

Neuroscience. 2011;31(41):14551-64.

40.   Nishimoto S, Vu AT, Naselaris T, Benjamini Y, Yu B, Gallant JL. Reconstructing

visual experiences from brain activity evoked by natural movies. Curr Biol.

2011;21(19):1641-6.

41.   Kriegeskorte N, Mur M, Bandettini P. Representational similarity analysis -

connecting the branches of systems neuroscience. Front Syst Neurosci. 2008;2:4.

42.   Grill-Spector K, Weiner KS. The functional architecture of the ventral temporal

cortex and its role in categorization. Nature Reviews Neuroscience. 2014;15(8):536-48.

43.   Haxby JV, Gobbini MI, Furey ML, Ishai A, Schouten JL, Pietrini P. Distributed and

overlapping representations of faces and objects in ventral temporal cortex. Science.

2001;293(5539):2425-30.

44.   Foster C, Zhao M, Bolkart T, Black MJ, Bartels A, Bülthoff I. Separated and

overlapping neural coding of face and body identity. Hum Brain Mapp. 2021;42(13):4242-60.

45.   Foster C, Zhao M, Romero J, Black MJ, Mohler BJ, Bartels A, et al. Decoding

subcategories of human bodies from both body- and face-responsive cortical regions.

NeuroImage. 2019;202:116085.

46.   Weiner KS, Grill-Spector K. Not one extrastriate body area: using anatomical

landmarks, hMT+, and visual field maps to parcellate limb-selective activations in human

lateral occipitotemporal cortex. Neuroimage. 2011;56(4):2183-99.

47.   Li B, Poyo Solanas M, Marrazzo G, de Gelder B. Connectivity and functional

diversity of different temporo-occipital nodes for action perception. bioRxiv.

2024:2024.01.12.574860.

48.   Zimmermann M, Mars RB, de Lange FP, Toni I, Verhagen L. Is the extrastriate body

area part of the dorsal visuomotor stream? Brain Structure and Function. 2018;223(1):31-46.

49.   Caspari N, Popivanov ID, De Mazière PA, Vanduffel W, Vogels R, Orban GA, et al.

Fine-grained stimulus representations in body selective areas of human occipito-temporal

cortex. NeuroImage. 2014;102:484-97.

50.   Kumar S, Popivanov ID, Vogels R. Transformation of Visual Representations Across

Ventral Stream Body-selective Patches. Cerebral Cortex. 2019;29(1):215-29.

51.   Urgesi C, Candidi M, Ionta S, Aglioti SM. Representation of body identity and body

actions in extrastriate body area and ventral premotor cortex. Nat Neurosci. 2007;10(1):30-1.

52.   Blake R, Shiffrar M. Perception of human motion. Annu Rev Psychol. 2007;58:47-73.

53.   Larkum M. A cellular mechanism for cortical associations: an organizing principle for

the cerebral cortex. Trends in Neurosciences. 2013;36(3):141-51.

54.   Spratling MW. A review of predictive coding algorithms. Brain Cogn. 2017;112:92-7.

55.   Bastos AM, Usrey WM, Adams RA, Mangun GR, Fries P, Friston KJ. Canonical

microcircuits for predictive coding. Neuron. 2012;76(4):695-711.

56.   Felleman DJ, Van Essen DC. Distributed hierarchical processing in the primate

cerebral cortex. Cereb Cortex. 1991;1(1):1-47.

57.   Rockland KS, Pandya DN. Laminar origins and terminations of cortical connections

of the occipital lobe in the rhesus monkey. Brain Res. 1979;179(1):3-20.

58.   Margalit E, Jamison KW, Weiner KS, Vizioli L, Zhang R, Kay KN, et al. Ultra-high-

resolution fMRI of Human Ventral Temporal Cortex Reveals Differential Representation of

Categories and Domains. The Journal of Neuroscience. 2020;40(15):3008-24.

59.   Dayan P, Berridge KC. Model-based and model-free Pavlovian reward learning:

revaluation, revision, and revelation. Cogn Affect Behav Neurosci. 2014;14(2):473-92.

60.   Giese MA, Poggio T. Neural mechanisms for the recognition of biological

movements. Nature Reviews Neuroscience. 2003;4(3):179-92.

61.   Kilner JM, Friston KJ, Frith CD. Predictive coding: an account of the mirror neuron

system. Cogn Process. 2007;8(3):159-66.

62.   Schubotz RI. Prediction of external events with our motor system: towards a new

framework. Trends Cogn Sci. 2007;11(5):211-8.

63.   Candidi M, Urgesi C, Ionta S, Aglioti SM. Virtual lesion of ventral premotor cortex

impairs visual perception of biomechanically possible but not impossible actions. Social

Neuroscience. 2008;3(3-4):388-400.

64.   Pobric G, de C. Hamilton AF. Action Understanding Requires the Left Inferior

Frontal Cortex. Current Biology. 2006;16(5):524-9.

65.   Li B, Solanas MP, Marrazzo G, Raman R, Taubert N, Giese M, et al. A large-scale

brain network of species-specific dynamic human body perception. Prog Neurobiol.

2023;221:102398.

66.   Brainard DH. The Psychophysics Toolbox. Spat Vis. 1997;10(4):433-6.

67.   Kleiner M, Brainard DH, Pelli D. What's new in Psychtoolbox-3? 2007.

68.   Pelli DG. The VideoToolbox software for visual psychophysics: Transforming

numbers into movies. Spatial vision. 1997;10:437-42.

69.   Avants BB, Tustison N, Song G. Advanced normalization tools (ANTS). Insight j.

2009;2(365):1-35.

70.   Moeller S, Pisharady PK, Ramanna S, Lenglet C, Wu X, Dowdle L, et al. NOise

reduction with DIstribution Corrected (NORDIC) PCA in dMRI with complex-valued

parameter-free locally low-rank processing. NeuroImage. 2021;226:117539.

71.   Fritz L, Mulders J, Breman H, Peters J, Bastiani M, Roebroeck A, et al., editors.

Comparison of EPI distortion correction methods at 3T and 7T. Annual Meeting of the

Organization for Human Brain Mapping; 2014.

72. Greve DN, Fischl B. Accurate and robust brain image alignment using boundary-

based registration. Neuroimage. 2009;48(1):63-72.

73. Manjón JV, Coupé P, Martí-Bonmatí L, Collins DL, Robles M. Adaptive non-local

means denoising of MR images with spatially varying noise levels. J Magn Reson Imaging.

2010;31(1):192-203.

74. Huber L, Poser BA, Bandettini PA, Arora K, Wagstyl K, Cho S, et al. LayNii: A

software suite for layer-fMRI. NeuroImage. 2021;237:118091.

75. Goebel R, Esposito F, Formisano E. Analysis of functional image analysis contest

(FIAC) data with brainvoyager QX: From single□subject to cortically aligned group general

linear model analysis and self□organizing group independent component analysis. Human

brain mapping. 2006;27(5):392-401.

76. Ross P, de Gelder B, Crabbe F, Grosbras MH. A dynamic body-selective area

localizer for use in fMRI. MethodsX. 2020;7:100801.

77. Allen EJ, Moerel M, Lage-Castellanos A, De Martino F, Formisano E, Oxenham AJ.

Encoding of natural timbre dimensions in human auditory cortex. NeuroImage. 2018;166:60-

70.

78. Hoerl AE, Kennard RW. Ridge Regression: Biased Estimation for Nonorthogonal

Problems. Technometrics. 1970;12(1):55-67.

Figure 1. Stimuli and experimental procedure

(a) The videos were generated by animating mocap data from the MoVi dataset ([37]). Sixty possible videos were created from 17 actors performing 4 actions: kicking, jumping, pointing, waving. Additionally, we modified the joint angles of the elbows and knees to create 60 biomechanically impossible videos. In panel (a) we show frame of possible videos and their equivalent impossible. **(b)** For each run 1/6 of the stimuli (20) where presented in a pseudo-randomized order following a fast event-related design. Each stimulus was repeated three times per run. Each run was repeated two times across sessions resulting in a total of 120 stimuli repeated six times. To identify body sensitive region, the localizer stimuli included videos of humans performing natural body movement, objects, and their scrambled version. We presented stimuli following a block-design with each block repeated three times per run. **(c)** During the main experiment participants fixated on the cross and were presented with the stimuli depicting possible and impossible body movement for 1-2 TRs (depending on the length of each video) followed by a blank screen which appeared for 2, 3 or 4 TRs. When the fixation cross turned to a circle, they had to press a button whether with the right index finger. TR= 2300ms.

Figure 2 Voxels selection and EBA definition

Voxels that were significantly (q(FDR)<0.05) responding to localizer stimuli [Body + Objects]>0 were selected for the encoding analysis. Although the analysis was performed at single-subject level, for visualization purposes we show the average t-map (in red-yellow) obtained by averaging the thresholded single-subjects maps projected on a group cortex-based aligned mesh.

EBA was defined within the localizer via the contrast [Body + Body Scramble] > [Objects + Objects Scramble]. Shown in white-green is a probabilistic map indicating the overlap between individually defined EBAs (q(FDR)<0.05).

Figure 3. Group-level encoding results.

(a) Group Prediction accuracy for the joint model (kp3d, categorical, simdist). Statistical significance was assessed via permutation test (subject wise sign-flipping, $2^N=2048$ times with N=11), and correction for multiple comparison was performed using FDR (q<0.05).

(b) RGB map in which each vertex is colour coded according to the relative contribution of each model to the accuracy of the joint model (red = 100% kp3d; blue = 100% simdist; green = 100% categorical) as shown in (a). For clarity, we overlay the outline of EBA as defined in the probabilistic map depicted in Figure 2 by selecting vertices shared by at least 40% of the subjects.

Figure 4. Joint model performance and variance partitioning in EBA across cortical depths.

Variance partitioning in the extrastriate body area (EBA) across 11 subjects, comparing left (LH) and right hemispheres (RH) across three cortical layers (Left to right → inner, middle, superficial). The top panel shows the group average R² values in the EBA, indicating overall joint model performance across hemispheres.

The bottom left panels display the variance explained (R²) in the LH and RH EBA across layers. The bottom right panel illustrates the percentage of R² explained by each model across layers. To check for differences in variance explained between models, we ran an ANOVA which showed a significant main effect of models ($F_{(2,180)}$ = 4.408, p=0.014) and a significant interaction between hemispheres and models ($F_{(2,180)}$ = 3.572, p=0.030) were found, indicating that models performance varies between hemispheres and that the effectiveness of each model differs across layers. Error bars represent standard errors.