Denitrification is a community trait with partial pathways dominating across

microbial genomes and biomes

Grace Pold [1], Aurélien Saghaï [2], Christopher M Jones [2], and Sara Hallin [2*]

[1] Department of Soil and Environment, Swedish University of Agricultural Sciences,

Uppsala

[2] Department of Forest Mycology and Plant Pathology, Swedish University of

Agricultural Sciences, Uppsala

* Corresponding author: sara.hallin@slu.se

Running title: Global ecology of denitrification trait

Keywords: comparative genomics; comparative metagenomics; community ecology;

division of labor; global biomes; nitrous oxide; split pathways

Abstract

Diverse microorganisms can execute one or more steps in denitrification, during which nitrate

or nitrite is successively reduced into nitric oxide, nitrous oxide, and ultimately dinitrogen.

Many of the best-characterized denitrifiers are "complete" denitrifiers capable of executing

all steps in the pathway, but whether they dominate in natural communities and what

metabolic traits and environmental factors drive the global distribution of complete vs. partial

denitrifiers remains to be determined. To address this, we conducted a comparative analysis

of denitrification genes in 61,293 genomes, 3,991 metagenomes covering all major biomes,

and 413 terrestrial and aquatic metatranscriptomes. We show that partial denitrifiers

outnumber complete denitrifiers and the potential to initiate denitrification is more common

than the potential to terminate it, both among genomes and at the community level across

most biomes, particularly in nutrient rich environments. These patterns were also reflected in

the metatranscriptomes. Our results further indicate that complete denitrifiers are more likely

to be fast-growing organisms, favoring organic acid over sugar metabolism, and encoding the

ability to oxidize and reduce a broader range of organic and inorganic compounds compared

to partial denitrifiers. This suggests complete denitrifiers are metabolically flexible

opportunists. Together, our results indicate an environmental footprint on the presence of

denitrification genes which favors the genomic potential for partial over complete

denitrification in most biomes and highlight that completion of the denitrification pathway is

a community effort.

Main

The potent greenhouse gas and ozone depleting agent nitrous oxide (N_2O) is produced in

several microbial N-cycle processes, of which denitrification plays a dual role in both

producing and consuming N_2O. In this facultative anaerobic microbial respiratory pathway,

nitrate (NO_3^-) or nitrite (NO_2^-) are used as electron acceptors and successively reduced into

nitric oxide (NO), N_2O, and dinitrogen (N_2) by a diverse range of predominantly bacterial

species. As the genes encoding the enzymes involved in these reductive steps can be

independently gained and lost [1-3], they are found in different combinations among genomes [4],

such that denitrification can be executed by complete denitrifiers with genes encoding

enzymes for all steps or, alternatively, split between multiple partial denitrifier community

members. Recent studies suggest that genetically complete denitrifiers are scarce among

genomes assembled from environmental data [5-10]. However, much of our understanding of

the physiology and regulation of denitrification is based on studies of a very phylogenetically

restricted set of complete denitrifiers [11-16]. Thus, there is pressing need for a broader look at

the prevalence, diversity and ecology of organisms genetically capable of complete and

partial denitrification, particularly considering their differential roles in serving as sources

and sinks of N_2O [17].

Split pathways divide the protein production costs for a pathway between multiple cells and

may enable higher ATP flux [18,19], increasing the specific growth rate. Although this could be a

fitness advantage under nitrite-rich conditions, ATP yields per molecule nitrate or nitrite

would be lower for partial compared to complete denitrification. This is expected to lead to

partial denitrification being favored when electron acceptor supply is not limiting, but

complete denitrification being favored when the flux of electron donors such as carbon (C)

compounds is high relative to electron acceptor availability. Splitting the denitrification

pathway also increases dependencies between community members, not only because the

denitrification product of one organism serves as the electron acceptor for another [20,21], but

also because the regulation of specific reactions may rely on intermediates that those

executing the reaction themselves cannot use [22]. Further, there is less buildup of toxic

intermediate compounds, which may allow microorganisms only capable of carrying out the

initial steps of denitrification avoid the cytotoxic effects of NO_2^-, particularly under low pH

environments [11,23]. This jointly indicates that split pathways may open more niches, thereby

supporting a greater diversity of denitrifiers [21]. If organisms encoding partial vs. complete

denitrification fulfill different niches, this should also be reflected in additional traits, such as

growth rate, abiotic optima, and the ability to metabolize different substrates and use electron

acceptors beyond those in the denitrification pathway. For instance, if complete

denitrification is indeed associated with C influx, complete denitrifiers would be expected to

be capable of higher growth and broader substrate catabolism compared to partial denitrifiers.

Addressing the linkages between the complete or partial denitrification and the capacity for

other traits can help identify the broader roles and conditions that favor each denitrifier type.

Here we combined comparative genomics (**Figure 1a**) with a global survey of

denitrification gene ratios in environmental metagenomes and metatranscriptomes to

evaluate the diversity of bacterial denitrifier types, their metabolic traits, and their

distribution across engineered, environmental and host-associated biomes. We posited

that environmental factors favoring or disfavoring execution of the different steps leave

a genetic footprint in the form of presence and absence of genes encoding the enzymes

responsible for them, as well as via differential associations between denitrification

enzyme genes and redox traits. Furthermore, considering organisms encoding a

complete denitrification pathway are able to use a broader range of nitrogenous

terminal electron acceptors than those encoding fewer, we hypothesized that the former

would have broader metabolic flexibility. We used a framework in which partial

denitrifiers include genetically defined "initiators," which encode genes involved in

NO_2^- but not N_2O reduction, and "terminators," which encode genes for N_2O but not

NO_2^- reduction (**Figure 1b**). We used NO_2^- reduction, the decisive step of

denitrification[24], as the first point of our analysis. We combined comparative genomics

of 61,293 non-redundant bacterial isolate, metagenome assembled (MAGs), and single

cell genomes with 3,991 metagenomes derived from all major biomes and 413

terrestrial and aquatic metatranscriptomes to determine the frequency of a complete vs.

partial denitrification pathway and the genetic potential to initiate vs. terminate

denitrification at the genome and community level. We interpret our results in the

context of resource availability and complementary genomic traits associated with the

denitrifier types in our framework.

Results

Partial denitrification dominates microbial genomes

We assessed the prevalence of partial and complete denitrification pathways among

61,293 bacterial genomes represented in GTDB release 214.1 (**Fig. 1a**). Genes for the

denitrification enzymes NirK (*nirK*) and NirS (*nirS*) catalyzing NO_2^- reduction to NO,

all proposed NO-reducing heme copper oxidases (i.e. Nor (*nor*) [25]), and NosZ (*nos*ZI,

*nos*ZII) catalyzing N_2O reduction were annotated using hidden Markov models trained

on manually curated databases of these genes [25]. Genomes carrying *nor* but not *nir* or

nosZ accounted for 5% of the total number of assemblies and were excluded in

downstream analyses because Nor in *nor*-only genomes are likely involved in

detoxification rather than respiration. With this conservative approach, 18% of the

genomes (n=11,126) were considered denitrifiers and included *nir* and/or *nosZ* with or

without *nor*. Among these, 23% encoded genes for complete NO_2^- reduction to N_2

("complete denitrifiers"; **Fig. 1b**), with the complete denitrification trait more prevalent

among isolates compared to MAGs (29% vs. 17%, respectively, **Supplementary**

Table 1). However, there was considerable variation between phyla (**Fig. 1c,**

Supplementary Fig. 1). Twenty-eight different bacterial phyla encoded the complete

denitrification trait, with an overwhelming dominance of Pseudomonadota in our

dataset (**Fig. 1c**). Among phyla represented by at least 100 denitrifier genomes,

complete denitrifiers were most prevalent within Campylobacterota (38% of

denitrifiers), Pseudomonadota (37%), and Myxococcota (26%, **Fig. 1d**).

Using our framework of initiators and terminators, we found that initiators encoding

nir with or without *nor* but not *nosZ* were broadly more prevalent than terminators

having *nosZ* with or without *nor* but not *nir* (49% vs. 24% of genomes with

denitrification genes, respectively). This may represent the potentially divergent

functions for NO_2^- versus N_2O reduction as a means of growth or detoxification vs.

maintenance [26]. Partial denitrifiers occurred in 68 bacterial phyla, with initiators being

overrepresented within Actinomycetota (98% initiators, 1% terminators) and

Nitrospirota (88%, 8%) (**Fig. 1d**). However, Nitrospirota accounted for just a small

fraction of initiator genomes (2.7%), while the majority were Pseudomonadota (47%)

and Actinomycetota (22%) (**Fig 1c**). We found a higher proportion of terminators than

initiators within Planctomycetota (76% terminators, 19% initiators) and Bacteroidota

(60%, 18%) (**Fig. 1d**), with the latter phylum accounting for the majority of

terminators in the dataset (53%; **Fig. 1c**). We also found genomes carrying *nir* and

nosZ but not *nor,* suggesting a third category: the "initiator-terminators". They

accounted for just 4% of genomes with denitrification genes, including 13% of

Gemmatimonadota, 13% of Chloroflexota, 9% of Bacteroidota and 5% of

Verrucomicrobiota (**Fig. 1c**). It is possible that these organisms either depend on NO

reduction by other organisms or produce N_2O via pathways that do not depend on Nor

[27–30].

Among all genomes carrying *nosZ* (i.e. terminators, initiator-terminators, and complete

denitrifiers), those encoding clade I *nosZ* were more likely to encode the complete

denitrification trait compared to genomes encoding clade II (74% in clade I vs. 26% in

clade II; **Supplementary Fig. 2**), as previously observed [4]. Within clade II, complete

denitrification was most prevalent in Pseudomonadota and Aquificota, with 80% of

Pseudomonadota and 94% of Aquificota encoding complete denitrification compared

to an average of 20% in the remaining phyla. This suggests that the difference in clade

I vs. clade II prevalence can be used as a proxy for complete vs. partial denitrifier

prevalence within the N_2O reducing community, but not if clade II is dominated by

Pseudomonadota or Aquificota.

Inferred ecological preferences of complete vs. partial denitrifiers

We next assessed whether the genetic potential for complete denitrification or for

initiating rather than terminating denitrification was associated with the predicted

growth rate and resource use patterns among bacteria. Using the codon bias-based

maximum growth rate prediction tool gRodon [31], we categorized genomes into fast-

(growth rate faster than 0.2 h^{-1}) and slow-growing (growth rate lower than 0.2 h^{-1}) [31].

Rapid growth was more common among complete denitrifiers (83%) compared to

initiators (70%), terminators (65%) and initiator-terminators (62%; **Fig. 2a upper**

panel) and for genomes encoding clade I (85%) compared to clade II NosZ (65%; **Fig.**

2a lower panel). Similarly, estimated median growth rates of complete denitrifiers

(0.46 h^{-1}) were greater than that of initiators (0.28 h^{-1}), terminators (0.23 h^{-1}) and

initiator-terminators (0.20 h^{-1}; overall Kruskall-Wallis $\chi^2 = 41$, Dunn $P < 0.001$ in all

cases). This faster potential growth rate was associated with a lower sugar to acid

preference of complete denitrifiers compared to all three of the categories of partial

denitrifiers, based on the total sum of genes involved in each pathway [32] (**Fig. 2b**). This

was accompanied by a higher density of transcription factors (**Fig. 2c**) and transporters

(68 Mbp^{-1} in complete denitrifiers vs. 65 Mbp^{-1} in initiators, 40 Mbp^{-1} in terminators,

and 42 Mbp^{-1} in initiator-terminators, $P < 2 \times 10^{-16}$). Furthermore, complete denitrifiers

were inferred to grow on more of the 56 organic substrates examined (median of 20

corresponding to 4.75 Mbp^{-1}), compared to initiators (13; 3.32 Mbp^{-1}), terminators (11;

2.90 Mbp^{-1}), and initiator-terminators (9; 2.55 Mbp^{-1}; **Fig. 2d**). This was driven by

complete denitrifiers using a broader range of organic and amino acids compared to

partial denitrifiers (median of 15 vs. 6-7), rather than sugars (median of 2 vs. 2-3).

We subsequently examined the potential for partial and complete denitrifiers to use a

range of inorganic compounds as electron donors and acceptors (**Fig. 2e**). While the

ability to oxidize inorganic compounds such as hydrogen may be associated with an

ability to tolerate energy-limited conditions [33], the ability to reduce inorganic

compounds may indicate that one type or another of denitrifiers is better able to

tolerate lower redox conditions. Although most genomes lacked the genetic potential to

oxidize inorganic substances or reduce non-nitrogenous terminal electron acceptors

(median = 0), complete denitrifiers had the genetic potential to both oxidize and reduce

more compounds than partial denitrifiers (Kolmogorov-Smirnov statistic $P < 0.0001$ in

all cases).

The inferred abiotic niche breadths of complete denitrifiers were not uniformly broader

than that of partial denitrifiers. Complete denitrifiers had slightly broader inferred

temperature ranges compared to initiators (26.84 vs. 25.12 ^{o}C; t = -24.583, $P < 2$ x 10^{-}

^{16}), terminators (25.54 ^{o}C, t = -18.569, $P < 2$ x 10^{-16}) and initiator-terminators (25.93

^{o}C, t = -8.038, $P = 1.02$ x 10^{-15}). pH ranges were narrower in complete denitrifiers

compared to initiators (3.75 vs. 3.83; t = 8.136, $P = 4.44$ x 10^{-16}) and broader compared

to terminators (3.58; t = -6.756, $P = 1.50$ x 10^{-11}). Salinity ranges were broader in

complete denitrifiers than initiators (5.64 vs. 5.31; t = -3.775, $P = 0.000162$) but

narrower than in initiator-terminators (5.92; t = -2.441, $P = 0.014665$).

We subsequently assessed the degree to which complete and partial denitrifiers were

associated with high or low resource availability using CoverM [34] to map reads from

the TARA Oceans [35] project to MAGs derived from the same data [36]. No subset of the

metagenome data we collected or other databases we screened had a sufficient number

of complete denitrifier MAGs and environmental data to allow for this analysis,

highlighting the rarity of complete denitrifiers in global ecosystems. Nitrate was used

as a proxy for denitrification electron acceptor availability and chlorophyll

concentration as a proxy for electron donor supply because of its positive association

with gross primary productivity [37]. The proportion of reads mapping to the collection of

170 MAGs in the dataset differed by sample and ranged from 5 to 30%. We calculated

a standardized environmental response by correlating the proportion of reads mapping

the denitrifier types among the MAGs (7 complete, 13 initiators, 14 terminators, 134

non-denitrifiers and 2 initiator-terminators) with chlorophyll and NO_3^- concentrations

in different samples and standardizing this to the maximum observed relative

abundance of each MAG. We found that terminators and complete denitrifiers tended

to have a greater increase in relative abundance with increasing NO_3^- than non-

denitrifiers, but there was no difference between complete denitrifiers and partial

denitrifiers (**Supplementary Fig. 3a**). Mean standardized chlorophyll and chlorophyll:

NO_3^- ratio responses were similar across all groups of organisms, independent of

denitrifier type (**Supplementary Figs. 3b,c**). One possible reason for the apparent lack

of correlation between denitrifier type and resource availability is that these bulk

measurements likely do not represent the microhabitat that denitrifiers experience. For

instance, pre-filtering of the samples depleted much of the particulate organic matter,

where denitrification activity is concentrated [38], and where modeling indicates that

complete denitrifiers are favored [21]. Nonetheless, the results from this pelagic marine

dataset do not indicate that measured resource availability affects the prevalence of

complete or partial denitrifier genomes.

Prevalence of denitrification initiators and terminators varies across global biomes

Next, we assessed variation in denitrification initiator and terminator prevalence and

complete and partial denitrifiers among terminators (*i.e.* N_2O reducers) at the community

level in a broad range of environments. We screened 3,991 metagenomes derived from soil,

aquatic, engineered and host-associated biomes for *nirK*, *nirS*, and *nosZ* clades I and II using

an HMM-based search and phylogenetic placement approach [39]. The difference in the copy

number of *nosZ* and *nir* genes normalized to Gbp sequenced in each metagenome (δnos-*nir*)

was used as a proxy for the genetic potential for termination compared to initiation of

denitrification, and *nosZ* clade I vs. clade II ($\delta nosZI$-*nosZII*) as a proxy for complete

denitrifier dominance within the N$_2$O reducing community. Delta values were chosen for

their ability to account for differences in overall denitrifier prevalence between biomes,

including cases where one or both of the denitrification genes being compared were not

detected.

Across the majority of biomes, *nir* gene fragments were typically more abundant than *nosZ*,

indicating a greater or similar potential to initiate than terminate denitrification within the

denitrifier communities (**Fig. 3a**). Marine mats served as an exception, and *nosZ* prevalence

exceeded that of *nir* by 60%. This would occur if there were higher rates of NO$_3^-$ assimilation

or low inputs of NO$_2^-$ from nitrification [40], as suggested by the overall low prevalence of *nir*

and lack of ammonifier associated *nirK* clades observed previously in these samples [25]. *nosZ*

Clade II was more prevalent than *nosZ* Clade I in nearly all biomes considered, (**Fig. 3b**).

This indicates that partial denitrifiers generally dominate terminator communities because

partial denitrification is more prevalent among Clade II *nosZ*. This is further supported by our

observation that phyla depleted in complete denitrifiers, such as Bacteroidetes, Chloroflexota

and Gemmatimonadota, dominated *nosZ* clade II across biomes (**Figs. 3c,d**). Therefore, we

can conclude that the majority of organisms capable of terminating denitrification in global

biomes do not also initiate this process. This is particularly the case in croplands, marshes,

and activated sludge from municipal wastewater treatment plants, which were the biomes

where δ*nosZ*I-*nosZ*II is lowest for terrestrial, marine, and engineered environments (**Fig. 3b**).

The dominance of *nosZ*II coincides with the highest total *nosZ* and *nir* gene abundances

(**Supplementary Data 1**), indicating conditions that are overall more favorable for

denitrification also promote N$_2$O reducers that are partial rather than complete denitrifiers. A

notable exception to this pattern is sewage communities, which were dominated by clade I

nosZ despite having high overall *nir* gene fragment prevalence, and seems to largely reflect

the preponderance of *Acidovorax* in these samples [41].

We also assessed the correlation between the diversity of denitrifiers based on phylogenetic

diversity of the dominant nitrite reductase gene, *nirK* and potential initiator-terminator or

clade I and II differential abundance [25]. We found that *nirK* phylogenetic diversity was

positively correlated with δnos-*nir* in soils (Spearman's $\rho = 0.46$, $P < 2.2 \times 10^{-16}$) and marine

samples ($\rho = 0.19$, $P = 1.18 \times 10^{-11}$), while *nirK* diversity was positively correlated with

$\delta nosZ$I-*nosZ*II in soils ($\rho = 0.29$, $P < 2.2 \times 10^{-16}$) and uncorrelated in marine samples ($\rho =$

0.03, $P = 0.35$). These results indicate that more diverse denitrifier communities occur where

capacities for initiating and terminating denitrification are relatively more balanced, such that

complete denitrification at the community level is associated with greater niche partitioning.

Environmental drivers of *nir* vs. *nosZ* and *nosZ* clade prevalence were assessed using

random forest modeling on the largest soil and marine datasets with complete metadata

represented in our analysis [42]. We generated accumulated local effect plots to show the

main effect of the target variable on predicted δnos-*nir* (**Fig. 4**) and $\delta nosZ$I-*nosZ*II (**Fig.**

5), while accounting for the other predictors. Models explained less than half of the

variance in gene differences, except for $\delta nosZ$I-*nosZ*II in soils. Increases in soil organic

carbon content up to ~1% positively affected δnos-*nir,* while the effect of soil NO_3^-

content was always negative (**Fig. 4a**). *nosZ* was also predicted to become less

prevalent compared to *nir* as available phosphorus increased, indicating that high

nutrient conditions in soils favor NO_2^- reducers over N_2O reducers. Within the marine

dataset, predicted δnos-*nir* was lower at the highest concentrations of the sum of NO_3^-

and NO_2^- compared to the lowest concentrations, and showed a dramatic shift in

prediction from a decreasing to increasing relationship with temperature at 22 ^{o}C (**Fig.**

4b).

Available NO_3^-, phosphorus, potassium, and zinc had a negative effect on predicted

δ*nosZ*I-*nosZ*II (**Fig. 5a**), which suggest that partial N_2O reducers increase with

increasing nutrient levels. pH and clay content also had a negative effect on predicted

δ*nosZ*I-*nosZ*II, though only pH had a negative effect over the entirety of measured

values. Increases in elevation at low altitude were associated with increased predicted

δ*nosZ*I-*nosZ*II, but additional increases in elevation were not associated with further

increases. In the marine study, temperature and ammonium concentrations had

opposing unimodal effects on predicted values over their observed ranges (**Fig. 5b**).

Expression of denitrification termination and initiation in the environment

We subsequently quantified the presence of *nosZ* and *nir* transcripts in 413

metatranscriptomes from soils and aquatic environments, distributed among five

biomes, to assess whether the biome-level differences and associations between

environmental variables and δ*nosZ*I-*nosZ*II and δ*nos*-*nir* gene abundance were also

apparent in gene expression. *nirK* transcripts were strongly dominated by archaeal

nitrifier reads (median 50%, range 0-100%), which we excluded in our analysis (see

methods). Consistent with the metagenome analysis, *nir* was more prevalent than *nosZ*

and clade II *nosZ* was more prevalent than clade I *nosZ* (**Fig. 6a**). The soil studies

either directly or indirectly manipulated carbon availability, and soils associated with

higher carbon availability had higher δ*nos*-*nir* expression than their paired lower-

carbon samples (SMD 0.75, p=0.015) but did not have higher δ*nosZ*I-*nosZ*II (SMD

0.06, p = 0.73). Among aquatic studies, which were all observational, correlation

coefficients were negative between δ*nos-nir* and the $NO_3^- + NO_2^-$ (mean -0.27, CI: -

0.53 - 0.00) and positive between δ*nos-nir* and chlorophyll concentrations (0.22 CI

0.04, 0.41; **Fig. 6b**). δ*nos-nir* was not correlated with dissolved oxygen or phosphate

concentrations, or with bacterial production, another proxy for resource availability.

δ*nos*ZI-*nos*ZII was positively correlated with dissolved oxygen content, which would

occur if the dominant organisms encoding clade I NosZ lowered oxygen concentrations

around their N_2O reductase better than those encoding clade II [43].

Discussion

By combining a comparative analysis of genomes with broad metagenomic and

metatranscriptomic surveys, we provide unprecedented insight into the prevalence of the

complete and partial denitrification trait among global microbial communities. In contrast to

a previous comparative genomics study based on a small set of isolated microbes [4] but

consistent with more recent studies of MAGs from different environments [5,9,44–47], we found

that the genetic potential for complete denitrification is less common than that for partial

denitrification, among both genome-sequenced microorganisms and environmental

communities from all major biomes. Our results show denitrification is primarily a

community trait based on division of labour, thereby relying cross feeding between

microorganisms producing and consuming NO_2^-, NO and N_2O to complete the pathway.

Furthermore, other genes and genomic traits implicate complete denitrifiers as metabolically

flexible generalists compared to any partial denitrifier type.

Various explanations exist for the dominance of partial denitrifiers. We show that partial

denitrification enables greater overall niche partitioning, which was supported by narrower

substrate ranges, lower capacity for oxidizing and reducing various inorganic electron donors

and acceptors, and a lower genomic density of transcription factors and transporters among

partial denitrifiers compared to complete denitrifiers. In the environment, higher *nirK*

phylogenetic diversity, an indication of greater functional diversity among nitrite reducers [25],

was associated with an increasing balance between capacity to initiate and terminate

denitrification at the community level. This would further support niche partitioning if the

pathways were split among partial denitrifiers, which is likely considering that that we mainly

observed negative δ*nos-nir* values in the environment and the fact that MAGs are

predominantly (84%) partial denitrifiers. Partial denitrifiers may also dominate communities

due to the presence of a rate-efficiency tradeoff, i.e. that energy flux is slower through long

pathways but potentially enables more complete usage of the terminal electron acceptor when

it is limiting [19]. However, the dominance of partial denitrifiers among N_2O reducers in the

environment was not coherently explained by a rate-efficiency tradeoff. If this were the case,

we would expect complete denitrification to decline as NO_3^- increases or as $C:NO_3^-$ and

$C:NO_2^-$ ratios decrease, assuming denitrifiers are predominantly heterotrophs in the

environment. This agrees with predictions based on modelling and observations in oxygen

minimum zones in the oceans, proposing that the prevalence of a complete pathway increases

as the limiting substrate shifts from C to N [21]. Accordingly, our random forest modeling

inferred that the N_2O reducing community in higher NO_3^- soils became more dominated by

partial denitrifiers, i.e. those carrying *nosZ* clade II, and fertilized croplands had more

negative δ*nosZ*I-*nosZ*II than other soils in our cross-biome study. However, a similar

reduction in δ*nosZ*I-*nosZ*II was not associated with increasing $NO_3^- + NO_2^-$ in aquatic

metagenomes and metatranscriptomes or in the analysis mapping reads to marine MAGs. In

addition, there was no relationship between increasing carbon content in soil or higher

chlorophyll concentrations in aquatic samples and higher prevalence of complete denitrifiers

in either metagenomes or metatranscriptomes. This is consistent with a recent qualitative

analysis of ~1600 MAGs [46], which concluded that a rate-efficiency tradeoff cannot explain

observed partial denitrifier dominance. Based on our findings of broader substrate ranges and

greater genomic allocation to substrate uptake and transcriptional regulation, we instead

propose that complete denitrifiers are adapted to flexibly take advantage of resources varying

in time and space, while partial denitrifiers are restricted to slower and therefore less variable

growth rates. This may explain the over-representation of complete denitrifiers among model

denitrifiers and isolates that grow readily under standard resource-rich lab conditions. Under

carbon-rich conditions supporting rapid growth in the environment, complete denitrifiers may

benefit from using available NO_3^-/NO_2^- more efficiently [21], while under rapidly fluctuating

conditions many have the flexibility to use whichever electron donor and terminal electron

acceptors are most readily available. This indicates that partial denitrifiers dominate because

they have cheaper metabolisms to run, that do not require such complex regulatory

mechanisms. This is consistent with the overall dominance of slow-growing organisms in

natural environments [31,48,49].

The community level genetic balance between initiation and termination of denitrification

indicates a higher capacity for N_2O production than reduction in nearly all biomes. The

variations between biomes and among the specific ecosystems within biome categories were

readily explained by environmental variables. We found that high NO_3^- levels were

associated with lower $\delta nos\text{-}nir$ in both soils and aquatic systems, indicating greater imbalance

between the two genes and a decrease in the relative importance of N_2O reducers. In soils, the

same pattern was observed for phosphorous. High concentrations of NO_3^- may also allow

denitrifiers to outcompete ammonifiers, which both theoretical [50,51] and empirical evidence

indicate prevail under high C:NO_3^- [52,53], including in the soil metagenomes used in this study

[54]. Overall, this shows that nutrient rich environments promote denitrifier communities

dominated by initiators and similar patterns were observed at the transcriptional level in

aquatic environments (no data for soils). The observed imbalance in the denitrifier

community could explain why nutrient loaded environments like agricultural soils are major

sources of N_2O. However, complete denitrifiers have been observed to express only enzymes

involved in denitrification initiation under high NO_3^- and/or NO_2^- [55,56], which also lead to

accumulation of N_2O. Furthermore, high fertilization rates commonly favor nitrifier-

denitrification [57], and hydroxylamine from nitrification and high NO_2^- concentrations can

both induce N_2O production from chemodenitrification [57,58]. Finally, N_2O may also be

produced during the detoxification of NO from host immune responses [59,60]. Therefore, there

are a wealth of pathways that are distinct from denitrification and can reduce NO_2^- and/or

produce N_2O, and it is plausible that inter-pathway competition for NO_2^- favors the presence

of initiators ready to consume NO_2^- in denitrification rather than letting it enter other

pathways. Terminators could conceivably take advantage of the accumulated N_2O from all

these possible sources, in particular obligate terminators (i.e. predominantly *nosZ* clade II)

which use N_2O as terminal electron acceptor and thereby serve as important N_2O sinks [17,61].

This reasoning agrees with the observed low *δnosZI-nosZII* in metagenomes from agricultural

soils and wastewater treatment plants. We conclude that high levels of NO_3^- allow both

initiation and termination specialists to prevail, but with a dominance of initiators which

altogether increase the risk of N_2O emissions.

Fixed carbon availability was associated with reduced dominance of *nir* over *nosZ* in aquatic

transcriptomes and soil metagenomes, in particular in soils with less than 1% SOC. The soil

metatranscriptomes also indicated increased carbon availability could help NosZ overcome a

low competitiveness for electrons [62–64]. However, the chemical composition of the available

carbon is expected to affect the balance between initiation and termination of denitrification

and complete vs. partial denitrifiers. Organic acids have short catabolic pathways and fewer

opportunities for enzyme bottlenecks to slow the flow of electrons into the electron transport

chain compared to sugars and should favor complete denitrifiers able to use an array of

terminal electron acceptors, in agreement with our comparative genome analysis. At the

community level however, organisms metabolize and co-metabolize [65] substrates from among

tens of thousands of molecularly-distinct potential electron donors of varying bioavailability

[66,67], leading to conflicting effects of sugar and organic acids addition on N_2O to N_2

production from soils [68–70]. High dissolved oxygen concentrations were also positively

associated with dominance of clade I over clade II *nosZ* I expression but not gene abundance

in aquatic samples. There is no clear consensus on clade I being less sensitive to high oxygen

than clade II NosZ [71,72], and oxygen niche of N_2O reducers may be a function of organism

oxygen consumption [43] or even indirectly their competition for NO_2^- with nitrite oxidizing

bacteria [21]. Thus, while the mechanism for oxygen preference cannot be inferred from our

data, our results nonetheless indicate that terminators and complete denitrifiers make distinct

contributions to N_2O reduction as a function of environmental conditions.

Our results indicate that the prevalence of denitrification functional types within and between

environments should be considered in light of not only the potential organism level

advantages to having a particular denitrification genotype under given environmental

conditions, but also through biotic interactions. Denitrification does not occur in isolation,

and its steps may occur in response to its intermediates being produced by other pathways.

Further, spatiotemporal variation in electron donor and acceptor ratios can enable coexistence

of organisms completing different steps of denitrification. Future work addressing phenotypic

plasticity of denitrifiers under realistic environmental conditions is necessary to resolve this

uncertainty and disentangle biotic and abiotic drivers of denitrifier community assembly. This

will be particularly enhanced by improved generation of high-quality assemblies of

representative environmental genomes, and by the ability to sample microbial communities

on the scale at which they interact with one another and the environment.

Methods

Denitrification enzyme databases

Alignments and phylogenies for NirK and NirS were obtained from Pold *et al.* [25]. The Nor

reference database was derived from Murali *et al.* [73] and consisted of 67 Nor sequences

(cNOR, qNOR, bNOR, eNOR, gNOR and sNOR, and nNOR) and 865 other heme copper

oxidases. A hidden Markov model for NosZ was generated by updating an alignment from

Graf *et al.* [4] with NosZ sequences present in RefSeq release 95, following the procedure

described in Pold *et al.* [25]. After manually checking the alignment for conserved residues [74]

and trimming the alignment to its conserved core, we built a phylogeny using IQ-TREE v.

2.1.3 with the best model identified by modelfinder (LG+R10) [75]. Nitrate reductases,

catalyzing the first step in the denitrification pathway, were excluded from the analyses

because they are found in a wide variety of microorganisms and their activity often occurs in

isolation from the characteristic gas-producing denitrification process itself [24,47,76].

Genome annotation

We used the representative set of genomes from GTDB v214.1 for our analysis

(**Supplementary Data 2**).Genomes less than 80% complete, more than 5%

contaminated, or which failed taxonomic consistency of contigs based on GUNC

v.1.0.6 [77] were excluded from our analysis, resulting in 61,293 genomes in our final

dataset. We included both isolate (49% of assemblies) and culture-independent genome

assemblies (metagenome assembled and single cell genomes; 51%) in our analysis to

balance genome quality with inclusion of genetic diversity representative of

environmental samples. We identified genes for NirK, NirS, Nor, and NosZ in each

genome using an HMMsearch [78] against the corresponding amino acid databases as

described above. After aligning potential proteins to the reference, we imported them

into ARB v.7.0 [79] and used a combination of phylogeny building with FastTreeMP

v.2.1 [80] and manual curation to exclude reads lacking conserved ligand binding

domains. Nor and NosZ were also classified according to protein class (i.e. bNOR,

cNOR, eNOR, gNOR, nNOR, qNOR and sNOR for Nor and halophiles, clade I and

clade II for NosZ). We then classified each genome into non-denitrifiers (lacking *nir*

and *nosZ*), complete denitrifiers (*nir*, *nor*, and *nosZ*), and partial denitrifiers (lacking

nir or *nosZ*), with partial denitrifiers further divided into "initiators" (*nir* with or

without *nor*), terminators (*nosZ* with or without *nor*) and initiator-terminators (*nir* and

nosZ; **Fig. 1b**). Anammox genomes carrying *nir* (n = 47) were classified as non-

denitrifiers because their nitrite reductase is not thought to be involved in

denitrification [81]. Similarly, genomes encoding just *nor* were denoted as non-

denitrifiers (n = 3,064) as a conservative measure since they are likely involved in

detoxification rather than respiration. Since we only used bacterial genomes, our

analysis includes ammonia oxidizing bacteria potentially capable of nitrifier-

denitrification, but not ammonia oxidizing archaea for which a respiratory function for

nitrite reductase has not yet been established [82,83].

A range of tools were used to identify and validate genes associated with potential electron

donors and acceptors used by microorganisms. Genes for proteins involved in arsenite

oxidation (AioBA; validated with alignment from Quemeneur *et al.* [84]), anammox (Hzs, Hdh)

and iron redox were identified in genomes using MagicLamp v.1.0 with final annotations

curated to match minimum subunit composition proposed for each enzyme [85]. We used AmoA

as a marker for ammonia oxidation in bacteria using the alignment of AmoA, HmoA and

PmoA from Diamond *et al.* [5]. Respiratory reductive dehalogenases were identified using the

RDaseDB alignment (v.2020 [86]) combined with a literature search to exclude catabolic

enzymes; we note that the TAT motif proposed to be indicative of respiratory enzymes [87] was

absent from the majority of sequences and could not be used. Sulfur redox genes were

identified using HMSS2 and categorized into oxidation or reduction-associated enzymes

following Tanabe and Dahl [88]. Sulfide: quinone reductase (SQR) enzymes were excluded

from our analysis, both because their physiological role is variable, and because we found the

HMMs were unspecific. Hydrogenases were identified using MagicLamp, refined using the

residues in Greening *et al.* [33], and verified as to their clade and probable physiological

function using the HydDB web server [89]. Clades 1a-d, 1g,h and j and clade 2 were considered

involved in hydrogen oxidation, and clades 4 a,b,c and h were considered involved in

reduction. ArrA and ArxA involved in arsenate reduction and arsenite oxidation, respectively,

were identified using an HMM built from sequences from Wells *et al.* [90], and validated using

the residues from Ospino *et al.* [91]. All redox gene annotations are reported in **Supplementary**

Data 3.

Sugar-acid preference of organisms was inferred using the sum of KEGG orthologues

identified by enrichM (v0.6.3, default settings) that were annotated to sugar vs. acid

metabolism [32], noting that the majority of the genomes fall outside the originally-calibrated

range of GC contents (**Supplementary Data 4**). GapMind carbon [92] (March 22nd, 2021

database version) was run to infer ability to grow on 62 compounds. We used custom

pathway completeness cutoff scores for each pathway based on annotations for organisms for

which growth had been reported on that substrate in BacDive (**Supplementary Data 5**). Our

positive controls dataset did not have data for 6 substrates, leaving 56 for analysis.

Transporters were annotated using DIAMOND against the TCDB database, with hits showing

at least 40% identity to the reference and 70% coverage of both reference and query

sequence, with no more than 10% difference in alignment length kept for analysis [93].

Transcription factors were annotated using DeepTFactor [94] (v. 2020-11-09) with default

settings (**Supplementary Data 6**). We used GenomeSPOT to infer temperature, pH, and

salinity ranges [95], and used the difference between the maximum and minimum values for

each genome in our analysis (**Supplementary Data 7**).

Unless otherwise noted, all subsequent analyses were completed in R v.4.1 [96]. The

maximum growth rate of each taxon was estimated using gRodon [31], which used codon

usage bias of ribosomal proteins annotated in the genomes using hmmearch against the

ribosomal proteins extracted from Uniprot PFAM v. 35.0 with profile-specific cutoffs

(**Supplementary Data 8**). Differences in growth rates between denitrifier types were

assessed using Kruskall-Wallis tests (stats::kruskal.test) followed by Dunn tests with

Benjamini-Hochberg correction for multiple testing (FSA::dunnTest), where all

maximum growth rates less than 0.2 doublings h^{-1} were given the same rank. This

cutoff was needed because codon use bias saturates with growth rates slower than this

[31].

To determine whether complete and partial denitrifiers differ in transcription factor,

transporter, carbohydrate-active enzyme, or catabolized substrate count or sugar acid

preference, we fit linear models, logging or taking the exponent of variables where

needed. However, we analyzed GapMind data with a generalized linear mixed model

with a Tweedie distribution in glmmTMB [97] (v.1.1.10) due to the poor fit of OLS and

high proportion (10-37%) of assemblies where no substrates within a chemical class

were predicted to be used. We subsequently ascertained that the model residuals were

not phylogenetically clustered using phytools::phylosig (v.2.3.0 [98]) and an organism

tree derived from the GTDB reference tree [99]. This tree was rerooted in Fusobacteriota

[100] and forced to be ultrametric using the castor::date_tree_red function [101] (v.1.8.2).

Blomberg's K [102] of model residuals was less than one in all instances ($P < 0.05$), so we

did not include phylogenetic relatedness as a random effect in our analysis. We

evaluated differences between complete and partial denitrifier genome content using

post-hoc multiple comparisons with the sandwich::vcovHC function (v.3.1.1 [103]) to

account for heteroskedasticity. Due to the large number of zeroes, we assessed the

hypothesis that partial denitrifiers encode the potential to oxidize and reduce a smaller

number of inorganic compounds than complete denitrifiers by comparing the

cumulative density distributions using a one-sided Kolmogorov-Smirnov test using

stats::ks.test.

Metagenome and metadata collection

To assess the dominance of complete vs. partial denitrifier communities across biomes,

we selected metagenomes with at least 100,000 reads ≥150 nt, primarily from larger

sampling campaigns where uniform metadata were available (**Supplementary Data 9**).

The majority of soil metagenomes came from the Australian Microbiome

Project/BASE [104], the National Ecological Observatory Network [105], Topsoils

Microbiome Project [106], Long-Term Soil Productivity experiment [107], and the Stordalen

Mire [108]. The majority of aquatic metagenomes came from the Australian Microbiome

Project [109], Linnaeus Microbial Observatory [110], the Amazon Continuum Project [111] and

BATS, GEOTRACE and HOTS [112]. Engineered metagenomes included those from

drinking water [113] sewage [41], and wastewater treatment plants [114]. The host-associated

metagenomes dataset consists primarily of plant-associated sequences, such as from

leafy greens [115], beans [116], citrus rhizosphere [117], switchgrass [118] and *Arabidopsis* [119],

though we also included a handful of sequences from invertebrates such as sponges and

gutless marine worms. For unpublished metagenomes where contact details were

provided, we emailed authors to request permission to use them.

We completed a similar search for metatranscriptomes, focusing on soil (including

rhizosphere) and marine studies where carbon and/or nutrient availability had either

been manipulated or logic or metadata indicated they differed between samples

(**Supplemental Data S10**). These metatranscriptomes were identified based on

searching the American Society for Microbiology journals website for

metatranscriptomes and google databases for the terms "soil" or "marine" and "carbon"

or "nutrient" and "metatranscript*".

Where multiple sequencing runs or files existed for the same sample, we combined the

fastq files prior to searching. We used only the forward read in the case of paired end

reads, and only the first 150 nt of any reads longer than that. In total, our dataset

comprised 3,991 metagenomes, with 1,489 from aquatic habitats, 1,642 from soil, 658

host-associated, and 202 from engineered habitats. Our final set of metatranscriptomes

only shares samples with a handful of samples from a single study in our metagenome

dataset and contains 413 metatranscriptomes (64 rhizosphere or soil with root exudates

added, 92 other soil, 185 primarily coastal and estuarine marine, 24 lake sediment, and

48 river metatranscriptomes).

Biomes were assigned to metagenomes based on The Nature Conservancy Terrestrial

Ecoregions for terrestrial samples [120] and latitude for marine samples (polar: latitude >

60°; westerlies: latitude 30-60°; trades: latitude 0-30°). Terrestrial biome assignment

required use of the packages sp v.1.4-5 [121], rgdal v.1.523 [122], and rgeos v0.5-5 [123]. Soils

under cultivation were excluded from the biome-based approach and instead

categorized as croplands.

Dominance of complete vs. partial denitrifier communities based on proxy gene

searches across biomes

We used GraftM [39] to identify denitrification genes (*nirK*, *nirS* and *nosZ*) in the

metagenomes and metatranscriptomes, as previously described [25]. Briefly, GraftM uses

a two-step process in which a HMM search identifies candidate reads, followed by

phylogenetic placement on a reference tree using pplacer (v.1.1.19; [124]). We used the

accumulate function in gappa[125] to find the position on the tree where at least 95% of

the mass for each read descended from, and excluded all reads with any mass placed in

the outgroup. We further excluded sequences placed in the non-denitrifying anammox

NirS clade 1h (median 0% of *nirS* reads in metagenomes and 0% in

metatranscriptomes), archaeal nitrifier NirK clades 2 and 4 (1.4% of *nirK* reads in

metagenomes and 50% in metatranscriptomes), as well as the eukaryotic NirK clades

1b and 1e (0% and 0%) [25]. Our analysis also excluded the 0.4% of reads placed in the

bioinformatically inferred "clade III" *nosZ* [10], which forms the outgroup in our

reference tree. This metagenome and metatranscriptome search depended on: hmmer

(v.3.2.1; [78]), OrfM (v.0.7.1; [126]), bbtools (v.38.90;

"https://sourceforge.net/projects/bbmap/"), lbzip2 (v.2.5; "https:// lbzip2.org/"), and

fxtract (v.2.3; "https://github.com/ctSkennerton/fxtract").

We validated these search and placement and post-processing parameters by generating 150

nt fragments of full-length sequences that were picked up by the NirK, NirS and NosZ

HMMs, which in addition to the target proteins also included homologous multicopper

oxidase proteins from Cyanobacteriota and Thermoprotetoa, NirN and NirF, and various yet

to be characterized NosZ-like proteins, respectively. Sensitivity is the fraction of total

ingroup fragments searched which were placed in the ingroup, and specificity is one minus

the fraction of outgroup fragments incorrectly placed in the ingroup (or, in the case of *nosZ*,

in a clade other than the target clade). Sensitivity was 76% for *nirK,* 93% for *nirS,* 91% for

clade I *nosZ* and 94% for clade II *nosZ*. Specificity was 97% for *nirK*, 100% for *nirS*, 100%

for clade I *nosZ* and 100% for clade II *nosZ*.

We compared gene prevalence across biomes using the absolute difference in counts of

clade I and II *nosZ* or *nir* and *nosZ* genes standardized by the number of Gbp

sequenced to account for differences in sequencing depth. For metatranscriptomes we

standardized based on Gbp mRNA sequenced, which was determined by excluding

rRNA reads from total RNA using SortMeRNA v.4.3.7 with the smr_v4.3_default_db

database and settings "--other --fastx --num-alignments 1 --no-best". For *nir* vs. *nosZ*

abundance we used Eq. 1 such that positive values denote *nosZ* is more abundant than

nir:

$$\delta nos - nir = 10^9 \text{ x } ((nosZI + nosZII) - (nirS + nirK))/(150 \text{ x reads}) \qquad (1)$$

For *nosZ*, we used Eq. 2 such that positive values denote clade I is more abundant than clade

II:

$$\delta nosZI\text{-}nosZII = 10^9 \text{ x } (nosZI - nosZII)/(150 \text{ x reads}) \qquad (2)$$

This method of calculating gene prevalence enables the abundance of both genes to be

considered concurrently. This is particularly relevant for the 14% of metagenomes and 26%

of metatranscriptomes where *nos*ZI was not detected but *nos*ZII was found in 1-13,041 (1-

10,926 in metatranscriptomes) copies, and the 0.8% of metagenomes where *nos*Z was not

found but 3-28 copies of *nir* were identified.

Environmental correlates with biome gene counts and denitrifier MAGs

We used the TARA Oceans [35] MAGs present in the OceanDNA MAG database to test

the hypothesis that complete denitrifiers are more abundant under conditions with high

electron donor to acceptor ratios. These MAGs were chosen because the database is

sufficiently large that multiple MAGs from each denitrifier type category were present,

and where multiple conspecific MAGs were present, those with inconsistent

denitrification gene repertoires could be excluded. After quality-controlling the raw

paired-end reads with cutadapt (-m 100 -q 20 –max-n 0 –trim-n), we mapped them to

all MAGs using CoverM (v. 0.6.1; [34]; -m relative abundance –min-read-aligned-

percent 0.75 –min-read-percent-identity 0.95 –min-covered-fraction 0) then determined

the total relative abundance of each MAG-OTU in each sample (95% ANI). Only the

reads for the 3 μm filter fraction were used in this analysis to allow for bigger cells or

cell aggregates than using the 0.22 μm fraction. We subsequently determined the

standardized relative abundance by dividing the observed relative abundance within

each sample to the maximum relative abundance observed across all samples. We then

fit a linear model predicting the standardized relative abundance of each MAG OTU

based on the combination of NO_3^- and chlorophyll concentrations (i.e. NO_3^- after

accounting for chlorophyll, chlorophyll after accounting for NO_3^-, and the logarithm of

chlorophyll: NO_3^-). The mean slope and standard error for each MAG OTU were

extracted and used to calculate the weighted mean response and 95% confidence

intervals for each denitrifier type. Differences in mean response to NO_3^-, chlorophyll

and their ratios between denitrifier types were established based on non-overlapping

95% confidence intervals.

We evaluated the relationship between differences in normalized gene counts and

resource availability using random forests. This analysis was restricted to

metagenomes from the BASE and Australian marine projects[104,112] as examples of

studies with a large number of metagenomes with uniformly-collected metadata for

variables associated with denitrification, including NO_3^-/NO_2^- levels, pH, carbon, and

micronutrients. A pre-selection of variables for inclusion in the model was completed

to exclude collinear variables (Spearman's ρ >0.7 or variance inflation factor > 4); in

the collinear groups, the variable hypothesized to be most strongly and directly

explicable of gene ratio was retained (Supplementary Fig. 3; Supplementary Tables 2

and 3). We subsequently ran VSURF v. 1.1.0[127] 100 times to identify the best

predictors for each ratio and biome, keeping only those variables retained in at least 95

of the 100 iterations. We then used the randomForest package v. 4.7.1-1[128] to model

the relationship between gene ratios and the final set of retained variables. A grid

search was used to find the combination of tuning parameters yielding lowest out-of-

bag root-mean-square error. Accumulated local effects plots were generated to

visualize results using iml v. 0.11 [129].

Finally, we verified whether high carbon and/or low nutrient availability favored the

transcription of *nosZ* over *nir* or clade I *nosZ* over clade II *nosZ* by calculating $\delta nos-$

nir and $\delta nosZ$I-*nosZ*II in the metatranscriptomes, respectively. We calculated Hedges

G in a meta-analysis of effect sizes for soil studies (meta::metacont), taking the lower

carbon soil as the reference and carbon-enriched soil as the treatment (e.g. bulk soil or

unamended soil compared to rhizosphere, glucose, glycine, or root exudate amended

soils). Aquatic studies were observational, so we fit a Spearman correlation coefficient

between δnos-nir or δnosZI-nosZII and environmental variables (NO_3^- + NO_2^-,

phosphate, bacterial production and chlorophyll A concentrations as proxies of

resource availability, and oxygen). Where communities were captured and analyzed on

two different filter sizes from the same sample, we determined the correlation

separately for the two size fractions but counted them as a single study in our overall

effect size calculations. We used psychmeta::ma_r to calculate overall correlations in

the aquatic data.

Data availability

All genomes, metagenomes, and metatranscriptomes used in this paper are publicly available

in NCBI or other sources under the identifiers denoted in **Supplementary Data 2, 9** and **10**.

Hidden Markov Models, and reference databases used for denitrification gene searches are

available in FigShare (10.6084/m9.figshare.23913078 for NirK and NirS and

10.6084/m9.figshare.24922776 for NosZ). Source Data are provided with this manuscript.

Code availability

Scripts used to generate figures are available in FigShare (10.6084/m9.figshare.24922776).

Acknowledgments

This work was supported by the Swedish University of Agricultural Sciences Senior Career

Grant 2019-2025 to SH and the Swedish Research Council Grant Agreements No. 2016-

03551 to SH and 2023-03627 to GP. Computing resources were provided by the Department

of Forest Mycology and Plant Pathology, Swedish National Infrastructure for Computing

project SNIC 2022/22-1119 and NAISS 2023/22-1278, funded by the Swedish Research

Council through Grant Agreement No. 2022-06725, and the Swedish National Infrastructure

for Computing (SNIC) at the PDC Center for High Performance Computing, KTH Royal

Institute of Technology, partially funded by the Swedish Research Council through grant

agreement no. 2016-07213 (NAISS 2024/22-133).

Author contributions

Conceptualization: S.H. Data curation: G.P., A.S. and C.J. Formal analysis: G.P. Funding

acquisition: S.H and G.P. Investigation: G.P. Methodology: G.P., A.S., C.J. and S.H. Project

administration: G.P. and S.H. Resources: S.H. Software: C.J., G.P and A.S. Visualization: GP,

AS. Writing – original draft preparation: G.P., A.S. and S.H. Writing – review and editing:

G.P., A.S., C.J. and SH.

Competing interests

The authors declare no competing interests.

References

1.   Alvarez, L., Bricio, C., Gómez, M. J. & Berenguer, J. Lateral transfer of the

denitrification pathway genes among *Thermus thermophilus* strains. *Appl. Environ.*

Microbiol. **77**, 1352–1358 (2011).

2.   Rees, E., Siddiqui, R. A., Köster, F., Schneider, B. & Friedrich, B. Structural gene (*nirS*)

for the cytochrome cd1 nitrite reductase of *Alcaligenes eutrophus* H16. *Appl. Environ.*

Microbiol. **63**, 800–802 (1997).

3.   Schwintner, C., Sabaty, M., Berna, B., Cahors, S. & Richaud, P. Plasmid content and

localization of the genes encoding the denitrification enzymes in two strains of

Rhodobacter sphaeroides. FEMS Microbiol. Lett. **165**, 313–321 (1998).

4.   Graf, D. R. H., Jones, C. M. & Hallin, S. Intergenomic comparisons highlight

modularity of the denitrification pathway and underpin the importance of community

structure for N_2O emissions. *PLoS ONE* **9**, e114118 (2014).

5.   Diamond, S. *et al.* Mediterranean grassland soil C–N compound turnover is dependent

on rainfall and depth, and is mediated by genomically divergent microorganisms. *Nat.*

Microbiol. **4**, 1356–1367 (2019).

6.   Orellana, L. H., Chee-Sanford, J. C., Sanford, R. A., Löffler, F. E. & Konstantinidis, K.

T. Year-round shotgun metagenomes reveal stable microbial communities in agricultural

soils and novel ammonia oxidizers responding to fertilization. *Appl. Environ. Microbiol.*

84, e01646-17 (2018).

7.   Dove, N. C., Torn, M. S., Hart, S. C. & Taş, N. Metabolic capabilities mute positive

response to direct and indirect impacts of warming throughout the soil profile. *Nat.*

Commun. **12**, 1–13 (2021).

8.   Karthikeyan, S. *et al.* Metagenomic characterization of soil microbial communities in

the Luquillo experimental forest (Puerto Rico) and implications for nitrogen cycling.

Appl. Environ. Microbiol. **87**, e00546-21 (2021).

9.   Pessi, I. S. *et al.* In-depth characterization of denitrifier communities across different

soil ecosystems in the tundra. *Environ. Microbiome* **17**, 30 (2022).

10.   Mosley, O. E. *et al.* Nitrogen cycling and microbial cooperation in the terrestrial

subsurface. *ISME J.* **16**, 2561–2573 (2022).

11. Lilja, E. E. & Johnson, D. R. Segregating metabolic processes into different microbial

cells accelerates the consumption of inhibitory substrates. *ISME J.* **10**, 1568–1578

(2016).

12. Lilja, E. E. & Johnson, D. R. Substrate cross-feeding affects the speed and trajectory of

molecular evolution within a synthetic microbial assemblage. *BMC Evol. Biol.* **19**, 1–11

(2019).

13. Dolinšek, J., Ramoneda, J. & Johnson, D. R. Initial community composition determines

the long-term dynamics of a microbial cross-feeding interaction by modulating niche

availability. *ISME Commun.* **2**, 1–10 (2022).

14. Vitale, A. *et al.* Mapping of the Denitrification Pathway in *Burkholderia thailandensis*

by Genome-Wide Mutant Profiling. *J. Bacteriol.* **202**, e00304-20 (2020).

15. Zumft, W. G. & Kroneck, P. M. H. Respiratory transformation of nitrous oxide (N_2O) to

dinitrogen by bacteria and archaea. in *Advances in Microbial Physiology* (ed. Poole, R.

K.) vol. 52 107–227 (Academic Press, 2006).

16. Takahashi, K. *et al.* Denitrification in low oxic environments increases the accumulation

of nitrogen oxide intermediates and modulates the evolutionary potential of microbial

populations. *Environ. Microbiol. Rep.* **16**, e13221 (2024).

17. Jones, C. M. *et al.* Recently identified microbial guild mediates soil N_2O sink capacity.

Nat. Clim. Change **4**, 801–805 (2014).

18. Heinrich, R., Schuster, S. & Holzhütter, H.-G. Mathematical analysis of enzymic

reaction systems using optimization principles. *EJB Rev. 1991* 167–187 (1991).

19. Kreft, J.-U., Griffin, B. M. & González-Cabaleiro, R. Evolutionary causes and

consequences of metabolic division of labour: why anaerobes do and aerobes don't.

Curr. Opin. Biotechnol. **62**, 80–87 (2020).

20.   Wang, Y., Gao, H. & F. Wells, G. Integrated omics analyses reveal differential gene

781   expression and potential for cooperation between denitrifying polyphosphate and

782   glycogen accumulating organisms. *Environ. Microbiol.* **23**, 3274–3293 (2021).

783   21.   Sun, X. *et al.* Ecological dynamics explain modular denitrification in the ocean. *Proc.*

784   *Natl. Acad. Sci.* **121**, e2417421121 (2024).

785   22.   LaSarre, B., Morlen, R., Neumann, G. C., Harwood, C. S. & McKinlay, J. B. Nitrous

786   oxide reduction by two partial denitrifying bacteria requires denitrification intermediates

787   that cannot be respired. *Appl. Environ. Microbiol.* **90**, e01741-23 (2023).

788   23.   Crocker, K. *et al.* Environmentally dependent interactions shape patterns in gene content

789   across natural microbiomes. *Nat. Microbiol.* **9**, 2022–2037 (2024).

790   24.   Mahne, I. & Tiedje, J. M. Criteria and methodology for identifying respiratory

791   denitrifiers. *Appl. Environ. Microbiol.* **61**, 1110–1115 (1995).

792   25.   Pold, G. *et al.* Phylogenetics and environmental distribution of nitric oxide-forming

793   nitrite reductases reveal their distinct functional and ecological roles. *ISME Commun.* **4**,

794   ycae020 (2024).

795   26.   Park, D., Kim, H. & Yoon, S. Nitrous oxide reduction by an obligate aerobic bacterium,

796   *Gemmatimonas aurantiaca* strain T-27. *Appl Environ Microbiol* vol. 83 e00502-17

797   (2017).

798   27.   Costa, C. *et al.* Regulation of the hexaheme nitrite/nitric oxide reductase of

799   *Desulfovibrio desulfuricans*, *Wolinella succinogenes* and *Escherichia coli*. *FEBS Lett.*

800   **276**, 67–70 (1990).

801   28.   Kim, S. O., Orii, Y., Lloyd, D., Hughes, M. N. & Poole, R. K. Anoxic function for the

802   *Escherichia coli* flavohaemoglobin (Hmp): reversible binding of nitric oxide and

803   reduction to nitrous oxide. *FEBS Lett.* **445**, 389–394 (1999).

29.   Xu, Z. *et al.* Unprecedented N_2O production by nitrate-ammonifying Geobacteraceae

with distinctive N_2O isotopocule signatures. *mBio* **15**, e02540-24 (2024).

30.   Simon, J. & Klotz, M. G. Diversity and evolution of bioenergetic systems involved in

microbial nitrogen compound transformations. *Biochim. Biophys. Acta BBA-Bioenerg.*

1827, 114–135 (2013).

31.   Weissman, J. L., Hou, S. & Fuhrman, J. A. Estimating maximal microbial growth rates

from cultures, metagenomes, and single cells via codon usage patterns. *Proc. Natl.*

Acad. Sci. **118**, e2016810118 (2021).

32.   Gralka, M., Pollak, S. & Cordero, O. X. Genome content predicts the carbon catabolic

preferences of heterotrophic bacteria. *Nat. Microbiol.* **8**, 1799–1808 (2023).

33.   Greening, C. *et al.* Genomic and metagenomic surveys of hydrogenase distribution

indicate H_2 is a widely utilised energy source for microbial growth and survival. *ISME*

J. **10**, 761–777 (2016).

34.   Aroney, S. T. N. *et al.* CoverM: read alignment statistics for metagenomics.

Bioinformatics **41**, btaf147 (2025).

35.   Sunagawa, S. *et al.* Structure and function of the global ocean microbiome. *Science* **348**,

1261359 (2015).

36.   Nishimura, Y. & Yoshizawa, S. The OceanDNA MAG catalog contains over 50,000

prokaryotic genomes originated from various marine environments. *Sci. Data* **9**, 305

(2022).

37.   Morin, A., Lamoureux, W. & Busnarda, J. Empirical models predicting primary

productivity from chlorophyll a and water temperature for stream periphyton and lake

and ocean phytoplankton. *J. North Am. Benthol. Soc.* **18**, 299–307 (1999).

38.   Wan, X. S. *et al.* Particle-associated denitrification is the primary source of N_2O in oxic

coastal waters. *Nat. Commun.* **14**, 8280 (2023).

39. Boyd, J. A., Woodcroft, B. J. & Tyson, G. W. GraftM: a tool for scalable,

phylogenetically informed classification of genes within metagenomes. *Nucleic Acids*

Res. **46**, e59 (2018).

40. Coban, O., Rasigraf, O., de Jong, A. E. E., Spott, O. & Bebout, B. M. Quantifying

potential N turnover rates in hypersaline microbial mats by using ^{15}N tracer techniques.

Appl. Environ. Microbiol. **87**, e03118-20 (2021).

41. Hendriksen, R. S. *et al.* Global monitoring of antimicrobial resistance based on

metagenomics analyses of urban sewage. *Nat. Commun.* **10**, 1–12 (2019).

42. Saghaï, A. *et al.* Diversity of archaea and niche preferences among putative ammonia-

oxidizing Nitrososphaeria dominating across European arable soils. *Environ. Microbiol.*

24, 341–356 (2022).

43. Wang, Z., Vishwanathan, N., Kowaliczko, S. & Ishii, S. Clarifying microbial nitrous

oxide reduction under aerobic conditions: tolerant, intolerant, and sensitive. *Microbiol.*

Spectr. **11**, e04709-22 (2023).

44. Hester, E. R., Jetten, M. S., Welte, C. U. & Lücker, S. Metabolic overlap in

environmentally diverse microbial communities. *Front. Genet.* **10**, 989 (2019).

45. Valk, L. C. *et al.* Exploring the microbial influence on seasonal nitrous oxide

concentration in a full-scale wastewater treatment plant using metagenome assembled

genomes. *Water Res.* 118563 (2022).

46. Roothans, N., van Loosdrecht, M. C. M. & Laureni, M. Metabolic labour division trade-

offs in denitrifying microbiomes. *ISME J.* **19**, wraf020 (2025).

47. Zhang, I. H. *et al.* Partitioning of the denitrification pathway and other nitrite

metabolisms within global oxygen deficient zones. *ISME Commun.* **3**, 1–14 (2023).

48. Stone, B. W. G. *et al.* Life history strategies among soil bacteria—dichotomy for few,

continuum for many. *ISME J.* **17**, 611–619 (2023).

49. Caro, T. A., McFarlin, J., Jech, S., Fierer, N. & Kopf, S. Hydrogen stable isotope

probing of lipids demonstrates slow rates of microbial growth in soil. *Proc. Natl. Acad.*

Sci. **120**, e2211625120 (2023).

50. Tiedje, J. M., Sexstone, A. J., Myrold, D. D. & Robinson, J. A. Denitrification:

ecological niches, competition and survival. *Antonie van Leeuwenhoek* vol. 48 569–583

(1983).

51. Jia, M., Winkler, M. K. H. & Volcke, E. I. P. Elucidating the competition between

heterotrophic denitrification and DNRA using the resource-ratio theory. *Environ. Sci.*

Technol. **54**, 13953–13962 (2020).

52. Stremińska, M. A., Felgate, H., Rowley, G., Richardson, D. J. & Baggs, E. M. Nitrous

oxide production in soil isolates of nitrate-ammonifying bacteria. *Environ. Microbiol.*

Rep. **4**, 66–71 (2012).

53. Yoon, S., Cruz-García, C., Sanford, R., Ritalahti, K. M. & Löffler, F. E. Denitrification

versus respiratory ammonification: environmental controls of two competing

dissimilatory NO_3^-/NO_2^- reduction pathways in *Shewanella loihica* strain PV-4. *ISME J.*

9, 1093–1104 (2015).

54. Saghaï, A., Pold, G., Jones, C. M. & Hallin, S. Phyloecology of nitrate ammonifiers and

their importance relative to denitrifiers in global terrestrial biomes. *Nat. Commun.* **14**,

8249 (2023).

55. Saleh-Lakha, S. *et al.* Effect of nitrate and acetylene on *nirS*, *cnorB*, and *nosZ*

expression and denitrification activity in *Pseudomonas mandelii*. *Appl. Environ.*

Microbiol. **75**, 5082–5087 (2009).

56. Thomas, K. L., Lloyd, D. & Boddy, L. Effects of oxygen, pH and nitrate concentration

on denitrification by Pseudomonas species. *FEMS Microbiol. Lett.* **118**, 181–186 (1994).

57.   Wrage-Mönnig, N. *et al.* The role of nitrifier denitrification in the production of nitrous

879   oxide revisited. *Soil Biol. Biochem.* **123**, A3–A16 (2018).

880   58.   Venterea, R. T. Nitrite☐driven nitrous oxide production under aerobic soil conditions:

881   kinetics and biochemical controls. *Glob. Change Biol.* **13**, 1798–1809 (2007).

882   59.   Poole, R. K. Flavohaemoglobin: the pre-eminent nitric oxide–detoxifying machine of

883   microorganisms. *F1000Research* **9**, F1000-Faculty Rev-7 (2020).

884   60.   Wonderen, J. H. van, Burlat, B., Richardson, D. J., Cheesman, M. R. & Butt, J. N. The

885   nitric oxide reductase activity of cytochrome c nitrite reductase from *Escherichia coli*. *J.*

886   *Biol. Chem.* **283**, 9587–9594 (2008).

887   61.   Onley, J. R., Ahsan, S., Sanford, R. A. & Löffler, F. E. Denitrification by

888   *Anaeromyxobacter dehalogenans*, a Common Soil Bacterium Lacking the Nitrite

889   Reductase Genes *nirS* and *nirK*. *Appl. Environ. Microbiol.* **84**, e01985-17 (2018).

890   62.   Wei, Q. *et al.* Molecular mechanisms through which different carbon sources affect

891   denitrification by *Thauera linaloolentis*: Electron generation, transfer, and competition.

892   *Environ. Int.* **170**, 107598 (2022).

893   63.   Pan, Y., Ni, B.-J., Bond, P. L., Ye, L. & Yuan, Z. Electron competition among nitrogen

894   oxides reduction during methanol-utilizing denitrification in wastewater treatment.

895   *Water Res.* **47**, 3273–3281 (2013).

896   64.   Schalk-Otte, S., Seviour, R. J., Kuenen, J. G. & Jetten, M. S. M. Nitrous oxide (N_2O)

897   production by *Alcaligenes faecalis* during feast and famine regimes. *Water Res.* **34**,

898   2080–2088 (2000).

899   65.   McGill, S. L. *et al.* Pseudomonas aeruginosa reverse diauxie is a multidimensional,

900   optimized, resource utilization strategy. *Sci. Rep.* **11**, 1457 (2021).

901   66.   Dufour, L. J. P. *et al.* Potential energetic return on investment positively correlated with

902   overall soil microbial activity. *Soil Biol. Biochem.* **173**, 108800 (2022).

903   67.   LaRowe, D. E. & Van Cappellen, P. Degradation of natural organic matter: A

904   thermodynamic analysis. *Geochim. Cosmochim. Acta* **75**, 2030–2042 (2011).

905   68.   Morley, N. & Baggs, E. M. Carbon and oxygen controls on N_2O and N_2 production

906   during nitrate reduction. *Soil Biol. Biochem.* **42**, 1864–1871 (2010).

907   69.   Curtright, A. J. & Tiemann, L. K. Chemical identity of carbon substrates drives

908   differences in denitrification and N_2O reduction within agricultural soils. *Soil Biol.*

909   *Biochem.* **184**, 109078 (2023).

910   70.   Henry, S. *et al.* Disentangling the rhizosphere effect on nitrate reducers and denitrifiers:

911   insight into the role of root exudates. *Environ. Microbiol.* **10**, 3082–3092 (2008).

912   71.   Sennett, L. B. *et al.* Determining how oxygen legacy affects trajectories of soil

913   denitrifier community dynamics and N2O emissions. *Nat. Commun.* **15**, 7298 (2024).

914   72.   Bertagnolli, A. D., Konstantinidis, K. T. & Stewart, F. J. Non-denitrifier nitrous oxide

915   reductases dominate marine biomes. *Environ. Microbiol. Rep.* **12**, 681–692 (2020).

916   73.   Murali, R. *et al.* Diversity and evolution of nitric oxide reduction in bacteria and

917   archaea. *Proc. Natl. Acad. Sci.* **121**, e2316422121 (2024).

918   74.   Sanford, R. A. *et al.* Unexpected nondenitrifier nitrous oxide reductase gene diversity

919   and abundance in soils. *Proc. Natl. Acad. Sci.* **109**, 19709–19714 (2012).

920   75.   Minh, B. Q. *et al.* IQ-TREE 2: new models and efficient methods for phylogenetic

921   inference in the genomic era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

922   76.   Sun, Y., De Vos, P. & Heylen, K. Nitrous oxide emission by the non-denitrifying, nitrate

923   ammonifier Bacillus licheniformis. *BMC Genomics* **17**, 68 (2016).

924   77.   Orakov, A. *et al.* GUNC: detection of chimerism and contamination in prokaryotic

925   genomes. *Genome Biol.* **22**, 178 (2021).

926   78.   Eddy, S. R. Accelerated Profile HMM Searches. *PLOS Comput. Biol.* **7**, e1002195

927   (2011).

79. Ludwig, W. *et al.* ARB: a software environment for sequence data. *Nucleic Acids Res.*

32, 1363–1371 (2004).

80. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 – approximately maximum-

likelihood trees for large alignments. *PLOS ONE* **5**, e9490 (2010).

81. van de Vossenberg, J. *et al.* The metagenome of the marine anammox bacterium

'Candidatus *Scalindua profunda*' illustrates the versatility of this globally important

nitrogen cycle bacterium. *Environ. Microbiol.* **15**, 1275–1289 (2013).

82. Qin, W. *et al.* Alternative strategies of nutrient acquisition and energy conservation map

to the biogeography of marine ammonia-oxidizing archaea. *ISME J.* **14**, 2595–2609

(2020).

83. Stein, L. Y. Insights into the physiology of ammonia-oxidizing microorganisms. *Curr.*

Opin. Chem. Biol. **49**, 9–15 (2019).

84. Quéméneur, M. *et al.* Diversity surveys and evolutionary relationships of *aoxB* genes in

aerobic arsenite-oxidizing bacteria. *Appl. Environ. Microbiol.* **74**, 4567–4573 (2008).

85. Garber, A. I. *et al.* FeGenie: a comprehensive tool for the identification of iron genes

and iron gene neighborhoods in genome and metagenome assemblies. *Front. Microbiol.*

11, (2020).

86. Molenda, O. *et al.* Insights into origins and function of the unexplored majority of the

reductive dehalogenase gene family as a result of genome assembly and ortholog group

classification. *Environ. Sci. Process. Impacts* **22**, 663–678 (2020).

87. Fisher, K. *et al.* Efficient NADPH-dependent dehalogenation afforded by a self-

sufficient reductive dehalogenase. *J. Biol. Chem.* **299**, 105086 (2023).

88. Tanabe, T. S. & Dahl, C. HMS-S-S: A tool for the identification of sulphur metabolism-

related genes and analysis of operon structures in genome and metagenome assemblies.

Mol. Ecol. Resour. **22**, 2758–2774 (2022).

89. Søndergaard, D., Pedersen, C. N. S. & Greening, C. HydDB: A web tool for

hydrogenase classification and analysis. *Sci. Rep.* **6**, 34212 (2016).

90. Wells, M., Kim, M., Akob, D. M., Basu, P. & Stolz, J. F. Impact of the dimethyl

sulfoxide reductase superfamily on the evolution of biogeochemical cycles. *Microbiol.*

Spectr. **11**, e04145-22 (2023).

91. Ospino, M. C., Kojima, H. & Fukui, M. Arsenite oxidation by a newly isolated

Betaproteobacterium possessing *arx* genes and diversity of the *arx* gene cluster in

bacterial genomes. *Front. Microbiol.* **10**, (2019).

92. Price, M. N., Deutschbauer, A. M. & Arkin, A. P. Filling gaps in bacterial catabolic

pathways with computation and high-throughput genetics. *PLOS Genet.* **18**, e1010156

(2022).

93. Alballa, M. & Butler, G. TooT-T: discrimination of transport proteins from non-transport

proteins. *BMC Bioinformatics* **21**, 25 (2020).

94. Kim, G. B., Gao, Y., Palsson, B. O. & Lee, S. Y. DeepTFactor: A deep learning-based

tool for the prediction of transcription factors. *Proc. Natl. Acad. Sci.* **118**, e2021171118

(2021).

95. Barnum, T. P. *et al.* Predicting microbial growth conditions from amino acid

composition. 2024.03.22.586313 Preprint at https://doi.org/10.1101/2024.03.22.586313

(2024).

96. R Core Team. R: a language and environment for statistical computing. R Foundation

for Statistical Computing (2023).

97. Brooks, M. *et al.* glmmTMB: generalized linear mixed models using template model

builder. (2025).

98. Revell, L. J. phytools: an R package for phylogenetic comparative biology (and other

things). *Methods Ecol. Evol.* **3**, 217–223 (2012).

99. Parks, D. H. *et al.* A complete domain-to-species taxonomy for Bacteria and Archaea.

Nat. Biotechnol. **38**, 1079–1086 (2020).

100. Coleman, G. A. *et al.* A rooted phylogeny resolves early bacterial evolution. *Science*

372, eabe0511 (2021).

101. Louca, S. castor: Efficient Phylogenetics on Large Trees. (2024).

102. Blomberg, S. P., Garland JR., T. & Ives, A. R. Testing for phylogenetic signal in

comparative data: behavioral traits are more labile. *Evolution* **57**, 717–745 (2003).

103. Zeileis, A., Köll, S. & Graham, N. Various versatile variances: an object-oriented

implementation of clustered covariances in R. *J. Stat. Softw.* **95**, 1–36 (2020).

104. Bissett, A. *et al.* Introducing BASE: the Biomes of Australian Soil Environments soil

microbial diversity database. *Gigascience* **5**, s13742-016 (2016).

105. National Ecological Observatory Network (NEON). Soil microbe metagenome

sequences. NEON https://doi.org/10.48443/fzzj-g053 (2021).

106. Bahram, M. *et al.* Structure and function of the global topsoil microbiome. *Nature* **560**,

233–237 (2018).

107. Wilhelm, R. C. *et al.* A metagenomic survey of forest soil microbial communities more

than a decade after timber harvesting. *Sci. Data* **4**, 1–8 (2017).

108. Woodcroft, B. J. *et al.* Genome-centric view of carbon processing in thawing

permafrost. *Nature* **560**, 49–54 (2018).

109. Brown, M. V. *et al.* Systematic, continental scale temporal monitoring of marine pelagic

microbiota by the Australian Marine Microbial Biodiversity Initiative. *Sci. Data* **5**, 1–10

(2018).

110. Hugerth, L. W. *et al.* Metagenome-assembled genomes uncover a global brackish

microbiome. *Genome Biol.* **16**, 1–18 (2015).

111. Satinsky, B. M. *et al.* The Amazon continuum dataset: quantitative metagenomic and

metatranscriptomic inventories of the Amazon River plume, June 2010. *Microbiome* **2**,

1–7 (2014).

112. Biller, S. J. *et al.* Marine microbial metagenomes sampled across space and time. *Sci.*

Data **5**, 1–7 (2018).

113. Hu, W. *et al.* Metagenomics unravels differential microbiome composition and

metabolic potential in rapid sand filters purifying surface water versus groundwater.

Environ. Sci. Technol. **54**, 5197–5206 (2020).

114. Singleton, C. M. *et al.* Connecting structure to function with the recovery of over 1000

high-quality metagenome-assembled genomes from activated sludge using long-read

sequencing. *Nat. Commun.* **12**, 1–13 (2021).

115. Bandla, A., Pavagadhi, S., Sridhar Sudarshan, A., Poh, M. C. H. & Swarup, S. 910

metagenome-assembled genomes from the phytobiomes of three urban-farmed leafy

Asian greens. *Sci. Data* **7**, 1–7 (2020).

116. Mendes, L. W., Raaijmakers, J. M., de Hollander, M., Mendes, R. & Tsai, S. M.

Influence of resistance breeding in common bean on rhizosphere microbiome

composition and function. *ISME J.* **12**, 212–224 (2018).

117. Xu, J. *et al.* The structure and function of the global citrus rhizosphere microbiome. *Nat.*

Commun. **9**, 1–10 (2018).

118. Howe, A. *et al.* Seasonal activities of the phyllosphere microbiome of perennial crops.

Nat. Commun. **14**, 1039 (2023).

119. Levy, A. *et al.* Genomic features of bacterial adaptation to plants. *Nat. Genet.* **50**, 138–

150 (2018).

120. Olson, D. M. & Dinerstein, E. The global 200: priority ecoregions for global

conservation. *Ann. Mo. Bot. Gard.* **89**, 199–224 (2002).

121. Bivand, R. S., Pebesma, E. & Gomez-Rubio, V. *Applied Spatial Data Analysis with R,*

Second Edition. (Springer, NY, 2013).

122. Bivand, R. *et al.* rgdal: bindings for the 'geospatial' data abstraction library. (2022).

123. Bivand, R. & Rundel, C. *Rgeos: Interface to Geometry Engine - Open Source ('GEOS').*

(2020).

124. Matsen, F. A., Kodner, R. B. & Armbrust, E. V. pplacer: linear time maximum-

likelihood and Bayesian phylogenetic placement of sequences onto a fixed reference

tree. *BMC Bioinformatics* **11**, 538 (2010).

125. Czech, L., Barbera, P. & Stamatakis, A. Genesis and Gappa: processing, analyzing and

visualizing phylogenetic (placement) data. *Bioinformatics* **36**, 3263–3265 (2020).

126. Woodcroft, B. J., Boyd, J. A. & Tyson, G. W. OrfM: a fast open reading frame predictor

for metagenomic data. *Bioinformatics* **32**, 2702–2703 (2016).

127. Genuer, R., Poggi, J.-M. & Tuleau-Malot, C. VSURF: variable selection using random

forests. (2022).

128. Breiman, L., Cutler, A., Liaw, A. & Wiener, M. randomForest: Breiman and Cutler's

random forests for classification and regression. (2022).

129. Molnar, C., Bischl, B. & Casalicchio, G. iml: an R package for interpretable machine

learning. *JOSS* **3**, 786 (2018).

1045

 Figure captions

 Fig. 1: Comparative genomics indicates denitrifier types are unequally common and

 dominated by distinct phyla. **a** Overview of experimental approach for comparative

 genomics. Metagenome-assembled, single cell and isolate genomes covering a broad range of

 environmental, host-associated, and engineered biomes were extracted from GTDB v.214.1 [99]

 (**Supplementary Data 2**). After quality filtering the genomes, denitrification genes were

 searched using a combination of HMM-based searches, phylogeny construction, and manual

 inspection of alignments. Denitrifier genomes were then searched for additional functional

 traits. Maximum growth rate was inferred based on codon use bias in highly expressed

 ribosomal proteins, while transcription factors were detected using DeepTFactor, and

 potential to use various organic and inorganic electron donors and acceptors with a range of

 tools. **b** Venn diagram of prevalence of denitrifier types. The number inside each circle

 denotes the percentage of denitrifier genomes with that genotype. Letters in italics denote

 whether genotype is considered an "initiator" (*i*), "terminator" (*t*) or both (*it*). Reactions

 completed by each enzyme and corresponding denitrifier type name are listed below.

 Genomes encoding *nor* without accompanying *nir* or *nosZ* genes are marked with an asterisk

 and classified as belonging to non-denitrifiers. **c** Stacked bar charts denoting relative

 proportion of genomes belonging to phyla within each denitrifier type, with non-denitrifiers

 defined as containing neither *nir* nor *nosZ*, or being archaeal nitrifiers or anammox bacteria

 (see methods). The number of genomes corresponding to each denitrifier type is indicated

 above the corresponding bar. **d** Proportion of genomes within dominant phyla encoding each

 denitrification enzyme gene, among denitrifiers. Numbers to the right in panel d indicate the

 number of denitrifier genomes belonging to each phylum denoted in graph. Phyla represented

1070   by fewer than 100 denitrifier genomes are aggregated under "other" and can be found in

1071   **Supplementary Data 11**.

1072

Fig. 2: Genome-inferred traits of complete and partial denitrifier bacteria. a Estimated

maximum growth rate of organisms based on codon usage bias. Stacked bar charts show the

maximum predicted growth rate for genomes separated by denitrifier type (upper panel) and

NosZ clade (lower panel). Slow-growing taxa (maximum growth rate < 0.2 h^{-1}) are depicted

to the left on the horizontal axis and fast-growing taxa to the right. Numbers to the right

denote number of genomes used in maximum growth rate predictions and exclude genomes

with fewer than 10 ribosomal proteins annotated. **b** Boxplots of sugar and organic acid

preference of denitrifiers, based on KEGG annotation ratios[32]. **c** Genomic density of

transcription factors inferred using DeepTFactor. **d** Boxplot of substrate use counts by

denitrifier type and substrate category based on GapMind, with colors following panels b-c. **e**

Heatmap of terminal electron acceptors and donors, with intensity of color and number

denoting proportion of genomes within denitrifier type encoding redox ability. In panels b-d,

box boundaries represent first and third quartiles, with midline denoting median. Whiskers

denote the 1.5 IQR, and outliers are shown as individual points. Asterisks above boxes

indicate a difference between complete denitrifiers and the partial denitrifier type and no

comparisons between partial denitrifier types were made. Numbers to the right in panel e

denote number of genomes used in panels b-e.

1090

Fig. 3: Balance in prevalence of denitrification genes shows dominance of initiators at

the community level across global biomes. **a, b** Boxplots showing difference in *nos* versus

nir counts (($nosZ$ I + $nosZ$ II) − ($nirK$ + $nirS$)) (**a**); and clade I and clade II *nosZ* counts (**b**)

per Gb sequenced. Biomes were compared using Benjamini-Hochberg FDR-corrected

pairwise ranked comparisons following Kruskall-Wallis, and common letters to the right in

each boxplot denotes biomes with similar median gene prevalences. Numbers to the right

denote the number of metagenomes included for each biome. Box boundaries represent first

and third quartiles, with midline denoting the median and whiskers the 1.5 IQR. **c, d** Mean

phylum level composition of clade I (**c**) and clade II *nosZ* reads (**d**), organized by biome.

Biomes represented by fewer than 20 metagenomes are excluded from the figure. The

percentage of genomes having clade II *nosZ* and a complete denitrification pathway are

indicated for each phylum in the legend and can be seen for both clades in **Supplementary**

Fig. 2. Standard errors corresponding for each phylum are available in Supplementary Data 6.

Host assoc. = host-associated; Engineer. = engineered.

1105

Fig. 4: Environmental predictors of balance between *nos* and *nir* counts using random

forest models. a, b Abiotic predictors of the difference in *nos* and *nir* gene counts, calculated

as ($nosZ$ I + $nosZ$ II) − ($nirK$ + $nirS$) and normalized per Gb sequenced in soil (**a**) and marine

metagenomes (**b**). Accumulated local effects plots show the differences in prediction of the

δ*nos-nir* (y-axis) compared to the mean prediction along the range of each predictor (x-axis),

while accounting for potential correlations among predictor values. Values above zero

indicate the model predicts higher than average dominance of *nosZ* over *nir* at a given value

of the predictor variable. The analyses were performed on a subset of the soil (n = 298) and

marine (n = 93) metagenomes for which relevant environmental metadata was available (see

methods). Root mean square error and variance explained were 37 and 29% in the soil (**a**)

and 20% and 20% in marine (**b**) models, respectively. Vertical marks on the x axis in each

panel denote data density. Five hundred trees were run, and mtry, nodesize, and sampsize

parameters were 3, 10, and 176 for soil and 3,10, 88 for the marine models, respectively.

1119

Fig. 5: Environmental predictors of balance between *nosZ* clade I and clade II using random forest models. a, b Abiotic predictors of the difference in *nosZ* clade I and II gene counts normalized per Gb sequenced in soil (**a**) and marine metagenomes (**b**). Accumulated local effects plots show the differences in prediction of the $\delta nosZ$I-*nosZ*II (y-axis) compared to the mean prediction along the range of each predictor (x-axis), while accounting for potential correlations among predictor values. Values above zero indicate the model predicts higher than average dominance of *nosZ* clade I over *nosZ* clade II at a given value of the predictor variable. The analyses were performed on a subset of the soil (n = 298) and marine (n = 93) metagenomes for which relevant environmental metadata was available (see methods). Root mean squared error and variance explained were 40 and 59% in the soil (**a**) and 14 and 33% in marine (**b**) models, respectively. Vertical marks on the x axis in each panel denote data density. Five hundred trees were run, and mtry, nodesize, and sampsize parameters were 4, 6 and 209 for soil and 3,6, and 88 for the marine models, respectively.

Fig. 6: Balance in prevalence of denitrification gene transcripts shows dominance of initiators at the community level in terrestrial and aquatic biomes. a δnos-*nir* and $\delta nosZ$I-*nosZ*II normalized per Gbp mRNA sequenced across different biomes. Numbers to the right denote number of metatranscriptomes included in each category. **b** Mean and 95 % confidence intervals of spearman correlation coefficients between δnos-*nir* or $\delta nosZ$I-*nosZ*II and environmental factors in aquatic metatranscriptomes. Coloring follows panel **a**. Circle size and numbers to the right denote the number of studies the effect size is calculated from. Soil samples with root exudate addition are categorized under rhizosphere in **a**. Box boundaries represent first and third quartiles, with midline denoting median and whiskers the 1.5 IQR. Values for all biomes can be found in **Supplementary Data 12**.

a

slow growth → rapid growth

Denitrifier type

Initiator	5476
Terminator	2683
Init-term	412
Complete	2499

NosZ clade

Clade I	2192
Clade II	3413

Proportion of genomes

Doubling time

< 0.2 hr^{-1}	0.2-1 hr^{-1}
1-2 hr^{-1}	> 2 hr^{-1}

b

Sugar-acid preference

c

Transcription factors (Mbp^{-1})

Initiator   Terminator   Init-term   Complete

d

Substrates (Mbp^{-1})

Sugars   Amino acids   Organic acids   Total

Substrate

e

	Halogens	Tetrathionate	Elemental S	Sulfite	Sulfate	Iron	Hydrogen	Arsenate	Ammonium	Sulfide	Sulfite	Tetrathionate	Thiosulfate	Iron	Hydrogen	Arsenite	
Initiator	0	0.01	0.02	0	0.03	0.02	0.01	0.01	0	0.01	0.2	0.01	0.06	0.07	0.13	0.04	5500
Terminator	0.01	0.02	0.09	0.02	0.03	0.01	0.02	0.02	0	0.03	0.1	0.01	0.05	0.03	0.22	0.03	2694
Init-term	0.01	0	0.03	0	0.02	0	0	0.02	0	0.03	0.14	0	0.04	0.03	0.09	0.04	416
Complete	0	0.09	0.03	0	0.07	0.02	0.01	0.01	0	0.06	0.29	0	0.18	0.07	0.22	0.08	2516

Reduction   Oxidation

Genome proportion

0.0   0.1   0.2

a

Category	Site	δ*nos-nir* label
Engineer.	Drinking water treatment	m
	Sewage	e
	Municipal WWTP	efg
Host assoc.	Root	kl
	Endosphere	hi
	Leaf	d
Aquatic	Rivers & Streams	ghi
	Ponds & Lakes	fgh
	Geysers	cd
	Marine mats	a
	Trades	b
	Westerlies	c
	Vents	ghi
	Marshes	cd
Terrestrial	Deserts & Xeric Shrublands	k
	Croplands	lm
	Tundra	ef
	Boreal Forests & Taiga	hi
	Temperate Conifer Forests	i
	Tropical & Subtropical Dry Broadleaf Forests	klm
	Tropical & Subtropical Moist Broadleaf Forests	jk
	Temperate Broadleaf & Mixed Forests	i
	Mediterranean Forests Woodlands & Scrub	gh
	Temperate Grasslands Savannas & Shrublands	m
	Tropical & Subtropical Grasslands Savannas and Shrublands	ij

b

Site	δ*nosZ*I-*nosZ*II label	count
Drinking water treatment	d	39
Sewage	a	78
Municipal WWTP	m	71
Root	i	250
Endosphere	hij	21
Leaf	a	181
Rivers & Streams	k	68
Ponds & Lakes	de	120
Geysers	jk	23
Marine mats	k	23
Trades	c	384
Westerlies	b	716
Vents	hij	23
Marshes	m	43
Deserts & Xeric Shrublands	ij	107
Croplands	l	134
Tundra	fg	87
Boreal Forests & Taiga	ef	83
Temperate Conifer Forests	hi	263
Tropical & Subtropical Dry Broadleaf Forests	ghi	24
Tropical & Subtropical Moist Broadleaf Forests	ij	84
Temperate Broadleaf & Mixed Forests	gh	542
Mediterranean Forests Woodlands & Scrub	ij	84
Temperate Grasslands Savannas & Shrublands	jk	183
Tropical & Subtropical Grasslands Savannas and Shrublands	ijk	42

x-axis (a): δ*nos-nir*
x-axis (b): δ*nosZ*I-*nosZ*II

c — x-axis: *nosZ*I copies per Gb

d — x-axis: *nosZ*II copies per Gb

Phylum

Acidobacteriota (8%) · Actinomycetota (9%) · Chloroflexota (9%) · Desulfobacterota (0%) · Gemmatimonadota (24%) · Myxococcota (34%) · Pseudomonadota (80%) · Other (28%)

a

Biome

- Rhizosphere — 64
- Rivers and Streams — 48
- Freshwater Sediment — 24
- Marine — 185
- Soil — 92

$\delta nos\text{-}nir$

$\delta nosZ\text{I-}nosZ\text{II}$

b

Environmental variable

- $(NO_3^- + NO_2^-)$ — 7
- Chlorophyll — 7
- Bacterial Production — 4
- Phosphate — 5
- Oxygen — 3

Mean ρ