Physiologically-relevant light exposure and light behaviour in Switzerland and Malaysia

Authors

Anna M Biller*^1,2 [0000-0002-3673-8838]

Johannes Zauner*1,2 [0000-0003-2171-4566]

Christian Cajochen3 [0000-0003-2699-7171]

Marisa A Gerle3 [0009-0000-9159-1093]

Vineetha Kalavally4 [0000-0002-3531-9076]

Anas Mohamed4 [0000-0001-5418-4501]

Lukas Rottländer3 [0009-0007-9062-0113]

Ming-Yi Seah4 [0009-0005-5409-5839]

Oliver Stefani5 [0000-0003-3854-3222]

Manuel Spitschan1,2,6 [0000-0002-8572-9268]

* Equal contributions
^ Corresponding author

Affiliations

Technical University of Munich, TUM School of Medicine and Health, Department Health and Sport Sciences, Chronobiology & Health, Munich, Germany

Translational Sensory & Circadian Neuroscience, Max Planck Institute for Biological Cybernetics, Tübingen, Germany

Centre for Chronobiology, University Psychiatric Clinics Basel (UPK), Basel, Switzerland

Monash University Malaysia, Electrical and Computer Systems Engineering, Intelligent Lighting Laboratory, Malaysia

Lucerne School of Engineering and Architecture, Lucerne University of Applied Sciences and Arts, Horw, Switzerland

TUM Institute for Advanced Study (TUM-IAS), Technical University of Munich, Garching, Germany

Abstract

Light synchronises the internal clock with the external light-dark cycle. Keeping this alignment benefits health and prevents diseases. Quantifying light exposure is, therefore, vital for effective prevention. Since light exposure depends on photoperiod, culture, and behaviour, we investigated objective light exposure and individual light-related behaviour in Switzerland and Malaysia. In this observational field study, participants (N=39) wore a calibrated melanopic light logger at chest level for 30 consecutive days. At baseline and study end, the Pittsburgh Sleep Quality Index was assessed, and every 3 to 4 days, the Light Exposure Behaviour Assessment (LEBA) was filled. Our pre-registered analyses reveal that participants in Switzerland experienced brighter days (+3.16 times the average mEDI) and spent more time (x1.9 times the duration) in daylight levels per hour of daylight, had ~1.5h later bright light exposure in the afternoon, and stayed over 1h longer in dim light conditions before bedtime. LEBA scores did not differ between Malaysia and Switzerland, and LEBA items were stable over time. LEBA items also correlated with objective light exposure variables in Switzerland but not Malaysia, with a medium effect size (range of absolute r=0.32-0.48). These results highlight cultural and geographical differences in light exposure. We showed that LEBA can be related to actual light exposure and is ecologically informative, but this varies by culture.

Keywords: Light Exposure; Cultural Characteristics; Lighting; Data Collection; Photometry; Wearable Technology; Radiometry; Subjective Exposure Assessment; Objective Exposure Assessment; Light Interaction

Introduction

Light is the main entrainment stimulus (zeitgeber) for synchronising human circadian rhythms to the natural light-dark rhythm of daylight given by Earth's rotation (e.g, Foster et al., 2020; Kuhlman et al., 2018; Roenneberg & Klerman, 2019; Roenneberg & Merrow, 2016). A predictable and strong zeitgeber is key for a functioning sleep-wake rhythm and for human physiology to best deal with daily demands and stressors (e.g., Dijk & Archer, 2009; Foster, 2020; Windred, Burns, Lane, et al., 2024; Windred, Burns, Rutter, et al., 2024). Since the invention of electrical light in the 19[th] century, the daily pattern of light exposure has rapidly altered from this robust light-dark cycle characterised by bright days and dark nights to a less predictable one, resulting from a potentially incalculable combination of artificial lights in the built environment and behavioural interactions with natural light and electrical light sources (Biller, Balakrishnan, et al., 2024). Additionally, bright days become dimmer due to a switch from daylight to weaker electrical sources indoors, while dark nights are brightened by electrical light (artificial light at night; ALAN). Factors that drastically influence natural exposure to daylight on a societal level are, to a large degree, the level of industrialisation and access to electricity (Pilz et al., 2018), and consequently, working hours such as shift work (Boivin et al., 2022).

Disrupting the entrainment of endogenous circadian rhythms by shifted or disordered light exposure can have various serious health consequences, such as disrupted sleep, shift work disorder, metabolic disorders, cardiovascular diseases, mood disorders, impaired immune system, or certain cancers (e.g., Ansu Baidoo & Knutson, 2023; Fishbein et al., 2021; Lunn et al., 2017; Neves et al., 2022; Roenneberg & Merrow, 2016). Recognising these negative consequences has led to a recent consensus statement on the optimal light exposure levels to best support physiology, sleep, and wakefulness in healthy adults (Brown et al., 2022). This statement was strengthened through a concerted call to apply these recommendations in policy-making, lighting design, and preventative measures for health and safety (Kervezee et al., 2024). Analysing light exposure is, therefore, the necessary basis for restoring healthy, strong, and reliable circadian rhythmicity to both mitigate the negatives and increase the benefits.

However, the current expert recommendations by Brown and colleagues, albeit a beneficial and important start, especially for public health, are based on isolated laboratory lighting conditions and do not factor in individual light *behaviour* (Biller, Balakrishnan, et al., 2024). Photic history (Chang et al., 2013; Hébert et al., 2002; Smith et al., 2004) and an individual's spectral diet (Webler et al., 2019) – the timing and spectral composition of prior light exposure – also play a role in subsequent sensitivity to light. Recommendations for healthier light exposure thus also need to factor in individual light exposure during day and night and how they interact with light sources. At present, there is minimal data about real-world light exposure and how much humans deviate from these target light levels. Previous studies often only

sampled for a limited amount of time (e.g., several days to one or two weeks) and
used wrist-based light loggers from actimeters (e.g., Didikoglu et al., 2023) or devices
worn at spectacle frames (Stefani et al., 2024), the latter being valuable but limited in
usability for more extended wear periods. Novel light logging devices at physiologically
more relevant wearing positions (e.g. chest, eye level) are now available, thus
quantifying ecological light exposure over long periods. One such device is the
SPECCY light logger (Mohamed et al., 2021), a chest-worn wearable light logger that
outputs numerous visual and non-visual metrics of light exposure following relevant
standards from the CIE (International Commission on Illumination) standards via a
spectral reconstruction algorithm (CIE, 2018).

Since individual light exposure-related behaviours also depend on an individual's
location (e.g., photoperiod, climate, temperature), culture, and the built environment,
quantifying light exposure should factor in such behavioural and structural
components (Biller, Balakrishnan, et al., 2024). The Light Exposure Behaviour
Assessment (LEBA) instrument (Siraji et al., 2023) has recently been developed to
quantify individual light-related behaviours. It categorises light exposure-related
behaviours into five broad categories (factors) based on the participant's recall of the
past four weeks: i) wearing blue light filters, ii) spending time outdoors, iii) using
phones and smartwatches in bed, iv) using light before bedtime, and v) using light in
the morning and during daytime. This questionnaire is well suited to map objectively
measured light exposure patterns to individual self-reported behaviours but also lends
itself to studying potential cultural differences in interactions with light.

This project had three objectives. The first objective was to investigate differences in
objectively measured physiologically-relevant light exposure light loggers at two sites
that vary in culture and climate – Malaysia compared to Switzerland. A second
objective was to investigate differences between the sites in subjectively measured
light exposure behaviour using the LEBA questionnaire. The third objective was to
examine which objectively derived light variables correlate with LEBA-derived items
and composite scores and whether this depended on the study location.

Knowing more about the scope of assessment methods and cultural differences in
light exposure and interactions with light can help inform future tailored and public
health interventions to improve and foster healthy light exposure (Biller, Balakrishnan,
et al., 2024).

Methods

The current study was preregistered (Biller, Zauner, et al., 2024). We describe deviations from the preregistration document in the "Deviation from the preregistration" section below.

Research questions and hypotheses

We addressed the following research questions and hypotheses as specified in the pre-registration document:

1. *Are there differences in objectively measured light exposure between Switzerland and Malaysia, and if so, in which light metrics?*
   - **H1:** There are differences in light logger-derived light exposure intensity levels and intensity duration between Malaysia and Switzerland.
   - **H0$_1$:** No differences between Malaysia and Switzerland.

   - **H2:** There are differences in light logger-derived timing of light exposure between Malaysia and Switzerland.
   - **H0$_2$:** No differences between Malaysia and Switzerland.

2. *Are there differences in self-reported light exposure patterns using LEBA across time or between the two sites, and if so, in which questions/scores?*
   - **H3:** There are differences in LEBA items and factors between Malaysia and Switzerland.
   - **H0$_3$:** No differences between Malaysia and Switzerland.

   - **H4:** LEBA scores vary over time within participants.
   - **H0$_4$:** No differences in LEBA scores over time among participants

3. *In general, how are light exposure and LEBA related, and are there differences in this relationship between the two sites?*
   - **H5:** LEBA items correlate with preselected light-logger-derived light exposure variables.
   - **H0$_5$:** No correlation.

   - **H6:** There is a difference between Malaysia and Switzerland in how well light-logger-derived light exposure variables correlate with subjective LEBA items.
   - **H0$_6$:** No differences between Malaysia and Switzerland.

Study design

This observational study took place in Kuala Lumpur, Malaysia, and Basel, Switzerland, and was conducted for one month per participant at each site. Participants were not randomly assigned to treatment nor blinded in any way and were

allowed to live their regular lives as usual. Day 0 and day 31 involved the collection
and return of equipment, while days 1 – 30 involved the collection of light exposure
and questionnaire data from the participants (**Fig. 1A**). Light data were collected via a
neck-worn portable light logger (SPECCY). Questionnaires were deployed online via
Qualtrics (Malaysia) and REDCap (Switzerland) (Harris et al., 2009, 2019)

– Insert Figure 1 here –
181
On day 0, participants collected the light logger and filled in the Pittsburgh Sleep
Quality Index (PSQI) questionnaire. The light logger was worn around the neck at
chest level, approximately 20 cm below the wearer's chin, with the sensing area facing
outwards towards the sight of light arriving on the wearer's eyes. From day 1 to day
14, participants wore the light logger continuously except when exposed to high
amounts of water (e.g., bath, swimming) or when the dosimeter was being charged.
Participants were asked to charge the sensor daily for an hour. The participants chose
this charging time, but it was recommended to the participants that it remained
consistent throughout the experiment and preferably at a time when they were
primarily sedentary. Participants were also instructed to ensure that the sensor area
of the light logger was not covered by any obstacles or objects (e.g., hands, hair,
seatbelts, etc.). No specific instructions were given on light logger placement while
participants were asleep. When the light logger was not worn, participants were
instructed to place it so that its sensor area was blocked and could not detect any light.
Participants were asked to answer the LEBA questionnaire twice a week before going
to bed at night. On day 15, the light exposure data recorded by the light logger was
downloaded from the device, and on day 31, participants returned the device, after
which the collected light exposure data were downloaded. Data collection was
completed after participants answered the Pittsburgh Sleep Quality Index (PSQI)
questionnaire.

Inclusion and exclusion criteria
In total, 20 (Switzerland) and 19 (Malaysia) participants were recruited to participate
in the data collection experiment with the following inclusion criteria:

1. Age between 19 and 65 years old
2. No history of drug abuse
3. Caffeine consumption ≤2 cups per day
4. Alcohol consumption <2 drinks per day or 14 drinks a week
5. Smoke <2 cigarettes per day

Data collection procedures
Data collection took place at the Psychiatric University Clinics (UPK), Basel,
Switzerland 47.5714° N, 7.5677° E) and at Monash University, Kuala Lumpur, Malaysia
(3.0650° N, 101.6009° E), but participants were allowed to roam freely (**Fig. 1B**). Data
collection in Switzerland was carried out between July and October 2023
(summer/autumn), while data collection in Malaysia was carried out between

November and December 2023 (**Fig. 1C**). Photoperiod change during the collection
period was considerable in Switzerland, but not Malaysia (**Fig. 1D**). Participants
stayed within a 50 km radius of the respective sites.

For participants in Malaysia, applicants were asked to answer questions related to the
inclusion/exclusion criteria using Google Forms. A total of 50 applicants were
registered. Fourteen applicants did not meet the inclusion criteria, and the remaining
applicants were either accepted, declined participation, or not contacted due to the
maximal sample size being reached. One person was later excluded from the analysis
due to a lost SPECCY device, leading to a final sample size of 19 participants in the
Malaysian dataset. Qualtrics links to the LEBA and PSQI questionnaires were sent to
the participants according to their schedules, and they would fill in the questionnaires
via the link.

Participants in Switzerland were recruited through an online advertisement posted on
the University's bulletin board. The advertisement included a brief description of the
study and inclusion and exclusion criteria. Those who expressed interest in the study
were contacted via email to verify their eligibility. Upon confirmation, they were
provided with a detailed study information sheet. Eligible and interested applicants
were scheduled to visit the lab on a Monday. Participants were briefed in person about
the study and registered using Qualtrics. Participants also completed the PSQI on-site
using their mobile phones and received instructions on how to use the light logger.

Wearable light logger

A wearable light logger, SPECCY, was used for data collection in Malaysia and
Switzerland. The light logger, modified from the light spectral sensor developed by
Mohamed et al. (2021) and developed at Monash University Malaysia's Intelligent
Lighting Laboratory (MMILL), captures spectral information across the visible range
between 380 and 780 nm. The Australian Photometry and Radiometry Laboratory
validated SPECCY, which has an effective measuring range from 1 lux to 130,000 lux.
The light logger system is constructed from three multi-channel sensors, which provide
14 optical sensing channels within the measurement range of 380 nm and 780 nm and
four channels in the infrared (IR) range. These channels capture low-resolution
spectral data through proprietary software layers adjusted for sensor saturation and
baseline signal calibration. Incorporated within a compact printed circuit board (PCB),
the device houses the three multi-channel sensors, optimally placed for temperature
consistency, an ambient temperature sensor, a microprocessor enabling Bluetooth
Low Energy (BLE) connectivity, 16 MB of onboard flash storage allowing for over
130,000 spectral measurements, vital indicators and controls such as LEDs and a
connectivity toggle button, and a micro-USB charging port. These components are
securely encased within a 3D-printed housing made of polylactic material. A cosine
corrector built into the case ensures appropriate sensor directionality. The output
measurements are consistent with the CIE S 026/E:2018, including melanopic
equivalent daylight illuminance (CIE, 2018).

Surveys

LEBA

The Light Exposure Behaviour Assessment instrument (LEBA; Siraji et al., 2023) was developed to examine how light exposure behaviours affect health and well-being. LEBA categorises five key behaviours: wearing blue light filtering glasses indoors and outdoors (Factor 1), spending time outdoors (Factor 2), using phones and smartwatches in bed before sleep (Factor 3), controlling and using ambient light before bedtime (Factor 4), and using light in the morning and during daytime (Factor 5). **Table S1** lists detailed computations of these factors.

PSQI

The Pittsburgh Sleep Quality Index (PSQI; Buysse et al., 1989) is a self-report questionnaire that assesses sleep quality over one month. It measures seven components, including sleep duration, disturbances, and daytime dysfunction, and produces a global score. A higher global score (≥5) indicates poor sleep. **Table S2** lists details on the computation of components and global scores.

Sample size calculation

No specific sample size was calculated for the study. The sample was based on convenience sampling and resource limitations. No specific stopping rule was implemented.

Measured variables

SPECCY-derived variables

Light data from the SPECCY device (melanopic EDI) were analysed using the R package *LightLogR v0.4.2* (Zauner et al., 2024) to perform cleaning steps, create visualisations, and calculate derived light exposure metrics. Melanopic EDI, the melanopic equivalent daylight illuminance (mEDI) measured in lux, measures how strongly light affects the melanopsin system, which is the primary driver of the non-image-forming pathways in humans (Brown et al., 2022). An overview of each calculated light metric is given in **Table S3.** The exact calculation equation for each metric is contained within the open-source code of LightLogR (https://github.com/tscnlab/LightLogR).

LEBA-derived variables

The Light Exposure Behaviour Assessment instrument (LEBA; Siraji et al., 2023) captures light exposure-related behaviours on a 5-point Likert-type scale ranging from 1 to 5 (1 = never; 2 = rarely; 3 = sometimes; 4 = often; 5 = always). The score of each factor is calculated by the summation of scores of items belonging to the corresponding factor (**Table S1**). Respondents are requested to respond to each item retrospectively to capture their propensity for different light exposure-related behaviours. Originally, this covers the past 4 weeks but was amended in the current

study to "past 3-4 days" (asked twice per week) for the Swiss site (the original framing of 4 weeks was kept for the Malaysian site). A list of all LEBA items is given in **Table S5**.

Composite scores

For the LEBA, five factors were calculated based on individual LEBA items (**see Table S1**). In scoring the PSQI, seven component scores are derived, each scored 0 (no difficulty) to 3 (severe difficulty). The component scores are summed to produce a global score (range 0 to 21). Higher scores indicate worse sleep quality, with a score >5 suggesting significant sleep difficulties.

Transformations of variables and preprocessing steps

LEBA-derived variables

Item 4 of the LEBA questionnaire was reversed scored for the F2 calculation (see section "Measured variables" for more information on items and factors of the LEBA). LEBA scores were stored in a dummy-coded way in the Switzerland dataset (exported from RedCap), where 1 to 5 encode "Never", "Rarely", "Sometimes", "Often", and "Always", respectively. Answers were directly coded in the Malaysia dataset (exported from Qualtrics). Upon import, all LEBA scores were converted to dummy-coded factors as described for the Switzerland dataset.

Melanopic EDI-derived variables

The base measurement was the time series of melanopic EDI (in lux). Several summary metrics are calculated from that measure (see **Table S3**). The period over which these metrics are calculated varies depending on the specific research question, hypothesis, and metric. Generally, metrics were calculated per participant and day, except interdaily stability (IS), which was calculated per participant. Whenever melanopic EDI was directly used in a statistical model, it was logarithmically transformed (base 10).

Time-of-Day

For some research questions, the distinction of daytime vs. evening or nighttime was relevant when calculating metrics, as either time of day was part of the statistical model or only a portion of the day was relevant. In those cases, metrics were calculated based on a filtered time series of melanopic EDI. These filters have three thresholds: dusk, dawn, and midnight. Dusk and dawn were calculated based on latitude, longitude (Switzerland: 47.5585° N, 7.5839° E; Malaysia: 3.0650° N, 101.6009° E), and calendar date with the *suntools* package (Bivand & Luque, 2023), with a solar depression angle of 6°, which yields civil dusk and dawn. Dawn to dusk is considered daytime, dusk to midnight as evening, and midnight to dawn as nighttime.

Datetimes

Since the two sites were in different time zones, directly including the datetime variable

in a statistical model or a combined visualisation was not sensible since we were
interested in differences depending on local time, not real-time differences due to
different time zones. When only the local clock time (not the date) was of interest,
datetime was transformed into seconds from midnight local time. When the local time,
including date, was of interest, datetime was transformed by overwriting the time zone
of both sites to UTC, thus forcing them to operate on the same timeline.
349

Correlating LEBA scores with light exposure metrics
Depending on the site, different timespans were used in the correlation analysis (**H5**)
due to how the items in the questionnaire were framed:
For Switzerland, the LEBA covered behaviour in the current respective period (i.e.,
about the past three or four days), and light exposure metrics were calculated from
the timespans between each LEBA questionnaire. These metrics were correlated with
the respective LEBA scores. For example, LEBA scores from day 7 were correlated
with light exposure metrics from days 5 to 7, as the prior LEBA questionnaire had been
collected at the end of day 4. All other time spans between assessments were
calculated accordingly.
For Malaysia, where the LEBA had asked about the behaviour in the past four weeks,
light exposure metrics were calculated for the entire four weeks of data collection and
correlated with the final round of LEBA scores collected on day 31.
An average value was calculated prior to correlation for light exposure metrics that
were calculated for each day. This avoided a one-to-many comparison (one LEBA
datapoint to many light exposure metric datapoints) that would have skewed
significance tests of the correlation.
367

368 # Descriptive statistics and statistical models
369 We also tested if light exposure and LEBA items were correlated (**RQ3**) by calculating
370 specific light variables using the R package *LightLogR* and correlated them to the
371 respective LEBA item. See **Table S5** for an overview of which light variables were
372 calculated for each LEBA item.
373

Exploratory analyses
We extended our analyses beyond the specified hypotheses described above in a
secondary analysis. Since we expected differences between the two sites regarding
objectively measured light exposure (using the SPECCY light logger; **RQ1**) and
subjectively measured light exposure (using the LEBA questionnaire; **RQ2**)**,** we also
wanted to test this thoroughly in a data-driven way. To this end, we did a cross-
correlation of all available light metrics in *LightLogR* for the objectively measured light
exposure with all available variables and factors from the LEBA questionnaire across
sites.

Inference criteria

Check for assumptions

Appropriate model diagnostics were performed for all statistical tests to ensure linearity of (generalised) relationship, normally distributed residuals and random effects, and homoscedastic residuals. In the case of generalised additive models, the number of knots (k) was tested to ensure sufficient degrees of freedom to adequately capture the nonlinear relationships.

Statistical tests

Table S4 links specific statistical tests to the research questions and hypotheses. Overall, one of five statistical model types was used:

1. (Generalised) linear mixed-effect models for continuous numeric dependent variables, using the lmer() and glmer() functions in the *lme4* package version 1.1-35.5 (Bates et al., 2015). Generally, a Gaussian error distribution was assumed for dependent variables. A Poisson error distribution was used for count data, such as the seconds above a given threshold.

2. Cumulative link mixed-effect models for ordinal dependent variables, using the clmm() function in the *ordinal* package version 2023.12-4.1 (Christensen, 2010).

3. Generalised additive mixed-effect models for non-linear relationships, using the gam() and bam() functions in the mgcv package version 1.9-1 (Wood, 2000)

4. Correlation matrices using Spearman-rank correlation for the correlation of continuous numeric and ordinal variables. These models use the cor() function in the *stats* package version 4.4.2 in base R (R Core Team, 2021) and the rcorr() function in the *Hmisc* package version 5.2-1 (Harrell, 2003).

5. Bootstrapping to determine 95% confidence intervals for variables. Bootstrapping was performed with base R functions such as sample() and the *tidyverse* package version 2.0.0 (Wickham, 2016; Wickham et al., 2019).

Accounting for multiple testing

A false-discovery-rate correction (Benjamini-Hochberg) was applied for every hypothesis based on the number of models/tests performed within that specific hypothesis.

Effect size

For the statistical tests in model types 1, 2, and 3, unstandardised effect sizes are reported as a result of the final models, i.e., beta-coefficients and non-linear dependency curves. For type 4, the correlation coefficients are effect sizes.

Significance levels

For statistical tests in model types 1, 2, and 4, p-values were calculated and corrected for multiple testing. Values equal to or below 0.05 were considered significant. For type 1 and type 2, p-values are calculated through a likelihood ratio test of models with and without a parameter in question. For type 3, model selection/significance was based on Akaike's Information Criterion (AIC), where the more complex model was

preferred if their AIC was two or more below the less complex model. For type 4, p-
values were calculated through the asymptotic t-approximation (Press et al., 1992).
For type 5, 95% confidence intervals were calculated based on 10^4 bootstrapping
samples. If a standard deviation of 0 was not within the confidence interval, the item
variance over time was considered significant.

Photoperiod correction
Figure 1D shows the differences in photoperiod between the two sites during data
collection. The photoperiod, as measured from civil dawn to civil dusk, was, on
average, 12.7 hours in Malaysia during data collection, while it was 14.4 hours in
Switzerland. Furthermore, while there is negligible difference in photoperiod within the
Malaysia site, there is a strong variation within the Switzerland site due to higher
latitude and a longer data collection period (not per participant but for overall data
collection). To account for this, metrics in H1 were corrected by their respective
periods: times above or below a given threshold per day were divided by the
(photo)period relevant for the metric. The resulting metrics should be interpreted such
that per hour of photoperiod, *x* seconds were spent above/below the threshold.
Similarly, this correction was applied for time metrics covering the evening, which were
divided by the evening length.

Data exclusion, outlier handling and awareness checks

Exclusion criteria of data
444 Measurement data were excluded if their values lied outside the measurement
445 equipment's plausible thresholds (≥130,000 lux for photopic illuminance). No
446 awareness checks were implemented during data collection, and outlier handling was
447 not necessary after viewing the data.

Missing data
The missing data percentages (implicit missing data) and percentage data where the
recorded photopic illuminance fell outside the light logger's effective measuring range
between 1 lux and 130,000 lux (explicit missing data) were recorded and analysed.
Values above the upper threshold were few (one data point for Malaysia and eight
data points for Switzerland) and negligible relative to the amount of available data.
Values below 1 lux were regular and expected, as this value indicates darkness.
Values below 1 lux were converted to 0.1 lux, which allowed for their inclusion in
logarithmically transformed models.
All implicit missing data were made explicit in the analysis, i.e., gaps in the time series
were filled with the respective date, and corresponding measurement values were set
to NA. Participant days with more than a fixed threshold of 6 hours of missing data per
day were excluded from further analysis. The threshold was determined by a
sensitivity analysis using three randomly chosen light exposure metrics and three
randomly selected participants without missing data. The threshold was set so that the
average of each metric stayed within a 95% confidence interval based on their original
variation (mean±2 SE) and 10^4 resamples for each threshold. This ensured a threshold

that did not significantly affect metric calculation based on the actual data (see **Fig.**
S1). While the threshold may seem rather large, it exemplifies that for a sufficiently
long time series, i.e. 30 days in this case, the chosen light exposure metrics (IV, last
timing below 10 lux, MP ratio) are rather robust to missing data, even for larger time
frames within a day. For an overview of missing data, see **Table S6**.
470

Deviations from the preregistration

In research question 1, H2 *("There are differences in light logger-derived timing of light*
exposure between Malaysia and Switzerland.") of the preregistration document
specified an additive model to describe light exposure patterns within participants and
between sites. This approach led to the conclusion that the light exposure patterns
were dominantly driven by individual patterns rather than an overarching pattern by
site (see **Results**). To explore differences by site, another model that only included
random intercepts by participant (rather than random smooths by participant) was
constructed. Both models are part of the analysis documentation in the
Supplementary Materials (Section 5.2.2.4), and their results will be discussed
according to their respective relevance.

The preregistration document specified a generalised mixed model approach to test
H6 ("*There are differences between Malaysia and Switzerland on how well light-logger*
derived light exposure variables correlate with subjective LEBA items"). As the
specified formula did not lead to the stated number of models, i.e. 23+5, and also did
not provide an overall overview of whether selected light exposure metrics correlate
with LEBA items and factors, we provide here an alternative approach by analysing
whether there are significant differences in correlation coefficients between the sites.
In this approach, a linear model is used, and the number of models equals the number
specified in the preregistration. The output is also more suitable for answering the
hypothesis. The stricter analysis is still included in section 5.4.2. of the analysis
documentation (link in the supplementary material).

We also had to correct $H0_4$, which originally stated "*No differences between Malaysia*
and Switzerland". However, this comparison is not sensible given that we were
investigating the LEBA scores within the Malaysian dataset over time instead of
investigating if these scores differed over time, dependent on the research site. The
updated $H0_4$ is thus: "*No differences in LEBA scores over time within participants*", as
already stated above at the beginning of the Methods section.

Results

Participant characteristics

The sample in Basel, Switzerland, consisted of five male and 15 female participants (age 31.35 ± 9.66 years old), and all participants completed their data collection successfully. In Malaysia, 8 males and 11 females (age 24 ± 6.74 years old) completed data collection successfully, and one participant failed to complete the experiment due to the loss of the light logger. PSQI scores were comparable between sites both on day 1 and day 31 and did not differ significantly (**Table S13-14**).

Individual differences in light exposure

Fig. 2. shows individual light profiles from both sites averaged across the recording period. While the results below go into detail about condensed quantifiers of light exposure metrics, individual patterns lead to valuable insights into the wide breadth of light exposure patterns and their variation. To highlight just a few, some participants, such as ID 16 or 18 in the Swiss dataset, show very *small deviations* from their daily light exposure patterns as indicated by narrow 95-, 75-, and 50-percentile ribbons. On the other hand, participants 15 and 17 in the Swiss dataset or participants 08, 13 or 16 in the Malaysian dataset show *strong deviations* from their median daily patterns. Interestingly, this is somewhat captured by the respective interdaily stability (IS) measures but not as much as expected. In general, the IS values are also rather small (all <0.3), indicating strong variation between days (higher numbers mean more stability).

Furthermore, the data also show a degree of within-day variation: while participant 08 in Malaysia shows a clear tendency for predominant nighttime exposure, there are regular "daytime patterns" visible in the 95- and 75-percentile ribbons. Participant 17 in Switzerland, on the other hand, has a clear daytime exposure profile but with regular evening and nighttime extensions and variations.

– Insert Figure 2 here –

Participants in Switzerland spent more time in daylight, experienced brighter days and stayed longer in recommended light levels during the day compared to participants in Malaysia

Fig. 1E shows the grand average of light exposure profiles for the site in Switzerland (yellow line) and the site in Malaysia (blue line). The model summary (**Table S7**) shows that participants in Switzerland spent significantly more time above threshold per hour of daylight for 250 lux (TAT250) and 1000 lux (TAT1000) than participants in Malaysia, i.e., x1.78 and x1.94, respectively (x1.77 over 250 lux melanopic EDI during daytime hours, TATd250). The second model summary (**Table S9**) also shows that the 10 brightest hours (M10m) of participants in Switzerland are significantly brighter than for participants in Malaysia, i.e., 661 lux and 229 lux, respectively. The frequency of crossing 250 lux is about twice as high for participants in Switzerland compared to

Malaysia (64 absolute times compared to 36 times, respectively). Lastly, participants in Switzerland experienced light levels of above 250 lux melanopic EDI about 1.5 hours longer across the day, compared to Malaysia (19:09 vs 17:41 respectively, **Table S9**) and about 2.5 times higher melanopic EDI illuminance levels daily (437 lux vs 170 lux respectively) as demonstrated in the interaction of site and time of day (**Table S8**).

Participants in Switzerland experienced earlier dim light exposure and darker evenings

Participants' last time above 10 lux after dusk (LLiT10) in Switzerland was about 1 hour earlier than participants in Malaysia (22:05 vs 23:16, respectively). There was no difference ($p<.9$) in time below threshold of 10 lux during the evening (TBTe10) between sites (**Table S7**). Exploring the interaction of site and time of day (day vs evening) in illumination levels revealed an average for participants in Switzerland of 3.5 lux during evening hours and a 5-fold increase (17.6 lux) for participants in Malaysia (**Table S8**).

Modelled light exposure patterns vary between sites, with strong differences between individuals

Fig. 3 shows model results when estimating light exposure patterns, with **Fig. 3A** showing individual patterns, **Fig. 3B** patterns by site, and **Fig. 3C** differences between the sites. While each site shows a characteristic pattern (**Fig. 3B**), these are composed of strongly varied individual patterns within each site (**Fig. 3A)**. In fact, when modelling each participant's pattern, an overall site-specific pattern does not significantly increase the model's explanatory power. This is indicative of strong differences between participants within each site. Overall, the light exposure patterns are consistent with the results derived from individual light metrics: Participants in Switzerland experience longer and brighter days and darker and earlier nights. While the longer photoperiod in Switzerland might partly explain the daytime effects, it does explain the described night-time effects.

- Insert Figure 3 here -

Despite this strong individual variation, we can still reject $H0_1$ and $H0_2$ of our first research question since we did find statistical differences in light-logger-derived light exposure intensity levels and duration of intensity, as well as timing of light exposure, between Malaysia and Switzerland (see also **Table S10**).

LEBA responses do not differ between Malaysia and Switzerland and are stable over time

To address the second research question, we compared LEBA items and scores across sites and found that under the strict p-value adjustment for H3 (n=28), none of the five LEBA factors or individual items were significantly different between sites (for all: $p >.3$; see section 5.3.1 in the full analysis document linked in supplementary

material). Thus, we cannot reject our $H0_3$ for this research question since we could not
find strong differences in Malaysian and Swiss LEBA scores.
Looking at the stability of LEBA scores across time within the Malaysia site (H4),
bootstrapping analyses showed that scores were very stable over one month. All 23
questions and 1 out of 5 factors did not vary significantly in more than 50% of
participants. We consequently cannot reject $H0_4$ and assume scores to be stable over
time, at least in this dataset (see Table "Average change in LEBA metrics across four
weeks of data collection" in section 5.3.2. of the full analysis document linked in the
supplementary material).
595

596 **LEBA items correlate with preselected light-logger-derived light**
597 **exposure variables in Switzerland but not in Malaysia**
598 While exploratory analysis shows correlations in both sites, only Switzerland shows
599 significant correlations after correction for multiple testing (**Fig. 4**). Out of 84
600 correlations hypothesised *a priori*, 10 were significant in Switzerland.
601

602 – Insert Figure 4 here –
603

604 The effect size of those correlations was medium on average (r = 0.39), and all
605 correlations showed the hypothesised direction (**Table 1**). We can thus reject $H0_5$
606 since we did indeed observe statistically significant correlations between LEBA and
607 light-logger-derived variables
608

609 – Insert Table 1 here –
610

There were also a few LEBA questions where the correlation coefficients differed
significantly between sites. Specifically, these were the questions *"I dim my mobile*
phone screen within 1 hour before attempting to fall asleep" (F4 item 15) and *"I dim*
my computer screen within 1 hour before attempting to fall asleep" (F4 item 18; **Fig.**
4). While the correlation is positive with preselected light exposure metrics in Malaysia
(r= .25 both, p=.019), it is zero or negative in Switzerland (r = -.01 and -.15,
respectively, p=.035; **Table S12**). We can, therefore, also reject $H0_6$, given that we
observed differences in the strength and directions of LEBA scores/items and light-
logger-derived variable correlations between Malaysia and Switzerland.
620

Discussion

In this observational study, we investigated light exposure and light exposure behaviour differences in 39 participants in their everyday lives over 30 consecutive days. Participants were recruited in Basel, Switzerland, and Kuala Lumpur, Malaysia, two locations that differ in culture, climate, urban environment and photoperiod. Given the multitude of available light metrics to quantify objective light exposure and a lack of standards which of those metrics are to be reported (Spitschan et al., 2019, 2023), we also investigated which specific light metrics differed between these sites (**RQ 1**).

Objective light exposure is different between Switzerland and Malaysia beyond the differences in photoperiod

We found that participants in Switzerland spent significantly more time in bright levels likely corresponding to daylight (i.e., >1000 lux melanopic EDI) than participants in Malaysia during the day. Participants in Switzerland also stayed longer and about twice as often in recommended light levels during the day (i.e., >250 lux melanopic EDI). This is generally expected, given the longer photoperiod in Switzerland compared to the Malaysian site (see **Fig 1D**). However, these findings still hold when adjusting for the longer photoperiod in Switzerland, indicating differences between the sites. Furthermore, the brightest time of day was significantly brighter for participants in Switzerland compared to participants in Malaysia, with a difference of about 0.5 $\log_{10}$ units, indicating that participants in Switzerland experienced their brightest time of day with an intensity approximately 3.16 times higher. This potentially hints towards differences in behaviour since theoretical access to daylight based on photoperiod was comparable, although shorter photoperiods also offer lower daylight levels. A more likely explanation might be, though, that in both sites, melanopic EDI levels during the day indicate predominantly indoor environments. Still, the significantly higher and longer exposure to bright light in participants in Switzerland suggests better access to daylight while being indoors, compared to Malaysia.

As expected, the last time of day with bright light exposure above 250 lux melanopic EDI was later by about 1.5 hours in Switzerland compared to Malaysia, which can partially be explained by the later dusk in Switzerland (about 0.8 hours), but also by daylight savings time (DST), which was in place in Switzerland during data collection, but not Malaysia. This means that participants in Switzerland experienced light until later in their day, which could contribute to later chronotypes (although the reverse could also be true). Unfortunately, chronotype was not assessed in this study, so we cannot verify this assumption, but previous studies have shown an interaction of chronotype and light exposure as well as sleep timings and light exposure (Wahl et al., 2019; Wright et al., 2013; Zerbini et al., 2020, 2021).

Participants in Switzerland in our study did, however, also avoid light exposure above 10 lux mel EDI significantly earlier than participants in Malaysia by about 1 hour and 10 minutes (i.e., the timing of brighter light exposure in the evening). This means that although they still experienced some light later in their 24-day, it was dimmer than the

light participants in Malaysia experienced before their bedtime. Additionally, participants in Switzerland averaged about 5 times (~0.6 $\log_{10}$ units) lower melanopic EDI levels during evenings, suggesting that participants in Switzerland seem to have actively avoided bright light exposure or preferred dimly lit environments. The latter could be explained by cultural differences in light preference, with a potential preference for brighter and whiter electric light in Asian cultures (Baniya et al., 2015). We cannot provide evidence beyond speculation here as we did not assess participants' lighting preferences.

Objective light exposure patterns reveal strong differences between individuals within the sites but do not follow recommendations for healthy light exposure in either site

Our analysis of light exposure patterns reveals that while, on average, the above-mentioned differences in light exposure hold up, the explanatory power of a single exposure pattern per site is negligible. Instead, each site consists of several individual patterns, with a within-site variation greater than between sites. This is encouraging, as it means people on an individual level have influence over their light exposure pattern beyond their environmental or cultural baseline. Thus, interventions to modify personal light exposure patterns might have a greater chance of success than they would have with very similar individual patterns across each site.

While light exposure was more favourable in the Swiss dataset, neither participants in Switzerland nor Malaysia spent much of their time in bright light above 250 lux melanopic EDI, as recommended by Brown and colleagues (Brown et al., 2022). For participants in Switzerland, it is about 12 minutes per hour of daylight, and 7 minutes for participants in Malaysia, i.e., 20% and 12% of the available daytime, respectively, which corroborates previous findings where participants in Switzerland spent only 14% of their light logger wearing time above 250 lux melanopic EDI during daytime (Stefani et al., 2024). In the evening, the participants in Switzerland in our study reduced their light exposure level earlier to the recommended 10 lux melanopic EDI or below. Still, both sites had times of elevated light exposure levels after dusk.

Self-reported light exposure behaviour does not differ between Switzerland and Malaysia and is stable over time

A second goal of this study was to understand if subjectively reported light exposure behaviour measured with the LEBA instrument showed cultural differences between the two locations (**RQ 2**). Since LEBA is a new instrument, cultural differences, also in specific items or scores of the questionnaire, are yet to be investigated and tested. In contrast to our expectations, LEBA items and factors did not show a significant difference between Malaysia and Switzerland, meaning that light exposure behaviour assessed with the LEBA was comparable across these cultures, at least in our sample. Due to the timeframe overlap in the Malaysian dataset (i.e., participants were asked every few days about the previous *four* weeks), we were also able to investigate if

items and scores were stable over time, which we confirmed for most of the LEBA
items. All 23 questions and 1 out of 5 factors did not vary significantly in more than
50% of participants. While the other factors showed variation over time, it was very
small, with an average sum of 3-to-4-point total differences across nine survey periods.
Unfortunately, we were unable to do this analysis in the Swiss dataset because the
timespans used for the instruction in the LEBA questionnaires were not comparable.
Therefore, we cannot be sure if the stability of LEBA items can be generalised beyond
the Malaysian site, but we would hypothesise so based on the overall lack of
differences between cultures in LEBA items and scores.

718 Self-reported light exposure behaviour is reflected in objective light
exposure but only in participants in Switzerland
Finally, we wanted to test if subjectively reported light exposure behaviour can be
related to objectively measured light exposure and, again, if these relationships might
depend on location (Research Question 3). The results clearly show that there are
strong correlations between certain light exposure metrics and answers to the LEBA
questionnaire, but only in the Swiss dataset. After adjustment for multiple
comparisons, none of the correlations in the Malaysian dataset remained significant.
Interestingly, the average (absolute) correlation was stronger in the Malaysia site than
in Switzerland, indicating a higher variance in either light exposure metrics or LEBA
answers in the Malaysia dataset compared to Switzerland. For the Swiss dataset, ten
of the hypothesised correlations were shown to be significant after adjustment with an
average medium effect size (Cohen, 1988, 1992).

We acknowledge that differences in the sites might also be due to the different
phrasing of the LEBA: whereas it questioned participants about the *past few days* in
Switzerland, it asked about the *past four weeks* in Malaysia. Thus, the differences in
correlative strength might also be due to participants being able to recollect light
exposure-related behaviour better for a few days compared to the last month.

736 Strength and weaknesses
Our study has several strengths and weaknesses. We sampled data from a relatively
small cohort of n=19 and n=20 per site, respectively, but given the longitudinal data of
four weeks per participant, we are confident to have enough data points to rely on
model results. Since we only collected data from one site per country, we cannot be
sure that these sites represent the entire country, but this was also not the overarching
research goal. Unfortunately, we lacked information on chronotype and weekday type
information (work day vs. free day), so we could not compute statistics that would have
included such information. The light logger we used, the SPECCY wearable, also did
not record activity (actimetry data) to save battery, so we could not use this information
to detect the non-wear time of the device.
Our study, though, adds to the novel literature on light exposure *behaviour* and is the
first to relate subjectively assessed light behaviour to objectively recorded light
exposure behaviour. Our data sampling is both longitudinal and takes place at high

frequency, allowing for fine-grained and highly reliable data analysis. By positioning
the light logger at chest level, we were able to better approximate physiologically
relevant light exposure than at wrist positions. Finally, we add to the growing literature
on cultural differences, which need to be factored in for better and fine-tuned
recommendations and provide here a dataset that generated novel hypotheses that
can be tested explicitly in the future.
756

Conclusion

This study offers a detailed investigation into the differences in light exposure and light
exposure behaviour between two distinct geographical and cultural contexts:
Switzerland and Malaysia. Key findings reveal significant differences in objective light
exposure between the two sites. Participants in Switzerland experienced brighter and
longer daylight exposure, even when accounting for photoperiod differences while
avoiding evening light exposure earlier and to a greater extent than participants in
Switzerland. These results suggest behavioural and cultural preferences or
differences in access to daylight, all of which influence individual light exposure
patterns. Additionally, our finding that subjective light exposure behaviour correlates
with objective light metrics only in the Swiss dataset underscores potential cultural or
methodological differences in self-reporting accuracy. Despite this, the LEBA
instrument proved stable over time, offering preliminary evidence for its reliability.

While our study advances the understanding of light exposure behaviours and cultural
nuances, it also highlights several areas for future exploration. Investigating
preferences for electrical light colour and intensity could also clarify the cultural
differences hypothesised in this study. Expanding the sample size and geographic
scope would improve the generalisability of these findings, while further validation of
the LEBA instrument in diverse populations is essential to confirm its cross-cultural
applicability and whether it could be used as an approximation for objectively assessed
light exposure.

From an applied perspective, these findings have implications for designing culturally
informed light exposure guidelines to promote healthier circadian alignment. The
apparent avoidance of evening light exposure in the participants in Switzerland aligns
with recommendations for better sleep and circadian health. In contrast, the
'preferences', or at least exposure patterns, observed among the participants in
Malaysia highlight the need for interventions aimed at reducing evening light exposure.
These could include promoting the use of warmer, dimmer lighting in the evening and
adjusting the timing of brighter light exposure during the day. Such strategies should
consider local preferences and location-specific constraints, such as Malaysia's
daytime heat, which often discourages outdoor activities and exposure during daylight
hours. Future studies should explore how environmental and behavioural interventions
could optimise light exposure for health outcomes across diverse populations and
climates to be feasible in the real world beyond Western locations and in light of global
warming.

In conclusion, this study provides a foundation for understanding the interaction of light
exposure behaviour, culture, and geography, opening new avenues for research and
intervention strategies. By addressing the identified limitations and building on the
strengths of our approach, future research can continue to uncover actionable insights
into the critical role of light exposure in human circadian and sleep health.

Data availability statement

Data and code are available on
https://github.com/tscnlab/BillerEtAl_RSocOpenSci_2025.

Contributions

Conceptualisation: AMB, CC, MS, VK, JZ

Data curation: JZ

Formal analysis: JZ

Funding acquisition: CC, MS, OS, VK

Methodology: AM, LR, MG, MS, MYS, OS, VK

Project administration: CC, OS, VK

Software: JZ

Resources: CC, VK

Supervision: CC, MS, OS, VK

Validation: -

Visualisation: JZ

Writing – original draft: AMB, JZ

Writing – review & editing: AMB, AM, CC, LR, MG, MS, MYS, OS, VK, JZ

Acknowledgements

We thank ETH Zurich and Leading House Asia for awarding the RPG Research Partnership Grant ASEAN Number 072022 ("Shaping global human light behaviour – from chaos to order") to OS and VK.

Competing interests

OS is listed as an inventor on the following patents: "Display system having a circadian effect on humans", US8646939B2; "Projection system and method for projecting image content", DE102010047207B4; "Adaptive lighting system", US8994292B2; "Projection device and filter therefor", WO2006013041A1; "Method for the selective adjustment of a desired brightness and/or colour of a specific spatial area, and data processing device", WO2016092112A1. OS is a member of the Daylight Academy, Good Light Group and Swiss Lighting Society. OS has had the following commercial interests in the last two years (2022–2024) related to lighting: investigator-initiated research grants from SBB (Swiss Federal Railways), Zumtobel, DFS (German air navigation service provider), and Porsche. MS is listed as an inventor in a European Patent application (23159999.4), "System and method for corneal-plane physiologically-relevant light logging with an application to personalised light interventions related to health and well-being". MS is a member of the Daylight Academy and is currently the Speaker of the Steering Committee and Chair of the CIE JTC20. The remaining authors declare no conflicts of interest relevant to this work.

References

Ansu Baidoo, V., & Knutson, K. L. (2023). Associations between circadian disruption
and cardiometabolic disease risk: A review. *Obesity*, *31*(3), 615–624.
https://doi.org/10.1002/oby.23666

Baniya, R. R., Tetri, E., & Halonen, L. (2015). A study of preferred illuminance and
correlated colour temperature for led office lighting. *Light & Engineering*, *23*,
39–47.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects
models using **lme4**. *Journal of Statistical Software*, *67*(1).
https://doi.org/10.18637/jss.v067.i01

Biller, A. M., Balakrishnan, P., & Spitschan, M. (2024). Behavioural determinants of
physiologically-relevant light exposure. *Communications Psychology*, *2*(1),
114. https://doi.org/10.1038/s44271-024-00159-5

Biller, A. M., Zauner, J., Cajochen, C., Gerle, M. A., Kalavally, V., Mohamed, A.,
Rottländer, L., Seah, M.-Y., Stefani, O., & Spitschan, M. (2024). *Light
exposure and light behaviour differences between Malaysia and Switzerland*.
OSF Registries. https://doi.org/10.17605/OSF.IO/53UQX

Bivand, R., & Luque, S. (2023). *suntools: Calculate sun position, sunrise, sunset,
solar noon and twilight*. 1.0.0.
https://doi.org/10.32614/CRAN.package.suntools

Boivin, D. B., Boudreau, P., & Kosmadopoulos, A. (2022). Disturbance of the
circadian system in shift work and its health impact. *Journal of Biological
Rhythms*, *37*(1), 3–28. https://doi.org/10.1177/07487304211064218

Brown, T. M., Brainard, G. C., Cajochen, C., Czeisler, C. A., Hanifin, J. P., Lockley,
S. W., Lucas, R. J., Münch, M., O'Hagan, J. B., Peirson, S. N., Price, L. L. A.,

Roenneberg, T., Schlangen, L. J. M., Skene, D. J., Spitschan, M., Vetter, C.,

Zee, P. C., & Wright, K. P. (2022). Recommendations for daytime, evening,

and nighttime indoor light exposure to best support physiology, sleep, and

wakefulness in healthy adults. *PLoS Biology*, *20*(3), e3001571.

https://doi.org/10.1371/journal.pbio.3001571

Buysse, D. J., Reynolds, C. F., Monk, T. H., Berman, S. R., & Kupfer, D. J. (1989).

The Pittsburgh sleep quality index: A new instrument for psychiatric practice

and research. *Psychiatry Research*, *28*(2), 193–213.

https://doi.org/10.1016/0165-1781(89)90047-4

Chang, A.-M., Scheer, F. A. J. L., Czeisler, C. A., & Aeschbach, D. (2013). Direct

effects of light on alertness, vigilance, and the waking electroencephalogram

in humans depend on prior light history. *Sleep*, *36*(8), 1239–1246.

https://doi.org/10.5665/sleep.2894

Christensen, R. H. B. (2010). *Ordinal: Regression models for ordinal data*. 2023.12-

4.1. https://doi.org/10.32614/CRAN.package.ordinal

CIE. (2018). *CIE S 026/E:2018 CIE System for metrology of optical radiation for

ipRGC-influenced responses to light*. International Commission on Illumination

(CIE). https://doi.org/10.25039/S026.2018

Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.).

Routledge, New York. https://doi.org/10.4324/9780203771587

Cohen, J. (1992). Statistical power analysis. *Current Directions in Psychological

Science*, *1*(3), 98–101. https://doi.org/10.1111/1467-8721.ep10768783

Didikoglu, A., Mohammadian, N., Johnson, S., Van Tongeren, M., Wright, P.,

Casson, A. J., Brown, T. M., & Lucas, R. J. (2023). Associations between light

exposure and sleep timing and sleepiness while awake in a sample of UK

adults in everyday life. *Proceedings of the National Academy of Sciences*,

120(42), e2301608120. https://doi.org/10.1073/pnas.2301608120

Dijk, D.-J., & Archer, S. N. (2009). Light, sleep, and circadian rhythms: Together

again. *PLoS Biology*, *7*(6), e1000145.

https://doi.org/10.1371/journal.pbio.1000145

Fernández-Martínez, M., Vicca, S., Janssens, I. A., Carnicer, J., Martín-Vide, J., &

Peñuelas, J. (2018). The consecutive disparity index, *D*: A measure of

temporal variability in ecological studies. *Ecosphere*, *9*(12), e02527.

https://doi.org/10.1002/ecs2.2527

Fishbein, A. B., Knutson, K. L., & Zee, P. C. (2021). Circadian disruption and human

health. *Journal of Clinical Investigation*, *131*(19), e148286.

https://doi.org/10.1172/JCI148286

Foster, R. G. (2020). Sleep, circadian rhythms and health. *Interface Focus*, *10*(3),

20190098. https://doi.org/10.1098/rsfs.2019.0098

Foster, R. G., Hughes, S., & Peirson, S. N. (2020). Circadian Photoentrainment in

Mice and Humans. *Biology*, *9*(7), 180. https://doi.org/10.3390/biology9070180

Harrell, F. E. J. (2003). *Hmisc: Harrell Miscellaneous*. 5.1-3.

https://doi.org/10.32614/CRAN.package.Hmisc

Harris, P. A., Taylor, R., Minor, B. L., Elliott, V., Fernandez, M., O'Neal, L., McLeod,

L., Delacqua, G., Delacqua, F., Kirby, J., & Duda, S. N. (2019). The REDCap

consortium: Building an international community of software platform partners.

Journal of Biomedical Informatics, *95*, 103208.

https://doi.org/10.1016/j.jbi.2019.103208

Harris, P. A., Taylor, R., Thielke, R., Payne, J., Gonzalez, N., & Conde, J. G. (2009).

Research electronic data capture (REDCap)—A metadata-driven

methodology and workflow process for providing translational research

informatics support. *Journal of Biomedical Informatics*, *42*(2), 377–381.

https://doi.org/10.1016/j.jbi.2008.08.010

Hébert, M., Martin, S. K., Lee, C., & Eastman, C. I. (2002). The effects of prior light

history on the suppression of melatonin by light in humans. *Journal of Pineal*

Research, *33*(4), 198–203. https://doi.org/10.1034/j.1600-079x.2002.01885.x

Kervezee, L., Blume, C., Brainard, G. C., Broszio, K., Brown, T. M., Dennemann, J.,

Fernandez-Alonso, M., Joyce, D. S., Lok, R., Lucas, R. J., McGlashan, E. M.,

Najjar, R. P., Price, L., Rabstein, S., Sliney, D., Stefani, O., Veitch, J.,

Weiskopf, D., Zauner, J., & Spitschan, M. (2024). *Call for a coordinated*

international communications effort on the effects of ocular light exposure on

health and well-being: White paper from the Ladenburg Roundtable. OSF.

https://doi.org/10.17605/osf.io/rxa35

Kuhlman, S. J., Craig, L. M., & Duffy, J. F. (2018). Introduction to chronobiology.

Cold Spring Harbor Perspectives in Biology, *10*(9), a033613.

https://doi.org/10.1101/cshperspect.a033613

Lunn, R. M., Blask, D. E., Coogan, A. N., Figueiro, M. G., Gorman, M. R., Hall, J. E.,

Hansen, J., Nelson, R. J., Panda, S., Smolensky, M. H., Stevens, R. G.,

Turek, F. W., Vermeulen, R., Carreón, T., Caruso, C. C., Lawson, C. C.,

Thayer, K. A., Twery, M. J., Ewens, A. D., … Boyd, W. A. (2017). Health

consequences of electric lighting practices in the modern world: A report on

the National Toxicology Program's workshop on shift work at night, artificial

light at night, and circadian disruption. *Science of The Total Environment*,

607–608, 1073–1084. https://doi.org/10.1016/j.scitotenv.2017.07.056

Mohamed, A., Kalavally, V., Cain, S. W., Phillips, A. J. K., McGlashan, E. M., & Tan,

C. P. (2021). Wearable light spectral sensor optimized for measuring daily α-

opic light exposure. *Optics Express*, *29*(17), 27612–27627.

https://doi.org/10.1364/OE.431373

Neves, A. R., Albuquerque, T., Quintela, T., & Costa, D. (2022). Circadian rhythm

and disease: Relationship, new insights, and future perspectives. *Journal of*

Cellular Physiology, *237*(8), 3239–3256. https://doi.org/10.1002/jcp.30815

Pilz, L. K., Levandovski, R., Oliveira, M. A. B., Hidalgo, M. P., & Roenneberg, T.

(2018). Sleep and light exposure across different levels of urbanisation in

Brazilian communities. *Scientific Reports*, *8*(1), 11389.

https://doi.org/10.1038/s41598-018-29494-4

Press, W. H., Teukolsky, S. A., Vetterling, W. T., & Flannery, B. P. (1992). *Numerical*

Recipes in C: The Art of Scientific Computing (2nd ed.). Cambridge University

Press.

R Core Team. (2021). *R: A Language and Environment for Statistical Computing*. R

Foundation for Statistical Computing. https://www.R-project.org/

Roenneberg, T., & Klerman, E. B. (2019). Chronobiology: A short introduction.

Somnologie, *23*(3), 142–146. https://doi.org/10.1007/s11818-019-00217-9

Roenneberg, T., & Merrow, M. (2016). The circadian clock and human health.

Current Biology, *26*(10), R432–R443.

https://doi.org/10.1016/j.cub.2016.04.011

Siraji, M. A., Lazar, R. R., van Duijnhoven, J., Schlangen, L. J. M., Haque, S.,

Kalavally, V., Vetter, C., Glickman, G. L., Smolders, K. C. H. J., & Spitschan,

M. (2023). An inventory of human light exposure behaviour. *Scientific*

Reports, *13*(1), 22151. https://doi.org/10.1038/s41598-023-48241-y

Smith, K. A., Schoen, M. W., & Czeisler, C. A. (2004). Adaptation of human pineal

melatonin suppression by recent photic history. *The Journal of Clinical*

Endocrinology and Metabolism, *89*(7), 3610–3614.

https://doi.org/10.1210/jc.2003-032100

Spitschan, M., Kervezee, L., Lok, R., McGlashan, E., Najjar, R. P., & ENLIGHT

Consortium. (2023). ENLIGHT: A consensus checklist for reporting laboratory-

based studies on the non-visual effects of light in humans. *EBioMedicine*, *98*,

104889. https://doi.org/10.1016/j.ebiom.2023.104889

Spitschan, M., Stefani, O., Blattner, P., Gronfier, C., Lockley, S., & Lucas, R. (2019).

How to report light exposure in human chronobiology and sleep research

experiments. *Clocks & Sleep*, *1*(3), 280–289.

https://doi.org/10.3390/clockssleep1030024

Stefani, O., Marek, R., Schwarz, J., Plate, S., Zauner, J., & Schrader, B. (2024).

Wearable Light loggers in field conditions: Corneal light characteristics, user

compliance, and acceptance. *Clocks & Sleep*, *6*(4), 619–634.

https://doi.org/10.3390/clockssleep6040042

Wahl, S., Engelhardt, M., Schaupp, P., Lappe, C., & Ivanov, I. V. (2019). The inner

clock—Blue light sets the human rhythm. *Journal of Biophotonics*, *12*(12),

e201900102. https://doi.org/10.1002/jbio.201900102

Webler, F. S., Spitschan, M., Foster, R. G., Andersen, M., & Peirson, S. N. (2019).

What is the 'spectral diet' of humans? *Current Opinion in Behavioral*

Sciences, *30*, 80–86. https://doi.org/10.1016/j.cobeha.2019.06.006

Wickham, H. (2016). *tidyverse: Easily install and load the 'Tidyverse'*. 2.0.0.

https://doi.org/10.32614/CRAN.package.tidyverse

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R.,

Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T.,

Miller, E., Bache, S., Müller, K., Ooms, J., Robinson, D., Seidel, D., Spinu, V.,

… Yutani, H. (2019). Welcome to the Tidyverse. *Journal of Open Source*

Software, *4*(43), 1686. https://doi.org/10.21105/joss.01686

Windred, D. P., Burns, A. C., Lane, J. M., Olivier, P., Rutter, M. K., Saxena, R.,

Phillips, A. J. K., & Cain, S. W. (2024). Brighter nights and darker days predict

higher mortality risk: A prospective analysis of personal light exposure in

>88,000 individuals. *Proceedings of the National Academy of Sciences*,

121(43), e2405924121. https://doi.org/10.1073/pnas.2405924121

Windred, D. P., Burns, A. C., Rutter, M. K., Yeung, C. H. C., Lane, J. M., Xiao, Q.,

Saxena, R., Cain, S. W., & Phillips, A. J. K. (2024). Personal light exposure

patterns and incidence of type 2 diabetes: Analysis of 13 million hours of light

sensor data and 670,000 person-years of prospective observation. *The*

Lancet Regional Health – Europe, *42*.

https://doi.org/10.1016/j.lanepe.2024.100943

Wood, S. (2000). *mgcv: Mixed GAM Computation Vehicle with Automatic*

Smoothness Estimation. 1.9-1. https://doi.org/10.32614/CRAN.package.mgcv

Wright, K. P., Jr., McHill, A. W., Birks, B. R., Griffin, B. R., Rusterholz, T., & Chinoy,

E. D. (2013). Entrainment of the human circadian clock to the natural light-

dark cycle. *Current Biology*, *23*(16), 1554–1558.

https://doi.org/10.1016/j.cub.2013.06.039

Zauner, J., Hartmeyer, S., & Spitschan, M. (2024). *LightLogR: Reproducible analysis*

of personal light exposure data with an open-source R package.

https://doi.org/10.31219/osf.io/qk924

Zerbini, G., Kantermann, T., & Merrow, M. (2020). Strategies to decrease social

jetlag: Reducing evening blue light advances sleep and melatonin. *The*

European Journal of Neuroscience, *51*(12), 2355–2366.

https://doi.org/10.1111/ejn.14293

Zerbini, G., Winnebeck, E. C., & Merrow, M. (2021). Weekly, seasonal, and

chronotype-dependent variation of dim-light melatonin onset. *Journal of Pineal*

Research, *70*(3). https://doi.org/10.1111/jpi.12723

1017

1018

Figures and Tables

Figure 1. Overview of the study design. A, Shows the study design and measurements included for both research sites. **B**, Both research locations shown on a world map. **C,** Individual recording periods per site and participant including implicit missing data in grey. Note that participant ID 06 is missing in the Malaysian dataset due to a lost Speccy device. **D,** Photoperiod duration in Singapore (blue), and Switzerland (yellow). **E,** Grand averages of light exposure of illuminance (melanopic EDI) for Singapore (blue) and Switzerland (yellow). Abbreviations: LEBA, Light Exposure Behaviour Assessment; MEDI, melanopic equivalent daylight illuminance; PSQI, Pittsburgh Sleep Quality Index.

Figure 2. **Double plots of average light exposure over participants across both research sites.** Median and percentiles of light exposure measured in illuminance (lux, melanopic EDI per 15 minutes (95%, 75%, and 50% of data

shown in ribbons). Grey background indicates civil nighttime. Abbreviations: IS, Interdaily stability. Melanopic EDI, Melanopic equivalent daylight illuminance.

Figure 3. **Model fits of light exposure differences between Malaysia and Switzerland. A,** Individual model fits per ID and research site for melanopic EDI (lux). **B,** Model fits over all IDs per research site with with 95% confidence interval. **C,** Fitted differences between research sites smoothed with 95% confidence interval with an indication of significant differences per time. Abbreviations: CI, confidence interval; mel EDI, melanopic equivalent daylight illuminance.

Figure 4. **Correlations between individual LEBA items and pre-selected light metrics for Malaysia and Switzerland.** Abbreviations: FcT250, Frequency crossing threshold at 250 lux; FLiT1000, Frequency of light exposure above 1000 lux; FLiT250, Frequency of light exposure above 250 lux; IS, Interdaily Stability; IV, Intradaily Variability; L5m, Mean light exposure during the least active 5 hours; L5mde, Mean duration of exposure during the least active 5 hours; L5mn, Number of exposures during the least active 5 hours; LE, Light Exposure; LEBA, Light Exposure Behaviour Assessment; LLiT10, Lowest light exposure intensity threshold at 10 lux; M10m, Mean light exposure during the most active 10 hours; MPratio, Ratio of mean photopic to melanopic light exposure; PAT1000, Period above threshold at 1000 lux; SCmelEDI, Spectral composition of melanopic equivalent daylight illuminance; TAT1000, Time above threshold at 1000 lux; TAT250, Time above threshold at 250 lux; TATd250, Time above dynamic threshold at 250 lux; TBTe10, Time below threshold at 10 lux.

Summary of significant results for Hypothesis 5						
Site	Question/Factor	Metric	LEBA item	Correlation coefficient[1]	p-value[2]	Interpretation
switzerland	I spend 30 minutes or less per day (in total) outside	PAT1000	F2, 04	-0.33	0.018	The more often 30 minutes or less are spent outside, the shorter the period above 1000 lx
switzerland	I spend more than 3 hours per day (in total) outside	PAT1000	F2, 07	0.46	<0.001	The more often >3 hours per day are spent outside, the longer the period above 1000 lx
switzerland	I spend more than 3 hours per day (in total) outside	TAT1000	F2, 07	0.45	<0.001	The more often > 3 hours per day are spent outside, the higher the time above 1000 lx
switzerland	I spend as much time outside as possible	M10m	F2, 08	0.39	0.003	The more spend time outside as possible, the brighter the brightest 10 hours (mean)
switzerland	I spend as much time outside as possible	PAT1000	F2, 08	0.47	<0.001	The more spend time outside as possible, the longer the period above 1000 lx
switzerland	I spend as much time outside as possible	TAT1000	F2, 08	0.48	<0.001	The more spend time outside as possible, the higher the time above 1000 lx
switzerland	I spend as much time outside as possible	TAT250	F2, 08	0.32	0.011	The more spend time outside as possible, the higher the time above 250 lx
switzerland	I look at my smartwatch within 1 hour before attempting to fall asleep	LLiT250	F3, 13	0.34	<0.001	The more often smartwatches are looked at within 1h before attempting to fall asleep, the later the last time above 250 lx
switzerland	I use an alarm with a dawn simulation light	SCmelEDI	F5, 22	-0.34	<0.001	The more an alarm with a dawn simulation light is used, the warmer the light colour temperature
switzerland	I turn on the lights immediately after waking up	FLiT250	F5, 23	-0.35	0.002	The more often the lights are turned on immediately after waking up, the earlier the first time above 250 lx

[1] Spearman correlation coefficient between the metric and the LEBA question/factor

[2] p-values are adjusted for multiple comparisons using the false-discovery-rate for n= 168 comparisons

1062

Table 1. **Significant correlations between individual LEBA items and**
hypothesised light metrics for Malaysia and Switzerland. P-values are
considered significant below 0.05. Abbreviation: FLiT250, First time above threshold
of 250 lux; LLiT250, Last time above threshold of 250 lux; M10m, Mean light
exposure during the most active 10 hours; PAT1000, Period above threshold at 1000
lux; SCmelEDI, Spectral composition of equivalent daylight illuminance; TAT1000,
Time above threshold at 1000 lux; TAT250, Time above threshold at 250 lux.

Supplementary material

Supplementary Tables

Factor name	Score
F1: Wearing blue light filters	01+02+03
F2: Spending time outdoors	04(R)+05+06+07+08+09
F3: Using phone and smartwatch in bed	10+11+12+13+14
F4: Using light before bedtime	15+16+17+18
F5: Using light in the morning and during daytime	19+20+21+22+23

Table S1. Scoring of Light Exposure Behaviour Assessment (LEBA) items. Note that R denotes items that are reversed-scored. F1-F5 denotes the calculated Factors.

Component	Question (Q)	Component Score
(1) Subjective sleep quality	#6	0-3
(2) Sleep latency	#2 #5a	Sum of Q2 (0-3)+Q5a (0-3): If 0 → 0 1-2 → 1 3-4 → 2 5-6 → 3
(3) Sleep duration	#4	0-3
(4) Habitual sleep efficiency (Sleep efficiency = hours slept / hours in bed) X 100%)	#1 #3 #4	If >85% → 0 75-84% → 1 65-74% → 2 <65% → 3
(5) Sleep disturbance	#5b-5j	Sum of Q5b (0-3) to Q5 (0-3): If 0 → 0 1-9 → 1 10-18 → 2 19-27 → 3
(6) Use of sleep medication	#7	0-3
(7) Daytime dysfunction	#8 #9	Sum of Q8 (0-3) +Q9 (0-3: If 0 → 0 1-2 → 1 3-4 → 2 5-6 → 3
Global PSQI Score		**Sum of Components 1-7**

1077
1078   **Table S2. Scoring of the Pittsburgh Sleep Quality Index (PSQI) items.**

Call to calculate metric	Light metric name	Description
bright_dark_period()	Brightest or darkest continuous period	Finds the brightest or darkest continuous period of a given timespan and calculates its mean light level, as well as the timing of the period's onset, midpoint, and offset. It is defined as the period with the maximum or minimum mean light level. Note that the data need to be regularly spaced (i.e., no gaps) for correct results
duration_above_threshold()	Duration above/below threshold or within threshold range	Calculates the duration spent above/below a specified threshold light level or within a specified range of light levels
frequency_crossing_threshold()	Frequency of crossing light threshold	Calculates the number of times a given threshold light level is crossed
intradaily_variability()	Intradaily variability (IV)	Calculates the variability of consecutive light levels within a 24h day. Calculated as the ratio of the variance of the differences between consecutive light levels to the total variance across the day. Calculated with mean hourly light levels. Higher values indicate more fragmentation
interdaily_stability()	Interdaily stability (IS)	This function calculates the variability of 24h light exposure patterns across multiple days. Calculated as the ratio of the variance of the average daily pattern to the total variance across all days. Calculated with mean hourly light levels. Ranges between 0 (Gaussian noise) and 1 (Perfect Stability)
period_above_threshold()	Length of longest continuous period above/below threshold	Length of the longest continous period above/below a specified threshold light level or within a specified range of light

		levels
pulses_above_threshold()	Pulses above threshold	Clusters the light data into continuous clusters (pulses) of light above/below a given threshold
threshold_for_duration()	Find threshold for given duration	Threshold for which light levels are above/below for a given duration. This function can be considered as the inverse of duration_above_threshold
timing_above_threshold()	Mean/first/last timing above/below threshold.	Mean, first, and last timepoint (MLiT, FLiT, LLiT) where light levels are above or below a given threshold intensity within the given time interval

1079

1080 **Table S3. Overview of objectively derived outcome variables from light logging**
1081 **data and their calculations.**

RQ #	Hypothesis	Variable(s)	Statistical test(s)	Wilkinson Formula (if appropriate)
1	**H1:** There are differences in light logger-derived light exposure **intensity levels** and **duration** of intensity between Malaysia and Switzerland. **H0₁:** No difference.	Intensity level light metrics - TAT250 - TAT1000 - Period above threshold 1000 - TAT250 (daytime hours) - TBT10 (evening hours)	Generalized mixed-effect analyses on the effect of site on various light parameters 5 Models	$$\text{metric} = \text{site} + (1|\text{site:participant})$$
	H2: There are differences in light logger-derived **timing of light** exposure between Malaysia and Switzerland. **H0₂:** No difference.	Timing light metrics - M10m - L5m - IS - IV - LLiT 10 - LLit 250 - Frequency crossing threshold	Generalized mixed-effect model (GAMM) on the effect of site on various parameters 7 models	$$\text{metric} = \text{site} + (1|\text{site:participant})$$
		Time-of-day (two-level factor: daytime / evening) Melanopic EDI	Linear mixed-effect model on the interaction of time-of-day and site on melanopic EDI	$$\text{melEDI} = \text{site} \cdot \text{time-of-day} + (1 + \text{time-of-day} | \text{site:participant})$$
		Time-of-day (seco from midnight) Melanopic EDI	Generalized mixed-effect model (GAMM) on the interaction of time-of-day and site on melanopic EDI	$$\text{melEDI} = s(\text{time-of-day, site}) + s(\text{time-of-day, by} = \text{participant}) + s(\text{participant, by} = \text{site})$$ Note on GAMM basis splines for smooth (*s()*) terms: 1. cyclic spline (cs), factor spline (fs) 2. cyclic spline (cs) 3. random effect (re)
2	**H3:** There are differences in LEBA items and factors between Malaysia and Switzerland.	- all individual LEBA items (**Table 5**) - all 5 LEBA Factors (**Table 2**)	Cumulative link mixed-effect analyses on the effect of site on each LEBA question and factor	$$\text{LEBA item/factor} = \text{site} + (1 | \text{site:participant})$$

	H0₃: No difference.			23 + 5 models	
	H4: LEBA scores vary over time within participants **H0₄:** no variance	- all individual LEBA items (**Table 5**) - all 5 LEBA Factors (**Table 2**)	Bootstrap analysis of the standard deviation of LEBA scores within participants for the Malaysia dataset 23 + 5 bootstraps	
3	**H5:** LEBA items correlate with pre-selected light logger-derived light exposure variables. **H0₅:** No correlation.	See **Table 5** for all relevant variables.	Correlation matrix with one matrix per site. 2x85 correlations	
	H6: There is a difference between Malaysia and Switzerland on how well light logger-derived light exposure variables correlate with subjective LEBA items. **H0₆:** No difference between Malaysia and Switzerland.	See **Table 5** for all relevant variables.	Generalized mixed-effect analyses to determine the effect of site on the dependency between light exposure metrics and LEBA items/factors 23 + 5 models	$\text{metric} =$ $\text{site} \cdot \text{LEBA item/factor} +$ $(1	\text{site:participant})$

Table S4. Overview of research questions, hypotheses and statistical tests to address the hypotheses. Abbreviations: GAMM, Generalised additive mixed-effect model; $H0_x$, Null hypothesis; HX, Hypothesis X; IS, Interdaily Stability; IV, Intradaily variability; LEBA, Light Exposure Behaviour Assessment; LLiT, Last time above threshold; LLiT 10, Low light intensity threshold at 10 lux; LLiT 250, Low light intensity threshold at 250 lux; L5m, Mean across darkest 5 hours; M10m, Mean across brightest 10 hours; melEDI, Melanopic equivalent daylight illuminance; RQ, Research question; TAT250, Time above threshold 250 lux; TAT1000, Time above threshold 1000 lux; TBT10, Time below threshold 10 lux.

LEBA Item	LEBA Question	List of calculated light variables
1	I wear blue-filtering, orange-tinted, and/or red-tinted glasses indoors during the day.	NA
2	I wear blue-filtering, orange-tinted, and/or red-tinted glasses outdoors during the day.	NA
3	I wear blue-filtering, orange-tinted, and/or red-tinted glasses 1 hour before attempting to fall asleep.	NA
4	I spend 30 minutes or less per day (in total) outside.	['M10', 'TAT250', 'TAT1000', 'Period above threshold 1000', 'IV', 'IS']
5	I spend between 30 minutes and 1 hour per day (in total) outside.	['M10', 'TAT250', 'TAT1000', 'Period above threshold 1000', 'IV', 'IS']
6	I spend between 1 and 3 hours per day (in total) outside.	['M10', 'TAT250', 'TAT1000', 'Period above threshold 1000', 'IV', 'IS']
7	I spend more than 3 hours per day (in total) outside.	['M10', 'TAT250', 'TAT1000', 'Period above threshold 1000', 'IV', 'IS']
8	I spend as much time outside as possible.	['M10', 'TAT250', 'TAT1000', 'Period above threshold 1000', 'IV', 'IS']
9	I go for a walk or exercise outside within 2 hours after waking up.	['first time above threshold of 1000 (FLitT 1000)', 'IV', 'IS']
10	I use my mobile phone within 1 hour before attempting to fall asleep.	['last time above threshold of 10 (LLiT 10)', 'last time above threshold of 250 (LLiT 250)', 'L5 (without sleep period)', 'IV', 'IS']
13	I look at my smartwatch within 1 hour before attempting to fall asleep.	['last time above threshold of 10 (LLiT 10)', 'last time above threshold of 250 (LLiT 250)', 'L5 (without sleep period)', 'IV', 'IS']
11	I look at my mobile phone screen immediately after waking up.	['first time above threshold of 250 (FLitT 250)', 'IV', 'IS']
12	I check my phone when I wake up at night.	['L5', 'IV', 'IS']
14	I look at my smartwatch when I wake up at night.	['L5', 'IV', 'IS']
15	I dim my mobile phone screen within 1 hour before attempting to fall asleep.	['last time above threshold of 10 (LLiT 10)', 'last time above threshold of 250 (LLiT 250)', 'L5 (without sleep period)', 'IV', 'IS']
16	I use a blue-filter app on my computer screen within 1 hour before attempting to fall asleep.	['last time above threshold of 10 (LLiT 10)', 'last time above threshold of 250 (LLiT 250)', 'L5 (without sleep period)', 'IV', 'IS']
17	I use as little light as possible when I get up during the night.	['L5 and L5 (only night)', 'IV', 'IS']
18	I dim my computer screen within 1 hour before attempting to fall asleep.	['last time above threshold of 10 (LLiT 10)', 'last time above threshold of 250 (LLiT 250)', 'L5 (without sleep period)', 'IV', 'IS']
19	I use tunable lights to create a healthy light environment.	['IV', 'IS', 'L5 (with/without night)', 'M10']
20	I use LEDs to create a healthy light environment.	['LE', 'Spectral contribution', 'Melanopic/photopic ratio']
21	I use a desk lamp when I do focused work.	['LE']
22	I use an alarm with a dawn simulation.	['Spectral contribution', 'first time above threshold of 10 (FLitT 10)', 'first time above threshold of 250 (FLitT 250)']
23	I turn on the lights immediately after waking up.	['Spectral contribution', 'first time above threshold of 10 (FLitT 10)', 'first time above threshold of 250 (FLitT 250)']

Table S5. List of Light Exposure Behaviour Assessment (LEBA) items and
corresponding objectively derived light variables from light logging data. In
yellow are the hypothesised and significant relationships in the Swiss dataset.
Abbreviations: FLiT 10, First time above threshold of 10 lux; FLiT 250, First time above
threshold of 250 lux; FLiT 1000, First time above threshold of 1000 lux; IS, Interdaily
Stability; IV, Intradaily Variability; LE, Light Exposure; L5, Mean across darkest 5
hours; L5 (only night), Mean across darkest 5 hours during the night; LLiT 10, Last
time above threshold of 10 lux; LLiT 250, Last time above threshold of 250 lux; M10,
Mean across brightest 10 hours; TAT250: Time above threshold of 250 lux; TAT1000:
Time above threshold of 1000 lux.
1103
1104

1105

A   Missing data in the Malaysia dataset overall

	min	max	median	mean	total
Number of gaps	0	7	0	1	22
Duration missed by available data points	0 days	14 days	0 days	1 day	29 days
Duration covered by available data points	11 days	30 days	30 days	26 days	509 days
Number of missing data points	0	21,040	0	2,200	41,797
Number of available data points	16,604	43,996	43,211	38,627	733,907
Percentage of missing data points	0.0%	48.7%	0.0%	5.4%	5.4%

min, max, median, mean, and total values for 19 participants

B   Missing data in the Swiss dataset overall

	min	max	median	mean	total
Number of gaps	1	4	1	2	35
Duration missed by available data points	<1 day	11 days	<1 day	1 day	28 days
Duration covered by available data points	7 days	30 days	29 days	27 days	554 days
Number of missing data points	21	16,017	44	2,035	40,698
Number of available data points	11,476	44,586	42,974	39,954	799,070
Percentage of missing data points	0.0%	40.4%	0.1%	5.9%	4.8%

min, max, median, mean, and total values for 20 participants

1106
1107
1108 **Table S6. Overview of missing data for the two datasets.** A, Shows the overview
1109 for Malaysia and B, for Switzerland. Abbreviations: Min, minimum; Max, maximum.
1110

1111

	Model Results for Hypothesis 1				
			Site coefficients		
	p-value[1]	**Intercept**[2]	**Malaysia**[3]	**Switzerland**[3]	
TAT250	**0.002**	418	1	1.780	
TAT1000	**0.011**	180	1	1.942	
PAT1000	0.15	78	—	—	
TATd250	**0.003**	410	1	1.772	
TBTe10	0.9	678	—	—	

[1] p-values are adjusted for multiple comparisons using the false-discovery-rate for n= 5 comparisons

[2] Model prediction for the intercept per hour of photo- or nighttime period, reference level for the site is Malaysia

[3] Exponentiated beta coefficients from the final model, denoting the multiplication factor for the intercept, conditional on the site

1112

1113 **Table S7. Model results for H1.** P-values are considered significant below 0.05.
1114 Abbreviations: PAT1000, Period above threshold 1000, TAT250, Time above
1115 threshold of 250 lux; TAT1000, Time above threshold of 1000 lux; TATd250, Time
1116 above dynamic threshold 250 lux; TBTe10, Time below threshold 10 lux.

1117

1118

Model Results for Hypothesis 2, Interaction							
			Site coefficients		Time coeff.		Interaction coeff.
	p-value[1,2]	Intercept	Malaysia	Switzerland	Day	Evening	Switzerland:Evening
log10(daily mean mel EDI)	<0.001	2.23	0	0.41	0	−0.98	−1.11

Values were log10 transformed before model fitting

[1] p-values are adjusted for multiple comparisons using the false-discovery-rate for n= 1 comparisons

[2] p-value refers to the interaction effect

Table S8. Model results for H2. P-values are considered significant below 0.05. Abbreviations: Mel EDI, melanopic equivalent daylight illumination.

Model Results for Hypothesis 2, Timing

	p-value[1]	Intercept	Site coefficients Malaysia	Switzerland	
M10m[2]	**0.004**	2.36	0	0.46	
L5m[2]	0.2	−0.88	—	—	
IV	0.9	1.28	—	—	
LLiT10	**0.005**	23:16:14	0	−01:10:42	
LLiT250	**<0.001**	17:41:23	0	01:27:54	
FcT250	**0.005**	35.79	0	28.71	
IS	0.9	0.16	—	—	

[1] p-values are adjusted for multiple comparisons using the false-discovery-rate for n= 9 comparisons

[2] Values were log10 transformed before model fitting

Table S9. Model results for H2. P-values are considered significant below 0.05. Abbreviations: FcT250, Frequency of crossing light threshold at 250 lux; FLiT 10, First time above threshold of 10 lux; FLiT 250, First time above threshold of 250 lux; FLiT 1000, First time above threshold of 1000 lux; IS, Interdaily Stability; IV, Intradaily Variability; L5m, Mean across darkest 5 hours; LE, Light Exposure; LLiT 10, Last time above threshold of 10 lux; LLiT 250, Last time above threshold of 250 lux; M10m, Mean across brightest 10 hours;

	Malaysia			Switzerland			Difference M-S	Boxplot
	Mean	SD	n	Mean	SD	n		
FLiT1000	38,664	8,857	413	35,909	8,818	489	2,755	
FLiT250	37,192	9,723	439	33,204	9,880	509	3,988	
FcT250	36	31	490	67	44	533	−31	
IS	0.162	0.0581	19	0.164	0.0664	20	−0.00129	
IV	1.29	0.483	472	1.28	0.504	504	0.00480	
L5m[1]	−0.781	0.524	490	−0.966	0.183	533	0.185	
L5mde[1]	0.836	0.785	490	0.971	0.912	533	−0.136	
L5mn[1]	−0.341	0.897	490	−0.797	0.489	533	0.456	
LE	5,117	7,132	490	18,765	26,249	533	−13,648	
LLiT10	23:18	01:04	483	22:18	01:52	528	01:00	
LLiT250	17:42	02:36	432	19:14	02:10	510	−01:31	
M10m[1]	2.37	0.578	490	2.87	0.706	533	−0.503	
MEDI[1]	0.469	1.42	704,123	0.547	1.61	765,617	−0.0788	
MPratio	0.932	0.119	490	0.947	0.103	533	−0.0150	
PAT1000	957	1,360	490	1,692	2,031	533	−736	
SCmelEDI	1.52×10^{-3}	1.36×10^{-4}	490	1.40×10^{-3}	1.19×10^{-4}	533	1.15×10^{-4}	
TAT1000	47m	1h 4m	490	1h 44m	1h 41m	533	−57m	
TAT250	1h 38m	1h 38m	490	3h 28m	2h 20m	533	−1h 50m	
TATd250	1h 36m	1h 36m	490	3h 25m	2h 18m	533	−1h 49m	
TBTe10	2h 32m	1h 5m	490	2h 39m	1h 1m	533	−7m	

[1] values were log10 transformed

Table S10. Differences in descriptive statistics for light exposure metrics between Malaysia and Switzerland. Mean, standard deviation (SD), sample size (n) and mean differences between the two locations (M-S) are presented for each variable. Abbreviations: FcT250, Frequency of crossing light threshold at 250 lux; FLiT250, First time above threshold of 250 lux; FLiT1000, First time above threshold of 1000 lux; IS, Interdaily Stability; IV, Intradaily Variability; LE, Light Exposure; L5m, Mean across darkest 5 hours; L5mde, Mean duration of exposure during the least active 5 hours; L5mn, Number of exposures during the least active 5 hours; LLiT10, Last time above threshold of 10 lux; LLiT250, Last time above threshold of 250 lux; M10m, Mean across brightest 10 hours; MEDI, Melanopic equivalent daylight illuminance; MPratio, Ratio between melanopic and photopic illuminance; PAT1000, Period above threshold of 1000 lux; SCmelEDI, Spectal composition equivalent daylight illuminance; TAT250, Time above threshold of 250 lux; TAT1000, Time above threshold of 1000 lux; TATd250, Time above dynamic threshold of 250 lux; TBTe10, Time below threshold of 10 lux.

1149

Hypothesis 4: Extent of stable LEBA items and factors over time
stable over time in 75-100% of participants
Factor 1: Item 01, Factor 1: Item 02, Factor 1: Item 03, Factor 2: Item 09, Factor 3: Item 10, Factor 3: Item 13, Factor 3: Item 14, Factor 5: Item 19, Factor 5: Item 21, Factor 5: Item 22, Factor 1
stable over time in 50-75% of participants
Factor 2: Item 04, Factor 2: Item 05, Factor 2: Item 06, Factor 2: Item 07, Factor 2: Item 08, Factor 3: Item 11, Factor 3: Item 12, Factor 4: Item 15, Factor 4: Item 16, Factor 4: Item 17, Factor 4: Item 18, Factor 5: Item 20, Factor 5: Item 23
stable over time in 25-50% of participants
Factor 3
stable over time in 25-0% of participants
Factor 2, Factor 4, Factor 5

1150

1151

1152   **Table S11. Stability of LEBA items and factors over time.** For a description of each
1153   LEBA item, see Table S5.

1154

1155

| | p-value[†] | Intercept | Site coefficients | |
			Malaysia	Switzerland
leba_F2_04	0.2	−0.07	—	—
leba_F2_05	0.9	−0.07	—	—
leba_F2_06	0.054	0.03	—	—
leba_F2_07	0.4	0.06	—	—
leba_F2_08	0.9	0.16	—	—
leba_F2_09	0.9	−0.08	—	—
leba_F3_10	0.9	0.03	—	—
leba_F3_11	0.9	0.05	—	—
leba_F3_12	0.9	0.00	—	—
leba_F3_13	0.9	0.01	—	—
leba_F3_14	0.9	−0.08	—	—
leba_F4_15	0.019	0.25	0	−0.40
leba_F4_16	0.9	0.09	—	—
leba_F4_17	0.9	−0.16	—	—
leba_F4_18	0.035	0.25	0	−0.26
leba_F5_19	0.9	−0.10	—	—
leba_F5_20	0.9	0.21	—	—
leba_F5_22	0.9	−0.20	—	—
leba_F5_23	0.9	−0.07	—	—

Model Results for Hypothesis 6

[†] p-values are adjusted for multiple comparisons using the false-discovery-rate for n= 28 comparisons

Table S12. Model results of H6. P-values are considered significant below 0.05. For a description of each LEBA item, see Table S5.

Characteristic	Malaysia N = 19[1]	Switzerland N = 20[1]	p-value[2]
Subjective sleep quality			>0.9
0	3 (16%)	3 (15%)	
1	15 (79%)	16 (80%)	
2	1 (5.3%)	1 (5.0%)	
3	0 (0%)	0 (0%)	
Sleep latency			>0.9
0	7 (37%)	6 (30%)	
1	9 (47%)	10 (50%)	
2	3 (16%)	3 (15%)	
3	0 (0%)	1 (5.0%)	
Sleep duration			0.2
0	4 (21%)	9 (45%)	
1	13 (68%)	11 (55%)	
2	1 (5.3%)	0 (0%)	
3	1 (5.3%)	0 (0%)	
Habitual sleep efficiency			0.020*
0	18 (95%)	12 (60%)	
1	1 (5.3%)	8 (40%)	
2	0 (0%)	0 (0%)	
3	0 (0%)	0 (0%)	
Sleep disturbance			0.5
0	2 (11%)	2 (10%)	
1	17 (89%)	16 (80%)	
2	0 (0%)	2 (10%)	
3	0 (0%)	0 (0%)	
Use of sleep medication			>0.9
0	18 (95%)	17 (85%)	
1	1 (5.3%)	2 (10%)	
2	0 (0%)	1 (5.0%)	
3	0 (0%)	0 (0%)	
Daytime dysfunction			0.4
0	3 (16%)	7 (35%)	
1	11 (58%)	9 (45%)	
2	4 (21%)	4 (20%)	
3	1 (5.3%)	0 (0%)	
Global PSQI	5.00 (2.00,9.00)	4.00 (1.00,10.00)	0.8

[1] n (%); Median (Min,Max)

[2] *p<0.05; **p<0.01; ***p<0.001

Table S13. Pittsburgh Sleep Quality Index comparisons between Malaysia and Switzerland at Day 0. P-values are considered significant below 0.05.

Characteristic	Malaysia N = 19[1]	Switzerland N = 20[1]	p-value[2]
Subjective sleep quality			0.8
0	2 (11%)	3 (17%)	
1	13 (68%)	13 (72%)	
2	4 (21%)	2 (11%)	
3	0 (0%)	0 (0%)	
missing	0	2	
Sleep latency			0.7
0	8 (42%)	5 (28%)	
1	7 (37%)	9 (50%)	
2	2 (11%)	3 (17%)	
3	2 (11%)	1 (5.6%)	
missing	0	2	
Sleep duration			0.2
0	8 (42%)	13 (72%)	
1	8 (42%)	5 (28%)	
2	2 (11%)	0 (0%)	
3	1 (5.3%)	0 (0%)	
missing	0	2	
Habitual sleep efficiency			>0.9
0	16 (84%)	17 (94%)	
1	2 (11%)	1 (5.6%)	
2	1 (5.3%)	0 (0%)	
3	0 (0%)	0 (0%)	
missing	0	2	
Sleep disturbance			0.6
0	0 (0%)	0 (0%)	
1	18 (95%)	16 (89%)	
2	1 (5.3%)	2 (11%)	
3	0 (0%)	0 (0%)	
missing	0	2	
Use of sleep medication			>0.9
0	18 (95%)	18 (100%)	
1	1 (5.3%)	0 (0%)	
2	0 (0%)	0 (0%)	
3	0 (0%)	0 (0%)	
missing	0	2	
Daytime dysfunction			0.3
0	3 (16%)	7 (39%)	
1	11 (58%)	7 (39%)	
2	4 (21%)	4 (22%)	
3	1 (5.3%)	0 (0%)	
missing	0	2	
Global PSQI	5.00 (1.00,11.00)	3.50 (2.00,8.00)	0.15
missing	0	2	

[1] n (%); Median (Min,Max)
[2] *p<0.05; **p<0.01; ***p<0.001

Table S14. Pittsburgh Sleep Quality Index comparisons between Malaysia and Switzerland on Day 31. P-values are considered significant below 0.05.

Supplementary Figures

1170

1171

Figure S1. Summary of the bootstrapping procedure to determine an acceptable threshold of daily missing. Data based on three individual participants from the Malaysia site (MY004, MY009, MY012). Three variables were randomly chosen (IV, last time below 10 lux MEDI, MP ratio). The blue horizontal line shows the average value of data across the full dataset, the blue rectangle a 95% confidence interval around that average. Dots show the average value across all bootstraps of that respective threshold value. A blue dot indicates that all 10^4 bootstraps lie within the 95% confidence interval of the full dataset. A red dot indicates that at least one bootstrap lies outside of the 95% confidence interval. Abbreviations: IV; Interdaily variability; MP ratio, Melanopic/photopic ratio.

1182

Full Analysis Document

1183

1184 The full analysis document can be found on our GitHub page:

1185 https://github.com/tscnlab/BillerEtAl_RSocOpenSci_2025