Genome structural variation in natural *Caenorhabditis elegans* populations

Kyle J. Lesack, James D. Wasmuth[*]

Faculty of Veterinary Medicine, University of Calgary, Calgary, Alberta, Canada

* Corresponding author: jwasmuth@ucalgary.ca

Abstract

Structural variation, involving large alterations in chromosome structure, drives genetic diversification and the emergence of new phenotypes. These changes are widespread in natural populations and play an important role in adaptation and speciation. For many species, research has been limited to laboratory adapted strains and experimental evolution, which may not reflect the diversity of structural variants in the wild. Furthermore, technological limitations have proved to be a major barrier to accurate and comprehensive variant calling. In this study, we used PacBio sequencing data from 14 wild *Caenorhabditis elegans* strains to characterize structural variants in a natural population. With long-reads, we overcame limitations associated with short-read approaches and leveraged population-level data to further refine the accuracy of variant calls. We found that large, rapidly evolving gene families, such as GPCR, F-Box, and C-Type lectin, were prominent among the variants predicted to have phenotypic consequences. These results shed light on the significant role of structural variants in the evolution of *Caenorhabditis elegans* in its wild habitats and the limitations of treating N2 as the reference wild-type.

Keywords: structural variation; *Caenorhabditis elegans*; environmental adaptation

Introduction

Structural variants (SVs) are large-scale alterations in a genome—such as deletions, duplications, and inversions—that affect gene content, order, or dosage. These variants can have profound phenotypic effects by disrupting genes, altering gene dosage, or rearranging regulatory elements. While often deleterious, SVs can also drive adaptive evolution through mechanisms such as gene duplication followed by neofunctionalization or chromosomal inversions that suppress recombination and maintain beneficial allele combinations (Faria et al., 2019; Katju and Bergthorsson, 2013; Kirkpatrick, 2010).

Despite their importance, SVs remain poorly characterized in most species, especially at the population level. This is partly due to the technical limitations of short-read DNA sequencing, which struggles to resolve large or complex variants, particularly in repetitive regions (Koboldt, 2020; Lesack et al., 2022). Long-read sequencing has substantially improved the resolution of SVs and providing more reliable alignments (Smolka et al., 2024). In *Caenorhabditis elegans*, much of our knowledge about SVs comes from laboratory strains and mutation accumulation experiments. These studies have revealed a high rate of spontaneous duplications and the formation of complex alleles (Farslow et al., 2015; Konrad et al., 2018). For example, a recombinant inbred line showed increased fitness due to a partial duplication and inversion of five different regions of *rcan-1*, a gene which regulates thermotactic behavior (Zhao et al., 2020). However, laboratory strains, such as N2, carry numerous derived alleles and pseudogenised genes, limiting their relevance for understanding variation in natural populations (Andersen and Rockman, 2022; Sterken et al., 2015). Comparative studies between N2 and wild isolates, particularly the Hawaiian strain CB4856, have revealed extensive structural divergence

44   (Lee et al., 2021; Vergara et al., 2014). In 12 natural strains, more than 5% of the N2 genome is

45   affected by SVs, with CB4856 showing over 79,000 insertion/deletion events, more than 800

46   unaligned segments, and 61 hyper-divergent regions (Lee et al., 2021; Thompson et al., 2015).

47   The *Caenorhabditis elegans* Natural Diversity Resource (CeNDR) has expanded access

to genomic data from wild strains (Cook et al., 2017). This collection has facilitated the

discovery of genetic loci underlying xenobiotic metabolism and anthelmintic drug resistance

(Hahnel et al., 2018; Zdraljevic et al., 2019). In this study, we used PacBio long-read sequencing

data from 14 wild *C. elegans* strains to generate a high-confidence, population-scale map of SVs.

We identified thousands of deletions, duplications, and inversions—many affecting genes

involved in environmental response, stress, and immune function.

Materials and Methods

Strains

We analysed PacBio sequencing data from 14 *C. elegans* strains (DL238, ECA36, ECA396, EG4725, JU1400, JU2526, JU2600, JU310, MY2147, MY2693, NIC2, NIC526, QX1794, XZ1516) sourced from CeNDR and collected from Hawaii, New Zealand, and Western Europe (Figure S1).

Sequence alignment and structural variant calling

Reads were aligned to the N2 reference genome using NGMLR (v0.2.7), and alignments were sorted using Picard (v2.27.5) (Broad Institute, 2025; Sedlazeck et al., 2018). SVs were called using Sniffles2 (v2.0.7), filtered to include only deletions, duplications, and inversions ≥ 100bp (Smolka et al., 2024). SV calls were refined using Iris (v1.0.04) for improved breakpoint resolution and merged across strains using Jasmine (v1.1.5) and Iris (v1.0.4) (Kirsche et al., 2023).

SV annotation and functional profiling

We used Ensmbl's Variant Effect Predictor (v.109.3) to predict the functional consequences of each SV (McLaren et al., 2016). SVs classified as "high impact" were selected for functional enrichment using easyGSEA (v1.3.1) and annotated using WormCat (v2) (Cheng et al., 2021; Holdorf et al., 2020). Genes were assigned to statistically significant terms across WormCat's hierarchical levels (FDR ≤ 0.05), though no single level was exclusively used due to trade-offs in resolution and coverage.

Data presentation

SV densities along chromosomes were plotted using karyoploteR (v1.26.0) (Gel and Serra, 2017). SV allele frequencies were visualized UpSetPlot (v0.8.0) (Lex et al., 2014). Only the largest intersections—25 duplications, 20 for deletions and inversions—were displayed. The remaining plots were created using ggplot2 (v.3.4.2) (Wickham, 2016).

Data availability

The code, and Snakemake pipeline, used in this project is freely available:

https://github.com/kyleLesack/sv_analysis_cendr_pacbio

Results and Discussion

We identified 14,473 deletions, 3,149 duplications, and 1,748 inversions in 14 wild *C. elegans* strains (Table S1). SVs were unevenly distributed across the genome (Figure S2). Deletions and duplications were enriched at the autosomal arms, whereas inversions were more evenly distributed. These patterns are consistent with previous findings; SVs preferentially occur in low gene density, high recombination regions (Cutter et al., 2009). Chromosome arms may promote SV formation through recombination-based mechanisms while protecting essential genes located in denser chromosome centres (Carvalho and Lupski, 2016). Among all SVs, 64% were singletons—variants unique to a single strain—suggesting high inter-strain diversity (Table S2). Hawaiian strains had the highest number and proportion of singleton variants while Western European strains had the fewest (Table 1). The high ancestral diversity in Hawaiian strains is known, with reduced diversity in non-Hawaiian lineages due to selective sweeps (Crombie et al., 2019). Conversely, 25 variants were found in all 14 strains: 15 deletions, two duplications, and eight inversions (Table S2; Figures S3, S4, S5).

To prioritise variants with likely functional consequences, we focused on high-impact SVs. This subset included 1,485 deletions, 214 duplications, and three inversions, collectively affecting 1,258 genes (Figure 1; Tables S3, S4, S5). Deletions were enriched for splicing disruptions, transcript ablations, and frameshift variants (Table 2). Duplications primarily resulted in transcript amplifications, and all three high impact inversions were classified as stop/loss mutations. Functional enrichment revealed strong overrepresentation in categories related to environmental stress, including GPCRs, F-box proteins, C-type lectins (CLECs), and detoxification enzymes (Figure 2; Table S6). Many of these genes were in hyper-variable

105 regions, which have been shown to maintain ancestral variation through balancing selection (Lee

106 et al., 2021).

Strain	Region	Deletions			Duplications			Inversions		
		Total	Unique	Unique Proportion	Total	Unique	Unique Proportion	Total	Unique	Unique Proportion
DL238	Hawaii	2343	804	0.34	485	209	0.43	333	95	0.29
ECA396	Hawaii	3412	1643	0.48	519	300	0.58	399	140	0.35
QX1794	Hawaii	2363	756	0.32	459	186	0.41	311	74	0.24
XZ1516	Hawaii	4007	2440	0.61	443	266	0.60	519	301	0.58
ECA36	New Zealand	3522	1671	0.47	586	308	0.53	398	147	0.37
EG4725	Western Europe	1913	290	0.15	452	94	0.21	280	42	0.15
JU1400	Western Europe	1692	225	0.13	417	120	0.29	240	34	0.14
JU2526	Western Europe	2257	452	0.20	442	96	0.22	309	55	0.18
JU2600	Western Europe	1519	290	0.19	385	87	0.23	238	39	0.16
JU310	Western Europe	1366	170	0.12	385	72	0.19	217	16	0.07
MY2147	Western Europe	1279	217	0.17	364	85	0.23	199	29	0.15
MY2693	Western Europe	1251	97	0.08	385	84	0.22	258	35	0.14
NIC2	Western Europe	457	80	0.18	248	77	0.31	150	12	0.08
NIC526	Western Europe	1544	179	0.12	402	78	0.19	257	22	0.09

108 **Table 1:** Singleton variants among individual *C. elegans* strains.

Figure 1: Number of structural variants determined to have high functional impact.

Figure 2: Statistically significant overrepresented WormCat terms for putative high impact SVs. The red, green, and blue colours represent the categories for the C1, C2, and C3 levels, respectively.

SV Type	Term	Description‡	Count
Deletion (n = 1,485)	splice_donor_variant	A splice variant that changes the 2-base region at the 5' end of an intron	733
	splice_acceptor_variant	A splice variant that changes the 2-base region at the 3' end of an intron	724
	transcript_ablation	A feature ablation whereby the deleted region includes a transcript feature	411
	frameshift_variant	A sequence variant which causes a disruption of the translational reading frame, because the number of nucleotides inserted or deleted is not a multiple of three	223
	stop_lost	A sequence variant where at least one base of the terminator codon (stop) is changed, resulting in an elongated transcript	120
	start_lost	A codon variant that changes at least one base of the canonical start codon	20
	stop_gained	A sequence variant whereby at least one base of a codon is changed, resulting in a premature stop codon, leading to a shortened transcript	2
Duplication (n = 241)	transcript_amplification	A feature amplification of a region containing a transcript	240
	start_lost	A codon variant that changes at least one base of the canonical start codon	1
Inversion (n=3)	start_lost	A codon variant that changes at least one base of the canonical start codon	3

Table 2: Predicted Consequences of High Severity Structural Variants.

‡ Consequence descriptions downloaded on May 22, 2023 from:
https://useast.ensembl.org/info/genome/variation/prediction/predicted_data.html

GPCRs and sensory behaviour

Seven-transmembrane G-protein-coupled receptors (GPCRs), involved in chemosensation and behaviour, were among the most affected gene families. Of the 253 genes in the "Transmembrane protein" category, 152 were GPCRs. The SRH family was enriched, with 36 genes, which is consistent with previous reports of rapid expansion and diversification in *C. elegans* (Thomas and Robertson, 2008; Vidal et al., 2018). A duplication of *srh-234* was present

in two strains. The expression of *srh-234* is regulated by the chemosensation of food (Gruner et

al., 2014). Two deletions are GPCRs were noteworthy: *sri-14* is involved in response to high and

toxic concentrations of diacetyl (Taniguchi et al., 2014); and *str-208* is implicated in erratic

movement (Daigle et al., 2022). While the loss of these genes may reduce responsiveness to

adverse stimuli or affect locomotion, it is not yet known whether the associated structural

variants confer an adaptive benefit, impose a cost, or are selectively neutral.

F-box genes and immune adaptation

SVs affected 128 proteolysis-related genes, including 74 F-box genes. The F-box gene

family is considerably expanded in *C. elegans* relative to other *Caenorhabditis* species with

several members under positive selection in the non-Hawaiian strains (Ma et al., 2021). One of

these is *fbxa-158*, which promoted a thermotolerance-associated intracellular pathogen response,

but we found it was lost in three non-Hawaiian strains, EG4725, JU1400, and MY2693, and one

Hawaiian strain, ECA396 (Panek et al., 2020). Additional immune-related SVs may reflect

adaptations to local microbial environments. For example, *hecw-1*, deleted in the Hawaiian

strains ECA396 and ZX1516, is an E3 ubiquitin ligase that modulate pathogen avoidance (Chang

et al., 2011).

C-type lectins and pathogen response

Thirty-one C-type lectin (CLEC) genes were affected by SVs. CLECs are a rapidly

evolving gene family involved in innate immunity and pathogen recognition (Pees et al., 2016).

Seven CLECs were affected by deletions—*clec-2, clec-10, clec-13, clec-67, clec-122, clec-143,*

and *clec-174*—all known to be modulated by bacterial pathogens (Pan et al., 2021; Sahu et al.,

2012; White and Herman, 2018). A further three CLECs were duplicated; *clec-70* is induced by

Microbacterium nematophilum and *Staphylococcus aureus*, while *clec-189* and *clec-258* await

functional annotation but may represent novel immune effectors (Pan et al., 2021).

Detoxification and xenobiotic response

Several families of detoxification genes were enriched in SVs, including cytochrome P450s

(CYPs), glutathione S-transferases (GSTs), UDP-glucuronosyltransferases (UGTs), and nuclear

hormone receptors (NHRs). Two caffeine inducible CYPs, *cyp-33e2* and *cyp-35a4* harbored

deletions; the latter gene also regulates fat storage (Aarnio et al., 2011; Min et al., 2015). Other

deletion-affected CYP genes included: *cyp-33c5* and *cyp-13a5*, which are involved in cadmium

detoxification; and *cyp-33d3*, which is inducible by ethidium bromide and low pH (Cong et al.,

2020; Cui et al., 2007). In other families, notable deletions included: *ugt-14* is upregulated in

response to benzimidazole-class anthelmintics (Stasiuk et al., 2019); and *nhr-196* is regulated by

exposure to the mycotoxin aflatoxin B1 (Karengera et al., 2022).

Evolutionary and Functional Implications

Our results highlight the utility of long-read sequencing for characterizing SVs at a

population scale. The high diversity of SVs in wild *C. elegans* populations—particularly those in

gene families related to sensory perception, immunity, and detoxification—emphasizes the

limitations of relying solely on N2 as a wild-type reference (Andersen and Rockman, 2022;

Sterken et al., 2015). Many of the identified SVs may confer adaptive advantages, particularly in

geographically and microbially diverse environments.

Importantly, many SVs overlap with hyper-divergent regions—genomic hotspots for

adaptive diversity—shown in *C. elegans* and the parasitic *Heligmosomoides polygyrus* to contain

genes involved in sensory perception and stress response (Lee et al., 2021; Stevens et al., 2023).

Going forward, hybrid approaches that integrate long- and short-read sequencing and

transcriptomic data from wild isolates will help resolve the regulatory and phenotypic

consequences of SVs (Zhang et al., 2022). In particular, eQTL mapping that includes SVs could

offer insights into complex traits not explained by single-nucleotide variation alone (Boyling et

al., 2022). Additionally, functional studies targeting SVs genes currently categorized as

"Unassigned" may reveal previously uncharacterized molecular mechanisms.

Acknowledgments

We thank Dr. Erik Andersen and members of the Andersen lab (Johns Hopkins University) for discussions on using CeNDR datasets. We also acknowledge the high-performance computing resources made available by Research Computing and the Faculty of Veterinary Medicine at the University of Calgary.

Funding

This work was funded through grants to JDW from the Natural Sciences and Engineering Research Council of Canada (NSERC; #04589-2020).

References

Aarnio, V., Lehtonen, M., Storvik, M., Callaway, J.C., Lakso, M., Wong, G., 2011.
Caenorhabditis Elegans Mutants Predict Regulation of Fatty Acids and Endocannabinoids
by the CYP-35A Gene Family. Front. Pharmacol. 2.
https://doi.org/10.3389/fphar.2011.00012

Andersen, E.C., Rockman, M.V., 2022. Natural genetic variation as a tool for discovery in
Caenorhabditis nematodes. Genetics 220, iyab156.
https://doi.org/10.1093/genetics/iyab156

Boyling, A., Perez-Siles, G., Kennerson, M.L., 2022. Structural Variation at a Disease Mutation
Hotspot: Strategies to Investigate Gene Regulation and the 3D Genome. Front. Genet. 13,
842860. https://doi.org/10.3389/fgene.2022.842860

Broad Institute, 2025. broadinstitute/picard.

Carvalho, C.M.B., Lupski, J.R., 2016. Mechanisms underlying structural variant formation in
genomic disorders. Nat Rev Genet 17, 224–238. https://doi.org/10.1038/nrg.2015.25

Chang, H.C., Paek, J., Kim, D.H., 2011. Natural polymorphisms in C. elegans HECW-1 E3
ligase affect pathogen avoidance behaviour. Nature 480, 525–529.
https://doi.org/10.1038/nature10643

Cheng, X., Yan, J., Liu, Y., Wang, J., Taubert, S., 2021. eVITTA: a web-based visualization and
inference toolbox for transcriptome analysis. Nucleic Acids Research 49, W207–W215.
https://doi.org/10.1093/nar/gkab366

Cong, Y., Yang, H., Zhang, P., Xie, Y., Cao, X., Zhang, L., 2020. Transcriptome Analysis of the

Nematode Caenorhabditis elegans in Acidic Stress Environments. Front. Physiol. 11,

1107. https://doi.org/10.3389/fphys.2020.01107

Cook, D.E., Zdraljevic, S., Roberts, J.P., Andersen, E.C., 2017. CeNDR, the Caenorhabditis

elegans natural diversity resource. Nucleic Acids Res 45, D650–D657.

https://doi.org/10.1093/nar/gkw893

Crombie, T.A., Zdraljevic, S., Cook, D.E., Tanny, R.E., Brady, S.C., Wang, Y., Evans, K.S.,

Hahnel, S., Lee, D., Rodriguez, B.C., Zhang, G., Van Der Zwagg, J., Kiontke, K.,

Andersen, E.C., 2019. Deep sampling of Hawaiian Caenorhabditis elegans reveals high

genetic diversity and admixture with global populations. eLife 8, e50465.

https://doi.org/10.7554/eLife.50465

Cui, Y., McBride, S.J., Boyd, W.A., Alper, S., Freedman, J.H., 2007. Toxicogenomic analysis of

Caenorhabditis elegans reveals novel genes and pathways involved in the resistance to

cadmium toxicity. Genome Biol 8, R122. https://doi.org/10.1186/gb-2007-8-6-r122

Cutter, A.D., Dey, A., Murray, R.L., 2009. Evolution of the Caenorhabditis elegans Genome.

Molecular Biology and Evolution 26, 1199–1234.

https://doi.org/10.1093/molbev/msp048

Daigle, A.T., Deiss, T.C., Melde, R.H., Bergthorsson, U., Katju, V., 2022. Bergerac strains of

Caenorhabditis elegans revisited: expansion of Tc *1* elements imposes a significant

genomic and fitness cost. G3 Genes|Genomes|Genetics 12, jkac214.

https://doi.org/10.1093/g3journal/jkac214

Faria, R., Johannesson, K., Butlin, R.K., Westram, A.M., 2019. Evolving Inversions. Trends in

Ecology & Evolution 34, 239–248. https://doi.org/10.1016/j.tree.2018.12.005

234 Farslow, J.C., Lipinski, K.J., Packard, L.B., Edgley, M.L., Taylor, J., Flibotte, S., Moerman,

235   D.G., Katju, V., Bergthorsson, U., 2015. Rapid Increase in frequency of gene copy-

236   number variants during experimental evolution in Caenorhabditis elegans. BMC

237   Genomics 16, 1044. https://doi.org/10.1186/s12864-015-2253-2

238 Gel, B., Serra, E., 2017. karyoploteR: an R/Bioconductor package to plot customizable genomes

239   displaying arbitrary data. Bioinformatics 33, 3088–3090.

240   https://doi.org/10.1093/bioinformatics/btx346

241 Gruner, M., Nelson, D., Winbush, A., Hintz, R., Ryu, L., Chung, S.H., Kim, K., Gabel, C.V., Van

242   Der Linden, A.M., 2014. Feeding State, Insulin and NPR-1 Modulate Chemoreceptor

243   Gene Expression via Integration of Sensory and Circuit Inputs. PLoS Genet 10,

244   e1004707. https://doi.org/10.1371/journal.pgen.1004707

245 Hahnel, S.R., Zdraljevic, S., Rodriguez, B.C., Zhao, Y., McGrath, P.T., Andersen, E.C., 2018.

246   Extreme allelic heterogeneity at a Caenorhabditis elegans beta-tubulin locus explains

247   natural resistance to benzimidazoles. PLoS Pathog 14, e1007226.

248   https://doi.org/10.1371/journal.ppat.1007226

249 Holdorf, A.D., Higgins, D.P., Hart, A.C., Boag, P.R., Pazour, G.J., Walhout, A.J.M., Walker,

250   A.K., 2020. WormCat: An Online Tool for Annotation and Visualization of

251   *Caenorhabditis elegans* Genome-Scale Data. Genetics 214, 279–294.

252   https://doi.org/10.1534/genetics.119.302919

253 Karengera, A., Sterken, M.G., Kammenga, J.E., Riksen, J.A.G., Dinkla, I.J.T., Murk, A.J., 2022.

254   Differential expression of genes in C. elegans reveals transcriptional responses to

255   indirect-acting xenobiotic compounds and insensitivity to 2,3,7,8-

tetrachlorodibenzodioxin. Ecotoxicology and Environmental Safety 233, 113344.

https://doi.org/10.1016/j.ecoenv.2022.113344

Katju, V., Bergthorsson, U., 2013. Copy-number changes in evolution: rates, fitness effects and

adaptive significance. Front. Genet. 4. https://doi.org/10.3389/fgene.2013.00273

Kirkpatrick, M., 2010. How and Why Chromosome Inversions Evolve. PLoS Biol 8, e1000501.

https://doi.org/10.1371/journal.pbio.1000501

Kirsche, M., Prabhu, G., Sherman, R., Ni, B., Battle, A., Aganezov, S., Schatz, M.C., 2023.

Jasmine and Iris: population-scale structural variant comparison and analysis. Nat

Methods 20, 408–417. https://doi.org/10.1038/s41592-022-01753-3

Koboldt, D.C., 2020. Best practices for variant calling in clinical sequencing. Genome Med 12,

91. https://doi.org/10.1186/s13073-020-00791-w

Konrad, A., Flibotte, S., Taylor, J., Waterston, R.H., Moerman, D.G., Bergthorsson, U., Katju, V.,

2018. Mutational and transcriptional landscape of spontaneous gene duplications and

deletions in *Caenorhabditis elegans*. Proc. Natl. Acad. Sci. U.S.A. 115, 7386–7391.

https://doi.org/10.1073/pnas.1801930115

Lee, D., Zdraljevic, S., Stevens, L., Wang, Y., Tanny, R.E., Crombie, T.A., Cook, D.E., Webster,

A.K., Chirakar, R., Baugh, L.R., Sterken, M.G., Braendle, C., Félix, M.-A., Rockman,

M.V., Andersen, E.C., 2021. Balancing selection maintains hyper-divergent haplotypes in

Caenorhabditis elegans. Nat Ecol Evol 5, 794–807. https://doi.org/10.1038/s41559-021-

01435-x

Lesack, K., Mariene, G.M., Andersen, E.C., Wasmuth, J.D., 2022. Different structural variant

prediction tools yield considerably different results in Caenorhabditis elegans. PLOS

ONE 17, e0278424. https://doi.org/10.1371/journal.pone.0278424

Lex, A., Gehlenborg, N., Strobelt, H., Vuillemot, R., Pfister, H., 2014. UpSet: Visualization of

Intersecting Sets. IEEE Trans Vis Comput Graph 20, 1983–1992.

https://doi.org/10.1109/TVCG.2014.2346248

Ma, F., Lau, C.Y., Zheng, C., 2021. Large Genetic Diversity and Strong Positive Selection in F-

Box and GPCR Genes among the Wild Isolates of Caenorhabditis elegans. Genome Biol

Evol 13, evab048. https://doi.org/10.1093/gbe/evab048

McLaren, W., Gil, L., Hunt, S.E., Riat, H.S., Ritchie, G.R.S., Thormann, A., Flicek, P.,

Cunningham, F., 2016. The Ensembl Variant Effect Predictor. Genome Biol 17, 122.

https://doi.org/10.1186/s13059-016-0974-4

Min, H., Kawasaki, I., Gong, J., Shim, Y.-H., 2015. Caffeine Induces High Expression of cyp-

35A Family Genes and Inhibits the Early Larval Development in Caenorhabditis elegans.

Molecules and Cells 38, 236–242. https://doi.org/10.14348/molcells.2015.2282

Pan, W., Huang, X., Guo, Z., Nagarajan, R., Mylonakis, E., 2021. Identification and Functional

Analysis of Cytokine-Like Protein CLEC-47 in Caenorhabditis elegans. mBio 12,

e02579-21. https://doi.org/10.1128/mBio.02579-21

Panek, J., Gang, S.S., Reddy, K.C., Luallen, R.J., Fulzele, A., Bennett, E.J., Troemel, E.R., 2020.

A cullin-RING ubiquitin ligase promotes thermotolerance as part of the intracellular

pathogen response in *Caenorhabditis elegans*. Proc. Natl. Acad. Sci. U.S.A. 117, 7950–

7960. https://doi.org/10.1073/pnas.1918417117

Pees, B., Yang, W., Zárate-Potes, A., Schulenburg, H., Dierking, K., 2016. High Innate Immune

Specificity through Diversified C-Type Lectin-Like Domain Proteins in Invertebrates. J

Innate Immun 8, 129–142. https://doi.org/10.1159/000441475

Sahu, S.N., Lewis, J., Patel, I., Bozdag, S., Lee, J.H., LeClerc, J.E., Cinar, H.N., 2012. Genomic

Analysis of Immune Response against Vibrio cholerae Hemolysin in Caenorhabditis

elegans. PLoS ONE 7, e38200. https://doi.org/10.1371/journal.pone.0038200

Sedlazeck, F.J., Rescheneder, P., Smolka, M., Fang, H., Nattestad, M., von Haeseler, A., Schatz,

M.C., 2018. Accurate detection of complex structural variations using single-molecule

sequencing. Nat Methods 15, 461–468. https://doi.org/10.1038/s41592-018-0001-7

Smolka, M., Paulin, L.F., Grochowski, C.M., Horner, D.W., Mahmoud, M., Behera, S., Kalef-

Ezra, E., Gandhi, M., Hong, K., Pehlivan, D., Scholz, S.W., Carvalho, C.M.B., Proukakis,

C., Sedlazeck, F.J., 2024. Detection of mosaic and population-level structural variants

with Sniffles2. Nat Biotechnol 1–10. https://doi.org/10.1038/s41587-023-02024-y

Stasiuk, S.J., MacNevin, G., Workentine, M.L., Gray, D., Redman, E., Bartley, D., Morrison, A.,

Sharma, N., Colwell, D., Ro, D.K., Gilleard, J.S., 2019. Similarities and differences in the

biotransformation and transcriptomic responses of Caenorhabditis elegans and

Haemonchus contortus to five different benzimidazole drugs. Int J Parasitol Drugs Drug

Resist 11, 13–29. https://doi.org/10.1016/j.ijpddr.2019.09.001

Sterken, M.G., Snoek, L.B., Kammenga, J.E., Andersen, E.C., 2015. The laboratory

domestication of Caenorhabditis elegans. Trends in Genetics 31, 224–231.

https://doi.org/10.1016/j.tig.2015.02.009

Stevens, L., Martínez-Ugalde, I., King, E., Wagah, M., Absolon, D., Bancroft, R., Gonzalez de la

Rosa, P., Hall, J.L., Kieninger, M., Kloch, A., Pelan, S., Robertson, E., Pedersen, A.B.,

Abreu-Goodger, C., Buck, A.H., Blaxter, M., 2023. Ancient diversity in host-parasite

interaction genes in a model parasitic nematode. Nat Commun 14, 7776.

https://doi.org/10.1038/s41467-023-43556-w

Taniguchi, G., Uozumi, T., Kiriyama, K., Kamizaki, T., Hirotsu, T., 2014. Screening of Odor-

Receptor Pairs in *Caenorhabditis elegans* Reveals Different Receptors for High and Low

Odor Concentrations. Sci. Signal. 7. https://doi.org/10.1126/scisignal.2005136

Thomas, J.H., Robertson, H.M., 2008. The Caenorhabditis chemoreceptor gene families. BMC

Biol 6, 42. https://doi.org/10.1186/1741-7007-6-42

Thompson, O.A., Snoek, L.B., Nijveen, H., Sterken, M.G., Volkers, R.J.M., Brenchley, R., Van'T

Hof, A., Bevers, R.P.J., Cossins, A.R., Yanai, I., Hajnal, A., Schmid, T., Perkins, J.D.,

Spencer, D., Kruglyak, L., Andersen, E.C., Moerman, D.G., Hillier, L.W., Kammenga,

J.E., Waterston, R.H., 2015. Remarkably Divergent Regions Punctuate the Genome

Assembly of the *Caenorhabditis elegans* Hawaiian Strain CB4856. Genetics 200, 975–

989. https://doi.org/10.1534/genetics.115.175950

Vergara, I.A., Tarailo-Graovac, M., Frech, C., Wang, J., Qin, Z., Zhang, T., She, R., Chu, J.S.,

Wang, K., Chen, N., 2014. Genome-wide variations in a natural isolate of the nematode

Caenorhabditis elegans. BMC Genomics 15, 255. https://doi.org/10.1186/1471-2164-15-

255

Vidal, B., Aghayeva, U., Sun, H., Wang, C., Glenwinkel, L., Bayer, E.A., Hobert, O., 2018. An

atlas of Caenorhabditis elegans chemoreceptor expression. PLoS Biol 16, e2004218.

https://doi.org/10.1371/journal.pbio.2004218

White, C.V., Herman, M.A., 2018. Transcriptomic, Functional, and Network Analyses Reveal

Novel Genes Involved in the Interaction Between Caenorhabditis elegans and

Stenotrophomonas maltophilia. Front. Cell. Infect. Microbiol. 8, 266.

https://doi.org/10.3389/fcimb.2018.00266

Wickham, H., 2016. ggplot2, Use R! Springer International Publishing, Cham.

https://doi.org/10.1007/978-3-319-24277-4

Zdraljevic, S., Fox, B.W., Strand, C., Panda, O., Tenjo, F.J., Brady, S.C., Crombie, T.A., Doench,

J.G., Schroeder, F.C., Andersen, E.C., 2019. Natural variation in C. elegans arsenic

toxicity is explained by differences in branched chain amino acid metabolism. eLife 8,

e40260. https://doi.org/10.7554/eLife.40260

Zhang, G., Roberto, N.M., Lee, D., Hahnel, S.R., Andersen, E.C., 2022. The impact of species-

wide gene expression variation on Caenorhabditis elegans complex traits. Nat Commun

13, 3462. https://doi.org/10.1038/s41467-022-31208-4

Zhao, Y., Long, L., Wan, J., Biliya, S., Brady, S.C., Lee, D., Ojemakinde, A., Andersen, E.C.,

Vannberg, F.O., Lu, H., McGrath, P.T., 2020. A spontaneous complex structural variant in

rcan-1 increases exploratory behavior and laboratory fitness of Caenorhabditis elegans.

PLoS Genet 16, e1008606. https://doi.org/10.1371/journal.pgen.1008606

362