Ecology

Article

Nonlinear effects of temperature on mosquito parasite infection across a large geographic climate

gradient

Johannah E. Farner[1*], Kelsey P. Lyberger[2], Lisa I. Couper[3], Mauricio Cruz-Loya[1], Erin A.

Mordecai[1]

[1]Department of Biology, Stanford University, Stanford, CA 94305, USA; [2] College of Integrative

Sciences and Arts, Arizona State University, Mesa, AZ 85212, USA; [3] Division of Environmental

Health Sciences, University of California, Berkeley, Berkeley, CA 94720, USA.

*Corresponding author: jfarner@stanford.edu

Data and code for this submission are provided on

Dryad: http://datadryad.org/stash/share/CfZkk4LsJzljetJJnFZMDMrjuciTXMxrkrc95I2J3tA.

Abstract

Temperature drives ectothermic host – parasite interactions, making them particularly sensitive to climatic variation and change. To isolate the role of temperature, lab-based studies are increasingly used to assess and forecast disease risk under current and future climate conditions. However, in the field, the effects of temperature on parasitism may be mediated by other sources of variation, including local adaptation. To address the key knowledge gaps of how temperature influences host – parasite interactions and whether thermal responses measured in controlled experiments capture infection across temperature gradients in nature, we paired an extensive field survey of parasitism—by the ciliate *Lambornella clarki* on its tree hole mosquito host, *Aedes sierrensis*—with laboratory experiments describing parasitism thermal performance curves (TPCs) for six host populations from varying climates. We also investigated the mechanisms underlying the thermal biology of the host – parasite interaction by separately measuring TPCs for infection, host immunity, and parasite growth rates. Along the west coast of North America, across an 1100 km climate gradient spanning 12ºC mean rainy season temperature variation, we found that parasitism peaked at intermediately cold temperatures, and was consistent both between field seasons and with the lab experiment results. The experiments produced no evidence of host intraspecific variation in temperature sensitivity to parasitism. Importantly, parasitism peaked at temperatures below the thermal optimum for free-living *L. clarki* due to the balance of temperature effects on parasite growth and reproduction against the strength of the host melanization immune response. The results suggest that nonlinear responses to temperature drive parasitism in nature, and that simple lab and field studies can accurately capture the thermal biology of multilayered host – parasite interactions.

Introduction

Climate warming is fundamentally important to disease risk because temperature drives many of the constituent biological processes of host – parasite interactions, including host and parasite growth and development, contact rates, and immune functions (Kutz et al. 2005; Labaude et al. 2017; Laughton, O'Connor, and Knell 2017; Franke et al. 2019; Mordecai et al. 2019). Strong relationships between temperature and parasitism are frequently observed in laboratory experiments and captured in thermal performance curves (TPCs) (Gehman, Hall, and Byers 2018; Kirk et al. 2018; Padfield, Castledine, and Buckling 2020), which researchers then use to assess disease risk based on current and forecasted climate and weather conditions (Gehman, Hall, and Byers 2018; Ryan et al. 2019; Aleuy et al. 2023). However, it is unclear whether TPCs measured in controlled laboratory settings fully capture the thermal response of disease in complex natural settings, where other abiotic and biotic factors may modify or outweigh parasitism - temperature relationships (Mordecai et al. 2019; Taylor et al. 2019; Brown et al. 2023).

Parasitism TPCs are typically measured for single laboratory populations, under constant temperatures, in the absence of other stressors, and are infrequently compared to disease burdens along temperature gradients in the field (e.g., Gehman, Hall and Byers, 2018; Mordecai *et al.*, 2019; Padfield, Castledine and Buckling, 2020). Although an increasing number of studies describe thermal responses for multiple populations or genotypes within host-parasite systems (Mazé-Guilmo et al. 2016; Ahrens et al. 2019; Franke et al. 2019; Aleuy et al. 2023; Ragonese et al. 2024), research on statistical or correlative relationships between TPCs for disease transmissibility (e.g., R_0) and natural infection patterns has thus far relied on laboratory measurements from single populations or clones, and field data collated from disparate sources

and/or collected over small geographic areas. To date, this research has focused primarily on

mosquito-borne human disease and fungal pathogens of zooplankton (Mordecai et al. 2013;

2017; Shocket et al. 2018; 2020; Tesla et al. 2018; Shah et al. 2019; Kirk et al. 2024). However,

TPCs can vary geographically within species due to local adaptation and population structure

(Sinclair, Williams, and Terblanche 2012; Gaitán-Espitia et al. 2014; Mazé-Guilmo et al. 2016;

Jurriaans and Hoogenboom 2019; Aleuy et al. 2023; Couper et al. 2024), and additional biotic

and abiotic environmental variability in field settings may outweigh or modify the temperature

dependence of parasitism (e.g., Boots 2000; Seppälä et al. 2008; Hall et al. 2009; Penczykowski

et al. 2014; Palmer-Young et al. 2019; Brown et al. 2023). Thus, there is a critical need to

address the questions of how temperature affects parasitism in the field, whether lab-based TPC

estimates can accurately, and mechanistically, capture these patterns, and whether local

adaptation modifies the temperature-dependence of parasitism.

Central to capturing the thermal biology of parasitism is understanding the host and

parasite mechanisms that drive the interaction. For example, host life history and defense traits

and parasite growth, attack, dispersal, and transmission traits may each respond nonlinearly and

distinctly to temperature. The combined effect of these traits determines the overall impact of

temperature on parasitism (Paull, LaFonte, and Johnson 2012; Shocket et al. 2018; Mordecai et

al. 2019; Sun et al. 2023). Moreover, host and parasite traits may differ across populations due to

local adaptation and across environmental conditions (Thomas and Blanford 2003; Sternberg and

Thomas 2014). Given that it is not feasible to measure TPCs for every possible combination of

host population, parasite population, and environmental context, it is important to establish

whether TPCs measured for a single set of populations and conditions can generalize to cover

entire species ranges, and which traits capture the essential thermal biology of the interaction.

Here, we address these questions in a highly tractable experimental model system of a mosquito

(*Aedes sierrensis*) and its facultative ciliate parasite (*Lambornella clarki*), which occur across a

large geographic gradient in western North America.

In this study, we pair field surveys along the entire west coast of the continental United

States—spanning the geographic range of the host species—with laboratory experiments

describing the temperature dependence of infection and key traits for host populations from

different climates across the survey area. Both species are short-lived ectotherms that are directly

affected by environmental temperatures, occur commonly across a large geographic climate

gradient in discrete tree hole habitats that are naturally replicated across landscapes, and can be

reared separately and together in the lab (Washburn and Anderson 1986; Broberg and Bradshaw

1997; Darsie and Ward 2005; Ismail et al. 2023; Lyberger et al. 2024). A 1986 field survey

found that infection occurrence increased from southern to northern California (Washburn and

Anderson 1986). Our recent work describing temperature effects on different populations of *Ae.*

sierrensis, L. clarki, and their interactions indicates that infection has a consistent nonlinear

relationship with temperature for both sympatric and allopatric pairs of host and parasite

populations collected across climates, despite evidence of local thermal adaptation in free-living

L. clarki populations (Ismail et al. 2023; Lyberger et al. 2024). This body of work generates the

hypothesis that parasitism responds to temperature as a function of the thermal biology of

multiple host and parasite traits. However, it is unknown whether the lab-measured temperature

impacts on parasitism across different *Ae. sierrensis* host populations are borne out in infection

occurrence across the environmentally complex species range.

Methods

Study system

        Ae. sierrensis and its facultative ciliate parasite *L. clarki* occur across western North

America in seasonally rainwater-filled tree holes, where mosquitoes undergo one reproductive

cycle annually in which larvae feed on detritus and microbes (including free-living *L. clarki*)

before pupating and emerging as adults, and free-living *L. clarki* ciliates feed on aquatic bacteria

(Washburn and Anderson 1986; Darsie and Ward, 2005). In the presence of *Ae. sierrensis* larvae,

free-living *L. clarki* irreversibly transform into parasites that attack larval mosquitoes and often

kill them after a few weeks (Washburn et al., 1988). Mosquitoes can resist infection through

multiple defenses that each likely respond nonlinearly to temperature: melanization (a

generalized immune response), molting between larval instars, developing into pupae, and

consuming free-living *L. clarki* in the water column (Clark and Brandl 1976; Washburn et al.

1988; Washburn, Anderson, and Mercer 1991; Suwanchaichinda and Paskewitz 1998;

Christensen et al. 2005; Murdock et al. 2012; Martin and Hillyer 2024). Beyond the previously

documented latitudinal gradient, *L. clarki* presence is not correlated with habitat characteristics

such as tree hole size, exposure, height above ground, or tree species (Washburn and Anderson

1986).

Field survey

    To document infection patterns over a large geographic temperature gradient and collect

materials for lab experiments, we sampled from over 500 water-filled tree hole habitats

containing *Ae. sierrensis* larvae spanning 1100 km north to south with a mean November – April

rainy season temperature gradient of 2.6ºC to 15.3ºC in California, Oregon, and Washington,

USA, during the rainy seasons of 2022 and 2023 (Figure 1) (PRISM Climate Group, 2021). Each

year, we started in the warmest and ended in the coolest locations to track the seasonal timing of

Ae. sierrensis egg hatching and development (Jordan and Bradshaw 1978; Jordan 1980a; 1980b),

collecting larval samples and examining them for infection at 10-40x magnification (Appendix

S1). We aimed to sample each tree hole in both years, but had to replace some locations in 2023

due to storm damage (Figure 1).

For each sampling site, we used PRISM climate data to characterize long-term mean

temperature and precipitation conditions in the November – April rainy season when mosquito

larvae and parasites are active in tree holes, and Rainy and Dry Season (RADS) data to

characterize the median wet season duration (Bombardi, Kinter, and Frauenfeld 2019; PRISM

Climate Group and Oregon State University 2021; Washburn and Anderson 1986; Luković et al.

2021; Praskievicz and Chang 2009). We also placed iButton temperature loggers (Maxim

Integrated, San Jose, CA, USA) in 84 arbitrarily selected tree holes across the survey area from

2022 to 2023 to assess how local temperatures compared to PRISM. Finally, we accessed NOAA

climate reports that compared regional temperatures and precipitation in each field season to

long-term averages (NOAA National Centers for Environmental Information 2022; 2023).

Additional details on the field survey methods are provided in Appendix S1.

Lab experiments

1. Temperature-dependent infection experiment

In September 2022, we measured the effects of temperature on the host – parasite interaction

with an experiment that used the F1 offspring of six host populations reared in the lab from

larvae collected in the 2022 field survey: two each from warm (14.8 – 15.1ºC rainy season mean

temperature; San Juan Capistrano and Trabuco Canyon, CA), intermediate (10.8 – 11.8ºC; Big

Sur and Woodside, CA), and cool (5.3 – 6.5ºC; Shaver Lake, CA and Eugene, OR) climates

(Figure 1, Appendix S2: Table S1). We exposed each host population to the same allopatric

parasite culture isolated from an infected larva collected in Marin County, CA (10.6ºC) (Figure

1), so that any variation observed in the temperature dependence of infection would be due to the

host population. Mosquito rearing and ciliate culturing occurred at 21.5ºC following Ismail *et al.*

(2023), Couper *et al.* (2024), and Lyberger *et al.* (2024).

For the experiment, mosquito eggs were hatched and day-old larvae were incubated in groups

of five with 7 mL parasite culture diluted to a standardized density of 35 ciliates per 100 uL at six

constant temperatures: 7ºC, 11ºC, 14ºC, 18ºC, 23ºC, and 28ºC (N = 6 microcosms per host

population per temperature, except for Big Sur, where N = 4 due to lower egg hatching). We

monitored the proportions of hosts with infections and parasite melanization spots, and the

survival of infected and uninfected hosts, up to the first pupation at each temperature. Because

Ae. sierrensis development speed is thermally sensitive, we set monitoring frequences that

ensured at least four checkpoints spanning the host's four larval instars at each temperature; these

ranged from every two days at 28ºC to every seven days at 7ºC and 11ºC (Appendix S2: Table

S2).

2. *Host and parasite thermal responses in isolation*

To compare the thermal performance of mosquito larvae and ciliates to that of parasitism, we

incubated mosquito-only and ciliate-only microcosms alongside the infection experiment,

following the same design as above. The mosquito-only microcosms were monitored on the

same timeline as the infection experiment. For the ciliate-only microcosms, we measured the

population growth rate by inoculating wells with culture diluted to 4 cells per 100 uL and then

counting the cells per 300 uL daily for four days of positive growth.

To assess the temperature sensitivity of the host melanization process (removing the

temperature sensitivity of parasite attack rates), we performed a mechanical melanization assay

on fourth instar larvae from one population at 4, 8, 12, 16, 20, 24, and 28ºC (N = 6 wells of 5

mosquitoes per temperature with no ciliates). Additional details are provided in Appendix S1.

Statistical analyses

Field survey

To investigate the nonlinear temperature dependence of parasitism, we tested for a quadratic

relationship between temperature and the presence of *L. clarki* infections in the surveyed tree

holes using a binomial generalized linear mixed effects model, with mean temperature, mean

temperature squared, and year as fixed effects, and tree hole ID as a random effect. The final

model did not include precipitation metrics because of strong correlations between temperature

and precipitation variables (R^2 = -0.8 for both cumulative precipitation and median wet season

duration), and because we expect temperature to more directly affect the host-parasite

interaction, as precipitation was sufficient across all sampled locations for tree holes to fill with

water and *Ae. sierrensis* to occur. We used the glmer function in the R package "lme4" with

scaled and centered environmental variables, and assessed model fit with the R package

"DHARMa" (Bates et al. 2003; Hartig 2017). Because interactions between survey year and the

temperature variables were not significant, we estimated the overall mean temperature where

infection was most likely to occur with a model that did not include year. We additionally used

general linearized models fit individually to the data from each survey year to describe the

temperatures where infection occurrence peaked within each field season. To evaluate the

relationship between within-tree hole habitat temperatures logged by iButtons and the PRISM air

temperature dataset used in the model, we tested for correlations and calculated the mean

differences between daily mean temperature time series recorded by iButtons and extracted from

PRISM. Because there were only two field seasons, the NOAA annual climate reports for each

field season were not included in statistical analyses, but used only to qualitatively compare the

two years.

Lab experiments

To quantify the thermal sensitivity of the host – parasite interaction measured from the

laboratory experiment, we used a Bayesian approach to fit TPCs for the proportion of infected

hosts, the proportion of hosts with melanization spots, and the proportion of hosts killed by

infections. The total number of infections observed in each microcosm over the course of the

experiment depended on both initial parasite attack success and subsequent rounds of parasite

transmission following infected host deaths that released new parasites as killed mosquitoes

decomposed. We fit separate TPCs for both initial infection for the period up to the first deaths of

infected hosts at each temperature, when all observed infections resulted from the first round of

host – parasite interactions (covering checkpoints 1-3, approximately corresponding to larval

instars 1-3), and total infection rates over the entire experiment (including all host-parasite

interaction cycles for all four larval instars); we used the same procedure to fit initial and total

TPCs for the proportions of hosts melanizing attacking parasites. We additionally fit TPCs for

free-living *L. clarki* population growth rate and for the host melanization immune response

induced by mechanical penetration of the cuticle. We planned to fit TPCs for host survival in the

absence of parasitism, but >99% of larvae incubated without parasites survived across all

temperatures. Because initial parasite success is likely to determine whether there is a non-zero

infection probability over the rest of the season (Washburn, Anderson, and Mercer 1991), this is

the laboratory-based TPC that we expect to most closely align with the field-observed response

of infection occurrence to temperature.

Unimodal traits were fit with a flexTPC functional form (Equation 1) which can describe

unimodal TPCs of any skewness (Cruz-Loya, Mordecai and Savage, *preprint*). Using this

flexible function to fit TPCs of different shapes across our traits of interest allowed for ease of

comparison of thermal responses among traits and populations.

231
$$r(T) = \begin{cases} r_{max}\left[\left(\dfrac{T - T_{min}}{\alpha}\right)^{\alpha}\left(\dfrac{T_{max} - T}{1 - \alpha}\right)^{1-\alpha}\left(\dfrac{1}{T_{max} - T_{min}}\right)\right]^{\frac{\alpha(1-\alpha)}{\beta^2}} & T_{min} < T < T_{max} \\ 0 & \text{otherwise} \end{cases} \quad (1)$$

232   In this equation, T_{min} and T_{max} correspond to the minimum and maximum temperature,

233   respectively, r_{max} to the maximum trait value, and α and β are parameters that determine the

234   curve skewness and rate of decrease from the maximum trait value (Appendix S1). Host

235   melanization increased monotonically with temperature, and was fit to a 4-parameter shifted

236   logistic model (Equation 2, see Appendix S1 for derivation).

237
$$m(T) = \begin{cases} \dfrac{1}{\left[1 + (2^{1/s} - 1)\left(\dfrac{T_{50} - T_{min}}{T - T_{min}}\right)^a\right]^s} & T > T_{min} \\ 0 & \text{otherwise} \end{cases} \quad (2)$$

where T_{min} is the minimum temperature, T_{50} is the temperature at which 50% of the population

is melanized, and a and s are constants that allow for different rates of change of melanization

near T_{50} and for the response to be asymmetric, which was necessary to fit the data.

We used a binomial likelihood for all traits that were measured as proportions. For the free-living

parasite population growth rate TPC, we used a Gamma likelihood with population growth rates

obtained from the raw time-course data with the R package "growthrates" (Petzoldt 2022).

Optimal temperatures were calculated analytically using the posterior samples of the model

parameters (Appendix S1).

To assess population variation in lab-measured thermal performance among host

populations and the responses measured in the infection experiment, we compared 95% credible

intervals for the thermal optimum (T_{opt}), maximum (T_{max}), and minimum (T_{min}) and interpreted

non-overlapping intervals as evidence of biologically significant variation. We additionally

compared the magnitudes of the 95% credible intervals over TPC plots for each population.

Since the credible intervals for T_{opt}, T_{min}, and T_{max} overlapped for all populations for all

measured traits, we combined the data to generate species-representative TPCs, which we used

for our main analyses and interpretations. Finally, we compared the parasitism TPC to the host

and parasite trait TPCs measured in isolation.

We used R version 4.2.1 for all statistical analyses and for figure generation. See

Appendix S1 for additional details and R packages used.

Results

Does infection occurrence vary with climate across the species range?

Infection occurrence followed a hump-shaped pattern along the surveyed temperature

gradient, peaking at 9.4ºC mean rainy season temperature (Figure 2). Mean temperature^2 and

year were significant predictors of *L. clarki* infection presence (Table 1). Quadratic models fit

individually for infection occurrence in each year peaked at 10.0ºC in 2022 and 9.2ºC in 2023,

but the interactions between year and both temperature variables were not statistically significant

(Table 1). Geographically, infections were common in northern California and southern Oregon

and rarer at the warm and cold ends of the surveyed temperature gradient in southern California

and northern Washington, respectively (Figure 1).

Of the two sampling years, 2023 was cooler and wetter and had a higher percentage of

infected tree holes (29.7%, N = 431) compared to 2022 (12.5%, N = 320) (Figure 2) (NOAA

National Centers for Environmental Information 2022; 2023). Tree hole habitat temperatures

logged by iButtons (N = 32 successfully retrieved) did not systematically differ from PRISM air

temperature estimates (mean temperature difference = 1.3ºC, SD = 2.6ºC; R^2 mean = 0.94, SD =

0.03) (Appendix S2: Figure S1). Additional details are provided in Appendix S2.

Does the thermal response of the host – parasite interaction depend on the host population?

We found similar thermal optima, minima and maxima across the six host populations in the

infection experiment within each of the three measured interaction outcomes of infection,

melanization of parasites, and mortality (Figure 3; Appendix S2: Figures S2-S11; Tables S3-S7).

Although the mean estimated T_{opt}, T_{min}, and T_{max} varied slightly among populations for each

trait, all 95% credible intervals across populations for each outcome x parameter combination

overlapped. Therefore, we combined the data for all host populations to estimate broadly

representative TPCs for the host – parasite interaction, with reduced uncertainty compared to the

population-specific TPCs (Figure 4; Appendix S2: Figure S12; Table S9). The only trait in which

we observed clear intraspecific variation was in mortality around 10ºC, which was higher in one

intermediate climate population (Big Sur (BIG)) compared to the other (Woodside (JR)) and to

one cold climate population (Eugene (EUG)) (Figure S13).

How do different components of the host – parasite interaction relate to the overall thermal

sensitivity of parasitism?

Although free-living parasite population growth was maximized at warm temperatures, infection and parasite-induced mortality peaked at significantly colder temperatures, corresponding to conditions that suppressed host defense mechanisms, namely the melanization immune response and larval development speed (Figure 4). Specifically, the initial infection rate peaked at 10ºC (95% CI: 8.7 ºC, 11.3 ºC), and the total infection rate peaked at 8.4ºC (95% CI: 6.8 ºC, 9.4 ºC), much lower temperatures than the 19.5ºC (95% CI: 18.9 ºC, 20.2 ºC) optimum for free-living parasite population growth (Figure 4A; Appendix S2: Figure S12; Tables S8-9). The mechanically assayed host melanization immune response increased with temperature up to 20ºC, where it plateaued (Figure 4A; Appendix S2: Table S10). At warmer temperatures, attacking parasites faced not only a stronger host immune response, but also shorter time windows in which to infect hosts without being shed with the cuticle during larval molting and pupation or eaten by later larval instars. The time to the first observed pupation shortened as the temperature increased, occurring after 9 days at 28ºC compared to 57 days at 7ºC (Figure 4B), consistent with previously measured *Ae. sierrensis* larval development rate TPCs that peaked at 27.7ºC (Couper *et al.*, 2024). Consequently, quick pupation interrupted upward trajectories of infection and mortality at intermediate temperatures, contributing to the decline in parasitism above 11ºC, and to the lower T_{opt} of total compared to initial infection.

Discussion

In this study, we used complementary field observations and lab experiments to understand how temperature affects a host-parasite interaction across a large, climatically variable species

range. We found that infection peaked at sites with intermediate temperatures (9.4ºC) in the field

and in a colder, wetter year, aligning with the unimodal temperature-dependence of infection

measured in the lab, which peaked at 10ºC. This reflects the interacting thermal biology of

multiple mosquito and ciliate traits, which together led to an optimal temperature for parasitism

(8-10ºC for total versus initial infection) that was colder than the optimal temperature for free-

living parasite performance (19.5ºC). Infection rates of mosquito populations from different

climates had similar thermal responses in the lab, and a single TPC described the infection

process across the entire surveyed area. These findings suggest that TPCs measured under

simplified lab conditions can capture both patterns of parasitism across large geographic

temperature gradients in complex natural settings and the underlying thermal biology

mechanisms.

Lab experiments capture nonlinear parasitism thermal responses in the field

The results add to evidence that temperature nonlinearly affects parasitism not only in the

lab, but also in nature, making laboratory and field studies that are designed to capture nonlinear

temperature effects valuable tools for understanding disease risk (Dudney et al. 2021; Kirk et al.

2024; Shocket, Ryan, and Mordecai 2018). However, despite the similarities between the field

observations and the lab-measured parasitism TPC, we cannot rule out the influence of other

factors. For example, the colder survey year with higher infection prevalence was also wetter,

suggesting that precipitation patterns that determined when and for how long tree holes were

water-filled may have also affected the potential for parasitism to occur. These limitations should

be addressed with field experiments that manipulate temperatures in natural tree holes at

different latitudes or elevations; lab experiments that incorporate additional environmental

complexity; and long term repeated yearly sampling of the same tree holes.

Intraspecific variation

Very few studies investigate the impact of local thermal adaptation on species distributions

and host – parasite interactions, a gap often noted in the literature (Sternberg and Thomas 2014;

Urban et al. 2016; Couper et al. 2021; Aleuy et al. 2023). In addressing this gap, we did not find

evidence of local temperature adaptation affecting infection patterns across temperatures in the

lab or the field, consistent with our previous work on sympatric and allopatric host – parasite

interactions over a narrower temperature range, and despite our previous findings that both *Ae.*

sierrensis pupal development and *L. clarki* free-living population growth rates are locally

adapted to temperature (Lyberger et al. 2024; Couper et al. 2024; Ismail et al. 2023). However,

this is one of the only ectothermic host – parasite systems for which interaction thermal

sensitivity has been assessed for populations from different climates, along with a stickleback –

tapeworm system in which effects of temperature on parasitism were similar among four study

populations collected from across a similar geographic temperature gradient (Franke et al. 2019).

Notably, intraspecific variation in parasitism temperature sensitivity has been shown for levels of

biological organization both above and below the population level that we assessed, for

Batrachochytrium dendrobatitis infection both among different frog host species, and among

salamander host color morphs within a single population (Cohen et al. 2017; Venesky et al.

2022). However, if population thermal sensitivities of parasitism are commonly consistent across

species ranges, broadly applicable TPCs should be easily attainable for the many host – parasite

systems that are amenable to laboratory studies, and field studies may also be able to capture

nonlinear responses of parasitism to temperature.

Mechanisms of parasitism temperature dependence

Our finding that parasitism peaked at temperatures that constrained host development and

defense—rather than at the thermal optimum for parasite growth and attack—aligns with theory

on the thermal biology of biotic interactions. Thermal asymmetry theory posits that parasitism

peaks at temperatures where parasites outperform hosts, which may differ from optimal

temperatures for the parasites themselves (Dell, Pawar, and Savage 2014; Cohen et al. 2017).

Our study supports the value of approaching this problem as the outcome of multiple interacting

host and parasite traits with varying thermal responses (Paull, LaFonte, and Johnson 2012; Marta

S. Shocket et al. 2018; Sun et al. 2023).

Conclusions

Overall, our characterization of the temperature – infection relationship of the *Ae. sierrensis* host

– *L. clarki* parasite interaction across settings demonstrates how controlled lab experiments can

be paired with large-scale field observations to understand the complex thermal biology of

parasitism. Climate warming is expected to cause geographic contractions, expansions, and shifts

in host thermal refuges from disease (Altizer et al. 2013; Gsell et al. 2023). Our results suggest

that parasitism burdens on *Ae. sierrensis* may decline at southern latitudes and low elevations as

thermally suitable conditions for parasitism shift northwards and upwards into areas where cold

conditions currently limit *L. clarki* infections. Whether the distribution of parasitism ultimately

tracks climate warming will depend on multiple factors, including parasite dispersal and the

effects of other environmental changes in conjunction with increasing mean temperatures, such

as increasingly extreme precipitation patterns (Swain et al. 2018). Understanding how

temperature affects parasitism in inherently complex and variable natural systems remains a

pressing problem with broad implications for current and future disease risk management.

Simple pairings of field surveys and lab experiments can offer straightforward means for

approaching this challenge.

Acknowledgments

Otto Seppälä, Jan Washburn, Nona Chiariello, and Angie Nakano played critical roles in

establishing the study system. Our field sampling and mosquito rearing efforts were made

possible by assistance from staff at vector control agencies for the California counties of San

Mateo, Santa Clara, Marin-Sonoma, Solano, Alameda, Shasta, Placer, Santa Barbara, and

Riverside; Jasper Ridge Biological Preserve 'Ootchamin 'Ooyakma; the UC Reserves at

Emerson Oaks, Dawson Los Monos Canyon, and Hopland; and Andrew Rivera of Clarke Pest

Control. Mordecai lab members helped with setting up the infection experiment. JEF was funded

by the Bing-Mooney Fellowship and the ARCS Foundation. EAM and MCL were supported by

the National Institutes of Health (R35GM133439 and R01AI168097). EAM was supported by

the National Science Foundation (DEB-2011147 with Fogarty International Center), the National

(R01AI102918), the Stanford Woods Institute for the Environment, the Stanford Center for

Innovation in Global Health, and the Stanford Doerr School of Sustainability. KL was funded by

the NSF Postdoctoral Research Fellowships in Biology Program under Grant No. 2208947. LIC

was funded by the NSF Postdoctoral Research Fellowship in Biology. MCL was supported by a

Stanford PRISM Baker Fellowship.

References

Ahrens, Collin W., Richard A. Mazanec, Trudy Paap, Katinka X. Ruthrof, Anthea Challis, Giles
    Hardy, Margaret Byrne, David T. Tissue, and Paul D. Rymer. 2019. "Adaptive Variation
    for Growth and Resistance to a Novel Pathogen along Climatic Gradients in a Foundation
    Tree." *Evolutionary Applications* 12 (6): 1178–90. https://doi.org/10.1111/eva.12796.

Aleuy, O. Alejandro, Stephanie J. Peacock, Péter K. Molnár, Kathreen E. Ruckstuhl, and Susan J.
    Kutz. 2023. "Local Thermal Adaptation and Local Temperature Regimes Drive the
    Performance of a Parasitic Helminth under Climate Change: The Case of *Marshallagia
    marshalli* from Wild Ungulates." *Global Change Biology* 29 (22): 6217–33.
    https://doi.org/10.1111/gcb.16918.

Altizer, Sonia, Richard S. Ostfeld, Pieter T. J. Johnson, Susan Kutz, and C. Drew Harvell. 2013.
    "Climate Change and Infectious Diseases: From Evidence to a Predictive Framework."
    Science 341 (6145): 514–19. https://doi.org/10.1126/science.1239401.

Bates, Douglas, Martin Maechler, Ben Bolker, and Steven Walker. 2003. "Lme4: Linear Mixed-
    Effects Models Using 'Eigen' and S4." https://doi.org/10.32614/CRAN.package.lme4.

Bombardi, Rodrigo J., James L. Kinter, and Oliver W. Frauenfeld. 2019. "A Global Gridded
    Dataset of the Characteristics of the Rainy And Dry Seasons." *Bulletin of the American
    Meteorological Society* 100 (7): 1315–28. https://doi.org/10.1175/BAMS-D-18-0177.1.

Boots, M. 2000. "Density-Independent Resource Limitation and the Transmission of an Insect
    Pathogen." *Oecologia* 124 (2): 172–75. https://doi.org/10.1007/s004420050004.

Broberg, Leonard E., and William E. Bradshaw. 1997. "*Lambornella clarki* (Ciliophora:
    Tetrahymenidae) Resistance in *Aedes sierrensis* (Diptera: Culicidae): Population

Differentiation in a Quantitative Trait and Allozyme Loci." *Journal of Medical*

Entomology 34 (1): 38–45. https://doi.org/10.1093/jmedent/34.1.38.

Brown, Joel J., Mercedes Pascual, Michael C. Wimberly, Leah R. Johnson, and Courtney C.

Murdock. 2023. "Humidity – The Overlooked Variable in the Thermal Biology of

Mosquito-Borne Disease." *Ecology Letters* 26 (7): 1029–49.

https://doi.org/10.1111/ele.14228.

Christensen, Bruce M., Jianyong Li, Cheng-Chen Chen, and Anthony J. Nappi. 2005.

"Melanization Immune Responses in Mosquito Vectors." *Trends in Parasitology* 21 (4):

192–99. https://doi.org/10.1016/j.pt.2005.02.007.

Clark, T.B., and D.G. Brandl. 1976. "Observations on the Infection of *Aedes sierrensis* by a

Tetrahymenine Ciliate." *Journal of Invertebrate Pathology* 28 (3): 341–49.

https://doi.org/10.1016/0022-2011(76)90009-4.

Cohen, Jeremy M., Matthew D. Venesky, Erin L. Sauer, David J. Civitello, Taegan A. McMahon,

Elizabeth A. Roznik, and Jason R. Rohr. 2017. "The Thermal Mismatch Hypothesis

Explains Host Susceptibility to an Emerging Infectious Disease." Edited by Richard

Ostfeld. *Ecology Letters* 20 (2): 184–93. https://doi.org/10.1111/ele.12720.

Couper, Lisa I., Johannah E. Farner, Jamie M. Caldwell, Marissa L. Childs, Mallory J. Harris,

Devin G. Kirk, Nicole Nova, et al. 2021. "How Will Mosquitoes Adapt to Climate

Warming?" Edited by George H Perry. *eLife* 10 (August):e69630.

https://doi.org/10.7554/eLife.69630.

Couper, Lisa I., Johannah E. Farner, Kelsey P. Lyberger, Alexandra S. Lee, and Erin A.

Mordecai. 2024. "Mosquito Thermal Tolerance Is Remarkably Constrained across a

Large Climatic Range." *Proceedings of the Royal Society B: Biological Sciences* 291

(2015): 20232457. https://doi.org/10.1098/rspb.2023.2457.

Cruz-Loya, Mauricio, Erin A. Mordecai, and Van M. Savage. 2024. "A Flexible Model for

Thermal Performance Curves." *bioRxiv*, January, 2024.08.01.605695.

https://doi.org/10.1101/2024.08.01.605695.

Darsie, Richard F., and Ronald A. Ward. 2005. *Identification and Geographical Distribution of*

the Mosquitoes of North America, North of Mexico. University Press of Florida.

Dell, Anthony I., Samraat Pawar, and Van M. Savage. 2014. "Temperature Dependence of

Trophic Interactions Are Driven by Asymmetry of Species Responses and Foraging

Strategy." *Journal of Animal Ecology* 83 (1): 70–84. https://doi.org/10.1111/1365-

2656.12081.

Dudney, Joan, Claire E. Willing, Adrian J. Das, Andrew M. Latimer, Jonathan C. B. Nesmith,

and John J. Battles. 2021. "Nonlinear Shifts in Infectious Rust Disease Due to Climate

Change." *Nature Communications* 12 (1): 5102. https://doi.org/10.1038/s41467-021-

25182-6.

Franke, Frederik, Nadja Raifarth, Joachim Kurtz, and Jörn P. Scharsack. 2019. "Consequences of

Divergent Temperature Optima in a Host–Parasite System." *Oikos* 128 (6): 869–80.

https://doi.org/10.1111/oik.05864.

Gaitán-Espitia, Juan Diego, Leonardo D. Bacigalupe, Tania Opitz, Nelson A. Lagos, Tania

Timmermann, and Marco A. Lardies. 2014. "Geographic Variation in Thermal

Physiological Performance of the Intertidal Crab *Petrolisthes violaceus* along a

Latitudinal Gradient." *Journal of Experimental Biology* 217 (24): 4379–86.

https://doi.org/10.1242/jeb.108217.

Gehman, Alyssa-Lois M., Richard J. Hall, and James E. Byers. 2018. "Host and Parasite Thermal
Ecology Jointly Determine the Effect of Climate Warming on Epidemic Dynamics."
Proceedings of the National Academy of Sciences 115 (4): 744–49.
https://doi.org/10.1073/pnas.1705067115.

Gsell, Alena S., Arjen Biere, Wietse de Boer, Irene de Bruijn, Götz Eichhorn, Thijs Frenken,
Stefan Geisen, et al. 2023. "Environmental Refuges from Disease in Host–Parasite
Interactions under Global Change." *Ecology* 104 (4): e4001.
https://doi.org/10.1002/ecy.4001.

Hall, Spencer R., Joseph L. Simonis, Roger M. Nisbet, Alan J. Tessier, and Carla E. Cáceres.
2009. "Resource Ecology of Virulence in a Planktonic Host-Parasite System: An
Explanation Using Dynamic Energy Budgets." *The American Naturalist* 174 (2): 149–62.
https://doi.org/10.1086/600086.

Hartig, Florian. 2017. "Package 'Dharma.'" *R Package*.

Ismail, Sherine, Johannah Farner, Lisa Couper, Erin Mordecai, and Kelsey Lyberger. 2023.
"Temperature and Intraspecific Variation Affect Host–Parasite Interactions." *Oecologia*,
November. https://doi.org/10.1007/s00442-023-05481-z.

Jordan, R. G., and W. E. Bradshaw. 1978. "Geographic Variation in the Photoperiodic Response
of the Western Tree-Hole Mosquito, *Aedes sierrensis*." *Annals of the Entomological
Society of America* 71 (4): 487–90. https://doi.org/10.1093/aesa/71.4.487.

Jordan, Robert G. 1980a. "Embryonic Diapause in Three Populations of the Western Tree Hole
Mosquito, *Aedes sierrensis*." *Annals of the Entomological Society of America* 73 (4):
357–59. https://doi.org/10.1093/aesa/73.4.357.

———. 1980b. "Geographic Differentiation in the Development *of Aedes sierrensis* (Diptera:

Culicidae) in Nature." *The Canadian Entomologist* 112 (2): 205–10.

https://doi.org/10.4039/Ent112205-2.

Jurriaans, S., and M. O. Hoogenboom. 2019. "Thermal Performance of Scleractinian Corals

along a Latitudinal Gradient on the Great Barrier Reef." *Philosophical Transactions of*

the Royal Society B: Biological Sciences 374 (1778): 20180546.

https://doi.org/10.1098/rstb.2018.0546.

Kirk, Devin, Natalie Jones, Stephanie Peacock, Jessica Phillips, Péter K. Molnár, Martin

Krkošek, and Pepijn Luijckx. 2018. "Empirical Evidence That Metabolic Theory

Describes the Temperature Dependency of Within-Host Parasite Dynamics." *PLOS*

Biology 16 (2): e2004608. https://doi.org/10.1371/journal.pbio.2004608.

Kirk, Devin, Samantha Straus, Marissa L. Childs, Mallory Harris, Lisa Couper, T. Jonathan

Davies, Coreen Forbes, et al. 2024. "Temperature Impacts on Dengue Incidence Are

Nonlinear and Mediated by Climatic and Socioeconomic Factors: A Meta-Analysis."

PLOS Climate 3 (3): e0000152. https://doi.org/10.1371/journal.pclm.0000152.

Kutz, S. J., E. P. Hoberg, L. Polley, and E. J. Jenkins. 2005. "Global Warming Is Changing the

Dynamics of Arctic Host-Parasite Systems." *Proceedings: Biological Sciences* 272

(1581): 2571–76.

Labaude, Sophie, Yannick Moret, Frank Cézilly, Charel Reuland, and Thierry Rigaud. 2017.

"Variation in the Immune State of *Gammarus pulex* (Crustacea, Amphipoda) According

to Temperature: Are Extreme Temperatures a Stress?" *Developmental & Comparative*

Immunology 76 (November):25–33. https://doi.org/10.1016/j.dci.2017.05.013.

Laughton, Alice M., Cian O. O'Connor, and Robert J. Knell. 2017. "Responses to a Warming

World: Integrating Life History, Immune Investment, and Pathogen Resistance in a

Model Insect Species." *Ecology and Evolution* 7 (22): 9699–9710.

https://doi.org/10.1002/ece3.3506.

Luković, Jelena, John C. H. Chiang, Dragan Blagojević, and Aleksandar Sekulić. 2021. "A Later

Onset of the Rainy Season in California." *Geophysical Research Letters* 48 (4):

e2020GL090350. https://doi.org/10.1029/2020GL090350.

Lyberger, Kelsey, Johannah Farner, Lisa Couper, and Erin A. Mordecai. 2024. "A Mosquito

Parasite Is Locally Adapted to Its Host but Not Temperature." *The American Naturalist*

204 (2): 121–32. https://doi.org/10.1086/730522.

Martin, Lindsay E., and Julián F. Hillyer. 2024. "Higher Temperature Accelerates the Aging-

Dependent Weakening of the Melanization Immune Response in Mosquitoes." *PLOS*

Pathogens 20 (1): e1011935. https://doi.org/10.1371/journal.ppat.1011935.

Mazé-Guilmo, Elise, Simon Blanchet, Olivier Rey, Nicolas Canto, and Géraldine Loot. 2016.

"Local Adaptation Drives Thermal Tolerance among Parasite Populations: A Common

Garden Experiment." *Proceedings of the Royal Society B: Biological Sciences* 283

(1830): 20160587. https://doi.org/10.1098/rspb.2016.0587.

Mordecai, Erin A., Jamie M. Caldwell, Marissa K. Grossman, Catherine A. Lippi, Leah R.

Johnson, Marco Neira, Jason R. Rohr, et al. 2019. "Thermal Biology of Mosquito-borne

Disease." Edited by James (Jeb) Byers. *Ecology Letters*, July, ele.13335.

https://doi.org/10.1111/ele.13335.

Mordecai, Erin A., Jeremy M. Cohen, Michelle V. Evans, Prithvi Gudapati, Leah R. Johnson,

Catherine A. Lippi, Kerri Miazgowicz, et al. 2017. "Detecting the Impact of Temperature

on Transmission of Zika, Dengue, and Chikungunya Using Mechanistic Models." *PLOS Neglected Tropical Diseases* 11 (4): e0005568. https://doi.org/10.1371/journal.pntd.0005568.

Mordecai, Erin A., Krijn P. Paaijmans, Leah R. Johnson, Christian Balzer, Tal Ben-Horin, Emily de Moor, Amy McNally, et al. 2013. "Optimal Temperature for Malaria Transmission Is Dramatically Lower than Previously Predicted." *Ecology Letters* 16 (1): 22–30. https://doi.org/10.1111/ele.12015.

Murdock, C. C., Krijn P. Paaijmans, Andrew S. Bell, Jonas G. King, Julián F. Hillyer, Andrew F. Read, and Matthew B. Thomas. 2012. "Complex Effects of Temperature on Mosquito Immune Function." *Proceedings of the Royal Society B: Biological Sciences* 279 (1741): 3357–66. https://doi.org/10.1098/rspb.2012.0638.

NOAA National Centers for Environmental Information. 2022. "Monthly National Climate Report for 2022." https://www.ncei.noaa.gov/access/monitoring/monthly-report/national/202408.

———. 2023. "Monthly National Climate Report for 2023." https://www.ncei.noaa.gov/access/monitoring/monthly-report/national/202408.

Padfield, Daniel, Meaghan Castledine, and Angus Buckling. 2020. "Temperature-Dependent Changes to Host–Parasite Interactions Alter the Thermal Performance of a Bacterial Host." *The ISME Journal* 14 (2): 389–98. https://doi.org/10.1038/s41396-019-0526-5.

Palmer-Young, Evan C., Lyna Ngor, Rodrigo Burciaga Nevarez, Jason A. Rothman, Thomas R. Raffel, and Quinn S. McFrederick. 2019. "Temperature Dependence of Parasitic Infection and Gut Bacterial Communities in Bumble Bees." *Environmental Microbiology* 21 (12): 4706–23. https://doi.org/10.1111/1462-2920.14805.

Paull, Sara H., Bryan E. LaFonte, and Pieter T. J. Johnson. 2012. "Temperature-Driven Shifts in

a Host-Parasite Interaction Drive Nonlinear Changes in Disease Risk." *Global Change*

Biology 18 (12): 3558–67. https://doi.org/10.1111/gcb.12018.

Penczykowski, Rachel M., Spencer R. Hall, David J. Civitello, and Meghan A. Duffy. 2014.

"Habitat Structure and Ecological Drivers of Disease." *Limnology and Oceanography* 59

(2): 340–48. https://doi.org/10.4319/lo.2014.59.2.0340.

Petzoldt, Thomas. 2022. "R Package Growthrates."

Praskievicz, Sarah, and Heejun Chang. 2009. "Winter Precipitation Intensity and ENSO/PDO

Variability in the Willamette Valley of Oregon." *International Journal of Climatology* 29

(13): 2033–39. https://doi.org/10.1002/joc.1838.

PRISM Climate Group, and Oregon State University. 2021. "PRISM Gridded Climate Data."

https://prism.oregonstate.edu.

Ragonese, Isabella G., Maya R. Sarkar, Richard J. Hall, and Sonia Altizer. 2024. "Extreme Heat

Reduces Host and Parasite Performance in a Butterfly–Parasite Interaction." *Proceedings*

of the Royal Society B: Biological Sciences 291 (2015): 20232305.

https://doi.org/10.1098/rspb.2023.2305.

Ryan, Sadie J., Colin J. Carlson, Erin A. Mordecai, and Leah R. Johnson. 2019. "Global

Expansion and Redistribution of *Aedes*-Borne Virus Transmission Risk with Climate

Change." *PLOS Neglected Tropical Diseases* 13 (3): e0007213.

https://doi.org/10.1371/journal.pntd.0007213.

Seppälä, Otto, Katri Liljeroos, Anssi Karvonen, and Jukka Jokela. 2008. "Host Condition as a

Constraint for Parasite Reproduction." *Oikos* 117 (5): 749–53.

https://doi.org/10.1111/j.0030-1299.2008.16396.x.

Shah, Melisa M., Amy R. Krystosik, Bryson A. Ndenga, Francis M. Mutuku, Jamie M. Caldwell,

Victoria Otuka, Philip K. Chebii, et al. 2019. "Malaria Smear Positivity among Kenyan

Children Peaks at Intermediate Temperatures as Predicted by Ecological Models."

Parasites & Vectors 12 (1): 288. https://doi.org/10.1186/s13071-019-3547-z.

Shocket, Marta S., Alexander T. Strauss, Jessica L. Hite, Maja Šljivar, David J. Civitello,

Meghan A. Duffy, Carla E. Cáceres, and Spencer R. Hall. 2018. "Temperature Drives

Epidemics in a Zooplankton-Fungus Disease System: A Trait-Driven Approach Points to

Transmission via Host Foraging." *The American Naturalist* 191 (4): 435–51.

https://doi.org/10.1086/696096.

Shocket, Marta S., Anna B. Verwillow, Mailo G. Numazu, Hani Slamani, Jeremy M. Cohen,

Fadoua El Moustaid, Jason Rohr, Leah R. Johnson, and Erin A. Mordecai. 2020.

"Transmission of West Nile and Five Other Temperate Mosquito-Borne Viruses Peaks at

Temperatures between 23°C and 26°C." Edited by Eduardo Franco, Talía Malagón, and

Alyssa Gehman. *eLife* 9 (September):e58511. https://doi.org/10.7554/eLife.58511.

Shocket, Marta S., Sadie J. Ryan, and Erin A. Mordecai. 2018. "Temperature Explains Broad

Patterns of Ross River Virus Transmission." *eLife* 7 (August):e37762.

https://doi.org/10.7554/eLife.37762.

Sinclair, Brent J., Caroline M. Williams, and John S. Terblanche. 2012. "Variation in Thermal

Performance among Insect Populations." *Physiological and Biochemical Zoology* 85 (6):

594–606. https://doi.org/10.1086/665388.

Sternberg, Eleanore D., and Matthew B. Thomas. 2014. "Local Adaptation to Temperature and

the Implications for Vector-Borne Diseases." *Trends in Parasitology* 30 (3): 115–22.

https://doi.org/10.1016/j.pt.2013.12.010.

Sun, Syuan-Jyun, Marcin K. Dziuba, Riley N. Jaye, and Meghan A. Duffy. 2023. "Temperature

Modifies Trait-Mediated Infection Outcomes in a *Daphnia*–Fungal Parasite System."

Philosophical Transactions of the Royal Society B: Biological Sciences 378 (1873):

20220009. https://doi.org/10.1098/rstb.2022.0009.

Suwanchaichinda, Chansak, and Susan M. Paskewitz. 1998. "Effects of Larval Nutrition, Adult

Body Size, and Adult Temperature on the Ability of *Anopheles gambiae* (Diptera:

Culicidae) to Melanize Sephadex Beads." *Journal of Medical Entomology* 35 (2): 157–

61. https://doi.org/10.1093/jmedent/35.2.157.

Swain, Daniel L., Baird Langenbrunner, J. David Neelin, and Alex Hall. 2018. "Increasing

Precipitation Volatility in Twenty-First-Century California." *Nature Climate Change* 8

(5): 427–33. https://doi.org/10.1038/s41558-018-0140-y.

Taylor, Rachel A., Sadie J. Ryan, Catherine A. Lippi, David G. Hall, Hossein A. Narouei-

Khandan, Jason R. Rohr, and Leah R. Johnson. 2019. "Predicting the Fundamental

Thermal Niche of Crop Pests and Diseases in a Changing World: A Case Study on Citrus

Greening." Edited by Luke Flory. *Journal of Applied Ecology*, July, 1365-2664.13455.

https://doi.org/10.1111/1365-2664.13455.

Tesla, Blanka, Leah R. Demakovsky, Erin A. Mordecai, Sadie J. Ryan, Matthew H. Bonds,

Calistus N. Ngonghala, Melinda A. Brindley, and Courtney C. Murdock. 2018.

"Temperature Drives Zika Virus Transmission: Evidence from Empirical and

Mathematical Models." *Proceedings of the Royal Society B: Biological Sciences* 285

(1884): 20180795. https://doi.org/10.1098/rspb.2018.0795.

Thomas, Matthew B., and Simon Blanford. 2003. "Thermal Biology in Insect-Parasite

Interactions." *Trends in Ecology & Evolution* 18 (7): 344–50.

https://doi.org/10.1016/S0169-5347(03)00069-7.

Urban, M. C., G. Bocedi, A. P. Hendry, J.-B. Mihoub, G. Pe'er, A. Singer, J. R. Bridle, et al.

2016. "Improving the Forecast for Biodiversity under Climate Change." *Science* 353

(6304): aad8466. https://doi.org/10.1126/science.aad8466.

Venesky, Matthew D., Joseph DeMarchi, Cari Hickerson, and Carl D. Anthony. 2022. "Does the

Thermal Mismatch Hypothesis Predict Disease Outcomes in Different Morphs of a

Terrestrial Salamander?" *Journal of Experimental Zoology Part A: Ecological and*

Integrative Physiology, 1–10. https://doi.org/10.1002/jez.2581.

Washburn, J., M. Gross, D. Mercer, and Anderson. 1988. "Predator-Induced Trophic Shift of a

Free-Living Ciliate: Parasitism of Mosquito Larvae by Their Prey." *Science* 240 (4856):

1193–95. https://doi.org/10.1126/science.3131877.

Washburn, Jan O., and John R. Anderson. 1986. "Distribution of *Lambornella clarki* (Ciliophora:

Tetrahymenidae) and Other Mosquito Parasites in California Treeholes." *Journal of*

Invertebrate Pathology 48 (3): 296–309. https://doi.org/10.1016/0022-2011(86)90058-3.

Washburn, Jan O., John R. Anderson, and David R. Mercer. 1991. "Parasitism of Newly-Hatched

Aedes sierrensis (Diptera: Culicidae) Larvae by *Lambornella clarki* (Ciliophora:

Tetrahymenidae) Following Habitat Flooding." *Journal of Invertebrate Pathology* 58 (1):

67–74. https://doi.org/10.1016/0022-2011(91)90163-K.

Tables

Predictor	Estimate	Standard error	Z-value	p-value
Mean annual temperature	-0.22	0.22	-1.01	3.13×10^{-1}
Mean annual temperature^2	-0.43	0.21	-2.01	**4.45×10^{-2}**
Survey year	1.31	0.34	3.90	**9.5×10^{-5}**
*Mean annual temperature * survey year*	-0.45	0.26	-1.72	8.5×10^{-2}
*Mean annual temperature^2 * survey year*	-0.127	0.247	-0.52	6.34×10^{-1}

649   **Table 1. Results of a generalized linear mixed-effects model with tree hole infection status**

650   **as the dependent variable and mean annual temperature, mean annual temperature^2, and**

651   **survey year as fixed effects, and tree hole ID as a random effect.** P-values < 0.05 are

652   displayed in bold to highlight statistical significance.

     Figure captions

Figure 1. Tree holes were surveyed, and experimental host and parasite populations reared,

from across a large geographic temperature gradient on the western coast of the United

States during the 2022 (left) and 2023 (right) rainy seasons. The mapped color gradient shows

mean temperature during the rainy season in Washington, Oregon, and California, USA, from the

4 km PRISM climate dataset. Points show the locations of 507 total surveyed tree holes. *L. clarki*

infection presence is indicated by black points, and absence by white points. Colored ovals

indicate the source locations of the six host populations used in the infection experiment (from

north to south: EUG = Eugene, OR; JR = Woodside, CA; FRS = Shaver Lake, CA; BIG = Big

Sur, CA; WHIT = Trabuco Canyon, CA; CAS = San Juan Capistrano, CA); the gray oval shows

the host population used in the melanization response experiment (PAR = Paso Robles, CA); and

the gray rectangle shows the source location of the ciliate parasite culture used in the infection

experiment (Marin = Novato, CA). Green outlines indicate tree holes in which iButtons logged

temperature data during the year between the two sampling visits.

Figure 2. Infection probability followed a consistent hump-shaped relationship with mean

temperature during the rainy season, and was higher overall in 2023 compared to 2022. (A)

Counts of infected (black) and uninfected (gray) tree holes across the climate gradient in each

survey year. **(B)** Generalized linear mixed-effects model predictions of infection probability by

temperature and year are plotted across the surveyed climate gradient for each year of the field

survey. The solid and dashed lines show the mean model predictions for 2022 and 2023,

respectively, and the shading shows the 95% confidence intervals.

Figure 3. Thermal responses of host – parasite interaction outcomes were similar among

host populations from warm, intermediate, and cold source climates. For each outcome

measured in the infection experiment, mean estimated TPCs for each of six host populations are

shown (lines) with 95% credible intervals (shading). The left column shows initial infection and

parasite melanization responses up to the third experiment checkpoint, when the first deaths of

infected larvae released additional parasites into the environment. The right column shows the

total infection, parasite melanization, and parasite-induced mortality rates across temperatures at

the end of the experiment. Mortality rates were significantly higher near 10 ºC for population

BIG compared to populations JR and EUG. The TPCs shown here are presented individually,

with the experiment data, in Appendix S2 (Figures S2, S4, S6, S8, S10).

Figure 4. Experimental infections were most likely to occur at cold incubation

temperatures where the host melanization response and development were suppressed. (A)

Points show mean values, error bars show +/- 1 SE, lines show mean estimated TPCs, and

shading shows 95% credible intervals for key host – parasitism interaction outcomes and

processes: free-living parasite population growth (green); the host mechanically stimulated

melanization immune response (black); the initial infection rate (light pink), calculated up to the

first fatality at each temperature; the total infection rate (dark pink); and total mortality (gray)

calculated up to the first pupation at each temperature. The infection data shown include all host

populations in the infection experiment. **(B)** Infection (top; dark pink), parasite-induced mortality

(middle; light gray), and parasite melanization (bottom; dark gray) counts are plotted over time

for each temperature used in the infection experiment. Colored points show data from individual

microcosms, and black points and lines show mean trajectories +/- 1 SE. For each temperature,

the final experiment checkpoint was at the time of the first observed pupation.

Figure 1.

Figure 2.

Figure 3.

Figure 4.