Ecology

Article

Nonlinear effects of temperature on mosquito parasite infection across a large geographic climate

gradient

Johannah E. Farner[1]*, Kelsey P. Lyberger[2], Lisa I. Couper[3], Mauricio Cruz-Loya[1], Erin A.

Mordecai[1]

[1]Department of Biology, Stanford University, Stanford, CA 94305, USA; [2] College of Integrative

Sciences and Arts, Arizona State University, Mesa, AZ 85212, USA; [3] Division of Environmental

Health Sciences, University of California, Berkeley, Berkeley, CA 94720, USA.

*Corresponding author: jfarner@stanford.edu

Upon acceptance, data and code will be provided via Dryad.

17
18 Key words/phrases: *Aedes sierrensis;* disease ecology, *Lambornella clarki*; parasitism; thermal

19 biology thermal performance curve

Abstract

Temperature drives ectothermic host – parasite interactions, making them particularly sensitive to climatic variation and change. To isolate the role of temperature, lab-based studies are increasingly used to assess and forecast disease risk under current and future climate conditions. However, in the field, the effects of temperature on parasitism may be mediated by other sources of variation, including local adaptation of hosts and parasites to one another and to the environment. To address the key knowledge gaps of how temperature influences host – parasite interactions and whether thermal responses measured in controlled experiments capture infection across temperature gradients in nature, we paired an extensive field survey of parasitism—by the ciliate *Lambornella clarki* on its tree hole mosquito host, *Aedes sierrensis*— with laboratory experiments describing parasitism thermal performance curves (TPCs) for six host populations from varying climates. We also investigated the mechanisms underlying the thermal biology of the host – parasite interaction by separately measuring TPCs for infection, host immunity, and parasite growth rates. Along the west coast of North America, across an 1100 km climate gradient spanning 12ºC mean rainy season temperature variation, we found that parasitism peaked at intermediately cold temperatures of 9.2 – 10ºC, and was consistent both between field seasons and with the lab experiment results. In the laboratory experiments, infection thermal responses were consistently nonlinear and peaked at 8.4 – 10ºC, showing no evidence of host intraspecific variation in temperature sensitivity to parasitism. Importantly, parasitism peaked at temperatures below the thermal optimum for free-living *L. clarki* due to the balance of temperature effects on parasite growth and reproduction against the strength of the host melanization immune response. The results suggest that nonlinear responses to temperature

43   drive parasitism in nature, and that simple lab and field studies can accurately capture the

44   thermal biology of multilayered host – parasite interactions.

45   **Introduction**
46
47   Climate warming is fundamentally important to infectious disease risk because

temperature drives many of the constituent biological processes of host – parasite interactions,

including host and parasite growth and development, contact rates, and immune functions (Kutz

et al., 2005; Labaude et al., 2017; Laughton et al., 2017; Franke et al., 2019; Mordecai et al.,

2019). Strong relationships between temperature and parasitism are frequently observed in

laboratory experiments and captured in thermal performance curves (TPCs) that quantify how

infection rates change with temperature, typically from data collected across a range of constant

incubation temperatures (Angilletta 2009; Gehman et al., 2018; Kirk et al. 2018; Padfield et al.,

2020). Following expectations from biochemical first principles, TPCs are commonly unimodal,

with biological rates increasing with temperature from their critical thermal minimum, peaking at

a thermal optimum, and then declining up to a critical thermal minimum (Angilletta, 2009).

Researchers increasingly use TPCs to assess nonlinear shifts in disease risk based on current and

forecasted climate and weather conditions (Gehman et al., 2018; Ryan et al., 2019; Aleuy et al.,

2023). However, it is unclear whether TPCs measured in controlled laboratory settings fully

capture the thermal response of disease in natural settings, where environmental complexity may

modify or outweigh parasitism - temperature relationships (Mordecai et al., 2019; Taylor et al.,

2019).

64   Parasitism TPCs are typically measured for single laboratory populations, in the absence

of other stressors, and are infrequently compared to disease burdens along temperature gradients

in the field (e.g., Gehman et al., 2018; Mordecai et al., 2019; Padfield et al., 2020). Although an

increasing number of studies describe thermal responses for multiple populations or genotypes

within host-parasite systems (Mazé-Guilmo et al., 2016; Ahrens et al., 2019; Franke et al., 2019;

Aleuy et al., 2023; Ragonese et al., 2024), research on statistical or correlative relationships

between TPCs for disease transmissibility (e.g., R_0) and natural infection patterns has thus far

relied on laboratory measurements from single host and parasite populations or clones, and field

data collated from disparate sources and/or collected over small geographic areas. To date, this

research has focused primarily on mosquito-borne human disease and fungal pathogens of

zooplankton (Mordecai et al., 2013; 2017; Shocket, Ryan et al., 2018; Shocket, Strauss et al.,

2018; Tesla et al., 2018; Shah et al., 2019; Shocket et al., 2020; Kirk et al., 2024). However,

TPCs for both hosts and parasites can vary geographically within species due to local adaptation

and population structure (Sinclair et al., 2012; Gaitán-Espitia et al., 2014; Mazé-Guilmo et al.,

2016; Jurriaans & Hoogenboom, 2019; Aleuy et al., 2023; Couper et al., 2024), and additional

environmental variability in field settings may outweigh or modify the temperature dependence

of parasitism (e.g., Boots, 2000; Seppälä et al., 2008; Hall et al., 2009; Penczykowski et al.,

2014; Palmer-Young et al., 2019; Brown et al., 2023). Thus, there is a critical need to address the

questions of how temperature affects parasitism in the field, whether lab-based TPC estimates

can accurately, and mechanistically, capture these patterns, and whether local thermal adaptation

modifies the temperature-dependence of parasitism.

Central to capturing the thermal biology of parasitism is understanding the host and

parasite mechanisms that drive the interaction. For example, host life history and defense traits

and parasite growth, attack, dispersal, and transmission traits may each respond nonlinearly and

distinctly to temperature. The combined effect of these traits determines the overall impact of

temperature on parasitism (Paull et al., 2012; Shocket et al., 2018; Mordecai et al., 2019; Sun et

al., 2023). Moreover, host and parasite trait thermal responses may differ across populations due

to local adaptation and across environmental conditions (Thomas & Blanford, 2003; Sternberg &

Thomas, 2014). Given that it is not feasible to measure TPCs for every possible combination of

host population, parasite population, and environmental context, it is important to establish

whether TPCs measured for a single set of populations and conditions can generalize to cover

entire species ranges, and which traits capture the essential thermal biology of the interaction.

Here, we address these questions in a model system of a mosquito (*Aedes sierrensis*) and its

facultative ciliate parasite (*Lambornella clarki*), which occur across a large geographic gradient

in western North America.

In this study, we pair field surveys along the entire west coast of the continental United

States—spanning the geographic range of the host species—with laboratory experiments

describing the temperature dependence of infection and key traits involved in host defenses and

parasite fitness, for host populations from different climates across the survey area. Both species

are short-lived ectotherms that are directly affected by environmental temperatures, occur

commonly across a large geographic climate gradient in discrete tree hole habitats that are

naturally replicated across landscapes, and can be reared separately and together in the lab

(Washburn & Anderson, 1986; Broberg & Bradshaw, 1997; Darsie & Ward, 2005; Ismail et al.,

2023; Lyberger et al., 2024). A 1986 field survey found that infection occurrence increased from

southern to northern California (Washburn & Anderson, 1986). Our recent work describing

temperature effects on different populations of *Ae. sierrensis, L. clarki*, and their interactions

indicates that infection has a consistent nonlinear relationship with temperature for both

sympatric and allopatric pairs of host and parasite populations collected across climates, despite

evidence of local thermal adaptation in free-living *L. clarki* populations (Ismail et al., 2023;

Lyberger et al., 2024). This body of work generates the hypothesis that parasitism responds to

temperature as a function of the thermal biology of multiple host and parasite traits. However, it

is unknown whether the lab-measured temperature impacts on parasitism across different *Ae.*

sierrensis host populations are borne out in infection occurrence across the environmentally

complex species range.

Methods

Study system

Ae. sierrensis and its facultative ciliate parasite *L. clarki* occur across western North

America in seasonally rainwater-filled tree holes, where mosquitoes undergo one reproductive

cycle annually in which larvae feed on detritus and microbes (including free-living *L. clarki*)

before pupating and emerging as adults, and free-living *L. clarki* ciliates feed on aquatic bacteria

(Washburn & Anderson, 1986; Darsie & Ward, 2005). In the presence of *Ae. sierrensis* larvae,

free-living *L. clarki* irreversibly transform into parasites that attack larval mosquitoes by

attaching to the larval cuticle, penetrating into the body cavity, and then replicating, often to high

densities that kill hosts after a few weeks (Washburn et al., 1988). Mosquitoes can resist

infection through multiple defenses that each likely respond nonlinearly to temperature:

melanization (a generalized immune response), molting between larval instars, developing into

pupae, and consuming free-living *L. clarki* in the water column (Clark & Brandl, 1976;

Washburn et al., 1988; Washburn et al., 1991; Suwanchaichinda & Paskewitz, 1998; Christensen

et al., 2005; Murdock et al., 2012; Martin & Hillyer, 2024). Beyond the previously documented

latitudinal gradient, *L. clarki* presence is not correlated with habitat characteristics such as tree

hole size, exposure, height above ground, or tree species (Washburn & Anderson, 1986).

Field survey

    To document infection patterns over a large geographic temperature gradient and collect

materials for lab experiments, we sampled from over 500 water-filled tree hole habitats

containing *Ae. sierrensis* larvae spanning 1100 km north to south with a mean November – April

rainy season temperature gradient of 2.6ºC to 15.3ºC in California, Oregon, and Washington,

USA, during the rainy seasons of 2022 and 2023 (Figure 1) (PRISM Climate Group, 2021). Each

year, we started in the warmest and ended in the coolest locations to track the seasonal timing of

Ae. sierrensis egg hatching and development (Jordan & Bradshaw, 1978; Jordan, 1980a; 1980b),

collecting larval samples and examining them for infection at 10-40x magnification (Appendix

S1). We aimed to sample each tree hole in both years, but had to replace some locations in 2023

due to storm damage (Figure 1).

    For each sampling site, we used PRISM climate data to characterize long-term mean

temperature conditions in the November – April rainy season when mosquito larvae and parasites

are active in tree holes (PRISM Climate Group, 2021; Washburn & Anderson, 1986; Luković et

al., 2021; Praskievicz & Chang, 2009). We chose the PRISM dataset for its long-term coverage

of the entire study region. To assess the match between the remotely sensed PRISM data and the

conditions in surveyed tree hole habitats, we placed iButton temperature loggers (Maxim

Integrated, San Jose, CA, USA) in 84 arbitrarily selected tree holes across the survey area during

the 2022 field season. These iButtons were retrieved the next year, and temperature logs were

compared to PRISM estimates for the same locations during the intervening time period and

found to be strongly correlated (R^2 mean = 0.94, SD = 0.03). Finally, we accessed NOAA

climate reports that compared regional temperatures and precipitation in each field season to

long-term averages (NOAA National Centers for Environmental Information, 2022; 2023).

Additional details on the field survey methods are provided in Appendix S1.

Lab experiments

1. Temperature-dependent infection experiment

In September 2022, we measured the effects of temperature on the host – parasite interaction

with an experiment that used the F1 offspring of six host populations reared in the lab from

larvae collected in the 2022 field survey: two populations each from warm (14.8 – 15.1ºC rainy

season mean temperature; San Juan Capistrano and Trabuco Canyon, CA), intermediate (10.8 –

11.8ºC; Big Sur and Woodside, CA), and cool (5.3 – 6.5ºC; Shaver Lake, CA and Eugene, OR)

climates (Figure 1, Appendix S2: Table S1). We exposed each host population to the same

allopatric parasite culture isolated from an infected larva collected in Marin County, CA (10.6ºC)

(Figure 1), so that any variation observed in the temperature dependence of infection would be

due to the host population. Mosquito rearing and ciliate culturing occurred at 21.5ºC following

Ismail et al. (2023), Couper et al. (2024), and Lyberger et al. (2024).

For the experiment, mosquito eggs were hatched and day-old larvae were incubated in groups

of five with 7 mL parasite culture diluted to a standardized density of 35 ciliates per 100 uL at six

constant temperatures: 7ºC, 11ºC, 14ºC, 18ºC, 23ºC, and 28ºC (N = 6 microcosms per host

population per temperature, except for Big Sur, where N = 4 due to lower egg hatching, totaling

34 microcosms per temperature, and 204 microcosms overall). We monitored the proportions of

hosts with infections and parasite melanization spots, the survival of infected and uninfected

hosts, and *L. clarki* cell density per 100 uL liquid up to the first pupation at each temperature.

Because *Ae. sierrensis* development speed is thermally sensitive, we set monitoring frequences

that ensured at least four checkpoints spanning the host's four larval instars at each temperature;

these ranged from every two days at 28ºC to every seven days at 7ºC and 11ºC (Appendix S2:

Table S2).

2. *Host and parasite thermal responses in isolation*

To compare the thermal performance of mosquito larvae and ciliates to that of parasitism, we

incubated mosquito-only and ciliate-only microcosms alongside the infection experiment,

following the same design as above. The mosquito-only microcosms were monitored on the

same timeline as the infection experiment. For the ciliate-only microcosms (N = 6 wells per

temperature, 36 total microcosms), we measured the population growth rate by inoculating wells

with culture diluted to 4 cells per 100 uL and then counting the cells per 300 uL daily for four

days of positive growth.

To assess the temperature sensitivity of the host melanization process (removing the

temperature sensitivity of parasite attack rates), we performed a mechanical melanization assay

on fourth instar larvae from one population at 4, 8, 12, 16, 20, 24, and 28ºC (N = 6 wells of 5

mosquitoes per temperature with no ciliates, for 42 total microcosms). Additional details are

provided in Appendix S1.

Statistical analyses

Field survey

To investigate the nonlinear temperature dependence of parasitism, we tested for a quadratic

relationship between temperature and the presence of *L. clarki* infections in the surveyed tree

holes using a binomial generalized linear mixed effects model, with mean temperature, mean

temperature squared (to account for hypothesized nonlinear effects of temperature), and year as

fixed effects, and tree hole ID as a random effect. We used the glmer function in the R package

"lme4" with scaled and centered environmental variables, and assessed model fit with the R

package "DHARMa" (Bates et al., 2003; Hartig, 2017). Because interactions between survey

year and the temperature variables were not significant, we estimated the overall mean

temperature where infection was most likely to occur with a model that did not include year. We

additionally used general linearized models fit individually to the data from each survey year to

describe the temperatures where infection occurrence peaked within each field season. To

evaluate the relationship between within-tree hole habitat temperatures logged by iButtons and

the PRISM air temperature dataset used in the model, we tested for correlations, calculated the

mean differences between daily mean temperature time series recorded by iButtons and extracted

from PRISM, and used linear regression with the mean temperature recorded within a tree hole

as the dependent variable and the mean PRISM temperature estimate for that location as the

independent variable. Because there were only two field seasons, the NOAA annual climate

reports for each field season were not included in statistical analyses, but used only to

qualitatively compare the two years.

Lab experiments

To quantify the thermal sensitivity of the host – parasite interaction measured from the

laboratory experiment, we used a Bayesian approach to fit TPCs for the proportion of infected

hosts, the proportion of hosts with melanization spots, and the proportion of hosts killed by

infections. The total number of infections observed in each microcosm over the course of the

experiment depended on both initial parasite attack success and subsequent rounds of parasite

transmission following infected host deaths that released new parasites as killed mosquitoes

decomposed. We fit separate TPCs for both initial infection for the period up to the first deaths of

infected hosts at each temperature, when all observed infections resulted from the first round of

host – parasite interactions (covering checkpoints 1-3, approximately corresponding to larval

instars 1-3), and total infection rates over the entire experiment (including all host-parasite

interaction cycles for all four larval instars); we used the same procedure to fit initial and total

TPCs for the proportions of hosts melanizing attacking parasites. We additionally fit TPCs for

free-living *L. clarki* population growth rate and for the host melanization immune response

induced by mechanical penetration of the cuticle. We planned to fit TPCs for host survival in the

absence of parasitism, but >99% of larvae incubated without parasites survived across all

temperatures. Because initial parasite success is likely to determine whether there is a non-zero

infection probability over the rest of the season (Washburn et al., 1991), this is the laboratory-

based TPC that we expect to most closely align with the field-observed response of infection

occurrence to temperature.

Unimodal traits were fit with a flexTPC functional form (Equation 1) which can describe

unimodal TPCs of any skewness (Cruz-Loya et al., *preprint*). Using this flexible function to fit

TPCs of different shapes across our traits of interest allowed for ease of comparison of thermal

responses among traits and populations.

243
$$r(T) = \begin{cases} r_{max} \left[\left(\dfrac{T - T_{min}}{\alpha} \right)^{\alpha} \left(\dfrac{T_{max} - T}{1 - \alpha} \right)^{1-\alpha} \left(\dfrac{1}{T_{max} - T_{min}} \right) \right]^{\frac{\alpha(1-\alpha)}{\beta^2}} & T_{min} < T < T_{max} \\ 0 & \text{otherwise} \end{cases} \quad (1)$$

244   In this equation, T_{min} and T_{max} correspond to the minimum and maximum temperature,

245   respectively, r_{max} to the maximum trait value, and α and β are parameters that determine the

246   curve skewness and rate of decrease from the maximum trait value, respectively (Appendix S1).

247   Host melanization increased monotonically with temperature, and was fit to a 4-parameter

248   shifted logistic model (Equation 2, see Appendix S1 for derivation).

249
$$m(T) = \begin{cases} \cfrac{1}{\left[1 + (2^{1/s} - 1)\left(\cfrac{T_{50} - T_{min}}{T - T_{min}}\right)^a \right]^s} & T > T_{min} \\ 0 & \text{otherwise} \end{cases} \quad (2)$$

where T_{min} is the minimum temperature, T_{50} is the temperature at which 50% of the population

is melanized, and a and s are constants that allow for different rates of change of melanization

near T_{50} and for the response to be asymmetric, which was necessary to fit the data.

We used a binomial likelihood for all traits that were measured as proportions. For the free-living

parasite population growth rate TPC, we used a Gamma likelihood with population growth rates

obtained from the raw time-course data with the R package "growthrates" (Petzoldt, 2022).

Optimal temperatures were calculated analytically using the posterior samples of the model

parameters (Appendix S1). Details on model fitting and diagnostics are provided in Appendix

S1.

To assess population variation in lab-measured thermal performance among host

populations and the responses measured in the infection experiment, we compared 95% credible

intervals for the thermal optimum (T_{opt}), maximum (T_{max}), and minimum (T_{min}) and interpreted

non-overlapping intervals as evidence of biologically significant variation. We additionally

compared the magnitudes of the 95% credible intervals over TPC plots for each population.

Since the credible intervals for T_{opt}, T_{min}, and T_{max} overlapped for all populations for all

measured traits, we combined the data to generate species-representative TPCs, which we used

for our main analyses and interpretations. Finally, we compared the parasitism TPC to the host

and parasite trait TPCs measured in isolation.

We used R version 4.2.1 for all statistical analyses and for figure generation (R Core

Team, 2022). See Appendix S1 for additional details and R packages used.

Results

Does infection occurrence vary with temperature across the species range?

      Infection occurrence followed a hump-shaped pattern along the surveyed temperature gradient, peaking at 9.4ºC mean rainy season temperature (Figure 2). Mean temperature^2 and year were significant predictors of *L. clarki* infection presence (Table 1). Quadratic models fit individually for infection occurrence in each year peaked at 10.0ºC in 2022 and 9.2ºC in 2023, but the interactions between year and both temperature variables were not statistically significant, indicating that the temperature – infection relationship did not vary between years (Table 1). Geographically, infections were common in northern California and southern Oregon and rarer at the warm and cold ends of the surveyed temperature gradient in southern California and northern Washington, respectively (Figure 1).

      Of the two sampling years, 2023 was cooler and wetter and had a higher percentage of infected tree holes (29.7%, N = 431) compared to 2022 (12.5%, N = 320) (Figure 2) (NOAA National Centers for Environmental Information, 2022; 2023). Mean tree hole habitat temperatures logged by iButtons (N = 32 successfully retrieved) were strongly correlated with and were not systematically warmer or colder than mean PRISM air temperature estimates (linear regression intercept = -0.41, t- SE = 1.76, t-value = -0.24, p-value = 0.82; slope = 0.95, SE = 0.1, t-value = 9.11, p-value = 3.86 x 10^{-10}) (mean temperature difference = 1.3ºC, SD = 2.6ºC) (Appendix S2: Figure S1). Additional details are provided in Appendix S2.

Does the thermal response of the host – parasite interaction depend on the host population?

We found similar thermal optima, minima and maxima across the six host populations in the infection experiment within each of the three measured interaction outcomes of infection,

melanization of parasites, and mortality (Figure 3; Appendix S2: Figures S2-S11; Tables S3-S7).

Although the mean estimated T_{opt}, T_{min}, and T_{max} varied slightly among populations for each

trait, all 95% credible intervals across populations for each outcome x parameter combination

overlapped. Therefore, we combined the data for all host populations to estimate broadly

representative TPCs for the host – parasite interaction, with reduced uncertainty compared to the

population-specific TPCs (Figure 4; Appendix S2: Figure S12; Table S9). The only trait in which

we observed clear intraspecific variation was in mortality around 10ºC, which was higher in one

intermediate climate population (Big Sur (BIG)) compared to the other (Woodside (JR)) and to

one cold climate population (Eugene (EUG)) (Figure S13).

How do different components of the host – parasite interaction relate to the overall thermal

sensitivity of parasitism?

Although free-living parasite population growth was maximized at warm temperatures, infection

and parasite-induced mortality peaked at significantly colder temperatures, corresponding to

conditions that suppressed host defense mechanisms, namely the melanization immune response

and larval development speed (Figure 4). Specifically, the experimentally measured initial

infection rate peaked at 10ºC (95% CI: 8.7 ºC, 11.3 ºC), and the total infection rate peaked at

8.4ºC (95% CI: 6.8 ºC, 9.4 ºC), much lower temperatures than the 19.5ºC (95% CI: 18.9 ºC, 20.2

ºC) optimum for free-living parasite population growth, but near the field-estimated 10.0ºC and

9.2ºC peaks in 2022 and 2023, respectively (Figure 4A; Appendix S2: Figure S12; Tables S8-9).

The mechanically assayed host melanization immune response increased with temperature up to

20ºC, where it plateaued (Figure 4A; Appendix S2: Table S10). At warmer temperatures,

attacking parasites faced not only a stronger host immune response, but also shorter time

windows in which to infect hosts without being shed with the cuticle during larval molting and

pupation or eaten by later larval instars. The time to the first observed pupation shortened as the

temperature increased, occurring after 9 days at 28ºC compared to 57 days at 7ºC (Figure 4B),

consistent with previously measured *Ae. sierrensis* larval development rate TPCs that peaked at

27.7ºC (Couper et al., 2024). Consequently, rapid pupation interrupted upward trajectories of

infection and mortality at intermediate temperatures, contributing to the decline in parasitism

above 11ºC, and to the lower T_{opt} of total compared to initial infection. At colder temperatures,

additional opportunities for parasitism were also generated as infected host deaths beginning in

the third week of the experiment maintained higher *L. clarki* densities throughout extended host

developmental periods (Figure S14). In comparison, at the warmest temperature with infections

(18ºC), the contributions of infected host deaths to subsequent transmission were limited by their

rarity and proximity to the onset of pupation.

Discussion

In this study, we used complementary field observations and lab experiments to understand

how temperature affects a host-parasite interaction across a large, climatically variable species

range. We found that infection peaked at sites with intermediate temperatures (9.4ºC) in the field

across two sampling years and was higher in a colder, wetter year, aligning with the unimodal

temperature-dependence of infection measured in the lab, which peaked at 10ºC. This reflects the

interacting thermal biology of multiple mosquito and ciliate traits, which together led to an

optimal temperature for parasitism (8-10ºC for total versus initial infection) that was colder than

the optimal temperature for free-living parasite performance (19.5ºC). Infection rates of

mosquito populations from different climates had similar thermal responses in the lab, and a

single TPC described the infection process across the entire surveyed area. These findings

suggest that for systems with strong temperature dependence, TPCs measured under simplified

lab conditions can capture patterns of parasitism across large geographic temperature gradients

in complex natural settings, by quantifying the underlying thermal biology mechanisms.

Lab experiments capture nonlinear parasitism thermal responses in the field

The results add to evidence that temperature nonlinearly affects parasitism not only in the

lab, but also in nature, making laboratory and field studies that are designed to capture nonlinear

temperature effects valuable tools for understanding disease risk (Dudney et al., 2021; Kirk et al.,

2024; Shocket, Ryan et al., 2018; Shocket, Strauss et al., 2018). However, despite the similarities

between the field observations and the lab-measured parasitism TPC, we cannot rule out the

influence of other factors. For example, while the role of precipitation in this system has not been

assessed, the colder survey year with higher infection prevalence was also wetter, suggesting that

precipitation patterns that determined when and for how long tree holes were water-filled may

have also affected the potential for parasitism to occur. Beyond mean temperature differences,

heatwaves or cold snaps occurring during key points in the host – parasite interaction may have

also contributed to interannual variation in infection presence (Claar & Wood, 2020; Altman et

al., 2016). These limitations should be addressed with field experiments that manipulate

temperatures in natural tree holes at different latitudes or elevations; lab experiments that

incorporate additional environmental complexity; and long term repeated yearly sampling of the

same tree holes.

Intraspecific variation

Very few studies investigate the impact of local thermal adaptation on species distributions

and host – parasite interactions, a gap often noted in the literature (Sternberg & Thomas, 2014;

Urban et al., 2016; Couper et al., 2021; Aleuy et al., 2023). In addressing this gap, we did not

find evidence of local temperature adaptation affecting infection patterns across temperatures in

the lab or the field. While our laboratory experiment specifically assessed host local adaptation,

the invariant temperature dependence of infection we observed is consistent with the similar

temperature-infection relationship in the field, and with our previous work comparing sympatric

and allopatric host – parasite interactions. Specifically, for nine *Ae. sierrensis* and *L. clarki*

populations with free-living parasite T_{opt} ranging from 16.5ºC to 23.5ºC, infection rates measured

at three temperatures peaked at the intermediate temperature of 12.5ºC, substantially cooler than

the free-living growth optima (Lyberger et al., 2024). Cumulatively, these findings suggest that

although both *Ae. sierrensis* pupal development and *L. clarki* free-living population growth rates

are locally adapted to temperature, this intraspecific variation is unlikely to directly affect host –

parasite interactions. (Lyberger et al., 2024; Couper et al., 2024; Ismail et al., 2023). Variability

in the infection T_{opt} should be further tested through TPC experiments with additional host –

parasite population pairs, but differences in factors such as host fitness and adaptation to

parasites may be more important sources of intraspecific variation in parasite impacts. For

example, in this experiment, the mosquito population with the lowest egg hatching rate, BIG, had

a higher mortality rate near the infection T_{opt} compared to two other populations; and in previous

experiments, individual host populations varied in their susceptibility to different parasite strains

(Ismail et al., 2023; Lyberger et al., 2024)

Importantly, this is one of the only ectothermic host – parasite systems for which interaction

thermal sensitivity has been assessed for populations from different climates, along with a

stickleback – tapeworm system in which effects of temperature on parasitism were similar

among four study populations collected from across a similar geographic temperature gradient

(Franke et al., 2019). Notably, intraspecific variation in parasitism temperature sensitivity has

been shown for levels of biological organization both above and below the population level that

we assessed, for *Batrachochytrium dendrobatitis* infection both among different frog host

species, and among salamander host color morphs within a single population (Cohen et al., 2017;

Venesky et al., 2022). However, if population thermal sensitivities of parasitism are commonly

consistent across species ranges, broadly applicable TPCs should be easily attainable for the

many host – parasite systems that are amenable to laboratory studies, and field studies may also

be able to capture nonlinear responses of parasitism to temperature.

Mechanisms of parasitism temperature dependence

Our finding that parasitism peaked at temperatures that constrained host development and

defense—rather than at the thermal optimum for parasite growth and attack—aligns with theory

on the thermal biology of biotic interactions. Thermal asymmetry theory posits that parasitism

peaks at temperatures where parasites outperform hosts, which may differ from optimal

temperatures for the parasites themselves (Dell et al., 2014; Cohen et al., 2017). Our study

supports the value of approaching this problem as the outcome of multiple interacting host and

parasite traits with varying thermal responses (Paull et al., 2012; Shocket et al., 2018; Sun et al.,

2023).

Conclusions

Overall, our characterization of the temperature – infection relationship of the *Ae. sierrensis* host

– *L. clarki* parasite interaction across settings demonstrates how controlled lab experiments can

be paired with large-scale field observations to understand the complex thermal biology of

parasitism. Climate warming is expected to cause geographic contractions, expansions, and shifts

in host thermal refuges from disease (Altizer et al., 2013; Gsell et al., 2023). Our results suggest

that parasite burdens on *Ae. sierrensis* may decline at southern latitudes and low elevations as

thermally suitable conditions for parasitism shift northwards and upwards into areas where cold

conditions currently limit *L. clarki* infections. Whether the distribution of parasitism ultimately

tracks climate warming will depend on multiple factors, including parasite dispersal, the pace of

thermal adaptation in host and parasite traits that allow local populations of each species to

persist individually, and the effects of other environmental changes in conjunction with

increasing mean temperatures, such as increasingly extreme precipitation patterns (Swain et al.,

2018). Understanding how temperature affects parasitism in inherently complex and variable

natural systems remains a pressing problem with broad implications for current and future

disease risk management. Simple pairings of field surveys and lab experiments can offer

straightforward means for approaching this challenge.

Acknowledgments

Otto Seppälä, Jan Washburn, Nona Chiariello, and Angie Nakano played critical roles in

establishing the study system. Our field sampling and mosquito rearing efforts were made

possible by assistance from staff at vector control agencies for the California counties of San

Mateo, Santa Clara, Marin-Sonoma, Solano, Alameda, Shasta, Placer, Santa Barbara, and

Riverside; Jasper Ridge Biological Preserve 'Ootchamin 'Ooyakma; the UC Reserves at

Emerson Oaks, Dawson Los Monos Canyon, and Hopland; and Andrew Rivera of Clarke Pest

Control. Mordecai lab members helped with setting up the infection experiment. JEF was funded

by the Bing-Mooney Fellowship and the ARCS Foundation. EAM and MCL were supported by

the National Institutes of Health (R35GM133439 and R01AI168097). EAM was supported by

the National Science Foundation (DEB-2011147 with Fogarty International Center), the National

(R01AI102918), the Stanford Woods Institute for the Environment, the Stanford Center for

Innovation in Global Health, and the Stanford Doerr School of Sustainability. KL was funded by

the NSF Postdoctoral Research Fellowships in Biology Program under Grant No. 2208947. LIC

was funded by the NSF Postdoctoral Research Fellowship in Biology. MCL was supported by a

Stanford PRISM Baker Fellowship.

Conflict of Interest Statement

The authors declare no conflict of interest.

References

Ahrens, Collin W., Richard A. Mazanec, Trudy Paap, Katinka X. Ruthrof, Anthea Challis, Giles

Hardy, Margaret Byrne, David T. Tissue, and Paul D. Rymer. 2019. "Adaptive Variation

for Growth and Resistance to a Novel Pathogen along Climatic Gradients in a Foundation

Tree." *Evolutionary Applications* 12 (6): 1178–90. https://doi.org/10.1111/eva.12796.

Aleuy, O. Alejandro, Stephanie J. Peacock, Péter K. Molnár, Kathreen E. Ruckstuhl, and Susan J.

Kutz. 2023. "Local Thermal Adaptation and Local Temperature Regimes Drive the

Performance of a Parasitic Helminth under Climate Change: The Case of *Marshallagia*

marshalli from Wild Ungulates." *Global Change Biology* 29 (22): 6217–33.

https://doi.org/10.1111/gcb.16918.

Altman, Karie A., Sara H. Paull, Pieter T. J. Johnson, et al. 2016. "Host and Parasite Thermal

Acclimation Responses Depend on the Stage of Infection." *Journal of Animal Ecology* 85

(4): 1014–24. https://doi.org/10.1111/1365-2656.12510.

Altizer, Sonia, Richard S. Ostfeld, Pieter T. J. Johnson, Susan Kutz, and C. Drew Harvell. 2013.

"Climate Change and Infectious Diseases: From Evidence to a Predictive Framework."

Science 341 (6145): 514–19. https://doi.org/10.1126/science.1239401.

Angilletta, Michael J. 2009. *Thermal Adaptation: A Theoretical and Empirical Synthesis*.

Oxford, UK: Oxford Univ. Press

Bates, Douglas, Martin Maechler, Ben Bolker, and Steven Walker. 2003. "Lme4: Linear Mixed-

Effects Models Using 'Eigen' and S4." https://doi.org/10.32614/CRAN.package.lme4.

Boots, M. 2000. "Density-Independent Resource Limitation and the Transmission of an Insect

Pathogen." *Oecologia* 124 (2): 172–75. https://doi.org/10.1007/s004420050004.

Broberg, Leonard E., and William E. Bradshaw. 1997. "*Lambornella clarki* (Ciliophora:

Tetrahymenidae) Resistance in *Aedes sierrensis* (Diptera: Culicidae): Population

Differentiation in a Quantitative Trait and Allozyme Loci." *Journal of Medical*

Entomology 34 (1): 38–45. https://doi.org/10.1093/jmedent/34.1.38.

Brown, Joel J., Mercedes Pascual, Michael C. Wimberly, Leah R. Johnson, and Courtney C.

Murdock. 2023. "Humidity – The Overlooked Variable in the Thermal Biology of

Mosquito-Borne Disease." *Ecology Letters* 26 (7): 1029–49.

https://doi.org/10.1111/ele.14228.

Christensen, Bruce M., Jianyong Li, Cheng-Chen Chen, and Anthony J. Nappi. 2005.

"Melanization Immune Responses in Mosquito Vectors." *Trends in Parasitology* 21 (4):

192–99. https://doi.org/10.1016/j.pt.2005.02.007.

Claar, Danielle C., and Chelsea L. Wood. 2020. "Pulse Heat Stress and Parasitism in a Warming

World." *Trends in Ecology & Evolution* 35 (8): 704–15.

https://doi.org/10.1016/j.tree.2020.04.002.

Clark, T.B., and D.G. Brandl. 1976. "Observations on the Infection of *Aedes sierrensis* by a

Tetrahymenine Ciliate." *Journal of Invertebrate Pathology* 28 (3): 341–49.

https://doi.org/10.1016/0022-2011(76)90009-4.

Cohen, Jeremy M., Matthew D. Venesky, Erin L. Sauer, David J. Civitello, Taegan A. McMahon,

Elizabeth A. Roznik, and Jason R. Rohr. 2017. "The Thermal Mismatch Hypothesis

Explains Host Susceptibility to an Emerging Infectious Disease." *Ecology Letters* 20 (2):

184–93. https://doi.org/10.1111/ele.12720.

Couper, Lisa I., Johannah E. Farner, Jamie M. Caldwell, Marissa L. Childs, Mallory J. Harris,

Devin G. Kirk, Nicole Nova, et al. 2021. "How Will Mosquitoes Adapt to Climate

Warming?" *eLife* 10 (August):e69630. https://doi.org/10.7554/eLife.69630.

Couper, Lisa I., Johannah E. Farner, Kelsey P. Lyberger, Alexandra S. Lee, and Erin A.

Mordecai. 2024. "Mosquito Thermal Tolerance Is Remarkably Constrained across a

Large Climatic Range." *Proceedings of the Royal Society B: Biological Sciences* 291

(2015): 20232457. https://doi.org/10.1098/rspb.2023.2457.

Cruz-Loya, Mauricio, Erin A. Mordecai, and Van M. Savage. 2024. "A Flexible Model for

Thermal Performance Curves." *bioRxiv*, January, 2024.08.01.605695.

https://doi.org/10.1101/2024.08.01.605695.

Darsie, Richard F., and Ronald A. Ward. 2005. *Identification and Geographical Distribution of*

the Mosquitoes of North America, North of Mexico. University Press of Florida.

Dell, Anthony I., Samraat Pawar, and Van M. Savage. 2014. "Temperature Dependence of

Trophic Interactions Are Driven by Asymmetry of Species Responses and Foraging

Strategy." *Journal of Animal Ecology* 83 (1): 70–84. https://doi.org/10.1111/1365-

2656.12081.

Dudney, Joan, Claire E. Willing, Adrian J. Das, Andrew M. Latimer, Jonathan C. B. Nesmith,

and John J. Battles. 2021. "Nonlinear Shifts in Infectious Rust Disease Due to Climate

Change." *Nature Communications* 12 (1): 5102. https://doi.org/10.1038/s41467-021-

25182-6.

Franke, Frederik, Nadja Raifarth, Joachim Kurtz, and Jörn P. Scharsack. 2019. "Consequences of

Divergent Temperature Optima in a Host–Parasite System." *Oikos* 128 (6): 869–80.

https://doi.org/10.1111/oik.05864.

Gaitán-Espitia, Juan Diego, Leonardo D. Bacigalupe, Tania Opitz, Nelson A. Lagos, Tania

Timmermann, and Marco A. Lardies. 2014. "Geographic Variation in Thermal

Physiological Performance of the Intertidal Crab *Petrolisthes violaceus* along a

Latitudinal Gradient." *Journal of Experimental Biology* 217 (24): 4379–86.

https://doi.org/10.1242/jeb.108217.

Gehman, Alyssa-Lois M., Richard J. Hall, and James E. Byers. 2018. "Host and Parasite Thermal

Ecology Jointly Determine the Effect of Climate Warming on Epidemic Dynamics."

Proceedings of the National Academy of Sciences 115 (4): 744–49.

https://doi.org/10.1073/pnas.1705067115.

Gsell, Alena S., Arjen Biere, Wietse de Boer, Irene de Bruijn, Götz Eichhorn, Thijs Frenken,

Stefan Geisen, et al. 2023. "Environmental Refuges from Disease in Host–Parasite

Interactions under Global Change." *Ecology* 104 (4): e4001.

https://doi.org/10.1002/ecy.4001.

Hall, Spencer R., Joseph L. Simonis, Roger M. Nisbet, Alan J. Tessier, and Carla E. Cáceres.

2009. "Resource Ecology of Virulence in a Planktonic Host-Parasite System: An

Explanation Using Dynamic Energy Budgets." *The American Naturalist* 174 (2): 149–62.

https://doi.org/10.1086/600086.

Hartig, Florian. 2017. "Package 'Dharma.'" *R Package*.

Ismail, Sherine, Johannah Farner, Lisa Couper, Erin Mordecai, and Kelsey Lyberger. 2023.

"Temperature and Intraspecific Variation Affect Host–Parasite Interactions." *Oecologia*,

November. https://doi.org/10.1007/s00442-023-05481-z.

Jordan, R. G., and W. E. Bradshaw. 1978. "Geographic Variation in the Photoperiodic Response

of the Western Tree-Hole Mosquito, *Aedes sierrensis*." *Annals of the Entomological

Society of America* 71 (4): 487–90. https://doi.org/10.1093/aesa/71.4.487.

Jordan, Robert G. 1980a. "Embryonic Diapause in Three Populations of the Western Tree Hole

Mosquito, *Aedes sierrensis*." *Annals of the Entomological Society of America* 73 (4):

357–59. https://doi.org/10.1093/aesa/73.4.357.

———. 1980b. "Geographic Differentiation in the Development *of Aedes sierrensis* (Diptera:

Culicidae) in Nature." *The Canadian Entomologist* 112 (2): 205–10.

https://doi.org/10.4039/Ent112205-2.

Jurriaans, S., and M. O. Hoogenboom. 2019. "Thermal Performance of Scleractinian Corals

along a Latitudinal Gradient on the Great Barrier Reef." *Philosophical Transactions of

the Royal Society B: Biological Sciences* 374 (1778): 20180546.

https://doi.org/10.1098/rstb.2018.0546.

Kirk, Devin, Natalie Jones, Stephanie Peacock, Jessica Phillips, Péter K. Molnár, Martin

Krkošek, and Pepijn Luijckx. 2018. "Empirical Evidence That Metabolic Theory

Describes the Temperature Dependency of Within-Host Parasite Dynamics." *PLOS*

Biology 16 (2): e2004608. https://doi.org/10.1371/journal.pbio.2004608.

Kirk, Devin, Samantha Straus, Marissa L. Childs, Mallory Harris, Lisa Couper, T. Jonathan

Davies, Coreen Forbes, et al. 2024. "Temperature Impacts on Dengue Incidence Are

Nonlinear and Mediated by Climatic and Socioeconomic Factors: A Meta-Analysis."

PLOS Climate 3 (3): e0000152. https://doi.org/10.1371/journal.pclm.0000152.

Kutz, S. J., E. P. Hoberg, L. Polley, and E. J. Jenkins. 2005. "Global Warming Is Changing the

Dynamics of Arctic Host-Parasite Systems." *Proceedings: Biological Sciences* 272

(1581): 2571–76.

Labaude, Sophie, Yannick Moret, Frank Cézilly, Charel Reuland, and Thierry Rigaud. 2017.

"Variation in the Immune State of *Gammarus pulex* (Crustacea, Amphipoda) According

to Temperature: Are Extreme Temperatures a Stress?" *Developmental & Comparative*

Immunology 76 (November):25–33. https://doi.org/10.1016/j.dci.2017.05.013.

Laughton, Alice M., Cian O. O'Connor, and Robert J. Knell. 2017. "Responses to a Warming

World: Integrating Life History, Immune Investment, and Pathogen Resistance in a

Model Insect Species." *Ecology and Evolution* 7 (22): 9699–9710.

https://doi.org/10.1002/ece3.3506.

Luković, Jelena, John C. H. Chiang, Dragan Blagojević, and Aleksandar Sekulić. 2021. "A Later

Onset of the Rainy Season in California." *Geophysical Research Letters* 48 (4):

e2020GL090350. https://doi.org/10.1029/2020GL090350.

Lyberger, Kelsey, Johannah Farner, Lisa Couper, and Erin A. Mordecai. 2024. "A Mosquito

Parasite Is Locally Adapted to Its Host but Not Temperature." *The American Naturalist*

204 (2): 121–32. https://doi.org/10.1086/730522.

Martin, Lindsay E., and Julián F. Hillyer. 2024. "Higher Temperature Accelerates the Aging-

Dependent Weakening of the Melanization Immune Response in Mosquitoes." *PLOS*

Pathogens 20 (1): e1011935. https://doi.org/10.1371/journal.ppat.1011935.

Mazé-Guilmo, Elise, Simon Blanchet, Olivier Rey, Nicolas Canto, and Géraldine Loot. 2016.

"Local Adaptation Drives Thermal Tolerance among Parasite Populations: A Common

Garden Experiment." *Proceedings of the Royal Society B: Biological Sciences* 283

(1830): 20160587. https://doi.org/10.1098/rspb.2016.0587.

Mordecai, Erin A., Jamie M. Caldwell, Marissa K. Grossman, Catherine A. Lippi, Leah R.

Johnson, Marco Neira, Jason R. Rohr, et al. 2019. "Thermal Biology of Mosquito-borne

Disease." *Ecology Letters*, July, ele.13335. https://doi.org/10.1111/ele.13335.

Mordecai, Erin A., Jeremy M. Cohen, Michelle V. Evans, Prithvi Gudapati, Leah R. Johnson,

Catherine A. Lippi, Kerri Miazgowicz, et al. 2017. "Detecting the Impact of Temperature

on Transmission of Zika, Dengue, and Chikungunya Using Mechanistic Models." *PLOS*

Neglected Tropical Diseases 11 (4): e0005568.

https://doi.org/10.1371/journal.pntd.0005568.

Mordecai, Erin A., Krijn P. Paaijmans, Leah R. Johnson, Christian Balzer, Tal Ben-Horin, Emily

de Moor, Amy McNally, et al. 2013. "Optimal Temperature for Malaria Transmission Is

Dramatically Lower than Previously Predicted." *Ecology Letters* 16 (1): 22–30.

https://doi.org/10.1111/ele.12015.

Murdock, C. C., Krijn P. Paaijmans, Andrew S. Bell, Jonas G. King, Julián F. Hillyer, Andrew F. Read, and Matthew B. Thomas. 2012. "Complex Effects of Temperature on Mosquito Immune Function." *Proceedings of the Royal Society B: Biological Sciences* 279 (1741): 3357–66. https://doi.org/10.1098/rspb.2012.0638.

NOAA National Centers for Environmental Information. 2022. "Monthly National Climate Report for 2022." https://www.ncei.noaa.gov/access/monitoring/monthly-report/national/202408.

———. 2023. "Monthly National Climate Report for 2023." https://www.ncei.noaa.gov/access/monitoring/monthly-report/national/202408.

Padfield, Daniel, Meaghan Castledine, and Angus Buckling. 2020. "Temperature-Dependent Changes to Host–Parasite Interactions Alter the Thermal Performance of a Bacterial Host." *The ISME Journal* 14 (2): 389–98. https://doi.org/10.1038/s41396-019-0526-5.

Palmer-Young, Evan C., Lyna Ngor, Rodrigo Burciaga Nevarez, Jason A. Rothman, Thomas R. Raffel, and Quinn S. McFrederick. 2019. "Temperature Dependence of Parasitic Infection and Gut Bacterial Communities in Bumble Bees." *Environmental Microbiology* 21 (12): 4706–23. https://doi.org/10.1111/1462-2920.14805.

Paull, Sara H., Bryan E. LaFonte, and Pieter T. J. Johnson. 2012. "Temperature-Driven Shifts in a Host-Parasite Interaction Drive Nonlinear Changes in Disease Risk." *Global Change Biology* 18 (12): 3558–67. https://doi.org/10.1111/gcb.12018.

Penczykowski, Rachel M., Spencer R. Hall, David J. Civitello, and Meghan A. Duffy. 2014. "Habitat Structure and Ecological Drivers of Disease." *Limnology and Oceanography* 59 (2): 340–48. https://doi.org/10.4319/lo.2014.59.2.0340.

Petzoldt, Thomas. 2022. "R Package Growthrates."

Praskievicz, Sarah, and Heejun Chang. 2009. "Winter Precipitation Intensity and ENSO/PDO

Variability in the Willamette Valley of Oregon." *International Journal of Climatology* 29

(13): 2033–39. https://doi.org/10.1002/joc.1838.

PRISM Climate Group, and Oregon State University. 2021. "PRISM Gridded Climate Data."

https://prism.oregonstate.edu.

Ragonese, Isabella G., Maya R. Sarkar, Richard J. Hall, and Sonia Altizer. 2024. "Extreme Heat

Reduces Host and Parasite Performance in a Butterfly–Parasite Interaction." *Proceedings*

of the Royal Society B: Biological Sciences 291 (2015): 20232305.

https://doi.org/10.1098/rspb.2023.2305.

R Core Team. 2022. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Ryan, Sadie J., Colin J. Carlson, Erin A. Mordecai, and Leah R. Johnson. 2019. "Global

Expansion and Redistribution of *Aedes*-Borne Virus Transmission Risk with Climate

Change." *PLOS Neglected Tropical Diseases* 13 (3): e0007213.

https://doi.org/10.1371/journal.pntd.0007213.

Seppälä, Otto, Katri Liljeroos, Anssi Karvonen, and Jukka Jokela. 2008. "Host Condition as a

Constraint for Parasite Reproduction." *Oikos* 117 (5): 749–53.

https://doi.org/10.1111/j.0030-1299.2008.16396.x.

Shah, Melisa M., Amy R. Krystosik, Bryson A. Ndenga, Francis M. Mutuku, Jamie M. Caldwell,

Victoria Otuka, Philip K. Chebii, et al. 2019. "Malaria Smear Positivity among Kenyan

Children Peaks at Intermediate Temperatures as Predicted by Ecological Models."

Parasites & Vectors 12 (1): 288. https://doi.org/10.1186/s13071-019-3547-z.

Shocket, Marta S., Alexander T. Strauss, Jessica L. Hite, Maja Šljivar, David J. Civitello,

Meghan A. Duffy, Carla E. Cáceres, and Spencer R. Hall. 2018. "Temperature Drives

Epidemics in a Zooplankton-Fungus Disease System: A Trait-Driven Approach Points to

Transmission via Host Foraging." *The American Naturalist* 191 (4): 435–51.

https://doi.org/10.1086/696096.

Shocket, Marta S., Anna B. Verwillow, Mailo G. Numazu, Hani Slamani, Jeremy M. Cohen,

Fadoua El Moustaid, Jason Rohr, Leah R. Johnson, and Erin A. Mordecai. 2020.

"Transmission of West Nile and Five Other Temperate Mosquito-Borne Viruses Peaks at

Temperatures between 23°C and 26°C." *eLife* 9 (September):e58511.

https://doi.org/10.7554/eLife.58511.

Shocket, Marta S., Sadie J. Ryan, and Erin A. Mordecai. 2018. "Temperature Explains Broad

Patterns of Ross River Virus Transmission." *eLife* 7 (August):e37762.

https://doi.org/10.7554/eLife.37762.

Sinclair, Brent J., Caroline M. Williams, and John S. Terblanche. 2012. "Variation in Thermal

Performance among Insect Populations." *Physiological and Biochemical Zoology* 85 (6):

594–606. https://doi.org/10.1086/665388.

Sternberg, Eleanore D., and Matthew B. Thomas. 2014. "Local Adaptation to Temperature and

the Implications for Vector-Borne Diseases." *Trends in Parasitology* 30 (3): 115–22.

https://doi.org/10.1016/j.pt.2013.12.010.

Sun, Syuan-Jyun, Marcin K. Dziuba, Riley N. Jaye, and Meghan A. Duffy. 2023. "Temperature

Modifies Trait-Mediated Infection Outcomes in a *Daphnia*–Fungal Parasite System."

Philosophical Transactions of the Royal Society B: Biological Sciences 378 (1873):

20220009. https://doi.org/10.1098/rstb.2022.0009.

Suwanchaichinda, Chansak, and Susan M. Paskewitz. 1998. "Effects of Larval Nutrition, Adult

Body Size, and Adult Temperature on the Ability of *Anopheles gambiae* (Diptera:

Culicidae) to Melanize Sephadex Beads." *Journal of Medical Entomology* 35 (2): 157–

61. https://doi.org/10.1093/jmedent/35.2.157.

Swain, Daniel L., Baird Langenbrunner, J. David Neelin, and Alex Hall. 2018. "Increasing

Precipitation Volatility in Twenty-First-Century California." *Nature Climate Change* 8

(5): 427–33. https://doi.org/10.1038/s41558-018-0140-y.

Taylor, Rachel A., Sadie J. Ryan, Catherine A. Lippi, David G. Hall, Hossein A. Narouei-

Khandan, Jason R. Rohr, and Leah R. Johnson. 2019. "Predicting the Fundamental

Thermal Niche of Crop Pests and Diseases in a Changing World: A Case Study on Citrus

Greening." *Journal of Applied Ecology*, July, 1365-2664.13455.

https://doi.org/10.1111/1365-2664.13455.

Tesla, Blanka, Leah R. Demakovsky, Erin A. Mordecai, Sadie J. Ryan, Matthew H. Bonds,

Calistus N. Ngonghala, Melinda A. Brindley, and Courtney C. Murdock. 2018.

"Temperature Drives Zika Virus Transmission: Evidence from Empirical and

Mathematical Models." *Proceedings of the Royal Society B: Biological Sciences* 285

(1884): 20180795. https://doi.org/10.1098/rspb.2018.0795.

Thomas, Matthew B., and Simon Blanford. 2003. "Thermal Biology in Insect-Parasite

Interactions." *Trends in Ecology & Evolution* 18 (7): 344–50.

https://doi.org/10.1016/S0169-5347(03)00069-7.

Urban, M. C., G. Bocedi, A. P. Hendry, J.-B. Mihoub, G. Pe'er, A. Singer, J. R. Bridle, et al.

2016. "Improving the Forecast for Biodiversity under Climate Change." *Science* 353

(6304): aad8466. https://doi.org/10.1126/science.aad8466.

Venesky, Matthew D., Joseph DeMarchi, Cari Hickerson, and Carl D. Anthony. 2022. "Does the

Thermal Mismatch Hypothesis Predict Disease Outcomes in Different Morphs of a

Terrestrial Salamander?" *Journal of Experimental Zoology Part A: Ecological and*

Integrative Physiology, 1–10. https://doi.org/10.1002/jez.2581.

Washburn, J., M. Gross, D. Mercer, and Anderson. 1988. "Predator-Induced Trophic Shift of a

Free-Living Ciliate: Parasitism of Mosquito Larvae by Their Prey." *Science* 240 (4856):

1193–95. https://doi.org/10.1126/science.3131877.

Washburn, Jan O., and John R. Anderson. 1986. "Distribution of *Lambornella clarki* (Ciliophora:

Tetrahymenidae) and Other Mosquito Parasites in California Treeholes." *Journal of*

Invertebrate Pathology 48 (3): 296–309. https://doi.org/10.1016/0022-2011(86)90058-3.

Washburn, Jan O., John R. Anderson, and David R. Mercer. 1991. "Parasitism of Newly-Hatched

Aedes sierrensis (Diptera: Culicidae) Larvae by *Lambornella clarki* (Ciliophora:

Tetrahymenidae) Following Habitat Flooding." *Journal of Invertebrate Pathology* 58 (1):

67–74. https://doi.org/10.1016/0022-2011(91)90163-K.

Tables

Table 1. Results of a generalized linear mixed-effects model with tree hole infection status

as the dependent variable and mean annual temperature, mean annual temperature^2, and

survey year as fixed effects, and tree hole ID as a random effect.

Predictor	Estimate	Standard error	Z-value	p-value
Mean annual temperature	-0.22	0.22	-1.01	3.13×10^{-1}
Mean annual temperature^2	-0.43	0.21	-2.01	**4.45×10^{-2}**
Survey year	1.31	0.34	3.90	**9.5×10^{-5}**
*Mean annual temperature * survey year*	-0.45	0.26	-1.72	8.5×10^{-2}
*Mean annual temperature^2 * survey year*	-0.127	0.247	-0.52	6.34×10^{-1}

699 P-values < 0.05 are displayed in bold to highlight statistical significance.

Figure captions

Figure 1. Tree holes were surveyed, and experimental host and parasite populations reared, from across a large geographic temperature gradient on the western coast of the United States during the (A) 2022 and (B) 2023 rainy seasons. The mapped color gradient shows mean temperature during the rainy season in Washington, Oregon, and California, USA, from the 4 km PRISM climate dataset. Points show the locations of 507 total surveyed tree holes. *L. clarki* infection presence is indicated by black points, and absence by white points. In (A), colored ovals indicate the source locations of the six host populations used in the infection experiment (from north to south: EUG = Eugene, OR; JR = Woodside, CA; FRS = Shaver Lake, CA; BIG = Big Sur, CA; WHIT = Trabuco Canyon, CA; CAS = San Juan Capistrano, CA); the gray oval shows the host population used in the melanization response experiment (PAR = Paso Robles, CA); and the gray rectangle shows the source location of the ciliate parasite culture used in the infection experiment (Marin = Novato, CA). In (B), green outlines indicate tree holes in which iButtons logged temperature data during the year between the two sampling visits.

Figure 2. Infection probability followed a consistent hump-shaped relationship with mean temperature during the rainy season, and was higher overall in 2023 compared to 2022. (A) Counts of infected (black) and uninfected (gray) tree holes across the climate gradient in each survey year. **(B)** Generalized linear mixed-effects model predictions of infection probability by temperature and year are plotted across the surveyed climate gradient for each year of the field survey. The solid and dashed lines show the mean model predictions for 2022 and 2023, respectively, and the shading shows the 95% confidence intervals.

Figure 3. Thermal responses of host – parasite interaction outcomes were similar among host populations from warm, intermediate, and cold source climates. For each outcome

724 measured in the infection experiment, mean estimated TPCs for each of six host populations are

725 shown (lines) with 95% credible intervals (shading). The left column shows **(A)** initial infection

726 and **(B)** parasite melanization responses up to the third experiment checkpoint, when the first

727 deaths of infected larvae released additional parasites into the environment. The right column

728 shows the **(C)** total infection, **(D)** parasite melanization, and **(E)** parasite-induced mortality rates

729 across temperatures at the end of the experiment. Mortality rates were significantly higher near

730 10 ºC for population BIG compared to populations JR and EUG. The TPCs shown here are

731 presented individually, with the experiment data, in Appendix S2 (Figures S2, S4, S6, S8, S10).

732 **Figure 4. Experimental infections were most likely to occur at cold incubation**

733 **temperatures where the host melanization response and development were suppressed. (A)**

734 Points show mean values, error bars show +/- 1 SE, lines show mean estimated TPCs, and

735 shading shows 95% credible intervals for key host – parasitism interaction outcomes and

736 processes: free-living parasite population growth (green); the host mechanically stimulated

737 melanization immune response (black); the initial infection rate (light pink), calculated up to the

738 first fatality at each temperature; the total infection rate (dark pink); and total mortality (gray)

739 calculated up to the first pupation at each temperature. The infection data shown include all host

740 populations in the infection experiment. **(B)** Infection (top; dark pink), parasite-induced mortality

741 (middle; light gray), and parasite melanization (bottom; dark gray) counts are plotted over time

742 for each temperature used in the infection experiment. Colored points show data from individual

743 microcosms, and black points and lines show mean trajectories +/- 1 SE. For each temperature,

744 the final experiment checkpoint was at the time of the first observed pupation.

Figure 1

2022

2023

Field data

● infected tree hole

○ uninfected tree hole

○ tree hole habitat with iButton temperature data

Field-collected populations used in lab experiments

infection experiment host

PAR melanization assay host

Marin infection experiment parasite

Map

18.5

PRISM rainy season mean temperature (°C)

-10

0   250   500 km

N

40°

-120°

-120°

A

B

EUG

Marin

JR

BIG

PAR

FRS

WHIT

CAS

Figure 2

Figure 3

Figure 4