Potato Late Blight Control with a Botanical Product and Reduced Copper Applications

Tomke Musa[1*], Andreas Kägi[1], Haruna Gütlin[1], Sylvain Schnee[2], Josep Massana-Codina[2], Karen E. Sullam[1,3*]

[1]Extension Arable Crops, Agroscope, Reckenholzstrasse 191, 8046 Zurich, Switzerland.
[2]Mycology, Agroscope, Rte de Duillier 60, 1260 Nyon, Switzerland.
[3]Present Address: Molecular Ecology, Agroscope, Reckenholzstrasse 191, 8046 Zurich, Switzerland.

*Corresponding author(s). E-mail(s): tomke.musa@agroscope.admin.ch;
karen.sullam@agroscope.admin.ch;
Contributing authors: andreas.kaegi@agroscope.admin.ch;
haruna.Guetlin@agroscope.admin.ch; sylvain.Schnee@agroscope.admin.ch;
josep.majs.massanacodina@agroscope.admin.ch;

Abstract

Potato late blight (PLB), due to the pathogenic oomycete, *Phytophthora infestans*, can cause extensive economic damage, particularly in organic potato production. Copper is used to combat PLB, but its reduction as a plant protection product is an increasing priority in Europe. Alternative control strategies, including botanicals, are a potential avenue to reduce copper and control PLB. We investigated the application of *Frangula alnus* bark, its pairing with a reduced copper application, and a reduced copper application alone in field and lab experiments. The effect of different dosages and preparations on efficacy, and the posited active ingredients were examined. *Frangula alnus* treatments decreased disease severity compared to the untreated control but showed differences in efficacy depending on dosage and disease pressure. No difference in disease severity between the full and reduced copper treatments was observed. Through in vitro and in planta experiments, we investigated whether *F. alnus* directly or indirectly controlled PLB. A bacterium (*Erwinia* spp.), originating from the *F. alnus* extract, colonized the media and accounted for most of the direct inhibition in vitro, but removing microorganisms through filtration had no effect on the extract's efficacy in planta. It is unclear whether extract-associated microorganisms contributed to PLB control and would require additional experimentation to assess. The presence of measured anthraquinones likely contributed to the effect of *F. alnus*. Yields from the field experiments within each year remained stable despite different treatments and dosages, suggesting that treatment effectiveness is intertwined with the timing of disease development and ambient environmental conditions.

Keywords: *Phytophthora infestans*, potato late blight, alternative control strategies, *Frangula alnus*, plant protection, oomycete, *Erwinia*

1 Introduction

Organic potato production relies on the use of copper to combat potato late blight (PLB) caused by the oomycete, *Phytophthora infestans* (Torre et al, 2018). Copper, however, is a heavy metal that has long-term consequences for soil health and toxicity (Shabbir et al, 2020). In the last decades, copper reduction in organic potato production has become a priority (Tamm et al, 2022), and much research has been focused on alternative plant protection products, including the application of microorganisms or natural products (Bangemann et al, 2014; Dorn et al, 2007). PLB is a polycyclic pathogen that typically requires multiple fungicide treatments within a growing season to control it, resulting in potato organic production receiving one of the highest fungicide and copper applications of all arable crops (Katsoulas et al, 2020; Andrivon and Savini, 2023). Therefore, the reduction of copper is tightly intertwined with the development of copper alternatives to control PLB.

Copper applications can also be reduced through the deployment of resistance varieties, but potato breeding requires a long-time horizon of at least 10 to 16 years (Keijzer et al, 2022; Pandey et al, 2023), which could be shortened by 3 years with the implementation of marker-assisted selection and genomic-estimated breeding values of complex traits (Slater et al, 2014). However, previous potato breeding efforts show that it is difficult to combine late blight resistance, tuber yield, starch content, and other characteristics (Reslow et al, 2022), and therefore, limited gains from breeding programs in Europe (Ortiz et al, 2022). Furthermore, the use of a single resistance gene is not a sustainable strategy, since *P. infestans* strains can evolve to overcome qualitative resistance (Gilroy et al, 2011). Genome-editing techniques could facilitate the development of resistant varieties, but the public, particularly organic producers and consumers, are very skeptical of genetically modified organisms' use in potato breeding (Pacifico and Paris, 2016). Therefore, adopting lower copper application levels and pairing them with an alternative product can help reduce copper application levels on a shorter-time scale.

Previous experiments with botanicals demonstrate efficacy against oomycetes (Forrer et al, 2017; Mulholland et al, 2017; Taillis et al, 2022; Dagostin et al, 2010), and buckthorn alder bark (*Frangula alnus*) is among the botanicals that show potential to combat potato late blight (Forrer et al, 2017). However, further optimizing the preparation and dosage are needed to enable the economic feasibility of the treatment.

The goal of this work was to determine the efficacy of *Frangula alnus*, gain information about its accumulative effects, evaluate its pairing with reduced copper treatments, and optimize its usage. In order to determine the efficacy of different *Frangula alnus* preparations in the field, we carried out field experiments over four growing seasons to test various preparations and dosages. In addition, we conducted lab experiments, including in planta and in vitro assays and analytical chemistry to gain more information about the nature of the active compounds, treatment dosage and timing, and applicability to practice. Overall, this work highlights the possibility of further reducing copper allowances and the application of sequentially combined treatments.

2 Methods

2.1 Field experiment set-up

From 2019 to 2022, four potato field experiments were conducted at Agroscope in Zurich Reckenholz, Switzerland. A split plot design was used in the experimental fields with six replications in 2019 and 2020 and five replications in 2021 and 2022. Each plot included two sub-plots: one sub-plot with two rows of the variety Agria and another sub-plot with two rows of the variety Victoria. Agria and Victoria are common varieties used in Swiss potato production, including organic production (Swisspatat, 2023). These varieties are classified as moderately susceptible to late blight according to the recommended Swiss Variety List of potatoes (Schwärzel et al, 2019). Varieties with moderate susceptibility are recommended for the evaluation of low-risk products as they are expected to be less effective than higher-risk products (Evenhuis, 2021). Each sub-plot was 7.5 m^2 and contained two rows that were each 75 cm wide and 5 m long. In each row, the 15 planted tubers were spaced 33 cm apart. The placement of Agria and Victoria subplots within the plots was randomly assigned. In the four rows between the plots, two rows of a low susceptible variety (Panda in 2019 and 2020, Jelly in 2021, and Innovator in 2022) were planted along the long plot edge. The highly susceptible variety, Bintje, was planted between the two rows of the low susceptible variety (see Supplementary Figure A1 for more information on field layout). The low susceptible variety was changed in different years due to availability. Treatments were randomly assigned to plots within each block.

For all field experiments, certified seeds were used, and the fields were managed according to the EPPO guidelines (EPPO, 2021) for conducting fungicide efficiency experiments against *P. infestans* on potatoes. The only deviations from these guidelines were regarding plot length and variety choice. Although a longer plot (8 m) is prescribed for yield evaluations, the shorter acceptable plot length (5 m) was used because of field size constraints. Additionally, varieties used in this study are considered moderately susceptible to PLB to better mimic conditions in organic potato production (Schwärzel et al, 2019). In all of the years except 2022, PLB occurred naturally in the field experiments. In 2022, untreated, infected Bintje leaves that had been attacked by a natural infection were laid in the Bintje spreader rows to promote the infection within the experiment. Haulms of Bintje spreader rows were destroyed when 5-15% of the leaves were diseased. More details on the experimental set-up and agronomic conditions can be found in the supplementary material (Supplementary Tables A1, A2, and Figure A2).

2.1.1 Treatments

All experiments contained an untreated (water) control, a copper full dose (4 kg/ha/year) treatment, a copper reduced dose (2 kg/ha/year) treatment, and one or more *F. alnus* treatments consisting of different dosages or preparations. The number of treatments increased across the years as more information about the dosages and extraction methods became available through laboratory experiments. Starting from the field experiment's second season, treatments with a reduced *F. alnus* dosage were added. Additionally, an *F. alnus* treatment with a lypholization step during its extraction (see section below for details on preparation) was included in the third and fourth season of the field experiment. For copper treatments, Kocide Opti$^®$ (Bayer Agrar Schweiz AG, Basel, Switzerland) was used, which contains copper hydroxide in the form of a water dispersible granule. The untreated control was sprayed with the same amount of water as the treated plots during the treatment applications. To treat all plots equally, plots that received the sequentially mixed treatment (treatment number 9, Table 1) were sprayed with water for the first four application dates before switching to the reduced copper dose.

Table 1: Treatments and dosages applied for potato field experiments conducted in the years 2019 - 2022. Dosages below show amount sprayed cumulatively (per ha) for all 10 sprayings. Equal dosages of active ingredients from the treatments were applied during each spraying. Each treatment was mixed with tap water based on a total spray volume of 450 L/ha for the first four applications. To account for the growing canopy, the amount of water was increased for subsequent applications to 700 L/ha in 2019-2021 and 630 L/ha in 2022.

Years	Treatment Number	Treatment	Amount per ha/year
2019-2022	1	Untreated control (water)	NA
2019-2022	2	Copper full dose	4 kg
2019-2022	3	Copper reduced dose	2 kg
2019-2021	4	*Frangula alnus* full dose	400 kg
2020-2022	5	*Frangula alnus* medium dose	250 kg
2021-2022	6	*Frangula alnus* low dose	25 kg
2021-2022	7	*Frangula alnus* low dose (FD)	5 kg[1]
2021-2022	8	4x *Frangula alnus* low dose, followed by 6x reduced dose copper	25 kg (*F. alnus*)/2 kg (copper)
2022	9	4x no treatment (water), followed by 6x reduced dose copper	2 kg

[1]The 5 kg of extracted freeze-dried material is the product of 25 kg of *F. alnus* ground bark that was used as a starting material. Therefore, the amount of original material is the same as the low dose treatment (treatment 6).

2.1.2 Preparation of *Frangula alnus* solution

For treatments 4, 5 and 6 (Table 1), dried and milled bark of *F. alnus* (Faulbaumrinde PhEur, Dixa AG, St. Gallen, Switzerland, country of origin Poland) was suspended in tap water and stirred at room temperature (see Supplementary Table A2 for water volumes). In 2019 and 2020, the *F. alnus* solution was stirred for 2 hours as previously published (Forrer et al, 2017). The *F. alnus* treatments used 2021 and 2022 were prepared with a reduced stirring time of 30 minutes because a laboratory experiment revealed that the efficacy was unchanged between stirring times (see section below on extraction procedure experiments for details). After stirring, the solution was filtered through a 0.2 mm cheesecloth and then directly used for the treatments in the field. For treatment 7, the extraction method of *F. alnus* was modified according to a previously described procedure (Massana Codina, 2020). In short, powder-ground bark of *Frangula alnus* was extracted at 20% w/v using aqueous extraction in nanopure water by stirring for three hours at room temperature. The resulting extract was then centrifuged (4000 rpm, 10 min, 20 °C) and the supernatant was filtered through a ⌀ 9 cm 589/3 cellulose filter (Schleicher & Schuell, Dassel, Germany) using a vacuum. The aqueous solution was then freeze-dried. This modified procedure will be referred to as the low dose, freeze-dried (FD) treatment (Table 1). The fine powder obtained was stored in a sealed box at 10 °C in darkness. Before field applications, the amount of the fine powder that corresponds to the original bark weight of the low dose treatment (i.e. 20% of the original weight) was suspended in tap water and stirred for 30 minutes in the same manner as the *F. alnus* treatments. All treatments were then directly used in the field following their preparation.

2.1.3 Spray application

Dosages during applications were consistent throughout the season (Table 1), although spraying volume increased to accommodate the growing plants (Supplementary Table A2). The first application was carried out according to the recommendation of the Swiss decision support system PhytoPRE (www.phytopre.ch) (Cao et al, 1997; Steenblock and Forrer, 2002). The system considers late blight occurrence (date, location and distance), growth stage of the crop as well as weather conditions favorable for the development of *P. infestans*. All subsequent applications were performed in a weekly routine and were generally applied in the morning unless rain postponed the spraying times. The applications were shifted according to the weather forecast to try to avoid application on days with rainfall, and treatments were aimed to be applied one day in advance when rain was forecasted (see Supplementary Figure A2 for meteorological conditions during each season and on treatment days). For some applications, spraying on days with rainfall was unavoidable. For those occurrences, it was attempted to postpone the treatment until after precipitation or ensured that the spray coating could dry before rainfall. A total of ten applications was made each year (see Supplementary Table A2).

In 2019 and 2020, the applications were conducted with a plot sprayer (Schachtner Gerätetechnik, Ludwigsburg, Germany) equipped with a 1.5 m spray-boom and three IDK compact air induction nozzles (IDK yellow, IDK 90 -02 C, Lechler GmbH, Metzingen, Germany) at 2.2 bar. In 2021 and 2022, the treatments were applied with a tractor (Fendt® F 345 GT, AGCO GmbH) with a 3 m spray-boom and 6 IDK nozzles (IDK red, IDK 120-04 C). For all years, a spraying volume of 450 L/ha was used for the first four applications. To cover the entire plant canopy, the 6 top down nozzles were complemented with 5 dropleg nozzles (Lechler tongue nozzles, FT 140-02) as soon as the plants reached BBCH 31-35 corresponding to a switch to the larger spraying volume (630 L/ha in all years except for 2022 when 700 L/ha). In 2020, because of extremely wet soil conditions, the fifth treatment was applied with a knapsack sprayer (Birchmeier, Stetten, Switzerland) outfitted with a 3 m spray-

boom and 6 IDK nozzles (IDK brown, IDK 120-05 C, 1.9 bar). The spraying pressure was adjusted according to the equipment used.

2.1.4 Data collection from field experiments

The infection severity was estimated based on the area of infected plant tissue within each variety sub-plot (Forbes et al, 2014). Observations were made every three to seven days depending on the progress of the disease. In the earlier stages of the experiment, more frequent observations were performed to better assess the early development of the disease. Potatoes were harvested and yield was calculated from the two internal rows of each plot. The potatoes were sorted into four size classes (<42 mm, 42-55 mm, 55-70 mm, >70 mm). The middle two size classes (42-55 mm and 55-70 mm) constitute the marketable potatoes and were weighed to calculate the marketable yield. During the harvest, the tubers were examined for tuber blight. Additionally, 40-50 tubers per treatment were then examined for signs of tuber blight one month after harvest. For plots in which fewer than 50 tubers were harvested, all tubers were examined (min. per sub-plot was 40 tubers).

2.2 Climate chamber and laboratory experiments

Laboratory experiments were concurrently conducted with field experiment to better understand the extraction procedure, dosages, spraying time and mechanism of *P. infestans* control with *F. alnus*. These experiments consisted of detached leaf assays and mycelium growth experiments as well as the chemical analyses of the *F. alnus* solutions.

2.2.1 Extraction procedure and application timing experiment on detached leaves

Detached leaves of the susceptible variety Bintje were used to compare the efficacy of *F. alnus* after 30 versus 120 minutes of extraction. A 4% concentration of *F. alnus* was prepared in the same manner as the field experiment explained above. The aqueous solution of *F. alnus* was stirred for 30 minutes in addition to the treatment stirred for 120 minutes.

Within this experiment, the timing of the *F. alnus* application in relation to *P. infestans* inoculation was also tested. The three treatment levels consisted of applications either: 1) one day before *P. infestans* inoculation 2) two days before *P. infestans* inoculation, or 3) both two days and one day before *P. infestans* inoculation. A control treatment composed of deionized water was included, and it was sprayed according to the same regime. A gravity feed spray gun with cup (DeVilbiss, England) was used to apply 3 mL of the treatments to each terminal leaf at a pressure of 1 bar. All treatments included eight detached leaves that were each placed separately on a 7 x 7 cm square of wire mesh that had been laid over a moistened filter paper in individual round dishes with a diameter of 125 mm and height of 34 mm (Semadeni AG, Switzerland). The dishes were closed with a lid and were set at a slight angle to allow for the water to collect at the bottom, and they were randomly distributed in a blocked design within a climate chamber. The experiment was run at 18 °C and a 16-8 hr day-night schedule of which 2 hours consisted of a gradual reduction or increase of light. During the night period, the humidity was 85%, and 75% during the day.

A *P. infestans* polyspore isolate (polyspore isolate no: 18-001) collected in 2018 from untreated Bintje leaves in Reckenholz, Zurich was used for this and all laboratory and climate chamber experiments presented in this study. Its SSR genotype (Martin et al, 2019) had been categorized to "Other" (David Cooke, *Pers. comm.*), and it is an A1 mating type. The sporangia inoculation solution prepared by scraping mycelia from a two week old rye agar plate and diluting the sample to a sporangia

concentration of 5 x 10^5/mL. The spore preparation was incubated at 4 °C for two hours in the dark prior to inoculation to promote the release of zoospores. Each detached leaf consisted of the terminal leaflet and two first primary leaflets. The terminal leaflets were inoculated with *P. infestans* two times while each primary leaflet was inoculated once, totaling four inoculation points across the three leaflets (See supplementary Figure A3). A volume of 30 μL of the *P. infestans* sporangia preparation was used for each inoculation point. The humidity within the climate chamber was increased to 100% for 48 hours following the *P. infestans* inoculation, during which the climate chamber was kept dark. After the 48 hour period, the prevous day-night (16-8 hr) schedule and humidity settings (day: 85% and night: 75%) were reinstated. The percentage of leaf area that was diseased with *P. infestans* was assessed seven days post infection.

2.2.2 *Frangula alnus* dosage experiment

Detached leaf assays were carried out in the same manner as described above to test different application amounts of *F. alnus* and to determine its half-maximal inhibitory concentration (IC50). Previous studies had used 4% *F. alnus* extracts, which corresponded to approximately 400 kg/ha/year (Forrer et al, 2017; Krebs et al, 2006). In order to obtain a more economically feasible *F. alnus* treatment, reduced *F. alnus* treatment amounts were tested in laboratory experiments to assess the efficacy of lower application amounts in a controlled environment. A five-point dilution series of *F. alnus* was used for the treatment amounts, ranging from 2500 kg/ha/year to 0.25 kg/ha/year. For this experiment, detached leaves of the variety Bintje were used. To obtain the treatment with the highest concentration, a 4% *F. alnus* bark was extracted in tap water with stirring for 30 min, and then strained with a 0.2 mm cheesecloth. From this concentration, the subsequent four treatments were serially diluted using tap water. The detached leaves were sprayed as described above with 3 mL of the treatment per leaf at 1 bar until the treatment was well distributed on both sides of the leaf. For the untreated control, tap water was applied to the detached leaves.

Each treatment included 12 detached leaves that were randomly distributed according to a block design within the climate chamber. The climate chamber conditions and *P. infestans* strain used were the same as those used for the extraction procedure and spray timing experiment. The leaves were sprayed two days prior to artificial infection with *P. infestans*, which was applied as 4 drops of 30 μL of sporangia solution, including two drops on the terminal leaf and one drop on each of the first two primary leaflets. The sporangia preparation was prepared in the same manner as in the experiment described above, and the sporangia concentration used was 1 x 10^5/mL. The disease severity was assessed by estimating the surface area of *P. infestans* infection at seven days post infection.

2.2.3 *In vitro* mycelial growth experiment

In this experiment, *F. alnus*'s direct control on *P. infestans* mycelial growth was tested in vitro applying the same treatments used in the field (treatment numbers 1-7, Table 1) on rye agar (200 g untreated rye seed boiled and strained with a normal kitchen strainer, 5 g D-glucose, 20 g agar in 1000 mL water). *Frangula alnus* powder was prepared in the same manner as described above with a 30 min extraction step, but the tap water was autoclaved prior to the aqueous extraction step. Due to observations that a characteristic bacterial culture would grow on rye agar from *F. alnus* extractions, each of the *F. alnus* treatments were applied with and without an additional filtration step to eliminate the direct influence of microorganisms associated with the plant extracts. The bacterium was later identified to be an *Erwinia* spp. from sequencing the 16S rRNA gene (see Supplementary Material for details). Subsequent experiments revealed that the *Erwinia* bacterium could be cultured from multiple

batches of *F. alnus*. Three batches were tested, and all three yielded an *Erwinia* spp. bacterium (see Supplementary Material for additional information). The extent of the bacterium's co-occurrence with *F. alnus* bark is not known.

For the filtered solutions, two supplementary filtration steps were made that included 1) a vacuum filtration step using a paper filter (Schleicher & Schuell AG, 8714 Feldbach ZH) and 2) a manual filtration step using a 20 mL Luer-Lok syringe (BD Plastipak, Becton Dickinson GmbH, Germany) equipped with a 0.2 μm filter (Minisart®, Sartorius stedim biotech, Germany). Following filtration, the solution was plated to verify that the bacteria that grew on non-filtered *F. alnus* had been removed.

A ø 5 mm *P. infestans* mycelial plug from a freshly cultivated colony (within 10-14 days) was placed upside down in the center of each Petri dish (ø 9 cm, Greiner Bio-one, Switzerland). Six concentric wells were constructed 15 mm from the dishes' center and were filled with 25 μL of the test substance, of which only one was used on each Petri dish. Each test substance was repeated in 8 dishes. Any dishes that showed contamination were later excluded from the analysis with the exception of the *Erwinia* spp. that grew in the treatments of the non-filtered aqueous extractions of *F. alnus*. A second iteration was run with a subset of the treatments, including the untreated control, Copper (full), Frangula low, Frangula low (filtered). After seven days, the mycelial growth of *P. infestans* from the original plug was quantified as the percentage of mycelial coverage on each dish. For the measurements, the whole plate area was quantified with Image J (Schneider et al, 2012), and a modified ImageJ macro (Laflamme et al, 2016) was used to measure the area of a defined color threshold in pixels (See Supplementary Material). The experiments were normalized based on a comparison between each treatment's mycelial growth to its untreated control within each experiment and then combined for statistical analysis. The difference between the mycelial coverage in the treatment and the untreated control was used to determine the treatments' growth inhibition.

2.2.4 Detached leaf assay with filtered and non-filtered *Frangula alnus* extracts

Because bacterial growth was observed following plating *F. alnus* solutions on rye agar (see the in vitro experiment described above and Supplemental Material for more information), detached leaf assays were run to determine whether filtered and non-filtered *F. alnus* extracts differed in their efficacy in controlling *P. infestans*. Only the high dosage of *F. alnus* from the field experiment was included in the experiment with a positive control (copper) and negative control (autoclaved tap water). A filtered bacterial free extraction of high dosage *F. alnus* was prepared as described in the in vitro experiment, and the same protocols for the detached leaf set-up was followed as described in the detached leaf experiments focusing on dosage and *F. alnus* extraction and application timing. The assay was conducted two times with replicates of ten leaves per treatment in each assay.

Leaves of the variety Bintje were placed individually in a Petri dish (125 mm, Semadeni AG, Switzerland) on a 7x7 plastic mesh lined with paper towels, and 8 mL of tap water was added to the bottom of each Petri dish to keep the leaves fresh. The upper side of the leaf was sprayed with 1.5 mL of the prepared treatments at 1 bar and incubated in a climate chamber at 18 °C with 14 h day light and 75% humidity followed by 10 h darkness and 85% humidity. One day after treatment application, the leaves were infected at four points with a 30 μL *P. infestans* sporangia suspension. The spore suspension was prepared as described above with a final concentration of 1.60 x 10^5 sporangia per mL for the first assay (Assay A) and 1.75 x 10^5 sporangia per mL for the second assay (Assay B). One week post inoculation, the percentage of the infected leaf area was visually assessed as described above.

2.3 Chemical profiling and anthraquinone quantification from *Frangula alnus* bark

In order to investigate the chemical profile and determine the putative active compounds that may be involved in the *F. alnus* extracts' control of *P. infestans* in the field, freshly prepared extracts used from field preparations were immediately frozen at -20 °C for later analysis. In total, five preparations of the medium dosage, low dosage and low-FD dosage from two applications in 2021 and three applications in 2022 were analyzed. Additionally, the preparations from the aqueous extraction time experiment were profiled.

All preparations were thawed, filtered through a 22µm filter and directly injected on a Vanquish Horizon UHPLC System consisting of a degasser, a mixing pump, an autosampler, a diode array detector (PDA) and a Vanquish Charged Aerosol detector (ThermoFisher, Waltham, MA, USA). Samples were separated on an Atlantis BEH C18 column (2.1 x 100 mm, 1.7µm, Waters, Milford, MA, USA) using a mobile phase consisting of nanopure water containing 0.1% formic acid (A) and acetonitrile containing 0.1% formic acid (B); to obtain a chromatographic profile of the extract, a separation was performed using a step gradient of 5% to 40% B in 30 minutes, from 40% to 100% in 5 minutes, and isocratic at 100% for 3 minutes at a flow rate of 0.3 ml/min. In order to quantify frangulins and emodin, the separation was performed using a step gradient from 15% to 30% B in 4 minutes, isocratic at 30% B for 13 minutes, from 40 to 100% in 10 minutes and held at 100% B for 4 minutes at a flow rate of 0.3 ml/min. Detection was performed at 210, 254, 280, 350 and 435 nm, and the CAD conditions were the following: evaporator temperature set at 35 °C and the Data Collection Rate set at 10 Hz.

2.4 Statistical analysis

All data analyses were conducted in R (version 4.4.0) (R Core Team, 2023). Area under the disease progress curve (AUDPC) was calculated using the R package agricolae (de Mendiburu, 2023). For field experiments, disease severity based on the AUDPC and marketable yield were compared separately for each year using linear mixed models with the experimental blocks as a random factor and variety, treatment, and their interaction as fixed factors using the R package nlme (Pinheiro et al, 2023). The residual values were checked for normality using a Q-Q plot. Pairwise comparisons between treatments were performed using the estimated marginal means, and Tukey's honest significant difference tests were used to adjust p-values in the emmeans package (Lenth et al, 2024). The compact letter display function (cld) in R package multcomp was used to display differences between groups (Hothorn et al, 2008). The measurements of compounds extracted from the *F. alnus* preparations were compared using linear mixed models as above. However, the data were logarithmically transformed prior to analysis to account for the deviation from the model's assumptions of homoscedascity and normality.

The climate chamber experiment on extraction duration and spraying regime was analyzed with Kruskal-Wallis Rank Sum Test to explore differences between *F. alnus* and control treatments as well as pair-wise differences within the experiment. In order to assess the effects of the *F. alnus*'s extraction time, treatment application regime, and their interaction, they were included in a zero-inflated negative binomial model using the glmmTMB package (Brooks et al, 2017) with experimental blocks as random factors. All other laboratory experiments, including the detached leaf tests focusing on effects of *F. alnus* dosage and *F. alnus* filtration as well as the mycelia growth inhibition tests were analyzed with Kruskal-Wallis tests, followed by Conover's all-pairs rank comparison tests using the R package PMCMRplus (Pohlert, 2018) with the Benjamini & Hochberg adjustment for multiple comparisons (Benjamini and Hochberg, 1995).

3 Results

3.0.1 Field Experiments

The 2021 potato growing season was the most conducive for disease development followed by 2020. Accordingly, disease severity was highest in 2021, followed by 2020 and 2019, and in 2022, there was very little PLB (Figure 1). In all years, the treatment had a significant effect on the AUDPC, while variety had a significant effect on the AUDPC in the years 2019 and 2021, and no interaction between the treatments and variety was found in any of the years (Table 2). The *F. alnus* treatments showed significantly reduced AUDPCs compared to the untreated control (Figure 2) except in 2021 when high amounts of precipitation led to increased disease pressure (Supplementary Figure A2, Supplementary Table A1). In the field, the reduced copper was not found to be significantly different from the full copper treatment. The marketable yield showed significant variety effects in all four years, but only significant treatment effects in 2019 and 2020. In 2019, the Victoria plots treated with *F. alnus* had a yield reduction of 15% compared to the copper treatments (Figure 3). In 2021, the full copper treatments on the Agria plants resulted in 16% more yield than the *F. alnus* treatments (Figure 3). Treatment and variety interactions did not have a significant effect on marketable yield (Table 2).

At the time of harvest, tubers infected with late blight were found only in 2021 (0 to 4 infected tubers per treatment and variety were found in the 240-250 tubers analyzed per treatment and variety). There was no difference in the number of infected tubers across the treatments ($\chi^2 = 10.88$, df = 7, p-value = 0.144) or varieties ($\chi^2 = 2.15$, df = 1, p-value = 0.142). In all years, no tuber blight was found in the 3941 tubers examined one month after harvest from all treatments.

Fig. 1: Disease severity of potato late blight epidemics in two varieties from field experiments between 2019 and 2022. Treatments are described in Table 1. The y-axis scale changes according to the year, showing the high variation in disease severity across years. Bars indicate standard error of the means.

Table 2: ANOVA table of the linear mixed-effects models of experimental treatment, variety and their interaction on disease pressure and marketable yield from field trials between 2019 and 2022.

Response variable	Year	Factor	Df (df1, df2)	F-value	p-value
AUDPC[1]	2019	Treatment	3, 35	140.821	< 0.001
		Variety	1, 35	7.334	0.010
		Treatment x Variety	3, 35	1.469	0.239
	2020	Treatment	4, 45	63.427	< 0.001
		Variety	1, 45	0.758	0.389
		Treatment x Variety	4, 45	0.686	0.605
	2021	Treatment	7, 60	40.441	< 0.001
		Variety	1, 60	3.461	0.068
		Treatment x Variety	7, 60	0.3428	0.931
	2022	Treatment	7,60	28.680	< 0.001
		Variety	1, 60	7.080	0.010
		Treatment x Variety	7, 60	0.832	0.565
Marketable yield	2019	Treatment	3, 35	4.781	0.007
		Variety	1, 35	6.533	0.015
		Treatment x Variety	3, 35	1.675	0.190
	2020	Treatment	4, 45	3.828	0.009
		Variety	1, 45	11.050	0.002
		Treatment x Variety	4, 45	0.707	0.592
	2021	Treatment	7, 60	0.534	0.805
		Variety	1, 60	65.779	< 0.001
		Treatment x Variety	7, 60	0.6341	0.726
	2022	Treatment	7, 60	1.471	0.195
		Variety	1, 60	11.541	0.001
		Treatment x Variety	7, 60	0.216	0.980

[1]Area under the disease progress curve

Fig. 2: Area under the disease progress curve (AUDPC) of potato late blight in two varieties from field experiments between 2019 and 2022, showing treatment by color as described in Table 1. The scale of the y-axis varies according to the year, reflecting the substantial variation in disease severity, particularly from 2021 to 2022. Treatments that do not share grouping letters were shown to be significantly different from each other. Bars indicate the standard error of the mean.

Fig. 3: Marketable yield (tons per ha) of tubers in two varieties (in size classes: 42-55 mm and 55-70 mm) from potato late blight field experiments between 2019 and 2022, showing treatment by color. Treatments are described in Table 1. Bars indicate the standard error of the mean.

3.0.2 *Frangula alnus* extraction duration experiment on detached leaves

The detached Bintje leaves that had been treated with *F. alnus* were significantly less infected with *P. infestans* compared to the untreated controls ($\chi^2 = 38.08$, df = 5, p-value < 0.001). There was no difference between *F. alnus* treatments when the extraction time was 30 min or 120 min based on pairwise comparisons (Figure 4). Between the *F. alnus* treatments within the experiment, neither the treatment timing (one day, two days, and both one and two days prior to inoculation) ($\chi^2 = 2.62$, df = 2, p-value < 0.270) nor the extraction time ($\chi^2 = 0.312$, df = 1, p-value < 0.576) affected the infection rate. There was also no interaction between treatment timing and the *F. alnus* extraction time ($\chi^2 = 2.44$, df = 2, p-value < 0.295). Furthermore, the UHPLC profile of the *F. alnus* treatments did not show major differences in extracted compounds between the different extraction times, although quantities of compounds extracted differed (see section below on the chemical analysis of *F. alnus* for more details). Because no difference was found in treatment efficacy based on extraction time, the extraction time for *F. alnus* treatments in the field experiments was reduced to 30 minutes starting in 2021.

Fig. 4: Disease severity on detached leaves when sprayed with *Frangula alnus* that had been extracted in an aqueous solution for 30 or 120 minutes compared to a control treatment of tap water. The treatments included spraying regimes consisting of one day before, two days before, and both one and two days before inoculation (dbi) of *Phytophthora infestans*. Bars indicate the standard error of the mean.

3.0.3 *F. alnus* dosage experiment on detached leaves

Based on detached leaf assays, the Kruskal-Wallis test revealed a difference among *F. alnus* extracts and resulting disease severity ($\chi^2 = 33.98$, df $= 5$, p-value < 0.001). Dosages less than 25 kg/ha (Figure 6) were significantly less effective then the dosages that were equal to or greater than 25 kg/ha. There were no significant differences in disease severity among the highest three dosages (25, 250 and 2500 kg/ha). The disease severity found in the treatments of the lowest two dosages (0.25 and 25 kg/ha) did not differ from the untreated control. The IC50, indicating the dosage at which *F. alnus* inhibits half of the *P. infestans* infection that it is maximally capable of, was found to be 12.53 kg/ha. At 25 kg/ha, 69.4% of the maximum inhibitory potential of *P. infestans* was reached based on this dosage experiment.

Fig. 5: Pictures from in vitro assays to assess treatment inhibition of *Phytophthora infestans* mycelial growth. Treatments shown include a) untreated control b) copper (full) dosage c) *Fragnula alnus* high dosage, non-filtered d) *F. alnus* high dosage, filtered e) *F. alnus* medium dosage, non-filtered f) *F. alnus* medium dosage, filtered g) *F. alnus* low dosage, non-filtered h) *F. aluns* low dosage, filtered. The bacterium associated with the *F. alnus* bark that colonized the plates in treatments shown in panels c) and e) was identified to be an *Erwinia* spp.

3.0.4 *In vitro* mycelial growth experiment

The mycelia of *P. infestans* showed significantly different levels of growth inhibition depending on the treatment ($\chi^2 = 88.35$, df = 10, p-value < 0.001). There were no differences in mycelial growth on plates that received the untreated control of autoclaved water, and the filtered *F. alnus* low treatment, and both filtered and non-filtered *F. alnus* freeze-dried (FD) treatments (Figure **??**). For the high, medium, and low *F. alnus* treatments that had been prepared by aqueous extraction, there was significantly greater mycelial inhibition compared to their respective filtered treatments of the same dosages. The non-filtered medium and high *F. alnus* treatments that resulted in the growth of the *Erwinia* spp. inhibited mycelial growth more than the filtered treatments of the same dosages (Figure 5). For filtered treatments prepared with an aqueous extraction, the low *F. alnus* dosage showed decreased efficacy compared to the medium and high *F. alnus* treatments. The full copper treatment showed an increase in mycelial inhibition compared to the reduced copper treatment but no difference from the medium and high non-filtered *F. alnus* treatment where bacterial colonization of the *Erwinia* spp. was observed.

Fig. 6: Disease severity of detached leaves when sprayed with different dosages of *Frangula alnus* compared to a control treatment of tap water shown with the standard error of the mean. The 250 kg/ha dosage corresponds to the medium dosage from the 2020-2022 field experiments while 25 kg/ha is the same dosage applied to the low *F. alnus* treatment in the field (Table 1).

3.0.5 Detached leaf assay with filtered *F. alnus*

A detached leaf assay (assay A) was run to test the difference between the filtered and non-filtered *F. alnus* treatments that had been prepared with an aqueous extraction method. The assay was then run a second time (Assay B) to determine the consistency of the results. There were differences among treatments in both assay A ($\chi^2 = 29.133$, df = 3, p-value = 0.001) and assay B ($\chi^2 = 35.648$, df = 3, p-value = 0.001). The untreated control showed significantly higher rates of disease severity in both

assays (Figure 7), while the copper treatment yielded a significantly lower disease severity than the *F. alnus* treatments in assay A. In assay B, copper and *F. alnus* treatments showed either low or no *P. infestans* infections, and the treatments were not different from each other.

Fig. 7: Two assays (Assay A and Assay B) were run on detached leaves in which *Frangula alnus* that had been filtered was compared to *F. alnus* that had not been filtered. The high dosage of *F. alnus* was used for this experiment. Copper was included as a positive control. Bars indicate the standard error of the mean.

3.0.6 Chemical analysis of *Frangula alnus* extracts

The Ultra High Performance Liquid Chromatography—charged aerosol detector (UHPLC-CAD) profile of the different dosages of the aqueous suspension of F. alnus (medium and low dosage) showed a similar chemical composition, differing only on the concentration of the extracted compounds. However, the FD-low dosage showed that the filtering and freeze-drying step concentrates the extract on polar compounds, thus reducing the relative quantity of semi-polar and apolar compounds (i.e. anthraquinones). Indeed, the quantification of the main constitutive anthraquinones differed among *F. alnus* dosage amounts (Figure 8, Frangulin A: $F_{(2,8)} = 299.190$, p < 0.001, Frangulin B: $F_{(2,8)} = 205.947$, p $<$ 0.001, Emodin: $F_{(2,8)} = 20.928$, p < 0.001). As expected, the medium dosage consistently yielded more anthraquinones than the low dosage. The medium dosage included 10X more ground *F. alnus* bark than the low dosage, but it proportionally yielded more Frangulin A and Frangulin B (17-18X more) than the low dosage, and proportionally less Emodin (only 5X more). The low dosages with the

lyophilization step (FD-Low) also yielded comparatively lower compounds than both aqueous extractions (2.5X less than the low dosage). Although the duration of the aqueous extraction did not change the efficacy of the *F. alnus* treatments (Figure 4), it seems as though the 30 min extraction yielded more Frangulin A and Frangulin B than the 120 min extraction but similar levels of Emodin (Figure A4). Based on two measurements of the *F. alnus* preparation following both extraction periods, the 30 min extraction yielded on average 4 times more Frangulin A (100.4 μg/mL versus 23.5 μg/mL) and almost 3 times more Frangulin B (52.7 μg/mL versus 18.2μg/mL) than the 120 min extraction.

Fig. 8: Chromatographic measurements of compounds extracted from *Frangula* preparations that were associated with potato late blight reductions. The medium dosage corresponds to 250 g/ha/year and the low dosage to 25 g/ha/year from the ground bark. The FD extraction included a lyophilization step while the other extractions were stirred at room temperature in tap water for 30 min. Bars indicate the standard error of the mean.

4 Discussion

Phytophthora infestans is difficult to control due to its ability to quickly spread and evolve resistance (Leesutthiphonchai et al, 2018). Copper is an effective control mechanism in organic farming (Torre et al, 2018), but due to residue persistence that can detrimentally affect soil organisms, governmental action plans are increasingly seeking avenues to reduce its use (Silva et al, 2022; Möhring et al, 2020). In 12 European countries, 56% of producers already use half of the allowable amount of copper (Tamm et al, 2022), showing that the reduction is feasible and could potentially be adopted more broadly.

Our research contributes additional evidence that shows using only half of the allowable copper is likely to result in comparable levels of disease control and potato yield that would result from the use of the full allowable amount in Switzerland (4 kg/ha/year). However, the timing of PLB development has a large impact on whether the yield is reduced (Wiik, 2014; González-Jiménez et al, 2023). In our experiments, the yield for both varieties, especially for Victoria, was noticeably reduced in 2021 compared to other years, which was likely due to the increased disease pressure that year. Although 2020 had greater disease pressure than 2019, the marketable yield did not vary among treatments of the same variety. The lack of treatment effect on yield and tuber infections, may have been due to the relatively late start of the epidemic, low disease pressure in three of the four years, or because the plot size of the experiment was not large enough to measure yield differences. Previous work focusing on copper reduction showed similar results, where copper reduction did not tend to affect yield, but rather yield was affected by year, site, and disease pressure (Bangemann et al, 2014).

Although the *F. alnus* treatments did not significantly increase the yield in the field experiments, they led to a significant decrease in disease severity compared to the untreated control in the years with low to moderate disease pressure (i.e. all years except 2021). In 2021, the PLB epidemic was particularly severe, and we found that the yield dropped within our experiment compared to the average yield of the other three years. However, compared to yields in 2019, 2020 and 2022, the Agria plots in our study had a lower yield reduction (35%) than the average yield reduction estimated across organic production (57%) in 2021 (Suisse, 2022). On the other hand, the yield in the Victoria plots in the present study were reduced by 63%. These varietal differences in yield reduction underscore the role that potato variety selection has in combating PLB's damage because the yield differences between varieties were much greater than any treatment differences in our study.

In 2020, both the *F. alnus* medium and high dosages were effective in reducing disease severity in the field. These results confirm the findings observed in former field studies (2010-2012) conducted by (Forrer et al, 2017), in which the control of potato late blight with a 4% *F. alnus* solution was equivalent to a copper treatment (eight times 0.3 kg/ha copper) and reduced disease severity based on the AUDPC by 37%. By 2022, our experiment no longer included the high dosage treatment due to the lack of economic feasibility. Instead a low dose, consisting of 25 kg/ha was included, which would be in the economic range of copper and fungicide treatments (430-580 CHF per season/ha). The low dosage treatment can inhibit 69.4% of the pathogen compared to the maximal inhibitory concentration of *F. alnus* based on the dosage experiment in this study. However, the inhibition is not effective at reducing disease severity (based on AUDPC) in a year with high disease pressure such as 2021. Because the *F. alnus* extraction has not undergone any subsequent formulation, its leaching may also have contributed to its inability to inhibit *P. infestans*. Nonetheless, all *F. alnus* dosages showed reduced disease severity based on the AUDPC in 2022, although it was a year with minimal PLB pressure. In 2021, only the treatments that contained copper showed a significant reduction in PLB, of which one treatment included a switch from low dosage *F. alnus* to copper on the fifth spraying application (treatment 8), resulting in overall less copper use.

The results from this mixed treatment, during which a *F. alnus* solution was first sprayed until disease pressure was elevated, followed by a switch to a low dose copper treatment mid-way through the season, may serve as an effective strategy to reduce overall copper application. Because the mixed treatments (treatments 8 and 9) represent a 70% reduction in copper from the allowable amount in Switzerland, delayed application of reduced dosages could be an effective means to reach governmental targets that aim to reduce copper application. Further work is needed to demonstrate whether the *F. alnus* gives an added benefit or copper's delayed application is sufficiently protective without any early *F. alnus* treatments. However, *F. alnus* clearly provides protection from *P. infestans* as demonstrated through the field studies in years of low to moderate PLB severity and reinforced by the multiple laboratory experiments presented in this work. Previous work has focused on mixed treatments that are sprayed at the same time (Liljeroth et al, 2010), and there is less information about sequentially mixed treatments.

The laboratory studies provided a controlled environment to gain further information about the mode of action and effect of different spraying regimes and application dosages. The in vitro work was performed to understand if *F. alnus* has a direct inhibition on *P. infestans* mycelial growth. When the non-filtered *F. alnus* aqueous extractions were plated, the *Erwinia* spp. associated with the *F. alnus* bark established on the plate and directly inhibited or competed with *P. infestans*. The bacterium did not grow from the *F. alnus* extracted through lyophilization (treatment 7, low dose, FD), the filtered extracts nor on the copper and water control plates. The effect of the bacterium originating from the *F. alnus* bark on *P. infestans* in the in vitro experiment also suggests that botanical applications may act not only through the chemical compounds from the plant, but depending on the extraction method, potentially also through their microbiota. The microbiota of plants can provide protection against pathogens (Vogel et al, 2021), and perhaps the application of botanical plant protection products could also transfer protective microbial constituents through targeted mycobiome transplants (Chock et al, 2021). However, it is unclear the extent to which the microbiota of *F. alnus* treatment actually establishes on the potato plant, and further work would need to directly assess the colonization ability of a treatment's microbiota. Additionally, it is also unclear how widespread *Erwinia* spp. co-occurs in *F. alnus* bark. Previous work, in which *F. alnus* extracts were plated mentioned no bacterial colonization (Forrer et al, 2017). Nonetheless, the possible transmission of microbiota could potentially have an influence on a botanical treatment's effect and should be considered in future studies.

In order to disentangle the inhibition from the bacterium and *F. alnus*'s chemical compounds, the extracts from all treatments were added to the in vitro assays after filtration to eliminate the bacterium or directly without filtration. These in vitro results suggest that the chemical compounds themselves provided minimal direct inhibition to *P. infestans* mycelial growth compared to the bacterium associated with the bark, which appears to provide most of the in vitro inhibition. These results corroborate previous findings that found that *F. alnus* itself only slightly inhibited mycelial growth (Forrer et al, 2017). However, it was found that *F. alnus* can inhibit sporangial germination of *P. infestans* (Forrer et al, 2017). Indeed, several anthraquinones and their analogues have been shown to possess antifungal activity against several phytopathogens (Kim et al, 2004; Barilli et al, 2022; Choi et al, 2004), and the presence of these compounds in the *F. alnus* extracts might account for decreased *P. infestans* through direct mechanism such as decreased sporangial germination and slightly decreased mycelial growth.

Detached leaves from potato plants were then used to see how filtration affects the *F. alnus* treatment's effect on plants. The high dosage was applied to detached leaves either directly with no filtration step or following filtration. Both *F. alnus* treatments, regardless of filtration, resulted in significantly reduced *P. infestans* disease severity. The filtered *F. alnus*, which had a much lower direct effect on *P. infestans* in vitro, showed as much efficacy as the non-filtered extraction in detached leaf assays,

pointing to the possible involvement of induced resistance (Forrer et al, 2017). Anthraquinones have also been shown to induce resistance in plants (Konstantinidou-Doltsinis and Schmit, 1998), and other chemical families present in the *F. alnus* extracts, such as polyphenols, have been implicated in defense induction (Gillmeister et al, 2019). In *Vitis vinifera*, the application of *F. alnus* bark extract induces the production of the phytoalexins piceid, resveratrol and pterostilbene, and protects against infection of the oomycete *Plasmopara viticola* (Godard et al, 2009). Plant elicitation can complement direct antifungal activity if deployed in the right situations to avoid reduced yields (Heil et al, 2000), and therefore genotypic and environmental variables should be evaluated before applying elicitors since these factors can influence their success (Walters et al, 2013). For example, previous research with *F. alnus* showed a varietal difference in yield (Forrer et al, 2017), which may be due to the way in which a potato genotype responds to an elicitor (Bruce, 2014)

Current obstacles to adoption include cost of the material, effort required for the preparations, and stability of the extraction. In order to further develop an effective and potentially cost-effective treatment, more information about the compounds involved could better elucidate the mode of action. Frangulin A, Frangulin B and Emodin were found in the *F. alnus* extract, but it is not clear which chemicals are responsible for the direct or indirect inhibition. Singularly testing each compound would help decipher how they individually and cumulatively control PLB. These steps could help determine whether a stable formulation can be achieved to obtain a cost-effective treatment. Therefore, many steps are still required (i.e. an understanding of kinetic degradation of the active compounds, stability in solution and on foliage, optimal formulations) to achieve a viable PLB spraying regime that incorporates *F. alnus* or its components.

Our work suggests that copper application amount could be substantially reduced and potentially supplemented with alternative products, such as *F. alnus*. Further testing of sequential deployments of different treatments (i.e. switch from an alternative treatment, such as a botanical or another plant elicitor to copper) can also help contribute to overall copper reduction. The success of such strategies would be dependent on the response of the variety to plant elicitation, and the timing of the epidemic may also very much impact their success. Reductions in copper application combined with the deployment of resistant varieties (Kessel et al, 2018) can help achieve goals of reduced pesticide targets (Silva et al, 2022; Purnhagen et al, 2021). Due to importance of potato production in Europe a combination of strategies are necessary to combat potato late blight.

Supplementary information. Detailed weather and agronomic information for all four years of the field experiment as well as additional information about sample and data analyses can be found in the supplement.

Author contributions. The study was conceived and designed by Tomke Musa, Karen Sullam, and Andreas Kägi. Material preparation, data collection and analysis were performed by Tomke Musa, Karen Sullam, Haruna Gütlin, Andreas Kägi, Sylvain Schnee and Josep Massana-Codina. Tomke Musa and Karen Sullam jointly wrote the first draft of the manuscript. All authors commented on previous versions of the manuscript and read and approved the final manuscript.

Acknowledgments. We thank Susanne Vogelgsang for the valuable support and constructive feedback. We also thank Damian Oswald, Nicole Togni, Lisa Besson, Arnaud Chotel, Anja Logo, Cecilia Panzetti, and Clara Chevalley for their assistance with laboratory and/or field experiments.

References

Andrivon D, Savini I (2023) Can organic agriculture cope without copper for disease control? Synthesis of the Collective Scientific Assessment report, èditions Quae, Versailles, https://doi.org/10.35690/978-2-7592-3156-0

Bangemann LW, Westphal A, Zwerger P, et al (2014) Copper reducing strategies for late blight (*Phytophthora infestans*) control in organic potato (*Solanum tuberosum*) production. Journal of Plant Diseases and Protection 121(3):105–116. https://doi.org/10.1007/BF03356498, URL https://doi.org/10.1007/BF03356498

Barilli E, Agudo FJ, Masi M, et al (2022) Anthraquinones and their analogues as potential biocontrol agents of rust and powdery mildew diseases of field crops. Pest Management Science 78(8):3489–3497. https://doi.org/https://doi.org/10.1002/ps.6989, URL https://scijournals.onlinelibrary.wiley.com/doi/abs/10.1002/ps.6989, https://scijournals.onlinelibrary.wiley.com/doi/pdf/10.1002/ps.6989

Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: A practical and powerful approach to multiple testing. Journal of the Royal Statistical Society Series B (Methodological) 57(1):289–300. URL http://www.jstor.org/stable/2346101

Brooks M, Kristensen K, van Benthem K, et al (2017) glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. The R Journal 9(2):378–400. https://doi.org/doi:10.32614/RJ-2017-066

Bruce TJA (2014) Variation in plant responsiveness to defense elicitors caused by genotype and environment. Frontiers in Plant Science 5. https://doi.org/10.3389/fpls.2014.00349, URL https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2014.00349

Cao K, Ruckstuhl M, Forrer H (1997) Crucial weather conditions for *Phytophthora infestans*: A reliable tool for improved control of potato late blight? In: Bouma E, Schepers H (eds) PAV-Special Report, 1:85-90

Chock MK, Hoyt BK, Amend AS (2021) Mycobiome transplant increases resistance to *Austropuccinia psidii* in an endangered hawaiian plant. Phytobiomes Journal 5(3):326–334. https://doi.org/10.1094/PBIOMES-09-20-0065-R, URL https://doi.org/10.1094/PBIOMES-09-20-0065-R, https://doi.org/10.1094/PBIOMES-09-20-0065-R

Choi GJ, Lee SW, Jang KS, et al (2004) Effects of chrysophanol, parietin, and nepodin of *Rumex crispus* on barley and cucumber powdery mildews. Crop Protection 23(12):1215–1221. https://doi.org/https://doi.org/10.1016/j.cropro.2004.05.005, URL https://www.sciencedirect.com/science/article/pii/S0261219404001371

Dagostin S, Formolo T, Giovannini O, et al (2010) Salvia officinalis extract can protect grapevine against *Plasmopara viticola*. Plant Disease 94(5):575–580. https://doi.org/10.1094/PDIS-94-5-0575, URL https://doi.org/10.1094/PDIS-94-5-0575, pMID: 30754462

de Mendiburu F (2023) agricolae: Statistical Procedures for Agricultural Research. URL https://CRAN.R-project.org/package=agricolae, r package version 1.3-6

Dorn B, Musa T, Krebs H, et al (2007) Control of late blight in organic potato production: evaluation of copper-free preparations under field, growth chamber and laboratory conditions. European Journal of Plant Pathology 119(2):217–240. https://doi.org/10.1007/s10658-007-9166-0, URL https://doi.org/10.1007/s10658-007-9166-0

EPPO (2021) Pp 1/002 (5) *Phytophthora infestans* on potato. EPPO Bulletin 51(1):79–82. https://doi.org/https://doi.org/10.1111/epp.12708, URL https://onlinelibrary.wiley.com/doi/abs/10.1111/epp.12708, https://onlinelibrary.wiley.com/doi/pdf/10.1111/epp.12708

Evenhuis A (2021) Efficacy to control potato late blight by applying biological crop protection products: EuroBlight field experiment AGV8005. No. WPR-OT 939 in Report Wageningen Plant Research, Wageningen Plant Research, Netherlands, https://doi.org/10.18174/570735

Forbes G, Perez W, Andrade-Piedra J (2014) Field assessment of resistance in potato to Phytophthora infestans: International cooperators guide. https://doi.org/10.4160/9789290604402

Forrer HR, Vogelgsang S, Musa T (2017) Botanicals and phosphonate show potential to replace copper for control of potato late blight. Journal of Fungi 3(4). https://doi.org/10.3390/jof3040065, URL https://www.mdpi.com/2309-608X/3/4/65

Gillmeister M, Ballert S, Raschke A, et al (2019) Polyphenols from rheum roots inhibit growth of fungal and oomycete phytopathogens and induce plant disease resistance. Plant Disease 103(7):1674–1684. https://doi.org/10.1094/PDIS-07-18-1168-RE, URL https://doi.org/10.1094/PDIS-07-18-1168-RE, pMID: 31095470, https://doi.org/10.1094/PDIS-07-18-1168-RE

Gilroy EM, Breen S, Whisson SC, et al (2011) Presence/absence, differential expression and sequence polymorphisms between piavr2 and piavr2-like in *Phytophthora infestans* determine virulence on r2 plants. New Phytologist 191(3):763–776. https://doi.org/https://doi.org/10.1111/j.1469-8137.2011.03736.x, URL https://nph.onlinelibrary.wiley.com/doi/abs/10.1111/j.1469-8137.2011.03736.x, https://nph.onlinelibrary.wiley.com/doi/pdf/10.1111/j.1469-8137.2011.03736.x

Godard S, Slacanin I, Viret O, et al (2009) Induction of defence mechanisms in grapevine leaves by emodin- and anthraquinone-rich plant extracts and their conferred resistance to downy mildew. Plant Physiology and Biochemistry 47(9):827–837. https://doi.org/https://doi.org/10.1016/j.plaphy.2009.04.003, URL https://www.sciencedirect.com/science/article/pii/S0981942809001077

González-Jiménez J, Andersson B, Wiik L, et al (2023) Modelling potato yield losses caused by *Phytophthora infestans*: Aspects of disease growth rate, infection time and temperature under climate change. Field Crops Research 299:108977. https://doi.org/https://doi.org/10.1016/j.fcr.2023.108977, URL https://www.sciencedirect.com/science/article/pii/S0378429023001703

Heil M, Hilpert A, Kaiser W, et al (2000) Reduced growth and seed set following chemical induction of pathogen defence: does systemic acquired resistance (sar) incur allocation costs? Journal of Ecology 88(4):645–654. https://doi.org/https://doi.org/10.1046/j.1365-2745.2000.00479.x, URL https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1046/j.1365-2745.2000.00479.x, https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1046/j.1365-2745.2000.00479.x

Heuer H, Krsek M, Baker P, et al (1997) Analysis of actinomycete communities by specific amplification of genes encoding 16s rrna and gel-electrophoretic separation in denaturing gradients. Applied and Environmental Microbiology 63(8):3233–3241. https://doi.org/10.1128/aem.63.8.3233-3241.1997, URL https://journals.asm.org/doi/abs/10.1128/aem.63.8.3233-3241.1997, https://journals.asm.org/doi/pdf/10.1128/aem.63.8.3233-3241.1997

Hothorn T, Bretz F, Westfall P (2008) Simultaneous inference in general parametric models. Biometrical Journal 50(3):346–363

Katsoulas N, Løes AK, Andrivon D, et al (2020) Current use of copper, mineral oils and sulphur for plant protection in organic horticultural crops across 10 european countries. Organic Agriculture 10(1):159–171. https://doi.org/10.1007/s13165-020-00330-2, URL https://doi.org/10.1007/s13165-020-00330-2

Keijzer P, van Bueren ETL, Engelen CJM, et al (2022) Breeding late blight resistant potatoes for organic farming—a collaborative model of participatory plant breeding: the bioimpuls project. Potato Research 65(2):349–377. https://doi.org/10.1007/s11540-021-09519-8, URL https://doi.org/10.1007/s11540-021-09519-8

Kessel GJ, Mullins E, Evenhuis A, et al (2018) Development and validation of IPM strategies for the cultivation of cisgenically modified late blight resistant potato. European Journal of Agronomy 96:146–155. https://doi.org/https://doi.org/10.1016/j.eja.2018.01.012, URL https://www.sciencedirect.com/science/article/pii/S1161030118300327

Kim JC, Choi GJ, Lee SW, et al (2004) Screening extracts of *Achyranthes japonica* and *Rumex crispus* for activity against various plant pathogenic fungi and control of powdery mildew. Pest Management Science 60(8):803–808. https://doi.org/https://doi.org/10.1002/ps.811, URL https://scijournals.onlinelibrary.wiley.com/doi/abs/10.1002/ps.811, https://scijournals.onlinelibrary.wiley.com/doi/pdf/10.1002/ps.811

Konstantinidou-Doltsinis S, Schmit A (1998) Impact of treatment with plant extracts from *Reynoutria sachalinensis* (f. schmidt) nakai on intensity of powdery mildew severity and yield in cucumber under high disease pressure. Crop Protection 17(8):649–656. https://doi.org/https://doi.org/10.1016/S0261-2194(98)00066-0, URL https://www.sciencedirect.com/science/article/pii/S0261219498000660

Krebs H, Dorn B, Forrer HR (2006) Control of late blight of potato with medicinal plant suspensions [in German]. Agrarforschung 13(1):16–21. URL https://www.agrarforschungschweiz.ch/2006/01/krautfaeulebekaempfung-mit-suspensionen-aus-heilpflanzen/

Laflamme B, Middleton M, Lo T, et al (2016) Image-based quantification of plant immunity and disease. Molecular Plant-Microbe Interactions® 29(12):919–924. https://doi.org/10.1094/MPMI-07-16-0129-TA, URL https://doi.org/10.1094/MPMI-07-16-0129-TA, pMID: 27996374, https://doi.org/10.1094/MPMI-07-16-0129-TA

Leesutthiphonchai W, Vu AL, Ah-Fong AMV, et al (2018) How does *Phytophthora infestans* evade control efforts? modern insight into the late blight disease. Phytopathology® 108(8):916–924. https:

//doi.org/10.1094/PHYTO-04-18-0130-IA, URL https://doi.org/10.1094/PHYTO-04-18-0130-IA, pMID: 29979126, https://doi.org/10.1094/PHYTO-04-18-0130-IA

Lenth R, et al (2024) emmeans: Estimated marginal means, aka least-squares means. R package version 1.10.1

Liljeroth E, Bengtsson T, Wiik L, et al (2010) Induced resistance in potato to *Phytphthora infestans*—effects of baba in greenhouse and field tests with different potato varieties. European Journal of Plant Pathology 127(2):171–183. https://doi.org/10.1007/s10658-010-9582-4, URL https://doi.org/10.1007/s10658-010-9582-4

Martin FN, Zhang Y, Cooke DEL, et al (2019) Insights into evolving global populations of *Phytophthora infestans* via new complementary mtDNA haplotype markers and nuclear ssrs. PLOS ONE 14(1):1–24. https://doi.org/10.1371/journal.pone.0208606, URL https://doi.org/10.1371/journal.pone.0208606

Massana Codina J (2020) Epidemiological insights, cultivar resistance and natural products to control black dot (*Colletotrichum coccodes*) and silver scurf (*Helminthosporium solani*) on potato (*Solanum tuberosum*) tubers. Phd dissertation, Université de Genève, https://doi.org/https://doi.org/10.13097/archive-ouverte/unige:146843

Mulholland DA, Thuerig B, Langat MK, et al (2017) Efficacy of extracts from eight economically important forestry species against grapevine downy mildew (*Plasmopara viticola*) and identification of active constituents. Crop Protection 102:104–109. https://doi.org/https://doi.org/10.1016/j.cropro.2017.08.018, URL https://www.sciencedirect.com/science/article/pii/S0261219417302491

Möhring N, Ingold K, Kudsk P, et al (2020) Pathways for advancing pesticide policies. Nature Food 1(9):535–540. https://doi.org/10.1038/s43016-020-00141-4, URL https://doi.org/10.1038/s43016-020-00141-4

Ortiz R, Reslow F, Cuevas J, et al (2022) Genetic gains in potato breeding as measured by field testing of cultivars released during the last 200 years in the nordic region of europe. The Journal of Agricultural Science 160(5):310–316. https://doi.org/10.1017/S002185962200034X

Pacifico D, Paris R (2016) Effect of organic potato farming on human and environmental health and benefits from new plant breeding techniques. is it only a matter of public acceptance? Sustainability 8(10). https://doi.org/10.3390/su8101054, URL https://www.mdpi.com/2071-1050/8/10/1054

Pandey J, Scheuring DC, Koym JW, et al (2023) Genomic selection and genome-wide association studies in tetraploid chipping potatoes. The Plant Genome 16(1):e20297. https://doi.org/https://doi.org/10.1002/tpg2.20297, URL https://acsess.onlinelibrary.wiley.com/doi/abs/10.1002/tpg2.20297, https://acsess.onlinelibrary.wiley.com/doi/pdf/10.1002/tpg2.20297

Pinheiro J, Bates D, R Core Team (2023) nlme: Linear and Nonlinear Mixed Effects Models. URL https://CRAN.R-project.org/package=nlme, r package version 3.1-163

Pohlert T (2018) PMCMRplus: Calculate Pairwise Multiple Comparisons of Mean Rank Sums Extended. URL https://CRAN.R-project.org/package=PMCMRplus

Purnhagen KP, Clemens S, Eriksson D, et al (2021) Europe's farm to fork strategy and its commitment to biotechnology and organic farming: Conflicting or complementary goals? Trends in Plant Science 26(6):600–606. https://doi.org/10.1016/j.tplants.2021.03.012, URL https://doi.org/10.1016/j.tplants.2021.03.012, doi: 10.1016/j.tplants.2021.03.012

R Core Team (2023) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria, URL https://www.R-project.org/

Reslow F, Carlson-Nilsson U, Crossa J, et al (2022) Public potato breeding progress for the nordic region of europe: evidence from multisite testing of selected breeding clones and available released cultivars. Acta Agriculturae Scandinavica, Section B — Soil & Plant Science 72(1):553–562. https://doi.org/10.1080/09064710.2021.2021279, URL https://doi.org/10.1080/09064710.2021.2021279, https://doi.org/10.1080/09064710.2021.2021279

Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. Nature Methods 9(7):671–675. https://doi.org/10.1038/nmeth.2089, URL https://doi.org/10.1038/nmeth.2089

Schwärzel R, Torche JM, Ballmer T, et al (2019) Swiss variety list for potatoes 2019 [in German]. Agroscope Transfer 9(11):1–6. URL https://link.ira.agroscope.ch/en-US/publication/40008

Shabbir Z, Sardar A, Shabbir A, et al (2020) Copper uptake, essentiality, toxicity, detoxification and risk assessment in soil-plant environment. Chemosphere 259:127436. https://doi.org/https://doi.org/10.1016/j.chemosphere.2020.127436, URL https://www.sciencedirect.com/science/article/pii/S0045653520316301

Silva V, Yang X, Fleskens L, et al (2022) Environmental and human health at risk – scenarios to achieve the farm to fork 50

Slater AT, Cogan NOI, Hayes BJ, et al (2014) Improving breeding efficiency in potato using molecular and quantitative genetics. Theoretical and Applied Genetics 127(11):2279–2292. https://doi.org/10.1007/s00122-014-2386-8, URL https://doi.org/10.1007/s00122-014-2386-8

Steenblock T, Forrer HR (2002) Phytopre+2000: a plot specific decision support system on the internet [in German]. Agrarforschung 9(5):207–214. URL https://www.agrarforschungschweiz.ch/2002/05/kartoffelanbau-krautfaeuleberatung-via-internet/

Suisse B (2022) Market reflections organic potatoes [in German]. URL https://www.bioaktuell.ch/fileadmin/documents/ba/Markt/Kartoffeln/2022_09_Marktspiegel_Kartoffeln.pdf, accessed 7 October 2024

Swisspatat (2023) Statistical information 2023: Potato cultivation and potato utilization [in German and French]. URL https://www.kartoffel.ch/_Resources/Persistent/c227210049cd7e4b19b602541e8e94c25cdf8381/Statistische_Angaben_2023_DE.pdf

Taillis D, Pébarthé-Courrouilh A, Lepeltier E, et al (2022) A grapevine by-product extract enriched in oligomerised stilbenes to control downy mildews: focus on its modes of action towards *Plasmopara viticola*. OENO One 56(3):55–68. https://doi.org/10.20870/oeno-one.2022.56.3.4911, URL https://

oeno-one.eu/article/view/4911

Tamm L, Thuerig B, Apostolov S, et al (2022) Use of copper-based fungicides in organic agriculture in twelve european countries. Agronomy 12(3). https://doi.org/10.3390/agronomy12030673, URL https://www.mdpi.com/2073-4395/12/3/673

Torre AL, Iovino V, Caradonia F (2018) Copper in plant protection: current situation and prospects. Phytopathologia Mediterranea 57(2):201–236. https://doi.org/10.14601/Phytopathol_Mediterr-23407, URL https://www.jstor.org/stable/26507086

Vogel CM, Potthoff DB, Schäfer M, et al (2021) Protective role of the arabidopsis leaf microbiota against a bacterial pathogen. Nature Microbiology 6(12):1537–1548. https://doi.org/10.1038/s41564-021-00997-7, URL https://doi.org/10.1038/s41564-021-00997-7

Walters DR, Ratsep J, Havis ND (2013) Controlling crop diseases using induced resistance: challenges for the future. Journal of Experimental Botany 64(5):1263–1280. https://doi.org/10.1093/jxb/ert026, URL https://doi.org/10.1093/jxb/ert026, https://academic.oup.com/jxb/article-pdf/64/5/1263/18044089/ert026.pdf

Wiik L (2014) Potato late blight and tuber yield: Results from 30 years of field trials. Potato Research 57(1):77–98. https://doi.org/10.1007/s11540-014-9256-2, URL https://doi.org/10.1007/s11540-014-9256-2

Appendix A   Supplementary Material

A.1   Agronomic and meteorological conditions

In order to clarify the conditions and set-up used in the field experiments, we provide the following supplementary information, including information on the plot set-up (Figure A1), weather during the field experiment (Figure A2), and agronomic conditions and treatment applications (Tables A1 and A2).

Fig. A1: Schematic diagram of the potato variety split-plot design used in the field experiment. Each sub-plot included 2 rows of one variety (Agria or Victoria) next to the other variety. These plots were sandwiched between a low susceptible variety (in black) and the high susceptible variety in gray. Each box represents a different row.

Fig. A2: Graphs of meteorological conditions during the field experiments between 2019 to 2022, showing the daily average temperature (°C) with blue lines (y-axis on the right side) and daily sum of precipitation (mm) with gray bar (y-axis on the left side). The dates during which treatments were applied are shown by the red point along the x-axis. Generally, the treatments were applied in the mornings unless rain postponed the treatment. The weather measurements were taken at the MeteoSwiss station Zurich/Affoltern.

A.2 Identification of bacterial isolated from *F. alnus* aqueous extracts

The bacterial isolate was directly amplified with the primers 27F and 1492R (Heuer et al, 1997). Reactions were run with Taq-&Go^{TM} Mastermix (MP Biomedicals, Heidelberg, Germany) in a reaction volume of 20 μL and primer concentration of 0.5 μM. Reaction conditions included an initial denaturation at 94 °C for 5 min followed by 32 cycles of 94 °C 30 sec, 55 °C for 1 min and 72 °C for 1.5 min and a final elongation step of 72°C for 5 min. Sanger sequencing was performed (Microsynth, Balgach, Switzerland) with both 27F and 1492R, and a contig was constructed using Geneious Prime$^{®}$ 2024.0.3. The sequence will be deposited in the NCBI nucleotide database (Accession #, to be provided).

A.3 Additional details on *in vitro* and detached leaf experiments

For detached leaf experiments, terminal leaves, consisting of three leaflets were used (See Figure A3). The chromatograms from the detached leaf experiment using 30 min and 120 min extraction times of *F. alnus* are also displayed (See Figure A4).

Table A1: Agronomic information about field and environmental conditions during the four years of field experiments included in this study.

Agronomic Information	2019	2020	2021	2022
Field site	Rümlang, Plot 153	Reckenholz, Plot 114A	Reckenholz Plot 119	Reckenholz Plot 159
Coordinates	47.438796, 8.524827	47.430933, 8.520306	47.428847, 8.515277	47.433979, 8.511500
Soil type	sandy loamy, very deep	clay, sandy loam	clay, clayey loam	clay, clayey loam
Pre-crop	buckwheat	canola	grass seed propagation	summer wheat, intercrop (green manure)
Soil management	12 August 2018 cultivator, 18 April 2019 harrowing	6 April 2020 harrowing	—	18 April 2022 harrowing
Fertilization	compost, total per ha: 68 N, 152 P, 315 K, 0 Mg	total per ha: 112 N, 0 P, 360 K, 26 Mg	total per ha: 106 N, 87.4 P, 120 K, 36 Mg	compost, Patentkali, Super Urea, total per ha: 83 N, 222 P, 425 K, 145 Mg
Herbicide treatment	7 May Bandur Boxer Sencor	17 April Bandur Boxer Sencor SC; 28 May Fusilade Max	30 April Stomp Aqua + Firebird Plus (some herbicide damage); 31 May Fusilade Max	10 May Boxer; Sencor; 30 May Fusilade Max
Irrigation	—	—	—	22 June, 24 June, 19 July
Control of Colorado beetle	24 June (Biscaya)	8 June (Biscaya)	17 June (Audienz), 10 July (Biscaya)	19 May (Audienz), 10 June (Audienz)
Planting	18 April	8 April	20 April	20 April
Emergence	8 May	1 May	21 May	10 May
BBCH15	20 May	11 May	26 May	20 May
Defoliation of Bintje spreader rows	—	6 July	12 July	—
Defoliation of test varieties	4 August	27 July	29 July	2 August
Harvest	29 August	19 August	18 August	23 August
Date yield measurement	16 September	21 August	29 August	21 August
First late blight reported in Switzerland	20 May	11 May	21 May	10 May
Location of first announced PLB in Switzerland	Schlatt (TG), 30 km distance	Gürbrü (BE)/Kerzers (FR), 110 km distance	Dorf (ZH), 23 km distance	Flaach (ZH), 19 km distance, Weite (SG), 109 km distance
First late blight in field experiment	11 June	24 June	6 July	30 June
Overall weather conditions during field season	wet	wet	very wet	dry
Sum of precipitation (mm) during entire spraying period	194.0	190.6	345.6	155.6
PLB Pressure in Switzerland	low	medium	high	very low

Table A2: Dates of treatment application, equipment used for the application, volume of water used for the treatment mixtures (dry ingredient amounts shown in Table 1), and the sum of precipitation (mm) between spraying dates as well as on application day in parentheses.

Year	Application date	Equipment	Spraying volume (L/ha)	Sum of precipitation between sprayings and on application day[1] (mm)
2019	27 May	plot sprayer	450	11.2
	4 June	plot sprayer	450	31.7
	11 June	plot sprayer	450	30.4 (13.2)
	17 June	plot sprayer	450	14.0
	25 June	drop leg[2]	630	0
	2 July	drop leg	630	54.4 (38.5)
	9 July	drop leg	630	16.8
	16 July	drop leg	630	3.1
	23 July	drop leg	630	17.0
	30 July	drop leg	630	15.3 (6.7)
2020	20 May	plot sprayer	450	12.6
	27 May	plot sprayer	450	0.6
	3 June	plot sprayer	450	58.5
	12 June	plot sprayer	450	34.7
	19 June	knapsack sprayer	630	Row 1.8 (1.8)
	24 June	tractor, drop leg	630	27.2
	1 July	tractor, drop leg	630	5 (1.2)
	7 July	tractor, drop leg	630	25.9
	14 July	tractor, drop leg	630	16.2
	21 July	tractor, drop leg	630	8.1
2021	2 June	tractor	450	13.6
	11 June	tractor	450	0
	17. June	tractor, drop leg	622	22.1
	23 June	tractor, drop leg	630	111 (36.2)
	3 July	tractor, drop leg	630	6.9 (2.7)
	6 July	tractor, drop leg	630	42.6 (11.3)
	12 July	tractor, drop leg	630	122
	20 July	tractor, drop leg	630	0
	23 July	tractor, drop leg	630	21.4
	27 July	tractor, drop leg	630	6.0 (1.0)
2022	23 May	tractor	450	6.2 (2.5)
	30 May	tractor	450	30.8
	8 June	tractor	450	9.3 (2.1)
	14 June	tractor, drop leg	630	4.5
	21 June	tractor, drop leg	630	27.6
	27 June	tractor, drop leg	630	40.2 (3.8)
	6 July	tractor, drop leg	630	0
	13 July	tractor, drop leg	630	0
	20 July	tractor, drop leg	630	20.9 (14.2)
	26 July	tractor, drop leg	630	16.1(1.2)

[1] Any precipitation on the application day is shown in parentheses

[2] The drop leg spraying equipment was pushed by hand in 2019 and by tractor in 2020-2022.

Fig. A3: Each terminal leaflet was inoculated twice, and each of the two adjacent leaflets were inoculated once. The four locations on which 30 μL of spore suspension were deposited for the *Phytophthora infestans* inoculations are indicated by the red markings in the picture. Leaves were placed on moistened paper towels and a wire rack in Petri dishes.

Fig. A4: CAD (A) and UV at 435 nm (B) chromatograms of the *Frangula alnus* bark aqueous extract after 30 minutes and 120 minutes of stirring. The numbers associated with peaks in the chromatograms correspond to the following compounds: 1—Frangulin B, 2—Frangulin A, and 3—Emodin.

A.3.1 Image J Macro scripts

The following Image J macro was modified from a version previously published (Laflamme et al, 2016) to quantify the effect of *F. alnus* in the in vitro mycelial growth experiment. It automatically measures the area in a defined color threshold and saves a painted picture with the selected area for visualization. Adjusted measurements of the Petri dish and the mycelium were used to calculate the percentage of mycelial coverage.

```
\\ originalName = getTitle();
\\run("Duplicate...", "title=Painted");
\\run("Duplicate...", "title=P.i");

\\// Color Thresholder 1.54d
\\// Autogenerated macro, single images only!
\\selectWindow("P.i")
\\min=newArray(3);
\\max=newArray(3);
\\filter=newArray(3);
\\a= originalName;
\\run("HSB Stack");
\\run("Convert Stack to Images");
\\selectWindow("Hue");
\\rename("0");
\\selectWindow("Saturation");
\\rename("1");
\\selectWindow("Brightness");
\\rename("2");
\\min[0]=0;
\\max[0]=255;
\\filter[0]="pass";
\\min[1]=10;
\\max[1]=97;
\\filter[1]="pass";
\\min[2]=130;
\\max[2]=254;
\\filter[2]="pass";
\\for (i=0;i<3;i++){
\\   selectWindow(""+i);
\\   setThreshold(min[i], max[i]);
\\   run("Convert to Mask");
\\   if (filter[i]=="stop")  run("Invert");
\\}
\\imageCalculator("AND create", "0","1");
\\imageCalculator("AND create", "Result of 0","2");
\\for (i=0;i<3;i++){
```

```
\\  selectWindow(""+i);
\\  close();
\\}
\\selectWindow("Result of 0");
\\close();
\\selectWindow("Result of Result of 0");
\\rename(a);
\\// Colour Thresholding-------------
\\
\\//define area to measure
\\makeOval(188, 164, 1035, 1028);
\\run("Analyze Particles...", "size=470-Infinity display clear \\summarize add");
\\
\\// paint it
\\roiManager("Combine");
\\run("Create Mask");
\\run("Create Selection");
\\selectWindow("Painted");
\\run("Restore Selection");
\\run("Fill", "slice");
\\
\\//save the results manually
\\//color threshold and the area to measure can be adjusted
```